**Systematics and geographical distribution of *Galba* species, a group of**

**cryptic and worldwide freshwater snails**

Pilar Alda[1,2,3], Manon Lounnas[2], Antonio A. Vázquez[2,4], Rolando Ayaqui[5], Manuel

Calvopiña[6], Maritza Celi-Erazo[7], Robert T. Dillon Jr.[8], Luisa Carolina González

Ramírez[9], Eric S. Loker[10], Jenny Muzzio-Aroca[11], Alberto Orlando Nárvaez[11,12], Oscar

Noya[13], Andrés Esteban Pereira[14], Luiggi Martini Robles[15], Richar Rodríguez-

Hidalgo[7,16], Nelson Uribe[14], Patrice David[17], Philippe Jarne[17], Jean-Pierre Pointier[18],

Sylvie Hurtrez-Boussès[2,19]

[1] Laboratorio de Zoología de Invertebrados I, Departamento de Biología, Bioquímica y

Farmacia, Universidad Nacional del Sur. San Juan No. 670, B8000ICN Bahía Blanca,

Buenos Aires, Argentina.

[2] MIVEGEC, University of Montpellier, CNRS, IRD, Montpellier, France.

[3] Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Argentina.

[4] Laboratory of Malacology, Institute of Tropical Medicine "Pedro Kourí", Autopista

Novia del Mediodía km 6, La Habana, Cuba.

[5] Departamento de Microbiología y Patología, Facultad de Medicina, Universidad

Nacional de San Agustín de Arequipa, Peru.

[6] Carrera de Medicina, Facultad de Ciencias de la Salud, Universidad de Las Américas,

Quito, Ecuador.

[7] Instituto de Investigación en Salud Pública y Zoonosis - CIZ, Universidad Central de

Ecuador, Quito, Ecuador.

[8] Freshwater Gastropods of North America Project, P.O. Box 31532, Charleston, SC

29417, USA.

[9] Grupo de Investigación "Análisis de Muestras Biológicas y Forenses" Laboratorio

Clínico, Facultad de Ciencias de la Salud, Universidad Nacional de Chimborazo,

Ecuador.

[10] Center for Evolutionary and Theoretical Immunology, Department of Biology,

University of New Mexico, Albuquerque, NM87131, USA.

[11] Instituto Nacional de Investigación en Salud Pública INSPI, Guayaquil, Ecuador.

[12] Universidad Agraria del Ecuador, Facultad de Medicina Veterinaria y Zootecnia,

Guayaquil, Ecuador.

[13] Sección de Biohelmintiasis, Instituto de Medicina Tropical, Facultad de Medicina,

Universidad Central de Venezuela. Centro para Estudios Sobre Malaria, Instituto de

Altos Estudios "Dr. Arnoldo Gabaldón"-Instituto Nacional de Higiene "Rafael Rangel"

del Ministerio del Poder Popular para la Salud. Caracas, Venezuela.

[14] Grupo de Investigación en Epidemiología Molecular (GIEM), Escuela de

Microbiología, Facultad de Salud, Universidad Industrial de Santander, Bucaramanga,

Colombia.

[15] Laboratorio de Parasitologia Luiggi Martini y colaboradores, Guayaquil, Ecuador.

[16] Facultad de Medicina Veterinaria y Zootecnia, Universidad Central del Ecuador,

Quito, Ecuador.

[17] Centre d'Ecologie Fonctionnelle et d'Evolution, UMR 5175, CNRS – Université de

Montpellier – Université Paul Valéry Montpellier – EPHE - IRD, 1919 route de Mende,

34293, Montpellier Cedex 5, France.

[18] PSL Research University, USR 3278 CNRS–EPHE, CRIOBE Université de

Perpignan, Perpignan, France.

[19] Département de Biologie–Ecologie, Faculté des Sciences, Université Montpellier,

Montpellier, France.

To whom correspondence should be addressed: **pilaralda@gmail.com** (P. Alda)

**Key words:** systematics, distribution, America, Lymnaeidae, *Galba*, vector snails,

biological invasions.

**Abstract**

Cryptic species can present a significant challenge to the application of systematic and

biogeographic principles, especially if they are invasive or transmit parasites or

pathogens. Detecting cryptic species requires a pluralistic approach in which molecular

markers facilitate the detection of coherent taxonomic units that can then be analyzed

using various traits (e.g., internal morphology) and crosses. In asexual or self-fertilizing

species, the latter criteria are of limited use. We studied a group of cryptic freshwater

snails (genus *Galba)* from the family Lymnaeidae that have invaded almost all

continents, reproducing mainly by self-fertilization and transmitting liver flukes to

humans and livestock. We aim to clarify the systematics, distribution and phylogenetic

relationships of these species with an integrative approach that includes morphology

(shell and reproductive anatomy), molecular markers, wide-scale sampling across

America, and data retrieved from GenBank (to include Old World samples). Our

phylogenetic analysis suggests that the genus *Galba* originated ca. 22 Myr ago and

today comprises six clusters of species. Four of these clusters (*G. truncatula*, *G.*

*cubensis*/*viator*, *G. humilis* and *G. schirazensis*) are morphologically cryptic and

constitute species or species complexes with wide variation in their genetic diversity,

geographic distribution and invasiveness. The other two clusters constitute a single

species or a species complex (*Galba cousini*/*meridensis*) that demonstrate more

geographically restricted distributions and exhibit an alternative morphology more

phylogenetically derived than the cryptic one. Further genetic studies are required to

clarify the status of both *G. cousini*/*meridensis* and *G. cubensis/viator*. We emphasize

that no *Galba* species should be identified without molecular markers and that

additional sampling is required, especially in North America, Eurasia and Africa to

clarify remaining questions in systematics and biogeography. We also discuss several

hypotheses that can explain crypsis in *Galba*, such as convergence and morphological

stasis, and hypothesize a role for stabilizing selection in amphibious (rather than strictly

freshwater) habitats.

## Introduction

Cryptic species are groups of populations in which the intrapopulation phenotypic variance exceeds interspecific phenotypic variance (see review in Bickford et al., 2007; Fišer et al., 2018; Struck et al., 2018). They have been described in all taxonomic kingdoms, although cryptic species seem to be more often reported in animals (Struck et al. 2018). Their frequency might reach a bit less than 1% of all animal species, an estimate that should be considered quite approximate given our lack of knowledge on the species totals. They also seem to be homogeneously distributed among taxa and biogeographical regions (Pfenninger and Schwenk 2007), although it has been suggested that caves and subterranean habitats may promote cryptic speciation. Such isolated, dark, low-energy habitats promote diversification but constrain morphological differentiation since very specialized adaptations are needed to survive in them (Katz et al. 2018). Four hypotheses have been proposed for the origin of cryptic species (Fig. S1; Bickford et al., 2007; Fišer et al., 2018; Struck et al., 2018): (1) recent divergence, when distinguishing traits have not as yet accumulated (e.g. cave fish, Niemiller et al. 2012); (2) parallelism, where independent phenotypic traits evolve in different taxa from a similar and shared ancestral trait (Struck et al. 2018); (3) convergence, where more distantly related species evolve from dissimilar ancestors (e.g. sea stars, Zulliger and Lessios, 2010), and (4) morphological stasis, where species remain similar over long periods of time constrained by limited genetic variation or stabilizing selection (e.g., Gomez et al., 2004; Struck et al., 2018).

Although interesting as models for study of the speciation process (Coyne and Orr 2004; De Queiroz 2007), cryptic species are problematic from two human perspectives: biological invasion and disease transmission. Cryptic species may exhibit wide differences in invasive ability and impact on invaded ecosystems and communities

(Fang et al. 2014). An accurate identification at the species level is required in such

situations to track invasions and mitigate any harmful consequences (Kolar and Lodge

2001; Dunn and Hatcher 2015; Jarić et al. 2019). Cryptic species may also exhibit

differences in disease transmission. This is the case in the *Anopheles gambiae* complex

which includes the most important vectors of malaria in Africa (Stevenson and Norris

2016). Some members of the complex are broadly zoophilic, while others feed more

strictly on humans. Accurate species identification is required for effective mosquito

control.

Snails, especially freshwater ones, are an interesting group for addressing

biogeographic issues in cryptic species. Although taxonomists have increasingly used

molecular markers over the last decades to identify snail species (Dayrat et al. 2011),

morphological characters, especially shell shape and sculpture, remain widely relied

upon—often resulting in large numbers of synonymous species (Qian et al. 2012). The

shells of some mollusk populations show significant phenotypic plasticity, however, in

reaction to temperature, pollution or predation (Bourdeau et al. 2015). Others stay stable

for millions of years (e.g., Weigand et al., 2011; Weiss et al., 2018). This has resulted

both in the proliferation of species names and descriptions (e.g., Taylor, 2003), most of

which are invalid (Jarne et al. 2010; Dillon et al. 2011), as well as the misidentification

of valid species, yielding errors in the assessment of species invasion ability and

distributional range (e.g., Pfenninger et al., 2006; Rama Rao et al., 2018). For example,

an invasive Asian clam of the genus *Sinanodonta* has been overlooked in Russia

because it is morphologically indistinguishable from another invasive Asian clam

(Bespalaya et al. 2018). Such problems call for an integrated approach to gastropod

systematics and biogeography in which phenotypic traits are studied together with

appropriate molecular tools (Dayrat 2005).

Here we focus on small-bodied basommatophoran pulmonate snails of the genus

*Galba* (Hygrophila, Lymnaeidae), common inhabitants of unstable freshwater habitats

worldwide. Baker (1911) recognized 30 species and subspecies in the subgenera *Galba*

(*s.s.*) and *Simpsonia* in North America, on the basis of minor shell morphological

variation, where Hubendick (1951) suggested that as few as four biological species

might be valid: *humilis*, *truncatula*, *cubensis* and *bulimoides*. The more recent work of

Burch (1982) proposed 22 North American species in the genus *Fossaria*, which we

here consider a junior synonym of *Galba*. In South America, Hubendick (1951)

recognized only two species of small-bodied, amphibious lymnaeids (*viator* and

*cousini*), but more recent work based on molecular approaches distinguishes seven

species (*viator*, *schirazensis*, *cousini*, *neotropica*, *meridensis*, *truncatula* and *cubensis*;

(Bargues et al. 2007, 2011b, 2011a; Correa et al. 2010, 2011; Lounnas et al. 2017,

2018).

Despite an estimated divergence time on the order of 20 Myr based on genomic

data (Burgarella et al. 2015), most of the nominal species of *Galba* share a similar shell

morphology and internal anatomy, as well as common levels of phenotypic plasticity in

shell, anatomy, and life history (Samadi et al. 2000; Correa et al. 2011). The exceptions

are *Galba cousini* (Paraense 1995) and *Galba meridensis* (Bargues et al. 2011b), which

are mutually similar but morphologically distinctive. These two groups are cryptic

species; they are often misidentified and often confused (Correa et al. 2010; Bargues et

al. 2011a).

A further challenge is that crossing cannot be used to distinguish species (Coyne

and Orr 2004), as has been done in other freshwater snails (e.g., *Physa* species, Dillon et

al., 2011), since *Galba* populations primarily reproduce by self-fertilization (Meunier et

al. 2004; Bargues et al. 2011a; Lounnas et al. 2017, 2018). Moreover, at least two

species, *G. schirazensis* and *G. truncatula*, have been shown to be extremely efficient

anthropogenic invaders, muddling our knowledge of species distribution. Populations of

*G. truncatula*, probably from Eurasia, have invaded South America, especially the

Bolivian Altiplano (Meunier et al. 2004). This is of special concern since *Galba*

populations are the main vectors of the liver fluke *Fasciola hepatica* which causes

fasciolosis in both livestock and humans (Mas-Coma et al. 2005) and transmission

efficiency and invasion ability differ among species (Vázquez et al. 2018).

Worldwide, the geographic distribution of *Galba* species is poorly known. The

dubious character of records based on morphological identification leaves us with a

small sample of molecular studies (Correa et al. 2010, 2011; Bargues et al. 2011b,

2011a, 2012; Lounnas et al. 2017, 2018) upon which to base a very large-scale

phenomenon. So, our objectives here are to characterize the geographic distribution of

*Galba* species at continental scale, based on an extensive sampling over America, to

reconstruct the genus phylogeny to delimit species, and to explore the origin of crypsis.

We aim to delineate species—the real scientific challenge of integrative taxonomy

(Dayrat 2005)—noting that in practice *Galba* species are very difficult to delineate due

to its crypsis, wide geographical distribution, and mating system. Previous studies

reconstructing *Galba* phylogeny have used single genes and analyzed fewer than 10

sequences per species, failing to account for the wide geographic distribution of this

genus (Correa et al. 2010, 2011; Bargues et al. 2011c, 2011a; Standley et al. 2013).

Here we employ morphological and molecular markers (microsatellite loci and DNA

sequences from two genes) to study more than 1,700 individual *Galba* from 161 sites.

Our data set was augmented with a complete sample of all the *Galba* DNA sequences

available in GenBank and multiple phylogenetic analyses conducted. We used both

gene trees and multispecies coalescent models to shed light on the phylogenetic

relationships and on the origin of crypsis in the genus *Galba*.

## Materials and methods

### Snail sampling and species identification

We conducted simple searches for *Galba* populations in suitable habitats throughout the

New World over a span of two decades, 1998–2017. Overall, *Galba* populations were

detected in 161 sites and 1,722 individuals were sampled from nine countries:

Argentina, Canada, Colombia, Cuba, Ecuador, France (French Guiana, Guadeloupe and

Martinique), Peru, Venezuela, and USA (Table S1). *Galba* populations have previously

been reported from some of these sites in Venezuela and Ecuador by the authors

(Pointier 2015; Orlando Narváez et al. 2017).

193            In most cases, we discovered our *Galba* populations in unstable habitats subject

194    to frequent flooding and droughts. The sampled habitats were characterized as brook,

195    irrigation canal, ditch, oxbow lake, pond, marsh, lake, tank, rice field, and river.

196    Individual snails were often collected above the water line, consistent with their

197    amphibious habit. Some sites were visited up to five times. Geographic coordinates

198    were recorded for most sites.

199            After collection, individuals were placed in water at 70 ℃ for 30–45 s. This

200    procedure allows fixation of individuals without contraction of soft parts and facilitates

201    a proper study of snail internal anatomy. The body was carefully withdrawn from the

202    shell using forceps and both body and shell stored in 70% ethanol until morphological

203    and DNA analyses (Pointier et al. 2004).

204            Species were characterized using a three-step procedure involving both

morphological and molecular markers (Fig. 1). Step 1 was an analysis of shell

morphology and reproductive anatomy. In step 2, we used a molecular tool that enables

us to distinguish *G. cubensis*, *G. schirazensis*, and *G. truncatula* (Alda et al. 2018). In

step 3, we sequenced mitochondrial and nuclear genes in individuals for which no PCR

amplification product was obtained in step 2. DNA from some of those individuals

identified in steps 1 and 2 were also sequenced in order to reconstruct the phylogeny of

*Galba*. Note that this three-step approach is less expensive than an approach based on

simply sequencing the same genes in all individuals.

*Step 1: morphology of the shell and of internal organs*

We photographed the shell of three to five adult snails from each site and dissected their

body under a stereoscopic microscope. We drew the anatomy of the penial complex,

prostate, and renal tube using a camera lucida attachment (Pointier et al. 2004). We did

not record any morphological measurements or perform any quantitative tests because

Correa et al. (2011) have shown that cryptic *Galba* species cannot be delimited by such

means. Our observations were qualitative only (Fig S2).

*Step 2: multiplex PCR of microsatellite loci*

We applied the multiplex PCR test designed by Alda et al. (2018) to all the 1,420

individuals that were not distinguishable based on shell and reproductive anatomy. This

method is based on species-specific primers amplifying three microsatellite loci (one

each per targeted species) and producing band sizes that are specific to these species

(179–200 pb in *G. cubensis*, 227–232 pb in *G. schirazensis* and 111–129 pb in *G.*

*truncatula*). DNA was extracted using a Chelex protocol following Estoup and Martin

(1996) as adapted for 96-well plates. Methods for DNA amplification and

electrophoretic resolution followed Alda et al. (2018).

*Step 3: identification by sequencing*

We amplified the internal transcribed spacer 2 (ITS2) and the cytochrome oxidase

subunit 1 (COI) genes in 35 individuals sampled from 15 sites in Argentina, Canada and

the USA (1 to 5 per population) where at least some individuals did not demonstrate an

amplification product in step 1, using the method of Lounnas et al. (2017, 2018). We

also amplified ITS2 and COI in 112 individuals that did return an amplification product

in step 1, including one individual identified as *G. cousini/meridensis*. To supplement

these results, we also amplified the internal transcribed spacer 1 (ITS1) and the ARN

ribosomal 16S in two individuals of *G. cubensis* from Bosque del Apache (USA) and in

one individual of *G. cousini/meridensis* from Ecuador (Table S1). With this approach,

we obtained at least one sequence from each hypothetical species represented by the

four genes and used them to delimit species. The total number of individuals amplified

was 151.

We used the primers NEWS2 (forward) 5' TGTGTCGATGAAGAACGCAG 3'

and ITS2-RIXO (reverse) 5' TTCTATGCTTAAATTCAGGGG 3' to amplify ITS2;

Lim1657 (forward) 5' CTGCCCTTTGTACACACCG 3' and ITS1-RIXO 5'

TGGCTGCGTTCTTCATCG 3' to amplify ITS1 (Almeyda-Artigas et al. 2000);

LCOI490 (forward) 5' GGTCAACAAATCATAAAGATATTGG 3' and HCO2198

(reverse) 5' TAAACTTCAGGGTGACCAAAAAATCA 3' to amplify COI (Folmer et

al. 1994) and forward 5' CGCCTGTTTATCAAAAACAT 3' and reverse 5'

CCGGTCTGAACTCAGATCACGT 3' to amplify 16S (Remigio and Blair 1997). In all

cases, PCR amplification was performed in a total volume of 25 µl containing 12.5 µl of

Taq PCR Master Mix (Qiagen), 2.5 µl of each primer (10 mM) and 2 µl of DNA in an

Eppendorf Thermal Cycler with an initial denaturation step at 95 ℃ for 15 minutes;

followed by 35 cycles of denaturation at 95 ℃ for 30 seconds, annealing at 50 ℃ for

one minute, extension at 72 ℃ for one minute; and a final elongation at 60 ℃ for 30

minutes. The presence and size of amplification products were electrophoretically

confirmed in 1% agarose gels stained with EZ-Vision. DNA sequencing was performed

by Eurofins Genomics (Ebersberg, Germany) using PCR-amplified products as

templates. All sequences were uploaded to GenBank (Table S1) and assigned to a

species using the phylogenetic reconstruction.

**Type localities**

Because of the longstanding confusion and uncertainty regarding the systematics of the

genus *Galba* worldwide, it is especially important to establish standard populations,

against which unknown populations can be compared. Type localities were specified by

the authors of all the more recently-described species, such as *neotropica* (Bargues et al.

2007) and *meridensis* (Bargues et al. 2011b), and others have been established by

subsequent use, for example *schirazensis* (Bargues et al. 2011a). But in his original

description of *Limnaeus viator*, D'Orbigny (1835) simply stated "Patagonia." And

Pfeiffer (1839) gave no locality data for his *Limnaeus cubensis* at all, beyond "Cuba".

In such circumstances, the ICZN code provides that subsequent authors may restrict

type localities to some more precise spot "within the known range of the taxon". Type

localities for all eight of the widely-recognized species in the genus *Galba*, either as

originally stated or as subsequently restricted, are listed in Table 1. COI sequences from

samples of all the populations inhabiting these localities have been previously uploaded

to GenBank, and most have ITS1, ITS2 or 16S sequences available as well.

**Retrieving data on *Galba* spp. distribution from published work**

We searched the literature and GenBank for sequence data at four genes (COI, ITS1,

ITS2, and 16S) apparently attributable to lymnaeids of the genus *Galba*. Coordinates

were provided for most sites by the authors. When coordinates were not provided, we

inferred them from the locality data using GoogleEarth. We found 132 New World sites

in which *Galba* species have been molecularly characterized (Table S2), and 45 sites in

the Old World (Table S3, Fig S3). The specific nomina attributed to these sequences by

their depositors in GenBank were 157 *truncatula*, 152 *schirazensis*, 70 *neotropica*, 57

*cubensis*, 44 *viator*, 20 *cousini*, 9 *humilis*, 6 *meridensis*, and 2 others. This corresponds

to 166 COI, 163 ITS2, 118 ITS1, and 70 16S sequences.

**Phylogenetic and ancestral reconstruction study**

Phylogenetic analyses were conducted on the ITS2 and COI sequences obtained in this

study, together with ITS2, COI, ITS1, and 16S sequences retrieved from GenBank.

Most of these sequences have been previously published (Tables S2–S3). Ultimately,

we included 796 sequences in our analyses: 90 for 16S, 251 for COI, 122 for ITS1, and

333 for ITS2. Some GenBank accession numbers appear more than once in Tables S2–

S3 because individuals with identical sequences have been registered under the same

GenBank accession number.

Alignment was performed individually for each gene using MAFFT (Katoh and

Standley 2013). Ambiguously aligned sites were excluded using GBLOCKS with

default settings for a less stringent selection (Castresana 2000). The number of positions

in the final sequences was 412 for 16S (83% of the original 493 positions), 609 for COI

(86% of 707), 435 for ITS1 (48% of 888), and 333 for ITS2 (23% of 1,429). We

examined levels of saturation for each gene and for the first and second *versus* third

codon positions of COI using DAMBE (Xia 2017). We did not find evidence of

saturation in the four genes analyzed, including all codon positions of COI. We

partitioned codon positions and unlinked substitution models in phylogenetic analyses.

We used Bayesian inference in Beast2 (Bouckaert et al. 2014) for five reasons:

(1) to assign sequences to species; (2) to validate, and to mend if necessary, species

identity for sequences retrieved from GenBank; (3) to delimit species; (4) to reconstruct

the phylogeny and estimate time divergence; and (5) to assess the ancestral phenotypic

state of *Galba* species. Because we did not have identical sets of individuals (or

populations) across genes it was necessary to build four unlinked gene trees to address

questions (1) and (2). The best-fitting models of sequence evolution for each gene were

selected using bModelTest (Bouckaert and Drummond 2017). We estimated a model for

each COI partition (1st, 2nd, and 3rd position). The best model describing the evolution of

16S was HKY+G+I, 123424+G+I for COI (1st codon), 121321+G+I for COI (2nd

codon), TN93+G+I for COI (3rd codon), 123424+G+I for ITS1, and 121323 for ITS2.

We linked trees for the three COI codon partitions. The analyses were run using four

gamma categories and a proportion of 0.5 invariant sites. Uncorrelated relaxed-clock

models were chosen for all loci. The relative clock mean priors were all lognormal (M =

0, S = 1). We used a birth-death model as priors with lognormal birth and death rates.

These gene trees allowed us to evaluate species names and validate or mend species

identity for all sequences, whether obtained in this study or retrieved from GenBank.

All the MCMC were run for 200,000,000 generations storing every 20,000 generations.

The names of the eight widely-recognized species in the genus *Galba* were ultimately

assigned by reference to their type localities. Since gene trees can be erroneously

inferred due to incomplete sampling, incomplete lineage sorting or introgression

between lineages, we also built haplotype networks for each gene using popART (Leigh

and Bryant 2015) and compared them with gene trees.

To address our question (3), species delimitation, we built ten multispecies

coalescent tree models in StarBeast2 (Ogilvie et al. 2017) differing in species

assignments, some models splitting species to as many as nine while others lumping to

as few as five. We assigned the species identity obtained from tasks 1 and 2 to each

individual and tested each scenario in turn. We also created an eleventh model in which

we separated the populations of *G. viator* from Argentina and from Chile to test whether

splitter models showed higher support than lumper models regardless of their biological

meaning. Since all the hypothetical species generated in these tree models must be

represented by all genes, we removed the sequences from Ethiopia in this analysis.

In all our multispecies coalescent tree models, we used the same site models as

for reconstructing the gene trees. We assigned to the mitochondrial loci a gene ploidy of

0.5 and to nuclear loci a gene ploidy of 2.0 (diploid). We used constant population sizes

(fixed to 1) and uncorrelated relaxed-clock models for all loci. For nuclear loci, we used

the molecular clock rate reported by Coleman and Vacquier (2002) for ITS in bivalves

(0.00255 per Myr). For mitochondrial loci, we retained the molecular clock rate

estimated by Wilke et al. (2009) for COI in invertebrates (0.0157 per Myr). We used the

birth-death model as tree prior with lognormal birth and death rates. We ran each model

using Multi-Threaded Nested Sampling analysis with 10 particles, 8 threads, a chain

length of 100,000,000 and sub-chain length of 5,000. Then, we compared the trees by

computing Bayes factor (BF), a model selection tool that is simple and well-suited for

comparing species-delimitation models (Leaché et al. 2014). BF is the difference

between the marginal likelihoods of two models: BF = model 1 – model 2. If BF is

larger than 1, model 1 is favored, and otherwise model 2 is favored. When BF is

between 20 and 150 the support strong and when the BF is above 150 the support is

overwhelming (Leaché et al. 2014). We used Nested Sampling implemented in the NS

package to calculate the marginal likelihoods necessary to obtain BF and also an

estimate of the variance of the marginal likelihoods (Russel et al. 2019).

We also applied two other species-delimitation methods: Automatic Barcode

Gap Detection (ABGD; Puillandre et al. 2012) and Species Tree and Classification

Estimation in Beast2 (STACEY, Jones 2017). ABGD was run for each gene using the

default settings (https://bioinfo.mnhn.fr/abi/public/abgd/). The prior on intraspecific

divergence defines the threshold between intra- and interspecific pairwise distances and

is iterated from minimum to maximum through ten steps (Puillandre et al. 2012). Given

the wide genetic diversity observed within some species (Lounnas et al. 2017), we

chose the partition that showed high prior on intraspecific divergence (the penultimate

partition).

For the STACEY delimitation method, we used only the ITS2 and COI

sequences because these both genes were better represented in the dataset than ITS1 and

16S. Since the hypothetical number of species in STACEY ranges from one to the

number of individuals, each of our 113 snail populations was considered as a minimal

cluster. The method distinguishes very shallow species divergences with a statistic

called "collapseHeight," which we set to a small value (0.0001) following Jones (2017).

Given the results obtained with ABGD and Nested Sampling analyses, we ran three

multispecies coalescence models with different collapseWeight parameters: (1) a

lumping model with five *Galba* species with 1/X distribution (initial value: 0.975,

between 0 and 1), (2) a splitting model with nine species with 1/X distribution (initial

value: 0.952 between 0 and 1), and (3) a no-prior-taxonomic model with a Beta

distribution ($\alpha = 2$, $\beta = 2$). The initial value of the lumping and splitting models were

calculated following Matos-Maravi et al. (2018). All other parameters were set as in the

Nested Sampling analysis. We ran the three models for 250,000,000 generations with

storing every 25,000 generations.

To address our question (4), regarding species topology and divergence time, we

reran the multispecies tree model that showed the highest Bayes Factor in StarBeast2.

We used the same parameters as in the Nested Sampling analysis, but we ran the

MCMC for a longer time (250,000,000 generations stored every 25,000 generations).

The MCMC output was visualized using Tracer (Rambaut et al. 2018) and tree samples

summarized by TreeAnnotator (utility program distributed with the Beast package)

using a 10% burn-in. The species tree was visualized and edited in FigTree, GIMP

(https://www.gimp.org), and DensiTree (Bouckaert and Heled 2014). Some analyses

were run in CIPRES Science Gateway (Miller et al. 2012). Note that we did not add an

outgroup to root trees because *Galba* monophyly has already been ascertained (Correa

et al. 2010, 2011).

Finally, to address our question (5), the ancestral phenotypic state, we applied

Bayesian Binary MCMC (BBM, Ronquist and Huelsenbeck 2003), statistical dispersal-

vicariance analysis (S-DIVA; Yu et al. 2010), and Statistical dispersal-extinction-

cladogenesis (S-DEC; Ree and Smith 2008) in the software Reconstruct Ancestral State

in Phylogenies (RASP, Yu et al. 2015). We used the splitting model (scenario A) under

default settings. We added two phenotypic states: one for all cryptic species and another

one for *G. cousini*/*meridensis*.

**Results**

**Morphology**

Most individuals (N = 1,420 from 133 sites) were not distinguishable based on shell and

reproductive anatomy (Fig. S4). The exception was a single group comprising all

individuals from *G. cousini* and *G. meridensis* (N = 302). These tended to demonstrate

more globose shells with shorter spires, adult sizes in excess of 10 mm standard shell

length. *Galba cousini* and *G. meridensis* also differed from the other species in their

internal anatomy—a ureter with two distinct flexures, a wider and more ovate prostate,

a larger penial complex, and a penis sheath approximately the same length as the

preputium (Fig. S4). We did not find any anatomical differences between *G. cousini* and

*G. meridensis*, however, comparing individuals from Ecuador, Colombia, and

Venezuela.

**Multiplex PCR of microsatellite loci**

DNA from the 1,420 American individuals with similar phenotypes was amplified using

the multiplex PCR procedure (step 2 from Fig. 1). We identified 541 individuals of *G.*

*cubensis*, 330 of *G. schirazensis*, and 349 of *G. truncatula* (Table S1). No amplification

was observed in 200 individuals sampled in one site from Argentina and 14 sites from

Canada and USA (Table S1).

**Identification by sequencing**

Phylogenetic analysis of COI sequences returned six clusters (Fig. S5). Clusters I–V

each contained a single type population: *truncatula* (I), *schirazensis* (II), *humilis* (III),

*cousini* (IV), and *meridensis* (V). Cluster VI contained the type populations of *cubensis*,

*neotropica*, and *viator*. The posterior probabilities (PP) of all clusters were 1.0, except

for cluster III (*humilis*, PP = 0.95).

Analysis of 16S, ITS1, and ITS2 sequences confirmed the COI results in almost

all respects, although sequences were missing for at least one or two type populations in

each tree (Figs. S6–S8). However, we detected discrepancies between the mitochondrial

and nuclear gene trees. A striking example are sequences from Bosque del Apache

(USA; cluster VI) that exhibited very long branches in the mitochondrial gene trees

(Figs. S5–S6) but clustered tightly with the *cubensis* type population and similar

populations in the nuclear gene trees (Figs. S7–S8). The mitochondrial sequences from

Ethiopia (one per gene) also formed a long branch, although nuclear sequences were not

reported in GenBank for that population. Within cluster VI, we found that the *cubensis*

and *neotropica* type populations were located in separate subclades for the nuclear

genes but clustered together in the mitochondrial trees (Figs. S5–S8).

Gene trees (Figs. S5–S8) and haplotype networks (Figs. S9–S12) showed that

genetic diversity varied among the six clusters and four genes. Cluster II (*schirazensis*)

showed reduced variation, while cluster VI (*cubensis*, *neotropica*, and *viator*) was larger

and more diverse. Mitochondrial genes seemed more diverse than nuclear genes. This

observation may be biased, however, by a structural correlation in our data between

genes sequenced and regions sampled.

Most of the sequences uploaded to GenBank identified as one of the eight

species of *Galba* were accurately clustered into the six clades containing their type

populations. However, there were some exceptions. Eight sequences of COI uploaded

as *G. truncatula* from France appeared in cluster II with the *G. schirazensis* type

population (Table S3; Fig. S5). We reidentified these sequences as belonging to *G.*

*schirazensis*. The COI sequence from Ethiopia, uploaded as *G. truncatula*, clustered at

the base of *truncatula* clade I with low posterior probability (Fig. S5). The other

Ethiopian sequence (16S), also identified as *G. truncatula*, did not cluster with any of

the *Galba* clades (Fig. S6).

**Species delimitation**

Figure 3 illustrates the results obtained using the three species-delimitation methods.

The Multi-Threaded Nested Sampling analysis (Fig. S13) suggested that scenario A

(nine species) is the best fit to the available data, demonstrating the largest maximum

likelihood estimate (Fig. 3; see Table S4 for Likelihoods). BF analysis preferred

scenario A over scenario D (current taxonomy) or scenario K, separating populations of

*G. viator* from Argentina and from Chile. This suggests that further splitting the

phylogeny of *Galba* (here to consider that *G. meridensis* and *G. cousini* are separate

species) is not required.

ABGD results varied depending on the gene analyzed (Fig. 3). Nine species

(scenario A) were suggested by ITS1, while six species only were returned by our

analysis of ITS2 and 16S, with *G. cubensis*, *G. viator*, *G. neotropica*, and *Galba* sp.

"Bosque del Apache" lumped. ABGD analysis of the COI gene indicated that *G. viator*

and *Galba* sp. "Bosque del Apache" are separate species, but that *G. cubensis* and *G.*

*neotropica* should be lumped together. The ITS2 and COI analyses also suggested that

some species (*G. cousini* and *G. truncatula*) might be represented by more than one

taxon.

The species-delimitation analysis implemented in STACEY suggested that six of

the nine clusters of scenario A might include more than one taxon. The exceptions were

*G. viator*, *G. meridensis*, and *Galba* sp. "Bosque del Apache," the last two species

including only one population. Our STACEY results converged towards similar

MCMCs regardless of which prior was used for the collapseWeight parameter (Fig. 3).

**Phylogeny, time of divergence, and state reconstruction**

Clusters III (*humilis*), IV (*cousini*), and V (*meridensis*) were grouped together in all

gene trees returned by our analysis. But inconsistent results among genes were obtained

for the remainder of the other clusters identified (Figs. S5–S8). The multilocus

multispecies tree returned three major groups: cluster I (*truncatula*) together with II

(*schirazensis*); cluster III (*humilis*) together with IV (*cousini*) and V (*meridensis*); and

the cluster VI group (*viator/cubensis/neotropica/sp.* "Bosque del Apache"). All groups

showed posterior probabilities of 1.0 except for the first one (0.44; Fig. 4). The

multilocus multispecies tree visualized in DensiTree (Fig. 3) confirmed that most tree

topologies united the clusters into the three major groups outlined above, although some

topologies placed clusters differently reflecting the incongruence found among the gene

trees.

The estimated divergence time from the most recent common ancestor of the

*Galba* group was 22.6 Mya [95% HPD interval: 14.6–33; Figs. 3–4]. Diversification

within the *G. cousini/meridensis* species complex and the *G. cubensis/viator* species

complex seems to have occurred 5 Mya ago, or less. The three phenotypic state

reconstruction analyses (S-DEC, S-DIVA, and BBM) suggested that the most recent

common ancestor of *Galba* species displayed the cryptic phenotype (Fig. 4). Thus, the

phenotype of *G. cousini* and *G. meridensis* should be considered as derived from the

cryptic phenotype.

**Discussion**

***Galba* comprises six clusters**

Here we report the largest study published to date of *Galba* systematics and distribution,

based on extensive sampling at a very large geographical scale and integration of

phenotypic and molecular approaches across all DNA sequences available in GenBank

for four genes (Dayrat 2005). The widespread occurrence of self-fertilization in these

populations voids the biological species concept (Coyne and Orr 2004), and the absence

of any reliable morphological distinction obviates the typological one, with the

exception of *G. cousini/meridensis*. Thus, we are left with a phylogenetic approach,

which suggests the existence of six clusters, perhaps corresponding to as many as eight

species, or as few as five. These findings reinforce several previously-published works

that have involved fewer genes and smaller sample sizes (Correa et al. 2010, 2011;

Bargues et al. 2011b, 2011a; Standley et al. 2013; Lounnas et al. 2017).

We suggest that three of our clusters are best understood as one species each.

The oldest taxonomic names for these are *Galba truncatula* (Müller 1774) for cluster I,

*Galba schirazensis* (Küster 1862) for cluster II, and *Galba humilis* (Say 1822) for

cluster III. The other three clusters (IV to VI) may in fact correspond to just two highly-

diverse species or species complexes.

Cluster IV (*Galba cousini* Jousseaume 1887) and cluster V (*Galba meridensis*

Bargues, Artigas, Khoubbane & Mas-Coma 2011) may be united into a single species

complex, with *cousini* the oldest name. All these populations inhabit northern regions of

South America, and together they exhibit a single phenotype not shared by other

populations of *Galba*. They always clustered together in our phylogenetic

reconstructions, with a divergence time of 4.7 Mya. However, *G. meridensis* has been

sampled in but a single locality, and more extensive sampling is required to ascertain

whether these are two species or a single species complex.

Cluster VI, including the nomina *Galba viator* (d'Orbigny 1835), *Galba*

*cubensis* (Pfeiffer 1839), *Galba neotropica* (Bargues, Artigas, Mera y Sierra, Pointier &

Mas-Coma, 2007), and a population from southern USA (Bosque del Apache) seem to

comprise *Galba* species number five. The distance separating *G. cubensis* and *G.*

*neotropica* is limited (1 Mya). Moreover, microsatellite markers defined in *G. cubensis*

amplified effectively in individuals of *G. neotropica* (Lounnas et al. 2017), suggesting a

very short genetic distance. We therefore suggest synonymy of these two species under

the name *G. cubensis*. *Galba viator* and *G. cubensis* have overlapping distributions in

Argentina, Chile, and Uruguay (Artigas et al. 2011; Sanabria et al. 2013; Standley et al.

2013; Medeiros et al. 2014), and additional sampling in this region would help resolve

their genetic relationships. Here any distinction between the two sets of populations

depends on both the genes and phylogenetic methodology employed. The population

from Bosque del Apache (USA) was also included in cluster VI, depending on the gene

analyzed. Its long mitochondrial branch may suggest rapid evolution, and here again

more extensive sampling is required before reaching any conclusions. On the whole, a

cautious position would be to suggest that cluster VI corresponds to a species complex

or a species with wide diversity, as has been found in other freshwater snails from the

clade Hygrophila (e.g., Ebbs et al., 2018; Mavárez et al., 2002; Pfenninger et al., 2006).

We also note that if we ultimately recognize a single species; its name should be *viator*,

and not *cubensis*, based on prior description.

The *Galba* species tree that we constructed based on a multispecies coalescent

model returned three groups: one group uniting *G. truncatula* and *G. schirazensis*,

another uniting *G. humilis* and *G. cousini/meridensis* and the last with *G.*

*cubensis/viator*. This result partially agrees with some gene trees published in previous

work (Correa et al. 2010, 2011; Bargues et al. 2011a). However, previous trees were

based on single genes and smaller sample sizes. Our phylogenetic analysis also revealed

some distinctive branches, including the mtDNA sequences from Ethiopia (Dayrat et al.

2011) and the samples from Bosque del Apache. These results may suggest undetected

species or accelerated molecular evolution (Fourdrilis et al., 2016; Pinceel et al., 2005;

Thomaz et al., 1996).

In North America, Burch (1982) recognized 22 species of small, mud-dwelling

lymnaeids, which he grouped into the genus *Fossaria* with two subgenera, *Fossaria*

(*s.s.*) with 11 species and *Bakerilymnaea* with 11 (see Table S5 for species names).

Johnson et al. (2013) transferred these species to the genus *Galba*, but otherwise

retained the Burch system. Included in the present analysis were topotypic samples of

*obrussa* from Philadelphia and *parva* from Cincinnati, both of which we here show to

be indistinguishable from topotypic *humilis*, collected at Owego, New York. Remigio

(2002) contributed to Genbank a 16S sequence from a Canadian population he

identified as *Fossaria obrussa*, also grouping with cluster III (*humilis*). We suggest that

*obrussa*, *parva*, and the seven other uniquely North American specific nomina listed

above ascribed by Burch (1982) to the subgenus *Fossaria* are junior synonyms of *G.*

*humilis*, setting aside American populations of *G. truncatula* as distinct. In addition to

his *obrussa* sequence, Remigio (2002) contributed a 16S sequence from Oklahoma to

Genbank which he labelled "*Fossaria bulimoides*". This sequence grouped with cluster

VI (*cubensis*/*viator*) in our analysis. We suggest that all 12 specific nomina ascribed by

Burch (1982) to the *Fossaria* subgenus *Bakerilymnaea* (Table S5) including *bulimoides*

(Lea 1841), are junior synonyms of *G. cubensis*/*viator*.

**A set of cryptic species**

Our study has confirmed the previous reports of Samadi et al. (2000) and Correa et al.

(2011) that most species of *Galba* cannot be distinguished on the basis of shell

morphology or internal anatomy. Trait variability within species seems to be greater

than variance among species, likely attributable to phenotypic plasticity (Correa et al.

2011). Reproductive and growth traits in *G. truncatula* have been shown to vary

according to habitat characteristics at small geographical scales (Chapuis et al. 2007),

suggesting both that life-history traits are phenotypically plastic, and that to rely on such

traits for specific identification is not advisable.

Our study also confirms that *G. cousini*/*meridensis* differs strikingly in adult

size, shell shape, and anatomy from all other *Galba* species. Our molecular evidence

suggests that *G. cousini/meridensis* evolved from a *Galba* ancestor demonstrating the

cryptic phenotype. Interestingly, *G. cousini*/*meridensis* is the largest species within the

genus *Galba*, occurs in a specialized habitat and displays the complex reproductive

anatomy of an outcrossing species (see Jarne et al. 2010; Escobar et al. 2011).

Among the freshwater pulmonates, crypsis has previously been documented in

*Ancylus* (Weiss et al. 2018) and *Radix* (Pfenninger et al. 2006). Our methods here were

strictly qualitative, as was the case for *Ancylus* and *Radix*, because previous studies

(Samadi et al. 2000; Correa et al. 2011) have shown that the dimensions of internal

organs depend on physiological state and hence that species cannot be distinguished by

means of such measurements. Future, more comprehensive approaches should include

other more discrete anatomical traits, such as radular morphology.

Four hypotheses have been offered to explain the occurrence of cryptic species:

recent divergence, parallelism, convergence, and stasis (Fig S1, Bickford et al. 2007;

Fišer et al. 2018; Struck et al. 2018). The recent divergence hypothesis seems unlikely

in this case. *Galba* has no closely related groups; its closest relatives are probably the

stagnicoline lymnaeids of North America and Eurasia, which demonstrate a very

distinctive morphology (Correa et al. 2010). Our analyses suggest that the several

species of *Galba* are separated by more than 20 Myr (Burgarella et al. 2015). And

indeed, the morphological divergence demonstrated by *G. cousini*/*meridensis* suggests

that time is not been a significant constraint. The parallelism hypothesis also seems

unlikely given that, based on our phylogenetic reconstruction, the cryptic morphology is

ancestral for *Galba*, and the only other morphology that has evolved in the group, as

demonstrated by *G. cousini*/*meridensis,* is derived. Nor does the topology of lymnaeid

phylogeny fit the convergence hypothesis (Correa et al. 2010). So, by default,

morphological stasis is left as the most likely hypothesis to explain crypsis in the genus

*Galba*, as has been proposed in other gastropod groups (e.g., Gomez et al., 2004; Struck

et al., 2018).

Strong stabilizing selection related to specialized environmental conditions is

one possible explanation for the morphological stasis demonstrated by our several

species of *Galba*. Populations have colonized habitats that are not exploited by other

freshwater snails, especially the transiently inundated margins of water bodies exposed

during the dry season(s) (see Table S1). In other words, *Galba* populations are more

amphibious than other freshwater snails, which might mitigate both predation and

interspecific competition (Burgarella et al. 2015). Adaptation to such habitats might

impose strong stabilizing selection for a shell morphology able to resist desiccation and

concomitant morphological stasis.

Among the many adaptations that have been correlated with life in transitory

environments such as the exposed margins of water bodies is the evolution of self-

fertilization (Escobar et al. 2011; Burgarella et al. 2015). The stasis in reproductive

anatomy we have documented in populations of *Galba* may simply reflect a shared

adaption to self-fertilization in unpredictable habitat patches (Jarne et al. 2010). Once

established, self-fertilization promotes the rapid erosion of genetic variation, as inbred

populations become progressively unable to generate new genetic combinations through

recombination (Noël et al. 2017). Thus, selfing could reinforce morphological stasis that

might have evolved for other reasons.

The challenge of identifying cryptic *Galba* species is aggravated by their wide

and poorly-known geographical distributions, recently scrambled by biological

invasion. For example, *G. schirazensis* and *G. truncatula* have broadly expanded their

distribution over recent decades (Brown 1994; Bargues et al. 2001, 2011a; Vinarski and

Kantor 2016; Lounnas et al. 2018). We have documented up to three *Galba* species

occurring in some South American sites (Table S1). Their identification is not possible

without molecular tools.

The specific identity of *Galba* populations is important because they are

involved in the transmission of fasciolosis caused by the liver fluke *F. hepatica*. Some

studies have shown that lymnaeid species demonstrate different patterns of

susceptibility, host-parasite compatibility and immunological resistance to *F. hepatica*

(Gutiérrez et al. 2003; Vázquez et al. 2014; Dreyfuss et al. 2015). Although all species

can be infected under laboratory conditions (Vázquez et al. 2018), field transmission

depends on ecological and sociological conditions. Cattle or wildlife do not occupy the

same grazing habitats as infecting snails in many parts of the world (Sabourin et al.

2018). Ecological studies should be performed to evaluate whether the several cryptic

*Galba* species differ with regard to habitat preference, since our current knowledge is

essentially limited to *G. truncatula* (Chapuis et al. 2007).

**Conclusions and future directions**

Lymnaeid populations of the genus *Galba* are of interest for addressing a variety of

questions, including wide-scale biogeography, biological invasions, evolution of mating

systems, and host-parasite interactions. Our work is a first attempt to clarify the

phylogeny, systematics, and biogeographical distribution of this interesting group in the

New World. We have constructed a variety of gene trees using classical approaches, as

well as a species tree based on a multispecies coalescent model that reconciles gene

trees and provides a much better estimation accuracy for species tree topology than, for

instance, concatenation (Heled and Drummond 2010). The inferred phylogenetic

relationships among species varied, depending on the genes analyzed and techniques

employed. Such a discordance is not unusual and has been reported in many different

studies (e.g., Kutschera et al., 2014; Stewart et al., 2014; Suh et al., 2015), including in

mollusks (Krug et al. 2013; Sales et al. 2013). Incomplete lineage sorting or

introgressive hybridization of specific genes may indeed lead to such a result

(Felsenstein 2004). Future studies could investigate which evolutionary processes (gene

duplication, horizontal gene transfer, incomplete lineage sorting, hybridization) gave

rise to the incongruence we have observed in gene and species trees.

The integration of fossil data would also likely provide more accurate insights

into macroevolutionary dynamics of *Galba* than molecular phylogenies alone. The

fossil record has proven very helpful in determining the phylogeny of groups containing

hard body parts like plants, mammals, and mollusks (Magallón and Sanderson 2005;

Agnarsson et al. 2011; Bolotov et al. 2016). But unlike marine gastropods that have a

diverse fossil record with many helpful shell characters (sculptures, ornamentation,

coloration, protoconch; Bouchet and Strong 2010), fossil freshwater pulmonates are

scarce and lack useful taxonomic characters. The fossils that have been attributed to

*Galba* are not strikingly distinct from those of other species belonging to other genera

of Lymnaeids. Baker (1911) reported *Galba*-like fossils in the Cretaceous of North

America, but these shells might also belong to other genera of Lymnaeids, or even an

ancestor of current Lymnaeids. When the generalized shell morphology is added to the

absence of anatomical data, the application of fossil data to *Galba* phylogenetic

reconstruction becomes excessively problematic.

Although our study was conducted at an extremely large geographic scale,

especially in America, *Galba* populations occur on almost all continents. Much more

extensive sampling and molecular analysis will be required to get a worldwide picture

of the phylogeny and distribution of the genus. Of special interest is North America,

where we have confirmed the occurrence of *G. humilis*, *G. cubensis/viator*, and *G.*

*schirazensis*. We did not, however, confirm *G. truncatula*, despite its otherwise

worldwide distribution (Bargues et al. 2001; Correa et al. 2011; Vinarski et al. 2011;

Novobilský et al. 2014). The recent arrival of *G. schirazensis* in Europe and the Middle

East (Bargues et al. 2011a; Lounnas et al. 2018) and the recent description of *Galba*

*robusta* from Yemen, based on shell and penial morphology alone (Vinarski 2018)

should be confirmed with molecular approaches.

Broader, worldwide sampling will be required to (i) confirm that the European,

African and Asian populations referred to as *G. truncatula* actually belong to this

species, (ii) clarify the biogeographic origin of *G. truncatula*, (iii) evaluate whether the

*G. cubensis/viator* and *G. cousini/meridensis* groups constitute species complexes or

single species with wide diversity, (iv) understand how the invasive species (especially

*G. truncatula* and *G. schirazensis*) are spreading, and (v) confirm the loss of genetic

variation in those populations as reported by other authors (Meunier et al. 2001;

Lounnas et al. 2018).

It might also be useful to construct maps of *Galba* absence, possibly associated

with detection probabilities. For example, *Galba* has not been found in Northern Brazil

(Paraense 1982), possibly because the acidic waters of the Amazon and Orinoco rivers

have blocked colonization (Paraense 1982, 1983). *Galba* populations are also absent

from the most southern regions of Argentina (authors' unpublished data). Negative

records in species distribution can be important when constructing accurate distribution

maps and inferring species-environment associations (Brotons et al. 2004). Models

based only on presence data are less predictive (Václavík and Meentemeyer 2009).

Absence data have been used to good effect modelling the spatial distribution of the

medically important freshwater snail species *Bulinus globosus*, *Biomphalaria pfeifferi*,

and *Radix natalensis* in Zimbabwe and predicting their distribution in a future climate

(Pedersen et al. 2014).

More detailed ecological studies, based on long-term surveys of sites and

analyses of life-history traits under laboratory conditions (e.g., Chapuis et al., 2007)

would facilitate our understanding of species interactions and the transmission of *F.*

*hepatica* (Sabourin et al. 2018). For example, *G. schirazensis* seems to be spreading

very efficiently and genetic studies suggest that this is mainly due to one genetic clone

(Lounnas et al. 2018). Studying the competitive and transmissive abilities of this clone,

compared to both other *G. schirazensis* and other *Galba* species would be worthwhile.

**717 Data and code accessibility**

718 Xml files for phylogenetic analyses are available from the Zenodo repository

719 (https://zenodo.org/record/3473937#.XZiPcC0ryTd).

**721 Acknowledgments**

We would like to express our gratitude to Nicolás Bonel for useful comments on earlier

drafts of the manuscript and Harry G. Lee for advice and assistance on the taxonomy.

We thank Jimena Guerrero, Björn Stelbrink and Thomas Wilke for suggestions on

phylogenetic analyses and for Graham R. Jones, Patricio Maturana Russel and Remco

R. Bouckaert for assistance in running STACEY and Multi-threaded nested sampling.

We thank the reviewers Pável Matos-Maraví and Christelle Fraïsse and the

recommender from PCI in Evolutionary Biology Fabien Condamine for their thoughtful

comments and suggestions. Fellowships granted by Erasmus Mundus PRECIOSA and

Méditerranée Infection supported research stays of PA at the Institute de Recherche

pour le Développement, MIVEGEC (Montpellier, France). AV was supported by a

grant from IRD (BEST) and ML by a doctoral fellowship from University of

Montpellier and a post-doctoral grant from Labex CeMeb. This study was financially

supported by IRD, CNRS, ECOS-SUD (A16B02) and Malacological Society of

London.

**737 Conflict of interest disclosure**

738 The authors of this preprint declare that they have no financial conflict of interest with

739 the content of this article. Philippe Jarne is one of the PCI Evolutionary Biology

740 recommenders.

## References

**References**

Agnarsson I., Zambrana-Torrelio C.M., Flores-Saldana N.P., May-Collado L.J. 2011. A

time-calibrated species-level phylogeny of bats (Chiroptera, Mammalia). PLoS Curr.

3:RRN1212.

Albrecht C., Wolff C., Glöer P., Wilke T. 2008. Concurrent evolution of ancient sister

lakes and sister species: the freshwater gastropod genus *Radix* in lakes Ohrid and

Prespa. Hydrobiologia. 615:157–167.

Alda P., Lounnas M., Vázquez A.A., Ayaqui R., Calvopiña M., Celi-Erazo M., Dillon

R.T., Jarne P., Loker E.S., Muñiz Pareja F.C., Muzzio-Aroca J., Nárvaez A.O., Noya

O., Robles L.M., Rodríguez-Hidalgo R., Uribe N., David P., Pointier J.-P., Hurtrez-

Boussès S. 2018. A new multiplex PCR assay to distinguish among three cryptic *Galba*

species, intermediate hosts of *Fasciola hepatica*. Veterinary Parasitology. 251:101–105.

Almeyda-Artigas R.J., Bargues M.D., Mas-Coma S. 2000. ITS-2 rDNA sequencing of

Gnathostoma species (Nematoda) and elucidation of the species causing human

gnathostomiasis in the Americas. The Journal of parasitology. 86:537–544.

Artigas P., Bargues M.D., Mera y Sierra R.L., Agramunt V.H., Mas-Coma S. 2011.

Characterisation of fascioliasis lymnaeid intermediate hosts from Chile by DNA

sequencing, with emphasis on *Lymnaea viator* and *Galba truncatula*. Acta Tropica.

120:245–257.

Baker F.C. 1911. The Lymnæidæ of North and Middle America, recent and fossil.

Chicago: The Academy.

Bargues M., Artigas P., Khoubbane M., Ortiz P., Naquira C., Mas-Coma S. 2012.

Molecular characterisation of *Galba truncatula*, *Lymnaea neotropica* and *L.*

*schirazensis* from Cajamarca, Peru and their potential role in transmission of human and

animal fascioliasis. Parasites & Vectors. 5:174.

Bargues M.D., Artigas P., Khoubbane M., Flores R., Glöer P., Rojas-García R., Ashrafi

K., Falkner G., Mas-Coma S. 2011a. *Lymnaea schirazensis*, an overlooked snail

distorting fascioliasis data: genotype, phenotype, ecology, worldwide spread,

susceptibility, applicability. PLoS ONE. 6:e24567.

Bargues M.D., Artigas P., Khoubbane M., Mas-Coma S. 2011b. DNA sequence

characterisation and phylogeography of *Lymnaea cousini* and related species, vectors of

fascioliasis in northern Andean countries, with description of *L. meridensis* n. sp.

(Gastropoda: Lymnaeidae). Parasites & Vectors. 4:132.

Bargues M.D., Artigas P., Mera y Sierra R., Pointier J.-P., Mas-Coma S. 2007.

Characterisation of *Lymnaea cubensis*, *L. viatrix* and *L. neotropica* n. sp., the main

vectors of *Fasciola hepatica* in Latin America, by analysis of their ribosomal and

mitochondrial DNA. Annals of Tropical Medicine & Parasitology. 101:621–641.

Bargues M.D., González C.L., Artigas P., Mas-Coma S. 2011c. A new baseline for

fascioliasis in Venezuela: lymnaeid vectors ascertained by DNA sequencing and

analysis of their relationships with human and animal infection. Parasites & Vectors.

4:1–18.

Bargues M.D., Vigo M., Horak P., Dvorak J., Patzner R.A., Pointier J.P., Jackiewicz

M., Meier-Brook C., Mas-Coma S. 2001. European Lymnaeidae (Mollusca:

Gastropoda), intermediate hosts of trematodiases, based on nuclear ribosomal DNA

ITS-2 sequences. Infection, Genetics and Evolution. 1:85–107.

Bespalaya Y.V., Bolotov I.N., Aksenova O.V., Gofarov M.Yu., Kondakov A.V.,

Vikhrev I.V., Vinarski M.V. 2018. DNA barcoding reveals invasion of two cryptic

*Sinanodonta* mussel species (Bivalvia: Unionidae) into the largest Siberian river.

Limnologica. 69:94–102.

Bickford D., Lohman D.J., Sodhi N.S., Ng P.K.L., Meier R., Winker K., Ingram K.K.,

791 Das I. 2007. Cryptic species as a window on diversity and conservation. Trends in

792 Ecology & Evolution. 22:148–155.

793 Bolotov I.N., Vikhrev I.V., Bespalaya Y.V., Gofarov M.Y., Kondakov A.V., Konopleva

794 E.S., Bolotov N.N., Lyubas A.A. 2016. Multi-locus fossil-calibrated phylogeny,

795 biogeography and a subgeneric revision of the Margaritiferidae (Mollusca: Bivalvia:

796 Unionoida). Molecular Phylogenetics and Evolution. 103:104–121.

797 Bouchet P., Strong E. 2010. Historical name-bearing types in marine molluscs: an

798 impediment to biodiversity studies? In: Polaszek A., editor. Systema Naturae 250 - The

799 Linnaean Ark. CRC Press. p. 63–74.

800 Bouckaert R., Heled J. 2014. DensiTree 2: Seeing trees through the forest.

801 https://doi.org/10.1101/012401.

802 Bouckaert R., Heled J., Kühnert D., Vaughan T., Wu C.-H., Xie D., Suchard M.A.,

803 Rambaut A., Drummond A.J. 2014. BEAST 2: A software platform for Bayesian

804 evolutionary analysis. PLoS Computational Biology. 10:e1003537.

805 Bouckaert R.R., Drummond A.J. 2017. bModelTest: Bayesian phylogenetic site model

806 averaging and model comparison. BMC Evolutionary Biology. 17.

807 Bourdeau P.E., Butlin R.K., Brönmark C., Edgell T.C., Hoverman J.T., Hollander J.

808 2015. What can aquatic gastropods tell us about phenotypic plasticity? A review and

809 meta-analysis. Heredity. 115:312–321.

810 Brotons L., Thuiller W., Araújo M.B., Hirzel A.H. 2004. Presence-absence versus

811 presence-only modelling methods for predicting bird habitat suitability. Ecography.

812 27:437–448.

813 Brown D.S. 1994. Freshwater snails of Africa and their medical importance. London:

814 Taylor & Francis.

815 Burch J.B. 1982. North American freshwater snails. Transactions of the POETS

Society. 1:217–365.

Burgarella C., Gayral P., Ballenghien M., Bernard A., David P., Jarne P., Correa A.,

Hurtrez-Boussès S., Escobar J., Galtier N., Glémin S. 2015. Molecular evolution of

freshwater snails with contrasting mating systems. Molecular Biology and Evolution.

32:2403–2416.

Castresana J. 2000. Selection of conserved blocks from multiple alignments for their

use in phylogenetic analysis. Molecular Biology and Evolution. 17:540–552.

Chapuis E., Trouve S., Facon B., Degen L., Goudet J. 2007. High quantitative and no

molecular differentiation of a freshwater snail (*Galba truncatula*) between temporary

and permanent water habitats. Molecular Ecology. 16:3484–3496.

Coleman A.W., Vacquier V.D. 2002. Exploring the phylogenetic utility of ITS

sequences for animals: a test case for *Abalone* (*Haliotis*). Journal of Molecular

Evolution. 54:246–257.

Correa A.C., Escobar J.S., Durand P., Renaud F., David P., Jarne P., Pointier J.-P.,

Hurtrez-Boussès S. 2010. Bridging gaps in the molecular phylogeny of the Lymnaeidae

(Gastropoda: Pulmonata), vectors of Fascioliasis. BMC Evolutionary Biology. 10:381.

Correa A.C., Escobar J.S., Noya O., Velásquez L.E., González-Ramírez C., Hurtrez-

Boussès S., Pointier J.-P. 2011. Morphological and molecular characterization of

Neotropic Lymnaeidae (Gastropoda: Lymnaeoidea), vectors of fasciolosis. Infection,

Genetics and Evolution. 11:1978–1988.

Coyne J.A., Orr A.H. 2004. Speciation. Sunderland, Massachusetts: Sinauer Associates,

Inc. Publ.

Dayrat B. 2005. Towards integrative taxonomy: integrative taxonomy. Biological

Journal of the Linnean Society. 85:407–415.

Dayrat B., Conrad M., Balayan S., White T.R., Albrecht C., Golding R., Gomes S.R.,

Harasewych M.G., de Frias Martins A.M. 2011. Phylogenetic relationships and

evolution of pulmonate gastropods (Mollusca): New insights from increased taxon

sampling. Molecular Phylogenetics and Evolution. 59:425–437.

De Queiroz K. 2007. Species concepts and species delimitation. Systematic Biology.

56:879–886.

Dillon R.T., Wethington A.R., Lydeard C. 2011. The evolution of reproductive isolation

in a simultaneous hermaphrodite, the freshwater snail *Physa*. BMC Evolutionary

Biology. 11.

D'Orbigny A. 1835. Synopsis terrestrium et fluviatilium molluscorum, in suo per

Americam meridionalem itinere, ab A. D'Orbigny, collectorum. Mag de Zoologie. 5:1–

44.

Dreyfuss G., Correa A.C., Djuikwo-Teukeng F.F., Novobilský A., Höglund J., Pankrác

J., Kašný M., Vignoles P., Hurtrez-Boussès S., Pointier J.P., Rondelaud D. 2015.

Differences in the compatibility of infection between the liver flukes *Fascioloides*

*magna* and *Fasciola hepatica* in a Colombian population of the snail *Galba* sp. Journal

of Helminthology. 89:720–726.

Dunn A.M., Hatcher M.J. 2015. Parasites and biological invasions: parallels,

interactions, and control. Trends in Parasitology. 31:189–199.

Ebbs E.T., Loker E.S., Brant S.V. 2018. Phylogeography and genetics of the globally

invasive snail *Physa acuta* Draparnaud 1805, and its potential to serve as an

intermediate host to larval digenetic trematodes. BMC Evolutionary Biology. 18.

Escobar J.S., Auld J.R., Correa A.C., Alonso J.M., Bony Y.K., Coutellec M.-A., Koene

J.M., Pointier J.-P., Jarne P., David P. 2011. Patterns of mating-system evolution in

hermaphroditic animals: correlations among selfing rate, inbreeding depression, and the

timing of reproduction: mating-system evolution in hermaphroditic animals. Evolution.

65:1233–1253.

Estoup A., Martin O. 1996. Marqueurs microsatellites: isolement à l'aide de sondes

non-radioactives, caractérisation et mise au point. .

Fang Y.-W., Liu L.-Y., Zhang H.-L., Jiang D.-F., Chu D. 2014. Competitive ability and

fitness differences between two introduced populations of the invasive whitefly *Bemisia*

*tabaci* Q in China. PLoS ONE. 9:e100423.

Felsenstein J. 2004. Inferring phylogenies. Sunderland, Mass: Sinauer Associates.

Fišer C., Robinson C.T., Malard F. 2018. Cryptic species as a window into the

paradigm shift of the species concept. Molecular Ecology. 27:613–635.

Folmer O., Black M., Hoeh W., Lutz R., Vrijenhoek R. 1994. DNA primers for

amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan

invertebrates. Molecular marine biology and biotechnology. 3:294–299.

Fourdrilis S., Mardulyn P., Hardy O.J., Jordaens K., de Frias Martins A.M., Backeljau

T. 2016. Mitochondrial DNA hyperdiversity and its potential causes in the marine

periwinkle *Melarhaphe neritoides* (Mollusca: Gastropoda). PeerJ. 4:e2549.

Gomez S., Fleeger J.W., Rocha-Olivares A., Foltz D. 2004. Four new species of

*Cletocamptus* Schmankewitsch, 1875, closely related to *Cletocamptus deitersi* (Richard,

1897) (Copepoda: Harpacticoida). Journal of Natural History. 38:2669–2732.

Gutiérrez A., Pointier J.-P., Fraga J., Jobet E., Modat S., Pérez R.T., Yong M., Sanchez

J., Loker E.S., Théron A. 2003. *Fasciola hepatica*: identification of molecular markers

for resistant and susceptible *Pseudosuccinea columella* snail hosts. Experimental

Parasitology. 105:211–218.

Heled J., Drummond A.J. 2010. Bayesian inference of species trees from multilocus

data. Molecular Biology and Evolution. 27:570–580.

Hubendick B. 1951. Recent Lymnaeidae, their variation, morphology, taxonomy,

nomenclature and distribution. Kungl Svenska Vetenskapsakademiens Handlingar. 3:1–

223.

Jarić I., Heger T., Castro Monzon F., Jeschke J.M., Kowarik I., McConkey K.R., Pyšek

P., Sagouis A., Essl F. 2019. Crypticity in biological invasions. Trends in Ecology &

Evolution.

Jarne P., Pointier J.-P., David P., Koene J.M. 2010. Basommatophoran gastropods. The

evolution of primary sexual characters in animals. Oxford University Press. p. 173–196.

Johnson P.D., Bogan A.E., Brown K.M., Burkhead N.M., Cordeiro J.R., Garner J.T.,

Hartfield P.D., Lepitzki D.A.W., Mackie G.L., Pip E., Tarpley T.A., Tiemann J.S.,

Whelan N.V., Strong E.E. 2013. Conservation status of freshwater gastropods of

Canada and the United States. Fisheries. 38:247–282.

Jones G. 2017. Algorithmic improvements to species delimitation and phylogeny

estimation under the multispecies coalescent. J. Math. Biol. 74:447–467.

Katoh K., Standley D.M. 2013. MAFFT multiple sequence alignment software version

7: improvements in performance and usability. Molecular Biology and Evolution.

30:772–780.

Katz A.D., Taylor S.J., Davis M.A. 2018. At the confluence of vicariance and dispersal:

Phylogeography of cavernicolous springtails (Collembola: Arrhopalitidae,

Tomoceridae) codistributed across a geologically complex karst landscape in Illinois

and Missouri. Ecology and Evolution. 8:10306–10325.

Kolar C.S., Lodge D.M. 2001. Progress in invasion biology: predicting invaders. Trends

in Ecology & Evolution. 16:199–204.

Krug P.J., Vendetti J.E., Rodriguez A.K., Retana J.N., Hirano Y.M., Trowbridge C.D.

2013. Integrative species delimitation in photosynthetic sea slugs reveals twenty

candidate species in three nominal taxa studied for drug discovery, plastid symbiosis or

biological control. Molecular Phylogenetics and Evolution. 69:1101–1119.

Kutschera V.E., Bidon T., Hailer F., Rodi J.L., Fain S.R., Janke A. 2014. Bears in a

forest of gene trees: phylogenetic inference is complicated by incomplete lineage

sorting and gene flow. Molecular Biology and Evolution. 31:2004–2017.

Leaché A.D., Fujita M.K., Minin V.N., Bouckaert R.R. 2014. Species delimitation

using genome-wide SNP data. Systematic Biology. 63:534–542.

Leigh J.W., Bryant D. 2015. PopART: full-feature software for haplotype network

construction. Methods Ecol Evol. 6:1110–1116.

Lounnas M., Correa A.C., Alda P., David P., Dubois M.-P., Calvopiña M., Caron Y.,

Celi-Erazo M., Dung B.T., Jarne P., Loker E.S., Noya O., Rodríguez-Hidalgo R., Toty

C., Uribe N., Pointier J.-P., Hurtrez-Boussès S. 2018. Population structure and genetic

diversity in the invasive freshwater snail *Galba schirazensis* (Lymnaeidae). Canadian

Journal of Zoology. 96:425–435.

Lounnas M., Vázquez A.A., Alda P., Sartori K., Pointier J.-P., David P., Hurtrez-

Boussès S. 2017. Isolation, characterization and population-genetic analysis of

microsatellite loci in the freshwater snail *Galba cubensis* (Lymnaeidae). Journal of

Molluscan Studies. 83:63–68.

Magallón S.A., Sanderson M.J. 2005. Angiosperm divergence times: the effect of

genes, codon positions, and time constraints. Evol. 59:1653.

Mas-Coma S., Bargues M.D., Valero M. a. 2005. Fascioliasis and other plant-borne

trematode zoonoses. International Journal for Parasitology. 35:1255–1278.

Matos-Maravi P., Wahlberg N., Antonelli A., Penz C. 2019. Species limits in butterflies

(Lepidoptera: Nymphalidae): Reconciling classical taxonomy with the multispecies

coalescent. Systematic Entomology, 44:745–756.

Mavárez J., Pointier J.-P., David P., Delay B., Jarne P. 2002. Genetic differentiation,

dispersal and mating system in the schistosome-transmitting freshwater snail

Biomphalaria glabrata. Heredity. 89:258–265.

Medeiros C., Scholte R.G.C., D'Ávila S., Caldeira R.L., Carvalho O.D.S. 2014. Spatial

distribution of Lymnaeidae (Mollusca, Basommatophora), intermediate host of *Fasciola*

*hepatica* Linnaeus, 1758 (Trematoda, Digenea) in Brazil. Revista do Instituto de

Medicina Tropical de São Paulo. 56:235–252.

Meunier C., Hurtrez-Bousses S., Durand P., Rondelaud D., Renaud F. 2004. Small

effective population sizes in a widespread selfing species, *Lymnaea truncatula*

(Gastropoda: Pulmonata). Molecular Ecology. 13:2535–2543.

Meunier C., Tirard, C., Hurtrez-Bousses S., Durand P., Bargues M.D., Mas-Coma S.,

Pointier J.P., Jourdane J., Renaud F. 2001. Lack of molluscan host diversity and the

transmission of an emerging parasitic disease in Bolivia. Molecular Ecology. 10:1333–

1340.

Miller M.A., Pfeiffer W., Schwartz T. 2012. The CIPRES science gateway: enabling

high-impact science for phylogenetics researchers with limited resources. :1.

Niemiller M.L., Near T.J., Fitzpatrick B.M. 2012. Delimiting species using multilocus

data: diagnosing cryptic diversity in the southern cavefish, *Typhlichthys subterraneus*

(Teleostei: Amblyopsidae): species delimitation in cavefish. Evolution. 66:846–866.

Noël E., Jarne P., Glémin S., MacKenzie A., Segard A., Sarda V., David P. 2017.

Experimental evidence for the negative effects of self-fertilization on the adaptive

potential of populations. Current Biology. 27:237–242.

Novobilský A., Engström A., Sollenberg S., Gustafsson K., Morrison D.A., Höglund J.

2014. Transmission patterns of *Fasciola hepatica* to ruminants in Sweden. Veterinary

Parasitology. 203:276–286.

Ogilvie H.A., Bouckaert R.R., Drummond A.J. 2017. StarBEAST2 brings faster species

tree inference and accurate estimates of substitution rates. Molecular Biology and

Evolution. 34:2101–2114.

Orlando Narváez A., Aroca J.M., Alda P., Macías V., Lounnas M., Hurtrez-Boussès S.,

Noya O., Martini Robles L., Pointier J.-P. 2017. Primer reporte de *Galba cubensis*

(Gastropoda: Lymnaeidae) en el Ecuador, hospedador potencial de *Fasciola hepatica* en

arrozales de la costa ecuatoriana. El Misionero del Agro. 13:36–47.

Paraense W.L. 1982. *Lymnaea rupestris* sp. n. from Southern Brazil (Pulmonata:

Lymnaeidae). Memórias do Instituto Oswaldo Cruz. 77:437–443.

Paraense W.L. 1983. *Lymnaea columella* in northern Brazil. Memórias do Instituto

Oswaldo Cruz. 78:477–482.

Paraense W.L. 1995. *Lymnaea cousini* Jousseaume, 1887, from Ecuador (Gastropoda:

Lymnaeidae). Memórias do Instituto Oswaldo Cruz. 90:605–609.

Pedersen U.B., Stendel M., Midzi N., Mduluza T., Soko W., Stensgaard A.-S.,

Vennervald B.J., Mukaratirwa S., Kristensen T.K. 2014. Modelling climate change

impact on the spatial distribution of fresh water snails hosting trematodes in Zimbabwe.

Parasites & Vectors. 7:536.

Pfeiffer L. 1839. Bericht über die Ergebnisse maine Reise nach Kuba im Winter 1838-

1839. Archiv für Naturgeschichte. 5:346–358.

Pfenninger M., Cordellier M., Streit B. 2006. Comparing the efficacy of morphologic

and DNA-based taxonomy in the freshwater gastropod genus *Radix* (Basommatophora,

Pulmonata). BMC Evolutionary Biology.:14.

Pfenninger M., Schwenk K. 2007. Cryptic animal species are homogeneously

distributed among taxa and biogeographical regions. BMC Evolutionary Biology.

7:121.

Pinceel J., Jordaens K., Backeljau T. 2005. Extreme mtDNA divergences in a terrestrial

slug (Gastropoda, Pulmonata, Arionidae): accelerated evolution, allopatric divergence

and secondary contact: Molecular divergence in a terrestrial slug. Journal of

Evolutionary Biology. 18:1264–1280.

Pointier J., Noya O., Amarista M., Théron A. 2004. *Lymnaea cousini* Jousseaume, 1887

(Gastropoda: Lymnaeidae): first record for Venezuela. Memórias do Instituto Oswaldo

Cruz. 99:567–569.

Pointier J.-P. 2015. Freshwater molluscs of Venezuela and their medical and veterinary

importance. Harxheim: ConchBooks.

Puillandre N., Lambert A., Brouillet S., Achaz G. 2012. ABGD, Automatic Barcode

Gap Discovery for primary species delimitation: ABGD, automatic barcode gap

discovery. Molecular Ecology. 21:1864–1877.

Qian Z., Yang J., Lu Y., He J. 2012. Description of three freshwater species

(Gastropoda) from China. Shell Discoveries. 1:30–31.

Rama Rao S., Liew T.-S., Yow Y.-Y., Ratnayeke S. 2018. Cryptic diversity: Two

morphologically similar species of invasive apple snail in Peninsular Malaysia. PLOS

ONE. 13:e0196582.

Rambaut A., Drummond A.J., Xie D., Baele G., Suchard M.A. 2018. Posterior

summarization in bayesian phylogenetics using Tracer 1.7. Systematic Biology.

Ree R.H., Smith S.A. 2008. Maximum likelihood inference of geographic range

evolution by dispersal, local extinction, and cladogene. Systematic Biology. 57:4–14.

Remigio E. 2002. Molecular phylogenetic relationships in the aquatic snail genus

*Lymnaea*, the intermediate host of the causative agent of fascioliasis: insights from

broader taxon sampling. Parasitology Research. 88:687–696.

Remigio E.A., Blair D. 1997. Molecular systematics of the freshwater snail family

Lymnaeidae (Pulmonata: Basommatophora) utilising mitochondrial ribosomal DNA

sequences. J Mollus Stud. 63:173–185.

Ronquist F., Huelsenbeck J.P. 2003. MrBayes 3: Bayesian phylogenetic inference under mixed models. Bioinformatics. 19:1572–1574.

Russel P.M., Brewer B.J., Klaere S., Bouckaert R.R. 2019. Model selection and parameter inference in phylogenetics using nested sampling. Systematic Biology. 68:219–233.

Sabourin E., Alda P., Vázquez A., Hurtrez-Boussès S., Vittecoq M. 2018. Impact of human activities on fasciolosis transmission. Trends in Parasitology. 34:891–903.

Sales J.B. de L., Shaw P.W., Haimovici M., Markaida U., Cunha D.B., Ready J., Figueiredo-Ready W.M.B., Schneider H., Sampaio I. 2013. New molecular phylogeny of the squids of the family Loliginidae with emphasis on the genus *Doryteuthis* Naef, 1912: Mitochondrial and nuclear sequences indicate the presence of cryptic species in the southern Atlantic Ocean. Molecular Phylogenetics and Evolution. 68:293–299.

Samadi S., Roumégoux A., Bargues M.D., Mas-Coma S., Yong M., Pointier J.-P. 2000. Morphological studies of Lymnaeid snails from the human fasciolosis endemic zone of Bolivia. Journal Molluscan Studies. 66:31–44.

Sanabria R., Mouzet R., Pankrác J., Djuikwo Teukeng F.F., Courtioux B., Novobilský A., Höglund J., Kašný M., Vignoles P., Dreyfuss G., Rondelaud D., Romero J. 2013. *Lymnaea neotropica* and *Lymnaea viatrix*, potential intermediate hosts for *Fascioloides magna*. Journal of Helminthology. 87:494–500.

Standley C.J., Prepelitchi L., Pietrokovsky S.M., Issia L., Stothard J., Wisnivesky-Colli C. 2013. Molecular characterization of cryptic and sympatric lymnaeid species from the *Galba*/*Fossaria* group in Mendoza Province, Northern Patagonia, Argentina. Parasites & Vectors. 6:304.

Stevenson J., Norris D. 2016. Implicating cryptic and novel anophelines as malaria

vectors in Africa. Insects. 8:1.

Stewart J.E., Timmer L.W., Lawrence C.B., Pryor B.M., Peever T.L. 2014. Discord

between morphological and phylogenetic species boundaries: incomplete lineage

sorting and recombination results in fuzzy species boundaries in an asexual fungal

pathogen. BMC Evolutionary Biology. 14:38.

Struck T.H., Feder J.L., Bendiksby M., Birkeland S., Cerca J., Gusarov V.I., Kistenich

S., Larsson K.-H., Liow L.H., Nowak M.D., Stedje B., Bachmann L., Dimitrov D. 2018.

Finding evolutionary processes hidden in cryptic species. Trends in Ecology &

Evolution. 33:153–163.

Suh A., Smeds L., Ellegren H. 2015. The dynamics of incomplete lineage sorting across

the ancient adaptive radiation of neoavian birds. PLOS Biology. 13:e1002224.

Taylor D.W. 2003. Introduction to Physidae (Gastropoda: Hygrophila); biogeography,

classification, morphology. Revista de Biología Tropical. 51:1–287.

Thomaz D., Guiller A., Clarke B. 1996. Extreme divergence of mitochondrial DNA

within species of pulmonate land snails. Proceedings of the Royal Society of London.

Series B: Biological Sciences. 263:363–368.

Václavík T., Meentemeyer R.K. 2009. Invasive species distribution modeling (iSDM):

Are absence data and dispersal constraints needed to predict actual distributions?

Ecological Modelling. 220:3248–3258.

Vázquez A.A., Alda P., Lounnas M., Sabourin E., Alba A., Pointier J.-P., Hurtrez-

Boussès S. 2018. Lymnaeid snails hosts of *Fasciola hepatica* and *Fasciola gigantica*

(Trematoda: Digenea): a worldwide review. CAB Reviews. 13:1–15.

Vázquez A.A., Sánchez J., Pointier J.-P., Théron A., Hurtrez-Boussès S. 2014. *Fasciola*

*hepatica* in Cuba: compatibility of different isolates with two intermediate snail hosts,

*Galba cubensis* and *Pseudosuccinea columella*. Journal of Helminthology. 88:434–440.

Vinarski M.V. 2018. *Galba robusta* sp. nov. from Yemen (Gastropoda: Lymnaeidae).

Zoosystematica Rossica. 27:2–10.

Vinarski M.V., Kantor J.I. 2016. Analytical catalogue of fresh and brackish water

molluscs of Russia and adjacent countries. Moscow: A.N. Severtsov Institute of

Ecology and Evolution of RAS.

Vinarski M.V., Schniebs K., Glöer P., Hundsdoerfer A.K. 2011. The taxonomic status

and phylogenetic relationships of the genus *Aenigmomphiscola* Kruglov and

Starobogatov, 1981 (Gastropoda: Pulmonata: Lymnaeidae). Journal of Natural History.

45:2049–2068.

Weigand A.M., Jochum A., Pfenninger M., Steinke D., Klussmann-Kolb A. 2011. A

new approach to an old conundrum-DNA barcoding sheds new light on phenotypic

plasticity and morphological stasis in microsnails (Gastropoda, Pulmonata,

Carychiidae): DNA barcoding. Molecular Ecology Resources. 11:255–265.

Weiss M., Weigand H., Weigand A.M., Leese F. 2018. Genome-wide single-nucleotide

polymorphism data reveal cryptic species within cryptic freshwater snail species-The

case of the *Ancylus fluviatilis* species complex. Ecology and Evolution. 8:1063–1072.

Wilke T., Schultheiß R., Albrecht C. 2009. As time goes by: a simple fool's guide to

molecular clock approaches in invertebrates. American Malacological Bulletin. 27:25–

45.

Xia X. 2017. DAMBE6: new tools for microbial genomics, phylogenetics, and

molecular evolution. Journal of Heredity. 108:431–437.

Yu Y., Harris A.J., Blair C., He X. 2015. RASP (Reconstruct Ancestral State in

Phylogenies): A tool for historical biogeography. Molecular Phylogenetics and

Evolution. 87:46–49.

Yu Y., Harris A.J., He X. 2010. S-DIVA (Statistical Dispersal-Vicariance Analysis): A

tool for inferring biogeographic histories. Molecular Phylogenetics and Evolution.

56:848–850.

Zulliger D.E., Lessios H.A. 2010. Phylogenetic relationships in the genus *Astropecten*

Gray (Paxillosida: Astropectinidae) on a global scale: molecular evidence for

morphological convergence, species-complexes and possible cryptic speciation.

Zootaxa.:1–19.

1097

1098 **Table 1.** Type localities of *Galba* species and sequences of individuals recovered in those localities. References mentioned are the ones in which

1099 sequences were provided and not in which species were described, except for articles describing *Galba meridensis* and *Galba neotropica* were

1100 the authors also provided sequences (Bargues et al. 2007, 2011b). The type localities for *Galba cubensis* and *Galba viator* were here restricted

1101 because original description of the species did not provide a locality, authors simply stated "Cuba" and "Patagonia", respectively (D'Orbigny

1102 1835; Pfeiffer 1839).

| Species | Country | Locality | Coordinates | ITS1 | ITS2 | 16S | COI | Reference |
|---|---|---|---|---|---|---|---|---|
| *Galba cousini* | Ecuador | Chanchu-Yacu | 00°18'55"S 78°34'02"W | FN598157 | FN598153 | - | FN598161 | Bargues et al. 2011b |
| *Galba cubensis* | Cuba | Vaqueria 21 | 23°01'N 82°32'W | AM412226 | AM412223 | - | AM494009 | Bargues et al. 2007 |
| *Galba humilis* | USA | Owego, New York | 42°06'01"N 76°15'40"W | FN182193, FN182194 | FN182191, FN182192 | - | FN182197, FN182198 | Correa et al. 2011 |
| *Galba meridensis* | Venezuela | Laguna Mucubají (Mérida) | 08°47'52"N 70°49'32"W | FN598159 | FN598154 | - | FN598164 | Bargues et al. 2011b |
| *Galba neotropica* | Peru | Lima, Rio Rimac | 12°02'S 76°56'–77°08'W | AM412228 | AM412225 | - | AM494008 | Bargues et al. 2007 |
| *Galba schirazensis* | Iran | Taleb Abad river, Bandar Anzali, Gilan province | 37°27'46"N 49°37'07"E | JF272603 | JF272601 | JF272605 | JF272607 | Bargues et al. 2011a |

| | | | | | | | | |
|---|---|---|---|---|---|---|---|---|
| *Galba truncatula* | Germany | Thuringia, Erfurt-Bindersleben | ND | - | - | - | EU818799 | Albrecht et al. 2008 |
| *Galba viator* | Argentina | Frias | 40°14' S 64°10' W | JN614428 | HQ283265, JN614465 | - | JN614397, JN614398 | Correa et al. 2011 |

 **Figures captions**

 **Figure 1.** The three-step procedure followed to identify the 1,722 individuals of *Galba*

 species. The number of individuals identified at each step is indicated in the left and the

 species identified are indicated on the right. In Step 1, we photographed the shell and

 dissected three to five adult snails from each of the 166 sites. Fragments of the ITS2 and

 COI genes were sequenced in 146 individuals: *Galba cousini/meridensis* (1), *Galba*

 *cubensis* (41), *Galba schirazensis* (41), *Galba truncatula* (30), *Galba humilis* (34) and

 *Galba viator* (1).

 **Figure 2.** Geographic distribution of *Galba* species in America, based on molecular

 identification. Coordinates are given in Tables S1–S2. The sites Perdriel (Argentina),

 Batallas and Tambillo (Bolivia), Ontario (Canada), San Rafael (Mexico), Canal Salinas

 (Puerto Rico) are not represented since coordinates are missing in the original

 publications.

 **Figure 3.** Time-calibrated phylogenetic hypotheses of the model that best approximate

 species status in *Galba* and species-delimitation methods. The trees showed are the

 most probable topologies based on the multispecies tree model that showed the highest

 Bayes Factor in StarBeast2 (visualized in Densitree), as a function of time (in Mya).

 Greater topological agreement is visualized by a higher density of trees (on the left),

 whereas uncertainty in the height and distribution of nodes are represented by increased

 transparency. The most common topologies are shown in blue, the second most

 common topologies in red and the third in green. The six clusters built with individual

 gene trees are indicated on the extreme right. The number of species recovered varied

 with eight approaches (ABGD with each gene, StarBeast with all genes, STACEY with

 a variable number of species included as prior). For each approach, the colored bars

 represent different species. Bars were striped when the groups included more than one

species. Clusters as appeared in text are shown in roman numbers. Scale bar represents

branch length expressed as number of substitutions per site.

**Figure 4.** Most common topology of the species tree and phenotype of the most recent

common ancestor of *Galba* inferred in StarBeast2. The probability of a cryptic

phenotype in the most recent common ancestor of *Galba* species is 100% according to S

-DEC and BBM and 78% according to S-DIVA. Clusters as appeared in text are shown

in roman numbers. Node values indicate posterior probability and blue bars indicate

95% credibility intervals. The mating system is indicated on the right.

**Supplementary Material**

**Table legends**

**Table S1.** Sampled sites from America in which *Galba* species were found. Individuals

were submitted to the three-step procedures for species identification (see text and Fig.

1). For each site, we provide the country, site name, geographic coordinates, sampling

date, and number of sampled individuals. Note that only a fraction of sampled

individuals was sequenced. For each step (and species), we indicate the number of

individuals considered. NA: not available. * indicate sites that have been resampled at

different dates. Accession names in GenBank (ITS2 and COI) are indicated into

parentheses. Note that in some cases a single sequence was obtained. The last column

show the 16S and ITS1 sequences that have been incorporated to the study in order to

test the species hypothesis with the multispecies coalescent models.

**Table S2.** Sites retrieved from literature and GenBank where *Galba* species were

molecularly identified in America. Both the *Galba* and *Lymnaea* names have been used

in the literature at genus level for the species considered in our study—we used *Galba*

here for this monophyletic group of small lymnaeids. For each site, we report the

country, site, geographical coordinates available sequences of mitochondrial (COI and

16S) and nuclear (ITS1 and ITS2) genes, species identification by specific

microsatellites, bibliographic reference, and the species name used in the reference.

Coordinates from Owego, New York were obtained from GoogleEarth and those from

Correa et al. (2010) from Correa et al. (2011). Some coordinates were corrected in order

to match the specific site: Rio Negro (Argentina) from Correa et al. (2010), Frias

(Argentina) from Correa et al. (2011) and Lounnas et al. (2017a), Estanque Lagunillas

(Venezuela) from Bargues et al. (2011c), Baños del Inca (Peru) from Bargues et al.

(2012), Paysandú (Uruguay) from Lounnas et al. (2017a) and Geffrier (Guadeloupe)

(provided by the authors). The KT461809 sequence was erroneously tagged as an ITS2

sequence, but is, in fact, a COI sequence. Sequences of the individuals molecularly

identified by (Medeiros et al. 2014) are missing in the original publication and were not

uploaded to GenBank. ND, no data available.

**Table S3.** Sites retrieved from literature and GenBank where *Galba* species were

molecularly identified in Europe, Asia, and Africa. Coordinates that were not given in

the original articles or in GenBank were best-guess estimated. The information reported

for each site is as in Table S2. ND, no data available.

**Table S4.** Nested sampling results for the eleven species-delimitation models shown in

Figure S13. The model with the higher Marginal Likelihood estimate is the top-ranked

model. All Bayes factor (BF) calculations are made against the current taxonomy model

(scenario D). Therefore, negative BF values indicate support for the current taxonomy

model, and positive BF values indicate support for the alternative model.

**Table S5.** Species of small, mud-dwelling lymnaeids recognized by Burch (1982) in

North America. The author grouped the 22 species into the genus *Fossaria* with two

subgenera, *Fossaria* (*s.s.*) and *Bakerilymnaea*.

1177

 **Figure captions**

 **Figure S1.** Four evolutionary processes that can lead to cryptic species illustrated by

 their phylogenetic histories. (A) Recent divergence: cryptic species are very closely

 related and only recently diverged from each other and, thus, the rate of morphological

 disparity between them (A1 and A2) is not substantially different from that for non-

 cryptic species. (B) Parallelism: the cryptic species are not very closely related to each

 other and the rate of morphological disparity for non-cryptic species is much greater

 than that for cryptic species. (C) Convergence: the cryptic species are also not closely

 related to each other. Initially, morphological disparity for cryptic species can change in

 a manner similar to that for the non-cryptic species pair. However, at some point,

 morphological disparity decreases for the cryptic species, while continuing to increase

 between non-cryptic taxa. (D) Stasis: the cryptic species are closely related to each

 other or are part of a species complex and diverged a long time ago. In comparison with

 non-cryptic species, the rate of morphological change is substantially reduced. (Figure

 and legend extracted and modified from Struck et al. 2018).

 **Figure S2.** First two planes of a principal component analysis on reproductive traits in

 Lymnaeid species. These planes represent 80.36% of the total variance. The analysis

 was conducted using the following measurements: prostate height, prostate width, penis

 length, preputium length, anterior and posterior widths of the penis sheath, and shell

 length and width. All variables were ln-transformed. This analysis included most of the

 species considered in our work, as well as *Pseudosuccinea columella* (one of the closest

 relative of *Galba* spp.), sampled in Argentina, Colombia, Cuba, France, Guadeloupe,

 Peru and Venezuela. Each point represents an individual (N = 81). (Figure and legend

 extracted and modified from Correa et al. 2011).

**Figure S3.** Geographic distribution of *Galba cubensis*, *Galba schirazensis* and *Galba*

*truncatula* in the European, Asian and African samples retrieved from GenBank.

Coordinates are given in Table S3.

**Figure S4.** Shells and reproductive and urinary systems of the six *Galba* species

studied. *Galba cousini*/*meridensis* is the only species that can be morphologically

identified.

**Figure S5.** Phylogenetic tree of *Galba* species based on Bayesian inference in Beast2 of

the COI gene. All sequences from the current study, as well as the ones retrieved from

GenBank, are included in this tree. Sequence coloration represents species. Arrows

indicate sequences belonging to a type locality (see Table 1 for details). Sequence data

are given in Tables S1, S2, and S3.

**Figure S6.** Phylogenetic tree of *Galba* species based on Bayesian inference in Beast2 of

the 16S gene. All sequences were retrieved from GenBank except for sequences from

individuals from Bosque del Apache (USA). Sequence coloration represents species.

Arrows indicate sequences belonging to a type locality (see Table 1 for details).

Sequence data are given in Tables S1, S2, and S3.

**Figure S7.** Phylogenetic tree of *Galba* species based on Bayesian inference in Beast2 of

the ITS1 gene. All sequences were retrieved from GenBank except for sequences from

individuals from Bosque del Apache (USA). Sequence coloration represents species.

Arrows indicate sequences belonging to a type locality (see Table 1 for details).

Sequence data is given in Tables S1, S2, and S3.

**Figure S8.** Phylogenetic tree of *Galba* species based on Bayesian inference in Beast2 of

the ITS2 gene. All sequences for the current study, as well as the ones retrieved from

GenBank, are included in this tree. Sequence coloration represents species. Arrows

indicate sequences belonging to a type locality (see Table 1 for details). Sequence data

are given in Tables S1, S2, and S3.

**Figure S9.** Haplotype network of *Galba* species based on 16S gene. Circle sizes are

proportional to haplotype frequencies and colors represent species. The number of

mutations separating circles are indicated by dashes. The six clusters detected in the

phylogenetic analysis are represented as grey shapes.

**Figure S10.** Haplotype network of *Galba* species based on COI gene. Circle sizes are

proportional to haplotype frequencies and colors represent species. The number of

mutations separating circles are indicated by dashes. The six clusters detected in the

phylogenetic analysis are represented as grey shapes.

**Figure S11.** Haplotype network of *Galba* species based on ITS1 gene. Circle sizes are

proportional to haplotype frequencies and colors represent species. The number of

mutations separating circles are indicated by dashes. The six clusters detected in the

phylogenetic analysis are represented as grey shapes.

**Figure S12.** Haplotype network of *Galba* species based on ITS2 gene. Circle sizes are

proportional to haplotype frequencies and colors represent species. The number of

mutations separating circles are indicated by dashes. The six clusters detected in the

phylogenetic analysis are represented as grey shapes.

**Figure S13.** Scenarios for species assignments used to run the multispecies tree models

using Multi-Threaded Nested Sampling in StarBeast2. The scenario K is an unreal

scenario that separates the populations of *G. viator* from Argentina and Chile to test

whether splitter models showed higher support than lumper models regardless its

biological sense.

**Step 1: Morphological study**

644 individuals

*Galba cousini*

1 individual

**Step 2: Multiplex PCR**

1,420 individuals

*Galba cubensis*
*Galba schirazensis*
*Galba truncatula*

112 individuals

**Step 3: ITS2 and/or COI sequencing**

120   130
GAT AAAT CT GG TCT T AT T TCC

35 individuals

*Galba humilis*
*Galba viator*

*Galba humilis*  III

*Galba cousini*  IV

*Galba meridensis*  V

*Galba* sp.
Bosque del Apache

*Galba viator*  VI

*Galba neotropica*

*Galba cubensis*

*Galba truncatula*  I

*Galba schirazensis*  II

ABGD in 16S
ABGD in COI
ABGD in ITS1
ABGD in ITS2
StarBeast & Nested Sampling
STACEY no species prior
STACEY 5 species-prior
STACEY 9 species-prior

40    30    20    10    5    0   Mya