1

Scale-eating specialists evolved adaptive feeding kinematics within a microendemic radiation of San Salvador Island pupfishes

Michelle E. St. John[1,2], Christopher H. Martin[1,2*]

[1]Department of Biology, University of North Carolina at Chapel Hill, NC, USA

[2]Department of Integrative Biology and Museum of Vertebrate Zoology, University of California, Berkeley, CA 94720, USA

[*]Corresponding Author: chmartin@berkeley.edu

Short title: Scale-eating specialists evolved adaptive feeding kinematics.

keywords: feeding kinematics, novelty, key innovation, lepidophagy, performance, postzygotic isolation, hybrid kinematics

Abstract: 269 words, Main text plus references/figures: 7,357, 5 Tables, 6 Figures, Supplemental Tables S1-S3, Figures S1-S3, and Supplemental video S1

Abstract

Adaptation to novel ecological niches often includes shifts in behaviors, such as new foraging

preferences or changes in kinematics. Investigating prey capture kinematics is an excellent way

to understand behavioral mechanisms underlying the origins of novel trophic specialization, in

which organisms begin to exploit novel resources. We investigated the contribution of

kinematics to the origins of a novel ecological niche for scale-eating within a microendemic

adaptive radiation of pupfishes on San Salvador Island, Bahamas. We compared the feeding

kinematics of scale-eating, generalist, snail-eating, and F1 hybrid pupfishes while they consumed

scales and shrimp in the lab and compared them to scale-eating kinematics observed in situ in the

wild. We then connected variation in feeding kinematics to scale-biting performance by

measuring the area removed per strike from standardized gelatin cubes. We found that scale-

eating pupfish exhibited divergent feeding kinematics compared to all other groups and that

these differences were consistent across food items. The peak gapes of scale-eaters were twice as

large as all other groups, but their gape angles were simultaneously 32% smaller, in both

laboratory and in situ wild observations. We also show that this kinematic combination of large

peak gape and small gape angle resides on a performance optimum for scale-biting. Finally, F1

hybrid kinematics and performance were not additive, and were instead closer to the generalist

pupfish. This suggests that impaired hybrid performance in the scale-eating niche may contribute

to extrinsic postzygotic reproductive isolation between species. Ultimately, our results suggest

that shifts in kinematics (i.e. peak gape and gape angle) are an adaptation to the novel niche of

scale-eating and contribute to reproductive isolation between species.

Introduction

Determining how organisms use resources for the first time and occupy novel niches is an outstanding question in evolutionary ecology. Many changes accompany adaptation to a novel niche, and previous studies have identified that shifts in behaviors (Bowman and Billeb 1965; Tebbich et al. 2010; Curry and Anderson 2012), morphologies (Ferry-Graham et al. 2001; Ferry-Graham 2002; Hata et al. 2011; Davis et al. 2018), physiologies (Arias-Rodriguez et al. 2011; Tobler et al. 2015, 2018), and kinematics (Janovetz 2005; Patek et al. 2006; Cullen et al. 2013; McGee et al. 2013) can all facilitate this transition.

Shifts in kinematic traits—particularly those which affect prey capture and feeding—are especially promising, because they can provide biomechanical insights into the origins of novel trophic niches. For example, some species of trap-jaw ants have evolved specialized latch, spring, and trigger structures in their jaws which allow for rapid, power-amplified bites used to capture prey items unavailable to other ant species (Patek et al. 2006; Larabee and Suarez 2014); power-amplified jaws in Syngnathiform fishes have evolved two different latch mechanisms for specialization on evasive prey items (Longo et al. 2018); and the Pacific leaping blenny (*Alticus arnoldorum)* uniquely uses axial tail twisting to improve propulsion and stability for greater jumping performance and terrestrial prey capture (Hsieh 2010).

Differences in prey capture kinematics between species may also contribute to postzygotic extrinsic reproductive isolation by reducing hybrid foraging performance (Higham et al. 2016b), which may lead to speciation (Henning et al. 2017; Matthews and Albertson 2017). For example, McGee et al. (2015) measured prey capture kinematics and performance in two sunfish species (Centrarchidae) and their naturally occurring hybrids. Hybrid sunfish had an intermediate gape compared to parental types and initiated strikes from an intermediate distance.

However, the force exerted on prey items and the maximum escape force needed to evade

attacks, estimated from the suction-induced flow field model (Holzman et al. 2012), was less

than the intermediate performance expected. Hybrid Lake Victoria cichlids (produced by

crossing *Haplochromis chilotes* (thick-lipped) and *Pundamilia nyererei* (thin-lipped) species)

also exhibited lower foraging performance compared to parental species, most likely due to

antagonistic pleiotropy and genetic correlations between head and lip morphology (Henning et

al. 2017). Despite these findings, few studies investigate how hybrid kinematics affects the

evolution of novelty or explicitly connect kinematics to performance consequences.

Investigating the kinematics of scale-eating is an excellent system for connecting a

mechanistic understanding of feeding kinematics with adaptation to a novel trophic niche. Scale-

eating (lepidophagy) is a novel behavioral trophic niche that is extremely rare which has

independently evolved only 19 times in approximately 100 fish species out of over 35,000

(Sazima 1983; Martin and Wainwright 2013a; Kolmann et al. 2018). However, not much is

known about its evolutionary origins or its kinematics. Current hypotheses for the origins of

scale-eating vary, but they all take a strict behavior-first approach (Greenwood 1965; Sazima

1983; St. John et al. 2018). It is therefore plausible that kinematic variation may have also

contributed to the origins of scale-eating.

Currently, only a few studies have investigated the feeding kinematics and performance

of scale-eating fishes. Janovetz (2005) measured feeding kinematics of the wimple piranha

(*Catoprion mento*) while consuming: 1) free floating scales, 2) whole fish, and 3) scales off the

sides of fish, and found that scale-eating kinematics were divergent from those used in either

suction-feeding or biting. Interestingly, scale-eating attacks produced gape angles that ranged

from 30-100% larger than those produced from consuming free-floating scales or whole fish

respectively— suggesting that a larger gape is necessary for scale-eating. Furthermore, this

variation in gape angle across food items was documented within individuals indicating that

scale-eating kinematics may be plastic (Janovetz 2005). The feeding kinematics of the

Lake Tanganyikan scale-eating cichlid, *Perissodus microlepis,* have also been examined

(Takeuchi et al. 2012; Takeuchi and Oda 2017); however, these studies primarily focused on

how kinematics interacted with *P. microlepis*' antisymmetric mouth morphology and not on

scale-eating kinematics *per se*. Nonetheless, these studies found a significant interaction between

kinematic traits, behavior, and morphology as these fish were able to perform more successful

scale-eating strikes using their dominant side (Takeuchi et al. 2012; Takeuchi and Oda 2017). A

similar oral jaw antisymmetry and behavioral laterality was documented in a scale-eating

characiform (*Exodon paradoxus;* Hata et al. 2011). While these studies provide valuable insights

into scale-eating kinematics and performance, the lack of kinematic measurements in closely

related non-scale-eating species or hybrids has so far limited further investigations of the origins

of scale-eating.

There are four additional comparisons and measurements that would help shed light on

the relationship between kinematic traits and occupation of a novel niche when investigating

scale-eating. First, comparisons of scale-eating kinematics across scale-eating and closely related

non-scale-eating outgroup species is necessary for investigating the origins of novelty. Without

the comparative method it is impossible to determine which kinematic variables are unique or

important for scale-eating. Second, very few kinematic studies investigate hybrid kinematics.

Understanding hybrid kinematics, especially in the context of novelty, is informative because 1)

impaired performance in hybrids is a form of extrinsic postzygotic isolation between species

(McGee et al. 2015; Higham et al. 2016b) and 2) it can allow the decoupling of morphology,

behavior, and kinematics making it easier to identify causative traits underlying performance

(Holzman and Hulsey 2017). Third, very few studies investigate kinematics in the wild (but see:

Keren et al. 2018; Whitford et al. 2019). While this is undoubtedly due to the difficulty of

collecting these data in the field, it does introduce a level of uncertainty regarding whether

kinematics observed in the lab are relevant to performance demands in nature (Shapiro et al.

2011; Stevens et al. 2011). Finally, few studies connect observed variation in kinematics to

variation in whole organism feeding performance (but see: Svanbäck et al. 2003; Takeuchi et al.

2012; Whitford et al. 2019). Making this connection is important because it can identify

kinematic traits associated with performance tasks relevant to evolutionary fitness rather than

simply describing phenotypic variation in kinematic traits, most of which may not be relevant to

performance or fitness (Hu et al. 2017).

The scale-eating pupfish (*Cyprinodon desquamator*) is an excellent organism to

investigate the interaction of kinematics and novelty for several reasons. First, the scale-eating

pupfish evolved within a recent sympatric radiation of pupfishes on San Salvador Island,

Bahamas. This radiation is endemic to a few hypersaline lakes on the island (Martin and

Wainwright 2013a), which were most likely dry during the last glacial maximum 10-15 kya

(Hagey and Mylroie 1995). Second, the radiation provides closely related sister taxa for

kinematic comparison. The radiation contains three species: 1) the scale-eating pupfish, 2) a

generalist pupfish (*C. variegatus*), and 3) a snail-eating pupfish (*C. brontotheroides*).

Phylogenetic evidence suggests that scale-eating pupfishes form a clade across all lakes where

they are found on San Salvador and that this clade is sister to a clade containing generalists and

snail-eaters (Martin and Feinstein 2014; Lencer et al. 2017). Generalists and snail-eaters,

however, cluster together within lakes suggesting either extensive gene flow between species or

multiple origins of snail-eating (Martin and Feinstein 2014; Martin 2016; Richards and Martin

2017). Third, all three pupfish species can be crossed in the lab to measure the kinematics and

performance of hybrid phenotypes.

We investigated the interaction between kinematics and novelty in San Salvador

generalist, snail-eating, and scale-eating pupfishes, and their F1 hybrids while performing high-

speed strikes on three different food items (scales, frozen mysis shrimp, and gelatin cubes). We

asked: 1) if scale-eating pupfish varied in their feeding kinematics compared to other groups, 2)

if scale-eating strikes differed from strikes on frozen mysis shrimp, 3) if scale-eating strikes in

the lab differed from scale-eating strikes in the wild, 4) whether variation in kinematics was

associated with bite performance, and 5) if F1 hybrid feeding kinematics differed from parental

species. Ultimately, we found that feeding kinematics of scale-eating pupfish resulted in bite

sizes 50% larger than all other species, suggesting that scale-eater kinematics are a recent

adaptation for scale-eating.

Methods

Collection and Husbandry

We used seine nets to collect generalist, snail-eating, and scale-eating pupfishes from Crescent

Pond, Little Lake, and Osprey Lake on San Salvador Island, Bahamas in July, 2017 and March,

2018. Wild-caught fish were maintained in 37-75L mixed-sex stock tanks at a salinity of 5-10

ppt and temperature of 23-27°C. While in stock tanks, fish were fed a diet of bloodworms, mysis

shrimp, and commercial pellet foods daily. In the lab, we crossed generalist and scale-eating

pupfishes from both Little Lake and Crescent Pond to produce F1 hybrid offspring. Prior to

filming, pupfish were isolated in a 2L tank to maintain individual IDs throughout the study.

Laboratory feeding kinematics

We recorded pupfishes feeding on three different food items: frozen mysis shrimp, scales, and

standardized gelatin cubes (dimensions: 1.5 cm x5cm X 1.5 cm x 1.5 cm cube; Repashy

Superfoods, Community Plus Omnivore Gel Premix; prepared following manufacturer's

instructions). In the lab, fish freely consumed mysis shrimp (Hikari, Inc.), but we had to train all

species to feed on scales from the sides of euthanized zebrafish (*Danio rerio*; stored frozen) and

to feed from gelatin cubes (stored at 4℃). For training, we isolated each fish in a 2 liter plastic

tank and presented a given food item (either euthanized zebrafish or gelatin cube) daily. If a

pupfish began feeding on the item, it was left in the tank until the pupfish stopped feeding. If a

pupfish did not begin feeding within one minute, the food item was removed from the tank. Any

pupfish that did not feed received a supplemental feeding of commercial pellet food (New Life

Spectrum Thera-A, medium sinking pellets). If an individual did not feed on a training item for

more than two days, we reduced supplemental feedings to once every two days to ensure that the

fish was sufficiently motivated. Once pupfish reliably began feeding on either scales or gelatin

cubes, we proceeded to film their feeding behaviors according to the filming protocol below.

Fish were never trained on more than one item at a time, and we instead ensured that all filming

was completed for a single food item before proceeding to train for the next item.

For all three food items, we followed the same filming protocol and used the same

equipment: 1) a Sony Cyber-shot DSC-RX10 III (480fps) or Sony Cyber-shot DSC-RX100 IV

20.1 MP (480fps) for high-speed video of foraging strikes and 2) a dimmable bi-color 480

LED light (Neewer) as an additional light source. Prior to filming, each pupfish—still in their 2

L tank—was placed in front of 0.5 cm grid paper. We placed our camera as close as possible to

the tank while still keeping both the pupfish and food item in frame. We placed the additional

LED light source approximately 0.3 m from the filming tank and allowed pupfish to acclimate to

the lighting until they moved around their tank freely after approximately 5 minutes. Once a

pupfish was acclimated we presented the pupfish with a given food item. For scale-eating we

used forceps to hold a euthanized zebrafish horizontally in the water column and perpendicular

to the front of an individual. For mysis shrimp and gelatin cubes, we dropped the food item a few

inches in front of an individual. All videos were taken from a lateral perspective. Once filming

for one food item was completed, the process was repeated until we filmed each individual

consuming all three food items.

Scale-eating kinematics in the wild

We also filmed seven usable lateral scale-eating strikes by scale-eaters in the wild. It was not

possible to film generalist or snail-eater scale-eating strikes, since these species do not naturally

strike scales at a detectable frequency in nature (although scales have been found in the guts of

both species at very low frequencies; Martin and Wainwright 2013). We used a Chronos camera

(Kron Technologies, model 1.4, 16 GB memory, Color image sensor) with an f1.4 zoom lens in a

custom underwater housing (Salty Surf, Inc. Krontech Chronos 1.4 housing with M80 flat port)

to record natural high-speed videos of scale-eating in Crescent Pond, San Salvador Island (see

Supplemental Video S1). We recorded videos while snorkeling along the shoreline in 0.3 – 0.5 m

depth in March, 2018. We set the frame rate to the minimum of 1080 fps on the camera,

however, our manual calibration of stopwatch videos filmed in the field indicated that the

effective frame rate was only 270 fps.

Standardized reference objects for calibration (i.e. grid paper) were not present in the

wild scale-eating videos and filming a reference object in a second video was a substantial

obstacle given the constantly shifting distance between the camera and subject and the necessity

of removing the camera from the underwater housing to store and reset after each video. Instead,

we measured the diameter of 20 mermaid's cup macroalgae (*Acetabularia acetabulum*) from a

photograph containing a ruler and estimated the mean cup diameter of adult cups (mean $\pm$ SE:

3.31 ± 2.86 mm). This macroalgae was present in all videos. We used this mean cup diameter to

convert wild measurements in pixels to meters.

Kinematic analyses

We analyzed videos by converting them to image stacks using Adobe Media Encoder (Version

13.0.2). Image sequences were then imported into image processing software (FIJI) for analysis

(Schindelin et al. 2012). To quantify feeding performance, we measured 8 kinematic trait

metrics: peak jaw protrusion, time to peak jaw protrusion, peak gape, time to peak gape, gape

angle at time of peak gape, starting distance from food item, time to impact, and ram speed

(Table 1).

All time and distance metrics were measured from the start of a strike defined as when

the lower jaw was opened to ~20% of peak gape, identified by visual inspection of each frame

(following: Oufiero et al. 2012; Staab et al. 2012). In addition to our kinematic metrics, we also

measured body length and lower jaw length (Table S1) using images from the video. We

calibrated each video using the background grid for laboratory strikes and macroalgae for wild

strikes. Kinematic profiles of a single representative scale-eating strike from each species are

shown in Figure 1.

Figure 1.

Figure 1. Kinematic profiles of pupfish species for scale-eating strikes observed in the lab and in situ in the wild. We produced kinematic profiles for a single representative strike for each group, and reported the proportional change in gape distance, jaw protrusion distance, and gape angle over the course of a single scale-eating strike. Time zero for each group represents the start of an attack (i.e. 20% of peak gape) and the dotted line indicates the time of impact with the prey. Point positions are relative to the highest value metric across groups (e.g. a value of 1 indicates the largest value of a given metric). Fill colors show raw values for each point.

Measuring bite performance

In order to connect variation in feeding kinematics to variation in bite size we recorded high-

speed strikes on gelatin meal replacement for fish in the shape of a 1.5 x 1.5 x 1.5 cm cube. We

filmed a feeding strike on a single cube and immediately removed the cube from the tank. The

gel cube retains its shape in water and therefore allowed us to precisely photograph and measure

the area removed by each bite. We used an Olympus Tough TG-5 camera to take photos of each

lateral surface of the cube –ensuring that we had photographed the entire bite—and measured the

total area removed from the cube (Figure 4).

Statistical analyses

Comparing strike kinematics

We collected and analyzed 101 feeding strikes from 31 individuals striking both shrimp and

scales. (7 generalists; 7 snail-eaters; 9 scale-eaters; 8 F1 hybrids). We used linear mixed models

in the lme4 package in R (Bates et al. 2014) and RStudio (R Core Team 2018) to determine if

any of our kinematic metrics varied between species or food item. In each model we included: 1)

the kinematic metric as the response variable, 2) species designation, food item, and their

interaction as fixed effects, 3) individual fish IDs and population as random effects, and 3) log

body size as a covariate (Table 3). Similarly, we used linear mixed models to determine if the

feeding strikes of scale-eaters in the wild differed from those in the lab. We used each kinematic

metric as the response variable, and modeled: 1) environment (e.g. lab or wild) as a fixed effect,

2) individual ID nested within environment as a random effect, and 3) log body length as a

covariate (Table 4).

We also performed a linear discriminant analysis (LDA) on the combined shrimp and

scales kinematic data to reduce dimensionality and identify which kinematic metrics contributed

most to differences between species (Table 2, Figure 2A). We used a MANOVA and Wilks' λ to

assess the significance of the LDA. Our MANOVA included 1) all 8 kinematic metrics as

response variables, 2) species designation as a predictor variable, and 3) individual ID as a

random effect.

Although we compared kinematic data across multiple species, very few genetic variants

are fixed between species (<1,000 SNPs out of 12 million) and generalists and molluscivores

cluster by lake rather than by species (McGirr and Martin 2017; Richards and Martin 2017).

Thus, is it appropriate to analyze species differences at these recent timescales as population-

scale data using mixed model analyses of independent populations (e.g. Hatfield and Schluter

1999; McGee et al. 2013), rather than phylogenetic comparative methods.

Determining how kinematic variables affect bite performance

We used a linear mixed model to investigate if the area removed from a cube (mm^2) was

associated with any of the kinematic variables measured during strikes by all three species and

F1 hybrids. We first performed principle components analysis (PCA) of our 8 kinematic

variables (Table S3) to obtain a set of orthogonal axes and subsequently used all eight PCs as

fixed effects in our mixed model. We included 1) area removed from a cube (mm^2) as the

response variable, 2) PC1-8 as fixed effects, 3) fish ID and species as random effects, and 4) log

body length as a covariate (Table 5).

We also used generalized additive models (GAMs) using the mgcv package (Wood 2011)

in R to further investigate how peak gape, peak protrusion, and gape angle affected bite size. We

specifically chose these kinematic variables because our previous linear mixed model indicated

that PC5 was a significant predictor variable of bite size which included major loadings of peak

gape and peak protrusion (Table 5). Our first GAM included 1) area removed from a cube (mm^2)

as the response variable and 2) a spline modeling the interaction between peak gape and peak

protrusion as predictor. Since peak gape and peak protrusion are on the same relative scale (mm)

we used an isotropic thin-plate smoothing spline for this model. In our second model, we

included 1) area removed from a cube (mm^2) as the response variable and 2) peak gape, gape

angle, and peak protrusion as predictor variables, and used univariate smoothing splines for each

kinematic variable within the GAM.

Finally, we predicted the area removed per bite for each fish from their peak gape and

gape angle kinematic measurements using a machine-learning algorithm from the caret package

using a spline-based method (Kuhn 2008). Thus, for all scale-eating and shrimp-feeding strikes

resulting in unknown bite sizes (unlike the gelatin cube strikes), we were able to build and tune a

GAM model connecting our two kinematic variables of interest to the area removed from gelatin

cubes (observed bite performance) in order to predict bite performance for all strikes in our

dataset.

We built the model using 1) area removed from a cube (mm^2) as the response variable

and 2) peak gape and gape angle as predictor variables. We trained the model using all strikes

observed on gelatin cubes (31 strikes across all three species and F1 hybrids) and 10-fold cross-

validations with three repeats as the resampling scheme. We tested the accuracy of this model by

comparing fitted values from the model to observed values from the data set and found that our

model was able to predict 68% of the variance in the dataset (df=1, F=63.84, P=8.2x10^{-9},

R^2=0.68). We then used this model to predict the bite area removed for each scale-eating and

shrimp-eating strike based on the kinematic measurements alone. We used bootstrap resampling

(20,000 iterations) to calculate mean bite size (predicted area removed) and 95% confidence

intervals for each species.

Determining if hybrid kinematics match additive predictions

For peak gape, gape angle, and bite size, we calculated the predicted values of F1 scale-eater by

generalist hybrids under the hypothesis that these kinematic traits would be additive and

therefore intermediate between generalist and scale-eater values. We used a one sample *t*-test to

test whether the observed values of peak gape, gape angle, and the predicted bite sizes for F1

hybrids deviated from additive predictions.

Results

Scale-eaters exhibited divergent feeding kinematics compared to other pupfishes

Scale-eaters exhibited divergent feeding kinematics, while consuming both shrimp and scales,

compared to other groups (Figure 2A). A MANOVA supported the significance of this

discriminant analysis and found species designation was a significant predictor of kinematics

(Wilks' $\lambda = 0.10$; F = 2.9471; $df = 3$; $P = 0.000394$). Species significantly varied in their peak

gape and gape angles during feeding strikes—regardless of the food item— in a linear mixed

model controlling for individual ID and body length (Table 3). This pattern was driven by scale-

eaters who had peak gapes that were twice as large as other species, but also had gape angles that

were one third smaller than other species (Figure 2B-C).

Pupfish ram speed was 48% faster when consuming shrimp

Ram speed was the only kinematic variable that varied between food items, increasing by ~50%

for shrimp strikes (0.098 m/s, CI: 0.082, 0.12) relative to scale-eating strikes (0.066 m/s, CI:

0.061, 0.070; Table 3; Figure S2).

347   **Figure 2.**

Figure 2. Divergent feeding kinematics in scale-eaters. A) Biplot of discriminant axes 1 (LD1) and 2 (LD2) describing overall kinematic differences among pupfish species (generalists, snail-eaters, scale-eaters, or F1 hybrids). Ellipses represent 95% CIs. B) Mean peak gape (mm) for each species with ± 95% CIs calculated via bootstrapping (20,000 iterations). C) Mean gape angle at peak gape (mm) for each species with ± 95% CIs calculated via bootstrapping (20,000 iterations).

Scale-eating in the wild differed from scale-eating in the lab

Scale-eating attacks in the wild started from over twice as far away as those observed in the lab

(*P* = 0.033; Table 4; Figure 3A) and ram speed was almost three times faster in the wild than in

the lab (*P* = 0.00099; Table 4; Figure 1, 3B). However, the range of strike kinematics for both

these variables overlapped between the lab and field, indicating substantial variation in scale-

eating kinematics in both environments.

Figure 3.

398 **Figure 3. Wild scale-eating strikes are more than twice as fast and start three**
399 **times farther away than lab strikes.** a) Mean starting distance from prey defined as
400 20% of peak gape and b) mean ram speed (± 95% CIs; bootstrapping 20,000 iterations)
401 for scale-eaters consuming scales in the lab and observed in situ in Crescent Pond, San
402 Salvador Island, Bahamas.

Variation in feeding kinematics affected bite size performance

Two out of eight principal component axes of strike kinematics were significantly associated

with the surface area removed from a gelatin cube per strike (Table 5). However, the significance

of PC2 was entirely driven by a single outlier, and when removed, this axis was no longer

significantly associated with bite size ($\chi^2 = 0.71$, *df*=1, *P* =0.40; Figure S3). PC5 remained

significantly associated with bite size regardless of whether this outlier was included in the

dataset. PC5 contained two major loadings (|loadings|>.4): peak gape and peak protrusion (Table

S3), which loaded in opposite directions (Table 5, Figure 4).

Figure 4.

Figure 4. Peak gape and peak protrusion affect bite size. Graph illustrating the positive relationship between PC5 and the area removed from a gelatin cube (mm^2) from a single strike. The major loadings of PC5 are peak gape and peak protrusion. Grey dots represent individual fish, the line represents the fitted prediction of a linear model along with a 95% CI ribbon in blue.

Our GAMs also indicated that the interaction between peak gape and peak protrusion, modeled as a thin-plate spline, was significantly associated with bite size (*edf*=6.61, *Ref.df*=29, F= 0.59, *P*= 0.027), and explained 47.6% of the observed deviance in bite size (Figure 5A). Our second GAM, which modeled peak gape, gape angle, and peak protrusion as independent univariate smoothing splines, explained 69% of the deviance in bite size and indicated that increased peak gape (*edf*=4.96, F= 7.88, *P*= 6.22×10^{-5}) and reduced gape angle were significantly associated with increased bite size (*edf*=1, F= 4.72, *P*= 0.04). We also fit a model including gape angle and peak protrusion as linear fixed effects instead of splines; however, this model provided a poorer fit to the data (ΔAIC = 2.13). Taken together, our results suggest that a larger gape and smaller gape angle are associated with larger areas removed per bite, but that jaw protrusion distance has only a small effect (increases explained deviance by 0.9%; change in ΔAIC = 1.39) on bite size (Figure 5B-D).

Our machine learning model also predicted that scale-eater kinematics would result in bite sizes that are 65% larger than the predicted bites of the other species (Figure 6). Interestingly, estimates for F1 hybrid bite sizes were smaller than expected if kinematic metrics and bite performance are additive traits; however, this effect was only marginal (T-test, μ= 340.73, mean= 289.18, *P = 0.059*).

Figure 5.

Figure 5. The interaction of peak gape and peak protrusion may result in a performance optimum for scale-biting. a) Visualization of a GAM model investigating how the thin-plate interaction between peak gape (mm) and peak protrusion (mm) affects bite size (area removed per strike from a gelatin cube). Yellow colors represent larger bites, while blues represent smaller bites. Grey areas represent bite sizes that cannot be predicted from the current data set. b) Visualizations of the smoothing splines for b) peak gape size, c) gape angle, and d) peak protrusion within the GAM model.

Figure 6.

Figure 6. Scale-eaters have larger predicted bite sizes compared to other species.
Predicted bite sizes for all strikes from each species using machine-learning
optimization of GAM models. Grey points represent predicted bite sizes for individuals,
color points represent means, and bars represent ± 95% CIs calculated via
bootstrapping (20,000 iterations).

F1 hybrid kinematics are not additive and more closely resemble generalist kinematics

F1 hybrid feeding kinematics, across both food items, differed from scale-eater kinematics

(TukeyHSD, $P = 4.59 \times 10^{-8}$), but were not significantly different from generalist kinematics

(Tukey's HSD, $P = 0.21$). Mean hybrid peak gape was 58% smaller than scale-eater peak gape

and 24% larger than generalist peak gape (Figure 2B). Similarly, mean hybrid gape angle was

56% larger than scale-eater peak gape angle, but was only 4.25% smaller than the mean

generalist gape angle (Figure 2C). Hybrids failed to match additive predictions of intermediate

kinematics for both peak gape (t-test, $\mu = 3.13$, mean= 2.25 mm, $P = 0.0032$) and gape angle (t-

test, $\mu = 42.43$, mean= 48.37 degrees, *P = 0.044*).

Discussion

Scale-eating pupfish have divergent, but not plastic, feeding kinematics

Scale-eating pupfish exhibited peak gapes during scale-eating strikes that were twice as large

and gape angles that were 32% smaller than other sympatric species within the San Salvador

Island radiation. Similarly, Janovetz (2005) found that the peak gape angle of the scale-eating

piranha *Cataprion mento* was 1.5 times larger during scale-eating strikes than while suction-

feeding on whole fish. Unlike *C. mento*, however, pupfish feeding kinematics remained

divergent between species, but consistent across prey items (Table 3). In fact, the only kinematic

variable that varied between prey items was ram speed (Table 3, Figure S2). However, this may

simply be due to the fact that shrimp were a moving target during feeding trials while scales

were stationary on the side of a euthanized zebrafish. Alternatively, phenotypic plasticity due to

rearing environment could produce a similar pattern where wild caught pupfish display plastic

kinematic traits, while lab reared pupfish do not. However, we find this scenario unlikely as all

fish were acclimated to the laboratory environment for several months before feeding trials

began, and we did not observe any difference in kinematic traits between these two groups.

Is jaw morphology solely responsible for kinematic variation?

The kinematic variables that varied the most between scale-eating and non-scale-eating

pupfishes were peak gape and gape angle—both related to the size of the oral jaws. Previous

work has documented that the oral jaws of scale-eating pupfish are three times larger than their

sister species (Holtmeier 2001; Martin and Wainwright 2013a; Martin 2016) and controlled by

four moderate-effect QTL with all positive effects on jaw size, consistent with directional

selection on this trait (Martin et al. 2017). It may be that increased oral jaw size is sufficient to

create variation in feeding kinematics without an accompanying shift in behavior. Previous

studies have documented how changes in morphology alone can alter feeding kinematics. For

example, kinematic studies have found that the scaling of the lower jaw in bluegill (Wainwright

and Shaw 1999) and body size in largemouth bass (*Micropterus salmoides;* Richard and

Wainwright 1995) both significantly affected prey capture kinematics. Furthermore, Ferry-

Graham et al. (2010) used the pike killifish (*Belonesox belizanus*) to show that simply doubling

the length of the jaws significantly affected key kinematics variables such as peak gape size—

even while keeping gape angle constant. Simply stated, the key innovation necessary for scale-

eating may be an enlarged jaw. If this hypothesis were true, we would expect that peak gape

would increase with jaw size, but that all other kinematics variables would remain constant

across species. Our results reject this hypothesis. Instead, the gape angle of scale-eaters is 32%

smaller than other groups resulting in a reduction in their potential peak gape size. This suggests

that scale-eaters have evolved a smaller gape angle possibly to increase feeding performance

(Figure 5 & 6). Furthermore, this reduced gape angle is consistent across food items and strikes,

indicating that it is not a plastic trait relative to individual foraging context.

Scale-eating performance optimum

Scale-eaters may have reduced their gape angles relative to other species in order to remain on a

performance optimum for scale-eating. Our models of bite performance supported this: peak

gapes larger than approximately 4.5 mm counterintuitively resulted in smaller bite sizes (Figure

5A). A reduced angle in scale-eating pupfish results in a lower jaw that is essentially parallel

with the ventral surface of its body—possibly resulting in greater stability for biting scales while

retaining a large gape. This large gape and parallel jaw alignment may allow scale-eaters to

attack prey from a roughly perpendicular angle (as frequently observed during field

observations) —appearing to wrap their large lower jaw under prey items and subsequently

scrape scales from their sides using their independently protrusible upper jaws (also observed in

a scale-eating characin: Hata et al. 2011). Interestingly, perpendicular angles of attack and large

gapes are associated with scraping in benthic feeding fish (Van Wassenbergh et al. 2008; O'Neill

and Gibb 2013). In fact, one prominent hypothesis for the origins of scale-eating is that it arose

from an algae-scraping ancestor (Sazima et al. 1983). One caveat for this hypothesis, however, is

that our current performance estimates do not include all possible combinations of peak gape and

gape angle. Future work should estimate performance across all combinations of peak gape and

gape angle (e.g. Stayton 2019).

Kinematics in the wild differs from the lab

Scale-eating in the lab is decidedly different from in the wild. In the wild, scale-eaters must

approach a prey pupfish while remaining unnoticed, attack them very quickly, hold on to the side

of the pupfish as it attempts to escape, and finally rip off the desired scales and protein-rich

mucus coat (Supplemental Video S1). In the lab, however, a scale-eating attack on euthanized

immobilized prey involves none of these evasive prey aspects of an attack, which may affect

kinematics. Despite these stark differences, many kinematic traits measured in both the lab and

the wild were similar. Importantly, the two kinematics which varied the most between scale-

eaters and other species in the lab, peak gape and gape angle, were consistent across

environments (Table 4).

However, we did find that wild scale-eating strikes were faster than those in the lab and

began from further away; however, the range of both these kinematics variables still overlapped

with strikes recorded in the lab (Table 4, Figure 3). This may indicate that pupfish can vary their

feeding kinematics based on evasive prey dynamics.

Non-additive F1 hybrid feeding kinematics may contribute to reproductive isolation of

scale-eaters

Given that complex performance traits are most likely highly polygenic and thus may exhibit

additive heritability on average, we expected F1 hybrids to exhibit intermediate kinematics and

performance relative to both parental species. Instead, we found that F1 hybrid kinematics did

not meet the predictions of additivity and that their kinematics and performance more closely

resembled that of the generalist (Table 3; Figure 2 & 6), suggesting that F1 hybrids may have

higher performance in a generalist trophic niche. Current evidence from field fitness experiments

supports this idea. One field experiment in these lakes measured F2 hybrid fitness in the wild and

found high mortality and low growth rates for hybrids most closely resembling the scale-eating

phenotype (Martin and Wainwright 2013b). Furthermore, for the few hybrids resembling scale-

eaters which did survive, only 36% had recently consumed any scales compared to 92% of wild-

caught scale-eaters (Martin and Wainwright 2013a,b). Impaired hybrid performance in the scale-

eating niche may contribute to extrinsic postzygotic isolation between species (McGhee et al.

2007; McGee et al. 2013; Higham et al. 2016). Reproductive isolation may also evolve more

quickly in species that occupy a more distant fitness peak with a larger fitness valley such as the

scale-eating pupfish due to stronger selection against hybrids and reinforced pre-mating isolation

(Martin and Feinstein 2014). Thus impaired hybrid scale-eating performance could also

contribute to increased diversification rates through the mechanism of a wider fitness valley.

Low hybrid performance may also be due to their morphological differences from purebred

scale-eaters and generalists. As mentioned above, it is possible that a shift in morphology – such

as enlarged oral jaws in scale-eaters—may be sufficient to change kinematic profiles alone. F1

hybrid kinematics clearly differed from scale-eater kinematics, but their jaw lengths were also

significantly smaller than the jaws of scale-eaters (Tukey's HSD, $P = 0.018$). Furthermore,

previous work has shown that hybrid pupfish offspring (produced from generalist x scale-eater

crosses) tend to develop along a more similar trajectory to their maternal parent (Holtmeier

2001). This could indicate that hybrid pupfish with scale-eating mothers are more likely to

develop jaws resembling a purebred scale-eater, but may also retain their generalist-like

kinematics. The resulting mismatch between morphology, kinematic traits, and ecological niche

may be driving low hybrid survival in the scale-eating niche and contributing to reproductive

isolation between generalist and scale-eating pupfish species.

Conclusion

In conclusion, this study suggests that shifts in kinematic traits may have preceded or facilitated

the origin of scale-eating in *Cyprinodon* pupfishes. Scale-eating pupfish exhibited peak gapes

that were twice as large as other pupfish species, but simultaneously had gape angles that were

significantly smaller. Surprisingly, we found that this unique combination of scale-eater

kinematics may reside on a performance optima, as large peak gapes and small gape angles result

in larger bite sizes. Impaired F1 hybrid kinematics and performance in the scale-eating niche also

suggests that kinematic traits contribute to reproductive isolation of the scale-eating pupfish and

the evolution of novelty. Future work should investigate if other performance optima exist on the

kinematic landscape and whether F2 hybrid fitness in the wild is due to a mismatch between

morphology and feeding kinematics.

Authors' Contributions

MES and CHM conceptualized the project, MES collected data and performed all analyses, MES

wrote the manuscript, MES and CHM revised drafts, and CHM funded the study.

Data Accessibility

All raw data will be deposited in the Dryad Digital Repository.

Acknowledgements

We thank the University of North Carolina at Chapel Hill, NSF CAREER 1749764, NIH

5R01DE027052-02, and BSF 2016136 for funding to CHM and the UNC Quality Enhancement

Plan for course-based undergraduate research (CURE) funding to CHM. The Bahamas

Environmental Science and Technology Commission and the Ministry of Agriculture kindly

provided permission to export fish and conduct this research. Rochelle Hanna, Velda Knowles,

Thomas Rothfus, and the Gerace Research Centre provided logistical assistance in the field; Kristi

Dixon, Casey Charbonneau, Gabriel Harris, Amelia Ward, Delaney O'Connell, and the

undergraduate students of BIO221L 'Evolution of Extraordinary Adaptations' assisted with high-

speed videography and kinematic analysis; and Roi Holzman provided helpful feedback on the

research. All animal care protocols were approved by the University of California, Berkeley and

the University of North Carolina at Chapel Hill Animal Care and Use Committees.

References

Arias-Rodriguez, L., M. Tobler, M. Palacios, F. J. García de León, D. Bierbach, M. Mateos, I.

Mitrofanov, M. Plath, and L. J. Chapman. 2011. Evolution in extreme environments:

replicated phenotypic differentiation in livebearing fish inhabiting sulfidic springs.

Evolution (N. Y). 65:2213–2228.

Bates, D., M. Mächler, B. Bolker, and S. Walker. 2014. Fitting Linear Mixed-Effects Models

using lme4. J. Stat. Softw. 67:1–48.

Bowman, R. I., and S. L. Billeb. 1965. Blood-eating in a Galápagos finch. Living Bird 4:29–44.

Cullen, J. A., T. Maie, H. L. Schoenfuss, and R. W. Blob. 2013. Evolutionary Novelty versus

Exaptation: Oral Kinematics in Feeding versus Climbing in the Waterfall-Climbing

Hawaiian Goby Sicyopterus stimpsoni. PLoS One 8:e53274. Public Library of Science.

Curry, R. L., and D. J. Anderson. 2012. Interisland Variation in Blood Drinking by Galápagos

Mockingbirds. Auk 104:517–521.

Davis, A. L., M. H. Babb, M. C. Lowe, A. T. Yeh, B. T. Lee, and C. H. Martin. 2018. Testing

Darwin's Hypothesis about the Wonderful Venus Flytrap: Marginal Spikes Form a "Horrid

Prison" for Moderate-Sized Insect Prey. Am. Nat. 193:309–317.

Ferry-Graham, L. A. 2002. Using Functional Morphology to Examine the Ecology and Evolution

of Specialization. Integr. Comp. Biol. 42:265–277.

Ferry-Graham, L. A., L. P. Hernandez, A. C. Gibb, and C. Pace. 2010. Unusual kinematics and

jaw morphology associated with piscivory in the poeciliid, Belonesox belizanus. Zoology

113:140–147.

Ferry-Graham, L. A., P. C. Wainwright, C. Darrin Hulsey, and D. R. Bellwood. 2001. Evolution

and mechanics of long jaws in butterflyfishes (Family Chaetodontidae). J. Morphol.

248:120–143.

Hata, H., M. Yasugi, and M. Hori. 2011. Jaw laterality and related handedness in the hunting

behavior of a scale-eating characin, exodon paradoxus. PLoS One 6.

Hatfield, T., and D. Schluter. 1999. Ecological Speciation in Sticklebacks: Environment-

Dependent Hybrid Fitness. Evolution (N. Y). 53:866. Mayr.

Henning, F., G. Machado-Schiaffino, L. Baumgarten, and A. Meyer. 2017. Genetic dissection of

adaptive form and function in rapidly speciating cichlid fishes. Evolution (N. Y). 71:1297–

1312.

Higham, T. E., S. M. Rogers, R. B. Langerhans, H. A. Jamniczky, G. V Lauder, W. J. Stewart,

C. H. Martin, and D. N. Reznick. 2016. Speciation through the lens of biomechanics:

locomotion, prey capture and reproductive isolation. Proc. R. Soc. B Biol. Sci.

283:20161294.

Holtmeier, C. L. 2001. Heterochrony, maternal effects, and phenotypic variation among

sympatric pupfishes. Evolution 55:330–8.

Holzman, R., D. C. Collar, R. S. Mehta, and P. C. Wainwright. 2012. An integrative modeling

approach to elucidate suction-feeding performance. J. Exp. Biol. 215:1–13. The Company

of Biologists Ltd.

Hu, Y., N. Nelson-Maney, and P. S. L. Anderson. 2017. Common evolutionary trends underlie

the four-bar linkage systems of sunfish and mantis shrimp. Evolution (N. Y). 71:1397–

1405.

Janovetz, J. 2005. Functional morphology of feeding in the scale-eating specialist *Catoprion*

mento. J. Exp. Biol. 208:4757–68.

Keren, T., M. Kiflawi, C. Martin, V. China, O. Mann, and R. Holzman. 2018. A complex

performance landscape for suction-feeding reveals constraints and adaptations in a

population of reef damselfish. bioRxiv 239418.

Kuhn, M. 2008. Building predictive models in R using the caret package. J. Stat. Softw. 1–26.

Larabee, F. J., and A. V. Suarez. 2014. the Evolution and Functional Morphology of Trap-Jaw

Ants. Myrmecological News 25–36.

Lencer, E. S., W. C. Warren, R. Harrison, and A. R. McCune. 2017. The Cyprinodon variegatus

genome reveals gene expression changes underlying differences in skull morphology among

closely related species. BMC Genomics 18:424.

Longo, S. J., T. Goodearly, and P. C. Wainwright. 2018. Extremely fast feeding strikes are

powered by elastic recoil in a seahorse relative, the snipefish, Macroramphosus scolopax. P.

20181078 *in* Proceedings of the Royal Society B: Biological Sciences. The Royal Society.

Martin, C. H. 2016. The cryptic origins of evolutionary novelty: 1000-fold faster trophic

diversification rates without increased ecological opportunity or hybrid swarm. Evolution

(N. Y). 70:2504–2519.

Martin, C. H., and L. C. Feinstein. 2014. Novel trophic niches drive variable progress towards

ecological speciation within an adaptive radiation of pupfishes. Mol. Ecol. 23:1846–1862.

Martin, C. H., and P. C. Wainwright. 2013a. A Remarkable Species Flock of Cyprinodon

Pupfishes Endemic to San Salvador Island, Bahamas. Bull. Peabody Museum Nat. Hist.

54:231–241.

Martin, C. H., and P. C. Wainwright. 2013b. Multiple fitness peaks on the adaptive landscape

drive adaptive radiation in the wild. Science 339:208–11.

Matthews, D. G., and R. C. Albertson. 2017. Effect of craniofacial genotype on the relationship

between morphology and feeding performance in cichlid fishes. Evolution (N. Y). 71:2050–

2061.

McGee, M. D., D. Schluter, and P. C. Wainwright. 2013. Functional basis of ecological

divergence in sympatric stickleback. BMC Evol. Biol. 13:277. BioMed Central.

McGhee, K. E., R. C. Fuller, and J. Travis. 2007. Male competition and female choice interact to

determine mating success in the bluefin killifish. Behav. Ecol. 18:822–830.

O'Neill, M. W., and A. C. Gibb. 2013. Does Feeding Behavior Facilitate Trophic Niche

Partitioning in Two Sympatric Sucker Species from the American Southwest? Physiol.

Biochem. Zool. 87:65–76.

Patek, S. N., J. E. Baio, B. L. Fisher, and A. V Suarez. 2006. Multifunctionality and mechanical

origins: Ballistic jaw propulsion in trap-jaw ants.

R Core Team. 2018. R: A Language and Environment for Statistical Computing. R Found. Stat.

Comput. Vienna, Austria. URL https//www.R-project.org/. 0:{ISBN} 3-900051-07-0.

Richards, E., and C. Martin. 2017. Adaptive introgression from distant Caribbean islands

contributed to the diversification of a microendemic radiation of trophic specialist

pupfishes. PLOS Genet., doi: 10.1101/115055.

Sazima, I., D. De Zoologia, U. E. De Campinas, and S. Paulo. 1983. Scale-eating in characoids

and other fishes. 9.

Shapiro, L. J., J. W. Young, and A. Souther. 2011. Quadrupedal Locomotion of Saimiri

boliviensis: A Comparison of Field and Laboratory-based Kinematic Data. Pp. 335–356 *in*

Primate Locomotion. Springer New York, New York, NY.

Stayton, C. T. 2019. Performance in three shell functions predicts the phenotypic distribution of

hard-shelled turtles. Evolution (N. Y). 73:720–734. John Wiley & Sons, Ltd (10.1111).

Stevens, N. J., J. H. Ratsimbazafy, and F. Ralainasolo. 2011. Linking Field and Laboratory

Approaches for Studying Primate Locomotor Responses to Support Orientation. Pp. 311–

333 *in* Primate Locomotion. Progress and Prospects.

Svanbäck, R., P. C. Wainwright, and L. A. Ferry-Graham. 2002. Linking Cranial Kinematics,

Buccal Pressure, and Suction Feeding Performance in Largemouth Bass. Physiol. Biochem.

Zool. 75:532–543.

Takeuchi, Y., M. Hori, and Y. Oda. 2012. Lateralized kinematics of predation behavior in a Lake

Tanganyika scale-eating cichlid fish. PLoS One 7:e29272. Public Library of Science.

Tebbich, S., K. Sterelny, and I. Teschke. 2010. The tale of the finch: adaptive radiation and

behavioural flexibility. Philos. Trans. R. Soc. Lond. B. Biol. Sci. 365:1099–109. The Royal

Society.

Tobler, M., J. L. Kelley, M. Plath, and R. Riesch. 2018. Extreme environments and the origins of

biodiversity: Adaptation and speciation in sulphide spring fishes.

Tobler, M., R. Riesch, and M. Plath. 2015. Extremophile fishes: An integrative synthesis. Pp.

279–296 *in* Extremophile Fishes: Ecology, Evolution, and Physiology of Teleosts in

Extreme Environments.

Van Wassenbergh, S., T. Lieben, A. Herrel, F. Huysentruyt, T. Geerinckx, D. Adriaens, and P.

Aerts. 2008. Kinematics of benthic suction feeding in Callichthyidae and Mochokidae, with

functional implications for the evolution of food scraping in catfishes. J. Exp. Biol.

212:116–125.

Whitford, M. D., R. W. Clark, G. A. Freymiller, and T. E. Higham. 2019. Determinants of

predation success : How to survive an attack from a rattlesnake. Funct. Ecol. 1–11. John

Wiley & Sons, Ltd (10.1111).

Wood, S. N. 2011. Fast stable restricted maximum likelihood and marginal likelihood estimation

of semiparametric generalized linear models. J. R. Stat. Soc. Ser. B Stat. Methodol. 73:3–

36.

745 **Table 1.** Description of the 8 kinematic metrics and 3 morphological metrics measured from

746 high-speed videos of feeding strikes.

Metric	Measurement
Peak Protrusion (mm)	The distance (mm) from the center of the eye to the anterior tip of the premaxilla.
Time to Peak Protrusion (s)	Time (s) from the start of an attack (20% of peak gape) to peak protrusion.
Peak Gape	The distance (mm) from the anterior tip of the premaxilla to the anterior tip of the dentary.
Time to Peak Gape (s)	Time (s) from the start of an attack to peak gape.
Gape Angle (degrees)	180° minus the angle produced at peak gape between the lower jaw, the quadrate-articular joint, and the ventral surface of the fish beneath the suspensorium (Fig. 2).
Time to Impact (seconds)	Time (s) from the start of an attack to first contact of oral jaws with the prey item.
Starting Distance from prey (mm)	The distance (mm) from the center of the orbit at the start of an attack to the center of the orbit at impact with prey item.
Ram speed (m/s)	Starting distance from prey (m) / time to impact (s)
Body length (mm)	The length (mm) from the anterior tip of the dentary with mouth closed to the posterior end of the hypural plate at the midpoint insertion of the caudal fin.
Lower jaw length (mm)	The length (mm) from the anterior tip of the dentary to the quadrate-articular joint (jaw joint).

756   **Table 2.** Results of a linear discriminant analysis for kinematic variables between strikes on

757   shrimp and scales.

Kinematic Metrics	LD1	LD2	LD3
Peak Protrusion (mm)	0.23	-0.44	0.14
Peak Gape (mm)	1.33	0.47	-0.83
Gape Angle (degrees)	0.08	0.00	0.02
Time to Peak Protrusion (s)	37.30	-12.88	9.90
Time to Peak Gape (s)	-5.10	-14.42	49.24
Time to Impact (s)	9.85	35.43	0.18
Starting Distance from prey (mm)	-614.27	28.85	-545.38
Ram speed (m/s)	25.12	14.23	45.50
Proportion of Trace	0.94	0.05	0.02
Kinematic Metrics	LD1	LD2	LD3
Peak Protrusion (mm)	0.23	-0.44	0.14
Peak Gape (mm)	1.33	0.47	-0.83
Gape Angle (degrees)	0.08	0.00	0.02
Time to Peak Protrusion (s)	37.30	-12.88	9.90
Time to Peak Gape (s)	-5.10	-14.42	49.24
Time to Impact (s)	9.85	35.43	0.18
Starting Distance from prey (mm)	-614.27	28.85	-545.38
Ram speed (m/s)	25.12	14.23	45.50
Proportion of Trace	0.94	0.05	0.02

765 **Table 3.** Results of linear mixed models investigating if strike kinematic variables vary between

766 1) species (generalists, snail-eaters, scale-eaters, or hybrids), 2) food item (shrimp or scales), or

767 3) the interaction between the two. Significant predictors are indicted in bold.

Response	Predictors	χ^2	df	P
Peak Protrusion (mm)				
	Species	4.5	3	0.21
	Food Item	1.2	1	0.27
	log(Body Length)	3	1	0.085
	Species:Food Item	2.3	3	0.51
Peak Gape (mm)				
	Species	**23**	**3**	**3.4×10^{-5}**
	Food Item	1.1	1	0.3
	log(Body Length)	1.3	1	0.26
	Species:Food Item	0.83	3	0.84
Peak gape angle (degrees)	**Species**	**18**	**3**	**0.00037**
	Food Item	0.016	1	0.9
	log(Body Length)	3.4	1	0.066
	Species:Food Item	3.6	3	0.3
Time to Peak Protrusion (s)	Species	4.1	3	0.25
	Food Item	1.1	1	0.3
	log(Body Length)	1.2	1	0.27
	Species:Food Item	3.3	3	0.34
Time to Peak Gape (s)				
	Species	2.5	3	0.48
	Food Item	0.87	1	0.35
	log(Body Length)	2	1	0.16
	Species:Food Item	1.6	3	0.65
Time to Impact (s)				
	Species	3.4	3	0.33
	Food Item	2.1	1	0.14
	log(Body Length)	1.5	1	0.21
	Species:Food Item	4.5	3	0.21
Starting Distance from prey (mm)				
	Species	2.8	3	0.42
	Food Item	0.032	1	0.86
	log(Body Length)	2.1	1	0.15
	Species:Food Item	0.35	3	0.95
Ram speed (m/s)				
	Species	3.7	3	0.3
	Food Item	**4**	**1**	**0.047**

log(Body Length)	1.2	1	0.27
Species:Food Item	2.2	3	0.54

789 **Table 4.** Results of linear mixed models investigating if scale-eating kinematics vary between

790 lab trials and wild strikes observed in situ in Crescent Pond, San Salvador Island, Bahamas.

791 Significant predictors are indicted in bold.

Response	Predictors	χ^2	df	P
Peak Protrusion (mm)				
	Environment	1.5	1	0.22
	log(Body Length)	5.9	1	0.016
Peak Gape (mm)				
	Environment	0.0018	1	0.97
	log(Body Length)	2.6	1	0.11
Gape Angle (degrees)				
	Environment	0.0027	1	0.96
	log(Body Length)	3.1	1	0.078
Time to Peak Protrusion (s)				
	Environment	0.017	1	0.9
	log(Body Length)	1.7	1	0.19
Time to Peak Gape (s)				
	Environment	0.27	1	0.6
	log(Body Length)	3.7	1	0.053
Time to Impact (s)				
	Environment	0.98	1	0.32
	log(Body Length)	2	1	0.16
Starting Distance from prey (mm)				
	Environment	**4.5**	**1**	**0.033**
	log(Body Length)	0.9	1	0.34
Ram speed (m/s)				
	Environment	**11**	**1**	**0.00099**
	log(Body Length)	0.1	1	0.75

799   **Table 5.** Results of linear mixed models investigating if the area removed from a cube by a

800   single bite was affected by any of the PCs from our PCA analysis. Significant predictors are

Response	Predictor	χ^2	df	P
Area Removed from Cube (mm^2)	PC1	1.60	1	0.21
	PC2	**5.49**	**1**	**0.02**
	PC3	0.11	1	0.74
	PC4	0.29	1	0.59
	PC5	**8.80**	**1**	**0.003**
	PC6	0.46	1	0.50
	PC7	1.86	1	0.17
	PC8	0.00	1	1.00
	log(Body Length)	2.21	1	0.14

 Supplemental Material

819 **Table S1. Jaw length varies between species.** Results of linear mixed model investigating if
820 any morphological traits varied between species. Significant predictors are indicated in bold.

Response	Predictors	χ^2	df	P
Log Body Length				
	Species	6.36	3	0.095
Log Jaw Length				
	Species	**45.87**	**3**	**6.039×10^{-10}**

Table S2. PCA Loadings describing kinematic variation in pupfish while feeding on shrimp
and scales. Major loadings (|loading|> 0.4) are indicated in bold.

	PC1	PC2	PC3	PC4	PC5	PC6	PC7	PC8
Peak Protrusion (mm)	0.376	0.142	-0.063	**-0.449**	**0.786**	-0.015	0.077	-0.095
Peak Gape (mm)	0.368	-0.036	-0.175	**-0.669**	**-0.575**	0.131	0.188	0.056
Gape Angle (degrees)	0.08	-0.277	**-0.925**	0.224	0.069	-0.077	-0.015	0.021
Time to Peak Protrusion (s)	**0.419**	-0.188	0.19	0.192	-0.096	**-0.784**	0.287	-0.1
Time to Peak Gape (s)	**0.434**	-0.249	0.123	0.035	-0.063	0.044	**-0.845**	-0.118
Time to Impact (s)	**0.412**	-0.23	0.137	0.39	0.012	0.588	0.391	-0.328
Starting Distance from prey (mm)	**0.407**	0.39	-0.002	0.292	-0.005	0.099	0.004	**0.766**
Ram speed (m/s)	0.135	**0.774**	-0.199	0.159	-0.184	-0.071	-0.099	**-0.518**
Standard Deviation	2.096	1.19	0.987	0.734	0.621	0.427	0.311	0.123
Proportion σ2	0.549	0.177	0.122	0.067	0.048	0.023	0.012	0.002
Cumulative Proportion σ2	0.549	0.726	0.848	0.915	0.963	0.986	0.998	1

825
826
827

828 **Table S3. PCA Loadings describing kinematic variation in pupfish while feeding on fish**
829 **cubes.** Major loadings for PC2 and PC5 (|loading|> 0.4) are indicated in bold.
830

	PC1	PC2	PC3	PC4	PC5	PC6	PC7	PC8
Peak Protrusion (mm)	-0.04	-0.16	0.72	0.43	**-0.51**	0.12	-0.04	0.00
Peak Gape (mm)	-0.30	-0.18	0.55	-0.09	**0.72**	-0.20	0.08	0.01
Gape Angle (degrees)	0.00	0.33	0.36	-0.82	-0.28	0.02	-0.08	0.02
Time to Peak Protrusion (s)	-0.55	0.06	-0.08	-0.03	0.05	0.80	0.21	-0.05
Time to Peak Gape (s)	-0.56	0.08	-0.12	0.08	-0.08	-0.19	-0.79	0.00
Time to Impact (s)	-0.53	0.15	-0.12	0.04	-0.29	-0.50	0.57	0.15
Starting Distance from prey (mm)	-0.12	**-0.62**	-0.10	-0.25	-0.18	-0.10	0.05	-0.69
Ram speed (m/s)	-0.04	**-0.64**	-0.10	-0.24	-0.12	0.06	-0.05	0.71
Standard Deviation	1.73	1.50	1.13	0.90	0.71	0.38	0.18	0.05
Proportion σ2	0.37	0.28	0.16	0.10	0.06	0.02	0.00	0.00
Cumulative Proportion σ2	0.37	0.65	0.81	0.91	0.98	1.00	1.00	1.00

832
Figure S1. Visualization of PC1 and PC2 describing kinematic variation in pupfish while feeding on shrimp and scales. Dots represent individuals, dotted lines represent 95% confidence intervals around species.

836
837
838

839 **Figure S2. Pupfishes approach shrimp more quickly than they do scales.** Colored points
840 represent mean ram speed (m/s) and 95% confidence intervals when consuming shrimp versus
841 scales for all species.
842

844   **Figure S3. Ram speed and starting distance from prey affects bite size.** Graph illustrating the
845   negative relationship between PC2 and the area removed from a fish cube (mm^2) from a single
846   bite. The major loadings of PC2 are ram speed and starting distance from a prey item. Grey dots
847   represent individual fish, the line represents the fitted prediction of a linear model along with a
848   standard error ribbon in blue.

849
850
851

852 **Supplemental Video 1**
853 **Scale-eating pupfish attacks a generalist pupfish in the wild.**
854
855
856