Rapid adaptive evolution of scale-eating kinematics to a novel ecological niche

Michelle E. St. John[1], Roi Holzman[2,3], Christopher H. Martin[1*]

[1]Department of Integrative Biology and Museum of Vertebrate Zoology, University of California, Berkeley, CA 94720, USA

[2]School of Zoology, Tel Aviv University, Eilat, Israel
[3]Inter-University Institute for Marine Sciences, Eilat, Israel

[*]Corresponding Author: chmartin@berkeley.edu

Short title: Scale-eating specialists evolved adaptive feeding kinematics.

keywords: feeding kinematics, novelty, key innovation, lepidophagy, performance, postzygotic

isolation, hybrid kinematics

Abstract: 218 words, Main text plus references: figures: 2 Tables, 4 Figures, Supplemental

Tables S1-S2, Figures S1, and 1 Supplemental video S1

Abstract

The origins of novel trophic specialization, in which organisms begin to exploit novel resources

for the first time, may be explained by shifts in behavior such as foraging preferences or feeding

kinematics. One way to investigate the behavioral mechanisms underlying ecological novelty is

by comparing prey capture kinematics between groups. In this study, we investigated the

contribution of kinematics to the origins of a novel ecological niche for scale-eating within a

microendemic adaptive radiation of pupfishes on San Salvador Island, Bahamas. We compared

prey capture kinematics across three species of pupfish while consuming shrimp and scales in the

lab and found that scale-eating pupfish exhibited peak gape sizes that were twice as large as all

other groups, but also attacked prey with a more obtuse angle between their lower jaw and

suspensorium. We then investigated how this variation in feeding kinematics could explain

scale-biting performance by measuring the surface area removed per strike from standardized

gelatin cubes. We found that a combination of larger peak gape and more obtuse lower jaw and

suspensorium angles resulted in 67% more surface area removed per strike, indicating that scale-

eaters may reside on a performance optimum for scale-biting. We also measured feeding

kinematics of F1 hybrids to test whether feeding performance could contribute to reproductive

isolation between species and found that F1 hybrid kinematics and performance more closely

resembled those of generalists, suggesting that they may have low fitness in the scale-eating

niche. Ultimately, our results suggest that the evolution of strike kinematics in this radiation is

an adaptation to the novel niche of scale-eating.

Introduction

Determining how organisms use resources for the first time and occupy novel niches is an outstanding question in evolutionary ecology. Many changes accompany adaptation to a novel niche and previous studies have investigated how shifts in behaviors (Bowman and Billeb 1965; Tebbich et al. 2010; Curry and Anderson 2012), morphologies (Ferry-Graham et al. 2001; Ferry-Graham 2002; Hata et al. 2011; Davis et al. 2018), physiologies (Arias-Rodriguez et al. 2011; Tobler et al. 2015, 2018), and kinematics (Janovetz 2005; Patek et al. 2006; Cullen et al. 2013; McGee et al. 2013) can all facilitate this transition.

Shifts in kinematic traits—particularly those which affect prey capture and feeding—are especially promising, because they can provide biomechanical insights into the origins of novel trophic niches. For example, the wimple piranha (*Catoprion mento*) uses a ram attack coupled with a uniquely large gape angle to knock scales free from its prey (Janovetz 2005); Syngnathiform fishes specialize on evasive prey items using power-amplified jaws (Longo et al. 2018); and the Pacific leaping blenny (*Alticus arnoldorum)* is able to feed and reproduce on land by using unique axial tail twisting to improve propulsion and stability for greater jumping performance (Hsieh 2010).

Differences in prey capture kinematics between species may also contribute to postzygotic extrinsic reproductive isolation by reducing hybrid feeding performance (Higham et al. 2016b), which may lead to speciation (Henning et al. 2017; Matthews and Albertson 2017). For example, McGee et al. (2015) measured prey capture kinematics and performance in two sunfish species (Centrarchidae) and their naturally occurring hybrids. Hybrid sunfish displayed intermediate gape size compared to parental types and initiated strikes from an intermediate distance, yet their actual feeding performance was less than predicted from additive traits.

Hybrid Lake Victoria cichlids (produced by crossing thick-lipped *Haplochromis chilotes* and

thin-lipped *Pundamilia nyererei* parents) also exhibited lower foraging performance compared to

parental species, most likely due to antagonistic pleiotropy and genetic correlations between

head and lip morphology (Henning et al. 2017). Despite these findings, few studies investigate

how hybrid kinematics affects the evolution of novelty or explicitly connect kinematics to

performance consequences.

Scale-eating (lepidophagy) is excellent for connecting a mechanistic understanding of

feeding kinematics with adaptation to a novel trophic niche. It is an extremely rare trophic niche

which has independently evolved only 19 times in approximately 100 fish species out of over

35,000 (Sazima 1983; Martin and Wainwright 2013a; Kolmann et al. 2018). However, not much

is known about its evolutionary origins or its kinematics. Current hypotheses for the origins of

scale-eating vary, but they all take a strict behavior-first approach (Greenwood 1965; Sazima

1983; St. John et al. 2018) suggesting that kinematic variation may also contribute to the origins

of scale-eating. However, only a few studies have investigated the feeding kinematics and

performance of scale-eating fishes. Janovetz (2005) measured feeding kinematics of *C. mento*

while consuming: 1) free floating scales, 2) whole fish, and 3) scales off the sides of fish, and

found that scale-eating kinematics were different from those used during suction-feeding or

biting. Interestingly, scale-eating attacks produced gape angles that ranged from 30-100% larger

than those produced from consuming free-floating scales or whole fish respectively— suggesting

that a larger gape is necessary for scale-eating. Furthermore, this variation in gape angle across

food items was documented within individuals, indicating that scale-eating kinematics may be

behaviorally mediated (Janovetz 2005). The feeding kinematics of the Lake Tanganyikan scale-

eating cichlid, *Perissodus microlepis,* have also been examined (Takeuchi et al. 2012; Takeuchi

and Oda 2017). While many studies of this species focus on how kinematics interacted with *P.*

microlepis' antisymmetric mouth morphology, and not on scale-eating kinematics *per se*, they

have still documented a significant interaction between kinematic traits, behavior, and

morphology. For example, *P. microlepis* were able to perform more successful scale-eating

strikes using their dominant side (Takeuchi et al. 2012; Takeuchi and Oda 2017). A similar oral

jaw antisymmetry and behavioral laterality was documented in a scale-eating characiform

(*Exodon paradoxus;* Hata et al. 2011). While these studies provide valuable insights into scale-

eating kinematics and performance, the lack of comparative data on the kinematics of closely

related non-scale-eating species or hybrids has so far limited further investigations of the origins

of scale-eating.

The overall goal of our study was to fill the following knowledge gaps and shed light on

the relationship between kinematic traits and occupation of a novel niche: First, comparisons of

scale-eating kinematics across scale-eating and closely related non-scale-eating outgroup species

is necessary for investigating the origins of novelty. Without the comparative method it is

impossible to determine which kinematic variables are unique or important for scale-eating.

Second, very few kinematic studies investigate hybrid kinematics. Understanding hybrid

kinematics, especially in the context of novelty, is informative because 1) impaired performance

in hybrids is a form of extrinsic postzygotic isolation between species (McGee et al. 2015;

Higham et al. 2016b) and 2) it can allow the decoupling of morphology, behavior, and

kinematics making it easier to identify causative traits underlying performance (Holzman and

Hulsey 2017). Finally, few studies connect observed variation in kinematics to variation in whole

organism feeding performance (but see: Svanbäck et al. 2003; Takeuchi et al. 2012; Whitford et

al. 2019). Making this connection is important because it can identify kinematic traits associated

with performance tasks relevant to evolutionary fitness rather than simply describing phenotypic

variation in kinematic traits, most of which may not be relevant to performance or fitness

(Arnold 1983; Hu et al. 2017).

The scale-eating pupfish (*Cyprinodon desquamator*) is an excellent organism to

investigate the interaction of kinematics and novelty for several reasons. First, the scale-eating

pupfish evolved within a recent sympatric radiation of pupfishes on San Salvador Island,

Bahamas. This radiation is endemic to a few hypersaline lakes on the island (Martin and

Wainwright 2013a), which were most likely dry during the last glacial maximum 10-15 kya

(Hagey and Mylroie 1995). Second, the radiation provides closely related sister taxa for

kinematic comparison including: 1) the scale-eating pupfish, 2) a generalist pupfish (*C.

variegatus*), and 3) a snail-eating pupfish (*C. brontotheroides*). Phylogenetic evidence suggests

that scale-eating pupfish form a clade across all lakes where they are found on San Salvador and

that this clade is sister to a clade containing generalists and snail-eaters (Martin and Feinstein

2014; Lencer et al. 2017), although gene flow is still ongoing among all three species (Richards

and Martin 2017). All three pupfish species can be crossed in the lab to measure the kinematics

and performance of hybrid phenotypes.

The morphological similarities and differences between San Salvador pupfishes have also

previously been described. Specifically, 1) all pupfish species exhibit vestigial ascending

processes of the premaxilla allowing for independent movement of the upper and lower jaws

during jaw protrusion (Hernandez et al. 2009, 2018), and 2) scale-eating pupfish have two-fold

larger, supra-terminal oral jaws compared to the smaller, terminal jaws of the generalist or snail-

eating pupfish (Martin and Wainwright 2011, 2013a; Martin 2016). Their divergent morphology,

along with the findings of Janovetz (2005), leads us to predict that scale-eating pupfish should

have larger gapes during scale-eating attacks compared to other species of pupfish, and that this

peak gape should be the result of a larger angle between the anterior tip of the premaxilla, the

quadrate-articular joint, and the anterior tip of the dentary.

We investigated the interaction between kinematics and novelty in pupfishes by

recording the high-speed feeding strikes of San Salvador generalist, snail-eating, and scale-eating

pupfishes, along with F1 hybrids. We asked: 1) if scale-eating pupfish differed in their feeding

kinematics compared to other groups during scale-eating and suction-feeding strikes, 2) whether

variation in kinematics was associated with bite performance, and 3) if F1 hybrid feeding

kinematics differed from parental species. Ultimately, we found that the feeding kinematics of

scale-eating pupfish diverged from all other species and were not solely due to their increased

oral jaw size. Instead, scale-eaters may be behaviorally mediating their feeding kinematics to

optimize the surface area removed per strike, suggesting that scale-eater kinematics are a recent

adaptation to scale-eating.

Methods

Collection and Husbandry

We used seine nets to collect generalist, snail-eating, and scale-eating pupfishes from Crescent

Pond, Little Lake, and Osprey Lake on San Salvador Island, Bahamas in July, 2017 and March,

2018. Wild-caught fish were maintained in 37-75L mixed-sex stock tanks at a salinity of 5-10

ppt and temperature of 23-27℃. While in stock tanks, fish were fed a diet of frozen bloodworms,

mysis shrimp, and commercial pellet foods daily. In the lab, we crossed generalist and scale-

eating pupfishes from both Little Lake and Crescent Pond to produce F1 hybrid offspring. Prior

to filming, pupfishes were isolated in 2L tanks to maintain individual IDs throughout the study.

Feeding kinematics

We recorded pupfishes feeding on three different food items: frozen mysis shrimp (Mysida,

Hikari Inc.), scales, and standardized gelatin cubes (dimensions: 1.5 cm x5cm X 1.5 cm x 1.5 cm

cube; Repashy Superfoods, Community Plus Omnivore Gel Premix; prepared following

manufacturer's instructions). We measured feeding kinematics while fish consumed both shrimp

and scales because it allowed us to ask if 1) scale-eating pupfish differed in their feeding

kinematics compared to other groups, 2) if the kinematics of scale-eating strikes differ from

those used during suction feeding (e.g. shrimp), and 3) if F1 hybrid feeding kinematics differed

from parental species. We explicitly examined F1 hybrid kinematics in this study because

lowered hybrid feeding performance may contribute to reproductive isolation between species

and may shed light on the underlying genetics of feeding kinematics. We additionally measured

feeding kinematics across all groups while fish consumed gelatin cubes to ask whether variation

in kinematic traits affected feeding performance (i.e. bite size).

In the lab, fish freely consumed mysis shrimp, but we had to train all species to feed on

scales from the sides of euthanized zebrafish (*Danio rerio*; stored frozen) and to feed from

gelatin cubes (stored at 4□). For training, we isolated each fish in a 2-liter plastic tank and

presented a given food item (either euthanized zebrafish or gelatin cube) daily. If a pupfish began

feeding on the item, it was left in the tank until the pupfish stopped feeding. If a pupfish did not

begin feeding within one minute, the food item was removed from the tank. Any pupfish that did

not feed received a supplemental feeding of commercial pellet food (New Life Spectrum Thera-

A, medium sinking pellets). If an individual did not feed on a training item for more than two

days, we reduced supplemental feedings to once every two days to ensure that the fish was

sufficiently motivated. Once pupfish reliably began feeding on either scales or gelatin cubes, we

proceeded to film their feeding behaviors according to the filming protocol below. Fish were

never trained on more than one item at a time, and we instead ensured that all filming was

completed for a single food item before proceeding to train for the next item.

For all three food items, we used a Sony Cyber-shot DSC-RX10 III (480fps) or Sony

Cyber-shot DSC-RX100 IV 20.1 MP (480fps) for high-speed video of foraging strikes.

Illumination was provided by a dimmable bi-color 480 LED light (Neewer) positioned ~0.3 m

from the filming tank. Pupfish were allowed to acclimate to the lighting before feeding

commenced. Fish were considered acclimated when they moved around their tank freely (usually

after ~5 minutes). For scale-eating we used forceps to hold a euthanized zebrafish horizontally in

the water column and perpendicular to the front of an individual. For mysis shrimp and gelatin

cubes, we dropped the food item a few inches in front of an individual. All videos were taken

from a lateral perspective. Once filming for one food item was completed, the process was

repeated until we filmed each individual consuming all three food items.

Kinematic analyses

Videos were converted to image stacks and analyzed using the image processing software (FIJI;

Schindelin et al. 2012). To quantify feeding performance, we measured 10 kinematic trait

metrics including 1) Peak jaw protrusion, defined as the distance (mm) from the center of the

orbit to the anterior tip of the premaxilla. 2) Time to peak protrusion, defined as the time (s) from

the start of an attack (defined as 20% of peak gape) to peak protrusion. 3) Peak gape, defined as

the distance (mm) from the anterior tip of the premaxilla to the anterior tip of the dentary. 4)

Time to peak gape, defined as the time (s) from the start of an attack at 20% of peak gape to peak

gape. 5) Gape angle was the angle (degrees) produced at peak gape between the anterior tip of

the premaxilla, the quadrate-articular joint, and the anterior tip of the dentary. 6) Lower jaw

angle was the angle produced at peak gape between the lower jaw, the quadrate-articular joint,

and the ventral surface of the fish beneath the suspensorium (Figure 1&2). 7) Time to impact

was the time (s) from the start of an attack (20% peak gape) to first contact of oral jaws with the

prey item. 8) Time from peak gape to impact was the difference between the time to impact (s)

and the time to peak gape (s). 9) Starting distance from prey was the distance (mm) from the

center of the orbit at the start of an attack to the center of the orbit at impact with prey item.

Finally, 10) ram speed was the starting distance from prey at 20% of peak gape (m) divided by

the time to impact (s). In addition to our kinematic metrics, we also measured body length and

lower jaw length (Table S1) using images from the video. We calibrated each video using a grid,

positioned at the back of the filming tank.

Measuring bite performance

In order to connect variation in feeding kinematics to variation in bite size we recorded high-

speed strikes on gelatin meal replacement for fish in the shape of a 1.5 x 1.5 x 1.5 cm cube. We

filmed a feeding strike on a single cube and immediately removed the cube from the tank. The

gel cube retains its shape in water and therefore allowed us to precisely photograph and measure

the area removed by each bite. We used an Olympus Tough TG-5 camera to take photos of each

lateral surface of the cube –ensuring that we had photographed the entire bite—and measured the

total surface area removed from the cube (Figure 3D). One caveat is we did not measure the

depth of the bite, which may be affected by additional kinematic variables during the strike.

However, scale-eating attacks observed in the lab and field do not typically produce deep

wounds where bite depth may be relevant, thus we expect that surface area is the best proxy for

scale-biting performance in this system. Although bites were removed from both the lateral

surface and edge of the gelatin cubes during strikes, there was no significant difference in surface

area removed (*t*-test, $P = 0.12$).

Statistical analyses

Comparing strike kinematics

We collected and analyzed 101 feeding strikes from 31 individuals striking both shrimp and

scales (7 generalists; 7 snail-eaters; 9 scale-eaters; 8 F1 hybrids). We used linear mixed models

(LMMs) in the lme4 package (Bates et al. 2014) in R (R Core Team 2018) to determine if any of

our kinematic metrics varied between species or food item. In each model we included: 1) the

kinematic metric as the response variable, 2) species designation, food item, and their interaction

as fixed effects, 3) individual fish IDs and population nested within species as random effects,

and 3) log body size as a covariate (Table 1). Although we compared kinematic data across

multiple species, very few genetic variants are fixed between species (<1,000 SNPs out of 12

million) and generalists and molluscivores cluster by lake rather than by species (McGirr and

Martin 2017; Richards and Martin 2017). Thus, is it appropriate to analyze species differences at

these recent timescales as population-scale data using mixed model analyses of independent

populations (e.g. Hatfield and Schluter 1999; McGee et al. 2013), rather than phylogenetic

comparative methods.

We also performed a linear discriminant analysis (LDA) on the combined shrimp and

scales kinematic data to reduce dimensionality and identify which kinematic metrics contributed

most to differences between species (Table 2, Figure 1A). We used a MANOVA and Wilks' ⍰ to

assess the significance of the LDA. We did not have enough degrees of freedom to perform these

analyses with all of our kinematic variables, so we excluded time to peak protrusion and time to

impact as they were highly correlated with time to peak gape (Table S2, $r^2 > .85$), and also

excluded distance from prey as it was highly correlated with ram speed (Table S2, $r^2 = .90$). Our

MANOVA ultimately included 1) peak protrusion, peak gape, time to peak gape, gape angle,

lower jaw angle, time from peak gape to impact, and ram speed as response variables, 2) species

designation as a predictor variable, and 3) individual ID as a random effect.

Determining how kinematic variables affect bite performance

We collected and analyzed 31 strikes on cubes across all three species and F1 hybrids. We used

generalized additive models (GAMs) from the mgcv package (Wood 2011) in R to investigate

how peak gape, peak protrusion, gape angle and lower jaw angle affected bite size. We used

GAMs for this analysis because they do not assume a linear relationship between performance

(i.e. bite size) and our given kinematic variables, but instead can fit smoothing splines to the data

to test for nonlinear associations. We used AIC scores to select our optimal model (Table 3). We

started with the most complete model which included 1) area removed from a cube (mm^2) as the

response variable 2) a spline modeling the interaction between two of our predictor variables,

and 3) a single fixed effect. There were insufficient degrees of freedom to test all four terms at

once in this model, therefore we tested all combinations of this model with our four predictor

variables (Table 3A). We also tested all nested versions of this complex model by 1) removing

the interaction term, but maintaining two splines and a fixed effect (Table 3B), 2) removing one

spline and including three fixed effects (Table 3C), and finally 3) by testing the model with all

four variables as only fixed effects (Table 3D). Ultimately, our best fitting model included area

removed from a cube (mm^2) as the response variable, univariate smoothing splines for peak gape

and Gape angle, and peak protrusion as a fixed effect (average ΔAIC=24.96).

Finally, we predicted the area removed per bite for each fish from their peak gape and

gape angle kinematic measurements using a machine-learning algorithm from the caret package

(Kuhn 2008) using a spline-based method. We could not measure the realized bite size for all

scale-eating strikes (feeding off a euthanized zebrafish) and suction-feeding strikes (mysis

shrimp). We therefore used a GAM model, estimating the effect of gape size and gape angle on

the area removed from gelatin cubes, to predict bite performance (surface area removed) for all

scale-biting and suction-feeding strikes in our dataset. Ideally, we would have used our best-

fitting GAM model, which also included peak protrusion as a fixed effect. However, the caret

package currently only accepts two fixed effects, and peak protrusion ultimately did not affect

bite size ($P = 0.078$). We trained the model using all strikes observed on gelatin cubes (31

strikes across all three species and F1 hybrids) and 10-fold cross-validations with three repeats as

the resampling scheme. We tested the accuracy of this model by comparing fitted values from

the model to observed values from the data set and found that our model was able to predict 70%

of the variance in the gelatin-strike dataset (df=1, F= 66.81 , P=5.17x10^{-9}, R^2=0.70). We then

used this model to predict the area removed for each scale-biting and suction-feeding strike

based on the kinematic measurements alone. We used bootstrap resampling (10,000 iterations) to

calculate mean bite size (predicted area removed) and 95% confidence intervals for each species.

Determining if hybrid kinematics match additive predictions

We calculated the predicted values for peak gape, lower jaw angle, and bite size for the scale-

eater x generalist F1 hybrids under the hypothesis that these kinematic traits would be additive

and therefore intermediate between generalist and scale-eater values. We used a one sample *t*-test

to test whether the observed values of the three traits (peak gape, lower jaw angle, predicted bite

sizes) for F1 hybrids deviated from additive predictions.

Results

Scale-eaters exhibited divergent feeding kinematics compared to other pupfishes

Scale-eaters exhibited divergent feeding kinematics, while consuming both shrimp and scales,

compared to other groups (Figure 1A). A MANOVA supported the significance of this

discriminant analysis and found species designation was a significant predictor of kinematics

(Wilks' λ = 0.13; F = 3.05; *df* = 3; *P*= 0.00036). Species significantly varied in their peak gape

and lower jaw angles during feeding strikes—regardless of the food item— in a linear mixed

model controlling for individual ID and body length (Table 1). This pattern was driven by scale-

eaters who had peak gapes that were twice as large as other species, but also had lower jaw

angles with their suspensorium that were 14% more obtuse than other species (Figure 1B, C).

Thus, the scale-eaters' unique angle of the jaw complex with respect to their body, along with

their greatly enlarged oral jaws, allows them to have increased peak gape while maintaining the

same gape angle as other species (Figure 2). This may allow their upper jaws to more effectively

'rake' scales from the prey surface. Ram speed was the only kinematic variable that marginally

varied between food items: strikes on shrimp were approximately 16% faster than those on scales

(Table 1, Figure S1; *P* = 0.053).

Variation in strike kinematics affected bite size performance

328 GAM modeling indicated that the univariate splines of peak gape and gape angle were both

329 significantly associated with bite size (peak gape: *edf*=5.30, *F*=8.96, *P*= 6.33x10^{-5}; gape angle:

330 *edf* = 7.48, *F*= 2.78, *P*=0.032). The fixed effect of peak protrusion, however, was only

331 marginally significant (*t*=-1.88, *P*=0.078). Ultimately this model explained 84.5% of the

332 observed deviance in bite size, and suggests that a larger gapes paired with gape angles of 80°

333 are associated with larger surface areas removed per bite (Figure 3A-C).

335 *F1 hybrid kinematics are not strictly additive and more closely resemble generalist kinematics*

336 F1 hybrid feeding kinematics differed from scale-eater kinematics (Tukey HSD, peak gape: *P* =

337 1.2x10^{-6}, lower jaw angle: *P* = 0.0090), but were not significantly different from generalist

338 kinematics (Tukey's HSD, peak gape: *P* = 0.21, lower jaw angle: *P* = 0.37). Mean hybrid peak

339 gape was 39% smaller than scale-eater peak gape and 32% larger than generalist peak gape

340 (Figure 1B). Similarly, mean hybrid lower jaw angle was 9.5% more acute than scale-eater peak

341 lower jaw angle, and 5.6% more obtuse than the mean generalist lower jaw angle (Figure 1C). F1

342 hybrids failed to match additive predictions of intermediate kinematics for peak gape (t-test, μ=

343 3.035, mean= 2.52 mm, *P* = 0.013), but did meet these predictions for lower jaw angle (t-test, μ=

344 136.5, mean= 133.92 degrees, *P = 0.18*). Our machine learning model also predicted that scale-

345 eater kinematics would result in bite sizes that are 60% larger than the predicted bites of the

346 other species (Figure 4). Interestingly, estimates for F1 hybrid bite sizes were slightly smaller

347 than expected based on additive predictions (*t*-test, μ= 320.5, mean= 255.24, *P* = 0.13).

348
349 **Discussion**
350 **Scale-eating pupfish have divergent feeding kinematics**

Scale-eating pupfish exhibited peak gapes that were twice as large as other groups, but

simultaneously displayed gap angles that were not different from other groups, and lower jaw

angles that were 12% more obtuse. Thus, scale-eaters kept their jaws more closed during strikes

compared to other species, resulting in smaller gape sizes than the maximum achievable gape

given their morphology. These counterintuitive results only partially support our prediction that

scale-eaters should have larger peak gapes, similar to the findings of Janovetz (2005) for the

scale-eating piranha. Increased gape size in scale-eating pupfish was not due to an increased

gape angle as we predicted. Instead, scale-eaters appear to maintain the same gape angle of their

oral jaws as in other species (~80°) and increased their lower jaw angle resulting in more closure

of their jaws during strikes, effectively lowering the maximum peak gape possible due to their

enlarged oral jaws (Figure 2). For example, if a scale-eater decreased their lower jaw angle from

150° to 130°, the mean lower jaw angle of generalists, they could increase their peak gape by

about 8%. One possibility is that this more obtuse lower jaw angle is an artifact of filming scale-

eating strikes in the lab. To investigate this, we analyzed four scale-eating strikes performed by

wild scale-eaters observed in Crescent Pond, San Salvador Island, Bahamas (filmed using a

Chronos camera (Kron Technologies, model 1.4, 16 GB memory, Color image sensor) with an

f1.4 zoom lens in a custom underwater housing (Salty Surf, Inc. Krontech Chronos 1.4 housing

with M80 flat port)) and compared their jaw angles to those seen in the lab. Wild strikes had an

even more obtuse mean lower jaw angle of 168° while scale-eating strikes in the lab had a mean

lower jaw angle of 153° suggesting that an obtuse lower jaw angle is also used during natural

scale-eating strikes in hypersaline lakes on San Salvador Island.

Strike kinematics did not vary across prey items (Table 1), contrary to Janovetz (2005).

In fact, the only kinematic variable that remotely varied between prey items was ram speed

(Table 1, Figure S1), but this may simply be due to the fact that shrimp were a moving target

during feeding trials while euthanized zebrafish were held stationary with forceps for scale-

eating strikes. Alternatively, phenotypic plasticity due to rearing environment could produce a

similar pattern; however, we did not observe any differences in strike kinematics between wild

caught and lab-reared fish.

Is jaw morphology solely responsible for kinematic variation?

The kinematic variables that varied the most between scale-eating and non-scale-eating

pupfishes were peak gape and lower jaw angle—both related to the size of the oral jaws.

Previous work has documented that the oral jaws of scale-eating pupfish are two-fold than their

sister species (Holtmeier 2001; Martin and Wainwright 2013a; Martin 2016) and may be

controlled by four moderate-effect QTL with all positive effects on jaw size, consistent with

directional selection on this trait (Martin et al. 2017). It may be that increased oral jaw size is

sufficient to create variation in feeding kinematics without an accompanying shift in behavior.

Previous studies have documented how changes in morphology alone can alter feeding

kinematics. For example, kinematic studies have found that the scaling of the lower jaw in

bluegill (Wainwright and Shaw 1999) and body size in largemouth bass (*Micropterus salmoides;*

Richard and Wainwright 1995) both significantly affected prey capture kinematics. Furthermore,

Ferry-Graham et al. (2010) used the pike killifish (*Belonesox belizanus*) to show that simply

doubling the length of the jaws significantly affected key kinematic variables such as peak gape

size—even while keeping lower jaw angle constant. Simply stated, the key adaptation necessary

for scale-eating may be enlarged, supra-terminal jaws. If this hypothesis were true, we would

expect that peak gape would increase with jaw size and that gape angle would increase with the

shift from terminal to supra-terminal jaws, but all other kinematics variables would remain

constant across species. Our results reject this hypothesis. Instead, scale-eaters maintain the gape

angle observed in other species and increase their lower jaw angle with the suspensorium by 12

degrees resulting in a reduction in their potential peak gape size (Figure 2). This suggests that

scale-eaters have evolved more obtuse lower jaw angles during strikes to increase feeding

performance (Figures 3&4).

Scale-eating performance optimum

Scale-eaters may have reduced their lower jaw angles relative to other species in order to remain

on a performance optimum for scale-eating. Our models of bite performance supported this: peak

gapes larger than approximately 4.5 mm counterintuitively resulted in smaller bite sizes (Figure

3A&B). An enlarged angle in scale-eating pupfish results in a lower jaw that points directly

towards the prey during strikes – possibly resulting in greater stability for biting scales while

retaining a large gape (Figure 2). This large gape and unique jaw alignment may allow scale-

eaters to attack prey from a roughly perpendicular angle (as frequently observed during field

observations) —appearing to wrap their large lower jaw under prey items and subsequently

scraping scales from their sides using their independently protrusible upper jaws (also observed

in a scale-eating characin: Hata et al. 2011). Interestingly, perpendicular angles of attack and

large gapes are associated with scraping in benthic feeding fish (Van Wassenbergh et al. 2008;

O'Neill and Gibb 2013). In fact, one prominent hypothesis for the origins of scale-eating is that it

arose from an algae-scraping ancestor (Sazima et al. 1983). One caveat for this hypothesis,

however, is that our current performance estimates do not include all possible combinations of

peak gape and lower jaw angle and few observations of the largest peak gape sizes. Future work

should estimate performance across multiple performance axes (e.g. Stayton 2019, Keren et al.

2018, Dickson and Pierce 2019), ideally using F2 hybrids. F2 hybrids are a useful tool for this

type of experiment, as they are the first generation of offspring where recombination has the

potential to produce new combinations of kinematic, morphological, and behavioral traits not

observed in the F0 or F1 generations.

Non-additive F1 hybrid feeding kinematics may contribute to reproductive isolation of

scale-eaters

It is well documented that complex performance traits, such as feeding kinematics, are most

likely controlled by numerous loci (i.e. polygenic), and can mostly be described as additive

(reviewed in Sella and Barton 2019). We therefore expected F1 hybrids to exhibit intermediate

kinematics and performance relative to both parental species. Instead, we found that F1 hybrid

kinematics more closely resembled generalists (Table 1; Figure 1) and that their estimated bite

sizes are slightly below additive estimates (additive prediction= 341.48, predicted = 289.72, $P =$

0.058) suggesting that F1 hybrids may have higher performance in a generalist trophic niche.

Current evidence from field fitness experiments supports the idea that hybrid pupfish exhibit

better performance in the generalist ecological niche compared to their performance in the scale-

eater niche. One field experiment in these lakes measured hybrid fitness in the wild and found

high mortality and low growth rates for hybrids most closely resembling the scale-eating

phenotype (Martin and Wainwright 2013b). Furthermore, for the few hybrids resembling scale-

eaters which did survive, only 36% had recently consumed any scales compared to 92% of wild-

caught scale-eaters (Martin and Wainwright 2013a,b). Impaired hybrid performance in the scale-

eating niche may contribute to extrinsic postzygotic isolation between species (McGhee et al.

2007; McGee et al. 2013; Higham et al. 2016). Reproductive isolation may also evolve more

quickly in species that occupy a more distant fitness peak with a larger fitness valley such as the

scale-eating pupfish due to stronger selection against hybrids and reinforced pre-mating isolation

(Martin and Feinstein 2014). Thus, impaired hybrid scale-eating performance could also

contribute to increased diversification rates through the mechanism of a wider fitness valley.

Low hybrid performance may also be due to the morphological differences between scale-

eaters and generalists. As mentioned above, it is possible that a shift in morphology – such as

enlarged oral jaws in scale-eaters—may be sufficient to change kinematic profiles alone. F1

hybrid kinematics clearly differed from scale-eater kinematics, but their jaw lengths were also

significantly smaller than the jaws of scale-eaters (Tukey's HSD, $P = 5.21 \times 10^{-5}$). Furthermore,

previous work has shown that F1 hybrid pupfish offspring (produced from generalist x scale-

eater crosses) tend to develop along a more similar trajectory to their maternal parent (Holtmeier

2001). This could indicate that F1 hybrid pupfish with scale-eating mothers are more likely to

develop jaws resembling a purebred scale-eater, but may also retain their generalist-like

kinematics. The resulting mismatch between morphology, kinematic traits, and ecological niche

may be driving low hybrid survival in the scale-eating niche and contributing to reproductive

isolation between generalist and scale-eating pupfish species.

Conclusion

In conclusion, this study suggests that shifts in kinematic traits may have preceded or facilitated

the origin of scale-eating in *Cyprinodon* pupfishes. Scale-eating pupfish exhibited peak gapes

that were twice as large as other pupfish species, but simultaneously had lower jaw angles that

were significantly larger. Surprisingly, we found that this unique combination of scale-eater

kinematics may reside on a performance optimum, as large peak gapes and large lower jaw

angles resulted in larger surface areas removed per strike. Impaired F1 hybrid kinematics and

performance in the scale-eating niche also suggests that kinematic traits contribute to

reproductive isolation of the scale-eating pupfish and the evolution of novelty. Future work

should investigate if other performance optima exist on the kinematic landscape and whether F2

hybrid fitness in the wild is reduced due to a mismatch between morphology and feeding

kinematics.

Data Accessibility

All analyses reported in this article can be reproduced using the data provided by St. John and

Martin (2019). Raw data will be deposited in Dryad.

Acknowledgements

We thank the University of North Carolina at Chapel Hill, NSF CAREER 1749764, NIH

5R01DE027052-02, and BSF 2016136 for funding to CHM and the UNC Quality Enhancement

Plan for course-based undergraduate research (CURE) funding to CHM. The Bahamas

Environmental Science and Technology Commission and the Ministry of Agriculture kindly

provided permission to export fish and conduct this research. Rochelle Hanna, Velda Knowles,

Troy Day, and the Gerace Research Centre provided logistical assistance in the field; Kristi

Dixon, Casey Charbonneau, Gabriel Harris, Amelia Ward, Delaney O'Connell, and the

undergraduate students of BIO221L 'Evolution of Extraordinary Adaptations' assisted with high-

speed videography and kinematic analysis. All animal care protocols were approved by the

University of California, Berkeley and the University of North Carolina at Chapel Hill Animal

Care and Use Committees.

References

Arias-Rodriguez, L., M. Tobler, M. Palacios, F. J. García de León, D. Bierbach, M. Mateos, I.

Mitrofanov, M. Plath, and L. J. Chapman. 2011. Evolution in extreme environments:

replicated phenotypic differentiation in livebearing fish inhabiting sulfidic springs.

Evolution (N. Y). 65:2213–2228.

Arnold, S. J. 1983. Morphology, Performance and Fitness. Am. Zool. 23:347–361.

Bates, D., M. Mächler, B. Bolker, and S. Walker. 2014. Fitting Linear Mixed-Effects Models

using lme4. J. Stat. Softw. 67:1–48.

Bowman, R. I., and S. L. Billeb. 1965. Blood-eating in a Galápagos finch. Living Bird 4:29–44.

Cullen, J. A., T. Maie, H. L. Schoenfuss, and R. W. Blob. 2013. Evolutionary Novelty versus

Exaptation: Oral Kinematics in Feeding versus Climbing in the Waterfall-Climbing

Hawaiian Goby Sicyopterus stimpsoni. PLoS One 8:e53274. Public Library of Science.

Curry, R. L., and D. J. Anderson. 2012. Interisland Variation in Blood Drinking by Galápagos

Mockingbirds. Auk 104:517–521.

Davis, A. L., M. H. Babb, M. C. Lowe, A. T. Yeh, B. T. Lee, and C. H. Martin. 2018. Testing

Darwin's Hypothesis about the Wonderful Venus Flytrap: Marginal Spikes Form a "Horrid

Prison" for Moderate-Sized Insect Prey. Am. Nat. 193:309–317.

Ferry-Graham, L. A. 2002. Using Functional Morphology to Examine the Ecology and Evolution

of Specialization. Integr. Comp. Biol. 42:265–277.

Ferry-Graham, L. A., L. P. Hernandez, A. C. Gibb, and C. Pace. 2010. Unusual kinematics and

jaw morphology associated with piscivory in the poeciliid, Belonesox belizanus. Zoology

113:140–147.

Ferry-Graham, L. A., P. C. Wainwright, C. Darrin Hulsey, and D. R. Bellwood. 2001. Evolution

and mechanics of long jaws in butterflyfishes (Family Chaetodontidae). J. Morphol.

248:120–143.

Hata, H., M. Yasugi, and M. Hori. 2011. Jaw laterality and related handedness in the hunting

behavior of a scale-eating characin, exodon paradoxus. PLoS One 6.

Hatfield, T., and D. Schluter. 1999. Ecological Speciation in Sticklebacks: Environment-

Dependent Hybrid Fitness. Evolution (N. Y). 53:866. Mayr.

Henning, F., G. Machado-Schiaffino, L. Baumgarten, and A. Meyer. 2017. Genetic dissection of

adaptive form and function in rapidly speciating cichlid fishes. Evolution (N. Y). 71:1297–

1312.

Hernandez, L. P., D. Adriaens, C. H. Martin, P. C. Wainwright, B. Masschaele, and M. Dierick.

2018. Building trophic specializations that result in substantial niche partitioning within a

young adaptive radiation. J. Anat. 232:173–185.

Hernandez, P. L., A. C. Gibb, and L. Ferry-Graham. 2009. Trophic apparatus in

cyprinodontiform fishes: Functional specializations for picking and scraping behaviors. J.

Morphol. 270:645–661.

Higham, T. E., S. M. Rogers, R. B. Langerhans, H. A. Jamniczky, G. V Lauder, W. J. Stewart,

C. H. Martin, and D. N. Reznick. 2016. Speciation through the lens of biomechanics:

locomotion, prey capture and reproductive isolation. Proc. R. Soc. B Biol. Sci.

283:20161294.

Holtmeier, C. L. 2001. Heterochrony, maternal effects, and phenotypic variation among

sympatric pupfishes. Evolution 55:330–8.

Hu, Y., N. Nelson-Maney, and P. S. L. Anderson. 2017. Common evolutionary trends underlie

the four-bar linkage systems of sunfish and mantis shrimp. Evolution (N. Y). 71:1397–

1405.

Janovetz, J. 2005. Functional morphology of feeding in the scale-eating specialist *Catoprion*

mento. J. Exp. Biol. 208:4757–68.

Kolmann, M. A., J. M. Huie, K. Evans, and A. P. Summers. 2018. Specialized specialists and the

narrow niche fallacy□: a tale of scale-feeding fishes. R. Soc. Open Sci. 5.1:171581.

Kuhn, M. 2008. Building predictive models in R using the caret package. J. Stat. Softw. 1–26.

Lencer, E. S., W. C. Warren, R. Harrison, and A. R. McCune. 2017. The Cyprinodon variegatus

genome reveals gene expression changes underlying differences in skull morphology among

closely related species. BMC Genomics 18:424.

Longo, S. J., T. Goodearly, and P. C. Wainwright. 2018. Extremely fast feeding strikes are

powered by elastic recoil in a seahorse relative, the snipefish, Macroramphosus scolopax. P.

20181078 *in* Proceedings of the Royal Society B: Biological Sciences. The Royal Society.

Martin, C. H. 2016. The cryptic origins of evolutionary novelty: 1000-fold faster trophic

diversification rates without increased ecological opportunity or hybrid swarm. Evolution

(N. Y). 70:2504–2519.

Martin, C. H., and L. C. Feinstein. 2014. Novel trophic niches drive variable progress towards

ecological speciation within an adaptive radiation of pupfishes. Mol. Ecol. 23:1846–1862.

Martin, C. H., and P. C. Wainwright. 2013a. A Remarkable Species Flock of Cyprinodon

Pupfishes Endemic to San Salvador Island, Bahamas. Bull. Peabody Museum Nat. Hist.

54:231–241.

Martin, C. H., and P. C. Wainwright. 2013b. Multiple fitness peaks on the adaptive landscape

drive adaptive radiation in the wild. Science 339:208–11.

Martin, C. H., and P. C. Wainwright. 2011. Trophic novelty is linked to exceptional rates of

morphological diversification in two adaptive radiations of cyprinodon pupfish. Evolution

(N. Y). 65:2197–2212.

Matthews, D. G., and R. C. Albertson. 2017. Effect of craniofacial genotype on the relationship

between morphology and feeding performance in cichlid fishes. Evolution (N. Y). 71:2050–

2061.

McGee, M. D., D. Schluter, and P. C. Wainwright. 2013. Functional basis of ecological

divergence in sympatric stickleback. BMC Evol. Biol. 13:277. BioMed Central.

McGhee, K. E., R. C. Fuller, and J. Travis. 2007. Male competition and female choice interact to

determine mating success in the bluefin killifish. Behav. Ecol. 18:822–830.

O'Neill, M. W., and A. C. Gibb. 2013. Does Feeding Behavior Facilitate Trophic Niche

Partitioning in Two Sympatric Sucker Species from the American Southwest? Physiol.

Biochem. Zool. 87:65–76.

Patek, S. N., J. E. Baio, B. L. Fisher, and A. V Suarez. 2006. Multifunctionality and mechanical

origins: Ballistic jaw propulsion in trap-jaw ants.

R Core Team. 2018. R: A Language and Environment for Statistical Computing. R Found. Stat.

Comput. Vienna, Austria. URL https//www.R-project.org/. 0:{ISBN} 3-900051-07-0.

Sazima, I., D. De Zoologia, U. E. De Campinas, and S. Paulo. 1983. Scale-eating in characoids

and other fishes. 9.

Stayton, C. T. 2019. Performance in three shell functions predicts the phenotypic distribution of

hard-shelled turtles. Evolution (N. Y). 73:720–734. John Wiley & Sons, Ltd (10.1111).

Svanbäck, R., P. C. Wainwright, and L. A. Ferry-Graham. 2002. Linking Cranial Kinematics,

Buccal Pressure, and Suction Feeding Performance in Largemouth Bass. Physiol. Biochem.

Zool. 75:532–543.

Takeuchi, Y., M. Hori, and Y. Oda. 2012. Lateralized kinematics of predation behavior in a Lake

Tanganyika scale-eating cichlid fish. PLoS One 7:e29272. Public Library of Science.

Tebbich, S., K. Sterelny, and I. Teschke. 2010. The tale of the finch: adaptive radiation and

behavioural flexibility. Philos. Trans. R. Soc. Lond. B. Biol. Sci. 365:1099–109. The Royal

Society.

Tobler, M., J. L. Kelley, M. Plath, and R. Riesch. 2018. Extreme environments and the origins of

biodiversity: Adaptation and speciation in sulphide spring fishes.

Tobler, M., R. Riesch, and M. Plath. 2015. Extremophile fishes: An integrative synthesis. Pp.

279–296 *in* Extremophile Fishes: Ecology, Evolution, and Physiology of Teleosts in

Extreme Environments.

Van Wassenbergh, S., T. Lieben, A. Herrel, F. Huysentruyt, T. Geerinckx, D. Adriaens, and P.

Aerts. 2008. Kinematics of benthic suction feeding in Callichthyidae and Mochokidae, with

functional implications for the evolution of food scraping in catfishes. J. Exp. Biol.

212:116–125.

Whitford, M. D., R. W. Clark, G. A. Freymiller, and T. E. Higham. 2019. Determinants of

predation success□: How to survive an attack from a rattlesnake. Funct. Ecol. 1–11. John

Wiley & Sons, Ltd (10.1111).

Wood, S. N. 2011. Fast stable restricted maximum likelihood and marginal likelihood estimation

of semiparametric generalized linear models. J. R. Stat. Soc. Ser. B Stat. Methodol. 73:3–

36.

Table 1. Results of linear mixed models investigating if strike kinematic variables vary between 1) species (generalists, snail-eaters, scale-eaters, or hybrids), 2) food item (shrimp or scales), or 3) the interaction between the two. Significant predictors are indicted in bold.

Response	Predictors	χ^2	df	P
Peak Protrusion (mm)				
	Species	4.01	3	0.26
	Food Item	1.10	1	0.29
	log(Body Length)	3.01	1	0.082
	Species:Food Item	2.03	3	0.57
Time to Peak Protrusion (s)				
	Species	3.80	3	0.27
	Food Item	0.73	1	0.39
	log(Body Length)	1.02	1	0.31
	Species:Food Item	4.03	3	0.26
Peak Gape (mm)				
	Species	**23.13**	**3**	**3.8×10^{-5}**
	Food Item	0.71	1	0.40
	log(Body Length)	1.24	1	0.27
	Species:Food Item	0.65	3	0.88
Time to Peak Gape (s)				
	Species	2.43	3	0.49
	Food Item	0.57	1	0.45
	log(Body Length)	2.80	1	0.17
	Species:Food Item	1.87	3	0.60
Gape Angle (degrees)				
	Species	3.28	3	0.35
	Food Item	0.032	1	0.86
	log(Body Length)	1.01	1	0.32
	Species:Food Item	3.43	3	0.33
Lower Jaw Angle (degrees)				
	Species	**18.62**	**3**	**0.00033**
	Food Item	0.0031	1	0.96
	log(Body Length)	3.53	1	0.060
	Species:Food Item	3.56	3	0.31
Time to Impact (s)				
	Species	2.55	3	0.47
	Food Item	2.05	1	0.15
	log(Body Length)	1.40	1	0.24
	Species:Food Item	4.69	3	0.20
Time from Peak Gape to Impact (s)				
	Species	2.44	3	0.48
	Food Item	0.97	1	0.32
	log(Body Length)	0.57	1	0.45

	Species:Food Item	1.39	3	0.71
Starting Distance from prey (mm)				
	Species	0.43	3	0.93
	Food Item	1.99	1	0.16
	log(Body Length)	2.77	1	0.10
	Species:Food Item	0.80	3	0.85
Ram speed (m/s)				
	Species	3.25	3	0.35
	Food Item	3.75	1	0.053
	log(Body Length)	1.55	1	0.21
	Species:Food Item	2.02	3	0.57

610 **Table 2.** Results of a linear discriminant analysis for kinematic variables for strikes on
611 shrimp and scales.
612

Kinematic Metrics	LD1	LD2	LD3
Peak Protrusion (mm)	-0.082	-0.49	-0.065
Peak Gape (mm)	1.55	0.39	-0.56
Time to Peak Gape (s)	-8.00	12.16	10.24
Gape Angle (degrees)	-0.032	-0.012	-0.033
Lower Jaw Angle (degrees)	0.069	0.0029	0.022
Time from Peak Gape to Impact (s)	-9.85	31.32	-33.03
Ram speed (m/s)	-7.98	17.27	10.67
Proportion of Trace	0.92	0.056	0.028

614   **Table 3.** Results of GAM model comparisons using AIC score. The best-fitting model is
615   indicated in bold and ΔAIC is indicated relative to this model.

		Model	ΔAIC
A	1	Area~s(Peak Gape, Peak Protrusion, bs="ts")+(Lower Jaw Angle)	23.11
	2	Area~s(Peak Gape, Peak Protrusion, bs="ts")+(Gape Angle)	25.09
	3	Area~s(Peak Gape, Gape Angle, bs="ts")+(Lower Jaw Angle)	30.76
	4	Area~s(Peak Gape, Gape Angle, bs="ts")+(Peak Protrusion)	30.55
	5	Area~s(Peak Gape, Lower Jaw Angle, bs="ts")+(Peak Protrusion)	30.27
	6	Area~s(Peak Gape, Lower Jaw Angle, bs="ts")+(Gape Angle)	30.7
	7	Area~s(Peak Protrusion, Lower Jaw Angle, bs="ts")+(Peak Gape)	28.79
	8	Area~s(Peak Protrusion, Lower Jaw Angle, bs="ts")+(Gape Angle)	31.36
	9	Area~s(Peak Protrusion, Gape Angle, bs="ts")+(Peak Gape)	30.64
	10	Area~s(Peak Protrusion, Gape Angle, bs="ts")+(Lower Jaw Angle)	34.44
	11	Area~s(Gape Angle,Lower Jaw Angle, bs="ts")+(Peak Gape)	28.79
	12	Area~s(Gape Angle,Lower Jaw Angle, bs="ts")+(Peak Protrusion)	31.96
B	13	Area~s(Peak Gape)+ s(Peak Protrusion)+(Lower Jaw Angle)	9.99
	14	Area~s(Peak Gape)+ s(Peak Protrusion)+(Gape Angle)	15.41
	15	Area~s(Peak Gape)+ s(Gape Angle)+(Lower Jaw Angle)	9.72
	16	**Area~s(Peak Gape)+ s(Gape Angle)+(Peak Protrusion)**	**-**
	17	Area~s(Peak Gape)+ s(Lower Jaw Angle)+(Peak Protrusion)	9.99
	18	Area~s(Peak Gape)+ s(Lower Jaw Angle)+(Gape Angle)	11.05
	19	Area~s(Peak Protrusion)+s(Lower Jaw Angle)+(Peak Gape)	30.45
	20	Area~s(Peak Protrusion)+s(Lower Jaw Angle)+(Gape Angle)	34.9
	21	Area~s(Peak Protrusion)+s(Gape Angle)+(Peak Gape)	29.32
	22	Area~s(Peak Protrusion)+s(Gape Angle)+(Lower Jaw Angle)	33.42
	23	Area~s(Gape Angle)+s(Lower Jaw Angle)+(Peak Gape)	30.66
	24	Area~s(Gape Angle)+s(Lower Jaw Angle)+(Peak Protrusion)	33.75
C	25	Area~s(Peak Gape)+Peak Protrusion+Lower Jaw Angle+Gape Angle	11.86
	26	Area~s(Peak Protrusion)+Peak Gape+Lower Jaw Angle+Gape Angle	32.4
	27	Area~s(Lower Jaw Angle)+Peak Protrusion+Peak Gape+Gape Angle	32.76
	28	Area~s(Gape Angle)+Peak Protrusion+Peak Gape+Lower Jaw Angle	32.02
D	29	Area~Peak Gape+Peak Protrusion+Lower Jaw Angle+Gape Angle	32.76

Figure 1. Divergent feeding
kinematics in scale-eaters. A)
Biplot of discriminant axes 1 (LD1)
and 2 (LD2) describing overall
kinematic differences among pupfish
species (generalists, snail-eaters,
scale-eaters, or F1 hybrids). Ellipses
represent 95% CIs. B) Mean peak
gape (mm) for each species with ±
95% CIs calculated via
bootstrapping (10,000 iterations). C)
Mean lower jaw angle at peak gape
(mm) for each species with ± 95%
CIs calculated via bootstrapping
(10,000 iterations).
633

634   **Figure 2.**

635
636
Figure 2. The large jaws of scale-eating pupfish allow them to double their gape size and
increase the angle between their lower jaw and suspensorium (lower jaw angle) while
maintaining the same gape angle as other species during feeding strikes. A) Hypothetical
measurements of a generalist's protrusion distance, peak gape, and lower jaw angle if they strike
a food item with an 80° gape angle. B) Hypothetical measurements of a scale-eater's protrusion
distance, peak gape, and lower jaw angle if they strike a food item with the same 80° gape angle.

Figure 3.

Figure 3. The interaction of peak gape and peak protrusion may result in a performance
optimum for scale-biting. Visualization of our GAM model investigating how the thin-plate
univariate splines of A) peak gape (mm), B) gape angle (degrees), and C) the fixed effect of peak
protrusion (mm) is associated with bite size (surface area removed per strike from a Repashy
gelatin cube), and D) representative scale-eating bites taken out of gelatin cubes.
650

    Figure 4.

Figure 4. Scale-eaters have larger predicted bite sizes compared to other species. Predicted bite sizes for all strikes from each species using machine-learning optimization of GAM models. Grey points represent predicted bite sizes for individuals, color points represent means, and bars represent ± 95% CIs calculated via bootstrapping (10,000 iterations).