A stress-induced block in dicarboxylate uptake and utilization in Salmonella

Steven J. Hersch[†], Bojana Radan, Bushra Ilyas, Patrick Lavoie, and William Wiley Navarre*

Department of Molecular Genetics, University of Toronto, Toronto, ON, Canada

Running Title: *Salmonella* represses its growth in dicarboxylates

[†]Current address: Department of Ecosystem and Public Health, University of Calgary, Calgary,

Canada

*Address correspondence to William W. Navarre, william.navarre@utoronto.ca

Abstract

Bacteria have evolved to sense and respond to their environment by altering gene

expression and metabolism to promote growth and survival. In this work we demonstrate that

Salmonella displays an extensive (>30 hour) lag in growth when subcultured into media where

dicarboxylates such as succinate are the sole carbon source. This growth lag is regulated in part

by RpoS, the RssB anti-adaptor IraP, translation elongation factor P, and to a lesser degree the

stringent response. We also show that small amounts of proline or citrate can trigger early

growth in succinate media and that, at least for proline, this effect requires the multifunctional

enzyme/regulator PutA. We demonstrate that activation of RpoS results in the repression of *dctA*,

encoding the primary dicarboxylate importer, and that constitutive expression of *dctA* induced

growth. This dicarboxylate growth lag phenotype is far more severe across multiple *Salmonella*

isolates than in its close relative *E. coli*. Replacing 200 nt of the *Salmonella dctA* promoter

region with that of *E. coli* was sufficient to eliminate the observed lag in growth. We hypothesize

that this *cis*-regulatory divergence might be an adaptation to *Salmonella*'s virulent lifestyle

where levels of phagocyte-produced succinate increase in response to bacterial LPS. We found

that impairing *dctA* repression had no effect on *Salmonella*'s survival in acidified succinate or in

macrophage but propose alternate hypotheses of fitness advantages acquired by repressing

dicarboxylate uptake.

Importance

Bacteria have evolved to sense and respond to their environment to maximize their

chance of survival. By studying differences in the responses of pathogenic bacteria and closely

related non-pathogens, we can gain insight into what environments they encounter inside of an

infected host. Here we demonstrate that *Salmonella* diverges from its close relative *E. coli* in its

response to dicarboxylates such as the metabolite succinate. We show that this is regulated by

stress response proteins and ultimately can be attributed to *Salmonella* repressing its import of

dicarboxylates. Understanding this phenomenon may reveal a novel aspect of the *Salmonella*

virulence cycle, and our characterization of its regulation yields a number of mutant strains that

can be used to further study it.

Introduction

Bacteria must adapt to changing environmental conditions by sensing their surroundings

and integrating signals to initiate rapid growth in nutrient rich situations or instigate defence

mechanisms and metabolic hibernation in response to stress (1). Pathogens like *Salmonella* have

adapted to survive and replicate within their particular host niche, which is reflected in a variety

of subtle differences in regulation compared to their non-pathogenic relatives. Understanding the

differences in related pathogenic and commensal bacteria *E. coli* and *Salmonella*, has yielded a

wealth of insight regarding the challenges pathogens face in the host environment. These

bacteria are closely related, yet *Salmonella* has acquired a number of adaptations that

accommodate its virulent lifestyle; for example allowing *Salmonella* to invade tissues and

survive within host cells such as macrophage, which is important for *Salmonella* virulence (2–6).

Metabolic modulation is an important component in the adaptation to multiple types of

stress. The bacterial stringent response, wherein the second messenger molecule, guanosine 5'-

disphosphate 3'-diphosphate (ppGpp) is produced by RelA or SpoT in response to amino acid

starvation or other cellular stress cues (7–10). Bacteria can also alter gene expression using the

general stress response sigma factor RpoS (σ^S), which has been linked to virulence in a number

of pathogenic bacteria by contributing to virulence gene expression and survival within an

infected host (11–13). RpoS can be activated in response to a variety of conditions including

starvation, hyper-osmolarity and oxidative stress, and can be regulated at all levels of synthesis

from transcription to protein degradation where it is recognized by the adaptor RssB (also known

as MviA, SprE, or ExpM) and chaperoned to the ClpXP protease (14–18) . In response to

specific stresses, the anti-adaptors IraP, RssC (IraM in *E. coli*) and IraD can be induced, which

impair RssB and thereby rapidly stabilize RpoS (19–21). Strains with reduced *rpoS* activity have

been demonstrated to grow faster than wild-type *E*. coli when grown using the dicarboxylate

succinate as a sole carbon source (22–24). Furthermore, aerobic growth using succinate relies on

the dicarboxylate importer, DctA, which is known to be regulated by the DcuSR two-component

system in response to dicarboxylates, as well as by DctR (YhiF) in *E. coli* strains lacking ATP

Synthase activity (25–27).

Our previous studies of *Salmonella* translation Elongation Factor P (EF-P), which

facilitates the ribosomal synthesis of difficult-to-translate polyproline/glycine motifs, revealed

that strains lacking functional EF-P show rapid growth using succinate as a carbon source (28–

31). Here we demonstrate that activation of RpoS, via the antiadaptor IraP or via the stringent

response, triggers wild-type *Salmonella* to shut down import of dicarboxylates like succinate,

fumarate, and malate. For reasons that remain unclear, Salmonella resumes import and growth

on succinate media after a stereotypical lag of approximately 30 hours. The severity of this lag

differs between the vast majority of *Salmonella* and *E. coli* strains, and this difference appears to

be due almost entirely to differences within the promoter of the dicarboxylate importer DctA.

Furthermore, we find that trace amounts of proline or citrate can "license" early *Salmonella*

growth on dicarboxylates. In the case of proline, this phenotype depends on the unsual

polyfunctional enzyme/regulator PutA. This work adds to a larger body of evidence that proline

is an essential signaling metabolite that modulates key aspects of *Salmonella* growth within the

animal host.

Results

***Salmonella* delays its growth using dicarboxylic acids as a sole carbon source**

Our previous work investigating the role of EF-P in *Salmonella* demonstrated that strains

deficient in EF-P activity display a 'hyper-active' metabolism relative to wild-type when grown

under specific nutrient limited conditions (28, 29). Biolog phenotype microarrays (31) revealed

that wild-type *Salmonella enterica* Sv. Typhimurim strain 14028s failed to grow under several

varied nutrient conditions whereas isogenic strains lacking functional EF-P grew rapidly.

Through a series of further growth assays we determined that the underlying cause of this growth

defect was due entirely to the fact that many of the varied growth conditions in the Biolog assay

employ a base media where succinate provides the sole carbon source.

Earlier literature reported that many wild-type strains of *E. coli* display an extended lag

when subcultured on succinate media and that this growth delay was dependent on the sigma

factor RpoS (22–24). Accordingly, we tested the growth of strain 14028s in minimal media

containing succinate as the sole carbon source. We found that the *efp* (EF-P) mutant, as well as

an *rpoS* mutant, were able to grow readily in succinate with minimal lag (Figure 1). In contrast,

wild-type *Salmonella* exhibited an extended lag phase where it failed to grow for over 30 hours

before initiating exponential (log) growth. The dicarboxylic acid transporter, DctA, was also

required for growth and an isogenic *dctA* deletion strain showed no sign of growth by 48 hours.

We note that the wild-type log growth rate, when it finally resumed, was similar to the

log growth rate of the *rpoS* mutant, suggesting that the growth delay was due to an extended lag

phase upon subculturing into succinate media and not due to slow growth on succinate *per se*.

The length of this extended lag phase was remarkably consistent, stereotypically between 30-32

hours across dozens of experiments. Furthermore, colonies of *Salmonella* that grew on succinate

media after 30 hours, when recultured, continued to display an extended lag in growth.

Occasional colonies that did show rapid growth on succinate were found to have acquired

spontaneous mutations in RpoS. Together these results indicate that the bacteria that grew on

succinate after 30 hours did not arise from mutation, but rather from a genetically programmed

physiological response.

DctA can also import other dicarboxylic acids including malate and fumarate. We

therefore tested whether the extended lag of wild-type *Salmonella* would also occur in media

with fumarate and malate as the sole carbon source (Figure 1C). We found that, similar to

succinate, wild-type strain 14028s displayed a similarly extended lag in growth using either

malate or fumarate, while the *efp* mutant grew significantly earlier. This suggests that the growth

repression instigated by wild-type *Salmonella* is not specific to succinate but also occurs when

subcultured into media using other dicarboxylic acids as the sole carbon source.

These data also indicate that the failure of strain 14028s to grow rapidly on succinate was

not due to an inherent inability to utilize this carbon source (i.e. the strain does not lack the

factors necessary to import or catabolize succinate), but rather that delayed growth was the result

of a regulatory block that prevented the expression of the necessary succinate utilization systems.

This regulation involves RpoS, perhaps through a protein(s) that requires *efp* for its efficient

translation. We believe this phenotype may have been overlooked in prior studies on *Salmonella*

metabolism as many of these studies employed strain LT2, most isolates of which harbor a

hypomorphic lesion in the gene encoding RpoS (32).

Many *Salmonella* but few *E. coli* strains delay growth using dicarboxylates

Prior literature suggested the severity of the growth lag phenotype was much less severe

in *E. coli* than what we observed in strain 14028s. To determine if this is indeed the case, or if it

was merely a difference in growth conditions, the growth of *Salmonella* 14028s was compared to

E. coli K12 (strain BW25113)*.* This revealed that, while *E. coli* did display a lag in growth – this

lag was much shorter than what was observed for *Salmonella* (Figure 2A).

To determine whether the length of the dicarboxylate lag was a characteristic broadly

shared among each these two species, or limited to specific strains of *E. coli* and *Salmonella*,

growth on succinate media was assessed for all 105 non-typhoidal strains in the *Salmonella*

Genetic Stock Centre (SGSC) collection, as well as all 72 strains of the *E. coli* Reference

(ECOR) collection. Though there are some exceptions, the lag displayed by the vast majority of

E. coli strains on succinate media was considerably shorter compared to most *Salmonella* strains

(Figure 2B-D). Once logarithmic growth was initiated, *Salmonella* also appeared to trend

towards a slightly longer doubling time than the majority of *E. coli* strains (Figure 2C and E).

Hypomorphic variants of *rpoS* accumulate readily in *E. coli* and *Salmonella* during

laboratory passage, and there remained the possibility that many strains in the ECOR or SGSC

collections acquired such mutations during their cultivation prior to being stored/archived. To

ensure that the observed effects were not due to variations in RpoS activity, each strain was also

screened for catalase activity as a surrogate indicator of having a functional RpoS. Regardless of

catalase activity, the trend was maintained that *E. coli* strains in general showed a shorter lag

phase than *Salmonella* when using succinate as the sole carbon source (Figure S1).

IraP contributes to growth repression in dicarboxylate media

RpoS is activated under specific conditions of stress or nutrient limitation and its

expression, synthesis, stability, and activity are regulated at multiple levels. Under ideal growth

conditions RpoS protein is destablized through its interaction with RssB, an 'adaptor' protein

that promotes RpoS degradation via the ClpXP proteases. In response to specific stressors or

nutrient limitations, one or more anti-adaptor proteins, IraP, RssC and IraD, can antagonize

binding by RssB to alleviate RpoS degradation (14, 19, 20, 33). Systematic deletion of these

anti-adaptors revealed that IraP is a primary upstream factor invoved in the dicarboxylate lag,

and the rapid growth phenotype of an *iraP* mutant strain could be partially complemented by

expressing IraP from its native promoter on a plasmid (Figure 3). In contrast, deletion of the

genes encoding the anti-adaptors, RssC and IraD, had comparatively minor effects. These

findings demonstrate that, under the laboratory growth conditions we employ, IraP-mediated

stabilization of RpoS plays a role in repressing *Salmonella*'s growth using succinate as the sole

carbon source.

Proline and citrate override the lag observed in succinate media

During these studies it was observed that supplementation of our minimal media with

small amounts of LB media (a.k.a. 'lysogeny broth' or 'Luria broth') would shorten or even

eliminate the observed lag phase on succinate (Figure S2). Given that the primary nutrient

components in LB are peptides and amino acids, the effects of each amino and and a few other

carbon sources in the TCA cycle were examined.

Unlike all other amino acids we found that the addition of 0.005% proline to succinate

media could 'override' the extended growth lag to enable rapid growth on succinate. The cells

did not appear to simply consume the proline as a carbon source since no other individual amino

acid showed similar growth induction (Figure S2D). Furthermore, although proline can be used

by *Salmonella* as a carbon/nitrogen source, there was no discernible growth on 0.01% proline in

the absence of succinate, suggesting the levels we were providing were too small (Figure S2E).

A similar early-growth phenomenon could be invoked through the addition of small amounts of

citrate. Growth on succinate in the presence of either proline or citrate occurred in a manner

resembling a diauxy wherein growth would cease again following exhaustion of the proline or

citrate (Figure 4A and B).

We considered the possibility that proline or citrate are limiting nutrients during growth

on dicarboxylates (e.g. proline synthesis is shut down and the cell becomes auxotrophic),

however several additional observations suggest that this is not the case. We note, for example,

that growth is temporarilty restored by either citrate or proline, and neither metabolite is known

to be an intermediate in the synthesis of the other. Therefore, these two compounds are likely

affecting growth through similar but independent routes.

Another observation relates to the effect of proline or citrate on the dicarboxylate growth

lag in a *Salmonella relA/spoT* double mutant that is unable to produce the secondary messenger

ppGpp to trigger the stringent response. We find that the *Salmonella* stringent mutant displays a

growth lag similar to wild-type when subcultured into succinate media from an overnight culture

grown in LB or M9 media. Whereas wild-type *Salmonella* rapidly shuts down growth upon

exhaustion of available proline or citrate, the stringent mutant did not (Figure 4C and D) – that is

the strain continued to grow on succinate.

This suggests that exhaustion of proline or citrate are not, in and of themselves, limiting

during growth in succinate media, but rather that proline and citrate might act as external cues

that allow growth on dicarboxylates to start. In wild-type *Salmonella* strains, growth continues

until these stimulating compounds are exhausted; at which point the production of ppGpp via the

stringent response stimulates the shutdown of dicarboxylate utilization, possibly through its

effects on RpoS. In the absence of ppGpp or RpoS, however, the exhaustion of available

proline/citrate is not sensed by the cell, and growth will continue until the dicarboxylate (or

another essential nutrient) are exhausted.

Proline-mediated stimulation in growth requires the multifunctional proline utilization

enzyme PutA.

Proline has been shown to play a key, but unclear, role in *Salmonella*'s adapation to life

in an animal host. For both *E. coli* and *Salmonella*, proline can act as a compatible solute to

regulate osmotic balance in the cells (34, 35). However, the key *Salmonella*-specific virulence

factors encoded by *mgtA* and *mgtCBR* are controlled by short proline-rich peptides (*mgtL* and

mgtP, respectively) that attenuate their operons in response to increased levels of uncharged

tRNA$_{Pro}$ (36). Also, loss of EF-P, which is critical to ensure the smooth translation of proteins

rich in proline, causes translational stalling that likely mimics the effects of proline-limitation

(30, 37, 38). Notably, *Salmonella* strains lacking EF-P are poorly virulent in animal models and

appear to have difficulty adapting to an intraceulluar lifestyle; e.g. failing to produce

stereotypical "sifs" when infecting host cells (29).

To explore the factors that might link proline availability to the rapid utilization of

dicarboxylates as carbon sources, the roles of several proline-dependent regulatory systems were

examined. While neither *mgtA* nor *mgtCBR* had an effect on proline's effect on the

dicarboxylate growth lag (data not shown), we found that strains lacking PutA, an unusual

polyfunctional proline utilization enzyme, were unresponsive to proline with respect to growth

on succinate (35, 39, 40).

PutA is a polyfunctional membrane-bound enzyme that contains the two enzymatic

activities required to oxidatively convert proline to glutamate: proline dehydrongease (PRODH),

which oxidizes proline to P5C; and P5C dehydrogenase (P5CDH) which is critical to convert

P5C to glutamate. During the course of this conversion, PutA transfers resulting electrons to

ubiquinone (via PRODH), and NAD+ (via P5CDH). PutA also contains a DNA binding domain

that controls, via repression, its own expression (*putA*) as well as that of the divergently

transcribed *putP* gene, encoding the proline importer PutP. An implication of this finding is that

proline, *per se*, may not be the proximal metabolite that regulates the utilization of

dicarboxylates, but instead may be due to PutA-mediated reduction of the quinone or NAD pool,

the availability of gluatamine/glutamate (although neither glutamine nor glutamate had the same

effect as proline) or PutA-mediated gene regulation via the PutA DNA-binding domain.

Repression of dicarboxylate import accounts for growth lag

Activation of RpoS not only upregulates several systems directly involved in mitigating

cellular stress (e.g. catalase, import and synthesis of compatible solutes, the production of

glycogen, etc.), but it has also been shown to down regulate the expression of genes encoding

enzymes in the TCA cycle (41–43). An analysis of *dctA* gene expression, encoding the primary

dicarboxylate transporter, on SalCom (44, 45) showed that it generally anti-correlated with the

expression of RpoS-activated genes like OsmY, where *dctA* was downregulated in response to

osmotic shock and in late stationary phase. We hypothesized that activation of the RpoS-regulon

may repress the expression of the *dctA* gene and thereby restrict *Salmonella* from taking up

dicarboxylates such as succinate for consumption in response to stresses encountered within the

host. To test if low *dctA* expression accounts for the *Salmonella* extended lag on dicarboxylates,

we constitutively expressed *dctA* from a plasmid. Indeed, constitutive expression of *dctA* (but not

of a similar *lacZ* control plasmid) eliminated the *Salmonella* lag in succinate media (Figure 5A

and B).

The *E. coli dctA* promoter is sufficient to induce *Salmonella* growth using dicarboxylates

Given the differences between *Salmonella* and *E. coli*, with respect to growth on

succinate, we compared the nucleotide sequences upstream of *dctA* in these bacteria (Figure S3).

The two regions are largely similar (78% identity overall across 500bp upstream), especially near

the core promoter. However, there are notable regions of significant sequence difference that are

specific and conserved to each species. A comparison of the *dctA* promoter of *S. bongori* and of

the various *S. enterica* subspecies (enterica, houtenae, arizonae, diarizonae, salmae) and

typhoidal isolates reveal that they are highly conserved to one another - but distinct from the

dctA promoters found in pathogenic and commensal *E. coli* strains. This suggests that changes

to the regulation of *dctA* may have occurred as, or shortly after, the two species diverged.

To examine whether the species-specific differences in the *dctA* promoter (P$_{dctA}$) played a

role in the differences in how these species respond to succinate, the *E. coli* P$_{dctA}$ was engineered

into the *Salmonella* chromosome using an upstream chloramphenicol resistance cassette to select

for successful recombination. Replacing the 500bp upstream of the *Salmonella dctA* start codon

with those of *E. coli* was sufficient to abolish *Salmonella*'s ability to repress its uptake of

dicarboxylic acids and this strain grew readily in succinate media (Figure 5C and D). As a

control, using the same method to insert *Salmonella*'s native *dctA* promoter yielded no difference

from wild-type *Salmonella*. Further reducing the swapped region to 200bp maintained the

growth phenotype, but swapping only 54bp upstream of the *Salmonella* with that of *E. coli*

(constituting the 5' untranslated region) resulted in only a slight restoration of growth.

It is possible that *Salmonella* contains a factor to repress *dctA* expression that is not

present in *E. coli*, so we generated transcriptional fusion plasmids and tested the two *dctA*

promoters when expressed in *E. coli*. We found that the *dctA* promoter from *Salmonella* was

expressed to a significantly lower degree than that of *E. coli* (Figure S4), suggesting that it

contains a distinctive region that is recognized by a common regulator that is also present in *E.*

coli.

Restricting dicarboxylate import does not influence survival in macrophage cell lines

To probe the question of why *Salmonella* may have acquired the trait of blocking

dicarboxylate utilization and what evolutionary advantage it may gain by it, we considered that

succinate levels increase significantly in activated macrophage, an environment that *Salmonella*

(but not *E. coli*) has adapted to survive in effectively (5, 6, 46). In the *Salmonella*-containing

vacuole, the pH reaches approximately 5.0 and the lower estimates reach pH 4.4, which is

comparable to the acid dissociation constants of succinate ($pK_{a1,2}$ = 4.2, 5.6) (47, 48). This

suggests that in the acidified phagosome, succinate may become protonated and potentially act as

a proton shuttle to acidify the bacterial cytoplasm. The ability of *Salmonella* to restrict its uptake

of succinate could therefore provide a possible survival advantage in this environment.

Constitutive overexpression of *dctA* but not *lacZ* led to decreased survival in both

acidifed succinate media and in the human monocyte THP-1 cell line (Figure S5). However,

overexpression of *dctA* has been demonstrated to be toxic to *E. coli* and we found that survival in

acidified succinate was just as low for a *dctA* point mutant (N301A) that is defective for

succinate transport (49). This implicated that the reduced survival was not due to dicarboxylate

uptake but rather was an artifact of *dctA* overexpression to toxic levels. To bypass this artifact,

we tested the *Salmonella* strain containing the chromosomal *dctA* promoter from *E. coli,* which

grows readily in dicarboxylate media yet does not constitutively overexpress *dctA* from a

plasmid and so does not exhibit the associated toxic effects. Using this strain we found no

decrease in survival relative to wild-type *Salmonella* in acidifed succinate media or in human

(THP-1) or mouse (J774) macrophage cell lines (Figure 6). Deletion of *dctA* or *iraP* genes also

did not appear to significantly influence *Salmonella* survival in THP-1 macrophage in these

short-term infection assays.

Discussion

In this work we explore a previously uncharacterized *Salmonella* phenotype whereby it

restricts its ability to import dicarboxylates in an RpoS and Elongation Factor P dependent

manner. Stationary phase cells subcultured into media with succinate, fumarate, or malate as the

sole carbon source fail to grow for >30 hours before resuming normal growth. The cause of this

extended lag phase appears to largely reflect low expression of the *dctA* gene encoding a major

dicarboxylate importer. The 'cue' that induces cells to suddenly begin rapid growth on

dicarboxylates after 30 hours remains unclear. Interestingly, an almost immediate exit from the

extended lag phase could be triggered by the addition of low amounts of proline or citrate. Upon

exhaustion of the available proline or citrate, growth would again cease in a process that required

the synthesis of ppGpp.

311   Our model suggests that activation of RpoS leads to a rapid shutdown in the transporter

that mediates the uptake and utilization of dicarboxylates (Figure 7). Since the stringent

response and ppGpp can impact the expression of *rpoS* and RssB anti-adaptors including *iraP*,

the stringent response is likely the means by which IraP- and RpoS-mediate the shutdown of

growth (14, 50–52). We also note the many curious links that have been found between

regulation of genes essential for *Salmonella* host infection and the availability of proline, notably

the attenuation peptides MgtL and MgtP and the essentiality of EF-P to alleviate translational

stalling at proline rich motifs. Our finding that the poly-functional proline metabolic enzyme,

PutA, is essential for the proline-mediated induction of growth on succinate, leaves open the

question of whether the inducing cue is due to changes in the abilty of PutA to bind DNA

(perhaps PutA directly represses *dctA* in Salmonella), alterations in cellular energetics via the

production of either reduced quinones or NADH, or through increases in the concentrations of

cellular glutamate (although we note addition of glutamate does not have the same effect as

proline).

325   The difference in the response of *Salmonella* and *E. coli* to dicarboxylic acids may offer

important clues to identifying the evolutionary advantage conveyed by this adaptation. The

divergent evolutionary paths between *Escherichia* and *Salmonella*, that began over 100MYr ago,

were fueled both by the acquisition of novel functions via horizontal gene transfer and extensive

regulatory re-wiring, often through changes in cis-regulatory elements (53–56). The differences

in the severity of their dicarboxylate lag, as evidenced by sampling a range of isolates in the

SGSC and ECOR collections, appears to be due to extensive differences in the 200 nucleotide

region upstream of the *dctA* dicarboxylate import gene including regions that lie upstream of the

previously mapped 5' UTR. While it remains possible that this factor could involve a small

RNA, our finding that swapping just the 5' untranslated region of *dctA* (P_{dctA} 54) was insufficient

to reverse growth repression suggests a protein factor acting on the promoter at the

transcriptional level. RpoS likely impacts *dctA* expression via an intermediate and yet

undetermined transcription factor.

Since many of the traits that *Salmonella* has acquired since their divergence are related to

its pathogenic lifestyle it follows that this phenotype may reflect a situation that *Salmonella*

encounters during infection of a host. Interestingly, it has recently been shown that *Salmonella*

utilizes microbiota-derived succinate in the lumen of the inflamed gut (57). This suggests that the

uptake of succinate in the gut employs anaerobic dicarboxylate transporters rather than DctA, or

that an inducing compound such as proline or citrate is present in this environment and can

stimulate succinate uptake. The recent finding that succinate accumulates to high levels in

activated macrophage suggests that *Salmonella*'s intracellular survival may represent the crucial

selective environment that has led to the *dctA*-repression phenotype (46). It is conceivable that

Salmonella recognizes the succinate produced by activated macrophage and restricts its uptake of

dicarboxylates in response. Our examination of *Salmonella* strains that are impaired in their

regulation of dicarboxylate uptake identified no survival defect in acidifed succinate or in

macrophage cell lines. However, it remains possible that this repression phenotype is related to

other aspects of the *Salmonella* virulence cycle beyond short-term survival in phagocytes. For

instance, *Salmonella* may potentially restrict its import and consumption of dicarboxylates to

maximize macrophage succinate levels leading to the production of the pro-inflammatory

citokine IL-1β (46). This would not grant *Salmonella* an advantage for survival in macrophage

per se, but would give *Salmonella* remaining in the gut lumen a growth advantage by

maximizing the immune system's oxidative burst and subsequent production of tetrathionate, a

compound that *Salmonella* is distinctively equipped to use as a terminal electron acceptor (58–

60). Thus, if *Salmonella* were to import and consume succinate in macrophage, the inflammatory

response might be deterred, restricting the ability of *Salmonella* to outcompete the gut microbtioa

during infection.

We did note a few *Salmonella* strains that grew early on succinate despite having a

catalase positive (and presumably *rpoS*+) phenotype. In all instances where multiple strains from

the same *Salmonella enterica* serovar were tested, at least one exhibited an extended lag phase

(Table S1). Thus, the earlier growth does not appear to be a trait of particular serovars but rather

may reflect individual strains having lost (or never acquired) the delayed growth phenotype. This

could occur by mutations in genes other than *rpoS,* such as *efp* or *iraP*, that permit early growth

in dicarboxylic acids. Other exceptions include multiple *E. coli* strains that show an extended lag

in their growth using succinate. While it is possible that these have mutations in genes required

for the uptake of succinate (such as *dctA* or *dcuSR*), these may reflect genuine variation in how

E. coli strains respond to dicarboxylates.

Finally, our findings have important implications for the interpretation of previous

metabolic studies in commonly used laboratory strains. For example, it is important to consider

that the Biolog assays primarily use succinate as the carbon source under many conditions (28,

29, 31). Thus, what we previously assumed to be improved growth of the *efp* mutant in a variety

of different conditions in fact resulted solely from its ability to rapidly utilize dicarboxylate

media. We also note the large number of metabolic studies carried out with the commonly used

Salmonella Typhimurium strain, LT2, which is a known *rpoS* mutant (32). Because of this,

studies using the LT2 strain would behave quite differently in Biolog assays than the other

commonly used 'wild-type' *Salmonella* strains SL1344 (of which many isolates are auxotrophic

for histidine) and 14028s.

Materials and Methods

Bacterial strains and plasmids

Bacterial strains, plasmids and primers are listed in Supplementary Tables S2-S4. As

described previously, lambda red recombination (28, 61) and subsequent P22 phage transduction

(62) was used to generate all of the gene knockout mutants in *Salmonella enterica subsp.*

enterica serovar Typhimurium (*S.* Typhimurium) strain 14028s. *E. coli* was from the Keio

collection K12 BW25113 strain background (63). To sample the genetic diversity of *Salmonella*

and *E. coli* isolates, the *Salmonella* genetic stock centre (SGSC) SARA (64), SARB (65), and

SARC (66) collections were employed and compared to the *E. coli* reference (ECOR) collection

(67).

The full length *DctA* ORF was expressed from the pXG10sf plasmid under the control of

the constitutively active PLtet0-1 promoter (68, 69). The IraP complementation plasmid was

generated by inserting the *iraP* ORF and the upstream 300bp into pXG10sf. For promoter

expression, the *dctA* promoter (500bp upstream of the *dctA* start codon) was inserted into

pXG10sf to drive expression of superfolder GFP (69, 70). To generate the chromosomal *dctA*

promoter swap strain, 500bp upstream of the *E. coli dctA* start codon, along with a

chloramphenicol resistance cassette for selection, was inserted into the corresponding location of

the *Salmonella* chromosome.

Growth using dicarboxylates as a sole carbon source

Overnight LB cultures inoculated from single colonies were resuspended in MOPS

minimal media with no carbon source to an optical density (OD_{600}) of approximately 1.75. This

suspension was used to inoculate (1/200 dilution) MOPS minimal media containing 0.2% carbon

source (succinate unless otherwise indicated). Growth was conducted in a TECAN Infinite M200

plate reader at 37ºC with shaking and OD_{600} was read every 15 minutes. For salts and hydrates of

carbon sources the final concentration reflects the percent of the carbon source itself (e.g., 0.2%

succinate was made as 0.47% sodium succinate dibasic hexahydrate).

For the SGSC and ECOR collections screen, 47 strains were assessed in duplicate per run

in a 96-well plate. Wild-type and *rpoS* mutant *Salmonella* were included on every plate as

quality controls. Each strain was tested on at least three separate days.

Catalase assay

414   For each replicate of the SGSC and ECOR collections screen, each strain was tested for

415 catalase activity as an analog for RpoS function (71). In parallel to the LB overnight cultures

416 used as inoculum, 10µl of each culture was spotted onto an LB plate. The next day the spots

417 were tested for catalase activity by the addition of 10µl hydrogen peroxide. Bubbling was scored

418 compared to wild-type (catalase positive) and *rpoS* mutant (catalase negative) *Salmonella*.

Acid survival

LPM media was made as described previously (72) and succinate was added to either

0.2% or 0.4% as indicated in figures. The pH of the media was then adjusted to 4.4. LB

overnight cultures were resuspended to an OD of 0.1 in acidified media and incubated in a 37ºC

water bath. At time points, samples were taken, serial diluted and plated for colony forming units

(CFU).

Intra-macrophage survival

The THP-1 human monocyte cell line and the J774 mouse macrophage cell line were

maintained in RPMI Medium 1640 (with L-glutamine) supplemented with 10% FBS and 1%

Glutamax, and grown at 37ºC and 5% CO_2. For infection assays, THP-1 cells were seeded in 96-

well plates at 50,000 per well with 50nM PMA (phorbol 12-myristate 13-acetate) added to the

media to induce differentiation to adherent macrophage. After 48h, the media was replaced with

normal growth media (no PMA) overnight. For infections with J774 macrophage the cells were

seeded in 96-well plates at 50,000 per well overnight. *Salmonella* in RPMI were added onto

seeded cells at a multiplicity of infection (MOI) of approximately 20 bacteria to 1 macrophage

and centrifuged for 10 minutes at 1000rpm for maximum cell contact. After centrifuging the

plate was placed at 37ºC (5% CO_2) and this was called 'time 0'. After 30 minutes, non-adherent

Salmonella were washed off by three washes with PBS followed by replacement with fresh

media containing 100 μg/ml gentamicin to kill extracellular *Salmonella*. At 2 hours the media

was replaced with media containing gentamicin at 10 μg/ml. At timepoints, intracellular bacteria

were recovered using PBS containing 1% Triton X-100 and vigorous pipetting. Samples were

serially diluted and five 10μl spots were plated for CFU counting. Each sample included three

separate wells as technical replicates (a total of 15 x 10μl spots counted per biological replicate).

Acknowledgements

We would sincerely like to thank Dr. Scott Gray-Owen and members of his lab, in

particular Dr. Ryan Gaudet, for their generous donation of technical expertise, macrophage cells

lines, and use of their equipment. WWN was supported by an Operating Grant from the Canada

Institutes for Health Research (MOP-86683) and a Natural Sciences and Engineering Research

Council (NSERC) of Canada Grant (RGPIN 386286-10). SJH was supported by an NSERC

Vanier Canada Graduate Scholarship.

References

1.   Shimizu K. 2016. Metabolic Regulation and Coordination of the Metabolism in Bacteria

in Response to a Variety of Growth Conditions. Adv Biochem Eng Biotechnol 155:1–54.

2.   Navarre WW, Halsey TA, Walthers D, Frye J, McClelland M, Potter JL, Kenney LJ,

Gunn JS, Fang FC, Libby SJ. 2005. Co-regulation of Salmonella enterica genes required

for virulence and resistance to antimicrobial peptides by SlyA and PhoP/PhoQ. Mol

Microbiol 56:492–508.

3.   Tu X, Latifi T, Bougdour A, Gottesman S, Groisman EA. 2006. The PhoP/PhoQ two-

component system stabilizes the alternative sigma factor RpoS in Salmonella enterica.

Proc Natl Acad Sci U S A 103:13503–13508.

4.   Dalebroux ZD, Miller SI. 2014. Salmonellae PhoPQ regulation of the outer membrane to

resist innate immunity. Curr Opin Microbiol 17:106–113.

5.   Fields PI, Swanson R V, Haidaris CG, Heffron F. 1986. Mutants of Salmonella

typhimurium that cannot survive within the macrophage are avirulent. Proc Natl Acad Sci

U S A 83:5189–5193.

6.   Bäumler AJ, Kusters JG, Stojiljkovic I, Heffron F. 1994. Salmonella typhimurium loci

involved in survival within macrophages. Infect Immun 62:1623–1630.

7.   Boutte CC, Crosson S. 2013. Bacterial lifestyle shapes stringent response activation.

Trends Microbiol 21:174–180.

8.   Magnusson LU, Farewell A, Nyström T. 2005. ppGpp: a global regulator in Escherichia

coli. Trends Microbiol 13:236–242.

9.   Dalebroux ZD, Svensson SL, Gaynor EC, Swanson MS. 2010. ppGpp conjures bacterial

virulence. Microbiol Mol Biol Rev 74:171–199.

10. Potrykus K, Cashel M. 2008. (p)ppGpp: still magical? Annu Rev Microbiol 62:35–51.

11. Dong T, Schellhorn HE. 2010. Role of RpoS in virulence of pathogens. Infect Immun

78:887–897.

12. Fang FC, Libby SJ, Buchmeier NA, Loewen PC, Switala J, Harwood J, Guiney DG. 1992.

The alternative sigma factor katF (rpoS) regulates Salmonella virulence. Proc Natl Acad

Sci U S A 89:11978–11982.

13. Kazmierczak MJ, Wiedmann M, Boor KJ. 2005. Alternative Sigma Factors and Their

Roles in Bacterial Virulence. Microbiol Mol Biol Rev 69:527–543.

14. Battesti A, Majdalani N, Gottesman S. 2011. The RpoS-Mediated General Stress

Response in Escherichia coli. Annu Rev Microbiol 65:189–213.

15. Bearson SM, Benjamin Jr WH, Swords WE, Foster JW. 1996. Acid shock induction of

RpoS is mediated by the mouse virulence gene mviA of Salmonella typhimurium. J

Bacteriol 178:2572–2579.

16. Muffler A, Fischer D, Altuvia S, Storz G, Hengge-Aronis R. 1996. The response regulator

RssB controls stability of the sigma(S) subunit of RNA polymerase in Escherichia coli.

EMBO J 15:1333–1339.

17. Pratt LA, Silhavy TJ. 1996. The response regulator SprE controls the stability of RpoS.

Proc Natl Acad Sci U S A 93:2488–2492.

18. Zhou Y, Gottesman S, Hoskins JR, Maurizi MR, Wickner S. 2001. The RssB response

regulator directly targets sigma(S) for degradation by ClpXP. Genes Dev 15:627–637.

19.   Bougdour A, Wickner S, Gottesman S. 2006. Modulating RssB activity: IraP, a novel

497   regulator of sigma(S) stability in Escherichia coli. Genes Dev 20:884–897.

498   20.   Bougdour A, Cunning C, Baptiste PJ, Elliott T, Gottesman S. 2008. Multiple pathways for

499   regulation of sigmaS (RpoS) stability in Escherichia coli via the action of multiple anti-

500   adaptors. Mol Microbiol 68:298–313.

501   21.   Bougdour A, Gottesman S. 2007. ppGpp regulation of RpoS degradation via anti-adaptor

502   protein IraP. Proc Natl Acad Sci U S A 104:12896–12901.

503   22.   Chiang SM, Dong T, Edge TA, Schellhorn HE. 2011. Phenotypic diversity caused by

504   differential RpoS activity among environmental Escherichia coli isolates. Appl Environ

505   Microbiol 77:7915–7923.

506   23.   Dong T, Chiang S, Joyce C, Yu R, Schellhorn H. 2009. Polymorphism and selection of

507   rpoS in pathogenic Escherichia coli. BMC Microbiol 9:118.

508   24.   Chen G, Patten CL, Schellhorn HE. 2004. Positive selection for loss of RpoS function in

509   Escherichia coli. Mutat Res Mol Mech Mutagen 554:193–203.

510   25.   Lo TC. 1977. The molecular mechanism of dicarboxylic acid transport in Escherichia coli

511   K 12. J Supramol Struct 7:463–480.

512   26.   Boogerd FC, Boe L, Michelsen O, Jensen PR. 1998. atp Mutants of Escherichia coli Fail

513   To Grow on Succinate Due to a Transport Deficiency. J Bacteriol 180:5855–5859.

514   27.   Davies SJ, Golby P, Omrani D, Broad SA, Harrington VL, Guest JR, Kelly DJ, Andrews

515   SC. 1999. Inactivation and Regulation of the Aerobic C4-Dicarboxylate Transport (dctA)

516   Gene ofEscherichia coli.

517   28.   Zou SB, Hersch SJ, Roy H, Wiggers JB, Leung AS, Buranyi S, Xie JL, Dare K, Ibba M,

518   Navarre WW. 2012. Loss of elongation factor P disrupts bacterial outer membrane

519   integrity. J Bacteriol 194:413–425.

520   29.   Navarre WW, Zou SB, Roy H, Xie JL, Savchenko A, Singer A, Edvokimova E, Prost LR,

521   Kumar R, Ibba M, Fang FC. 2010. PoxA, YjeK, and elongation factor P coordinately

522   modulate virulence and drug resistance in Salmonella enterica. Mol Cell 39:209–221.

523   30.   Hersch SJ, Wang M, Zou SB, Moon K-M, Foster LJ, Ibba M, Navarre WW. 2013.

524   Divergent protein motifs direct elongation factor P-mediated translational regulation in

525   Salmonella enterica and Escherichia coli. MBio 4.

526   31.   Bochner BR, Gadzinski P, Panomitros E. 2001. Phenotype MicroArrays for High-

527   Throughput Phenotypic Testing and Assay of Gene Function. Genome Res 11:1246–1255.

528   32.   Lee IS, Lin J, Hall HK, Bearson B, Foster JW. 1995. The stationary-phase sigma factor

529   sigma S (RpoS) is required for a sustained acid tolerance response in virulent Salmonella

530   typhimurium. Mol Microbiol 17:155–167.

531   33.   Hengge-Aronis R. 2002. Signal Transduction and Regulatory Mechanisms Involved in

532   Control of the σS (RpoS) Subunit of RNA Polymerase. Microbiol Mol Biol Rev 66:373–

533   395.

534   34.   Metris A, George SM, Mulholland F, Carter AT, Baranyi J. 2014. Metabolic shift of

535   Escherichia coli under salt stress in the presence of glycine betaine. Appl Environ

536   Microbiol 80:4745–4756.

537   35.   Liang X, Zhang L, Natarajan SK, Becker DF. 2013. Proline mechanisms of stress

538   survival. Antioxidants Redox Signal 19:998–1011.

539   36.   Lee E-J, Choi J, Groisman EA. 2014. Control of a Salmonella virulence operon by

540   proline-charged tRNA(Pro). Proc Natl Acad Sci 111:3140–3145.

541   37.   Peil L, Starosta AL, Lassak J, Atkinson GC, Virumäe K, Spitzer M, Tenson T, Jung K,

542   Remme J, Wilson DN. 2013. Distinct XPPX sequence motifs induce ribosome stalling,

543   which is rescued by the translation elongation factor EF-P. Proc Natl Acad Sci

544   110:15265–15270.

545   38.   Ude S, Lassak J, Starosta AL, Kraxenberger T, Wilson DN, Jung K. 2013. Translation

546   elongation factor EF-P alleviates ribosome stalling at polyproline stretches. Science (80-)

547   339:82–85.

548   39.   Liu LK, Becker DF, Tanner JJ. 2017. Structure, function, and mechanism of proline

549   utilization A (PutA). Arch Biochem Biophys 632:142–157.

550   40.   Arentson BW, Sanyal N, Becker DF. 2012. Substrate channeling in proline metabolism.

551   Front Biosci 17:375–388.

552   41.   Weber H, Polen T, Heuveling J, Wendisch VF, Hengge R. 2005. Genome-wide analysis

553   of the general stress response network in Escherichia coli: sigmaS-dependent genes,

554   promoters, and sigma factor selectivity. J Bacteriol 187:1591–1603.

555   42.   Patten CL, Kirchhof MG, Schertzberg MR, Morton RA, Schellhorn HE. 2004. Microarray

556   analysis of RpoS-mediated gene expression in Escherichia coli K-12. Mol Genet

557   Genomics 272:580–591.

558   43.   Jung IL, Kim SK, Kim IG. 2006. The RpoS-mediated regulation of isocitrate

559   dehydrogenase gene expression in Escherichia coli. Curr Microbiol 52:21–26.

560   44.   Srikumar S, Kröger C, Hébrard M, Colgan A, Owen S V, Sivasankaran SK, Cameron

561   ADS, Hokamp K, Hinton JCD. 2015. RNA-seq Brings New Insights to the Intra-

562   Macrophage Transcriptome of Salmonella Typhimurium. PLoS Pathog 11:e1005262.

563   45.   Kröger C, Colgan A, Srikumar S, Händler K, Sivasankaran SK, Hammarlöf DL, Canals R,

564   Grissom JE, Conway T, Hokamp K, Hinton JCD. 2013. An infection-relevant

565   transcriptomic compendium for Salmonella enterica Serovar Typhimurium. Cell Host

566   Microbe 14:683–695.

46.   Tannahill GM, Curtis AM, Adamik J, Palsson-McDermott E, McGettrick AF, Goel G,

Frezza C, Bernard NJ, Kelly B, Foley NH, Zheng L, Gardet A, Tong Z, Jany SS, Corr SC,

Haneklaus M, Caffrey BE, Pierce K, Walmsley S, Beasley FC, Cummins E, Nizet V,

Whyte M, Taylor CT, Lin H, Masters SL, Gottlieb E, Kelly VP, Clish C, Auron PE,

Xavier RJ, O'Neill LAJ. 2013. Succinate is an inflammatory signal that induces IL-1β

through HIF-1α. Nature 496:238–242.

47. Alpuche Aranda C.M., Swanson JA, Loomis WP, Miller SI. 1992. Salmonella

typhimurium activates virulence gene transcription within acidified macrophage

phagosomes. Proc Natl Acad Sci U S A 89:10079–10083.

48. Rathman M, Sjaastad MD, Falkow S. 1996. Acidification of phagosomes containing

Salmonella typhimurium in murine macrophages. Infect Immun 64:2765–2773.

49. Steinmetz PA, Wörner S, Unden G. 2014. Differentiation of DctA and DcuS function in

the DctA/DcuS sensor complex of Escherichia coli: function of DctA as an activity switch

and of DcuS as the C4-dicarboxylate sensor. Mol Microbiol 94:218–229.

50. Durfee T, Hansen AM, Zhi H, Blattner FR, Jin DJ. 2008. Transcription profiling of the

stringent response in Escherichia coli. J Bacteriol 190:1084–1096.

51. Traxler MF, Summers SM, Nguyen H-T, Zacharia VM, Hightower GA, Smith JT,

Conway T. 2008. The global, ppGpp-mediated stringent response to amino acid starvation

in Escherichia coli. Mol Microbiol 68:1128–1148.

52. Merrikh H, Ferrazzoli AE, Lovett ST. 2009. Growth phase and (p)ppGpp control of IraD,

a regulator of RpoS stability, in Escherichia coli. J Bacteriol 191:7436–7446.

53. Perez JC, Groisman EA. 2009. Evolution of Transcriptional Regulatory Circuits in

Bacteria. Cell 138:233–244.

54. Zwir I, Latifi T, Perez JC, Huang H, Groisman EA. 2012. The promoter architectural

landscape of the Salmonella PhoP regulon. Mol Microbiol 84:463–485.

55.   Osborne SE, Walthers D, Tomljenovic AM, Mulder DT, Silphaduang U, Duong N,

Lowden MJ, Wickham ME, Waller RF, Kenney LJ, Coombes BK. 2009. Pathogenic

adaptation of intracellular bacteria by rewiring a cis-regulatory input function. Proc Natl

Acad Sci U S A 106:3982–3987.

56.   Will WR, Brzovic P, Trong I Le, Stenkamp RE, Lawrenz MB, Karlinsey JE, Navarre

WW, Main-Hester K, Miller VL, Libby SJ, Fang FC. 2019. The evolution of SlyA/RovA

transcription factors from repressors to countersilencers in enterobacteriaceae. MBio

10:1–22.

57.   Spiga L, Winter MG, Furtado de Carvalho T, Zhu W, Hughes ER, Gillis CC, Behrendt

CL, Kim J, Chessa D, Andrews-Polymenis HL, Beiting DP, Santos RL, Hooper L V,

Winter SE. 2017. An Oxidative Central Metabolism Enables Salmonella to Utilize

Microbiota-Derived Succinate. Cell Host Microbe.

58.   Winter SE, Thiennimitr P, Winter MG, Butler BP, Huseby DL, Crawford RW, Russell

JM, Bevins CL, Adams LG, Tsolis RM, Roth JR, Bäumler AJ. 2010. Gut inflammation

provides a respiratory electron acceptor for Salmonella. Nature 467:426–429.

59.   Hensel M, Hinsley AP, Nikolaus T, Sawers G, Berks BC. 1999. The genetic basis of

tetrathionate respiration in Salmonella typhimurium. Mol Microbiol 32:275–287.

60.   Thiennimitr P, Winter SE, Winter MG, Xavier MN, Tolstikov V, Huseby DL, Sterzenbach

T, Tsolis RM, Roth JR, Bäumler AJ. 2011. Intestinal inflammation allows Salmonella to

use ethanolamine to compete with the microbiota. Proc Natl Acad Sci U S A 108:17480–

17485.

61.   Datsenko KA, Wanner BL. 2000. One-step inactivation of chromosomal genes in

Escherichia coli K-12 using PCR products. Proc Natl Acad Sci 97:6640–6645.

62.   Schmieger H. 1971. A method for detection of phage mutants with altered transducing

ability. Mol Gen Genet MGG 110:378–381.

63.   Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, Datsenko KA, Tomita M,

Wanner BL, Mori H. 2006. Construction of Escherichia coli K-12 in-frame, single-gene

knockout mutants: the Keio collection. Mol Syst Biol 2:2006.0008.

64.   Beltran P, Plock SA, Smith NH, Whittam TS, Old DC, Selander RK. 1991. Reference

collection of strains of the Salmonella typhimurium complex from natural populations.

Microbiology 137:601–606.

65.   Boyd EF, Wang F-S, Beltran P, Plock SA, Nelson K, Selander RK. 1993. Salmonella

reference collection B (SARB): strains of 37 serovars of subspecies I. Microbiology

139:1125–1132.

66.   Boyd EF, Wang FS, Whittam TS, Selander RK. 1996. Molecular genetic relationships of

the salmonellae. Appl Environ Microbiol 62:804–808.

67.   Ochman H, Selander RK. 1984. Standard reference strains of Escherichia coli from

natural populations. J Bacteriol 157:690–693.

68.   Urban J, Vogel J. 2009. A green fluorescent protein (GFP)-based plasmid system to study

post-transcriptional control of gene expression in vivo, p. 301–319. *In* Serganov, A (ed.),

Riboswitches. Humana Press.

69.   Corcoran CP, Podkaminski D, Papenfort K, Urban JH, Hinton JCD, Vogel J. 2012.

Superfolder GFP reporters validate diverse new mRNA targets of the classic porin

regulator, MicF RNA. Mol Microbiol 84:428–445.

70.   Pédelacq J-D, Cabantous S, Tran T, Terwilliger TC, Waldo GS. 2006. Engineering and

characterization of a superfolder green fluorescent protein. Nat Biotechnol 24:79–88.

71.   Loewen PC, Triggs BL. 1984. Genetic mapping of katF, a locus that with katE affects the

synthesis of a second catalase species in Escherichia coli. J Bacteriol 160:668–675.

72.   Cooper CA, Zhang K, Andres SN, Fang Y, Kaniuk NA, Hannemann M, Brumell JH,

641   Foster LJ, Junop MS, Coombes BK. 2010. Structural and biochemical characterization of

642   SrcA, a multi-cargo type III secretion chaperone in Salmonella required for pathogenic

643   association with a host. PLoS Pathog 6:e1000751.

Figure Legends

Figure 1: *Salmonella* **displays an extended lag phase using dicarboxylic acids as a sole carbon source** but *efp* and *rpoS* mutants do not. **A)** Representative growth curve of *Salmonella* Typhimurium 14028s strains grown in MOPS minimal media with 0.2% succinate as the sole carbon source. Data shows optical density at 600nm (OD_{600}) for 48 hours from the time of inoculation. The data shown is representative of greater than three biological replicates. An OD_{600} of 0.1 is emphasized by a dashed line. **B)** Graphs showing the average time in hours that wild-type (WT) or mutant *Salmonella* takes to reach an OD of 0.1 as an analog of the length of lag phase using succinate as the sole carbon source. The data shows the average of at least three biological replicates and error bars show one standard deviation. >48h indicates the strain did not grow by 48 hours. **C)** As in (B) but comparing the use of three different dicarboxylic acids as the sole carbon source.

Figure 2: *Salmonella* **grows significantly later than *E. coli* using succinate** as the sole carbon source. Growth in MOPS minimal media with 0.2% succinate as the sole carbon source. Growth was conducted in a TECAN Infinite M200 plate reader and reads were taken every 15 minutes. **A)** Representative growth curve. An OD_{600} of 0.1 is emphasized by a dashed line. **B)** Percentage of SGSC *Salmonella* (green line) or ECOR *E. coli* (blue line) strains that (on average) had surpassed an OD_{600} of 0.1 by indicated times post inoculation. Red line is for *Salmonella* but excludes the 14 tested strains of the SARC collection that do not belong to subspecies *enterica*. **C)** Overview of 105 *Salmonella* strains (SGSC collection) and 72 *E. coli strains* (ECOR collection). Each strain is plotted by average time it takes to reach OD 0.1 (x-axis) compared to doubling time during growth from OD 0.1 to OD 0.2. All *Salmonella* are coloured black and all *E. coli* strains are coloured red. Strains with at least one replicate that did not grow by 72 hours

are shown as triangles; these values are the average of the remaining replicates. Inset at top right

shows zoomed out view to include outliers. **D)** Average time in hours for strains to reach an OD

of 0.1. Data compares all 105 *Salmonella* strains to all 72 *E. coli* strains tested. Line indicates the

median, boxes show the 25th to 75th percentiles, and whiskers show the 10th to 90th percentiles.

An unpaired t-test with Welch's correction indicated a p-value < 0.0001. **E)** As in (D) but

comparing doubling time in hours. An unpaired t-test with Welch's correction indicated a p-

value < 0.05 with all data points included or p-value < 0.0001 when excluding slow growing

outliers (doubling time > 5h).

Figure 3: Deletion of the IraP results in early growth on succinate. Growth of *Salmonella* in

MOPS minimal media with 0.2% succinate as the sole carbon source. **A)** The three known RssB

anti-adaptors were deleted from the *Salmonella* chromosome and growth is shown along with an

rssB mutant. Data is representative of at least three biological replicates. **B)** Average time in

hours for wild-type (WT) or mutant *Salmonella* to reach an OD of 0.1 as an analog of the length

of lag phase using succinate as the sole carbon source. The data shows the average of at least

three biological replicates and error bars show one standard deviation. **C)** Plasmid expression of

the *Salmonella iraP* gene partially complements the growth delay phenotype. **D)** As for B.

Figure 4: Specific nutrients induce growth using succinate and diauxic repression involves

the stringent response. A and C) Representative growth curves of *Salmonella* growth in MOPS

minimal media with 0.2% succinate as the primary carbon source and supplemented with

0.005% proline (+ Pro) or 0.005% citrate (+ Cit) where indicated. **B and D)** Graphs showing the

average time in hours for *Salmonella* to reach an OD of 0.1 as an analog of the length of lag

phase using succinate as the sole carbon source. The data shows the average of at least three

biological replicates and error bars show one standard deviation.

Figure 5: Expression of *dctA* induces growth using succinate. Growth of *Salmonella* in

MOPS minimal media with 0.2% succinate as the sole carbon source. **A)** Overexpression of *dctA*

from a plasmid. WT pLacZ and pDctA indicate wild-type *Salmonella* containing a pXG10sf

plasmid encoding full length *lacZ* or *dctA* with expression driven by the constitutively active

PLtet0-1 promoter. The *rpoS* mutant is shown for comparison. **B)** Graph showing the average

time in hours for wild-type (WT) or mutant *Salmonella* to reach an OD of 0.1 as an analog of the

length of lag phase using succinate as the sole carbon source. The data shows the average of at

least three biological replicates and error bars show one standard deviation. **C)** Stretches of the

E. coli dctA promoter (P$_{dctA}$) were inserted into the *Salmonella* chromosome replacing the native

dctA promoter. The length inserted (in base-pairs counting back from the *dctA* start codon) are

indicated. Replacement with *Salmonella*'s native *dctA* promoter controls for effects due to the

insertion method (P$_{dctA}$ *Salm.*). Wild-type *Salmonella* and *E. coli* K12 are shown for comparison.

D) As for B.

Figure 6: Replacement of the *Salmonella dctA* promoter does not influence survival in

acidified succinate or macrophage. **A)** Survival of *Salmonella* treated with LPM media

containing 0.2% succinate and acidified to pH 4.4. Wild-type (WT) is compared to *Salmonella*

with its chromosomal *dctA* promoter replaced with 500bp of the *E. coli dctA* promoter (P$_{dctA}$ *E.*

coli) or *Salmonella*'s native *dctA* promoter as a control (P$_{dctA}$ *Salm*). An *rpoS* mutant is shown as

a positive control. CFU recovered at 3 hours were normalized to input CFU at 0h and expressed

as percent survival. Data shows the average across three biological replicates and error bars

indicate one standard deviation. **B)** Infection of THP-1 human macrophage comparing survival

of wild-type *Salmonella* (WT) with various mutant strains and the *dctA* promoter swap strain. A

phoP mutant is included as a positive control. Data shows a logarithm of CFU recovered at 24

hours post infection and is the average of three biological replicates. Error bars show one

standard deviation. **C)** As in (B) but showing CFU recovered from mouse J774 macrophage.

Figure 7: Model of the link between stress and dicarboxylate utilization in *Salmonella*.

Under conditions of stress RpoS (σ^S) levels are elevated due to the production of anti-adaptors

that block the degradation of RpoS by the RssB adaptor and ClpXP protease system. RpoS

directs RNA polymerase (RNAP) to several promoters involved in stress protection as well as to

a series of downstream regulatory genes (including small RNAs) that decrease the production of

enzymes involved in the TCA cycle and other metabolic pathways. One of these downstream

regulatory factors, yet to be identified, impinges on the expression of *dctA*. The presence of

either citrate or proline is sufficient to alleviate the block on dicarboxylate utilization. The

proline mediated effect requires PutA, a polyfunctional enzyme that converts proline to

glutamate with the concomitant production of reducing equivalents that are transferred to NAD+

and ubiquinone (figure created with BioRender). The factors that mediate the citrate-dependent

effect remain unclear.

A

LPM pH 4.4 + 0.2% Succinate

B

Human THP-1 Macrophage

C

Mouse J774 Macrophage

Supplemental Materials and Methods

Promoter induction fluorescence

      Growth was conducted in a TECAN Infinite M200 plate reader at 37ºC with shaking and OD_{600} and GFP fluorescence (475nm and 511nm excitation and emission wavelengths respectively) were read every 15 minutes. For clarity, bar graphs show fluorescence at 16h post inoculation. Chloramphenicol was included in all media at a concentration of 20μg/ml to maintain the plasmids.

Supplemental Figures

Figure S1: SGSC *Salmonella* trend towards longer lag phase than ECOR *E. coli* regardless of catalase activity. Comparing time to reach OD 0.1 in succinate media with catalase result for all strains tested from the SGSC and ECOR collections. + or − indicate that the strain showed positive or negative (respectively) catalase result in all replicates. +/− indicates either delayed positive or inconsistent catalase results across replicates. Data is the average of at least three biological replicates with the exception of strains shown as triangles that had at least one replicate that did not reach OD 0.1 by 72h (shown as average of the remaining replicates).

Figure S2: Growth in succinate media is induced by dilute amino acids. Growth of

Salmonella in MOPS minimal media using **A)** 0.2% succinate as the sole carbon source, **B)** 1/10

diluted LB as the sole source of carbon, **C)**, both 0.2% succinate and 1/10 diluted LB. Legend for

A-C is indicated at right. **D)** Growth of wild-type *Salmonella* in MOPS minimal media with

0.2% succinate as the primary carbon source and supplemented with individual or pooled amino

acids at a concentration approximately 1/10 of that in LB media (1). **E)** Growth of *Salmonella*

wild-type (WT) or *rpoS* mutant (Δ*rpoS*) in MOPS minimal media with 0.2% succinate (Succ.)

and/or 0.01% proline (Pro.) as the sole carbon sources. Graphs are representative of at least two

independent replicates.

Figure S3: Alignment of *Salmonella* and *E. coli dctA* promoters conducted using

Clustal Ω software (2). Asterisks indicate identity and regions of consistent differences between

E. coli and *Salmonella* are highlighted in yellow. The 200bp from *E. coli* that were swapped into

the *Salmonella* chromosome in Figure 5C and D are indicated (-200), as are the predicted -35

and -10 promoter elements identified by RegulonDB (3) and supplemental reference (4). The

transcriptional start site (TSS) is indicated in blue with red text indicating the 5' UTR as

identified by RegulonDB (3) and in references (4, 5). The ATG start codons are underlined.

Figure S4: The *Salmonella dctA* promoter shows reduced expression in *E. coli* relative to the *E. coli dctA* promoter. Trancriptional fusions to GFP on a plasmid of the *Salmonella* or *E. coli dctA* promoters. GFP fluorescence was examined in *E. coli* K12 and data shows relative expression at 48h post-inoculation into MOPS minimal media with 0.2% succinate as the sole carbon source. Data is the average of three biological replicates and error bars show one standard deviation.

Figure S5: Overexpression of DctA from a plasmid decreases survival in macrophage due to toxicity. A) Infection of THP-1 macrophage examining survival of wild-type *Salmonella* with no plasmid (WT) compared to isogenic strains constitutively expressing LacZ (pLacZ) or DctA (pDctA) from the pXG10sf plasmid. Survival of *E. coli* and Δ*phoP* mutant *Salmonella* are included as controls for comparison. Data shows CFU recovered at 24 hours post-infection relative to wild-type *Salmonella* and is the average of three biological replicates. Error bars show one standard deviation. **B)** Survival of wild-type or *dctA* mutant *Salmonella* treated with LPM media containing 0.4% succinate and acidified to pH 4.4. CFU recovered at 3 hours were normalized to input CFU at 0h and expressed as percent survival. Data shows the average across three biological replicates and error bars indicate one standard

deviation. Where indicated, genes were expressed from the ampicillin resistant version of the pXG10sf plasmid under the control of the constitutively active PLtet0-1 promoter. **C)** Representative image showing overnight growth of wild-type *Salmonella* on an LB agar plate comparing colony size when constitutively expressing *dctA* point mutants. Black arrow indicates a small colony of the strain expressing DctA^{S380D}; white arrow indicates a large colony likely resulting from a suppressor mutation.

Table S2: Strains*

Strain	Genotype	ID	Description	Reference
Salmonella Typhimurium 14028S		WN150	Wild-type *Salmonella enterica* serovar Typhimurium strain 14028S	ATCC
	Δ*efp*	WN1405	Deletion of *efp* gene in WN150 background. Resistance cassette was been removed by FLP recombination	(6)
	Δ*rpoS::kan-FRT*	WN2881	Deletion of *rpoS* gene in WN150 background	This work
	Δ*dctA*	WN2914	Deletion of *dctA* gene in WN150 background. Resistance cassette was been removed by FLP recombination	This work
	Δ*iraP*	WN3186	Deletion of *iraP* gene in WN150 background. Resistance cassette was been removed by FLP recombination	This work
	Δ*rssB::cm-FRT*	WN3162	Deletion of *rssB* gene in WN150 background	This work
	Δ*rssC::cm-FRT*	WN3163	Deletion of *rssC* gene in WN150 background	This work
	Δ*iraD::cm-FRT*	WN3165	Deletion of *iraD* gene in WN150 background	This work
	Δ*putA::kan-FRT*	WN2963	Deletion of *putA* gene in WN150 background	This work
	Δ*relA::cm-FRT* Δ*spoT30 (sms-211)::Tn10*	WN138	Deletion of *relA* and *spoT* genes in 14028S background. Called ppGpp^0 in text and figures	This work
	Δ*phoP::cm-FRT*	WN152	Deletion of *phoP* gene in 14028S background. Also known as CS015	(7)
	P$_{dctA}$(*Salm.*)	WN3188	*Salmonella dctA* promoter with upstream chloramphenicol resistance cassette in the chromosome of the WM150 background.	This work
	P$_{dctA}$(*E. coli*) 500	WN3189	*E. coli dctA* promoter (500bp upstream of the start codon) with upstream chloramphenicol resistance cassette in the chromosome of the WM150 background.	This work
	P$_{dctA}$(*E. coli*) 200	WN3239	*E. coli dctA* promoter (200bp upstream of the start codon) in the chromosome of the WM150 background. -500 to -200bp remain from the *Salmonella* promoter preceded by an upstream chloramphenicol resistance cassette.	This work
	P$_{dctA}$(*E. coli*) 54	WN3238	*E. coli dctA* 5' UTR in the	This work

| | | | chromosome of the WM150 background. -500 to -54bp remain from the *Salmonella* promoter preceded by an upstream chloramphenicol resistance cassette. | |
| *E. coli* K12 BW25113 | | WN0970 | Wild-type *E. coli* strain K12 | (8) |

* Additional strains from the SGSC and ECOR collections are listed in Table S1

Table S3: Plasmids

Plasmid	Shorthand in Figures	Description	Reference
pXG10sf-PLtet0-1-LacZ-sfGFP	pLacZ	Constitutive expression of LacZ-sfGFP fusion	(9)
pXG10sf-PLtet0-1-DctA-LacZ-sfGFP		Full length DctA ORF inserted into pXG10sf	This work
pXG10sf-PLtet0-1-DctA	pDctA	Constitutive expression of DctA. Inserted stop codon into pXG10sf-PLtet0-1-DctA-LacZ-sfGFP	This work
pXG10sf-PLtet0-1-DctA-N301A	pDctA^{N301A}	Constitutive expression of DctA^{N301A}.	This work
pXG10sf-PLtet0-1-DctA-S380D	pDctA^{S380D}	Constitutive expression of DctA^{S380D}.	This work
pXG10sf-IraP	pIraP	Expression of the *iraP* ORF from its native promoter (300bp upstream of the start codon)	This work
pXG10sf-P$_{dctA}$ (*Salm.*)-LacZ-sfGFP	pP$_{dctA}$(*Salm.*)	For measuring expression from the *Salmonella dctA* promoter (500bp upstream of the start codon).	This work
pXG10sf-P$_{dctA}$ (*E. coli.*) 500-LacZ-sfGFP	pP$_{dctA}$(*E. coli*) 500	For measuring expression from the *E. coli dctA* promoter (500bp upstream of the start codon).	This work
pXG10sf-P$_{dctA}$ (*Salm.*)-DctA (*Salm.*)	P$_{dctA}$(*Salm.*)	-Inserted the 500bp upstream of the *dctA* start codon from *Salmonella* into pXG10sf-PLtet0-1-DctA-LacZ-sfGFP (replacing PLtet0-1). -Used as template for insertion (by lambda red recombination) of the *Salmonella dctA* promoter (with the upstream chloramphenicol resistance cassette taken from the plasmid) into the *Salmonella* chromosome.	This work
pXG10sf-P$_{dctA}$ (*E.	P$_{dctA}$(*E. coli*)	-Inserted the 500bp upstream of the *dctA*	This work

coli.) 500-DctA (*Salm.*)	500	start codon from *E. coli* into pXG10sf-PLtet0-1-DctA-LacZ-sfGFP (replacing PLtet0-1). -Used as template for insertion (by lambda red recombination) of the *E. coli dctA* promoter (with the upstream chloramphenicol resistance cassette taken from the plasmid) into the *Salmonella* chromosome.	
pXG10sf-P$_{dctA}$ (*E. coli.*) 200-DctA (*Salm.*)	P$_{dctA}$(*E. coli*) 200	-Inserted the 200bp upstream of the *dctA* start codon from *E. coli* into the pXG10sf-P$_{dctA}$ (*Salm.*)-DctA (*Salm.*) plasmid. (-500 to -200bp remain from the *Salmonella* promoter). -Used as template for insertion (by lambda red recombination) of the *E. coli dctA* 200bp promoter (with the upstream chloramphenicol resistance cassette taken from the plasmid) into the *Salmonella* chromosome.	This work
pXG10sf-P$_{dctA}$ (*E. coli.*) 54-DctA (*Salm.*)	P$_{dctA}$(*E. coli*) 54	-Inserted the 5' UTR from *E. coli dctA* into the pXG10sf-P$_{dctA}$ (*Salm.*)-DctA (*Salm.*) plasmid. (-500 to -54bp remain from the *Salmonella* promoter). -Used as template for insertion (by lambda red recombination) of the *E. coli dctA* 5' UTR (with the upstream chloramphenicol resistance cassette taken from the plasmid) into the *Salmonella* chromosome.	This work

Table S4: Primers

Name	Description	Sequence (length)
DctA NsiI F	For inserting full length DctA into pXG10sf	GTTTTTATGCATGCAGGTTACTCATAACCTTACAAGACCTGTGG (44)
DctA FL NheI R		GTTTTTGCTAGCGGAGGAAATTTCGTGCGTTTTGCCATC (39)
DctA STOP SDM F	For site-directed mutagenesis insertion of stop codon into pXG10sf-PLtet0-1-DctA-LacZ-sfGFP	AAATTTCCTCCTAAGCTAGCGGATCCGCTGGCTCCGCTGCTGGTTCTGGCG (51)
DctA STOP SDM R		GATCCGCTAGCTTAGGAGGAAATTTCGTGCGTTTTGCCATCCGGCGCACG (50)

DctA S380D SDM F	For site-directed mutagenesis of pXG10sf-PLtet0-1-DctA	GACCGCTTTATGGACGAAG CGCGCGCGCTGACTAACCT GGTGGGTAACGG (50)
DctA S380D SDM R		CGCGCGCTTCGTCCATAAA GCGGTCGATACCGAGGATT AACGCCAGTCCCGC (52)
DctA N301A SDM F	For site-directed mutagenesis of pXG10sf-PLtet0-1-DctA	GGTATTCGTTCGCCCTCGA TGGCACCTCTATCTACCTG ACGATGGCGGCG (50)
DctA N301A SDM R		TGCCATCGAGGGCGAACG AATACCCCGTCGGGATCAC CAGCCCTACAACC (50)
PiraP_salm F	For inserting full length IraP and upstream 300bp for native promoter into pXG10sf	CGTATCACGAGGCCCTTTC GTCTTCACCTCAGGCCAAT AGCGGTGCATC (49)
IraPsalm FL R		GGATCCGCTAGCGAGGCG GTTTTCTCCTTAGTGCCGG GGGTGTCTC (46)
pXG10 prom ATW F	PCR amplification of pXG10sf vector excluding the promoter region for insertion of alternate promoters as fusions to LacZ-sfGFP	ATGACCATGATTACGGATT CACTGG (25)
pXG10 prom ATW R	PCR amplification of pXG10sf vector excluding the promoter region for insertion of alternate promoters	GAGGTGAAGACGAAAGGG CC (20)
PdctA F (for pXG)	For amplifying *dctA* promoter (from 500bp before ATG start codon to 15bp into ORF) for insertion into pXG10sf.	CGTATCACGAGGCCCTTTC GTCTTCACCTCTAATGCTG GCGTGCGTATCG (50)
PdctA R (for pXG)	For amplifying *Salmonella dctA* promoter (from 500bp before ATG start codon to 15bp into ORF) for insertion into pXG10sf as fusion to LacZ-sfGFP	GACGGCCAGTGAATCCGTA ATCATGGTCATCAGAGAGG TTTTCATAGGGTGTCC (54)
PdctA e coli F (for pXG)	For amplifying *dctA* promoter from *E. coli* (from 500bp before ATG start codon to 15bp into ORF) for insertion into pXG10sf as fusion to LacZ-sfGFP. Used with PdctA (for pXG R)	CGTATCACGAGGCCCTTTC GTCTTCACCTCGGAGTTCG GGTGGCGCTGG (49)
pDctA ATW F	PCR amplification of pXG10sf-PLtet0-1-DctA vector for insertion of *Salmonella* or *E. coli dctA* promoter. Used with pXG10sf Prom ATW R.	AGGACACCCTATGAAAACC TCTCTG (25)
PdctAcoli200 R	For vector PCR using pXG10sf-P_dctA (*Salm.*)-DctA (*Salm.*) template.	CTTTTCTTCCAGGTACCGTT CTTCAAAGATATCCGCGGG (39)
new sfGFP qPCR F		GTTCCATGGCCAACACTTG TCACT (24)

PdctAcoli 200 F	For insert PCR using pXG10sf-P$_{dctA}$ (*E. coli.*) 500-DctA (*Salm.*) as template.	CTTTGAAGAACGGTACCTG GAAGAAAAGTAGCTACCC CAAAC (42)
new sfGFP qPCR R		TACATAACCTTCGGGCATG GCACT (24)
PdctAcoli UTR SDM F	For site directed mutagenesis using pXG10sf-P$_{dctA}$ (*Salm.*)-DctA (*Salm.*) template to replace insert *E. coli dctA* 5' UTR.	TAACCTTACAAGACCTGTG GTTTTACTAAAGGACACCC TATGAAAACCTCTCTGTTC (57)
PdctAcoli UTR SDM R		AGTAAAACCACAGGTCTTG TAAGGTTATGGGGTACCTG CGGCCTTAAAATAACACCC (57)
CmR-PdctA-DctA lambda red	For inserting PdctA-DctA(*Salm.*) constructs into *Salmonella* chromosome with upstream chloramphenicol resistance cassette. Used with pXG10sf-PdctA (Salm.)-DctA (Salm.),	AACCTTACAAGACCTGTGG TTTTTACTTCAAGGACACC CTTTACGCCCCGCCCTGCC ACT (60)
dctA FL lambda red R	pXG10sf-PdctA (E. coli.) 500-DctA (Salm.), pXG10sf-PdctA (E. coli.) 200-DctA (Salm.), or pXG10sf-PdctA (E. coli.) 54-DctA (Salm.) templates.	GCAGTCCTGGAAGGGACTA CGGGCATAAGTGGCGAGA TTAGGAGGAAATTTCGTGC GTTT (60)
RpoS KO F	For deletion of *rpoS* by lambda-red recombination.	TTGCTAGTTCCGTCAAGGG ATCACGGGTAGGAGCCACC TTGTGTAGGCTGGAGCTGC TTC (60)
RpoS KO R		GTCGGTAATCCCCGCCGGG CTTGAGCCCGGATCCAGTT CCCATATGAATATCCTCCT TAG (60)
DctA KO F	For deletion of *dctA* by lambda-red recombination.	AACCTTACAAGACCTGTGG TTTTTACTTCAAGGACACC CTGTGTAGGCTGGAGCTGC TTC (60)
DctA KO R		CCCGCAGTCCTGGAAGGGA CTACGGGCATAAGTGGCGA GACATATGAATATCCTCCT TAG (60)
IraP KO F	For deletion of *iraP* by lambda-red recombination.	CTTCATAGTGATAACGTCA CCCTGGAACTAATAAGGAA ATGTGTAGGCTGGAGCTGC TTC (60)
IraP KO R		TAAAAGTAACGTTATAACA ACTGTGTTGTTTTAAATAC GACATATGAATATCCTCCT TAG (60)

RssB KO F	For deletion of *rssB* by lambda-red recombination.	ACCATGCCACTATTGATTAAAGCCAGTCAGGGGAGAGAACGTGTAGGCTGGAGCTGCTTC (60)
RssB KO R		AAAGGCCGGTAAAGCAATTTCCGCTCACTCTTCCGTTTGGCATATGAATATCCTCCTTAG (60)
RssC KO F	For deletion of *rssC* by lambda-red recombination.	AATTTACACCATTGACTATTTCACTTAAAAGGTTTCTGTAGTGTAGGCTGGAGCTGCTTC (60)
RssC KO R		TACACTTAATCAGTAGTATGGCTAATGCATACTTTATTTCCATATGAATATCCTCCTTAG (60)
IraD KO F	For deletion of *iraD* by lambda-red recombination.	CGTCAGTATGGGATCTGGCGATGAACCAGATAGGGAAATGGTGTAGGCTGGAGCTGCTTC (60)
IraD KO R		ATAATGTTAAACGCCCGGCCCGGCGTTCCTTCCTGCATTACATATGAATATCCTCCTTAG (60)
PutA KO F	For deletion of *putA* by lambda-red recombination.	TATTGGTTAGCAGATAGAGCGAATAACAGGAGTAGAGGGTGTGTAGGCTGGAGCTGCTTC (60)
PutA KO R		CCATTGACGGAGCCTTAATTAACCACAGTATTAGCCAATTCATATGAATATCCTCCTTAG (60)

Supplementary References

1.   Sezonov G, Joseleau-Petit D, D'Ari R. 2007. Escherichia coli physiology in Luria-Bertani broth. J Bacteriol 189:8746–8749.

2.   Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H, Remmert M, Soding J, Thompson JD, Higgins DG. 2011. Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. Mol Syst Biol 7:539.

3.   Salgado H, Peralta-Gil M, Gama-Castro S, Santos-Zavaleta A, Muñiz-Rascado L, García-Sotelo S. J, Weiss V, Solano-Lira H, Martínez-Flores I, Medina-Rivera A, Salgado-Osorio G, Alquicira-Hernández S, Alquicira-Hernández K, López-Fuentes A, Porrón-Sotelo L, Huerta AM, Bonavides-Martí-nez C, Balderas-Martí-nez I. Y, Pannier L, Olvera M, Labastida A, Jiménez-Jacinto V, Vega-Alvarado L, del Moral-Chávez V, Hernández-Alvarez A, Morett E, Collado-Vides J. 2012. RegulonDB v8.0: omics data sets, evolutionary conservation, regulatory phrases, cross-validated gold standards and more. Nucleic Acids Res 41:D203–D213.

4.   Abo-Amer A, Munn J, Jackson K, Aktas M, Golby P, Kelly DJ, Andrews SC. 2004. DNA Interaction and Phosphotransfer of the C4-Dicarboxylate- Responsive DcuS-DcuR Two-Component Regulatory System from Escherichia coli. J Bacteriol 186:1879–1889.

5.   Kröger C, Dillon SC, Cameron ADS, Papenfort K, Sivasankaran SK, Hokamp K, Chao Y, Sittka A, Hébrard M, Händler K, Colgan A, Leekitcharoenphon P, Langridge GC, Lohan AJ, Loftus B, Lucchini S, Ussery DW, Dorman CJ, Thomson NR, Vogel J, Hinton JCD. 2012. The transcriptional landscape and small RNAs of Salmonella enterica serovar Typhimurium. Proc Natl Acad Sci 109:E1277–E1286.

6.   Hersch SJ, Wang M, Zou SB, Moon K-M, Foster LJ, Ibba M, Navarre WW. 2013. Divergent protein motifs direct elongation factor P-mediated translational regulation in Salmonella enterica and Escherichia coli. MBio 4.

7.   Miller SI, Mekalanos JJ. 1990. Constitutive expression of the phoP regulon attenuates Salmonella virulence and survival within macrophages. J Bacteriol 172:2485–2490.

8.   Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, Datsenko KA, Tomita M, Wanner BL, Mori H. 2006. Construction of Escherichia coli K-12 in-frame, single-gene knockout mutants: the Keio collection. Mol Syst Biol 2:2006.0008.

9.   Corcoran CP, Podkaminski D, Papenfort K, Urban JH, Hinton JCD, Vogel J. 2012. Superfolder GFP reporters validate diverse new mRNA targets of the classic porin regulator, MicF RNA. Mol Microbiol 84:428–445.