Type 2 Diabetes: One Disease, Many Pathways

Joon Ha

Arthur Sherman

Laboratory of Biological Modeling

National Institutes of Health

Bethesda, MD USA

Abstract (249 words)

Diabetes is a chronic, progressive disease that calls for longitudinal data and analysis. We introduce a

longitudinal mathematical model that is capable of representing the metabolic state of an individual at any

point in time during their progression from normal glucose tolerance to type 2 diabetes (T2D) over a

period of years. As an application of the model, we account for the diversity of pathways typically

followed, focusing on two extreme alternatives, one that goes through impaired fasting glucose (IFG)

first, and one that goes through impaired glucose tolerance (IGT) first. These two pathways are widely

recognized to stem from distinct metabolic abnormalities in hepatic glucose production and peripheral

glucose uptake, respectively. We confirm this but go beyond to show that IFG and IGT lie on a

continuum ranging from high hepatic insulin resistance and low peripheral insulin resistance to low

hepatic resistance and high peripheral resistance. We show that IFG generally incurs IGT, and IGT

generally incurs IFG on the way to T2D, highlighting the difference between innate and acquired defects

and the need to assess patients early to determine their underlying primary impairment and appropriately

target therapy. We also consider other mechanisms, showing that IFG can result from impaired insulin

secretion, that non-insulin dependent glucose uptake can also mediate or interact with these pathways, and

that impaired incretin signaling can accelerate T2D progression. We consider whether hyperinsulinemia

can cause insulin resistance in addition to being a response to it and suggest that this is a minor effect.

Introduction

28
29
Diabetes is by definition a state of hyperglycemia, but its natural history is diverse. For example, some

individuals experience fasting hyperglycemia first (Impaired Fasting Glucose, IFG), followed by

hyperglycemia at the two-hour time point (2hPG) of an oral glucose tolerance test (OGTT), defined as

Impaired Glucose Tolerance (IGT), and some experience these in the opposite order. Eventually, all

people with diabetes will have both IFG and IGT (combined glucose impairment, CGI), so we refer to

these two pathways as "IGT-first" and "IFG-first", respectively. An important implication of these

observations is that the best period for determining differences in the underlying physiology of these

pathways is during the pre-diabetic stage, when the phenotypes are still distinct.

38
Prediabetes is also the stage in which progression to type 2 diabetes (T2D) can be markedly delayed or

prevented (45), and interventions can plausibly be made even more effective by targeting the specific

metabolic defects of the patient. For example, IFG is generally thought to reflect insulin resistance at the

liver, resulting in elevated hepatic glucose production (HGP), whereas IGT is thought to reflect peripheral

insulin resistance, mainly in muscle, resulting in reduced glucose disposal. One would like to know

whether using drugs that primarily affect hepatic or peripheral insulin resistance makes a material

difference for patients on the IFG-first or IGT-first pathways, and whether any such benefit carries over

once T2D has begun.

47
Moreover, diabetes is not a state that one enters and exits, like an infection, but a chronic condition that is

the culmination of a series of gradual changes. Understanding of this progression is best obtained by

longitudinal studies over a period of years or decades. Here we will focus on one such study, the

Baltimore Longitudinal Study of Aging (BLSA) (51), which asked whether CGI is an obligatory stage

between IFG and T2D and between IGT and T2D (Supplemental Fig. S1; all supplemental material is

available at Figshare, https://doi.org/10.6084/m9.figshare.10792412).

These features of diabetes suggest that a longitudinal mathematical model for disease progression could

be a valuable adjunct to clinical studies. Here we establish such a model and demonstrate that it can

distinguish hepatic and peripheral insulin resistance and can simulate the IGT-first and IFG-first pathways

to T2D over a period of years. We will use the model to address the clinical questions raised by the

BLSA and Perrault studies. By tracking virtual patients continuously in time, the model can interpolate

between clinical observations, which are necessarily sparsely sampled, and indicate what is likely to have

happened in the interim.

The model provides broader insight into the different insulin-resistance phenotypes, showing that a wide

array of pathways, ranging from isolated IFG to isolated IGT to combinations of the two, can be obtained

by combining different degrees of hepatic and peripheral insulin resistance. Thus, the apparent clinical

diversity of individual paths comprises a set of quantitative variants within a unified process of metabolic

dysfunction.

There is also diversity in the relative contributions of insulin resistance and beta-cell dysfunction to

diabetes pathogenesis. This is particularly apparent in comparing diabetes risk factors in populations with

a high prevalence of obesity, such as those of African descent and Native Americans, to populations in

which diabetes risk is seen among lean individuals and beta-cell function is weaker, such as South and

East Asians; populations of European descent tend to lie in between (46). Furthermore, beta-cell

dysfunction can be subdivided into defects in first- and second-phase insulin secretion. As a first

installment on this large and complex set of problems, we examine whether the loss of first-phase

secretion early in pre-diabetes plays a fundamental causal role in T2D development or is just a useful

marker.

General modeling approach

The diabetes field is fortunate to have a strong tradition of mathematical modeling (20, 49). A common feature of those models is that they provide snapshots in time of the metabolic state of an individual, including insulin resistance and beta-cell function. These models have been used to track progression by taking a series of snapshots, but they do not contain mechanisms of progression and do not describe trajectories of progression.

The model developed here belongs to a different family of models that seek to explain disease progression mechanistically, rather than assess the current state. It is an extension of our previously published model of beta-cell mass and function, which successfully simulated progression to diabetes in rodents over months and humans over years (37). That model, in turn, was based on the seminal model of (75), which expressed mathematically the hypothesis that beta-cell mass provides negative feedback on a slow time scale to compensate for insulin resistance. If that compensation is inadequate, however, the toxic effects of very high glucose overcome the stimulatory effects of moderately elevated glucose. The normal homeostatic negative feedback is converted to positive feedback, leading to deterioration in glucose tolerance and culminating in diabetes. This fundamental concept has been incorporated into other models that broadly agree but emphasize different details (23, 24, 35, 38, 43, 79). Notably, the model of (38) was shown to be able to account for the progression of fasting hyperglycemia and T2D observed in the Diabetes Prevention Program (DPP)(45).

A third class of modeling studies has fit longitudinal data from clinical studies with non-linear mixed effects statistical approaches to assess the magnitude of treatment effects on glucose, insulin and HbA1c (18, 25) or, by fitting to a modified form of the model of (75), on beta-cell mass and insulin sensitivity (67).

See (24) for further comment on the models cited here and other models, and see (54) for a perspective on

modeling T2D.

The initial wave of mechanistic longitudinal models simulated fasting or average daily glucose and were

therefore unable to describe post-prandial responses or responses to glucose challenges such as OGTTs

and IVGTTs. A recent model (24) added the capability of following daily and post-challenge glucose

variations and was fit to OGTT data from the DPP. It also introduced a distinction between peripheral

and hepatic insulin resistance in order to account for the effects of drug and lifestyle interventions on

these parameters.

We also, using a different methodology, introduce simulations of OGTTs at selected points during

progression, and separate representations for hepatic and peripheral insulin resistance. We use these new

features to differentiate progression by either 2hPG or FPG. We perform IVGTTs as well to illustrate the

evolution of the acute insulin response to glucose (AIRg), often used to assess beta-cell function. This

requires the model to simulate first and second phase secretion, which we accomplish by incorporating a

previously published model for insulin granule exocytosis (17).

This study focuses on mechanism and insight, rather than assessment. Instead of fitting parameters to

particular data sets, we assume parameters and investigate the trajectories of glycemia and insulin

secretion that result. We demonstrate that the model captures known features of diabetes pathogenesis

data and provides novel insights and interpretations of the data. We conceptualize this as building a

factory, not a product. Once the utility of the model is established, we expect that a wide variety of

applications to clinical data will become possible.

Materials and Methods

We briefly review the previous version of the model (37) and then describe the enhancements introduced

here. The enhanced version has been used to study clinical implications of differences in glucose time

courses during an OGTT (19), but was not documented in detail. Parameter values and details of

functions not given here are in the Appendix (Supplementary Material).

Previous version of the model

The core element, retained in the new version, is a glucose-insulin feedback loop, represented by two

differential equations adapted from the Minimal Model of Bergman and Cobelli (8), who built on models

going back to the 1960's (2, 11), as modified in (75):

$\frac{dG}{dt} = MEAL + HGP - (E_{GO} + S_I I)G$ (1)

$\frac{dI}{dt} = \frac{\beta}{V} ISR - kI$ (2)

The glucose (G) equation (Eq. 1) says that G increases on a time scale of hours as a result of meal influx

and hepatic glucose production (HGP) and decreases as a result of uptake, which has both insulin-

independent and insulin-dependent components. The factor S_I in the insulin-dependent term is closely

related to the well-known sensitivity to insulin reported by the Minimal Model. The insulin (I) equation

(Eq. 2) says that I decreases due to removal, mainly in the liver, with rate constant k, and increases due to

secretion by beta cells, where β is the beta-cell mass, ISR is the insulin secretion rate per unit mass, and V

is the volume of distribution. In the original version, ISR depended only on G, through the rate of beta-

cell metabolism M:

$ISR = \sigma \frac{(M+\gamma)^{kISR}}{\alpha_{ISR}^{kISR} + (M+\gamma)^{kISR}}$ (3)

where M was assumed to be a sigmoidally-increasing function of G:

$M = \frac{G^{kM}}{\alpha_M^{kM} + G^{kM}}$ (4)

The parameter γ in Eq. (3) represents the effect of K(ATP) channel density to shift the glucose

dependence of secretion (the triggering pathway (39)); when the channel density is low, γ is high, and

shifts the dependence to the left, increasing secretion for the same level of M because Ca^{2+} is higher.

Experiments in (34) showed that mouse beta cells in vitro adjust the K(ATP) channel density down in

response to sustained (overnight) elevated glucose. This has also been observed in vivo in humans (41,

42, 44), along with evidence of reduced insulin clearance, k, (19–21). For the most part, we omit this for

simplicity, but we consider the implications of reduced clearance towards the end of Results. This can be

viewed as the first line of defense through enhanced beta-cell function against insulin resistance over a

time scale of days (e.g. holiday overeating).

The value of γ depends on glucose, which is taken into account by adding a third differential equation to

the system:

$$\frac{d\gamma}{dt} = \frac{\gamma_\infty(G) - \gamma}{\tau_\gamma} \tag{5}$$

where γ_∞ is an increasing sigmoidal function of G, and τ_γ is the time constant.

Insulin resistance that persists over longer periods (months in humans) despite reduced K(ATP) channel

density, is assumed to trigger a further level of compensatory increased beta-cell function via σ, the

maximal insulin secretion capacity (Eq. 3). This corresponds to the amplifying effects of metabolism

and/or modulators such as GLP-1 and ACh on the efficacy of Ca^{2+} to drive insulin granule exocytosis.

This second aspect of beta-cell functional compensation entails a fourth differential equation:

$$\frac{d\sigma}{dt} = \frac{\sigma_\infty(ISR, M) - \sigma}{\tau_\sigma} \tag{6}$$

We assume that increased *ISR* (workload in the sense of (64)) leads to an increase in σ whereas

increased *M* leads to a decrease in σ.

The slowest and final form of compensation for insulin resistance is increased beta-cell mass, β, which

develops over years in humans. We assume that β is increased by proliferation, *P*, and decreased by

apoptosis, *A*. Following the data of (64), we assume that *P* increases when *ISR* increases. We further

assume that apoptosis is largely driven by metabolic stress (e.g. through increased production of reactive

oxygen species) when glucose is high, so we make *A* an increasing function of *M* :

$$\frac{d\beta}{dt} = \frac{(P(ISR) - A(M))\beta}{\tau_\beta}$$ (7)

The parameters defining *P* and *A* are chosen such that modest increases in *G* result in a net increase in β,

but large increases in *G* result in a net decrease in β. As in the predecessor model (75), this leads to a

shift from compensation (negative feedback) to decompensation (positive feedback). In (37), we showed

that this can account for the threshold behavior observed in both rodents and humans, that is, nearly

steady *G* followed by a sharp, essentially irreversible increase (50, 73). We proposed this as an

explanation for why prevention of T2D is much easier than reversing it once it is established. The same

dynamic properties carry over in this study with the model enhanced as described next.

New features in the model

Modeling glucose flux during daily meals and glucose tolerance tests

In the previous version of the model (37), *G* represented average daily glucose and insulin levels in

response to steady glucose input. To address IFG and IGT, we need to be able to dissociate fasting

glucose from post-challenge glucose. The first step is to introduce variable glucose influx from meals,

represented by the term *MEAL* in Eq. (1). Timing of meals is standardized to 6:00 AM, 12:00 Noon, and

6:00 PM. The expression we use is:

$MEAL = meal_{max} \left(\frac{t^{mk}}{\alpha_{meal}+t^{mk}} \right)^{\eta} \exp(-\mu t)$ (8)

and Fig. S3 shows the glucose flux (panel A) and the resulting plasma glucose concentrations (panel B).

We do not yet account for other nutrients (amino acids, fats). Flux during an OGTT is modeled with a

more rapid rise and decay of flux (Eq. (A2) and shown in Fig. S3, C, D). Flux during an IVGTT rises and

decays still more rapidly.

Modeling hepatic glucose production (HGP)

In order to distinguish peripheral and hepatic insulin resistance and describe how they are related to each

other, we need to refine the model description of hepatic glucose production (*HGP* in Eq. 1). In the first

version of the model (37), *HGP* was assumed to be constant, which is an acceptable approximation as

long as fasting plasma insulin is adequate to compensate completely for any hepatic insulin resistance.

To study the failure of compensation, however, we need to make *HGP* a decreasing function of *I*:

208                 $HGP = \frac{hepa_{max}}{\alpha_{HGP} + hepa_{SI}I} + HGP_{bas}$ (9)

209   This is sufficient to give the typical drop and recovery of *HGP* after a meal (Fig. S4A).

210   Less obvious, but also important, we need to account for the correlation between hepatic and peripheral

211   insulin resistance, which we do by making two of the parameters in Eq. 8, *hepa_{max}* and α_{HGP}, functions of

212   S_I:

213                 $hepa_{max} = \frac{hepa_{bar}}{hepa_{k}+S_{I}} + hepa_{sh}$ (10)

$$\alpha_{HGP} = \frac{\alpha_{bar}}{\alpha_k + S_I} + \alpha_{sh} \qquad (11)$$

Both $hepa_{max}$ and α_{HGP} decrease with S_I, as shown in Figs. S4B, C, and the net effect is that *HGP*

decreases as S_I increases (Fig. S4D).

If this, or something like this, is not done, then severe peripheral resistance combined with strong

compensatory insulin secretion can result in fasting hypoglycemia, which is not the typical pattern (see

Fig. S5).

In more typical cases of progression to pre-diabetes or diabetes, the relative impairment in insulin

secretion would mask this effect: the level of glycemia would be reduced but hypoglycemia would not

result. We chose the parameters such that HGP remains normal when insulin is elevated unless there is a

defect in beta-cell mass or function.

Figure S6, E – H shows the response of HGP and glucose disposal to a simulated hyperinsulinemic,

euglycemic clamp, with steady-state values in good agreement with experimental data (6).

To represent hepatic insulin resistance over and above the component related to peripheral insulin

resistance, we decrease the parameter *hepa$_{SI}$* in Eq. A4, which increases HGP at any value of *I*.

Equations 9 – 11 are phenomenological expressions that achieve the goals of making *HGP* decrease with

I and increase when insulin resistance is present, but other expressions could also possibly work. We

considered making the value of *hepa$_{SI}$* depend on *I* as an alternative way to avoid hypoglycemia, but this

did not work well. It is also possible that glucagon, which is not in the model, would help to prevent

hypoglycemia (see *Limitations of the study* in the Discussion).

Modeling insulin granule exocytosis

To study the dynamics of glucose and insulin under glucose challenges such as meals, OGTT, and

IVGTT, the model needs to account for the multiple kinetic components of insulin secretion. We adapted

an existing model of insulin granule exocytosis (17), which was designed to capture the biphasic pattern

of ISR in response to a glucose step in vitro or a hyperglycemic clamp in vivo. The first phase is

characterized by a sharp peak of ISR during the first 10 minutes and the second phase by a steady increase

of ISR over the next hour. Figure S6 shows a simulated OGTT (panels A, B) and a simulated IVGTT

(panels, C, D) compared to experimental data.

These phases are mediated by progression of vesicles through a sequence of stages culminating in

exocytosis (fusion with the plasma membrane and release of insulin to the circulation; Fig. S2A). A large

reserve pool, treated as an inexhaustible reservoir, feeds the docked pool (vesicles at binding sites on the

plasma membrane). Once docked, vesicles are primed and enter the readily releasable pool (RRP).

Primed vesicles join the immediately releasable pool by becoming closely associated with voltage-

dependent Ca^{2+} channels. First-phase secretion depends mainly on the size of the RRP at basal glucose,

and second phase secretion is controlled by the rate of mobilization of vesicles to the docked pool.

250
The insulin secretion rate ISR in the first version (37) of the model (Eq. 3) is in the new version no longer

a function of glucose, but is calculated as an output of the exocytosis model (Eqs. A11), ISR thus now

depends on the history of exposure to G, as it should, not just the current value. The exocytosis model

requires as input the cytosolic Ca^{2+} concentration, which is modeled as a sigmoidal function of the beta-

cell metabolic rate M (Eq. A7), and the much higher Ca^{2+} concentration in the microdomains of Ca^{2+}

channels, which is modeled as a function of cytosolic Ca^{2+} (Eq. A8). The dose response curve shift γ,

previously included in Eq. 3, represents the dynamic changes in K(ATP) channel density as before, but

now explicitly alters cytosolic Ca^{2+} at a given level of M (Eq. A7). Cytosolic calcium enhances the rates

of mobilization of the reserve pool to the plasma membrane (Eq. A10) and priming of docked vesicles (r_2

in Eqs. A12), whereas vesicle fusion is primarily controlled by microdomain Ca^{2+}. The amplifying effect

of glucose (39) is incorporated as a multiplicative factor in the rate of vesicle mobilization (G_F in Eq. A9).

The effect of the incretin GLP-1and drugs that mimic it to increase insulin secretion is modeled as

enhancing the amplifying effect of glucose by increasing the parameter G_{Fmax} in Eq. A9 and as enhancing

vesicle priming by increasing the parameter r_2^0 in Eq. A12. This is in accord with a key feature of these

agonists, that they have little or no effect on secretion in the absence of glucose, minimizing the risk of

hypoglycemia that plagues sulfonylureas. Fig. S2C shows the effect of a two-fold increase in the incretin

effect on insulin secretion rate during a hyperglycemic clamp, in agreement with the effect of the GLP-1

receptor agonist liraglutide in the RISE study (68). Modeling the effect of incretins on glucose is more

difficult, as it would require including their effects to inhibit glucagon secretion or promote weight loss,

which is not feasible in the current model.

For IVGTT and hyperglycemic clamp simulations, we reduce G_F by about a factor of two to represent the

lack of the incretin effect on vesicle mobilization and reduce r_2^0 (Eq. A12) about 10-fold to represent the

lack of incretin effect on vesicle priming. The rate of mobilization (Eq. A11) is also assumed to be

proportional to the variable σ (Eq. 6), which thus controls the magnitude of second-phase insulin

secretion. As a consequence, *ISR* implicitly includes σ as a multiplicative factor, as in Eq. (3) of the

simpler, first version of the model (37), and the dynamic evolution of the maximal secretory capacity over

long time scales (months) is essentially equivalent.

Equations for insulin sensitivity

With one exception we model peripheral insulin sensitivity as an intrinsic property of the target tissues,

muscle, liver and adipose, determined implicitly by obesity and genetics. In the model, this makes it an

external, time-dependent parameter of the system, and we examine how hyperinsulinemia compensates

for insulin resistance. We model insulin sensitivity, S_I, as an exponentially decreasing function of time,

following (75):

$$\frac{dS_I}{dt} = \frac{(tar_{SI} - S_I)}{\tau_{SI}} \qquad (12)$$

In Fig. 9, we consider a hypothesis under current discussion in the literature (35, 65, 71, 74), that insulin

resistance may be caused by hyperinsulinemia, rather than the other way around. We think it is

implausible that this is the sole cause of insulin resistance, and propose instead that the core component of

insulin resistance is due to obesity and genetics but modified by the level of insulin I as a result of the

negative feedback generated in the insulin signaling cascade (70). We represent the effect, which we term

induced resistance, as follows:

$$S_I = S_{I,0}E \qquad (13)$$

293 where

$$\frac{dS_{I,0}}{dt} = \frac{(tar_{SI} - S_{I,0})}{\tau_{SI}} \qquad (14)$$

295 and

$$\frac{dE}{dt} = \frac{(E_\infty(I) - E)}{\tau_E} \qquad (15)$$

297 Equation (15) says that E relaxes towards E_∞ with a time constant τ_E, where E_∞ is a sigmoidally

298 decreasing function of I,

$$E_\infty(I) = \bar{E}\left[I - \frac{(I - I_{sh})^{kE}}{\alpha_E^{kE} + (I - I_{sh})^{kE}}\right] + E_b, \qquad (16)$$

several examples of which are shown in Fig. 9F. The time constant τ_E is set to two days so that E, and

hence S_I, will respond to sustained hyperinsulinemia, not to normal insulin fluctuations following meals

or glucose tolerance tests. The effect is propagated into hepatic insulin resistance through the terms

containing S_I. Other parameters are in the legend for Fig. 9 and Table S17.

Criteria of pre-diabetes and diabetes

Following ADA criteria, we define IFG as FPG > 100 mg/dl but < 126 mg/dl, IGT as 2hPG > 140 mg/dl

but < 200 mg/dl, and T2D as FPG ≥ 126 mg/dl or 2hPG ≥ 200 mg/dl. Combined glucose impairment

(CGI) is defined as co-occurrence of IFG and IGT.

Software

The model equations are solved using xppaut (5) and Matlab (The Mathworks, Natick, MA). Input files

defining the parameters and initial conditions are available on Figshare at:

https://doi.org/10.6084/m9.figshare.10792412).

Results

In Figs. 1 – 4, we carry out longitudinal simulations over a period of five years starting in the NGT state

in which either peripheral insulin resistance is dominant, in which case the first stage of hyperglycemia is

IGT, or hepatic insulin resistance is dominant, in which case the first stage of hyperglycemia is IFG.

These assumptions are applied by modeling peripheral and hepatic insulin sensitivity as exponentially

decreasing, using Eqs. A13 and A14, respectively. The initial values and rates of decline are the same for

all figures, but the steady state (target) values, tar_{SI} and tar_{hepaSI}, are varied. The simulations calculate the

daily responses to meals, but we plot only the results of OGTTs carried out periodically over the five-year

time span by pausing the longitudinal simulation. The corresponding peak postprandial glucose is shown

in Fig. S7. The capacity of beta-cell function to compensate for insulin resistance is assumed to be

limited, except in Fig. 2.   The defect consists of a right shift in γ_∞ relative to Fig. 2, which can be

interpreted as a slight gain of function mutation in KCNJ11, the Kir6.2 component of the KATP channels.

Alterations in σ_∞ representing a mild defect in insulin granule mobilization would have a similar effect.

Parameters that were varied to make the figures are listed in Table S12A in the Appendix, and the initial

conditions for Figs. 1 – 4 are in Table S12B.

IGT-first pathway

Figure 1 shows a longitudinal simulation of the effects of a strong decrease in peripheral insulin

sensitivity S_I (Fig. 1A) combined with a mild decrease in hepatic insulin sensitivity, *hepa$_{SI}$* (Fig. 1B). As

insulin resistance progresses, 2hPG increases rapidly while FPG increases more slowly (Fig. 1C),

resulting in progression from normal glucose tolerance (NGT) to IGT, CGI and ultimately T2D (Fig. 1C).

Fasting plasma insulin (FPI) and 2-hour plasma insulin (2hPI) rise as the beta cells initially compensate

partially for the insulin resistance, then fall as the beta cells fail (Fig. 1D), following the classic "Starling

law" of the pancreas (27). The initial rise in secretion results from an increase in beta-cell sensitivity to

glucose (the variable γ increases, not shown), and the decline results from a fall in the slower component

of beta-cell function, σ (Fig. 1E).   Beta-cell mass (β) also rises and falls, but the variation is limited

because of the slowness of β, and the fall occurs only after T2D is already underway (Fig. 1F). This

accords with observations that beta-cell mass is elevated in insulin-resistant pre-diabetics but reduced in

long-standing diabetes (15, 66, 69).

A subtle but important point of this simulation is that insulin resistance in both the liver and peripheral

tissues reaches saturation (Fig. 1A, B) well before the advent of T2D. It is rather the continuing fall in

beta-cell function, σ, that drives conversion to T2D. The same sequence was seen in the simulation of

T2D progression in Zucker diabetic fatty rats (Fig. 6 in (37)) and in data from monkeys (10).

The fall in σ is triggered by the hyperglycemia and glucotoxicity that follows the early loss of insulin

sensitivity (Fig. 1D) but would not lead to T2D if the pre-existing capacity of beta-cell function to

compensate were stronger. This is illustrated by a simulation with the same degree of insulin resistance

as in Fig. 1, but a milder beta-cell defect, which mimics a non-diabetic subject with insulin resistance

(Fig. 2). FPG and 2hPG increase modestly in response to insulin resistance and reach a plateau in the

IGT state as σ levels off (Fig. 2E) and plasma insulin plateaus (Fig. 2D).   The limitation of the rise in

glucose reduces the effects of glucotoxicity and buys time for beta-cell mass to increase and stabilize the

IGT state (Fig. 2F).

Figures 1 and 2 together paint a picture in which not only is a combination of insulin resistance and

impaired secretion necessary for T2D, but insulin resistance develops and saturates first, and T2D

develops only if the beta cells fail. Some defect in beta-cell function is required even for pre-diabetes, in

agreement with (28). Conversely, a pre-existing defect in insulin secretion would be silent in the absence

of insulin resistance (Fig. S8), unless it were severe (Fig. S9).

IFG-first pathway

In Fig. 3 we illustrate a contrasting case to Fig. 1, dominant hepatic insulin resistance with minor

peripheral resistance (Fig. 3A, B); the same beta-cell function defect is assumed as in Fig. 1. Hepatic

insulin resistance drives FPG across the threshold for IFG, while 2hPG remains below the threshold for

IGT (Fig. 3C). After the initial threshold crossing, however, FPG and 2hPG continue to rise, and IFG

progresses to CGI as in Fig. 1. Insulin again rises with the help of γ (not shown) and falls when the drop

in σ becomes too great (Fig. 3E). As in the IGT-first pathway, the conversion to T2D is driven mainly

by reduced beta-cell function, σ, because insulin resistance in both the liver and peripheral tissues

saturates well before T2D (or even CGI) begins. Beta-cell mass again plays a minor role (Fig. 3F).

In the BLSA some subjects who had progressed from NGT to IGT went on to T2D at the next follow-up,

which prompted the authors to ask whether CGI could be skipped (51). Figure 4 demonstrates that this

can happen if peripheral insulin resistance is made much greater than hepatic resistance (compare Fig. 4A

to Fig. 1A). Extreme loss of peripheral insulin sensitivity causes 2hPG to rise dramatically, while FPG

remains in the normal range, converting NGT to IGT. 2hPG continues to deteriorate without a substantial

increase in FPG, resulting in progression of IGT to T2D without passing through CGI. Eventually, FPG

crosses the thresholds for IFG and T2D, but after the individual has already reached T2D based on 2hPG.

Figures 1 and 3, respectively, consider extreme cases of peripheral insulin resistance (PIR), where S_I

dominates, and hepatic insulin resistance (HIR), where $hepa_{SI}$ dominates . However, most people on the

path to T2D will have both. A composite view of the cases of Figs. 1 and 3 together with intermediate

phenotypes is given in Fig. 5, where trajectories are plotted in the FPG – 2hPG plane. The only

difference among the trajectories is the degree of HIR and PIR, varied inversely from lower right to upper

left; the compensatory capacity of beta-cell function is identical for all traces. Fig. 5A shows that the

trajectories diverge markedly as the defects in HIR and PIR set in but converge as hyperglycemia

worsens; this happens because IFG induces IGT and IGT induces IFG. Thus, all the virtual patients end

up looking the same as time goes on. The latter part of the NGT stage and the early part of the IGT stage,

shown expanded in Fig. 5B, are when the underlying pathologies give rise to the most distinct behavior.

This is important both for designing clinical studies and for stratifying patients for treatment. The figure

suggests that the slope of the trajectory from two or more OGTTs spaced suitably far apart in time could

give a good indication of the future path of the patient.

We next look more closely at the pathogenesis process as it would appear clinically by simulating OGTTs

(Fig. 6) and IVGTTs (Fig. 7) at representative times for each stage of glucose tolerance, indicated by the

black circles on the time axes in Figs. 1C and 3C.

Simulation of OGTTs

Figure 6A, C shows representative glucose and insulin profiles during OGTTs at each stage of the IGT-

first pathway of Fig. 1. Insulin concentrations at the IGT stage (Fig. 6C, dotted curve) are increased

compared to NGT (Fig. 6C, solid curve) but are inadequate to maintain normoglycemia at two-hours

because of the decreased peripheral insulin sensitivity (Fig. 1A). In contrast, the level of fasting insulin at

IGT (Fig. 6C, dotted curve) is sufficient to maintain fasting glucose within the normal range because

hepatic insulin resistance is relatively mild (Fig. 1B). During CGI, glucose (Fig. 6A, dashed curve) is

increased at all time points compared to IGT, while the insulin level (Fig. 6C, dashed curve) is slightly

diminished at the early time points (relative to glucose, however, secretion is impaired at all time points).

This indicates that even progression to CGI from IGT is mainly due to impaired beta-cell function.

Further and more marked decreases in insulin, due to impaired secretion at all time points during the

OGTT, lead to diabetes (dot-dashed curve). Thus, during the progression towards T2D, insulin at all time

points during the OGTT first increases and then decreases, in accord with the Starling Law (27).

Figure 6B, D shows glucose and insulin at each stage of the IFG-first pathway of Fig. 3. Severe hepatic

insulin resistance increases glucose at all time points during the OGTT (Fig. 6B). The increase in FPG is

greater than for 2hPG, so the threshold for IFG is crossed first in Fig. 3. Even though fasting insulin at

the IFG stage (Fig. 6D, dotted curve) is very high, it is not enough to maintain normal FPG, because of

the severe hepatic insulin resistance (Fig. 3B). However, 2hPG is maintained in the normal range because

peripheral insulin resistance is mild (Fig. 3A). Since both peripheral and hepatic insulin sensitivity

saturate before the onset of CGI (Fig. 3A, B), the decrease in insulin at all time points during the OGTT

(Fig. 6D, dashed curve) due to falling beta-cell function (Fig. 3E) is the main contributor to the

progression to CGI from IFG and then to T2D.

Simulation of IVGTTs

Figures 1 – 6 highlight the importance of secretion defects, in the context of insulin resistance, in all the

pathways to T2D, but now we break out the contributions of early vs. late secretion. First-phase secretion

is widely considered a key early marker of future progress. For example, the classic paper (13) reported a

cross-sectional study of IVGTTs, and showed that AIRg declines as FPG rises and is nearly gone by the

time FPG reaches 115 mg/dl, well below the threshold for T2D. This supports the use of AIRg, and by

implication first-phase insulin secretion, as an early marker for T2D. We now show that the model can

reproduce the negative correlation between FPG and AIRg, but that rising FPG is not necessarily the sole

or proximal cause of the decline in AIRg.

Figures 7A, B show simulations of the insulin responses during IVGTTs performed during the IGT-first

and IFG-first pathways, respectively. AIRg is blunted and then vanishes in both pathways as FPG rises,

as found in (13), but 2hPG also rises at the same time, albeit to different degrees relative to FPG in the

two pathways. The decline of AIRg is more rapid than seen experimentally along both pathways (14), but

the main point is that it results from the decline of RRP size (Figs. 7C, D), which is more fundamental

than the level of glycemia, as the next two paragraphs explain.

RRP size is controlled by two factors, the rate of secretion, which determines the rate of vesicle efflux

from the RRP and the rate of vesicle influx into the RRP from the docked pool. High FPG increases the

rate of efflux in the basal state, so the RRP will already be depleted when the IVGTT commences, and

AIRg will consequently be reduced. The rate of influx depends on the size of the docked pool and the

rate of priming of docked vesicles. The rate of priming does not vary much in our simulations, but the

size of the docked pool does, depending mainly on the rate of docking, which is proportional to one of our

beta-cell function variables, σ. Although modest increases in glucose stimulate vesicle docking, larger

increases cause glucose toxicity, which reduces σ and hence docking.

Because σ is slow, it responds to the average daily glucose, including the contributions of both fasting

and post-prandial glucose, which differ according to pathway. Along the IGT-first pathway (Figs. 1 and

7A, C), AIRg initially declines primarily because of reduced σ (Fig. 1E), which is mainly determined by

high post-prandial glucose, rather than FPG. Positive feedback in turn drives post-prandial glucose

higher as σ decreases (Figs. 1C, E). Thus, the loss of AIRg and first-phase secretion is an indirect

consequence of diminished second-phase secretion capacity. As the subject progresses to CGI, FPG also

rises and further reduces AIRg due to pool depletion. In contrast, during the IFG-first pathway (Figs. 3

and 7B, D), the early rise of FPG reduces AIRg by depleting the RRP, and positive feedback from the

later rise of 2hPG further diminishes AIRg by driving down σ.

In summary, reduced AIRg can be a marker of impairment in either first- or second-phase secretion, and,

as glycemia progresses towards T2D, is likely to indicate both.

Targeted Drug Therapy

With the previous, simpler model (37) we showed that NGT and T2D were bistable states separated by a

threshold. This accounted for the well-known observation that it is easier to prevent T2D than to reverse

it. The new model retains these characteristics but raises the possibility that the response to therapeutic

interventions may vary depending on which pathway a patient follows to T2D. Figure 8 shows that this is

indeed the case, and that knowledge of a patient's subtype of insulin resistance can in principle lead to

more effective drug therapy.

Figures 8A, B contrast two drug therapies targeted to peripheral vs. hepatic insulin resistance in patients

in the early stages of diabetes. Figure 8A shows glucose for a patient on the IGT-first pathway with

dominant peripheral insulin resistance as in Fig. 1, in the absence of therapy (control, black curves) or in

response to a high dose of a drug targeted to peripheral insulin resistance (dashed curves) or a drug

targeted to hepatic insulin resistance (dotted). The high dose of the appropriately targeted drug only

transiently improves FPG and 2hPG and ultimately fails to reverse T2D. Nonetheless, it is more effective

at delaying progression than the mistargeted drug.

Figure 8B represents the complementary case, a patient on the IFG-first pathway with dominant hepatic

insulin resistance, as in Fig. 3. The same treatments are applied, and, as in panel A, both drugs only

transiently improve FPG and 2hPG, but the appropriately targeted drug is more effective at delaying

progression.

Figures 8C, D show the same drug therapies and progression pathways as in panels A and B, but with the

drugs applied before the onset of diabetes. A low dose of a drug targeted to the patient's specific insulin

resistance pathology is in both cases now able to prevent progression to diabetes, while a low dose of a

mistargeted drug only delays progression. These examples suggest that the effectiveness of drug therapy

depends on both early initiation of treatment and detection of the major metabolic abnormality.

The study in (24) found that it was necessary to assume that the efficacy of treatment wanes with time in

order to fit the data from lifestyle and drug interventions in the DPP. Here we have shown that even if the

efficacy of treatment is maintained, the intrinsic dynamics of progressive beta-cell dysfunction can cause

treatment to fail.

Caution should be used in interpreting the simulated treatments in Fig. 8 in terms of currently used drugs.

For example, in the DREAM study (29) it was found that rosiglitazone was effective in cases of isolated

IFG (IIFG), that is IFG in the absence of IGT. Although rosiglitazone is often thought of as primarily

improving peripheral insulin sensitivity, it also improves hepatic insulin sensitivity (81). In addition,

IIFG does not necessarily imply pure hepatic insulin resistance. For example, a person with FPG = 115

and 2hPG = 135 would be classified as IIFG but may have significant peripheral insulin resistance. Most

individuals with pre-diabetes likely have a mixture of peripheral and hepatic insulin resistance.

Hepatic Insulin Clearance

Another way to generate a compensatory increase in insulin is to reduce clearance. There is considerable

evidence that insulin clearance is positively correlated with peripheral insulin sensitivity (3, 63), but the

direction of causation is not clear. If clearance is reduced secondary to reduced overall insulin sensitivity,

then the increased insulin would contribute to the compensatory response, along with increased secretion.

However, it has been suggested that the increased insulin resulting from reduced clearance may also

contribute to insulin resistance (61). In Fig. 9, we explore the consequences of reduced clearance,

assuming that the increased insulin reduces S_I is reduced by varying degrees as I is increased, as described

in Methods. We assume for simplicity that clearance itself is constant throughout each simulation rather

than adapting to glucose, insulin or insulin sensitivity. The control case, shown in black, is equivalent to

Fig. 1, an example of an IGT-first pathway to T2D. If clearance is reduced, and a modest degree of

additional insulin resistance is assumed (green curve in panel F), such that the S_I is only slightly lower

than the control case (panel A), then the progression to T2D is delayed (compare black and green curves

in panel C). (The smallness of the difference in S_I is due to the fact that glucose is elevated to levels

where the induced effect is triggered only for a brief period of each day.) Note that σ is slightly larger

(green vs. black, panel E), indicating that reducing clearance partially spared beta-cell function.

If the effect of insulin to induce resistance is made larger (blue curves), the beneficial effect of higher I

due to reduced clearance and the harmful effect of reduced insulin sensitivity approximately cancel out.

Finally, if the induced resistance effect is made extreme, such that even a slight increase in I reduces S_I in

half (red curves), the net effect is somewhat accelerated progression. We conclude that induced insulin

resistance makes at best a modest contribution to the progression to T2D. If an even more extreme

degree of induced resistance is assumed, such that even a slight increase in I reduces S_I to near 0, then it is

possible for frictional resistance to be the main driver of T2D, but we consider such a scenario

implausible.

Non-Insulin-Mediated Glucose Uptake

We have focused up to this point on the contributions of hepatic and peripheral insulin resistance in

driving the IFG-first and IGT-first pathways. However, non-insulin-mediated glucose uptake, also known

as glucose effectiveness, is comparable in magnitude to insulin-dependent uptake (9) and has been found

to be impaired in T2D (30). It is thus a potential target of drug therapy. We model this by varying the

model parameter E_{G0} (similar to S_g in Minimal Model analysis) in Eq. 1. In Fig. S10 we demonstrate that

cutting glucose effectiveness in half can result in T2D in someone otherwise not at risk, such as the case

shown in Fig. 2. Conversely, in Fig. S11 we show that doubling glucose effectiveness can prevent T2D

in someone otherwise predisposed, such as the case shown in Fig. 1.

Discussion

In recognition of the fundamental character of type 2 diabetes (T2D) as a progressive disease that

develops over many years, we have established a longitudinal model for its pathogenesis. We follow in

the footsteps of other longitudinal models (23, 24, 75) but offer new clinical applications and insights.

We apply the model to analyze the diverse presentation of hyperglycemia, which may manifest first in

fasting glucose (IFG-first pathway) or two-hour glucose during an OGTT (IGT-first pathway). To carry

out this program, we modified the representation for beta-cell response to insulin resistance in the first

version of the model (37), enhancing it to differentiate between hepatic and peripheral insulin resistance.

The simulations show that heterogeneity in the degree of the two forms of insulin resistance can account

for a wide variety of observed patterns, supporting the idea of T2D as a unitary disease with quantitative

variants. We have focused on extreme cases to highlight the differences (e.g. Fig. 1 vs. Fig. 3), but the

family of trajectories in the FPG-2hPG plane (Fig. 5) shows that these lie on a continuum. Figure 5 also

highlights that differences in insulin resistance phenotype are most evident in the late NGT and early pre-

diabetes stages, which are thus most amenable to differential phenotyping and therapeutic stratification.

We have incorporated a description of insulin granule dynamics sufficient to account for both first- and

second-phase insulin secretion. This made it possible to simulate OGTT and IVGTT time courses and

show how they are transformed systematically during progression along the two canonical pathways to

diabetes (Figs. 6, 7). We also showed that sufficiently strong beta-cell function can prevent T2D even

when insulin resistance is severe, allowing individuals to maintain a permanent state of IGT (Fig. 2) or

even revert from IGT to NGT (not shown). Conversely, sufficient insulin sensitivity can prevent T2D

even when beta-cell function is somewhat impaired (37). As discussed below under *Secretion defects*, a

fuller treatment of the differences in the balance of insulin secretion and insulin action defects is needed

to account fully for the diverse patterns of T2D progression.

We summarize below the specific lessons learned and questions answered by this study and give a

preview of the clinical applications we anticipate for the model.

Questions raised in the BLSA and other studies

550
The BLSA study (51) asked whether subjects who enter the IGT state necessarily pass through CGI on the

way to T2D. The model suggests (Fig. 4) that this is not the case, but skipping CGI happens only if the

peripheral insulin resistance is markedly greater than hepatic insulin resistance, which may be relevant for

adolescents, who experience extreme obesity and insulin resistance. An intermediate possibility predicted

by the model is a short, but not absent, interval of CGI that could escape detection if the follow-up

interval is too long.

A parallel question is whether individuals can go directly from IFG to T2D without passing through CGI.

This is harder than going directly from IGT to T2D because 2hPG is much more labile than FPG; it is

difficult to get an increase in FPG sufficient to cross the threshold for T2D (125 mg/dl) without at the

same time having 2hPG cross the threshold for IGT (140 mg/dl). Indeed, we have not been able to

simulate this scenario just by choosing an appropriate mixture of hepatic and peripheral insulin sensitivity

using the other parameters as in Figs. 1 – 4, but model simulations (not shown) predict that it can happen

if a more severe beta-cell defect (in γ_∞, Eq. A15) is assumed.

The BLSA study also asked whether individuals can pass directly from NGT to T2D without passing

through any pre-diabetic state. Because glucose is in quasi-steady state with the much slower variables

representing beta-cell mass, beta-cell function and insulin sensitivity, this is not possible in the model

unless one of those slow variables undergoes a catastrophic, virtually discontinuous, change.

Pancreatectomy would be an example of this, but even type 1 diabetes, triggered by a rapid fall in beta-

cell mass, has a distinct prediabetes phase.   The cases observed in the BLSA in which subjects were NGT

at baseline and T2D at first follow-up most likely reflected rapid progression or a long gap between visits.

Similarly, the model simulations indicate that it is unlikely for individuals to go from NGT to CGI

without passing through IFG or IGT.

The BLSA reported that IFG is generally followed by IGT and IGT is generally followed by IFG, and the

model suggests that each state *induces* the other. This happens because the initial rise in glucose during

IFG impairs beta-cell function, which causes 2hPG to rise, and vice versa. Another study raised the

question of whether CGI is a progressed state of IFG (57); the model simulations together with the BLSA

data show that CGI may instead be a progressed state of IGT, to which IFG has been added.

One can also ask whether crossing the threshold for FPG or 2hPG of pre-diabetes predicts whether T2D

will be reached by crossing the corresponding threshold.   In Fig. 1, IGT is followed by T2D diagnosed

through 2hPG, and further simulations with the model (Fig. 5) suggest that this is typical.   In Fig. 3, IFG

is followed by T2D diagnosed through FPG, but further simulations (not shown) indicate that this may or

may not be the case, depending on the degree of discrepancy between HIR and PIR and the strength of γ

to control FPG.

Both the IFG- and IGT-first pathways exhibit elevated fasting insulin. This may account for the

observation that elevated fasting insulin was a better predictor of future diabetes in a prospective study

than fasting glucose, which is elevated early on only in the IFG-first pathway (22); it may not be

necessary to hypothesize a major causative role for high fasting insulin itself. Indeed, in the IGT-first

pathway, which is more common, fasting glucose may be suppressed by the compensatory increase in

beta-cell function (our variable σ) induced by high post-load glucose (compare Figs. 1 and 4 to Fig. 3).

The suppression of fasting glucose in the context of a predominance of peripheral insulin resistance has

particular importance for pre-diabetes screening in populations that are prone to IGT but not IFG. This

applies notably to people of African descent, for whom fasting glucose has markedly reduced sensitivity

for detecting pre-diabetes and diabetes (77). The problem is exacerbated for Africans living in Africa,

where measuring 2hPG with OGTTs is prohibitively expensive. The model suggests that in this and

similar cases, lowering the threshold for diagnosing pre-diabetes based on fasting glucose could be a cost-

effective strategy. More generally, the model points to the need for population- and patient-specific

thresholds for diagnosis, which may contribute to resolving current debate on whether prediabetes is a

useful diagnosis (62).

Insulin action defects

We have varied peripheral and hepatic insulin resistance independently to study their contributions to

T2D progression, but in reality, they are related. Statistically, they are correlated with a coefficient of

about 0.7 (1). This is expected for several reasons. For one, they share major components of the insulin

signaling pathway. Also, there is evidence that an important determinant of hepatic insulin resistance is

excess supply of free fatty acid (FFA) substrate from adipose tissue (58). Thus, insulin resistance in

adipose cells would increase lipolysis and FFA flux to the liver, which would drive increased

gluconeogenesis (59). On the other hand, the liver has unique roles in glucose and lipid production not

shared with muscle, which may account for the fact that the correlation is imperfect.

To address hepatic insulin resistance above and beyond the component correlated with peripheral insulin

resistance, we have independently varied the affinity of HGP for insulin (parameter $hepa_{SI}$ in Eq. 9), the

effect of which is shown in Fig. S4. We have obtained similar results (not shown) by varying the

maximal rate of HGP (parameter $hepa_{max}$).

Although we have primarily treated insulin resistance as an external influence on the model, representing

genetic and obesity, we have considered the much discussed hypothesis (65, 71, 74) that hyperinsulinemia

can contribute to insulin resistance in addition to being a compensatory response to insulin resistance

(Fig. 9). Any effect of increased insulin to induce insulin resistance has to be weighed against the effect

to enhance glucose uptake, and the outcome would depend on the net effect on the term $S_I I$ in Eq. 1. We

find that incorporating induced insulin resistance can either advance or delay the rate of progression to

T2D modestly depending on the strength of the two effects. We have not found support for the

hypothesis that it is the sole or primary initiating factor: to trigger T2D in the absence of intrinsic,

insulin-independent insulin resistance, we had to assume that the insulin-induced resistance is implausibly

strong. Similar results would apply if hyperinsulinemia resulted from increased secretion rather than

reduced clearance, but one would also have to consider whether in this case the increased beta-cell

workload might have an additional deleterious effect.

We are aware of only one other modeling study (35) that has addressed the role of hyperinsulinemia as a

driver of diabetes pathogenesis. More specifically, that thoughtful analysis considered a variant of the

model of Topp et al (75) in which *hypersecretion* would reduce S_I. Under that assumption, and an

additional assumption that high insulin would impair beta-cell mass, hypersecretion was found to be able

to drive progression to T2D. However, increased insulin itself that has been shown to induce insulin

resistance (71), so we prefer to have insulin act directly on S_I. As we have said above, the effect of

hyperinsulinemia governed by that mechanism is modest.

The effects of induced insulin resistance are limited fundamentally because insulin both promotes glucose

uptake and contributes to resistance to that uptake. This is analogous to friction in mechanics. The

higher the insulin concentration, the more resistance is generated through increased negative feedback in

the insulin signaling cascade. Similarly, the faster a car goes, the more wind and rolling resistance it

generates. This resistance is not the motive force driving the car forward but a by-product of forward

motion that limits its speed. However, resistance is generally not strong enough to prevent acceleration, it

just requires more gas to achieve the desired speed. Similarly, the insulin resistance induced by insulin

can in most cases be overcome by secreting more insulin.

Insulin secretion defects

The simulations of progression to T2D (Figs. 1, 3, and 4) require some degree of secretion defect in

addition to the various combinations of insulin resistance. We assumed a defect only in the triggering

pathway (γ), but we have obtained similar results by assuming defects in the amplifying pathway (σ).

The assumed triggering defect represents a right shift in the glucose dose response curve, which

correspond to a mild gain of function (GOF) mutation of the K(ATP) channels, a prominent GWAS hit

(55).

Assuming a more significant GOF mutation defect in the model leads to T2D with a lesser degree of

insulin resistance (Fig. S8). This is in line with the finding that K(ATP) defects are generally mild (for

example, the E23K mutation in the Kir6.2 component of K(ATP), KCNJ11, has an odds ratio of only 1.18

(33)), but they are fairly common and therefore may make significant contributions on the population

level in combination with insulin action defects.

If the K(ATP) defect is sufficiently severe, no insulin resistance is needed at all (Fig. S9). This is the

case in neonatal diabetes mellitus (NDM) (4, 47), which is of particular interest because it can be treated

effectively with sulfonylureas. This differs from typical T2D in adults, possibly because the drug brings

the genetically insufficiently active beta cells to a normal level of activity instead of overdriving them.

The examples of Figs. 2, 1, S8 and S9, in that order, lie on a continuum of progressively greater K(ATP)

channel function and lesser insulin resistance. As we showed previously in Fig. 5 of Ref. #6, the greater

the degree of beta-cell dysfunction, the lesser the degree of insulin resistance required for T2D. In

addition to the degree of insulin resistance, the rate of decline of insulin sensitivity is important. Fig. S12

shows cases in which a slow decline is compensated whereas a fast decline is not, as shown previously in

(75). We emphasize that the requirement for insulin resistance for typical T2D progression is not in

contradiction with genome wide studies showing a predominance of beta-cell genes: most people who are

overweight or obese become insulin resistant, but only those with beta-cell defects become diabetic. Lean

individuals who are not insulin resistant can also develop diabetes, but only if they have severe and rare

secretion defects.

Although we have focused on the role of insulin resistance in promoting IFG, secretion defects can also

lead to IFG with little contribution of insulin resistance. Fig. S13 shows a case with a mild GOF mutation

in K(ATP) channels, and Fig. S14 shows a case with a small readily releasable pool due to reduced

vesicle priming rate. With more extreme secretion defects, no insulin resistance is needed at all (not

shown). These patterns may be of particular relevance to Asian and possibly African populations where

secretion defects play a bigger role than insulin resistance in diabetes compared to those of European

descent.

Another secretory defect that may contribute to diabetes risk is impaired incretin signaling. Figure S15

illustrates that such an impairment would accelerate progression in a person predisposed to T2D.

The model shows that defects in first- and second-phase secretion are not independent. Impairment in one

leads to impairment in the other because of the harmful effects of elevated glucose. The model accounts

for classic data (13) showing that the acute insulin response to glucose (AIRg), a surrogate for first phase

secretion, declines with even modest increases in FPG during the NGT and pre-diabetic stages. The

model shows, however, that second-phase insulin secretion declines in parallel, belying a privileged role

for first-phase secretion, as also shown in another modeling study (36). In our model, reduced AIRg

results mainly from reduced size of the RRP, which can be caused by impairment in either first-phase or

second-phase secretion or both. More generally, any unanswered rise in either fasting or post-prandial

glucose impairs both first and second phase secretion, and any defect in either first or second phase

secretion raises both fasting and post-prandial glucose. This creates a vicious cycle that drives down both

first- and second-phase secretion regardless of which defect is primary. Empirical studies that do not

select for early isolated IFG or IGT cases do not show a clear prominence of first phase secretion loss

(31), in agreement with the model.

Non-insulin mediated glucose uptake

We have also considered briefly glucose effectiveness, the ability of glucose to promote its own uptake

independent of insulin (Figs. S10, S11). For simplicity, we have assumed the differences in E_{G0} to be pre-

existing and constant in time. However, there is evidence that it is a regulated process; it has been

reported to be enhanced by FGF19 acting in the brain and may be related to the glucose intolerance

resulting from leptin deficiency in ob/ob mice (52). It may also account for the rescue by leptin of

glucose tolerance in insulin-deficient rodents treated with streptozotocin (32). Further progress in

modeling this process would benefit from identifying the molecular mechanism(s) underlying it, which

are currently not well understood.

Limitations of the study

The model as presented here is oversimplified in that it only considers glucotoxicity and neglects other

factors that are likely to play a role, such as lipotoxicity (48). We are thus unable at present to address the

possible role of hyperinsulinemia to promote hepatic fat accumulation (76). If prolonged, this could lead

to liver damage and thus to a worsening of insulin resistance that could outlast the induced resistance we

have modeled here (Eqs. 9 – 11), which would resolve if normal insulin levels were restored.

In addition, we believe that prolonged high secretion rate is probably harmful beyond the negative effect

incorporated in the equation for σ, possibly because of ER stress and/or calcium toxicity (53). Those

factors were not included here because they were not needed to account for the pathways considered, but

they may be necessary to explain other pathways to T2D and will be addressed elsewhere.

To capture the full range of patterns, it is necessary to consider as well pre-existing variation in beta-cell

function, not the moment-to-moment beta-cell function, which evolves in response to hyperglycemia, but

the innate, genetic capacity of beta-cell function to adapt to hyperglycemia. For example, changing the

parameters defining σ_∞ (Eqs. 6, A16) has marked effects on the speed of progression in IGT; we

neglected this because it doesn't change the likelihood of entering the IGT state much.

We have modeled the suppression of hepatic glucose production as a direct effect of insulin on the liver.

However, much if not all of the acute effect of insulin is indirect, mediated by suppression of lipolysis in

adipose tissue, which reduces the supply of free fatty acid (FFA) to the liver (58). We consider this an

acceptable approximation for our purposes, as post-prandial suppression of lipolysis is roughly a mirror

image of the post-prandial rise in insulin. In future, if we want to account for FFA dynamics or adipose-

tissue insulin resistance, the model would have to be augmented.

We have modeled insulin-dependent glucose uptake as linear in insulin, but it is likely to be non-linear

(sigmoidal) (12) and has been suggested to exhibit hysteresis (78). We have not found these features to

be necessary to explain the data under consideration, but they can be easily added in the future if the need

arises.

We have neglected the role of excess glucagon secretion, which results from intrinsic defects in alpha

cells or from loss of paracrine regulation of alpha cells due to a relative lack of insulin (21, 56). This has

long been recognized to exacerbate post-prandial hyperglycemia in diabetes and pre-diabetes and is a

target of GLP-1-based insulin secretagogues. Without glucagon in the model, we are not able to simulate

fully the efficacy of that class of drugs, only their effect on insulin secretion. Hyperglucagonemia may

also contribute to hepatic insulin resistance, as proposed in recent reviews (26, 80) suggesting that

glucagon is part of a liver-alpha cell axis that is primarily devoted to regulating amino acid metabolism.

A primary form of glucagon resistance at the liver in this cycle could lead to hyperaminoacidemia and

hyperglucagonemia. The latter would contribute to excess hepatic glucose production and IFG in

addition to or in place of hepatic insulin resistance. This makes glucagon a tempting therapeutic target.

Hyperaminoacidemia and alpha-cell hyperplasia have thwarted many efforts, but some work in this

direction continues (72).

Other recent developments have shown that glucagon amplifies insulin secretion and thus under normal

conditions has a net hypoglycemic effect post-prandially (16, 82). Glucagon is thus now seen to be

homeostatic in both low and high glucose under normal conditions but can become anti-homeostatic in

diabetes or pre-diabetes. The emerging story is complex, and inclusion in the model is deferred to future

studies.

We have included only a very simple representation of insulin clearance, assuming a first-order

dependence on insulin concentration. We have not distinguished portal from peripheral insulin (63) or

considered possible regulation by glucose (60) or free fatty acids (7).

Future directions

This paper has demonstrated that if the insulin resistance phenotype of an individual is known, their

future trajectory of hyperglycemia can be predicted (Figs. 1 – 4) and drug choice can potentially be

optimized for the patient's insulin resistance phenotype (Fig. 8). Another longitudinal model has shown

similarly the results of targeting insulin resistance vs. beta-cell replication (38). Our model can provide

similar predictions if the relative impairments in insulin secretion and insulin resistance are known. For

example, plotting 1hPG vs. 2hPG obtained by simulating OGTTs results in a family of trajectories similar

to the ones for FPG vs. 2hPG shown in Fig. 5. The differences in trajectories due to the contributions of

insulin secretion and insulin action can be read off from the difference between 1hPG and 2hPG and are

similarly most pronounced during prediabetes (Ha et al, ADA poster 1490-P, June, 2019). This

prediction may provide deeper insight into the diagnostic information that can be extracted from different

time points during the OGTT, an issue that is currently receiving much attention (19, 40).

Future versions of the model should include roles for glucagon, as described above. In addition, a more

complex model for insulin signaling would permit the investigation of the effects of particular defects in

those pathways as well as possible mechanisms for the insulin resistance induced by insulin, which was

treated phenomenologically here.

In addition, to be a useful tool for patient stratification and treatment planning, one needs to ascertain the

patient's phenotype. We plan to investigate whether the model can be used, with suitable modifications,

to solve the inverse problem of inferring the individual's parameters of insulin resistance and beta-cell

function from the observed behavior.

Acknowledgments

The work was supported by the Intramural Research Program of the National Institutes of Health,

NIDDK. We thank Stephanie Chung, Richard Bertram, Over Cabrera, and Cecilia Diniz Behn for helpful

suggestions on the manuscript.

References

775 1. **Abdul-Ghani MA**, **Matsuda M**, **DeFronzo RA**. Strong association between insulin resistance in liver and skeletal muscle in non-diabetic subjects. *Diabet Med* 25: 1289–1294, 2008. doi: 10.1111/j.1464-5491.2008.02597.x.

2. **Ackerman E**, **Rosevear JW**, **McGuckin WF**. A Mathematical Model of the Glucose-tolerance test. *Phys Med Biol* 9: 203–213, 1964. doi: 10.1088/0031-9155/9/2/307.

3. **Ader M**, **Stefanovski D**, **Kim SP**, **Richey JM**, **Ionut V**, **Catalano KJ**, **Hucking K**, **Ellmerer M**, **Van Citters G**, **Hsu IR**, **Chiu JD**, **Woolcott OO**, **Harrison LN**, **Zheng D**, **Lottati M**, **Kolka CM**, **Mooradian V**, **Dittmann J**, **Yae S**, **Liu H**, **Castro AVB**, **Kabir M**, **Bergman RN**. Hepatic insulin clearance is the primary determinant of insulin sensitivity in the normal dog. *Obesity (Silver Spring)* 22: 1238–1245, 2014. doi: 10.1002/oby.20625.

4. **Ashcroft FM**, **Puljung MC**, **Vedovato N**. Neonatal Diabetes and the KATP Channel: From Mutation to Therapy. *Trends Endocrinol Metab* 28: 377–387, 2017. doi: 10.1016/j.tem.2017.02.003.

5. **Bard Ermentrout**. Simulating, Analyzing, and Animating Dynamical Systems [Online]. SIAM. http://www.math.pitt.edu/~bard/xpp/xpp.html.

6. **Basu R**, **Barosa C**, **Jones J**, **Dube S**, **Carter R**, **Basu A**, **Rizza RA**. Pathogenesis of prediabetes: role of the liver in isolated fasting hyperglycemia and combined fasting and postprandial hyperglycemia. *J Clin Endocrinol Metab* 98: E409-417, 2013. doi: 10.1210/jc.2012-3056.

7. **Bergman RN**, **Ader M**. Free fatty acids and pathogenesis of type 2 diabetes mellitus. *Trends Endocrinol Metab* 11: 351–356, 2000.

8. **Bergman RN**, **Ider YZ**, **Bowden CR**, **Cobelli C**. Quantitative estimation of insulin sensitivity. *Am J Physiol* 236: E667-677, 1979.

9. **Best JD**, **Kahn SE**, **Ader M**, **Watanabe RM**, **Ni TC**, **Bergman RN**. Role of glucose effectiveness in the determination of glucose tolerance. *Diabetes Care* 19: 1018–1030, 1996. doi: 10.2337/diacare.19.9.1018.

10. **Bodkin NL**, **Metzger BL**, **Hansen BC**. Hepatic glucose production and insulin sensitivity preceding diabetes in monkeys. *Am J Physiol* 256: E676-681, 1989.

11. **Bolie VW**. Coefficients of normal blood glucose regulation. *J Appl Physiol* 16: 783–788, 1961. doi: 10.1152/jappl.1961.16.5.783.

12. **Bonadonna RC**, **Groop L**, **Kraemer N**, **Ferrannini E**, **Del Prato S**, **DeFronzo RA**. Obesity and insulin resistance in humans: a dose-response study. *Metab Clin Exp* 39: 452–459, 1990. doi: 10.1016/0026-0495(90)90002-t.

13. **Brunzell JD**, **Robertson RP**, **Lerner RL**, **Hazzard WR**, **Ensinck JW**, **Bierman EL**, **Porte D**. Relationships between fasting plasma glucose levels and insulin secretion during intravenous glucose tolerance tests. *J Clin Endocrinol Metab* 42: 222–229, 1976. doi: 10.1210/jcem-42-2-222.

14. **Brunzell JD**, **Robertson RP**, **Lerner RL**, **Hazzard WR**, **Ensinck JW**, **Bierman EL**, **Porte D**. Relationships
between fasting plasma glucose levels and insulin secretion during intravenous glucose tolerance
tests. *J Clin Endocrinol Metab* 42: 222–229, 1976. doi: 10.1210/jcem-42-2-222.

15. **Butler AE**, **Janson J**, **Bonner-Weir S**, **Ritzel R**, **Rizza RA**, **Butler PC**. Beta-cell deficit and increased
beta-cell apoptosis in humans with type 2 diabetes. *Diabetes* 52: 102–110, 2003. doi:
10.2337/diabetes.52.1.102.

16. **Capozzi ME**, **Wait JB**, **Koech J**, **Gordon AN**, **Coch RW**, **Svendsen B**, **Finan B**, **D'Alessio DA**, **Campbell**
JE. Glucagon lowers glycemia when β-cells are active. *JCI Insight* 5, 2019. doi:
10.1172/jci.insight.129954.

17. **Chen Y**, **Wang S**, **Sherman A**. Identifying the targets of the amplifying pathway for insulin secretion
in pancreatic beta-cells by kinetic modeling of granule exocytosis. *Biophys J* 95: 2226–2241, 2008.
doi: 10.1529/biophysj.107.124990.

18. **Choy S**, **Kjellsson MC**, **Karlsson MO**, **de Winter W**. Weight-HbA1c-insulin-glucose model for
describing disease progression of type 2 diabetes. *CPT Pharmacometrics Syst Pharmacol* 5: 11–19,
2016. doi: 10.1002/psp4.12051.

19. **Chung ST**, **Ha J**, **Onuzuruike AU**, **Kasturi K**, **Galvan-De La Cruz M**, **Bingham BA**, **Baker RL**,
Utumatwishima JN, **Mabundo LS**, **Ricks M**, **Sherman AS**, **Sumner AE**. Time to glucose peak during
an oral glucose tolerance test identifies prediabetes risk. .

20. **Cobelli C**, **Dalla Man C**, **Toffolo G**, **Basu R**, **Vella A**, **Rizza R**. The oral minimal model method.
Diabetes 63: 1203–1213, 2014. doi: 10.2337/db13-1198.

21. **D'Alessio D**. The role of dysregulated glucagon secretion in type 2 diabetes. *Diabetes Obes Metab*
13 Suppl 1: 126–132, 2011. doi: 10.1111/j.1463-1326.2011.01449.x.

22. **Dankner R**, **Chetrit A**, **Shanik MH**, **Raz I**, **Roth J**. Basal-state hyperinsulinemia in healthy
normoglycemic adults is predictive of type 2 diabetes over a 24-year follow-up: a preliminary
report. *Diabetes Care* 32: 1464–1466, 2009. doi: 10.2337/dc09-0153.

23. **De Gaetano A**, **Hardy T**, **Beck B**, **Abu-Raddad E**, **Palumbo P**, **Bue-Valleskey J**, **Pørksen N**.
Mathematical models of diabetes progression. *Am J Physiol Endocrinol Metab* 295: E1462-1479,
2008. doi: 10.1152/ajpendo.90444.2008.

24. **De Gaetano A**, **Hardy TA**. A novel fast-slow model of diabetes progression: Insights into mechanisms
of response to the interventions in the Diabetes Prevention Program. *PLoS ONE* 14: e0222833,
2019. doi: 10.1371/journal.pone.0222833.

25. **de Winter W**, **DeJongh J**, **Post T**, **Ploeger B**, **Urquhart R**, **Moules I**, **Eckland D**, **Danhof M**. A
mechanism-based disease progression model for comparison of long-term effects of pioglitazone,
metformin and gliclazide on disease processes underlying Type 2 Diabetes Mellitus. *J*
Pharmacokinet Pharmacodyn 33: 313–343, 2006. doi: 10.1007/s10928-006-9008-2.

26. **Dean ED**. A Primary Role for α-Cells as Amino Acid Sensors. *Diabetes* 69: 542–549, 2020. doi:
10.2337/dbi19-0021.

27. **DeFronzo RA**. Lilly lecture 1987. The triumvirate: beta-cell, muscle, liver. A collusion responsible for
NIDDM. *Diabetes* 37: 667–687, 1988.

28. **DeFronzo RA**, **Tripathy D**. Skeletal Muscle Insulin Resistance Is the Primary Defect in Type 2
Diabetes. *Diabetes Care* 32: S157–S163, 2009. doi: 10.2337/dc09-S302.

29. **DREAM (Diabetes REduction Assessment with ramipril and rosiglitazone Medication) Trial**
Investigators, **Gerstein HC**, **Yusuf S**, **Bosch J**, **Pogue J**, **Sheridan P**, **Dinccag N**, **Hanefeld M**,
Hoogwerf B, **Laakso M**, **Mohan V**, **Shaw J**, **Zinman B**, **Holman RR**. Effect of rosiglitazone on the
frequency of diabetes in patients with impaired glucose tolerance or impaired fasting glucose: a
randomised controlled trial. *Lancet* 368: 1096–1105, 2006. doi: 10.1016/S0140-6736(06)69420-8.

30. **Dube S**, **Errazuriz-Cruzat I**, **Basu A**, **Basu R**. The forgotten role of glucose effectiveness in the
regulation of glucose tolerance. *Curr Diab Rep* 15: 605, 2015. doi: 10.1007/s11892-015-0605-6.

31. **Gerich JE**. Is reduced first-phase insulin release the earliest detectable abnormality in individuals
destined to develop type 2 diabetes? *Diabetes* 51 Suppl 1: S117-121, 2002.

32. **German JP**, **Thaler JP**, **Wisse BE**, **Oh-I S**, **Sarruf DA**, **Matsen ME**, **Fischer JD**, **Taborsky GJ**, **Schwartz**
MW, **Morton GJ**. Leptin activates a novel CNS mechanism for insulin-independent normalization of
severe diabetic hyperglycemia. *Endocrinology* 152: 394–404, 2011. doi: 10.1210/en.2010-0890.

33. **Gloyn AL**, **Weedon MN**, **Owen KR**, **Turner MJ**, **Knight BA**, **Hitman G**, **Walker M**, **Levy JC**, **Sampson**
M, **Halford S**, **McCarthy MI**, **Hattersley AT**, **Frayling TM**. Large-scale association studies of variants
in genes encoding the pancreatic beta-cell KATP channel subunits Kir6.2 (KCNJ11) and SUR1
(ABCC8) confirm that the KCNJ11 E23K variant is associated with type 2 diabetes. *Diabetes* 52:
568–572, 2003. doi: 10.2337/diabetes.52.2.568.

34. **Glynn E**, **Thompson B**, **Vadrevu S**, **Lu S**, **Kennedy RT**, **Ha J**, **Sherman A**, **Satin LS**. Chronic Glucose
Exposure Systematically Shifts the Oscillatory Threshold of Mouse Islets: Experimental Evidence for
an Early Intrinsic Mechanism of Compensation for Hyperglycemia. *Endocrinology* 157: 611–623,
2016. doi: 10.1210/en.2015-1563.

35. **Goel P**. Insulin resistance or hypersecretion? The βIG picture revisited. *J Theor Biol* 384: 131–139,
2015. doi: 10.1016/j.jtbi.2015.07.033.

36. **Grespan E**, **Giorgino T**, **Arslanian S**, **Natali A**, **Ferrannini E**, **Mari A**. Defective Amplifying Pathway of
β-cell Secretory Response to Glucose in Type 2 Diabetes: Integrated Modeling of in vitro and in
vivo Evidence. .

37. **Ha J**, **Satin LS**, **Sherman AS**. A Mathematical Model of the Pathogenesis, Prevention, and Reversal of
Type 2 Diabetes. *Endocrinology* 157: 624–635, 2016. doi: 10.1210/en.2015-1564.

38. **Hardy T**, **Abu-Raddad E**, **Porksen N**, **De Gaetano A**. Evaluation of a mathematical model of diabetes
progression against observations in the Diabetes Prevention Program. *Am J Physiol Endocrinol*
Metab 303: E200-212, 2012. doi: 10.1152/ajpendo.00421.2011.

39. **Henquin JC**. Triggering and amplifying pathways of regulation of insulin secretion by glucose.
Diabetes 49: 1751–1760, 2000.

40. **Hulman A**, **Witte DR**, **Vistisen D**, **Balkau B**, **Dekker JM**, **Herder C**, **Hatunic M**, **Konrad T**, **Færch K**,
Manco M. Pathophysiological Characteristics Underlying Different Glucose Response Curves: A
Latent Class Trajectory Analysis From the Prospective EGIR-RISC Study. *Diabetes Care* 41: 1740–
1748, 2018. doi: 10.2337/dc18-0279.

41. **Jones CN**, **Pei D**, **Staris P**, **Polonsky KS**, **Chen YD**, **Reaven GM**. Alterations in the glucose-stimulated
insulin secretory dose-response curve and in insulin clearance in nondiabetic insulin-resistant
individuals. *J Clin Endocrinol Metab* 82: 1834–1838, 1997. doi: 10.1210/jcem.82.6.3979.

42. **Jung S-H**, **Jung C-H**, **Reaven GM**, **Kim SH**. Adapting to insulin resistance in obesity: role of insulin
secretion and clearance. *Diabetologia* 61: 681–687, 2018. doi: 10.1007/s00125-017-4511-0.

43. **Karin O**, **Swisa A**, **Glaser B**, **Dor Y**, **Alon U**. Dynamical compensation in physiological circuits. *Mol*
Syst Biol 12: 886, 2016.

44. **Kim MK**, **Reaven GM**, **Kim SH**. Dissecting the relationship between obesity and hyperinsulinemia:
Role of insulin secretion and insulin clearance. *Obesity (Silver Spring)* 25: 378–383, 2017. doi:
10.1002/oby.21699.

45. **Knowler WC**, **Barrett-Connor E**, **Fowler SE**, **Hamman RF**, **Lachin JM**, **Walker EA**, **Nathan DM**,
Diabetes Prevention Program Research Group. Reduction in the incidence of type 2 diabetes with
lifestyle intervention or metformin. *N Engl J Med* 346: 393–403, 2002. doi:
10.1056/NEJMoa012512.

46. **Kodama K**, **Tojjar D**, **Yamada S**, **Toda K**, **Patel CJ**, **Butte AJ**. Ethnic differences in the relationship
between insulin sensitivity and insulin response: a systematic review and meta-analysis. *Diabetes*
Care 36: 1789–1796, 2013. doi: 10.2337/dc12-1235.

47. **Koster JC**, **Remedi MS**, **Dao C**, **Nichols CG**. ATP and sulfonylurea sensitivity of mutant ATP-sensitive
K+ channels in neonatal diabetes: implications for pharmacogenomic therapy. *Diabetes* 54: 2645–
2654, 2005. doi: 10.2337/diabetes.54.9.2645.

48. **Lim EL**, **Hollingsworth KG**, **Aribisala BS**, **Chen MJ**, **Mathers JC**, **Taylor R**. Reversal of type 2 diabetes:
normalisation of beta cell function in association with decreased pancreas and liver triacylglycerol.
Diabetologia 54: 2506–2514, 2011. doi: 10.1007/s00125-011-2204-7.

49. **Mari A**, **Ferrannini E**. Beta-cell function assessment from modelling of oral tests: an effective
approach. *Diabetes Obes Metab* 10 Suppl 4: 77–87, 2008. doi: 10.1111/j.1463-1326.2008.00946.x.

50. **Mason CC**, **Hanson RL**, **Knowler WC**. Progression to type 2 diabetes characterized by moderate then
rapid glucose increases. *Diabetes* 56: 2054–2061, 2007. doi: 10.2337/db07-0053.

51. **Meigs JB**, **Muller DC**, **Nathan DM**, **Blake DR**, **Andres R**, **Baltimore Longitudinal Study of Aging**. The
natural history of progression from normal glucose tolerance to type 2 diabetes in the Baltimore
Longitudinal Study of Aging. *Diabetes* 52: 1475–1484, 2003.

52. **Morton GJ**, **Matsen ME**, **Bracy DP**, **Meek TH**, **Nguyen HT**, **Stefanovski D**, **Bergman RN**, **Wasserman**
DH, **Schwartz MW**. FGF19 action in the brain induces insulin-independent glucose lowering. *J Clin*
Invest 123: 4799–4808, 2013. doi: 10.1172/JCI70710.

53. **Nichols CG**, **Remedi MS**. The diabetic β-cell: hyperstimulated vs. hyperexcited. *Diabetes Obes Metab*
14: 129–135, 2012. doi: 10.1111/j.1463-1326.2012.01655.x.

54. **Nyman E**, **Rozendaal YJW**, **Helmlinger G**, **Hamrén B**, **Kjellsson MC**, **Strålfors P**, **van Riel NAW**,
Gennemark P, **Cedersund G**. Requirements for multi-level systems pharmacology models to reach
end-usage: the case of type 2 diabetes. *Interface Focus* 6: 20150075, 2016. doi:
10.1098/rsfs.2015.0075.

55. **Odgerel Z**, **Lee HS**, **Erdenebileg N**, **Gandbold S**, **Luvsanjamba M**, **Sambuughin N**, **Sonomtseren S**,
Sharavdorj P, **Jodov E**, **Altaisaikhan K**, **Goldfarb LG**. Genetic variants in potassium channels are
associated with type 2 diabetes in a Mongolian population. *J Diabetes* 4: 238–242, 2012. doi:
10.1111/j.1753-0407.2011.00177.x.

56. **Omar-Hmeadi M**, **Lund P-E**, **Gandasi NR**, **Tengholm A**, **Barg S**. Paracrine control of α-cell glucagon
exocytosis is compromised in human type-2 diabetes. *Nat Commun* 11: 1896, 2020. doi:
10.1038/s41467-020-15717-8.

57. **Perreault L**, **Bergman BC**, **Playdon MC**, **Dalla Man C**, **Cobelli C**, **Eckel RH**. Impaired fasting glucose
with or without impaired glucose tolerance: progressive or parallel states of prediabetes? *Am J*
Physiol Endocrinol Metab 295: E428-435, 2008. doi: 10.1152/ajpendo.90354.2008.

58. **Perry RJ**, **Camporez J-PG**, **Kursawe R**, **Titchenell PM**, **Zhang D**, **Perry CJ**, **Jurczak MJ**, **Abudukadier A**,
Han MS, **Zhang X-M**, **Ruan H-B**, **Yang X**, **Caprio S**, **Kaech SM**, **Sul HS**, **Birnbaum MJ**, **Davis RJ**, **Cline**
GW, **Petersen KF**, **Shulman GI**. Hepatic acetyl CoA links adipose tissue inflammation to hepatic
insulin resistance and type 2 diabetes. *Cell* 160: 745–758, 2015. doi: 10.1016/j.cell.2015.01.012.

59. **Petersen MC**, **Shulman GI**. Mechanisms of Insulin Action and Insulin Resistance. *Physiol Rev* 98:
2133–2223, 2018. doi: 10.1152/physrev.00063.2017.

60. **Piccinini F**, **Dalla Man C**, **Vella A**, **Cobelli C**. A Model for the Estimation of Hepatic Insulin Extraction
After a Meal. *IEEE Transactions on Biomedical Engineering* 63: 1925–1932, 2016. doi:
10.1109/TBME.2015.2505507.

61. **Piccinini F**, **Polidori DC**, **Gower BA**, **Bergman RN**. Hepatic but Not Extrahepatic Insulin Clearance Is
Lower in African American Than in European American Women. *Diabetes* 66: 2564–2570, 2017.
doi: 10.2337/db17-0413.

62. **Piller C**. Dubious diagnosis. *Science* 363: 1026–1031, 2019. doi: 10.1126/science.363.6431.1026.

63. **Polidori D**, **Bergman RN**, **Sumner AE**. A New Model-Based Method for Estimating Hepatic and
Peripheral Insulin Clearance from Intravenous Glucose Tolerance Test Data (2015 ADA abstract
254-LB). [date unknown].

64. **Porat S**, **Weinberg-Corem N**, **Tornovsky-Babaey S**, **Schyr-Ben-Haroush R**, **Hija A**, **Stolovich-Rain M**,
Dadon D, **Granot Z**, **Ben-Hur V**, **White P**, **Girard CA**, **Karni R**, **Kaestner KH**, **Ashcroft FM**, **Magnuson**
MA, **Saada A**, **Grimsby J**, **Glaser B**, **Dor Y**. Control of pancreatic β cell regeneration by glucose
metabolism. *Cell Metab* 13: 440–449, 2011. doi: 10.1016/j.cmet.2011.02.012.

65. **Pories WJ**, **Dohm GL**. Diabetes: have we got it all wrong? Hyperinsulinism as the culprit: surgery
provides the evidence. *Diabetes Care* 35: 2438–2442, 2012. doi: 10.2337/dc12-0684.

66. **Rahier J**, **Guiot Y**, **Goebbels RM**, **Sempoux C**, **Henquin JC**. Pancreatic beta-cell mass in European
subjects with type 2 diabetes. *Diabetes Obes Metab* 10 Suppl 4: 32–42, 2008. doi: 10.1111/j.1463-
1326.2008.00969.x.

67. **Ribbing J**, **Hamrén B**, **Svensson MK**, **Karlsson MO**. A model for glucose, insulin, and beta-cell
dynamics in subjects with insulin resistance and patients with type 2 diabetes. *J Clin Pharmacol* 50:
861–872, 2010. doi: 10.1177/0091270009349711.

68. **RISE Consortium**. Lack of Durable Improvements in β-Cell Function Following Withdrawal of
Pharmacological Interventions in Adults With Impaired Glucose Tolerance or Recently Diagnosed
Type 2 Diabetes. *Diabetes Care* 42: 1742–1751, 2019. doi: 10.2337/dc19-0556.

69. **Saisho Y**, **Butler AE**, **Manesso E**, **Elashoff D**, **Rizza RA**, **Butler PC**. β-cell mass and turnover in
humans: effects of obesity and aging. *Diabetes Care* 36: 111–117, 2013. doi: 10.2337/dc12-0421.

70. **Sedaghat AR**, **Sherman A**, **Quon MJ**. A mathematical model of metabolic insulin signaling pathways.
Am J Physiol Endocrinol Metab 283: E1084-1101, 2002. doi: 10.1152/ajpendo.00571.2001.

71. **Shanik MH**, **Xu Y**, **Skrha J**, **Dankner R**, **Zick Y**, **Roth J**. Insulin resistance and hyperinsulinemia: is
hyperinsulinemia the cart or the horse? *Diabetes Care* 31 Suppl 2: S262-268, 2008. doi:
10.2337/dc08-s264.

72. **Sharma AX**, **Quittner-Strom EB**, **Lee Y**, **Johnson JA**, **Martin SA**, **Yu X**, **Li J**, **Lu J**, **Cai Z**, **Chen S**, **Wang**
M-Y, **Zhang Y**, **Pearson MJ**, **Dorn AC**, **McDonald JG**, **Gordillo R**, **Yan H**, **Thai D**, **Wang ZV**, **Unger RH**,
Holland WL. Glucagon Receptor Antagonism Improves Glucose Metabolism and Cardiac Function
by Promoting AMP-Mediated Protein Kinase in Diabetic Mice. *Cell Rep* 22: 1760–1773, 2018. doi:
10.1016/j.celrep.2018.01.065.

73. **Tabák AG**, **Jokela M**, **Akbaraly TN**, **Brunner EJ**, **Kivimäki M**, **Witte DR**. Trajectories of glycaemia,
insulin sensitivity, and insulin secretion before diagnosis of type 2 diabetes: an analysis from the
Whitehall II study. *Lancet* 373: 2215–2221, 2009. doi: 10.1016/S0140-6736(09)60619-X.

74. **Thomas DD**, **Corkey BE**, **Istfan NW**, **Apovian CM**. Hyperinsulinemia: An Early Indicator of Metabolic
Dysfunction. *J Endocr Soc* 3: 1727–1747, 2019. doi: 10.1210/js.2019-00065.

75. **Topp B**, **Promislow K**, **deVries G**, **Miura RM**, **Finegood DT**. A model of beta-cell mass, insulin, and
glucose kinetics: pathways to diabetes. *J Theor Biol* 206: 605–619, 2000. doi:
10.1006/jtbi.2000.2150.

76. **Tricò D**, **Galderisi A**, **Mari A**, **Polidori D**, **Galuppo B**, **Pierpont B**, **Samuels S**, **Santoro N**, **Caprio S**.
Intrahepatic fat, irrespective of ethnicity, is associated with reduced endogenous insulin clearance
and hepatic insulin resistance in obese youths: a cross-sectional and longitudinal study from the
Yale Pediatric NAFLD cohort. .

77. **Utumatwishima JN**, **Chung ST**, **Bentley AR**, **Udahogora M**, **Sumner AE**. Reversing the tide - diagnosis
and prevention of T2DM in populations of African descent. *Nat Rev Endocrinol* 14: 45–56, 2018.
doi: 10.1038/nrendo.2017.127.

78. **Wang G**. Raison d'être of insulin resistance: the adjustable threshold hypothesis. *J R Soc Interface*
11: 20140892, 2014. doi: 10.1098/rsif.2014.0892.

79. **Wang Y-F**, **Khan M**, **van den Berg HA**. Interaction of fast and slow dynamics in endocrine control
systems with an application to β-cell dynamics. *Math Biosci* 235: 8–18, 2012. doi:
10.1016/j.mbs.2011.10.003.

80. **Wewer Albrechtsen NJ**, **Pedersen J**, **Galsgaard KD**, **Winther-Sørensen M**, **Suppli MP**, **Janah L**,
Gromada J, **Vilstrup H**, **Knop FK**, **Holst JJ**. The Liver-α-Cell Axis and Type 2 Diabetes. *Endocr Rev* 40:
1353–1366, 2019. doi: 10.1210/er.2018-00251.

81. **Yki-Järvinen H**. Thiazolidinediones and the liver in humans. *Curr Opin Lipidol* 20: 477–483, 2009. doi:
10.1097/MOL.0b013e3283321d37.

82. **Zhu L**, **Dattaroy D**, **Pham J**, **Wang L**, **Barella LF**, **Cui Y**, **Wilkins KJ**, **Roth BL**, **Hochgeschwender U**,
Matschinsky FM, **Kaestner KH**, **Doliba NM**, **Wess J**. Intra-islet glucagon signaling is critical for
maintaining glucose homeostasis. *JCI Insight* 5, 2019. doi: 10.1172/jci.insight.127994.

1007

 Figure Legends

Figure 1: IGT-first pathway to diabetes. (A) Assumed severe decline in peripheral insulin sensitivity. (B)
Assumed mild decline in hepatic insulin sensitivity. (C) Simulated longitudinal changes based on the
assumptions in (A) and (B) in fasting plasma glucose (FPG) and two-hour glucose (2hPG) during OGTTs
performed at each time point. The virtual subject experiences first high two-hour glucose (IGT), then
high fasting glucose (CGI), and finally crosses the 2hPG threshold for T2D. (D) Simulated longitudinal
changes in fasting plasma insulin (FPI) and two-hour insulin (2hPI) during the OGTTs. Insulin increases
early on but decreases later. (E) The component of β-cell function represented by σ decreases
progressively throughout. (F) The β-cell mass, β, first increases during prediabetes, then decreases after
diabetes onset.

Figure 2: Insulin resistance (same as Fig. 1) does not lead to diabetes if beta-cell function is sufficiently
responsive. The γ-dynamics is made stronger than in Fig. 1 by decreasing γ_s to 90 *mg/dl* from 100 *mg/dl*
(see other parameters in Table S12A). (A) Assumed severe decline in peripheral insulin sensitivity. (B)
Assumed mild decline in hepatic insulin sensitivity. (C) Simulated longitudinal changes in FPG and 2hPG
during OGTTs performed at each time point. The virtual subject experiences modest rises in glucose and
crosses the threshold for IGT but never crosses the thresholds for IFG, CGI or T2D. (D) Simulated
longitudinal changes in fasting plasma insulin (FPI) and two-hour insulin (2hPI) during the OGTTs.
Insulin concentration increases and saturates but never declines. (E) The β-cell function component σ first
decreases, but then levels off. Increase in the β-cell function component represented by γ (not shown)
helps limit the rise in glucose, which allows the β-cell mass β to increase gradually throughout (F).

Figure 3: IFG-first pathway to diabetes. (A) Assumed mild decline in peripheral insulin sensitivity. (B)
Assumed severe decline in hepatic insulin sensitivity component *hepa$_{SI}$*. (C) Simulated longitudinal
changes in FPG and 2hPG during OGTTs performed at each time point. The virtual subject experiences
first high FPG (IFG), then high 2hPG (CGI), and finally crosses the FPG threshold for T2D first.
Simulated longitudinal changes in fasting plasma insulin (FPI) and two-hour insulin (2hPI) during the
OGTTs. Insulin concentration increases early on but decreases later. (E) The β-cell function component σ
decreases progressively throughout. (F) The β-cell mass β increases during prediabetes then decreases
after diabetes onset.

Figure 4: The CGI state is not obligatory. (A), (B): Extreme discrepancy between peripheral and hepatic
(A – B) insulin resistance results in progression directly from IGT to T2D without passing through CGI
based on OGTTs performed at each time point (C). FPG does not exceed the threshold for IFG until after
T2D onset. (D) Simulated longitudinal changes in fasting plasma insulin (FPI) and two-hour insulin
(2hPI) during the OGTTs. Insulin increases early on, then decreases. (E) The β-cell function component σ
decreases more rapidly than in Figs. 1, 3. (F) The β-cell mass β increases during prediabetes, then
decreases after diabetes onset.

Figure 5: (A) Two-hour glucose (2hPG) plotted vs. fasting glucose (FPG) for varying degrees of
peripheral insulin sensitivity, S_I, and hepatic insulin sensitivity *hepa$_{SI}$*, starting from the same initial
values as in Figs. 1 – 4 and evolving to the target values, *tar$_{SI}$* (Eq. A13) and *tar$_{hepaSI}$* (Eq. A14) as
follows: from upper left to lower right, the *tar$_{SI}$* values increase: 0.1, 0.17, 0.22, 0.33, 0.4 and 0.5; and the

tar$_{hepaSI}$ values decrease: 0.85, 0.6, 0.45, 0.35, 0.25 and 0.1. All other parameters are fixed. (B)
expanded view of (A) to highlight the prediabetes region.

1049

1050   Figure 6: Glucose during OGTTs performed at the times indicated by the black dots in (A) Fig. 1 (IGT-
1051   first pathway) and (B) Fig. 3 (IFG-first pathway). (C, D) insulin corresponding to A and B, respectively.
1052   See text for details.

1053

1054   Figure 7: Insulin during IVGTTs performed at the time points indicated by the black dots in (A) Fig. 1
1055   (IGT-first pathway) and (B) Fig. 3 (IFG-first pathway). (C, D): RRP size (numbers of vesicles)
1056   corresponding to A and B, respectively. Decline of AIRg parallels the reduction of RRP for each case.

1057

1058

Figure 8: Drug therapies targeting either peripheral insulin sensitivity (modeled as a rapid increase in S_I)
or hepatic insulin sensitivity (modeled as a rapid increase in *hepa$_{SI}$*; see Tables S13 – S16 for details).
The drugs are applied in the early stages of T2D (A, B) or during prediabetes (C, D). (A, C) Dominant
peripheral insulin resistance, leading to IGT-first pathway as in Fig. 1. (B, D) Dominant hepatic insulin
resistance, leading to IFG-first pathway as in Fig. 3. The appropriately targeted drug is more effective
than the inappropriately targeted drug in each case, but therapy is more effective when initiated during
prediabetes.

1066

Figure 9: Effects of insulin clearance and insulin-induced insulin resistance. The case of Fig. 1 (IGT first)
is re-simulated with varying rates of insulin clearance and induced insulin resistance, defined by Eqs. 13 –
16. Panel A: $S_I = S_{I,0}E$, where $S_{I,0}$ is calculated as in Fig. 1 and E is the induced effect plotted in Panel
F. Legend: *black* (control), same as Fig. 1, but with insulin clearance $k = 0.4861$ min^{-1}, essentially no
induced effect ($\alpha_E = 300$ µU/ml);   *green*: reduced clearance ($k = 0.37499$ min^{-1}), modest induced effect
($\alpha_E = 50$ µU/ml); *blue*: $k = 0.37499$ min^{-1}, stronger induced effect ($\alpha_E = 15$ µU/ml); *red*: $k = 0.37499$
min^{-1}, very strong induced effect ($\alpha_E = 5$ µU/ml). Other parameters are listed in Table S17.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

A

B

Fig. 6

Fig. 7

Fig. 8

Fig. 9