The *Salmonella* LysR family regulator, RipR, activates the SPI-13 encoded

itaconate degradation cluster.

*Steven J. Hersch † and William Wiley Navarre ***

Department of Molecular Genetics, University of Toronto, Toronto, Canada

†, Current address: Department of Ecosystem and Public Health, University of Calgary, Calgary,

Canada

*, For correspondence: william.navarre@utoronto.ca

Abstract

Itaconate is a dicarboxylic acid that inhibits the isocitrate lyase enzyme of the bacterial glyoxylate shunt. Activated macrophages have been shown to produce itaconate, suggesting that these immune cells may employ this metabolite as a weapon against invading bacteria. Here we demonstrate that, *in vitro*, itaconate can exhibit bactericidal effects under acidic conditions similar to the pH of a macrophage phagosome. In parallel, successful pathogens including *Salmonella* have acquired a genetic operon encoding itaconate degradation proteins, which are induced heavily in macrophages. We characterize the regulation of this operon by the neighbouring gene, *ripR*, in specific response to itaconate. Moreover, we develop an itaconate biosensor based on the operon promoter that can detect itaconate in a semi-quantitative manner and, when combined with the *ripR* gene, is sufficient for itaconate-regulated expression in *E. coli*. Using this biosensor with fluorescence microscopy, we observe bacteria responding to itaconate in the phagosomes of macrophage and provide additional evidence that interferon-γ stimulates macrophage itaconate synthesis and that J774 mouse macrophages produce substantially more itaconate than the human THP-1 monocyte cell line. In summary, we examine the role of itaconate as an antibacterial metabolite in mouse and human macrophage, characterize the regulation of *Salmonella*'s defense against it, and develop it as a convenient itaconate biosensor and inducible promoter system.

Importance

In response to invading bacteria, immune cells can produce a molecule called itaconate, which can inhibit microbial metabolism. Here we show that itaconate can also directly kill

Salmonella when combined with moderate acidity, further supporting itaconate's role as an

antibacterial weapon. We also discover how *Salmonella* recognizes itaconate and activates a

defense to degrade it, and we harness this response to make a biosensor that detects the

presence of itaconate. This biosensor is versatile, working in *Salmonella enterica* or lab strains

of *Escherichia coli*, and can detect itaconate quantitatively in the environment and in immune

cells. By understanding how immune cells kill bacteria and how the microbes defend

themselves, we can better develop novel antibiotics to inhibit pathogens such as *Salmonella*.

Introduction

The mammalian immune system includes a multitude of weapons to defend against

invading microbes and successful pathogens have evolved a plethora of mechanisms to evade,

manipulate, or even benefit from these immune responses. One such pathogen, *Salmonella*

*enterica s*erovar Typhimurium (hereafter referred to as *Salmonella*), has acquired a number of

Salmonella pathogenicity islands (SPI) that support its survival inside of a host organism. For

instance, *Salmonella* employs SPI-1 to invade non-phagocytic cells, and SPI-2 allows the bacteria

to survive intracellularly – including in macrophage – which is important for *Salmonella*

virulence [1–4]. These traits allow *Salmonella* to invade the gut epithelium and induce intestinal

inflammation resulting in the characteristic gastroenteritis disease. Moreover, the induced

inflammation is not merely a threat that *Salmonella* must survive, but it has adapted to thrive in

the oxidative environment of the inflamed intestine and utilize inflammation-derived

metabolites to outcompete resident microbiota [5–7].

Itaconate (2-Methylenesuccinic acid, 2-Methylidenebutanedioic acid) is a metabolite

originally recognized in fungal species such as *Aspergillus terreus* and produced commercially

for use in polymer production [8–10]. As an unsaturated diacaboxylate that is somewhat similar in

structure to succinate, itaconate is a potent inhibitor of the glyoxylate shunt enzyme AceA

(isocitrate lyase) [11–13]. As such, itaconate inhibits bacterial growth on carbon sources such as

acetate and fatty acids, conditions that necessitate the glyoxylate shunt to generate the 4-

carbon skeletons that are critical for amino acid biosynthesis and central metabolism.

Interestingly, it has been demonstrated that activated macrophage employ the IRG1 protein to

produce itaconate, with higher concentrations being produced by mouse macrophage than

human [13–15]. Moreover, IRG1 closely associated with vesicles containing *Legionella pneumophila*

and itaconate showed bactericidal activity against this pathogen *in vitro* [15]. Itaconate was also

found to inhibit *Salmonella* growth by reducing media pH and itaconate levels in *Salmonella*-

infected mice correlated with splenomegaly [16]. Cumulatively, these works suggest that

itaconate acts as a weaponized metabolite that the immune system employs to inhibit the

growth of, or kill, invading bacteria.

If itaconate is an immune-derived antibacterial metabolite then it follows logically that

successful pathogens must have methods to evade its effects. Indeed an operon has been

identified in *Yersinia* (*ripABC* for 'required for intracellular proliferation') that encodes three

enzymes catalyzing the ATP/succinyl-CoA-dependent degradation of itaconate into pyruvate

and acetyl-CoA [17]. The operon is not restricted to *Yersinia* and a variety of other bacteria

including *Pseudomonas* encode homologs or functional analogs of these enzymes. Several

lineages of *Salmonella enterica* harbor a cluster of genes (e.g. genes STM3120-STM3117 in S.

Typhimurium strain LT2) within SPI-13 that we refer to here as the 'itaconate response operon'

(IRO). Interestingly, the IRO genes of *Salmonella* have been shown to be induced heavily in

macrophage but not under any other condition tested, supporting a role in degrading

macrophage-produced itaconate [18–20]. High throughput screens have suggested that genes from

this operon are important for *Salmonella* survival in mice [21–23]. Furthermore, it has also been

shown that SPI-13 is present in many generalist *S. enterica* serovars but not in some human-

restricted serovars of *Salmonella* (e.g. serovars Typhi and Paratyphi A, which encode SPI-8)

possibly due to reduced itaconate synthesis by human macrophages [24].

In this work we show that itaconate is bactericidal at low but not neutral pH and

elucidate the regulation of the *Salmonella* IRO and its induction in mouse and human

macrophage. We show that the promoter of the IRO (P_{IRO}) is specifically induced by itaconate

and that the LysR family transcriptional regulator encoded by the upstream gene, STM3121

(which we propose to name *ripR*), is both necessary and sufficient for this induction.

Furthermore, using P_{IRO} with a GFP reporter, we develop a semi-quantitative itaconate

biosensor and employ it to show that the IRO is induced heavily in the J774 mouse macrophage

cell line but requires interferon-γ (IFN-γ) stimulation to show a detectable response in the THP-1

human monocyte cell line.

Results

The* Salmonella *IRO is induced specifically by itaconate in a* RipR-*dependent manner

The *Salmonella* pathogenicity island-13 includes genes encoding RipC/Ccl (STM3120 in

strain LT2), RipB/Ich (STM3119) and RipA/Ict (STM3118), whose homologs have been

demonstrated to degrade itaconate into pyruvate and acetyl-CoA [17]. This operon, which we

refer to as the itaconate response operon (IRO), appears to also include the STM3117 gene

(encoding a predicted glyoxalase-domain containing protein) and is adjacent to the STM3121

(*ripR*) gene on the reverse DNA strand (Figure 2).

The IRO is strongly induced in cultured mouse macrophages but, according to

expression data from SalCom and other published work, is seemingly independent of known

virulence regulators (PhoP, RpoS, SsrA/B, OmpR, SlyA, etc.). We hypothesized that itaconate

may act as an inducer of IRO expression via the adjacent LysR-family regulator encoded by

STM3121. To assess this, we constructed a plasmid-borne fusion of the operon's promoter

(P_{IRO}) to superfolder GFP (sfGFP) as a reporter[25]. Indeed, we found that the P_{IRO} promoter was

induced highly in the presence of itaconate and this response was entirely dependent on the

presence of RipR as neither a *Salmonella ripR* deletion mutant nor the same reporter plasmid in

E. coli K12 (which does not encode *ripR*) showed induction (Figure 3A). In contrast, when the

ripR gene was included on the reporter plasmid, itaconate-induced P_{IRO} expression was restored

in both the Δ*ripR Salmonella* strain and in *E. coli*. These data not only demonstrate that P_{IRO} is

induced in response to itaconate, but also that the neighbouring gene, *ripR*, is both necessary

and sufficient for this induction.

To assess if the IRO promoter is induced specifically by itaconate, we examined

induction of the P_{IRO} reporter plasmid in media supplemented with a panel of similar

metabolites. While mesaconate, citramalate and methylsuccinate (in order of induction

strength) slightly induced expression, induction by itaconate was drastically more pronounced,

suggesting that it is the principal inducer (Figure 3B). Notably, similar results were obtained in

complex media (LB) and in MOPS minimal media with either glucose or glycerol as a carbon

source, suggesting that the induction only requires itaconate and not additional factors in the

media (Figure S2). Furthermore, induction by itaconate occurred in a dose dependent manner

indicating that the reporter can be used to semi-quantitatively assess itaconate concentrations

encountered by the bacteria (Figure S3).

Itaconate import is independent of the dicarboxylate transporter DctA

125   Another important question was whether or not itaconate required the primary aerobic

dicarboxylate transporter, DctA, for import into the cell. We compared itaconate-mediated

induction of the P_{IRO}-sfGFP reporter in wild-type or $\Delta dctA$ *Salmonella*. Interestingly, we found

that the IRO promoter was still heavily induced in the *dctA* mutant in the presence of itaconate

(Figure S4). This demonstrates that the DctA dicarboxylate transporter is not required for

itaconate-mediated induction of P_{IRO}, suggesting that itaconate import is independent of *dctA*.

Itaconate is bactericidal at low pH

133   Previous work has demonstrated that itaconate can inhibit the function of the

glyoxylate shunt enzyme, AceA, and act as a bacteriostatic agent when bacteria rely on carbon

sources such as acetate that require this pathway [11–13]. It has also been suggested that

itaconate can inhibit bacteria by influencing media pH and at least one publication has

demonstrated that it can have bactericidal activity [15,16]. To clarify this later phenotype we

hypothesized that the dicarboxylic acid chemistry of itaconate would allow it to act in a

bactericidal fashion at low pH by acting as a proton shuttle. In brief, the carboxyl groups of

itaconate (pK_a = 5.5 and 3.8) protonate and lose their charge at lower pH allowing them to

traverse the bacterial membrane and release the protons in the more neutral pH of the

cytoplasm, thereby exacerbating acid stress (Figure 1A).

To emulate the intracellular conditions that *Salmonella* may encounter in a *Salmonella*

containing vacuole (SCV) of a macrophage we added itaconate to LPM media and then acidified

to pH 4.4, 5.0, or 5.8 to cover a range from the most acidic to more regular estimates of SCV pH

[26–28]. Indeed we found that wild-type *Salmonella* showed a 1000-fold decrease in survival after

3h hours at pH 4.4 with itaconate (Figure 1B). This lethality was alleviated at higher pH and also

occurred using a similar dicarboxylic acid, succinate. Importantly, the bactericidal effect was

also dependent on the presence of itaconate or succinate, as pH 4.4 LPM did not kill *Salmonella*

in the absence of a dicarboxylic acid (Figure S1A). Interestingly, deletion of the entire IRO

(STM3120-STM3117) or *aceA* had no effect, but deletion of the general stress response sigma

factor, RpoS (σ^{32}, σ^{S}), exacerbated *Salmonella's* sensitivity at both pH 4.4 and 5.0 (Figures 1C

and S1B). Cumulatively, these data demonstrate that itaconate or other dicarboxylic acids can

act in a bactericidal fashion under acidic conditions by exacerbating acid stress.

The* Salmonella *IRO does not significantly contribute to short-term survival in a mouse

macrophage cell line

The inhibitory effect of itaconate on AceA, its bactericidal activity under acidic

conditions, and the synthesis of itaconate in macrophage combine to support the concept that

these immune cells may be employing itaconate as an antibacterial compound. As a successful

pathogen, *Salmonella* has adapted to survive in activated macrophage and multiple previous

works have examined how IRO genes may influence *Salmonella* survival and virulence [21–23]. In

our hands, we found no significant reduction in survival of Δ*IRO* or Δ*ripR* strains in the mouse

J774 macrophage cell line (Figure S5). When the macrophages were pre-stimulated with IFN-γ,

there was a slight reduction in survival relative to wild-type, but this was not significant when

compared to an *aceA* mutant that showed no survival defect. In contrast, the growth of a *phoP*

deletion control strain was inhibited by macrophages even without IFN-γ.

Salmonella *encounter itaconate in the phagosomes of mouse and IFN-γ-stimulated human*

macrophage

Multiple high throughput studies have demonstrated that the IRO genes are induced

heavily in mouse macrophage [18–20]. Additional studies have identified itaconate in both mouse

and human macrophage but the mouse cells appear to produce significantly more of the

metabolite [13,14]. Furthermore, it has been demonstrated that interferon-β (IFN-β) and IFN-γ can

stimulate itaconate production in mouse macrophage [15,29–31].

To examine itaconate levels encountered by intracellular bacteria inside macrophage,

we employed our P_{IRO}-sfGFP reporter plasmid as a biosensor. By including a constitutively

expressed mCherry gene on the same plasmid, we could microscopically observe individual

bacteria inside macrophage and obtain semi-quantitative data by generating a GFP/mCherry

fluorescence ratio as an indicator of P_{IRO} induction and accordingly itaconate levels. Using this

system, we observed strong induction of the P_{IRO} promoter for wild-type *Salmonella* in J774

mouse macrophage and this signal was absent in the Δ*ripR* control (Figure 4). Furthermore, the

response could also be observed in *E. coli* if RipR was encoded on the same plasmid,

demonstrating that bacteria that are poorly adapted to intracellular survival also encounter

itaconate in macrophage.

In contrast to the mouse cell line, unstimulated human THP-1 monocytes showed

negligible itaconate levels as very few of the bacterial reporters showed any green fluorescence

above background levels (Figure 4). The bacteria did express the constitutive mCherry and

could be observed in the macrophage, suggesting that the lack of green fluorescence was not

due to decreased bacterial survival or protein expression. Stimulation of THP-1 cells with IFN-γ

(M1 activation), however, led to a significant increase in the green fluorescence of the reporter

bacteria in a RipR-dependent fashion. In contrast, itaconate levels in THP-1 cells induced with

IL-4 and IL-13 (M2 activation) resembled unstimulated cells (Figure S6).

Discussion

In this work we demonstrate that itaconate becomes bactericidal at acidic pH,

suggesting an additional mechanism for itaconate to act as an antibacterial metabolite beyond

inhibition of AceA. Thus, elevated itaconate levels in macrophage may act to inhibit bacterial

metabolism while also exacerbating acid stress on microbes in the phagosome. Protonation of

itaconate under acidic conditions may also grant it increased access to the bacterial cytoplasm

where de-protonation would trap the charged form close to its AceA target. This organic acid

killing effect has been demonstrated previously, including in a recent work showing propionate

inhibition of *Salmonella* in mice [32,33]. Of note, we also find that bacterial killing occurs with

succinate, a metabolite similar to itaconate that similarly increases in concentration in activated

macrophage [34]. Our findings that these dicarboxylic acids can kill *Salmonella* at pH 4.4 but not

higher may contribute to why *Salmonella* manipulates the SCV to maintain a pH closer to 5.0 in

order to avoid this organic acid stress. Moreover, they imply that organic acids such as

itaconate and succinate may contribute to the antibacterial activity of acidified phagosomes.

The antibacterial potential of itaconate, its synthesis in activated macrophage, and the

localization of IRG1 to bacteria-containing vacuoles, support its potential role as a weapon

against intracellular bacteria. Here we examined survival of *Salmonella* IRO or *ripR* deletion

strains in the mouse macrophage J774 cell line but saw no significant decrease in survival,

similar to a recent study examining SPI-13 in RAW264.7I macrophage [24]. However, in that work,

Espinoza *et al.* discovered that SPI-13 does play a role in *Salmonella* internalization into mouse

– but not human – macrophage [24]. Combined with previous works showing reduced survival of

IRO mutants in mice, this operon may play a more significant survival role in the context of

infection in animals that produce copious amounts of itaconate [20–23].

We find *Salmonella* responds to itaconate *in vitro* and intracellularly by strongly inducing

an operon encoding itaconate degradation proteins. This response is largely specific to

itaconate and is entirely dependent on the neighbouring gene (*STM3121* in the prototypical

Salmonella enterica Sv. Typhimurium strain LT2), which we propose to rename *ripR* ('rip operon

regulator') to conform to earlier nomenclature. The *ripR* gene product is predicated to be a LysR

family transcriptional regulator, suggesting that itaconate induces IRO expression by interacting

directly with the substrate binding domain of RipR to activate it. Moreover, we find that RipR is

sufficient for itaconate induction of the P_{IRO} promoter in *E. coli*, demonstrating its potential as a

novel inducible expression system with over 50-fold higher transcription in the presence of the

inexpensive and readily available inducer. A limitation of this expression system would be a

requirement for growth on carbon sources independent of the glyoxylate shunt and also

growth at neutral or alkaline pH, as we demonstrate that itaconate is bactericidal under acidic

conditions. However, for many studies these conditions are met, adding P_{IRO} to the repertoire

of available inducible promoter systems.

Using our P_{IRO}-sfGFP itaconate biosensor, we showed a pronounced response in

unstimulated mouse macrophage whereas no induction was observed in the THP-1 human

macrophage cell line without stimulation, suggesting that these cells are not producing

itaconate to the same degree. Interferon has previously been demonstrated to stimulate

itaconate production in mouse macrophage and we found that our biosensor was induced in

human cells stimulated with IFN-γ [15,29–31]. Thus, while the human cell line was able to produce

itaconate, it required auxiliary induction to do so and still produced less than the uninduced

mouse macrophage. While it is possible that this reflects an artifact of the cell lines employed, it

aligns well with previous works that quantified itaconate in both mouse and human cells [13,14].

Furthermore, Espinoza *et al.* recently determined that SPI-13 is abundant in generalist

Salmonella serovars but not human-restricted ones (which instead encode SPI-8), suggesting

that low itaconate levels in humans render the IRO dispensable in these strains [24].

A recent study demonstrated that small molecules can inhibit the activity of the IRO

proteins and sensitize *Salmonella* to itaconate inhibition in minimal media [35]. Such drugs could

also sensitise other bacteria to itaconate including *Yersinia, Pseudomonas* and *Mycobacteria*

species, which also encode an IRO [17,35]. Moreover, if human-restricted pathogens lack an IRO

because human cells truly produce less itaconate, then they are potentially sensitive to it and

itaconate itself could potentially be used as an antimicrobial against them. Our biosensor could

prove invaluable in such studies for determining how much itaconate the bacteria are

encountering, and the self-sufficiency of the biosensor allows it to be employed in a variety of

species, providing added versatility.

In summary, here we present data that itaconate can act as a bactericidal metabolite at

acidic but physiologically relevant pH. We identify the regulatory mechanism of an itaconate

response operon in *Salmonella* and employ its promoter as a novel biosensor of relative

itaconate concentrations in macrophage phagosomes. Finally, we provide further evidence that

IFN-γ stimulates itaconate synthesis and moreover that human cells produce less of the

metabolite than their mouse equivalents.

Materials and Methods

Bacterial strains and plasmids

All *Salmonella* strains used in experiments are derivatives of *Salmonella enterica subsp.*

enterica serovar Typhimurium (*S.* Typhimurium) strain 14028s. As described previously, lambda

red recombination and subsequent P22 phage transduction was used to generate all of the

gene knockout mutants in this background [36–38]. To allow for subsequent recombinations, the

antibiotic resistance cassette was removed from the chromosome using the pCP20 plasmid

encoding FLP recombinase [39]. The heat-unstable pCP20 plasmid was eliminated by passaging

overnight at 42°C and loss was confirmed by antibiotic treatment.

A reporter fusion of the IRO promoter to sfGFP (P$_{IRO}$-sfGFP) was generated using Gibson

cloning to insert the 333bp upstream of the STM3120 start codon (thereby including 25bp

272 upstream of the predicted -35 box and the 5' untranslated region) into the pXG10sf plasmid

273 (replacing the existing promoter)[40–42]. For the reporter construct including *ripR*, the same

274 region was extended to 1570bp upstream of the STM3120 start codon to include the entire

275 STM3121 ORF and a predicted transcriptional terminator following it. For fluorescence

276 microscopy, constitutively expressed (PLtet0-1 promoter) mCherry was inserted into a

277 transcriptionally independent region of the same plasmid. This variation of the plasmid was

278 renamed 'independent constitutive mCherry' or pICM.

280 ***Metabolite induction of*** P_{IRO} ***assay***

281  Induction of the *Salmonella* P_{IRO} promoter was assessed using a transcriptional fusion to

282 sfGFP in either the pXG10sf or pICM plasmids. Data from the two plasmids were combined as

283 the inducible region is identical and the plasmids only differ in the constitutively active mCherry

284 expressed independently on pICM. Overnight LB cultures were used to inoculate (1/200

285 dilution) either LB or MOPS minimal media containing 0.2% of the indicated carbon source.

286 Itaconate or other metabolites at neutral pH were supplemented to a concentration of 0.2%. Of

287 note, for salts and hydrates the final 0.2% concentration reflects the percent of the carbon

288 source itself; e.g. 0.2% succinate was made as 0.47% sodium succinate (dibasic) hexahydrate.

289 Growth was conducted in a TECAN Infinite M200 plate reader at 37°C with shaking and OD_{600}

290 and GFP fluorescence (475nm and 511nm excitation and emission wavelengths respectively)

291 were read every 15 minutes. For clarity, bar graphs show fluorescence at 16h post inoculation.

292 Chloramphenicol was included in all media at a concentration of 20μg/ml to maintain the

293 plasmids.

Acidified media survival

296   LPM media was made as described previously [43,44]. Succinate or itaconate were added

297   to 0.4% and the pH was then adjusted to 4.4, 5.0 or 5.8 as indicated. LB overnight cultures were

298   resuspended in acidified media to an OD of 0.1 and incubated in a 37°C water bath. At time

299   points, samples were taken, serial diluted and plated for colony forming units (CFU).

Intra-macrophage survival

The THP-1 human monocyte cell line and the J774 mouse macrophage cell line were

maintained in RPMI Medium 1640 (with L-glutamine) supplemented with 10% FBS and 1%

Glutamax and grown at 37°C and 5% CO_2. THP-1 cells were seeded in 96-well plates at 50,000

per well with 50nM PMA (phorbol 12-myristate 13-acetate) added to induce differentiation to

adherent macrophage. After 48h, media was replaced with no-PMA growth media overnight

with 100 U/ml human IFN-γ or IL-4 and IL-13 added if indicated. For J774 macrophage the cells

were seeded in 96-well plates at 50,000 per well overnight with 100 U/ml mouse IFN-γ added if

indicated. *Salmonella* in RPMI were added onto seeded cells at a multiplicity of infection (MOI)

of approximately 20:1 and centrifuged for 10 minutes at 1000 rpm to maximize cell contact.

After centrifuging the samples were incubated at 37°C and 5% CO_2 (time 0). After 30 minutes,

cells were washed three times with PBS followed by fresh media containing 100 µg/ml

gentamicin to kill extracellular *Salmonella*. At 2 hours the media was replaced with media

containing gentamicin at 10 µg/ml. At timepoints, intracellular bacteria were recovered using

PBS containing 1% Triton X-100 and vigorous pipetting. Samples were serially diluted and five

10μl spots were plated for CFU counting. Each sample included three separate wells as

technical replicates (a total of 15 x 10μl spots counted per biological replicate).

Fluorescence microscopy

Fluorescence microscopy was conducted similarly to the macrophage survival assay with

some exceptions: Cells were seeded in 24-well plates containing glass coverslips at 125,000 per

well. Bacteria were infected at an MOI of approximately 100 to maximize the instances of

macrophage containing bacteria. At timepoints the media was removed and cells were washed

three times with PBS. They were then fixed for 10 minutes at room temperature in PBS + 4%

paraformaldehyde (PFA). Following three more PBS washes the cells were permeabilized for 10

minutes in PBS + 0.2% Triton X-100 + 1% BSA. Coverslips were washed again, mounted on slides

using 3μl mounting media containing DAPI and allowed to dry overnight in the dark. Slides

where viewed using a Zeiss Observer.z1 microscope using a 100x oil immersion objective and

the Zeiss Zen microscopy software. Images were taken with a Zeiss Axiocam 506 mono camera

mounted on the microscope. For all samples a 2s exposure was used for mCherry and 1s

exposure for sfGFP. ImageJ software was employed for quantification to calculate fluorescence

intensities in the red and green channels relative to a neighbouring background region for each

bacterium and a GFP/mCherry ratio was generated.

Acknowledgements

We would sincerely like to thank Dr. Scott Gray-Owen and members of his lab, in

particular Dr. Ryan Gaudet, for their generous donation of technical expertise, macrophage

cells lines, and use of their equipment. WWN was supported by an Operating Grant from the

Canada Institutes for Health Research (MOP-86683) and a Natural Sciences and Engineering

Research Council (NSERC) of Canada Grant (RGPIN 386286-10). SJH was supported by an NSERC

Vanier Canada Graduate Scholarship.

Figure Legends

Figure 1: Itaconate is bactericidal at low pH. A) At neutral pH, the charges on itaconate inhibit

diffusion across a lipid membrane (left). At acidic pH, itaconate protonates to itaconic acid,

which can traverse the membrane (right). In the cytoplasm, itaconic acid dissociates and

releases protons, trapping it in the cell and acidifying the cytoplasm. **B)** Survival (relative to 0h

time point) of wild-type (WT) or *rpoS* mutant *Salmonella* after 3h in LPM media supplemented

with 0.4% itaconate or succinate (Succ) and adjusted to pH 4.4, 5.0, or 5.8 as indicated. **C)** As in

A but showing survival of *Salmonella aceA* and itaconate response operon (IRO) mutants in LPM

+ itaconate media at pH 4.4. All data are the average of at least three biological replicates and

error bars show one standard deviation. A one-way ANOVA with Sidak's multiple comparison

test was conducted comparing each sample to the same strain at pH 5.8 (B). WT in succinate

was compared to WT at pH 5.8 in itaconate. Mutants in panel C were not significantly different

than their parental strain (WT or Δ*rpoS*). ***, $p < 0.001$.

Figure 2: Overview of the itaconate degradation operon (IRO) and the itaconate-responsive

regulator, *ripR*. *Salmonella* gene loci based on the genome of *Salmonella* Typhimurium LT2 are

shown below. Identities of *ccl*, *ich*, and *ict* are shown above as previously reported [17]. STM3117

appears to be encoded in the same operon however its function remains unknown. In this work

we designate the names '*ripR*' for STM3121, 'itaconate response operon' (IRO) for STM3120-

STM3117, and 'P$_{IRO}$' for the IRO promoter.

Figure 3: RipR is necessary and sufficient to induce P$_{IRO}$ expression in response to itaconate.

Expression of P$_{IRO}$-sfGFP in wild-type (WT) or *ripR* knockout (Δ*ripR*) *Salmonella*, or in *E. coli*.

Figures show GFP fluorescence normalized to optical density at 600nm (OD$_{600}$) after 16h of

growth in LB alone or supplemented with 0.2% itaconate (A) or similar metabolites (B). Data are

the average of at least three biological replicates and error bars show one standard deviation.

pP$_{IRO}$, plasmid-borne transcriptional fusion of P$_{IRO}$ to sfGFP; pRipR-P$_{IRO}$, pP$_{IRO}$ with the *ripR* gene

and its native promoter included on the plasmid. A Games-Howell ANOVA was conducted

comparing with and without itaconate for each strain (A), or comparing WT to Δ*ripR* (B) for

each added metabolite. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$.

Figure 4: P$_{IRO}$ is activated in J774 macrophage and THP-1 macrophage stimulated with IFN-γ.

Expression of P$_{IRO}$-sfGFP in intracellular wild-type (WT) or *ripR* knockout (Δ*ripR*) *Salmonella*

containing the pICM-P$_{IRO}$ plasmid (constitutive mCherry expression). For expression in *E. coli,*

the *ripR* gene was also encoded on the reporter plasmid. '+ IFN-γ' samples were pre-treated

with human IFN-γ. **A)** Representative fluorescence microscopy images. **B)** Relative fluorescence

quantification of individual bacterial particles at 8h post-infection. Number of bacteria

quantified is indicated above and totalled from at least two biological replicates (≥ 3 for all WT

Salmonella samples). Boxes indicate first and third quartiles; central line, median; X, mean;

whiskers, 95[th] percentile; dots, all non-zero data points outside 95[th] percentile. The y-axis

changes scale at 1.5 to better show outliers. A Games-Howell ANOVA was conducted

comparing indicated samples. ***, $p < 0.001$.

Supplemental Figure Legends

Figure S1: pH 4.4 without itaconate was not bactericidal. A) Survival (relative to 0h time point)

of wild-type (WT) or *rpoS* mutant *Salmonella* after 3h in LPM media adjusted to pH 4.4, 5.0, or

5.8 as indicated. **B)** Survival of *Salmonella aceA* and itaconate response operon (IRO) mutants in

LPM + itaconate media at pH 5.0. All data are the average of at least three biological replicates

and error bars show one standard deviation.

Figure S2: Itaconate induction of P$_{IRO}$ occurs in minimal media. Expression of P$_{IRO}$-sfGFP in

wild-type (WT) or *ripR* knockout (Δ*ripR*) *Salmonella* as in Figure 1 except grown in MOPS

minimal media with 0.2% Glucose (**A**; MGlu) or Glycerol (**B**; MGly) instead of LB. Data show GFP

fluorescence normalized to OD$_{600}$ after 16h of growth and are the average of at least three

biological replicates. Error bars show one standard deviation. pP$_{IRO}$, plasmid-borne

transcriptional fusion of P$_{IRO}$ to sfGFP. A Games-Howell ANOVA was conducted comparing WT

to Δ*ripR* for each added metabolite. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$.

Figure S3: P$_{IRO}$ is induced by itaconate in a dose-dependent manner. A) Example replicate

showing *Salmonella* optical density (dashed lines) and P$_{IRO}$-sfGFP fluorescence (solid lines) for a

range of itaconate concentrations across 16h of growth. All conditions were in MOPS minimal

media with 0.2% glucose as a carbon source and indicated concentration of itaconate (%,

g/100ml). **B)** Average P_{IRO}-sfGFP fluorescence (normalized to OD_{600}) at 8h or 16h across two

biological replicates. Error bars indicate one standard deviation. The x-axis changes scale at

0.0075% to show lower concentrations more clearly.

Figure S4: DctA is not required for the response to itaconate. Expression of P_{IRO}-sfGFP in wild-

type (WT) or *dctA* knockout (Δ*dctA*) *Salmonella*. Figure shows GFP fluorescence normalized to

optical density at 600nm (OD_{600}) after 16h of growth in MOPS minimal media with 0.2% glycerol

as the carbon source. Data are the average of at least two biological replicates and error bars

show one standard deviation. pP_{IRO}-sfGFP, plasmid-borne transcriptional fusion of P_{IRO} to sfGFP.

Figure S5: Survival of *Salmonella* mutants in J774 macrophage with and without IFN-γ

stimulation relative to wild-type. The fold change in CFU recovered (24h relative to 2h post-

infection) is shown and data from each replicate was normalized to the 24h/2h ratio obtained

for wild-type *Salmonella*. Columns show the average of three biological replicates and error

bars show one standard error of the mean. Since samples were normalized to wild-type in each

replicate, a one-way ANOVA with Sidak's multiple comparison test was conducted comparing

each strain to the Δ*aceA* strain (A) or to the Δ*rpoS* strain (B). Only the Δ*phoP* mutant was found

to have a $p < 0.05$. *, $p<0.05$; **, $p < 0.01$.

Figure S6: P_{IRO} shows little-to-no activation in THP-1 macrophage stimulated with IL-4 and IL-

13. As in Figure 4, expression of P_{IRO}-sfGFP in intracellular wild-type (WT) or *ripR* knockout

(Δ*ripR*) *Salmonella* containing the pICM-P$_{IRO}$ plasmid (constitutive mCherry expression). THP-1

macrophage were pre-treated with no stimulant, human IFN-γ, or IL-4 and IL-13 and infected

with *Salmonella* for 8h. Data for WT *Salmonella* in THP-1 with no treatment or IFN-γ are shown

in Figure 4 and are shown again here for comparison. As in Figure 4, data shows relative

fluorescence quantification of individual bacterial particles. Number of bacteria quantified is

indicated above and totalled from at least three biological replicates. Boxes indicate first and

third quartiles; central line, median; X, mean; whiskers, 95th percentile; dots, all non-zero data

points outside 95th percentile. The y-axis changes scale at 1.5 to better show outliers.

References

1.   Hansen-Wester, I. & Hensel, M. Salmonella pathogenicity islands encoding type III

secretion systems. *Microbes Infect.* **3**, 549–559 (2001).

2.   Fields, P. I., Swanson, R. V, Haidaris, C. G. & Heffron, F. Mutants of Salmonella

typhimurium that cannot survive within the macrophage are avirulent. *Proc. Natl. Acad.*

Sci. U. S. A. **83**, 5189–5193 (1986).

3.   Bäumler, A. J., Tsolis, R. M., Ficht, T. A. & Adams, L. G. Evolution of Host Adaptation in

Salmonella enterica. *Infect. Immun.* **66**, 4579–4587 (1998).

4.   Bäumler, A. J., Kusters, J. G., Stojiljkovic, I. & Heffron, F. Salmonella typhimurium loci

involved in survival within macrophages. *Infect. Immun.* **62**, 1623–1630 (1994).

5.   Thiennimitr, P. *et al.* Intestinal inflammation allows Salmonella to use ethanolamine to

compete with the microbiota. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 17480–17485 (2011).

6.   Spiga, L. *et al.* An Oxidative Central Metabolism Enables Salmonella to Utilize Microbiota-

Derived Succinate. *Cell Host & Microbe* **22**, 291-301.e6 (2017).

7. Winter, S. E. *et al.* Gut inflammation provides a respiratory electron acceptor for

Salmonella. *Nature* **467**, 426–429 (2010).

8. Cordes, T., Michelucci, A. & Hiller, K. Itaconic Acid: The Surprising Role of an Industrial

Compound as a Mammalian Antimicrobial Metabolite. *Annu. Rev. Nutr.* **35**, 451–473

(2015).

9. Yang, Z., Wang, H., Wang, Y., Ren, Y. & Wei, D. Manufacturing Multienzymatic Complex

Reactors In Vivo by Self-Assembly To Improve the Biosynthesis of Itaconic Acid in

Escherichia coli. *ACS Synth. Biol.* **7**, 1244–1250 (2018).

10. Zhao, M., Lu, X., Zong, H., Li, J. & Zhuge, B. Itaconic acid production in microorganisms.

Biotechnol. Lett. **40**, 455–464 (2018).

11. Hoyt, J. C., Robertson, E. F., Berlyn, K. A. & Reeves, H. C. Escherichia coli isocitrate lyase:

properties and comparisons. *Biochim. Biophys. Acta - Gen. Subj.* **966**, 30–35 (1988).

12. McFadden, B. A. & Purohit, S. Itaconate, an isocitrate lyase-directed inhibitor in

Pseudomonas indigofera. *J. Bacteriol.* **131**, 136–144 (1977).

13. Michelucci, A. *et al.* Immune-responsive gene 1 protein links metabolism to immunity by

catalyzing itaconic acid production. *Proc. Natl. Acad. Sci.* **110**, 7820–7825 (2013).

14. Strelko, C. L. *et al.* Itaconic Acid Is a Mammalian Metabolite Induced during Macrophage

Activation. *J. Am. Chem. Soc.* **133**, 16386–16389 (2011).

15. Naujoks, J. *et al.* IFNs Modify the Proteome of Legionella-Containing Vacuoles and

Restrict Infection Via IRG1-Derived Itaconic Acid. *PLoS Pathog.* **12**, e1005408 (2016).

16. Zhu, X. *et al.* Systemic responses of BALB/c mice to Salmonella typhimurium infection. *J.*

Proteome Res. **13**, 4436–4445 (2014).

17. Sasikaran, J., Ziemski, M., Zadora, P. K., Fleig, A. & Berg, I. A. Bacterial itaconate

degradation promotes pathogenicity. *Nat Chem Biol* **10**, 371–377 (2014).

18. Eriksson, S., Lucchini, S., Thompson, A., Rhen, M. & Hinton, J. C. Unravelling the biology

of macrophage infection by gene expression profiling of intracellular Salmonella enterica.

Mol. Microbiol. **47**, 103–118 (2003).

19. Srikumar, S. *et al.* RNA-seq Brings New Insights to the Intra-Macrophage Transcriptome

of Salmonella Typhimurium. *PLoS Pathog* **11**, e1005262 (2015).

20. Shi, L. *et al.* Proteomic analysis of Salmonella enterica serovar typhimurium isolated from

RAW 264.7 macrophages: identification of a novel protein that contributes to the

replication of serovar typhimurium inside macrophages. *J. Biol. Chem.* **281**, 29131–29140

(2006).

21. Santiviago, C. A. *et al.* Analysis of pools of targeted Salmonella deletion mutants

identifies novel genes affecting fitness during competitive infection in mice. *PLoS Pathog.*

5, e1000477 (2009).

22. Haneda, T., Ishii, Y., Danbara, H. & Okada, N. Genome-wide identification of novel

genomic islands that contribute to Salmonella virulence in mouse systemic infection.

FEMS Microbiol. Lett. **297**, 241–249 (2009).

23. Elder, J. R. *et al.* The Salmonella pathogenicity island 13 contributes to pathogenesis in

streptomycin pre-treated mice but not in day-old chickens. *Gut Pathog.* **8**, 16 (2016).

24. Espinoza, R. A. *et al.* Differential roles for pathogenicity islands SPI-13 and SPI-8 in the

interaction of Salmonella Enteritidis and Salmonella Typhi with murine and human

macrophages. *Biol. Res.* **50**, 5 (2017).

25.   Pédelacq, J.-D., Cabantous, S., Tran, T., Terwilliger, T. C. & Waldo, G. S. Engineering and

characterization of a superfolder green fluorescent protein. *Nat. Biotechnol.* **24**, 79–88

(2006).

26.   Alpuche Aranda C.M., Swanson, J. A., Loomis, W. P. & Miller, S. I. Salmonella

typhimurium activates virulence gene transcription within acidified macrophage

phagosomes. *Proc. Natl. Acad. Sci. U. S. A.* **89**, 10079–10083 (1992).

27.   Rathman, M., Sjaastad, M. D. & Falkow, S. Acidification of phagosomes containing

Salmonella typhimurium in murine macrophages. *Infect. Immun.* **64**, 2765–2773 (1996).

28.   Chakraborty, S., Mizusaki, H. & Kenney, L. J. A FRET-based DNA biosensor tracks OmpR-

dependent acidification of Salmonella during macrophage infection. *PLoS Biol.* **13**,

e1002116 (2015).

29.   Mills, E. L. *et al.* Itaconate is an anti-inflammatory metabolite that activates Nrf2 via

alkylation of KEAP1. *Nature* **556**, 113–117 (2018).

30.   Jha, A. K. *et al.* Network integration of parallel metabolic and transcriptional data reveals

metabolic modules that regulate macrophage polarization. *Immunity* **42**, 419–430

(2015).

31.   Tallam, A. *et al.* Gene Regulatory Network Inference of Immunoresponsive Gene 1 (IRG1)

Identifies Interferon Regulatory Factor 1 (IRF1) as Its Transcriptional Regulator in

Mammalian Macrophages. *PLoS One* **11**, e0149050 (2016).

32.   Jacobson, A. *et al.* A Gut Commensal-Produced Metabolite Mediates Colonization

Resistance to Salmonella Infection. *Cell Host & Microbe* **24**, 296-307.e7 (2018).

33.   Ricke, S. C. Perspectives on the use of organic acids and short chain fatty acids as

515   antimicrobials. *Poult. Sci.* **82**, 632–639 (2003).

516   34.   Tannahill, G. M. *et al.* Succinate is an inflammatory signal that induces IL-1β through HIF-

517   1α. *Nature* **496**, 238–242 (2013).

518   35.   Hammerer, F., Chang, J. H., Duncan, D., Castaneda Ruiz, A. & Auclair, K. Small Molecule

519   Restores Itaconate Sensitivity in Salmonella enterica: A Potential New Approach to

520   Treating Bacterial Infections. *Chembiochem* **17**, 1513–1517 (2016).

521   36.   Schmieger, H. A method for detection of phage mutants with altered transducing ability.

522   *Mol. Gen. Genet. MGG* **110**, 378–381 (1971).

523   37.   Zou, S. B. *et al.* Loss of elongation factor P disrupts bacterial outer membrane integrity. *J.*

524   *Bacteriol.* **194**, (2012).

525   38.   Datsenko, K. A. & Wanner, B. L. One-step inactivation of chromosomal genes in

526   Escherichia coli K-12 using PCR products. *Proc. Natl. Acad. Sci.* **97**, 6640–6645 (2000).

527   39.   Cherepanov, P. P. & Wackernagel, W. Gene disruption in Escherichia coli: TcR and KmR

528   cassettes with the option of Flp-catalyzed excision of the antibiotic-resistance

529   determinant. *Gene* **158**, 9–14 (1995).

530   40.   Gibson, D. G. *et al.* Enzymatic assembly of DNA molecules up to several hundred

531   kilobases. *Nat Meth* **6**, 343–345 (2009).

532   41.   Corcoran, C. P. *et al.* Superfolder GFP reporters validate diverse new mRNA targets of the

533   classic porin regulator, MicF RNA. *Mol. Microbiol.* **84**, 428–445 (2012).

534   42.   Urban, J. & Vogel, J. A green fluorescent protein (GFP)-based plasmid system to study

535   post-transcriptional control of gene expression in vivo. in *Riboswitches* (ed. Serganov, A.)

540, 301–319 (Humana Press, 2009).

43.   Cooper, C. A. *et al.* Structural and biochemical characterization of SrcA, a multi-cargo

type III secretion chaperone in Salmonella required for pathogenic association with a

host. *PLoS Pathog* **6**, e1000751 (2010).

44.   Coombes, B. K., Brown, N. F., Valdez, Y., Brumell, J. H. & Finlay, B. B. Expression and

secretion of Salmonella pathogenicity island-2 virulence genes in response to

acidification exhibit differential requirements of a functional type III secretion apparatus

and SsaL. *J. Biol. Chem.* **279**, 49804–49815 (2004).

Figures

Figure 1: Itaconate is bactericidal at low pH. A) At neutral pH, the charges on itaconate inhibit diffusion across a lipid membrane (left). At acidic pH, itaconate protonates to itaconic acid, which can traverse the membrane (right). In the cytoplasm, itaconic acid dissociates and releases protons, trapping it in the cell and acidifying the cytoplasm. **B)** Survival (relative to 0h time point) of wild-type (WT) or *rpoS* mutant *Salmonella* after 3h in LPM media supplemented with 0.4% itaconate or succinate (Succ) and adjusted to pH 4.4, 5.0, or 5.8 as indicated. **C)** As in A but showing survival of *Salmonella aceA* and itaconate response operon (IRO) mutants in LPM + itaconate media at pH 4.4. All data are the average of at least three biological replicates and error bars show one standard deviation. A one-way ANOVA with Sidak's multiple comparison test was conducted comparing each sample to the same strain at pH 5.8 (B). WT in succinate was compared to WT at pH 5.8 in itaconate. Mutants in panel C were not significantly different than their parental strain (WT or Δ*rpoS*). ***, $p < 0.001$.

Figure 2: Overview of the itaconate degradation operon (IRO) and the itaconate-responsive regulator, *ripR*. *Salmonella* gene loci based on the genome of *Salmonella* Typhimurium LT2 are shown below. Identities of *ccl*, *ich*, and *ict* are shown above as previously reported (17). STM3117 appears to be encoded in the same operon however its function remains unknown. In this work we designate the names '*ripR*' for STM3121, 'itaconate response operon' (IRO) for STM3120-STM3117, and 'P$_{IRO}$' for the IRO promoter.

Figure 3: RipR is necessary and sufficient to induce P_{IRO} expression in response to itaconate. Expression of P_{IRO}-sfGFP in wild-type (WT) or *ripR* knockout (Δ*ripR*) *Salmonella*, or in *E. coli*. Figures show GFP fluorescence normalized to optical density at 600nm (OD_{600}) after 16h of growth in LB alone or supplemented with 0.2% itaconate (A) or similar metabolites (B). Data are the average of at least three biological replicates and error bars show one standard deviation. pP_{IRO}, plasmid-borne transcriptional fusion of P_{IRO} to sfGFP; $pRipR$-P_{IRO}, pP_{IRO} with the *ripR* gene and its native promoter included on the plasmid. A Games-Howell ANOVA was conducted comparing with and without itaconate for each strain (A), or comparing WT to Δ*ripR* (B) for each added metabolite. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$.

Figure 4: P_{IRO} is activated in J774 macrophage and THP-1 macrophage stimulated with IFN-γ.
Expression of P_{IRO}-sfGFP in intracellular wild-type (WT) or *ripR* knockout (Δ*ripR*) *Salmonella* containing the pICM-P_{IRO} plasmid (constitutive mCherry expression). For expression in *E. coli,* the *ripR* gene was also encoded on the reporter plasmid. '+ IFN-γ' samples were pre-treated with human IFN-γ. **A)** Representative fluorescence microscopy images. **B)** Relative fluorescence quantification of individual bacterial particles at 8h post-infection. Number of bacteria quantified is indicated above and totalled from at least two biological replicates (≥ 3 for all WT *Salmonella* samples). Boxes indicate first and third quartiles; central line, median; X, mean; whiskers, 95th percentile; dots, all non-zero data points outside 95th percentile. The y-axis changes scale at 1.5 to better show outliers. A Games-Howell ANOVA was conducted comparing indicated samples. ***, p < 0.001.

Figure S1: pH 4.4 without itaconate was not bactericidal. A) Survival (relative to 0h time point) of wild-type (WT) or *rpoS* mutant *Salmonella* after 3h in LPM media adjusted to pH 4.4, 5.0, or 5.8 as indicated. **B)** Survival of *Salmonella aceA* and itaconate response operon (IRO) mutants in LPM + itaconate media at pH 5.0. All data are the average of at least three biological replicates and error bars show one standard deviation.

Figure S2: Itaconate induction of P$_{IRO}$ occurs in minimal media. Expression of P$_{IRO}$-sfGFP in wild-type (WT) or *ripR* knockout (Δ*ripR*) *Salmonella* as in Figure 1 except grown in MOPS minimal media with 0.2% Glucose (**A**; MGlu) or Glycerol (**B**; MGly) instead of LB. Data show GFP fluorescence normalized to OD$_{600}$ after 16h of growth and are the average of at least three biological replicates. Error bars show one standard deviation. pP$_{IRO}$, plasmid-borne transcriptional fusion of P$_{IRO}$ to sfGFP. A Games-Howell ANOVA was conducted comparing WT to Δ*ripR* for each added metabolite. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$.

Figure S3: P$_{IRO}$ is induced by itaconate in a dose-dependent manner. A) Example replicate showing *Salmonella* optical density (dashed lines) and P$_{IRO}$-sfGFP fluorescence (solid lines) for a range of itaconate concentrations across 16h of growth. All conditions were in MOPS minimal media with 0.2% glucose as a carbon source and indicated concentration of itaconate (%, g/100ml). **B)** Average P$_{IRO}$-sfGFP fluorescence (normalized to OD$_{600}$) at 8h or 16h across two biological replicates. Error bars indicate one standard deviation. The x-axis changes scale at 0.0075% to show lower concentrations more clearly.

Figure S4: DctA is not required for the response to itaconate. Expression of P_{IRO}-sfGFP in wild-type (WT) or *dctA* knockout (*ΔdctA*) *Salmonella*. Figure shows GFP fluorescence normalized to optical density at 600nm (OD_{600}) after 16h of growth in MOPS minimal media with 0.2% glycerol as the carbon source. Data are the average of at least two biological replicates and error bars show one standard deviation. pP_{IRO}-sfGFP, plasmid-borne transcriptional fusion of P_{IRO} to sfGFP.

Figure S5: Survival of _Salmonella_ mutants in J774 macrophage with and without IFN-γ stimulation relative to wild-type. The fold change in CFU recovered (24h relative to 2h post-infection) is shown and data from each replicate was normalized to the 24h/2h ratio obtained for wild-type _Salmonella_. Columns show the average of three biological replicates and error bars show one standard error of the mean. Since samples were normalized to wild-type in each replicate, a one-way ANOVA with Sidak's multiple comparison test was conducted comparing each strain to the Δ_aceA_ strain (A) or to the Δ_rpoS_ strain (B). Only the Δ_phoP_ mutant was found to have a $p < 0.05$. *, $p<0.05$; **, $p < 0.01$.

Figure S6: P$_{IRO}$ shows little-to-no activation in THP-1 macrophage stimulated with IL-4 and IL-13. As in Figure 4, expression of P$_{IRO}$-sfGFP in intracellular wild-type (WT) or *ripR* knockout (Δ*ripR*) *Salmonella* containing the pICM-P$_{IRO}$ plasmid (constitutive mCherry expression). THP-1 macrophage were pre-treated with no stimulant, human IFN-γ, or IL-4 and IL-13 and infected with *Salmonella* for 8h. Data for WT *Salmonella* in THP-1 with no treatment or IFN-γ are shown in Figure 4 and are shown again here for comparison. As in Figure 4, data shows relative fluorescence quantification of individual bacterial particles. Number of bacteria quantified is indicated above and totalled from at least three biological replicates. Boxes indicate first and third quartiles; central line, median; X, mean; whiskers, 95th percentile; dots, all non-zero data points outside 95th percentile. The y-axis changes scale at 1.5 to better show outliers.