1 *Malpighamoeba* infection compromises fluid secretion

2 and P-glycoprotein detoxification in Malpighian tubules

3 Marta Rossi[1], Swidbert R. Ott[2] and Jeremy E. Niven[1,3]

4
[1]School of Life Sciences, University of Sussex, Falmer, Brighton BN1 9QG, UK
[2]Department of Neuroscience, Psychology and Behaviour, University of Leicester, Adrian Building,
University Road, Leicester LE1 7RH, UK
[3]Centre for Computational Neuroscience and Robotics, University of Sussex, Falmer, Brighton BN1
9QG, UK
*Authors for correspondence (M.Rossi@sussex.ac.uk and J.E.Niven@sussex.ac.uk)

Abstract

Malpighian tubules, analogous to vertebrate nephrons, play a key role in insect
osmoregulation and detoxification. Tubules can become infected with a protozoan,
Malpighamoeba, which damages their epithelial cells, potentially compromising their function.
Here we quantify the impact of *Malpighamoeba* infection on fluid secretion and P-glycoprotein-
dependent detoxification by Malpighian tubules of the desert locust using a modified Ramsay
assay. Infected tubules have a greater surface area and a higher fluid secretion rate than
uninfected tubules. Infection also impairs P-glycoprotein-dependent detoxification reducing
the net extrusion of rhodamine B per unit of surface area. However, infected tubules' increased
surface area and fluid secretion rate means they have similar total net extrusion per tubule to
uninfected tubules. Increased fluid secretion rate of infected tubules likely subjects locusts to
greater water stress and imposes increased energy costs. Coupled with reduced efficiency of
P-glycoprotein detoxification, *Malpighamoeba* infection is likely to reduce insect survival within
natural environments.

KEY WORDS: Osmoregulation, Excretion, ABC transporter, Xenobiotic, Honeybee.

28 INTRODUCTION

Osmoregulation and excretion are essential for the survival of animals through the homeostatic
maintenance of the internal environment and the removal of harmful toxins that impair cellular
processes. In insects the Malpighian tubules, which are analogous to the nephrons of the vertebrate
kidney, play an important role in osmoregulation as well as in the removal of metabolic waste and
xenobiotic substances (Maddrell and Gardiner, 1976). Consequently, compromised tubule function may
disrupt osmoregulation and impair detoxification, preventing insects from maintaining their internal
osmotic environment, and exposing their cells to the waste products of metabolism and to
environmental toxins. The impact of such impairment may be particularly severe in those insects that
are exposed to environmental toxins as a consequence of feeding directly on plant tissues or their

products (such as nectar or pollen) because of the presence of secondary metabolites (Despres *et al.*,

2007). Moreover, feeding on plant tissues or their products can expose insects to xenobiotics such as

insecticides, fungicides or antibiotics that may act as toxins (Goulson *et al.*, 2015). Consequently,

compromised tubule function may prevent extrusion of these xenobiotics increasing their effects on

insects.

Insect pathogens can interact with xenobiotics such as pesticides affecting survival (reviewed

in Goulson *et al.*, 2015). Interactions between pathogens and xenobiotics, which have been studied

primarily in honeybees and bumblebees (e.g. Alaux *et al.*, 2010; Vidau *et al.*, 2011), suggest that

pesticide exposure can exacerbate the effects of pathogens. Moreover, pathogens may themselves

exacerbate the impacts of xenobiotics, possibly due to trade-offs between the activity of the immune

system and detoxification pathways (McMillan *et al.*, 2018). In most cases, the mechanisms linking the

susceptibly of insects to xenobiotics with the presence of pathogens remain unknown, even in the case

of diseases of insect pollinators that provide commercially-important ecosystem services (reviewed in

Goulson *et al.*, 2015). Consequently, identifying mechanisms by which pathogens could affect the

xenobiotic compound removal is important for understanding insect immune responses, detoxification

and health.

In insects, including grasshoppers and honey bees, the Malpighian tubules can become infected

by the protozoan *Malpighamoeba* (King and Taylor, 1936; Taylor and King, 1937; Bailey, 1968; Harry

and Finlayson, 1975; Harry and Finlayson, 1976; Liu, 1985). This protozoan develops and multiplies

primarily in the Malpighian tubules before cysts pass into the gut and spread to uninfected individuals

through faeces and food contamination (King and Taylor, 1936; Harry and Finlayson, 1976). Infected

tubules appear swollen and cloudy, and their lumen is packed with cysts (King and Taylor, 1936; Harry

and Finlayson, 1975). As the disease progresses, tubule diameter increases in conjunction with the

thinning of the epithelium and the destruction of the brush border of the epithelial cells (King and

Taylor, 1936; Liu, 1985). Yet, despite the initial characterisation of *Malpighamoeba* infection over 80

years ago (King and Taylor, 1936; Taylor and King, 1937) and descriptions of the disease progression

(Bailey, 1968; Harry and Finlayson, 1975; Liu, 1985), the effect upon the physiology of Malpighian

tubules remains unknown.

Here we compare Malpighian tubule performance of gregarious desert locusts infected with

Malpighamoeba locustae with that of their uninfected counterparts. We focussed on fluid secretion as

well as the removal of alkaloids and hydrophobic organic cations from haemolymph because this

requires active transport into the tubule lumen by P-glycoprotein (P-gp), a member of the ABC

transporter family (Wright and Dantzler, 2004). Desert locust Malpighian tubules express P-gp (Rossi

et al., 2019), likely in the brush border on the apical side of the tubule epithelium (Murray, 1996).

Consequently, the destruction of the brush border caused by *Malpighamoeba* infection may

compromise xenobiotic substance removal. To test this hypothesis, we used a modified Ramsay assay

(Rossi *et al.*, 2019) to assess the performance of locust Malpighian tubules.

MATERIALS AND METHODS

Animals

78 Adult desert locusts (*Schistocerca gregaria*) were taken from a crowded colony maintained at the
79 University of Leicester or were pruchased from Peregrine Livefoods (Essex, UK). Locusts were fed
80 with organic lettuce, fresh wheat seedlings and wheat germ *ad libitum*.

Malpighian tubules, saline and Ramsay assay

Malpighian tubules were removed by cutting the proximal end at ~5 mm from the gut and moved
immediately into a 30 µL drop of saline on a 5 cm Sylgard® coated Petri dish, covered with paraffin
oil, and fixed with steel pins (Supplemental Figure 1) (Rossi *et al*., 2019). Each tubule was punctured
near the proximal end to allow the fluid secretion. The first droplet secreted immediately after
puncturing was discarded. Subsequently, tubules were incubated in saline containing 60 µM rhodamine
B, and secreted droplets were removed every 30 minutes using a 10 µL pipettor and photographed with
a digital camera (Canon EOS 7D; Canon, Tokyo, Japan) mounted on a stereoscopic microscope (Nikon
SMZ-U; Nikon Corp., Tokyo, Japan) (Supplemental Figure 2). Photobleaching of rhodamine B was
prevented using a custom dark box. We measured the volume of each droplet, and estimated the
rhodamine B concentration from the colour intensity using a calibration curve (Rossi *et al*., 2019). At
the end of the assay, we measured tubule diameter and length, and the length of tubule in contact with
the treatment bath. From these measurements, we calculated the surface area of the tubule and the
surface area in contact with the treatment bath. The sample consisted of eight healthy locusts and nine
locusts infected with *Malpighamoeba locustae*. Three Malpighian tubules were removed from each
locust (Supplemental Figure 1). One infected tubule was damaged during the experiment and therefore
excluded.

Statistical Analysis

All statistical analyses were conducted in R version 3.4.1 (R Core Team, 2018). We fitted Linear Mixed
Effect Models (LMEM) by restricted maximum likelihood (REML) estimation (`lmer` function,
package 'lme4' (Bates *et al*., 2015)). To investigate the effect of the infection and incubation time on
the fluid secretion rate (per tubule and per surface area), rhodamine B concentration and rhodamine B
net extrusion (per tubule and per surface area), we analysed the interaction between health (categorical)
and time (categorical). To account for the nested structure of data, we included the individual locust
and date as random intercepts in the model. We also included tubule identity as a random intercept and
time as random slope to account for the repeated measurements on the same individual tubule.
Significances of the fixed effects were determined using Satterthwaite's method for estimation of
degrees of freedom (`anova` function, package 'lmerTest' (Kuznetsova *et al*., 2017)) and non-

significant interactions (P>0.05) were removed. We retained all main effects even if they were not

statistically significant to avoid an increase in the type I error rate (Forstmeier and Schielzeth, 2011).

Estimated marginal means and standard errors (Table 1, Supplemental Table 1) were obtained using the

'emmeans' package in R.

RESULTS

Within the lumen of infected Malpighian tubules, the cysts of the protozoan *Malpighamoeba locustae*

(Amoebidae, Sarcodina) were visible (Figure 1A,B). The wall of infected tubules was thinner and less

clearly defined than that of uninfected tubules (Figure 1A–D). Infected tubules were significantly longer

and broader with a larger surface area than uninfected tubules (Supplemental Table 1A; Figure 1E–G).

The rhythmic movements of the tubules were also slower and less pronounced in infected compared

with uninfected locusts (Supplemental Video 1,2).

Fluid secretion rate was significantly higher in infected than in uninfected tubules, but showed

a similar decrease with time (Table 1A; Figure 2A). A greater surface area was exposed to the

incubation bath in infected compared to their uninfected counterparts (Supplemental Table 1B).

Nevertheless, the fluid secretion rate per surface area was near identical in infected and uninfected

tubules (Table 1B; Figure 2B). The P-glycoprotein substrate Rhodamine B (Eytan *et al.*, 1997) was

actively transported into the lumen of the tubules (Figure 2C) and its concentration in the droplets

secreted increased significantly during incubation in both infected and uninfected tubules, but this

increase depended upon the infection status: the increase in rhodamine B concentration was more

pronounced in droplets collected from uninfected tubules than in droplets from infected tubules (Table

1C; Figure 2D; Supplemental Figure 2). The rhodamine B net extrusion rate per unit of surface area

(moles extruded / (time × area)) was higher in uninfected than in infected tubules; although it increased

during the incubation in both infected and uninfected tubules, this increase over time was more

pronounced in uninfected tubules (Table 1D; Figure 2E). Accounting for the larger surface area of

infected tubules, however, there was no significant difference in the total extrusion per tubule (Table

1E; Figure 2F).

DISCUSSION

By comparing xenobiotic detoxification in infected and uninfected gregarious desert locusts, we

determined how *Malpighamoeba* infection affects the performance of insect Malpighian tubules. In

doing so, our results directly quantify the impact of a common pathogen upon major components of

insect osmoregulation and xenobiotic detoxification, namely the Malpighian tubules and the P-

glycoproteins they express. Infected tubules were longer, wider, had a greater surface area, and showed

less pronounced and slower rhythmic movements than uninfected tubules (Supplemental Video 1,2).

We assessed two aspects of tubule performance; fluid secretion and net extrusion of rhodamine B, a

substrate for P-glycoproteins (Eytan *et al*., 1997). Infected tubules secreted more fluid per time than

uninfected ones. Conversely, the net extrusion of rhodamine B per surface area was reduced in infected

compared with uninfected tubules. The total net extrusion per tubule, however, was similar in both

infected and uninfected tubules.

The greater fluid secretion rate of the infected tubules may be a consequence of their greater

surface area. The tubule epithelium expresses transporters that move Na^+, K^+, and Cl^- ions into the

lumen followed by the passive movement of water (O'Donnell, 2008), so greater surface area may

permit increased ion and water movement. The transepithelial water movement is transcellular, through

aquaporin channels located both on the basolateral and apical epithelial membranes (O'Donnell, 2008).

The greater fluid secretion rate decreases the luminal concentration of rhodamine B and may increase

its net extrusion by preventing back diffusion in the bath solution. In uninfected tubules the fluid

secretion rate positively correlates with the tubule surface area, which in turn correlates with the net

extrusion of rhodamine B (Rossi *et al*., 2019). However, it is possible that loss of the brush border in

infected tubules compromises the septate junctions between epithelial cells (Beyenbach *et al*., 2010)

promoting water movement and increasing fluid secretion.

Trans-tubular rhodamine B transport is predominantly dependent on P-glycoprotein

transporters in the brush border of the tubule epithelial cells (Murray, 1996; Ugwu *et al*., 2016). The

decreased net extrusion of rhodamine B per tubule surface area in infected tubules may therefore reflect

a reduced density of P-glycoproteins caused by the destruction of the brush border by the

Malpighamoeba trophozoites (King and Taylor, 1936; Liu, 1985). This reduced P-glycoprotein density

in infected tubules may be compensated by their greater total surface area compared to uninfected

tubules. The overall consequence of the reduced net extrusion coupled with the increased fluid secretion

rate is that the total net extrusion per tubule is similar in infected and uninfected tubules. This is

consistent with the finding that infection with *Malpighamoeba* does not increase the susceptibility of

grasshoppers (*Melanoplus sanguinipes*) to the insecticide cypermethrin, a P-glycoprotein substrate

(Hinks and Ewen, 1986; Sreeramulu *et al*., 2007).

Changes in feeding behaviour may also contribute to the ability of insects to compensate for

Malpighamoeba impairment of P-glycoprotein detoxification; infection reduces the amount of food that

locusts eat (M. Rossi pers. obs.; King and Taylor, 1936). Reducing food intake may reduce the exposure

of infected insects to noxious substances that their compromised detoxification pathways cannot

remove. This may be particularly important for generalist herbivores like desert locusts that feed on a

broad variety of plant species, many of which contain toxins (Bernays and Chapman, 1978).

Alternatively, reduced food intake in infected locusts may be driven by the activation of the immune

system. Both their immune system and their ability to detoxify noxious food substances rely on

pathways that involve the antioxidant glutathione (Habig *et al*., 1974; Stahlschmidt *et al*., 2015; Jeschke

et al., 2016). Consequently, competition for glutathione between these two physiological pathways

could lead to impaired detoxification when the immune system is already activated (McMillan *et al*.,

2018). Whether due to a direct reduction in feeding or as an indirect consequence of a trade-off with
the immune response, reduced food intake lowers the risk of ingesting toxins but also prevents locusts
acquiring water and nutrients.

A reduced food intake combined with an increased fluid secretion rate poses infected locusts a
difficult challenge. In adaptation to the dry and hot environment that they occupy, desert locusts must
balance their water secretion to avoid dehydration (Phillips, 1964a). The primary urine secreted by the
Malpighian tubules is isosmotic and the composition must be adjusted prior to excretion (O'Donnell,
2008). Most of the filtered water, ions and metabolites are reabsorbed in the anterior hindgut and in the
rectum (O'Donnell, 2008). The reabsorption of Na^+, K^+, Cl^- ions and hence water is an active process
consuming metabolic energy (Phillips, 1964a,b). To prevent desiccation, locusts infected with
Malpighamoeba would need to enhance the water reabsorption to counterbalance the increased fluid
secretion rate from the Malpighian tubules. Thus, infected locusts may be subjected to the increased
energy costs of reabsorbing the excess of water and ions transported into the lumen tubule. Thus,
Malpighamoeba infection may be highly debilitating for desert locusts in their natural environment,
imposing stresses in terms of water, ions, and energy even if they can avoid ingesting plant toxins.

Malpighamoeba infections are not limited to locusts and grasshoppers, occurring in other
insects including honeybees (Bailey, 1968). Although the exact details of *Malpighamoeba* infection
likely depend upon the specific host and pathogen species combination, honeybee Malpighian tubules
infected with *Malpighamoeba mellificae* show striking similarities to locust tubules infected with
Malpighamoeba locustae. In both cases, the lumen of the tubules is swollen and packed with cysts, the
epithelium thins, and the brush border of the epithelial cells is destroyed (King and Taylor, 1936; Bailey,
1968; Harry and Finlayson, 1975; Liu, 1985). Such similarities in tubule pathology despite the
separation of the insect hosts by approximately 380 million years (Misof *et al*., 2014) suggests that
Malpighamoeba infection may have similar consequences in other insects susceptible to infection.
Moreover, infection in honeybees and other insects may similarly compromise detoxification of
xenobiotic compounds and increase water and ion loss. Even though xenobiotic compounds may be
removed from insect haemolymph by infected tubules, our results show a substantial increase in fluid
secretion by infected tubules, which implies greater energy costs for reabsorption in the insect hind gut.
Such impacts are highly injurious and can lead to the premature death of the insect.

Acknowledgements
We thank Laura Corona for comments on an earlier version of the manuscript.

Data accessibility
All supporting data can be found in the electronic supplementary material, tables.

Authors' contributions

M.R. and J.E.N. designed the experiments and data analysis and wrote the paper. M.R.
performed the experiments and data analysis. S.R.O. provided the infected locusts, advised
on data analysis and contributed to writing the manuscript.

Competing interests
The authors declare no competing or financial interests.

Funding
M.R. was funded by a University of Sussex School studentship. S.R.O. was funded by
research project grant BB/L02389X/1 by the BBSRC.

Ethics statement
There are no ethical considerations.

Supplementary information
Supplementary information available online.

References

Alaux, C., Brunet, J.-L., Dussaubat, C., Mondet, F., Tchamitchan, S., Cousin, M., Brillard,
J., Baldy, A., Belzunces, L.P., Le Conte, Y. (2010). Interactions between *Nosema*
microspores and a neoniotinoid weaken honeybees (*Apis mellifera*). *Environ. Microbiol*. **12**,
774-782.

Bailey, L. (1968). Honey bee pathology. *Annu. Rev. Entomol.* **13**, 191–212.

Bates, D., Mächler, M., Bolker, B. and Walker, S. (2015). Fitting linear mixed-effects models
using lme4. *J. Stat. Softw.* **67**, 1–48. (doi:10.18637/jss.v067.i01)

Bernays, E.A. and Chapman, R.F. (1978). Plant chemistry and acridoid feeding behaviour.
In Harborne, J.B. (ed.) Biochemical Aspects of Plant and Animal Coevolution. Academic
Press, London, 99–141.

Beyenbach, K.W., Skaer, H. and Dow, J.A. (2010). The developmental, molecular, and
transport biology of Malpighian tubules. *Annu. Rev. Entomol.* **55**, 351–374.

Despres, L., David, J.P. and Gallet, C. (2007). The evolutionary ecology of insect resistance
to plant chemicals. *Trends Ecol. Evol.* **22**, 298–307.

Eytan, G.D., Regev, R., Oren, G., Hurwitz, C.D. and Assaraf, Y.G. (1997). Efficiency of P-
glycoprotein–mediated exclusion of ehodamine dyes from multidrug-resistant cells is
determined by their passive transmembrane movement rate. *Eur. J. Biochem.* **248**, 104–112.

Forstmeier, W. and Schielzeth, H. (2011). Cryptic multiple hypotheses testing in linear
models: overestimated effect sizes and the winner's curse. *Behav. Ecol. Sociobiol.* **65**, 47–55.

Goulson, D., Nicholls, E., Botias, C., Rotheray, E.L. (2015). Bee declines driven by
combined stress from parasites, pesticides, and lack of flowers. *Science* **347**, 1255957.

SHORT COMMUNICATION

Habig, W.H., Pabst, M.J., Jakoby, W.B. (1974). Glutathione S-transferases the first enzymatic step in mercapturic acid formation. *J. Biol. Chem*. **249**, 7130-7139.

Harry, O.G. and Finlayson, L.H. (1975). Histopathology of secondary infections of *Malpighamoeba locustae* (Protozoa, Amoebidae) in the desert locust, *Schistocerca gregaria* (Orthoptera, Acrididae). *J. Invertebr. Pathol.* **25**, 25–33.

Harry, O.G. and Finlayson, L.H. (1976). The life-cycle, ultrastructure and mode of feeding of the locust amoeba *Malpighamoeba locustae*. *Parasitology*, **72**, 127–135.

Hinks, C.F. and Ewen, A.B. (1986). Pathological effects of the parasite *Malameba locustae* in males of the migratory grasshopper *Melanoplus sanguinipes* and its interaction with the insecticide, cypermethrin. *Entomol. Exp. Appl.* **42**, 39–44.

Jeschke, V., Gershenzon, J., Vassão, D.G. (2016). A mode of action of glucosinolate-derived isothiocyanates: detoxification depletes glutathione and cysteine levels with ramifications on protein metabolism in *Spodoptera littoralis. Insect Biochem. Mol. Biol*. **71**, 37-48. doi:10.1016/j.ibmb.2016.02.002

King, R.L., Taylor, A.B. (1936). *Malpighamœba locustae*, n. sp. (Amoebidae), a protozoan parasitic in the Malpighian tubes of grasshoppers. *Trans. Am. Microsc. Soc.* **55**, 6–10.

Kuznetsova A, Brockhoff P, Christensen R. (2017). lmerTest package: tests in linear mixed effects models. *J. Stat. Softw.* **82**, 1–26.

Liu, T.P. (1985). Scanning electron microscope observations on the pathological changes of Malpighian tubules in the worker honeybee, *Apis mellifera*, infected by *Malpighamoeba mellificae. J. Invertebr. Pathol.* **46**, 125–132.

McMillan, L.E., Miller, D.W., Adamo, S.A. (2018). Eating when ill is risky: immune defense impairs food detoxification in the caterpillar *Manduca sexta. J. Exp. Biol.* **221**, jeb173336.

Maddrell, S.H. and Gardiner, B.O. (1976). Excretion of alkaloids by Malpighian tubules of insects. *J. Exp. Biol.* **64**, 267–281.

Misof, B., Liu, S., Meusemann, K., Peters, R.S., Donath, A., Mayer, C., Frandsen., P.B., Ware, J., Flouri, T., Beutel, R.G. *et al*. (2014) Phylogenomics resolves the timing and pattern of insect evolution**.** *Science* **346**, 763-767. DOI: 10.1126/science.1257570

Murray, C.L. (1996). A P-glycoprotein-like mechanism in the nicotine-resistant insect, *Manduca sexta*. PhD thesis, University of Ottawa (Canada).

O'Donnell, M. (2008). Insect excretory mechanisms. *Adv. Insect Physiol.* **35**, 1–122.

Phillips, J.E. (1964a). Rectal absorption in the desert locust, *Schistocerca gregaria* Forskål. I. Water. *J. Exp. Biol*. **41**, 15–38.

Phillips, J.E. (1964b). Rectal absorption in the desert locust, *Schistocerca gregaria* Forskål: II. Sodium, potassium and chloride. *J. Exp. Biol.* **41**, 39–67.

Rossi, M., De Battisti, D. and Niven, J.E. (2019). P-glycoprotein detoxification by the Malpighian tubules of the desert locust. *bioRxiv*, (doi: 10.1101/529750).

Sreeramulu, K., Liu, R. and Sharom, F.J. (2007). Interaction of insecticides with mammalian P-glycoprotein and their effect on its transport function. *Biochim. Biophys. Acta, Biomembr.* **1768**, 1750–1757.

296 **Stahlschmidt, Z.R., Acker, M., Kovalko, I., Adamo, S.A.** (2015). The double-edged sword
297 of immune defence and damage control: do food availability and immune challenge alter the
298 balance? *Funct. Ecol*. **29**, 1445-1452. doi:10.1111/1365-2435.12454
299
Taylor, A.B., King, R.L. (1937). Further studies on the parasitic amebae found in
grasshoppers. *Trans. Am. Microsc. Soc.* **56**, 172–176.

Ugwu, M.C., Oli, A., Esimone, C.O. and Agu, R.U. (2016). Organic cation rhodamines for
screening organic cation transporters in early stages of drug development. *J. Pharmacol.*
Toxicol. Methods **82**, 9–19.

Vidau, C., Diogon, M., Aufauvre, J., Fontbonne, R., Brunet, J.-L., Texier, C., Biron, D.G.,
Blot, N., El Alaoui, H., Belzunces, L.P., Delbac, F. (2011). Exposure to sublethal doses of
fipronil and thiacloprid highly increases mortality of honeybees previously infected by
Nosema ceranae. PLoS ONE **6**, e21550.

Wright, S.H. and Dantzler, W.H. (2004). Molecular and cellular physiology of renal organic
cation and anion transport. *Physiol. Rev.* **84**, 987–1049.

Figure 1. The Malpighian tubules of Malpighamoeba-infected locusts differ from those of uninfected locusts in appearance, size, and surface area. (A) Infected tubules are swollen and cloudy. (B) The lumen of infected tubules is filled with cysts. (C) Uninfected tubules are thinner and more transparent. (D) The lumen of uninfected tubules is free from cysts. A tracheole (dark line) wraps around the uninfected tubule. Infected locusts had longer (E) and wider (F) tubules than uninfected locusts. (G) Consequently, the surface area of infected tubules was

317   greater than that of uninfected tubules. Large filled circles with error bars represent mean

318   estimates from mixed effect model fits with the standard error. Small circles represent the raw

319   data.

Figure 2. Malpighamoeba infection affects the fluid secretion rate and rhodamine B extrusion of Malpighian tubules. Infected tubules secreted more fluid than the uninfected ones (A), whereas the secretion rate (SR) per unit of surface area did not differ between infected and uninfected Malpighian tubules (B). (C) The uninfected tubule shown in 1D transporting rhodamine B. (D) The rhodamine B concentration of the droplets secreted increased over time. Uninfected tubules secreted more concentrated droplets than the infected tubules. (E,F) The net extrusion rate of rhodamine B increased over time. The net extrusion rate per unit of surface area was higher in the healthy tubules than in the infected ones (E), however,

considering the whole surface area, the net rhodamine B extrusion rate per tubule was similar

in healthy and infected tubules (F). Large filled circles with error bars represent means

estimates from mixed effect model fits with the standard error. Small circles represent the raw

data.

Table 1. ANOVA results and estimated marginal means of fluid secretion rate per tubule (A) and per unit of surface area (B); rhodamine B

concentration of the droplet secreted (C), and net extrusion of rhodamine B per unit of surface area (D) and per tubule (E) for infected and

uninfected tubules at each time point of observation.

		Type II ANOVA (Satterthwaite's method)					Group means ± s.e. (model estimates)		
			num. df	den. df	F value	p		30 min	60 min	90 min
A	Fluid secretion rate per tubule (nL/min)	Time:	2	53.05	8.32	**<.001**	Uninfected	0.65 ± 0.30	0.36 ± 0.29	0.28 ± 0.29
		Health:	1	11.67	9.11	**0.011**	Infected	1.62 ± 0.25	1.34 ± 0.24	1.26 ± 0.24
B	Fluid secretion rate per unit of surface (nL/min mm^2)	Time:	2	53.91	14.49	**<.001**	Uninfected	0.27 ± 0.05	0.19 ± 0.05	0.17 ± 0.05
		Health:	1	10.33	0.07	0.803	Infected	0.28 ± 0.05	0.20 ± 0.04	0.18 ± 0.04
C	Rhodamine concentration (µM)	Time x Health:	2	62.03	16.21	**<.001**	Uninfected	77.6 ± 11.0	210.0 ± 16.7	299.9 ± 26.0
							Infected	19.2 ± 9.4	50.1 ± 15.3	86.7 ± 24.7
D	Net rhodamine extrusion per unit of surface (fmol/min mm^2)	Time x Health:	2	52.47	8.13	**<.001**	Uninfected	23.0 ± 5.4	38.0 ± 5.8	49.4 ± 6.5
							Infected	3.0 ± 4.5	8.1 ± 5.0	12.6 ± 5.7
E	Net extrusion of rhodamine per tubule (fmol/min)	Time:	2	51.72	31.62	**<.001**	Uninfected	53.7 ± 14.8	85.1 ± 15.6	113.7 ± 17.1
		Health:	1	12.78	3.04	0.105	Infected	24.6 ± 12.3	55.9 ± 13.3	84.5 ± 15.0

336