Two different epigenetic pathways detected in wild three-spined sticklebacks

are involved in salinity adaptation

Melanie J. Heckwolf1*†, Britta S. Meyer1†‡, Robert Häsler2, Marc P. Höppner2, Christophe

Eizaguirre3 and Thorsten B. H. Reusch1.

1 Marine Evolutionary Ecology, GEOMAR Helmholtz Centre for Ocean Research Kiel, Kiel,

Germany.

2 Institute of Clinical Molecular Biology, Kiel University, Kiel, Germany.

3 School of Biological and Chemical Sciences, Queen Mary University of London, London,

UK.

* corresponding author. Email: mheckwolf@geomar.de.

† These authors contributed equally to this work.

‡ present address: Max Planck Research Group Behavioural Genomics, Max Planck Institute

for Evolutionary Biology, Plön, Germany.

Keywords

DNA methylation, epigenetics, *Gasterosteus aculeatus*, local adaptation, phenotypic

plasticity, reduced representation bisulfite sequencing, transgenerational acclimation

Abstract

While environmentally inducible epigenetic marks are discussed as one mechanism of transgenerational plasticity, environmentally stable epigenetic marks emerge randomly. When resulting in variable phenotypes, stable marks can be targets of natural selection analogous to DNA sequence-based adaptation processes. We studied both postulated pathways in natural populations of three-spined sticklebacks (Gasterosteus aculeatus) and sequenced their methylomes and genomes across a salinity cline. Consistent with local adaptation, populations showed differential methylation (pop-DMS) at genes enriched for osmoregulatory processes. In a two-generation experiment, 62% of these pop-DMS were insensitive to salinity manipulation, suggesting that they could be stable targets for natural selection. Two-thirds of the remaining inducible pop-DMS became more similar to patterns detected in wild populations from the corresponding salinity, and this pattern accentuated over consecutive generations, indicating a mechanism of adaptive transgenerational plasticity. Natural DNA methylation patterns can thus be attributed to two epigenetic pathways underlying the rapid emergence of adaptive phenotypes in the face of environmental change.

Introduction

Recent advances in epigenetics have started to challenge our understanding of genetic diversity, inheritance and adaptive evolution (1-3). It has been suggested that epigenetic modification, for example via DNA methylation, histone modification or small RNAs, provides an additional pathway to create phenotypic diversity and ultimately contribute to rapid evolutionary adaptation (4-6). Several theoretical models posit that the heritable proportion of these molecular modifications can contribute to adaptation via two distinct information pathways (5, 7, 8). Firstly, selection-based epigenetic marks would emerge as spontaneous epimutations that remain stable across subsequent generations, although their overall stability seems to be 2-3 orders of magnitude lower as DNA base changes (7, 9). Similar to

adaptation from DNA sequence-based variation these epimutations can underlie phenotypes

that are targets of natural selection and thereby carry information on past selection regimes

without directly responding to the current environment (5, 8, 10). Secondly, detection-based

effects describe inducible epigenetic marks at defined genomic locations which are under

environmental control (7). This way, the inheritance of information on environmental cues

experienced by previous generations represents one rapid and reliable mechanism

underlying transgenerational plasticity which is hypothesized to buffer populations under

sudden environmental change (7, 11). Distinguishing between these mechanisms is crucial,

since they have very different evolutionary implications. While stable epigenetic marks are in

line with key tenets of the Modern Synthesis including random variation and genetic

inheritance, directional processes via inducible epigenetic marks are better captured by the

Extended Evolutionary Synthesis and its assumptions on non-random phenotypic variation

and inclusive inheritance (12). Given an ongoing and sometimes controversial discussion of

whether or not there is a need of an extended evolutionary synthesis (2), providing empirical

evidence of both pathways and assessing their proportional contribution to epigenetic

inheritance in nature is paramount. While the significance and principal difference of both

transmission pathways is clear (4, 5, 7), empirical evidence on the proportional existence and

adaptive value of these pathways is lacking. One prime objective of this study was thus to

assess whether these two epigenetic pathways can be detected in nature and to test whether

short term acclimation responses match patterns of DNA methylation variation of natural

populations. If transgenerational experiments result in DNA methylation profiles closer to

those of locally adapted natural populations, this would provide evidence that DNA

methylation is mechanistically involved in adaptive transgenerational plasticity.

Studying the adaptation to ocean salinity is particularly well suited to identify

selection-based and detection-based effects because spatio-temporal patterns in ocean

salinity are rather stable compared to for instance temperature. Since salinity change

imposes strong physiological stress with well-defined cellular effects (13), natural salinity

gradients offer unparalleled opportunities to use local patterns of epigenetic variation as

background against which direction and magnitude of experimental salinity manipulations

can be tested. One suitable ecosystem to follow such a space-for-time approach is the Baltic

Sea, a semi-enclosed marginal sea that has been dubbed a "time machine" for many

predicted perturbations associated with global change (14).

Taking advantage of the Baltic Sea salinity gradient, we sequenced the methylomes

(reduced representation bisulfite sequencing, RRBS) as well as whole genomes of three-

spined sticklebacks (Gasterosteus aculeatus) from three locally adapted populations (15) in-

and outside the Baltic Sea (6, 20 and 33 PSU). In this paper we focus on the patterns of

(epi)genomic variation, while transgenerational phenotypic effects in the same populations

have been described previously (16). Some key response variables are presented here to

validate adaptive transgenerational plasticity.

Baltic stickleback populations are genetically differentiated from one another (genome

wide average pairwise FST = 0.028 (15)) and show patterns consistent with local adaptation

to salinity regimes in controlled common garden experiments (16, 17). Furthermore,

sticklebacks are known for their adaptive transgenerational plasticity in response to variation

in temperature (18) and changes in DNA methylation levels at osmoregulatory genes in

response to within generation salinity acclimation (19, 20). However, it remains unclear

whether DNA methylation mediate adaptive transgenerational plasticity, one possible

mechanism allowing adaptive phenotypes to rapidly emerge in the face of environmental

change. To address this question, we complemented our field survey with a two-generation

salinity acclimation experiment using the mid salinity population (20 PSU) to quantify the

proportion of stable (potentially selection-based) and inducible (potentially detection-based)

DNA methylation within- and across generations (Figure 1). We focused on the methylation

of cytosines at cytosine-phosphate-guanine dinucleotides (CpG sites), the most common

methylation motif in vertebrates (21) with partial inheritance potentially facilitating adaptation

in natural populations (11).

We tested three inter-related hypotheses: (i) Stickleback populations originating from

different salinity areas (6, 20 and 33 PSU) show differentially methylated CpG sites

(hereafter pop-DMS), consistent with patterns of local adaptation. (ii) Such pop-DMS include

both types of methylation sites: experimentally stable sites (potentially selection-based) and

experimentally inducible sites (potentially detection-based). (iii) Upon transgenerational

salinity acclimation, inducible DNA methylations become more similar to the patterns of

natural populations at corresponding salinities. When associated with beneficial phenotypic

effects, the latter findings would be evidence of a mechanism of adaptive transgenerational

plasticity.

Results

Identifying differentially methylated CpG sites between stickleback populations along

a natural salinity cline

Differentially methylated CpG sites between stickleback populations (hereafter pop-DMS)

were determined via reduced representation bisulfite sequencing (RRBS) in 46 wild caught

sticklebacks from 3 different sites that varied in average salinity (Sylt, 33 practical salinity

units = PSU; Kiel, 20 PSU; Nynäshamn, 6 PSU; Figure 1). After quality and coverage

filtering, we obtained 525,985 CpG sites present in all groups, corresponding to ~4% of all

CpG sites in the stickleback genome. Between pairs of wild caught populations, we detected

1,470 (comparison 20 vs. 6 PSU) and 1,158 (20 vs. 33 PSU) pop-DMS according to our

significance threshold. The distribution of these sites was random with regard to the genomic

features (promoter, exon, intron, and intergenic; 20 vs. 6 PSU: $X2_3$ = 3.36, P = 0.340; 20 vs.

33 PSU: $X2_3$ = 1.61, P = 0.656; Supplementary Material: Table S1) and chromosomal

regions (Supplementary Figure S1A). Among these pop-DMS, 1,098 (20 vs. 6 PSU) and 871

(20 vs. 33 PSU) were located close to (< 10 kb from transcription start sites) or within genes

and thereby associated with 655 and 510 genes, respectively. Many of these genes were

involved in basic biological processes such as DNA repair and strand renaturation, as well as

chromosome condensation and separation (Supplementary Figure S2). Particularly relevant

and concordant with previous findings of local salinity adaptation (15), these genes were

enriched for osmoregulatory processes such as ion transport and channel activity, renal

water homeostasis and absorption, as well as urine volume regulation (Figure 2). Genes

associated with ≥ 10 pop-DMS are listed in Table 1 (for all genes, see Supplementary Table

S2A and S2B).

Characterizing stable and inducible DNA methylation in a two-generation salinity

acclimation experiment

In order to distinguish between stable and inducible DNA methylation we then conducted a

two-generation salinity acclimation experiment with laboratory bred sticklebacks from the mid

salinity population (Figure 1). We considered pop-DMS to be stable when both the within-

and the transgenerational acclimation group were not differentially methylated compared to

the control group (q-value ≥ 0.0125). On the other hand, if a pop-DMS was differentially

methylated between at least one of the acclimation groups (within- and transgenerational)

compared to the control group (q-value < 0.0125; methylation difference ≥ 15%) this site was

considered inducible. Pop-DMS with a significant q-value not exceeding the threshold of

differential DNA methylation (15%) were treated as a separate category, hereafter

inconclusive. After two generations of salinity acclimation, we found that the majority of the

pop-DMS remained stable, regardless of the direction of salinity change (926 pop-DMS =

63% at decreased salinity; 694 pop-DMS = 60% at increased salinity). A smaller number of

pop-DMS (13%) were inducible, as they showed a significant change in CpG methylation

upon experimental salinity decrease (198 pop-DMS) or increase (148 pop-DMS). An

additional 24% and 27% (346 and 316 pop-DMS respectively) differed significantly between

experimental treatment groups, but did not exceed the minimum threshold in differential DNA

methylation of 15% employed in this study. The number of inducible pop-DMS (13%) derived

from comparisons between natural populations was significantly higher compared to what

would be expected from a random subset of CpG sites across the genome (< 1%; 1000

replicates; salinity decrease: X_2^2 = 1090.7, P < 0.001; salinity increase: X_2^2 = 967.7, P <

0.001).

Stable and inducible methylated sites are associated with different functional gene categories

We then assessed both stable and inducible pop-DMS for associations with different putative gene function (Figure 2, Supplementary Figure S2). Genes associated with stable pop-DMS (452 and 329 under salinity decrease and increase, respectively) were enriched for basic biological processes (e.g. DNA repair and chromosome separation; Supplementary Figure S2), but also for osmoregulatory functions (e.g. ion channel activity; Figure 2). Furthermore, under increased salinity, many metabolic processes were among the stable pop-DMS (Figure 2). Inducible pop-DMS were associated with genes (100 and 82 under salinity decrease and increase, respectively) that were primarily enriched for other osmoregulatory functions regulating ion transmembrane transport (Figure 2, Supplementary Figure S2). Therefore, stable and inducible pop-DMS are not only affecting different genes but also different gene ontologies with very little overlap (Figure 2).

Assessing the role of inducible DNA methylation in nature

Next, we investigated whether multiple pop-DMS associated with the same gene showed a correlated response to experimental salinity acclimation, which would result in a non-random distribution of pop-DMS within genes among the categories 'stable', 'inducible' and 'inconclusive'. We found that in 13 out of 20 genes with ≥ 10 pop-DMS, these pop-DMS responded in a synchronous way across the gene upon salinity acclimation (Table 1, Fisher's exact test, $P < 0.05$), suggesting that genomic sites are pre-defined and not randomly hit by epi-mutations. Secondly, we tested whether a change at inducible pop-DMS in experimental fish increased the similarity to methylation patterns in natural populations locally adapted to their salinity conditions. Of the 198 (decreased salinity) and 148 (increased salinity) inducible pop-DMS, 130 (66%) and 101 (68%), respectively, became more similar to methylation levels of the field population locally adapted to the corresponding salinity (hereafter

'expected' direction). Conversely, at 68 (34%, decreased salinity) and 47 (32%, increased

salinity) inducible pop-DMS, experimental fish showed methylation changes in the opposite

direction, meaning the similarity to methylation levels observed in the natural populations

was reduced (hereafter 'opposite' direction).

Since genomic variation can have a strong cis-regulatory impact on epigenomic

variation (22), populations may respond with similar DNA methylation changes to salinity at

regions of low genetic population divergence, while at regions of high genetic population

divergence DNA methylation changes might differ between populations. Thus, we

hypothesized that opposite inducible pop-DMS are associated with higher genomic (DNA

sequence-based) differentiation, while we anticipated the reverse at expected inducible pop-

DMS. Accordingly, we re-sequenced whole genomes of the same wild caught individuals we

used for RRBS and calculated the degree of genomic differentiation per inducible pop-DMS

as mean FST value (± 5 kb window) between populations. In line with our hypothesis, the

populations from Kiel and Nynäshamn (decreased salinity) were genetically more

differentiated at opposite inducible pop-DMS than at expected sites (δ.mean.FST = -0.014, P

= 0.002; Figure 3A and 3C). A similar trend, yet not significant, was found between the

populations from Kiel and Sylt (increased salinity: δ.mean.FST = -0.005, P = 0.153; Figure

3B and 3D). To understand if selection might have played a role in shaping the differences

between increased and decreased salinity we tracked survival rates from fertilized eggs to

the three months old offspring and compared them between treatment groups. Mortality

differed significantly between the treatment groups (GLMM, X^2_4 = 66.159, P < 0.001; Figure

4A, Supplementary Material: Table S3A) with increased mortality under increased salinity,

while mortality under decreased salinity was generally low and did not differ from the control

group (Figure 4A, Supplementary Material: Table S3A).

Comparing within- and transgenerational acclimation effects on inducible DNA

methylation

We then tested whether or not DNA methylation followed patterns predicted under adaptive

transgenerational plasticity, namely that salinity acclimation over two instead of only one

consecutive generations would enhance the similarity at inducible pop-DMS with patterns

found among wild populations at corresponding salinities. To do so, we calculated the

percentage match (δ.meth.diff, Figure 3E-H) between the within- and transgenerational

acclimation groups in relation to the anticipated adaptive methylation level at inducible pop-

DMS. In line with our hypothesis, we found that transgenerational compared to within-

generation salinity manipulation increased the δ.meth.diff (for 'expected' inducible

methylation: decreased salinity: $F_{1,256} = 30.42$, $P < 0.001$; increased salinity: $F_{1,198} =$

10.39, $P = 0.001$; Figure 3E and F). Remarkably, under decreased experimental salinity, we

found an interaction of 'methylation direction' (hyper- or hypomethylation) and 'acclimation'

(within- and transgenerational) affecting the δ.meth.diff (ANOVA, δ.meth.diff ~ methylation

direction * acclimation, $F_{1,256} = 7.69$, $P = 0.006$; Figure 3E). Here, transgenerational

acclimation increased the similarity of hypomethylated sites to methylation levels found in

natural populations, while hypermethylated sites showed similar values within- and across

generations (Figure 3E). While for 'expected' inducible sites this effect was only present

under decreased salinity, at 'opposite' inducible sites transgenerational acclimation to

decreased and increased salinity enhanced the δ.meth.diff at hypomethylated sites (Figure

3G and H; ANOVA, δ.meth.diff ~ methylation direction * acclimation, decreased salinity:

$F_{1,132} = 19.89$, $P < 0.001$; increased salinity: $F_{1,90} = 9.85$, $P = 0.002$).

To infer the effect of DNA methylation differences on offspring fitness, we compared

fitness correlated proxies between control, within- and transgenerational acclimation groups.

Given that osmoregulation is energetically very costly we assessed the total weight, standard

length (SDL) and the hepatosomatic index (HSI) as a proxy for energy reserves in form of

liver glycogen storage. SDL (GLMM, $X^2_4 = 9.965$, $P = 0.041$; Figure 4B, Supplementary

Material: Table S3B) and total weight (GLMM, $X^2_4 = 11.518$, $P = 0.021$; Figure 4D,

Supplementary Material: Table S3D) differed only slightly between the treatment groups.

Highly significant differences between the treatment groups were detected for the HSI

(GLMM, X^2_4 = 22.688, P < 0.001; Figure 4C, Supplementary Material: Table S3C) with an

increased HSI under decreased salinity compared to the control group. This supports

previous findings, showing that osmoregulation at 6 PSU is energetically less demanding

(16, 17). Under increased salinity the HSI was lower compared to decreased salinity in the

within-generation acclimation group, while a transgenerational acclimation to increased

salinity partially removed this difference. Even though this effect was not significant, we

observed a trend towards higher mean HSI in the transgenerational acclimation group

compared to the within-generational acclimation group at the same salinity (Figure 4C,

Supplementary Material: Table S3C).

Discussion

This study investigated whether two postulated pathways of epigenetic inheritance

(selection-based and detection-based) can be identified in natural populations, and whether

the functions of the associated genes indicate a contribution to salinity adaptation in three-

spined sticklebacks. Consistent with modeled selection dynamics for selection-based DNA

methylation sites (7), we identified stable pop-DMS between populations enriched for

osmoregulatory functions. As suggested by models on the role of epigenetic and genetic

changes in adaptive evolution (5), such randomly originating DNA methylation sites could

serve as potential targets for natural selection, resulting in differential DNA methylation of

osmoregulatory genes between locally adapted populations. Phenotypic variation resulting

from stable DNA methylation sites with expected epimutation rates of approximately 10^{-4}

(estimated for A. thaliana (9)) could allow populations to explore the fitness landscape faster

than under DNA sequence based genetic variation alone (mutation rate $\sim 10^{-8}$) (5, 23).

Despite the random nature of such epimutations and the high gene flow among stickleback

populations (15), the regionally synchronized signal of methylation differentiation in proximity

to and within osmoregulatory genes suggests to be driven by local salinity environments.

Furthermore, since local adaptation is 10-times more likely to involve changes at the gene

expression than at the amino acid sequence level (24, 25), it is conceivable that differential

DNA methylation and consequently different regulation of osmoregulatory genes may

contribute to local adaptation to salinity. Remarkably, one of the top candidate genes

differentially methylated between populations from 20 and 6 PSU was eda (Ectodysplasin A),

a well-described gene involved in lateral plate formation (26). Salinity and calcium are

significant drivers of plate morphology (27) in proposed conjunction with predation (28). Our

findings suggest that repeated and parallel selection for the low plated eda allele in response

to low saline habitats (29-31), including the Baltic Sea (15, 32), may not only be due to DNA

base pair changes but also involve methylation mechanisms. Previous studies have

suggested that energetic cost for Baltic sticklebacks increase with increasing difference

between treatment and isosmotic salinity conditions (~11 PSU (33)) (16, 17). In line with

these findings, we observed many metabolic processes to be associated with stable pop-

DMS under increased salinity, which is also reflected in the lower hepatosomatic index under

increased salinity. Taken together, our results suggest that, along with genetic differentiation,

the stable fraction of differentially methylated genes follow patterns consistent with local

salinity adaptation across stickleback populations (Figure 2, Table 1; Supplementary

Material: Figure S2, Table S2A and S2B). Previous studies have shown that patterns of local

adaptation in DNA methylation can have a genomic basis in form of cis- and trans-acting

genomic loci (22, 34). Whether the pop-DMS in this study represent an independent

mechanism for local adaptation or whether they are a consequence of DNA-sequence based

genetic differentiation would need further functional genomic investigation.

For the second postulated pathway on detection-based epigenetic inheritance (7), we

identified a significantly higher number of experimentally inducible pop-DMS than expected

by chance. Interestingly, multiple DMS associated with the same gene were mostly

synchronized in their response (Table 1). Their association with different osmoregulatory

genes compared to stable pop-DMS reflects a salinity-mediated plastic response potentially

contributing to the molecular basis of adaptive phenotypic plasticity by allowing individuals to

regulate their ion balance relative to the seawater medium instantaneously without requiring

any further genetic adaptation at the population level. Interestingly, more than two-thirds of

the inducible pop-DMS changed methylation to become more similar to the population locally

adapted to the corresponding salinity. Remarkably, the similarity of these pop-DMS

methylation levels between the naturally adapted and the experimentally acclimated

population increased across generations. This strongly suggests that adaptive

transgenerational plasticity via DNA methylation plays a role in salinity acclimation in wild

animal populations that is changing incrementally with increasing exposure time to the

environmental cue. Since we used a split-clutch design for the breeding experiment and

mortality levels within the salinity decrease experiment were not different, we can assume

that these groups have a similar genomic background. Therefore, we can rule out that most

of the inducible pop-DMS simply follow patterns of genomic differentiation and assume that

these sites are independently responding to the existing environmental pressure.

The induction of methylation sites has been widely discussed as potential buffer for

environmental changes (11, 18, 35), which would allow populations to persist in the face of

disturbances by moving phenotypes in the direction of optimal fitness faster than genetic

changes alone (5, 23). Further work should be directed as to whether the inducible

methylation changes would also become persistent and thus move effectively into the stable

pop-DMS category, or whether these changes are reversible. We observed that the potential

for adaptive transgenerational plasticity differed among methylation directions (hypo- and

hypermethylated sites, Figure 3E, G and H), with a higher potential for transgenerational

plasticity at hypomethylated sites. Generally, the spontaneous addition of a methyl-group to a

cytosine is 2.5 times more likely than the removal (23) making a targeted de-methylation

more difficult. In the transgenerational acclimation group, the methylation reprogramming

including extensive methylation erasure and de novo methylation during gamete formation

and zygote development could serve as a mechanistic basis to enhance de-methylation of

CpG sites (36, 37).

The genetic background is considered one important source for epigenomic variation

via cis- and trans-regulatory mechanisms (22, 38, 39). Genomic characterization around

each experimentally inducible pop-DMS revealed a relationship between the level of

population genetic differentiation (FST) and the propensity of the experimental population to

approach the methylation level of the low salinity population (NYN) under experimental

salinity decrease (Figure 3A and C). Here, experimentally induced DNA methylation

becomes more similar to the methylation in natural populations only in genomic regions with

low genetic differentiation. On the other hand, when experimentally induced methylation

differences to the low salinity population increase (Figure 3A and C), this occurs in a more

divergent genomic background, suggesting that the genome has undergone past selection

leading to DNA-based local adaptation, rendering epigenetic modifications less relevant (5).

This finding underlines the importance of incorporating the genomic background in the

interpretation of DNA methylation patterns. Nevertheless, it remains to be tested what

happens to all induced DNA methylation sites with selection over multiple generations.

Taken together, our study provides the first empirical evidence that stable and

inducible DNA methylation exist in wild animal populations and follow predictions from

evolutionary theory according to a selection- and a detection-based epigenetic pathway to

promote adaptation to novel environments (5, 7) (Figure 5). The latter process, in particular,

may provide an extremely fast (only two generations) mechanism to promote adaptive

phenotypes which may be particularly important under predicted climate change (Figure 5).

While the selection-based pathway follows the core assumptions of the Modern Synthesis

(random variation and genetic inheritance), the detection-based pathway is better captured

by the Extended Evolutionary Synthesis (non-random variation and inclusive inheritance).

Because their evolutionary implications are very different, any future transgenerational or

epigenetic studies should distinguish between these pathways, which will also contribute

urgently needed data to the debate on an extended evolutionary synthesis. Whether

epigenetic marks such as differentially methylated sites can permanently be attributed to one

of the two categories or rather represent a continuum of stability levels and directionality will

need further experimental testing over multiple generations.

Materials and Methods

Animal Welfare

All catches were performed under legal authorization issued by the German 'Ministry of Energy Transition, Agriculture, Environment, Nature and Digitalization' in Schleswig-Holstein (MELUR – V242-7224.121-19), by the Danish 'Ministry of Food, Agriculture and Fisheries of Denmark' (Case no: 14-7410-000227), by the Estonian 'Ministry of the Environment' (Keskkonnaministeerium - eripüügiluba nr 28/2014) and by the Swedish Sea and Water Authority (Havs och Vattenmyndigheten). Ethical permission for the experiments required by German law was given by the MELUR: V312-7224.121-19).

Survey and experimental design

For the field survey, we collected juvenile three-spined sticklebacks (Gasterosteus aculeatus; 31.68 ± 14.25 mm) from three different salinity regimes inside and outside the Baltic Sea (Sylt (SYL), Germany (55°00'58.3''N, 8°26'22.0''E), 33 PSU, N = 16; Kiel (KIE), Germany (54°26'11.8"N 10°10'20.2"E), 20 PSU, N = 16; Nynäshamn (NYN), Sweden (58°52'44.7"N 17°56'06.2"E), 6PSU, N = 16) in September 2014. Once collected, fish were immediately euthanized using tricaine methane sulfonate solution (MS222), photographed, measured (length and total weight) and stored in RNA-later (24h at 7°C, afterwards at -20°C). A cut along the ventral side ensured that the RNA-later solution would diffuse into all tissues. Conserved specimen were later dissected in the lab and gill tissue was separated, since gills represent a main osmoregulatory organ in fishes. For the acclimation experiment, we collected live adult fish from Kiel (20 PSU), which were crossed in our facilities at GEOMAR to obtain ten F1 laboratory bred families, herein referred to as 'parental generation'. At nine months post-hatch we split each family into three salinity treatment groups of 10 fish each: one at 33 PSU, one at 6 PSU, and one control group at 20 PSU. The salinity transition was performed within 10 days by 3 PSU steps every second day. Over the entire time each group

was fed ad libitum and kept in a 20-L aquarium connected to one of three filter tanks per

salinity treatment. After 5 months under treatment conditions, six pure crosses per salinity

treatment group were performed in vitro, herein referred to as 'offspring generation' (F2).

Offspring and parental generation were kept at 18 °C water temperature and a 15:9 light/dark

cycle (L/D). The last eight weeks before performing the F2 crosses, the F1 generation

underwent an artificial winter to trigger reproduction (2 weeks at 12 °C, 12:12 L/D; 4 weeks 6

°C, 8:16 L/D, 2 weeks at 12 °C, 12:12 L/D). Upon fertilization, clutches were split and

separated into different treatments (Figure 1). At three months post-hatch, laboratory bred F2

sticklebacks were euthanized using MS222, photographed, dissected and their gill tissue

was stored in RNA-later. The age at sampling matched the estimated age of the wild caught

juveniles (3 months). Additionally, to the 48 wild caught individuals from Kiel, Nynäshamn

and Sylt that were used in above field survey, we sequenced whole genomes from gill tissue

of an additional three populations of sticklebacks, namely from Falsterbo, Sweden

(55°24'46.6"N 12°55'52.3"E; 10 PSU; N = 16), Letipea (59°33'07.6"N 26°36'29.7"E; 4 PSU; N

= 16) and Barsta (62°51'47.1"N 18°23'51.0"E; 5 PSU; N = 16).

Mortality and hepatosomatic index

Mortality was monitored throughout the experiment to account for possible non-random

mortality. Three months post-hatch, we assessed the standard length (SDL), total weight and

liver weight of the experimental F2 generation and calculated the hepatosomatic index (HSI =

liver weight / total weight * 100), which is a proxy for energy reserves in form of glycogen

storage in the liver. We analyzed the effect of treatment (5 treatment groups, Figure 1) on the

survival rate per family as a ratio of 'alive' vs 'dead' fish, using glmer implemented in the R

package 'lme4' (40) with Binomial error and 'crossing' as well as 'climate chamber' as

random effects. The effect of treatment on HSI, SDL and total weight was analyzed fitting

three individual linear mixed effect models, using lmer in 'lme4' (40) with Gaussian error and

'crossing' as well as 'tank' nested within 'climate chamber' as random effects. Tukey HSD

post hoc tests were run using the glht function implemented in the package 'multicomp' (41)

to identify significant differences between treatment groups.

DNA extraction

For the field survey, DNA extraction of gill tissue (N = 16 individuals per population) was

performed using the DNeasy Blood & Tissue Kit (Qiagen). Further purification of the

extracted DNA was done with NucleoSpin® gDNA Clean-up (Macherey-Nagel). For

laboratory bred F2 offspring of the two-generation acclimation experiment, dual extraction of

whole RNA and DNA was performed from gill tissue (N = 11-12 individuals per treatment

group, Figure 1) stored in RNAlater using the AllPrep DNA/RNA mini kit (Qiagen). Purity and

quality of the extracted DNA was estimated using a NanoDrop ND-1000 spectrophotometer

(Thermo Fisher Scientific) and a standard agarose gel (1% Agarose/TAE). DNA

concentration was assessed using the Qubit® 2.0 Fluorometer (Thermo Fisher Scientific). To

obtain a balanced sex ratio (50/50), we determined the gender of the individuals using a sex-

specific genetic polymorphism in isocitrate dehydrogenase (IDH) with a modified protocol

from Peichel et al. (2004) (42). For the PCR (settings: once 94°C for 3 minutes; 30 cycles of

94°C for 30 sec, 54°C for 20 sec, 72°C for 30 sec; once 72°C for 5 minutes), 1 µL forward

and reverse primer (5µM) was used with 4.9 µL water, 1 µL 10x buffer, 1 µL dNTPs (0.5 µM),

and 0.1 µL DreamTaq (5 U/µL). The resulting PCR products were visualized with a capillary

electrophoresis on the 3100 ABI sequencer and a LIZ500 size standard. While males show a

heterogametic signal with two bands (at approximately 300 bp and 270 bp), females lack the

band at 270 bp.

Library preparation and sequencing (Whole genome sequencing, WGS)

For whole genome sequencing, the 'TruSeq Nano DNA' (Illumina) library preparation kit was

used according to the manufacturer's protocol by the Sequencing Facility of the IKMB,

University of Kiel. Ultrasonication was conducted with a 'Covaris E220' (Covaris) to shear the

input DNA (100 ng per sample and 350 bp insert size). Before the enrichment with a PCR

step (8 cycles), fragmented and bead purified DNA was ligated with adenylate at the blunt 3'

ends (End repair and A-tailing) and with indexing adapters. Fragments were cleaned with

MagSi-NGS Prep Plus Beads (Steinbrenner). Paired-end sequencing of the quality-controlled

and multiplexed libraries was performed on the Illumina Hiseq 4000 platform (2 x 150 bp

reads).

Quality assessment, data filtering and mapping (WGS)

The command line tools of Picard v.2.7.1 (Broad Institute 2016) were used to (i) reformat the

Fastq to uBAM file format and to add further values (read group etc.) to the SAM header

using FastqToSam), to (ii) mark the location of adapter sequences using

MarkIlluminaAdapters, and to (iii) reconvert the sequences to Fastq format with SamToFastq.

The stickleback genome (here Broad/gasAcu1) was indexed with bwa index and used as a

reference for the mapping with bwa mem (43) v.07.12-r1044. To retain the meta-information

from the uBAMs we used MergeBamAlignment. Picard was also used to identify duplicates

with MarkDuplicates. Basic statistics were generated with CollectWgsMetrics,

CollectInsertSizeMetrics and AlignmentSummaryMetrics and summarized with MultiQC

v1.0.dev0 (44). A total number of 4,463,070,154 high-quality reads (mapping quality > Q20)

was mapped resulting in a mean depth of 13.84x (sd. 2.02x) and mean insert size 383.07 bp

(sd. 9.40 bp, Supplementary Table S3). GATK v 3.7 HaplotypeCaller (45) was run to

determine the likelihoods of the haplotypes per sample, i.e. to call SNPs and indels, which

were than processed with GenotypeGVCFs for a joint genotyping. SNPs were selected using

hard filters for quality and extracted from the raw genotypes with a combination of the

SelectVariants, VariantsToTable and VariantFiltration commands. VCFtools (46) was used in

a next step, removing SNPs with a minimum quality score (minQ) below 20 and a minor

allele frequency (maf) greater than or equal 0.0049.

Library preparation and sequencing (reduced representation bisulfite sequencing,

RRBS)

The library preparation for methylation analyses followed the Smallwood and Kelsey reduced

representation bisulfite sequencing (RRBS) protocol (47). A total of 100-250 ng purified DNA

was digested with the methylation-insensitive MspI restriction enzyme, which cuts at the

"CCGG" motif and thereby enriches for CpG regions. DNA end-repair and A-tailing was

conducted and un-tailed CEGX spike-in controls (Cambridge Epigenetix) were added. These

are DNA oligos of known sequence and with known cytosine modification, which can be used

for downstream assessment of bisulfite conversion efficiency. After adapter ligation, bisulfite

conversion was conducted using the EZ-96 DNA Methylation-Gold Kit (Zymo Research)

according to the manufacturer's protocol. PCR amplification with 19 cycles were performed.

Quality control of purified PCR products was performed on a 2200 TapeStation System

(Agilent) and high-quality libraries were pooled and diversified with 15% PhiX. Single-end

sequencing with 100 bp read length was conducted on a HiSeq 2500 sequencer (Illumina).

Quality assessment, data filtering and mapping (RRBS)

In total 106 individuals (48 wild caught and 58 experimental fish) of balanced sex ratio were

DNA sequenced at an average of 19.8 ± 3.5 million reads for experimental fish and 11.4 ±

2.1 million reads for wild-caught fish (Supplementary Table S4). De-multiplexed fastq files

were quality checked using FastQC v0.11.5 (48) and Multiqc v1.3 (44). Adapters were

removed with cutadapt v1.9.1 (49) using multiple adapter sequences

(NNAGATCGGAAGAGCACAC, AGATCGGAAGAGCACAC, ATCGGAAGAGCACAC) with a

minimum overlap of one base pair between adapter and read. This was necessary to remove

primer dimers and avoid false methylation calls systematically caused by the RRBS end-

repair step during library preparation, if the end repair step adds artificial cytosins.

Simultaneously, cutadapt was used to trim low quality bases (-q 20) from the 3'-end and

remove trimmed reads shorter than 10 bases. An air bubble during sequencing caused the

bases 66-72 of ten tiles of one lane (affecting 12 individuals) to have low quality values,

which were removed in a custom awk script. Two poor quality individuals (a Sylt and a

Nynäshamn female) did not meet our strict quality requirements (e.g.: ≥ 5 million reads,

mapping efficiency > 52%) and showed biases in the proportion of bases per position

compared to other individuals (plot in fastqc "per base sequence content"). Therefore, we

excluded these two libraries from downstream analysis resulting in 15 instead of 16

individuals from Sylt and Nynäshamn (Figure 1). Bisulfite conversion efficiency was

assessed from the spike-in controls (Cambridge Epigenetix), using the cegxQC software

(50). Overall conversion levels were 2.4 ± 1.8% conversion of methylated cytosines and 99.6

± 0.5% conversion of un-methylated cytosines, which is in line with expected conversion

rates (Supplementary Table S4). We used Bismark v0.17.0 (51) to index the UCSC

stickleback reference genome (Broad/gasAcu1) and to generate the bisulfite alignments with

Bowtie2 v2.3.3 at default settings. Bismark was also used to extract the methylation calls.

Average mapping efficiency was 63.7 ± 2.4% (Supplementary Table S4).

Identification of differentially methylated sites

Identification of differentially methylated sites

The methylation calls were analyzed in R v3.4.1 (52) using the package methylKit v1.3.8

(53). CpG loci were filtered for a minimum coverage of 10 reads per site. To account for

potential PCR bias, we additionally excluded all sites in the 99.9th percentile of coverage. To

improve the methylation estimates, we corrected for SNPs, which could have led to a wrong

methylation call. The excluded positions were derived with custom written perl scripts from C-

to-T and G-to-A-SNPs with genotype quality of 20 and a minimum allele frequency of 0.005

(see above) from the 96 wild caught individuals with a combination of custom written Perl

and R-scripts using packages from methylkit (53) and GenomicRanges (54) (Supplementary

File Summary_scripts.txt). After normalizing coverage values between samples, using

normalizeCoverage implemented in methylKit, we excluded all sites that were present in

fewer than 9 individuals per treatment group from downstream analysis. As previously

shown, sex specific methylation affects < 0.1% of CpG sites on autosomal chromosomes, but

> 5% of CpGs on the sex chromosome (19). Therefore, to exclude a potential sex bias, we

removed all CpG sites located on the sex chromosomes (chromosome 19), resulting in a

high-quality dataset with 525,985 CpG sites. Finally, by checking the first six principal

components of the resulting PCA and running an ANOVA on the filtered dataset, we

confirmed the absence of an effect of sex on global methylation pattern ($F_{124,1}$ = 2.611, P =

0.109). However, the PCA revealed a bias in methylation pattern by families over all

experimental groups. Therefore, to identify differentially methylated CpG sites (DMS)

between treatment groups, we performed pairwise comparisons (Supplementary Table S5)

fitting a logistic regression model per CpG site with calculateDiffMeth in methylKit using

family as covariate for the experimental groups. A Chi-square test was applied to assess

significance levels of DMS and P-values were corrected to q-values for multiple testing using

the SLIM (sliding linear model) method (55). Additionally, we accounted for multiple use of

groups in pairwise comparisons and adjusted the alpha for the q-value according to

Bonferroni correction to 0.0125 (= 0.05 / 4). Ultimately, CpG sites were considered to be

differentially methylated with a q-value < 0.0125 and a minimum weighted mean methylation

difference of 15%. To ensure that the DMS obtained are not laboratory artefacts, we used

calculateDiffMeth implemented in methylKit and compared the wild population from Kiel to

the experimental control group (Kiel population from 20 PSU at 20 PSU). The resulting

11,828 DMS were excluded from the DMS obtained by the pairwise comparisons mentioned

above (Supplementary Table S5). DMS were plotted across the genome for the comparison

between Kiel vs Nynäshamn (20 vs. 6 PSU, blue fish) and Kiel vs Sylt (20 vs. 33 PSU, yellow

fish) using ggplot2 (56) und hypoimg (57) (Supplementary Figure S1).

541   Assessment of inducibility and gene association of DMS

By comparing wild caught individuals from the mid salinity population (20 PSU, KIE) to the

populations sampled at low (6 PSU, NYN) and high (33 PSU, SYL) salinity in the field, we

obtained 1,470 (KIE-NYN) and 1,158 (KIE-SYL) pairwise pop-DMS. We first tested whether

these pop-DMS distinguishing natural populations are inducible or stable at the respective

salinity in the experiment. A pop-DMS was considered stable when the within- and the

transgenerational acclimation group did not significantly differ in methylation to the control

group (q-value ≥ 0.0125). On the other hand, pop-DMS were considered inducible when at

least one of the acclimation groups was differentially methylated compared to the control

group (q-value < 0.0125; methylation difference ≥ 15%). Pop-DMS with a significant q-value

not exceeding the threshold of differential DNA methylation (15%) will be referred to as

'inconclusive' hereafter. We used a randomization test to ensure that the number of inducible

sites obtained did not occur by chance. To this end, we randomly sampled 1,470 (KIE-NYN)

and 1,158 (KIE-SYL) pop-DMS from the complete dataset (1,000 replicates). A Chi-square

test was used to assess whether our observed number of inducible, stable and inconclusive

sites differs from a random set of sites (averaged over replicates). Finally, we tested whether

the weighted mean methylation difference (meth.diff, in percentage) between wild

populations matches the inducible methylation difference by subtracting the 'meth.diff' in the

experiment (exp) from the 'meth.diff' between wild caught populations (wild):

$\delta.meth.diff = 100 - (meth.diff.wild - meth.diff.exp)$

As we subtracted this difference from 100, values closer to 100 indicated higher similarity of

experimentally inducible methylation with the postulated adaptive DNA methylation pattern in

natural populations. By comparing the 'δ.meth.diff' for within- and transgenerational

acclimation using an ANOVA, we can assess whether there is a difference in inducibility of

methylation to match patterns found in wild caught populations. All analyses were run

separately for decreased (6 PSU; KIE-NYN) and increased (33 PSU; KIE-SYL) salinity.

In order to detect potential functional associations of the observed changes in DNA

methylation state, we classified the genomic region of a pop-DMS based on their nearest

transcription start site (TSS) using annotateWithGeneParts and getAssociationWithTSS

implemented in genomation v1.4.2 (58). We distinguished between promoter (1500 bp

upstream and 500 bp downstream of TSS), exon, intron and intergenic regions. To be

associated to a gene, the pop-DMS had to be either inside the gene or, if intergenic, not

further than 10 kb away from the TSS. We excluded three pop-DMS that were on a different

reference scaffold then the gene they were associated to on the chrUn linkage group (that

merges scaffolds into one large artificial chromosome). Using the genes with associated pop-

DMS, we applied a conditional hypergeometric GO term enrichment analysis (FDR corrected

P-value ≤ 0.05) with the ensembl stickleback annotation dataset 'gaculeatus_gene_ensembl'

and all genes that were associated to any sequenced CpG site as universe. We identified

overrepresented biological processes, molecular functions and cellular components using the

packages GOstats v2.5 (59) and GSEABase v1.46 (60) and corrected for multiple testing

using the FDR method implemented in goEnrichment v1.0 (61) in R v3.6 (52). Figures were

produced using ggplot2 v3.2 (56).

Estimation of DNA sequence based genetic differentiation at differentially methylated

sites

In order to evaluate the genetic differentiation up- and downstream (in sum 10 kb) of the pop-

DMS position, we calculated the mean FST values (≤ 60% missing data and depth ≥ 5) from

whole genome sequencing data of the exact same individuals with vcftools v.0.1.15 (62). We

hypothesized that inducible CpG positions matching the methylation difference expected

from the profile of the wild populations are genetically more similar between the populations

than sites that changed in the opposite direction. To test this one-tailed hypothesis we

applied a randomization test (with 10,000 bootstraps) on the mean FST difference between

the two groups (expected and opposite):

δ.mean.FST = 'expected' mean FST – 'opposite' mean FST

We plotted the 10,000 delta mean FST values and calculated a P-value by dividing the

proportion of values smaller than the true difference by the number of bootstraps. Figures

were produced using ggplot2 v3.2 (56).

Acknowledgments

Many thanks to Fabian Wendt, Jakob Gismann, Linda Sartoris, Christoph Giez, Zuzanna

Zagrodzka, Laura Niewendick, Max Bettendorff, Bastian Poerschke, and Tobias Strickmann

for taking care of the experimental animals and assisting the sampling procedures. Thanks to

Florian Rüppel for planning and constructing the recirculating aquaria systems. We thank the

many field helpers for catching wild sticklebacks. Many thanks also to Susanne Landis who

visualized our ideas in Figure 1 and 5. We kindly acknowledge the financial support for the

BAMBI project (Grant Agreement number: call 2012-76) by BONUS, the joint Baltic Sea

research program, funded by the European Union and the Federal Ministry of Education and

Research in Germany to TBHR and CE (reference number 03F0680A). Further, BSM's

research was generously supported by the Excellence Cluster 'The Future Ocean' (EXC 80)

and the sequencing partly by the Excellence Cluster 'Inflammation at Interfaces' (EXC 306)

and 'Origins and Function of Metaorganisms' (CRC1182), subprojects Z3 & INF, by the

Deutsche Forschungsgemeinschaft (DFG). Both 'The Future Ocean' and 'Inflammation at

Interfaces' are funded within the framework of the Excellence Initiative by the Deutsche

Forschungsgemeinschaft (DFG) on behalf of the German federal and state governments.

Author contributions

CE and TBHR conceived and designed the study with contributions for the bisulfite

sequencing strategy from BSM. BSM and MJH planned and carried out the fieldwork at the

German locations, BSM supervised the sampling in Estonia and Sweden, which was carried

out by a great team of the BONUS-BAMBI project. MJH, with the help from CE and TBHR,

planned and supervised the breeding and acclimatization experiment. MJH and BSM

conducted wet laboratory work (DNA extractions and quality assessment). MPH and RH and

conducted the library preparation and sequencing. MJH and BSM analyzed the data and

drafted the manuscript together with equal contributions. All co-authors discussed and

interpreted the results and contributed to the final version of the manuscript.

Competing interests

The authors declare no competing interests.

Data Availability

Fastq raw reads of genomes and methylation sequencing will be deposited in GenBank. Raw
data on mortality and phenotypic effects are available at PANGAEA
(https://doi.org/10.1594/pangaea.892493).

Figures and Tables

Fig. 1: Experimental space-for-time approach. We characterized DNA methylation profiles (via
Reduced Representation Bisulfite Sequencing, RRBS) and whole genomes (Whole Genome
Sequencing, WGS) of fish from three populations of wild caught three-spined sticklebacks locally
adapted to 6 (blue, N = 15), 20 (green, N = 16) and 33 PSU (yellow, N = 15). We also collected
sticklebacks from the mid salinity location (20 PSU) and acclimated laboratory bred offspring of these
fish within one ('within-generational') or over two ('transgenerational') generations to decreased (6
PSU) or increased (33 PSU) salinity, and maintained a control group at its original salinity (N = 11-12
per group, see details in Figure). Differential methylation within and across generations was assessed

and compared to natural populations locally adapted to the corresponding salinity, serving as the

hypothetical future DNA methylation state to capture long term adaptation processes.

Fig. 2: Gene ontology terms for biological processes and molecular functions.

Gene ontology (GO) terms for biological processes and molecular functions under salinity increase

(yellow, 20 vs. 33 PSU) and decrease (blue, 20 vs. 6 PSU) associated with differentially methylated

sites between populations (pop-DMS) are presented. The graph is split into GO terms associated with

pop-DMS from natural stickleback populations across a salinity cline (wild) and their experimental

inducibility (inducible and stable) in a two-generation acclimation experiment. The size of the circles

 refers to the number of genes of this term present in our groups (in %) and the transparency to the

 FDR corrected P-value (darker circles refer to a lower adjusted P-value). This subset is filtered for GO

 terms including the following keywords: "channel", "transport", "water", "chloride", "potassium",

 "homeostasis", "ion-dependent", "urine", "ATP", "metabolic", see Figure S2 (Supplementary Material)

 for the full figure.

 Fig. 3: The duration of acclimation (within- vs. transgenerational) and level of genomic

 differentiation between the populations influence DNA methylation at inducible sites. Figure 3A

 and 3B show mean FST values for pop-DMS (with a ± 5 kb window) inducible under experimental

 salinity decrease (top, blue) and increase (bottom, yellow), that either shifted methylation levels

 towards the values observed in the field (expected) or the opposite direction (opposite). A

 randomization test (with 10,000 bootstraps) was performed for the difference between expected and

 opposite mean FST value (δ.mean.FST = 'expected' mean FST – 'opposite' mean FST; Figure 3C and

 3D). Under the one tailed hypothesis of increased genetic differentiation at opposite sites and an alpha

 of 0.05 the P-value was calculated as values smaller than the true difference divided by 10,000

 bootstraps. In Figure 3E-H the y-axis shows the percentage match between the within- and

transgenerational acclimation group in relation to the methylation differentiation level found in natural

populations at inducible pop-DMS. This value was obtained by calculating the difference between the

methylation change in the experiment (meth.diff.exp in %; control vs. within-generational or control vs.

transgenerational) and the difference in methylation between natural populations (meth.diff.wild in %)

as δ.meth.diff = 100 − (meth.diff.wild − meth.diff.exp). Mean values ± 95% confidence interval are

shown for within- and transgenerational acclimation to decreased and increased salinity at expected

and opposite inducible sites. Colors refer to the direction of DNA methylation change (hypomethylation

or hypermethylation). Values closer to 100 indicate a shift in methylation pattern towards adaptive

methylation levels found in natural populations and stars indicate the significance level ($P \leq 0.001$ '***';

$P \leq 0.01$ '**') for the comparison between within- and transgenerational acclimation. 'Main effect' refers

to an effect of acclimation (within- or transgenerational) and 'interaction effect' to an interaction of

acclimation and methylation direction (hypo- or hypermethylation).

Fig. 4: Effects of salinity acclimation on fitness correlated factors. For all five acclimation groups

(control group (20 PSU), within- and transgenerational acclimation to 6 or 33 PSU) survival rates in

percent (A), standard length in cm (B), hepatosomatic index (C) and total weight in g (D) are

694   displayed. Letters indicate significant differences resulting from Tukey HSD post hoc tests

695   (Supplementary Material: Table S3).

697

Fig. 5: Graphical summary of experimental design and main results. We used the Baltic Sea

salinity gradient to study the role of DNA methylation in local salinity adaptation and the response to

salinity change in a space-for-time approach. To assess the potential future acclimatization and

adaptation processes of the natural stickleback population from 20 PSU (Kiel / green) to the predicted

desalination(63), we compared differences in DNA methylation at CpG sites between wild caught and

laboratory bred sticklebacks. Following the experiment timeline (bottom), we compared methylation

levels of the experimental control group from 20 PSU, to within- and transgenerational acclimation of

20 PSU sticklebacks to 6 PSU (DNA from left to right). The population locally adapted to 6 PSU serves

as the hypothetical future state in which salinities will decrease (blue, DNA on the right). The three

main results are written in the circles with schematically and horizontally corresponding DNA

methylation changes. (i) 63% of the DMS between the populations remained stable under

experimental salinity change. (ii) The direction of experimental methylation change was dependent not

only on the treatment but also on the degree of genetic differentiation between the populations (see

Figure 3A-D for results). (iii) Transgenerational salinity acclimation shifted DNA methylation patterns

closer to the anticipated adaptive state found in the hypothetical future population (see Figure 3E-H for

results). For clarity, only one (6 PSU) of the two foreign salinity regimes tested (6 and 33 PSU) is

shown, indicated by the yellow fish on the top left (see Figure 1 for full experimental design).

Ensembl gene ID	chromosome	start position	end position	gene name	wild	inducible	'expected' inducible	'opposite' inducible	stable	inconclusive	Fisher´s exact (*P*)
Salinity decrease:											
ENSGACG00000008328	Chr10	12860144	12863850	si:dkey-166k12.1	24	0	0	0	9	15	0.005
ENSGACG00000019416	**Chr7**	**4451892**	**4453656**	**HMX1 orthologue**	**17**	**0**	**0**	**0**	**9**	**8**	**0.033**
ENSGACG00000013229	**Chr18**	**15327717**	**15352321**		**15**	**0**	**0**	**0**	**3**	**12**	**0.011**
ENSGACG00000017287	Chr3	13454527	13465167	mmp16b	12	0	0	0	12	0	0.001
ENSGACG00000017584	Chr3	14690814	14694448	CCNY	12	12	12	0	0	0	0.001
ENSGACG00000018249	Chr4	12141625	12143011	si:ch211-153b23.5	12	1	1	0	3	8	0.188
ENSGACG00000008034	Chr6	9368187	9380941		11	10	10	0	0	1	0.014
ENSGACG00000009469	Chr1	9166576	9173856	egln2	11	0	0	0	11	0	0.001
ENSGACG00000004433	Chr17	2127457	2211376	igsf21a	10	10	10	0	0	0	0.003
ENSGACG00000007343	Chr10	10666995	10679875	col9a2	10	0	0	0	6	4	0.227
ENSGACG00000018407	Chr4	13828336	13837518	Sncb	10	2	2	0	5	3	0.848
Salinity increase:											
ENSGACG00000020323	Chr7	17010160	17011176		23	0	0	0	22	1	<0.001
ENSGACG00000013229	**Chr18**	**15327717**	**15352321**		**15**	**10**	**10**	**0**	**1**	**4**	**0.125**
ENSGACG00000013359	Chr11	12960883	12968110	sec14l1	15	0	0	0	12	3	0.011
ENSGACG00000019416	**Chr7**	**4451892**	**4453656**	**HMX1 orthologue**	**15**	**3**	**3**	**0**	**5**	**7**	**0.745**
ENSGACG00000002948	Chr8	218240	221355	ddx10	14	0	0	0	6	8	0.077
ENSGACG00000016350	Chr14	3603545	3604923		14	1	0	1	7	6	0.277
ENSGACG00000006636	Chr18	4780893	4786820	ZC3H12D	13	0	0	0	3	10	0.034
ENSGACG00000004667	Chr12	4273498	4286193	tti1	12	0	0	0	12	0	0.001
ENSGACG00000015566	Chr2	9043062	9051779	casc4	10	0	0	0	10	0	0.003

Table 1: Differentially methylated genes across natural populations along a salinity cline

Genes derived from DNA methylation comparisons between natural populations associated with ≥ 10 pop-DMS (decreased salinity: KIE (20 PSU) vs. NYN (6

PSU); increased salinity: KIE (20 PSU) vs. SYL (33 PSU)). Ensembl gene ID and name as well as the position on the chromosome are listed. The numbers refer

to the numbers of DMS in the population comparison (wild). These DMS were classified into 'inducible', 'inconclusive' and 'stable' sites according to their behavior

in a two-generation salinity acclimation experiment with laboratory bred sticklebacks from the mid salinity population (20 PSU) exposed to experimental salinity

increase or decrease (33 and 6 PSU respectively). Further, inducible sites were distinguished whether they matched methylation levels of the locally adapted

population ('expected') or not ('opposite'). Genes written in bold vary in both population comparisons. We used a Fisher's exact test to assess whether pop-DMS

associated to the same gene are correlated in their response to experimental salinity change (non-random distribution among the categories stable, inducible,

inconclusive) and reported corresponding P-values. For a full table on all genes associated with 1 or more pop-DMS see Table S2A and S2B (Supplementary

Material).

References

1. E. Jablonka, The evolutionary implications of epigenetic inheritance. *Interface focus* 7, 20160135 (2017).

2. K. N. Laland *et al.*, Does evolutionary theory need a rethink? *Nature News* 514, 161 (2014).

3. M. I. Lind, F. Spagopoulou, Evolutionary consequences of epigenetic inheritance. *Heredity* 121, 205-209 (2018).

4. O. Bossdorf, C. L. Richards, M. Pigliucci, Epigenetics for ecologists. *Ecology Letters* 11, 106-115 (2008).

5. F. D. Klironomos, J. Berg, S. Collins, How epigenetic mutations can affect genetic evolution: model and mechanism. *Bioessays* 35, 571-578 (2013).

6. P. A. Jones, Functions of DNA methylation: islands, start sites, gene bodies and beyond. *Nature Reviews Genetics* 13, 484 (2012).

7. N. Shea, I. Pen, T. Uller, Three epigenetic information channels and their different roles in evolution. *Journal of Evolutionary Biology* 24, 1178-1187 (2011).

8. I. Kronholm, S. Collins, Epigenetic mutations can both help and hinder adaptive evolution. *Molecular Ecology* 25, 1856-1868 (2016).

9. R. J. Schmitz *et al.*, Transgenerational epigenetic instability is a source of novel methylation variants. *Science* 334, 369-373 (2011).

10. F. Johannes *et al.*, Assessing the impact of transgenerational epigenetic variation on complex traits. *PLoS Genetics* 5, e1000530 (2009).

11. T. Ryu, H. D. Veilleux, J. M. Donelson, P. L. Munday, T. Ravasi, The epigenetic landscape of transgenerational acclimation to ocean warming. *Nature Climate Change* 8, 504-509 (2018).

12. K. N. Laland *et al.*, The extended evolutionary synthesis: its structure, assumptions and predictions. *Proceedings of the Royal Society B: Biological Sciences* 282, 20151019 (2015).

13.   D. H. Evans, Cell signaling and ion transport across the fish gill epithelium. *Journal of*

Experimental Zoology 293, 336-347 (2002).

14.   T. B. H. Reusch *et al.*, The Baltic Sea as a time machine for the future coastal ocean.

Science Advances 4, eaar8195 (2018).

15.   B. Guo, J. DeFaveri, G. Sotelo, A. Nair, J. Merilä, Population genomic evidence for

adaptive differentiation in Baltic Sea three-spined sticklebacks. *BMC Biology* 13, 19

(2015).

16.   M. J. Heckwolf, B. S. Meyer, T. Döring, C. Eizaguirre, T. B. H. Reusch, Transgenerational

plasticity and selection shape the adaptive potential of sticklebacks to salinity change.

Evolutionary Applications 11, 1873-1885 (2018).

17.   J. DeFaveri, J. Merilä, Local adaptation to salinity in the three-spined stickleback? *Journal*

of Evolutionary Biology 27, 290-302 (2014).

18.   L. N. S. Shama *et al.*, Transgenerational effects persist down the maternal line in marine

sticklebacks: gene expression matches physiology in a warming ocean. *Evolutionary*

applications 9, 1096-1111 (2016).

19.   D. C. H. Metzger, P. M. Schulte, The DNA Methylation Landscape of Stickleback Reveals

Patterns of Sex Chromosome Evolution and Effects of Environmental Salinity. *Genome*

Biol Evol 10, 775-785 (2018).

20.   A. V. Artemov *et al.*, Genome-wide DNA methylation profiling reveals epigenetic

adaptation of stickleback to marine and freshwater conditions. *Molecular biology and*

evolution 34, 2203-2213 (2017).

21.   A. Bird, Perceptions of epigenetics. *Nature* 447, 396 (2007).

22.   A. Taudt, M. Colomé-Tatché, F. Johannes, Genetic sources of population epigenomic

variation. *Nature Reviews Genetics* 17, 319 (2016).

23.   A. Van Der Graaf *et al.*, Rate, spectrum, and evolutionary dynamics of spontaneous

epimutations. *Proceedings of the National Academy of Sciences* 112, 6676-6681 (2015).

24.   H. B. Fraser, Gene expression drives local adaptation in humans. *Genome Research* 23,

1089-1096 (2013).

25. T. L. Lenz, C. Eizaguirre, B. Rotter, M. Kalbe, M. Milinski, Exploring local immunological

adaptation of two stickleback ecotypes by experimental infection and transcriptome-wide

digital gene expression analysis. *Molecular Ecology* 22, 774-786 (2013).

26. P. F. Colosimo *et al.*, Widespread parallel evolution in sticklebacks by repeated fixation of

Ectodysplasin alleles. *Science* 307, 1928-1933 (2005).

27. R. Spence *et al.*, Calcium and salinity as selective factors in plate morph evolution of the

three-spined stickleback (*Gasterosteus aculeatus*). *Journal of Evolutionary Biology* 25,

1965-1974 (2012).

28. A. Paccard *et al.*, Adaptation in temporally variable environments: stickleback armor in

periodically breaching bar-built estuaries. *Journal of evolutionary biology* 31, 735-752

(2018).

29. A. L. Ferchaud *et al.*, A low-density SNP array for analyzing differential selection in

freshwater and marine populations of threespine stickleback (*Gasterosteus aculeatus*).

BMC Genomics 15, 867 (2014).

30. P. A. Hohenlohe *et al.*, Population genomics of parallel adaptation in threespine

stickleback using sequenced RAD tags. *PLoS Genetics* 6, e1000862 (2010).

31. N. V. Terekhanova *et al.*, Fast evolution from precast bricks: genomics of young

freshwater populations of threespine stickleback *Gasterosteus aculeatus. PLoS Genetics*

10, e1004696 (2014).

32. J. DeFaveri, P. R. Jonsson, J. Merilä, Heterogeneous genomic differentiation in marine

threespine sticklebacks: adaptation along an environmental gradient. *Evolution* 67, 2530-

2546 (2013).

33. T. Schaarschmidt, E. Meyer, K. Jürss, A comparison of transport-related gill enzyme

activities and tissue-specific free amino acid concentrations of Baltic Sea (brackish water)

and freshwater threespine sticklebacks, Gasterosteus aculeatus, after salinity and

temperature acclimation. *Marine Biology* 135, 689-697 (1999).

34. M. J. Dubin *et al.*, DNA methylation in Arabidopsis has a genetic basis and shows

evidence of local adaptation. *elife* 4, e05255 (2015).

35.   G. Torda *et al.*, Rapid adaptive responses to climate change in corals. *Nature Climate*

Change 7, 627–636 (2017).

36.   M. E. Potok, D. A. Nix, T. J. Parnell, B. R. Cairns, Reprogramming the maternal zebrafish

genome after fertilization to match the paternal methylation pattern. *Cell* 153, 759-772

(2013).

37.   C. Labbé, V. Robles, M. P. Herraez, Epigenetics in fish gametes and early embryo.

Aquaculture 472, 93-106 (2017).

38.   J. Gertz *et al.*, Analysis of DNA methylation in a three-generation family reveals

widespread genetic influence on epigenetic regulation. *PLoS Genetics* 7, e1002228

(2011).

39.   J. R. Gibbs *et al.*, Abundant quantitative trait loci exist for DNA methylation and gene

expression in human brain. *PLoS Genetics* 6, e1000952 (2010).

40.   D. Bates, M. Maechler, B. Bolker, S. Walker, lme4: Linear mixed-effects models using

Eigen and S4. *R package version* 1, 1-23 (2014).

41.   T. Hothorn, F. Bretz, P. Westfall, Simultaneous Inference in General Parametric Models.

Biometrical Journal 50, 346-363 (2008).

42.   C. L. Peichel *et al.*, The master sex-determination locus in threespine sticklebacks is on a

nascent Y chromosome. *Current biology* 14, 1416-1424 (2004).

43.   H. Li, R. Durbin, Fast and accurate short read alignment with Burrows–Wheeler

transform. *Bioinformatics* 25, 1754-1760 (2009).

44.   P. Ewels, M. Magnusson, S. Lundin, M. Käller, MultiQC: summarize analysis results for

multiple tools and samples in a single report. *Bioinformatics* 32, 3047-3048 (2016).

45.   A. McKenna *et al.*, The Genome Analysis Toolkit: a MapReduce framework for analyzing

next-generation DNA sequencing data. *Genome Research*, (2010).

46.   P. Danecek *et al.*, The variant call format and VCFtools. *Bioinformatics* 27, 2156-2158

(2011).

47.   S. A. Smallwood, G. Kelsey, in *Genomic Imprinting: Methods and Protocols,* N. Engel,

Ed. (Humana Press, Totowa, NJ, 2012), pp. 187-197.

48.   S. Andrews, FastQC: a quality control tool for high throughput sequence data.

842   (http://www.bioinformatics.babraham.ac.uk/projects/fastqc). (2010).

843   49.   M. Martin, Cutadapt removes adapter sequences from high-throughput sequencing

844   reads. *EMBnet. Journal* 17, 10-12 (2011).

845   50.   CEGX Bioinfomatics Team, Cambridge Epigenetix (CEGX), Babraham Research

846   Campus, Cambridge. (2015).

847   51.   F. Krueger, S. R. Andrews, Bismark: a flexible aligner and methylation caller for Bisulfite-

848   Seq applications. *Bioinformatics* 27, 1571-1572 (2011).

849   52.   R Core Team, R: A language and environment for statistical computing. *R Foundation for*

850   *Statistical Computing, Vienna, Austria.*, (2017).

851   53.   A. Akalin *et al.*, methylKit: a comprehensive R package for the analysis of genome-wide

852   DNA methylation profiles. *Genome Biology* 13, (2012).

853   54.   M. Lawrence *et al.*, Software for computing and annotating genomic ranges. *PLoS*

854   *computational biology* 9, e1003118 (2013).

855   55.   H. Q. Wang, L. K. Tuominen, C. J. Tsai, SLIM: a sliding linear model for estimating the

856   proportion of true null hypotheses in datasets with dependence structures. *Bioinformatics*

857   27, 225-231 (2011).

858   56.   H. Wickham, ggplot2: Elegant Graphics for Data Analysis. *Springer-Verlag New York*,

859   (2016).

860   57.   K. Hench, https://github.com/k-hench/hypoimg. (2019).

861   58.   A. Akalin, V. Franke, K. Vlahoviček, C. E. Mason, D. Schübeler, Genomation: a toolkit to

862   summarize, annotate and visualize genomic intervals. *Bioinformatics* 31, 1127-1129

863   (2014).

864   59.   S. Falcon, R. Gentleman, Using GOstats to test gene lists for GO term association.

865   *Bioinformatics* 23, 257-258 (2006).

866   60.   M. Morgan, S. Falcon, R. Gentleman, GSEABase: Gene set enrichment data structures

867   and methods. (2019).

868   61.   A. Hallab, goEnrichment: Helper functions to compute GO enrichment tests using

869   GOstats and GSEABase. (2015).

870   62.   B. S. Weir, C. C. Cockerham, Estimating F-statistics for the analysis of population

871   structure. *Evolution* 38, 1358-1370 (1984).

872   63.   H. E. M. Meier, Baltic Sea climate in the late twenty-first century: a dynamical

873   downscaling approach using two global models and two emission scenarios. *Clim Dynam*

874   27, 39-68 (2006).

875