Transcranial focused ultrasound enhances behavioral and network mechanisms underlying response inhibition in humans

Justin M. Fine, Maria E. Fini, Archana S. Mysore, William "Jamie" Tyler, Marco Santello

Address for correspondence:
Justin M. Fine
justfine@iu.edu
480-299-8845
Psychological and Brain Sciences
1101 E 10th St
Indiana University
Bloomington, IN 47405

Keywords: Transcranial Focused Ultrasound; Response inhibition; Dynamic Causal Modeling; Inferior frontal gyrus; Cognitive control; Pars Opercularis; Neuronavigation

Abstract

To prevent erroneous actions, individuals must often inhibit prepared behavioral

responses. The right inferior frontal gyrus (rIFG) and its connectivity patterns are

prominently implicated as key to behavioral inhibition. However, previous studies have

applied neurostimulation methods with low spatial resolution that impede simultaneous

network modeling of neural activity. Therefore, direct evidence for inhibitory control in

rIFG is lacking, while the accompanying network mechanisms remain unknown. We

addressed this gap using a Stop Signal task and transcranial focused ultrasound

(TFUS) to pars opercularis in rIFG. TFUS improved stopping performance by enhancing

stopping speed. Electroencephalographic dynamic causal modeling indicated inhibition

performance increased by TFUS modulating pars opercularis pyramidal neuron

connectivity to subcortex. By combining TFUS and network modeling, our results

provide causal evidence that response inhibition is implemented along two pathways

originating from a direct rIFG to subcortical pathway and a parallel pathway that

modulates pre-SMA inhibition onto subcortical nodes.

Introduction

Behavioral inhibition is necessary to suppress impending actions that become contextually inappropriate (Aron et al., 2014; Baddeley, 1996; Logan and Cowan, 1984). The control over inhibitory capacities is dramatically reduced in pathologies dominated by aberrant impulse control, e.g. ADHD (Bari and Robbins, 2013). The stop-signal task has been widely used as a paradigm for probing inhibition (Logan and Cowan, 1984). This task involves cueing action execution (Go signal) on every trial. On a percentage of trials, individuals are cued (Stop signal) to attempt inhibiting responses at a delay after a Go. This task allows deriving the stop signal reaction time (SSRT), a latent measure of stopping speed.

The predominant conceptual framework implicates a right-lateralized prefrontal stopping circuit driving inhibition (Aron, Robbins, and Poldrack, 2014; Chambers et al., 2006), with an anatomical locus in the posterior portion of the right inferior frontal gyrus (rIFG), pars opercularis (Aron and Poldrack, 2006). It has been argued this area directly implement motor braking via projections to subcortical nodes (Aron et al., 2014). Supporting evidence is based on demonstrations that rIFG neural activity is larger in successful compared to failed stopping (Aron et al., 2006; Boehler et al., 2010; Li et al., 2006), and both inhibition and SSRTs being altered in individuals with rIFG lesions (Aron et al., 2003) and ADHD (Morein-Zamir et al., 2014). rIFG is considered a core node for response inhibition, nevertheless successful inhibition also engages a broader network that includes pre-supplementary motor area (pre-SMA; Duann et al., 2009), subthalamic nucleus (STN) and striatum (Aron et al., 2007; Mallet et al., 2016).

Although modulation of RIFG activity typically accompanies inhibitory control,

several researchers have proposed RIFG's involvement is indirect (Duann et al., 2009;

Sharp et al., 2010; Xu et al., 2017). This indirect role involves rIFG registering a stop-

signal context, and signaling the context to pre-SMA, which has been argued to

explicitly trigger inhibition (Duann et al., 2009; Sharp et al., 2010; Rae et al., 2015). At a

network level, this hypothesis has gained support from fMRI connectivity in stop-signal

studies wherein pre-SMA alone exhibited modulated connectivity with subcortical

structures during successful inhibition (Duann et al., 2009; Rae et al., 2015). However,

other connectivity studies have implicated both rIFG and pre-SMA connectivity to STN

and striatal pathways as predicting inhibition speed (Jahfari et al., 2011; Xu et al.,

2017). Additionally, primate research implies that direct neural projections to the STN

originate in both the rIFG and pre-SMA, with the STN acting as an integrator (Haynes

and Haber, 2013). These dual-pathway models leave open the possibility that rIFG can

directly trigger inhibition in parallel with pre-SMA (Aron et al., 2016).

In contrast to the above conclusions, an alternative framework posits that neither

rIFG or pre-SMA directly implement inhibition, with inhibition emerging from attentional

orienting and biased competition processes (Hampshire and Sharp, 2015; Chatham et

al., 2012). This proposition is based on findings indicating sectors of rIFG are

equivalently active during stop-signal and other putatively non-inhibitory tasks (Erika-

Florence et al., 2014; Hampshire et al., 2010; Xu et al., 2017). For example, rIFG

activity scales with stimulus probability (Shulman et al., 2009) and regularly during tasks

requiring attentional re-orienting (Vossel et al., 2006; Levy and Wagner, 2011). The

claim that attentional orienting drives response inhibition is supported primarily by

demonstrating rIFG fMRI activity is equivalent during both stop-signal tasks and other

tasks with no apparent inhibitory demands (Sharp et al., 2010). In this framework

(reviewed in Hampshire and Sharp, 2015), inhibition could occur through rIFG top-down

signals that bias attentional processing through increasing synaptic efficacy of sensory

cortices (Hampsire, 2015; Feldman and Friston, 2010).

Given the prevalence of findings supporting either a direct or indirect role of rIFG

in response inhibition, or the absence of behavioral inhibition processes altogether,

delineating between alternative mechanisms has remained inconclusive. For example,

an established way to control for attentional demands is comparing neural activity

during a stop-signal and putatively non-inhibitory tasks. However, a notable issue with

this comparison is that other tasks may still induce unaccounted for cognitive processes

or latent inhibitory demands not directly measurable in behavior (Aron et al., 2014).

Given the ambiguity introduced by comparing tasks to parcellate neural activity

underlying inhibitory versus other cognitive demands, direct approaches are needed to

circumvent these issues.

Neurostimulation during inhibitory tasks offers a potential in-route to identify how

rIFG, and particularly pars opercularis, is causally involved in motor braking, while

detailing its role in the broader inhibition network. Respectively, several studies have

applied transcranial magnetic (TMS; Cai et al., 2012; Obeso et al., 2013; Verbruggen et

al., 2010) or direct current (Jacobson et al., 2011) stimulation during inhibition tasks.

Some studies found offline TMS applied to either rIFG or pre-SMA impaired or improved

inhibition performance (Chambers et al., 2006; Verbruggen et al., 2010), respectively.

However, others have shown pre-SMA TMS can either improve inhibition with no effect

on rIFG (Obeso, 2013). Limitations of previous human neurostimulation response

inhibition studies include either lack of neural activity measurements or limited spatial

accuracy (Opitz et al., 2013). For example, TMS applied to rIFG during inhibition tasks

most likely engaged ventral premotor areas involved also in action switching (Buch et

al., 2010). The implication is that, if rIFG is potentially involved in attentional orienting,

inhibitory control, or both, these neurostimulation approaches likely elicited broad

effects on these processes. Therefore, it remains to be causally established that rIFG

and, more importantly, pars opercularis, engage an explicit motor inhibition mechanism

embedded in its connectivity that operates alongside attentional mechanisms (Munkata

et al., 2011; Wiecki and Frank, 2013).

Here, we employed MRI-guided, neuronavigated transcranial focused ultrasound

stimulation (TFUS) directly to the pars opercularis of the rIFG while humans performed

a stop-signal task. TFUS is a stimulation technique with a millimeter spatial resolution

(Fini and Tyler, 2017). Neural activity underlying response inhibition was assessed with

EEG event-related potentials (ERPs) and source analysis. Using this approach allowed

us to delineate the specific role of pars opercularis, while detailing which ERPs and

network functions are directly related to inhibition success and its speed (SSRT). TFUS

to pars opercularis significantly improved response inhibition through a targeted effect of

shortening SSRT. To determine how TFUS altered biophysical mechanisms generating

neural activity underlying inhibitory mechanisms, we built dynamic causal models

(DCM) of ERPs using microcircuit models. The main network hypothesis was that an

explicit rIFG inhibition mechanism would be embodied in direct rIFG-to-subcortical

connectivity weighting that directly reflects TFUS-induced changes in stopping efficiency

(SSRT). We confirmed this hypothesis by demonstrating TFUS directly altered rIFG

connectivity to a hidden subcortical node. In addition, DCM also indicated that only No-

TFUS successful versus failed stopping, rather than successful stopping in general,

were differentiated by mechanisms linked to network mechanisms associated with

attentional modulation, i.e., recurrent synaptic superficial gain of visual cortex.

Importantly, these results support the proposal that rIFG is directly involved in

implementing an explicit response inhibition function and stopping efficiency.

Methods

Participants

Participants consisted of healthy adult volunteers and were divided into one of

three experimental groups. The main experimental group received transcranial focused

ultrasound (TFUS) stimulation to the right inferior frontal gyrus (rIFG) (n = 25; 19 males,

mean age 24.1 yrs SD 3.2 yrs). A second group was used as cortical site, active control

group. These participants received stimulation to the ipsilateral somatosensory cortex

(S1) (n = 23; 15 males, mean age 22.4 yrs. SD = 3.3 yrs). A third group received a

sham stimulation near the right temple (n = 15; 8 male, mean age 24.2 yrs SD = 2.8 yrs)

and was used as control for possible auditory effects of TFUS over rIFG (sham rIFG).

All individuals were right-handed and received financial compensation for participation

in the study. Before being enrolled, each subject was screened for neurological

disorders and previous history of epilepsy, stroke, or brain injury. Furthermore, a

neurologist from the Barrow Neurological Institute (Phoenix, AZ) screened all subjects'

T1 scans after structural MRI acquisition, and before participation in the study.

Stop Signal Task and TFUS design

The current study used the conventional Stop Signal Task that involved both 'Go'

and 'Stop' trials (Fig. 1A). We presented the experiment using Opensesame (Mathôt et

al., 2012). Each trial started with a centrally-located fixation cross on the monitor. In

both trial types, the fixation cue was replaced by a green 'Go' circle (3° x 3° visual

angle), with an exponentially-distributed time interval (mean: 500 ms; standard

deviation: 50 ms). Subjects were instructed "to press the up key as soon as they

detected the Go circle" (top panel, Fig. 1A). In 'Go' trials, the circle vanished when the

button was pressed or after 800 ms had passed from the fixation cross stimulus. In

'Stop' trials, the stop was a red square which appeared around the green circle (middle

and bottom panel, Fig. 1A). If the subject successfully inhibited his/her response with

respect to the Stop cue within 800 ms, the red square was extinguished, and the trial

was considered a successful inhibition. The time required to inhibit a response following

the Stop signal is defined as stop signal reaction time (SSRT) (see below). The timing of

the Stop cue relative to Go cue, i.e., the stop signal delay (SSD), was presented at one

of four fixed, but subject-specific SSDs. The SSDs were designated by having each

subject first perform a practice block of 50 Go trials only to determine their baseline Go

reaction time (RT). After this block, the 4 SSD levels were set to 25, 35, 75 and 95% of

the mean Go RT. These SSDs were fixed throughout the experimental session. Using a

set of fixed SSDs allowed us to calculate the SSRT using routines that are less

susceptible to low trial numbers (Matzke et al., 2013; see Data processing). Additionally,

we sought to determine the effects of online TFUS at different SSDs to estimate the

effects of stimulation on neural and behavioral responses at different stages of a Go

process predicted by response inhibition models (Verbruggen and Logan, 2009). All

trials were separated by an inter-trial interval of 2000 ms (±300 ms randomly drawn

jitter).

TFUS was delivered either simultaneously with (1) the Go signal in both Go and

Stop trials, or (2) the Stop signal (Fig. 1A). The purpose of delivering TFUS during Go

trials was to determine the neural and behavioral effects of TFUS to rIFG independent

of a stopping signal. Specifically, this allowed us to assess whether any effects of TFUS

on stopping behavior are related merely to alteration of the timing of an underlying Go

process. Therefore, we used 5 types of trials. The first two consisted of Go trials with no

TFUS or with TFUS locked to the Go cue (No-TFUS and Go-TFUS trials, respectively).

The other three trials consisted of Stop trials: No-TFUS trials, Go-TFUS, and TFUS

locked to the Stop signal (Stop-TFUS). These three types of Stop trials were examined

across the four SSDs.

TFUS delivery for Stop trials was evenly distributed across the 4 SSD levels. The

overall probability of a stop trial was set to 35% of all trials (Fig. 1B). We chose this level

to accommodate the need for large amounts of Stop trials required to examine the

effects of TFUS on Stop trials across all SSD levels, while still making Go trials more

frequent. Each experimental session consisted of 1200 trials distributed across 12

blocks. Blocks were segmented into stimulation and no-stimulation blocks; the former

containing trials with and without stimulation, and the later containing no stimulation.

Trial types (Go and Stop trials) were randomly and evenly distributed throughout the

experiment. The trial numbers were chosen to enable the comparison between TFUS

and non-stimulation trials across successful and failed inhibition trials, while allowing a

reasonable number of trials to be performed without inducing significant fatigue to the

participants. The block design, as well as the use of an active-stimulation and sham

control groups was chosen to mitigate any possible carry-over effects of the stimulation

across trials.

EEG acquisition

EEG was recorded using a 64-channel ActiCap system (BrainVision, Morrisville,

NC), with a standard 10–20 layout. Data was recorded at a sampling rate of 5 kHz, with

resolution 0.1 µV and bandpass filter of 0.1–100 Hz. Impedances were always kept < 5

kΩ. Online recordings utilized a ground at AFz and left mastoid reference. At the

beginning of each session, electrode layouts with respect to each individual's head

shape were registered using a CapTrak camera system (BrainVision, Morrisville, NC)

with the left and right preauricular, and nasion as fiducial landmarks. This allowed for

later co-registration with each individuals T1 structural MRI scan and for source-

localized analysis (see Data Processing).

Structural MRI acquisition (T1) and processing

For purposes of TFUS neuronavigation and co-registering EEG electrode

placement, we obtained a structural T1 MRI scan for each participant. T1 volumes were

collected using an 3D MPRAGE sequence (TR = 2300 ms, TE = 4.5 ms, 1 x 1 x 1.1

mm^3 voxels, field of view 240 x 256 mm^2, 180 sagittal slices) in a Philips Ingenia 3T

scanner with a 32-channel head coil. Brainsuite was used to process T1s, which

included cortical extraction sequence and a surface label-registration procedure with the

BCI-DNI atlas. After labeling, we checked the locations and created a mask of either

pars opercularis (rIFG group) or the centroid of ipsilateral S1 (control group). This

volume labeling and mask creation procedure was used for guiding TFUS target

identification.

TFUS targeting, setup and parameters

All stimulation targets were planned prior to subject arrival. We used a Brainsight

neuronavigation system (Rogue industries) with subjects' T1 scans to guide placement

of the transducer beam profile with respect to each individual's neuroanatomy. First, we

created a subject-specific mask from the cortical atlas registration and projected it into

the Montreal Neurologic Institute (MNI) coordinate system (Evans et al., 1994). When

planning the TFUS target, we considered both MNI coordinates and individual anatomy.

For example, neuroimaging studies (Boehler et al., 2010) and metanalysis (Chikazoe et

al., 2009; Levy and Wagner, 2011) have shown specific activation of the pars

opercularis (around x=48, y=16, x=18) for contrasts of successful inhibition versus Go

trials and successful versus failed inhibition trials. In the case of the RIFG group, we first

identified these MNI coordinates. Notably, the pars opercularis is an anatomical

definition and is often referred to as ventro-lateral prefrontal cortex in neuroimaging

studies focused on localization of activity that is functionally related to response

inhibition and cognitive control (Levy and Wagner, 2011). During target planning, we

confirmed the identified MNI coordinates were inside the anatomical region of the pars

opercularis, identified from registering atlas maps to individual anatomy. We also

performed visually confirmation that the TFUS target was indeed rostral to the inferior

precentral sulcus, dorsal to the sylvian fissure, caudal to the ascending rhomulus of the

syvian fissure, and ventral to the inferior frontal sulcus (Tomaiuolo et al., 1999).

Because significant anatomical variation exists in this region, individual anatomy rather

than coordinates were prioritized when planning the TFUS focus. For the S1 group,

stimulation was targeted near x=-43, y=-29, z=54 and within the left post-central gyrus.

Because we used a single element transducer, with a fixed focal depth of 30mm and a

5mm silicon spacer, all stimulation was done at a penetration depth of 25mm and

normal the surface of the scalp.

After EEG setup, we used an infrared optical tracking system (Polars Vicra, NDI

Medical) to register the subjects' structural MRI scans in virtual space, with their head

and the ultrasound transducer in real space. The alignment and cortical registration

were accomplished by registering the individual's T1 derived anatomy using the nasion,

tip of the nose, philtrum, and left and right periauricular notch and tragus. To visualize

the TFUS target in the cortex, we created a custom design in Solidworks that rendered

the transducer housing and ellipsoidal beam profile projection into the registered cortex

(Fig. 1C). A 3D printed housing was made for the transducer to hold the optical tracking

unit and silicon spacer (ss-6060 Silicon Solutions, Cuyahoga Falls, OH

http://siliconesolutions.com/ss-6060.html). Acoustic conductive gel was applied to both

the transducer and the scalp. After correct placement of the transducer using the

neuronavigation, we recorded the coordinates of the stimulation target. Figure 1C

shows the rendering from one subject's T1 and scalp in the rIFG group, along with the

3D rendering of the transducer housing (green object) and the pars opercularis mask

(white anatomical structure). In the auditory rIFG control group, we employed a sham

TFUS (similar to Legon et al., 2018) by placing the gel coated transducer perpendicular

to the rIFG target. This sham procedure was done to ensure there was still an auditory

effect of the ultrasound (from the pulse repetition frequency) without active stimulation.

The TFUS transducer was held flush to the head with a custom-made,

lightweight, elastic mesh cap, which did not interfere with EEG recording. To ensure

accurate TFUS placement throughout the experimental session, the rotational and

cartesian displacement of the beam profile from the cortical target was tracked. The

overall accuracy was measured as deviation from the original alignment of the beam

with the anatomical target. During the experimental session, we sampled the position of

the TFUS transducer during each break. Accuracy was very high, with an average

deviation of ±1.5 mm displacement across all subjects and sessions.

The setup and parameters used for TFUS in this experiment were nearly

identical to those used by Legon et al. (2014). We used a broadband, single-element

focused ultrasound transducer with a center frequency of 0.5 MHz, a fixed focal depth of

30mm, and a lateral spatial resolution of 4.5 mm^2 and axial spatial resolution of 18mm^2

(Blatek, Inc., State College, PA) (Legon et al., 2014). Prior water tank testing through

cadaver skull revealed transcranial spatial-peak pulse average intensity (I_{sppa}) of 5.8

W/cm^2, and the optimal frequencies for TFUS transmission while minimizing cranial

attenuation are 0.2- 0.65 MHz (Hayner and Hynynen, 2001; White et al., 2006).

The TFUS waveforms were generated using a two-channel, 2 MHz function

generator (BK Precision) (Legon et al, 2014). Channel 1 was triggered by the

presentation computer and produced the pulse repetition frequency (PRF) of 1.0 kHz.

This was used to trigger channel 2, which produced short burst at the 0.5 MHz acoustic

frequency. The result produced a ultrasound waveform with a carrier frequency of 0.5

Mhz, PRF of 1.0Khz, and duty cycle 24%. Each stimulation duration was 0.5 s. The

transducer power was driven by sending channel 2's output to a 40-W linear RF

amplifier (E&I 240L; Electronics and Innovation). The waveforms were triggered in

alignment with experimentally-relevant temporal events (see description below and Fig.

1A). It has been previously verified that the resulting waveform does not incur any

heating of skin or skull bone (Legon et al., 2014).

Computational simulation of TFUS propagation

We quantified peak pressure amplitude, peak intensity and accuracy of the beam

distribution with TFUS target to rIFG using the pseudospectral simulation method in K-

wave (Treeby and Cox, 2010). Reference peak pressure planes for the simulations

were derived from a water tank test and previous data (Legon et al., 2014). Simulation

parameters were first validated by simulating the transducer in water to compare the

simulation results with those from the water tank test. The max pressure plane at the

30-mm focus in the water tank was used as a source input pressure for the transducer

during the simulation. The transducer was modeled to have a 30-mm radius of

curvature. Water simulations used a homogenously medium of water density (1000

kg/m^3) and speed of sound (1482 m/s). We created a computational grid over a 256 x

256 x 256 with 1-mm spacing. The points per wavelength were 6, Courant–Friedrichs–

Lewy = 0.1, and simulation time was set to 6 pulses (duration = 250 µs) to ensure

simulation stability.

Simulation of ultrasound through water predicted a max pressure of 1.05 Mpa

and spatial peak pulse average intensity (Isppa) of 22.4 W/cm^2 at the focus. This

prediction closely aligns with previous studies and simulations (Legon et al., 2014) of

the same transducer. Comparison of simulations and water data indicated a 97% match

of pressure/intensity at the focus taken over a 5 mm^3 voxel section in all 3 planes at the

focus. The lateral full-width at half maximum of the max pressure at the beam was 4.39

mm in simulation (Fig. 1C).

For simulating transcranial US, we extracted 3-dimensional maps of the skull

from a CT (1-mm resolution) and brain from T1 MRI scan (1-mm resolution) from three

preoperative patients at Barrow Neurological institute. The MRI and CT were both co-

registered and normalized to the MNI space in SPM12. To mimic our approach of

targeting used in the experiments, we surface registered the gray matter volume to the

BCI-DNI atlas and identified the centroid of pars opercularis. This allowed us to map

from world coordinates of the scan to MNI coordinates of the target (Fig. 1D). The

average stimulation location for these three subjects was x = 48, y = 18, and z = 6.

Conversion from Hounsfield units in the CT to sound speed and density were done

using the relations described in Aubry et al. (2003). All skull materials were set using

these parameters, while other tissues were treated as homogeneous with parameters

set to that of water. Attenuation was modeled as a power law with a $\beta = 0.5$ and

absorption was also modeled with a $b = 1.08$ (Treeby and Cox, 2010). Results for this

simulation are presented in the Results section.

335   **Data processing and experimental variables**

Behavioral variables

The main variables under consideration were the Go trial reaction time (Go RT),

percentage of successfully inhibited responses on Stop trials (successful stopping) per

SSD, failed inhibition reaction time, and SSRT. The SSRT was estimated using a

hierarchical Bayesian parametric approach (Matzke et al., 2013a) that allows estimation

of the distribution of SSRTs while assuming an ex-gaussian distribution of SSRTs.

Importantly, we chose this approach as Matzke et al. (2013a) showed that it performs

well even when there are only a few trials available per SSD level. This SSRT

estimation procedure was run separately per group (rIFG and S1) and trial types (No-

TFUS Stop trials, Go-TFUS Stop trials, and Stop- TFUS Stop trials). As we report in the

Results section, we used combined RTs from Go trials with and without TFUS because

stimulation did not alter the Go RT.

EEG artifact removal

Continuous EEG data were first down-sampled to 250 Hz, then high-pass filtered

(1 Hz) and re-referenced to the scalp average. Any channels that displayed artifacts for

more than 25% of the total session were removed before further processing but were

later interpolated after artifact rejection. We removed channels that were designated

unsuitable for analysis by visual inspection and absolute temporal standard deviation (>

5) across channels. It is important to note that, in each of the stimulation groups, the

cortical sites of rIFG and S1 were close to the F8 and CP4 electrodes. Therefore, these

electrodes could not be used for EEG recording in their respective groups and were

always interpolated after artifact removal. The remaining data processing involved

creating epochs from Stop trials locked to stop signal onset (-100 to 500 ms

peristimulus) because our analysis focused on this epoch. Individual epochs were then

rejected from further analysis if they contained large scalp EMG or rare events (< 8% of

all trials). The ERPs baseline corrected by subtracting the activity from -100 ms to the

stop signal. Out of the 53 participants, 3 subjects were excluded from analyses due to

EEG recording issues (impedance >25 kΩ across channels). The remaining data were

bandpass filtered from 1-25 Hz. EOG artifacts related to eye movements and blinks

were removed using Independent Components Analysis using eeglab (ICA; Delorme

and Makeig, 2004). On average, 3.35 components were removed per participant.

Because we later applied a Hanning taper to the edges of the event-related potentials

(ERP) before dynamic causal modeling, we applied the same procedure to the ERPs

after cleaning using a Hanning taper.

370   **QUANTIFICATION AND STATISTICAL ANALYSIS**

Behavior analysis

To quantify how the probability of response inhibition changed over levels of

SSD, P(respond|signal), and across TFUS conditions within and across groups we fit a

2-parameter logistic model. This was done to analyze P(respond|signal) as a curve.

This was achieved using by fitting the 2-parameter linear mixed-effects model with

random intercepts and slopes to obtain subject and condition specific model

parameters. P(respond|signal), denoted as p, were converted to a negative logit (log((1-

p)/p) before fitting. As our main goal was to estimate the logistic curve slope (**β)**, we ran

the mixed-effects model (using LME4 in R) with the full interaction of SSD and

stimulation condition (no-TFUS, Go-TFUS, Stop-TFUS). Logistic slopes per subject

were estimated by combining fixed and random coefficients. **β** parameters were

analyzed using a mixed-design ANOVA on **β** with factors of Group (3 levels) and TFUS

(3 levels: No, Go, Stop).

Separating neural components of Go-responses and response-inhibition

Our analysis of ERPs was based on the premises of the Independent Horse

Race model (Logan and Cowan, 1984) – which posits independent accumulation of Go

and Stop activity till one of them reaches threshold. Due to the nature of measuring

these processes through EEG and the inhibitory processes, there is overlap of neural

processes related to stopping and going during stop trials. Therefore, we removed Go-

related activity from both successfully (SS) and unsuccessfully (US) inhibited Stop trials

through subtracting the Go ERPs from the Stop ERPs. We used the approach

employed by Mattia et al. (2012). On a per-subject basis, we found Go-trial ERPs (No-

TFUS and TFUS) that had RTs that were latency-matched to SS trials based on each

subject's SSRT. These Go trials had to have RTs either equal to or greater than the

SSRT. For US trials, we found latency-matched Go trials with RTs with a different

procedure. We first calculated each subject's mean signal-respond RT for each of the

two highest SSDs. We then calculated the difference in SSD (ms) and searched for Go

RT trials for each SSD that fell within the mean signal-respond RT ± half the difference

of the SSD (ms). This was done to prevent overlap of activity from both faster and

slower Go RTs and signal-respond RTs. These steps were performed separately for the

highest and second highest SSD. This procedure was done separately for SS and US

trials for both TFUS conditions. After correcting the SS and US stop trial, the corrected

ERPs were averaged across the two highest SSDs per subject (corresponding to the

85% and 105% mean Go RT of each subject). These ERPs were used for the remaining

analysis.

Analysis of inhibition-related ERP

Our analysis focused on event-related potentials (ERPs) from source-localized

analysis and Dynamic Causal Modeling (DCM; David et al., 2006). The primary

motivation for these analyses is that previous work has revealed a set of ERPs that

often accompany response inhibition following a Stop signal (reviewed in Huster et al.,

2013 and Kenemans, 2015). The main ERPs found in inhibitory tasks include a N2/P3

complex that has a fronto-central and radial topography. This component has been

hypothesized to be generated mainly by the pre-SMA/SMA in the medial frontal cortex

(Huster et al., 2013) and has been considered to reflect a critical signature of reactive

stopping elicited by stop signal tasks (Kenemans, 2015; Wessel and Aron, 2017).

Furthermore, the P300 of this complex has been proposed to be a relevant marker of

stopping efficiency, given that it predicts the SSRT (Wessel and Aron, 2015) and

successful versus failed inhibition (Kok et al., 2003). The ERP associated with stopping

in rIFG is typically associated with a negative amplitude difference comparing

successful and failed stopping that emerges around 200 ms (N200; Schmajuk et al.,

2006). We examined these ERPs and P/N100 responses which are sometimes elicited

over sensory areas, depending on whether the stimuli used for Go and Stop signals are

in the same sensory modality (Kenemans, 2015). By combining TFUS and EEG, we can

examine some of the issues that are addressed by ongoing debate, i.e., which of these

potentials are stopping-relevant, and how their activity is generated by a network model

through DCM – all without signal interference induced by stimulation.

Scalp space analysis.

To examine the standard ERP effects typically found in the SST, we first

examined activity at the sensor level. This was done using permutation-based

dependent samples t-tests. Spatiotemporal activity was examined and multiple-

comparison corrected for using a cluster-based p-value correction of $p < 0.01$ and 5000

permutations for each contrast considered. The contrasts included comparison of (1)

successful stop (SS) – unsuccessful stopp (US) trials over Go-TFUS and Stop-TFUS

conditions, a (2) SS (No-TFUS) – SS (Stop-TFUS) contrast, and (3) and interaction

contrast comparing SS – US difference between the No-TFUS and Stop-TFUS

conditions. The first contrast is typically used to determine which areas exhibit ERPs (or

brain areas) that differentiate successful inhibition (Swaan et al., 2012). The third

contrast (interaction) was used to determine how the SS-US contrast differed between

No-TFUS and Stop-TFUS conditions. The second contrast was the main focus of our

analysis and was used to determine which scalp ERPs differentiated successful

stopping in the No-TFUS and Stop-TFUS conditions. We anticipated that this contrast

would reflect processes that mainly include those responsible for both the success and

efficiency of inhibitory processes, e.g., SSRT.

P300 onset and SSRT TFUS effects

Recent work has indicated that the frontocentral P300 onset latency is related to

the SSRT (Wessel and Aron, 2015). Therefore, we hypothesized that TFUS altered

inhibition through the SSRT and expected a shift in this latency as well. We calculated

the shift in P300 onset crossings between TFUS conditions in two steps. First, we took

the across-subject mean frontocentral ERP waveform in a time-window of ±50 ms

around the zero-crossing. To calculate each subject's zero-crossing time, we calculated

the dynamic time warping distance from the template mean ERP to the subject's ERP.

Second, this distance was added to the median zero-crossing time to obtain an

individual subject crossing for both the No-TFUS- and Stop TFUS-locked conditions.

Source localization

To estimate the activity in source space from the sensor recordings, we used a

group inversion with the multiple sparse-priors approach as implemented in SPM12.

The individual subject data passed to the inversion routine were mean sensor ERPs per

condition. We performed the group inversion only for the rIFG groups and analysis

because this was the only group to exhibit behavioral effects from TFUS. In this

procedure, the individual's recorded electrode locations and individual T1 were warped

to the default MNI anatomical brain and cortical mesh provided in SPM. These meshes

were used to calculate the forward solution using a boundary element head model. All

conditions were used in the group inversion routine. Because the multiple sparse-priors

approach attempts to fit the sensor data with respect to the lead-field matrix, we

performed the inversion over a window starting from the stop signal up to 500 ms. While

narrow windows are considered better for time-resolved estimation, we wanted to

estimate overall changes in activity using the same window we would employ later for

dynamic causal modeling.

Source analysis: Whole-brain contrasts and regression

Based on examination of the sensor level data, we first identified time windows

surrounding the ERPs discussed above (N100, N200, and P300). To balance the

number of points contributing to the source estimate used for analysis, we used the

same window size for estimating the source activity of each ERP. The time windows

were centered around the across-subject mean peak activity of each ERP with a

window of ±40 ms. These time windows were used to create 3D source image activity

interpolated into MNI voxel space for each subject and condition. The resulting images

were spatially smoothed (6 mm full-width half maximum) evoked power from 1-25 Hz.

We used evoked power because we were concerned with 'activation' and time window,

but not the direction of voltage deflection. This choice was driven by the fact that we

already established the canonical inhibition related ERPs at the scalp level. These

evoked images were analyzed used a flexible factorial design to implement a repeated-

measures ANOVA, including all main effects and interactions. The factors included (1)

inhibition success (SS or US trial), and (2) stimulation condition (No-TFUS or Stop

TFUS). The resulting statistical parametric maps were analyzed with a threshold set at p

< 0.005 (peak-level, uncorrected) and cluster-wise FWE $p < 0.05$. For expositional

brevity, these SPMs are presented in the supplementary material except for the

conjunction F-contrast showing the overlap of the SS-US and TFUS F-contrasts. This

conjunction shown in the main text both confirms the differential SS-US effect in rIFG

and the spatially precise impact of TFUS.

We also conducted a whole-brain SPM linear regression using the same time

windows identified above. We regressed the difference in evoked activity between No-

TFUS and Stop-TFUS SS trials against the difference in each subject's SSRT for these

conditions. Using this approach, we examined both positive and negative contrasts (t-

tests) in each window. This analysis was done to (1) determine prior source locations

for the DCM analysis (see below) and (2) identify which areas predicted the change in

SSRT due to TFUS. Because a primary goal was determining DCM priors, we used a

lenient uncorrected cluster threshold of $P < 0.01$ and a minimum cluster-extant

threshold of 20 voxels.

Dynamic Causal Modeling (DCM)

To model the connectivity of the network involved in successful stopping, we

used the canonical microcircuit model (Bastos et al., 2012) for all areas except the

contralateral motor cortex. For left motor cortex (M1), we used a recently developed

version built-off the canonical microcircuit that more closely resembles the agranular

structure of M1 (Bhatt et al., 2016). Our sequential model building and analysis focused

both on a priori areas of interest that have been well described in response inhibition

literature (e.g., Wessel and Aron, 2017) – including rIFG, rpre-SMA, rDLPFC – and

those areas that we identified in comparing the effects of Stop TFUS and No-TFUS

changes in SSRT on evoked power. Given the typically right-lateralized areas found to

be modulated by inhibition, we focused mainly on pathways on the right hemisphere.

We used source locations identified in both (1) the regression of change in SSRT and

(2) the whole-brain SPM interaction F-contrasts examining locations for which Stop-

TFUS and No-TFUS were different for the SS-US comparison. The significant source

cluster peaks ($P < 0.01$ threshold) revealed by these analysis were labeled using the

AAL atlas labeling toolbox, and used to identify source coordinates in MNI space

([x,y,x]: rIFG [48,28,4]; pre-SMA [6, 24, 54]; rDLPFC [30, 28, 40]; rParietal [10,-74,56];

rTemporal [52,-18,-12]; LM1 [-37,-25,-62]; right inferior occipital gyrus (rIOG) [46, -

76,10]). These source locations are in general agreement with previous literature in both

fMRI, EEG, and MEG studies of SSTs (Aron et al., 2006; Boehler et al., 2010; Rae et

al., 2014) and meta-analyses of cortical locations and boundaries (Chikazoe et al.,

2009). In building the model, we also included a hidden deep source to model the

potential connectivity effects to and from cortical sources. Because the main output of

the basal ganglia mediating inhibition and responding is the thalamus, we used a source

location [4, -16, -2] identified during a previous fMRI stop signal study (Boehler et al.,

2010). This approach of using a deep source node in DCM for EEG has been previously

employed by David et al. (2011). In the DCM, we used an equivalent current dipole

model and allowed the inversion process to optimize the source locations.

To determine the appropriate model connectivity structure, we first examined the

grand mean ERP data in SS No-TFUS trials from a window spanning 0 – 500 ms.

Recent work has shown that estimating the model structure from grand means (across

subjects) is sufficient and provides a close approximation to fixed-effects selection when

using ERP data (Litvak et al., 2015) rather than fitting DCMs across all subjects for each

variation in connection structures.

To optimize the model structure, we performed Bayesian model selection using

family-wise fixed effects in several iterations. In the first set of iterations, we inverted 48

different models from the grand mean data, and then partitioned it into several families

(Penny et al., 2010). These families were based on (1) the structure of the pre-frontal

hierarchy (Fig. S1), (2) structure of the lower hierarchy (Fig. S1), (3) whether exogenous

inputs were supplied to right inferior occipital gyrus (rIOG), rIFG, or both, and (4)

whether the cortical nodes projecting to the hidden deep source were of the forward or

backward type. Inputs were modeled as a Gaussian bump with a prior onset of 60 ms.

With respect to comparison (4), pulling evidence from previous inhibitory control,

primate tracing, and tractography studies and reviews, the areas we examined for

different connections projecting to the deep node included rIFG, pre-SMA, right

dorsolateral prefrontal cortex (rDLPFC) and rIOG. A backward connection from M1 to

the deep node was also included based on putative M1 to basal ganglia connections

(Nambu et al., 2002). Because the efferent connections from Thalamus output is

excitatory and has been found to target deep pyramidal layers (Yamawaki et al., 2014),

the above listed areas all received forward connections from the deep source.

In this modeling, we assumed hierarchically connected areas always entailed a

backward and forward connection between nodes. Nodes that were lateral in a

hierarchy were supplied with both forward and backward nodes (Fig. S1). A final note is

on the connections between lower areas and prefrontal areas. Rather than optimize all

possible connection permutations, we chose to instantiate connections that reflect the

cognitive and attentional control-related differences typically postulated to operate in

dorsal and ventral pathways (Corbetta and Shulman, 2001). Both streams received

forward input from rIOG. The ventral stream included a forward connection from

rTemporal cortex to rIFG. The dorsal stream included forward projections from rParietal

to both rDLPFC and pre-SMA.

After inversion of each model, we used a fixed-effects Bayesian model selection

to perform family inference and calculate the model posterior probability to determine

the winning model in each family. The family model probability results from these four

different family comparisons are shown in Figure S2. This analysis indicated that,

across families, the winning model had (1) a prefrontal hierarchy with laterally

connected rDLPFC and rIFG above pre-SMA, (2) a parallel structure of rParietal and

rTemporal cortices without a lateral connection, (3) exogenous inputs to both rIOG and

rIFG, and (4) backward cortical to deep connections. A diagram of the final model

structure is shown in the DCM results section.

The next step in model building involved determining which connections were

modulated by conditions specific changes between No-TFUS SS and US trials and,

primarily, between No-TFUS SS and Stop-TFUS SS trials. For our purposes, this

included extrinsic connections between areas and intrinsic gain connections within an

area. Instead of testing an expansive set of permutations of condition-specific

modulations of extrinsic connections (B matrix in DCM), we a priori opted to use the

recently developed Parametric Empirical Bayesian modeling framework for DCM

(Friston et al., 2016). As will be explained below, this involves doing Bayesian statistics

on full DCMs that will have all condition modulatory parameters of interest entered a

hierarchal-general linear model from which hypotheses can be tested. This obviates the

need for conventional statistics and model reduction over all extrinsic connections. If a

DCM model is referred to as full, this includes condition-driven modulations of all

extrinsic connections. Otherwise it includes an explicitly stated set of connections.

After determining a model structure, we inverted this model for each subject's

ERPs. This inversion was performed twice, using different combinations of trials to

assess different hypotheses. The DCM was first inverted using the No-TFUS US and

SS trial ERPs. The US trial was set as the baseline. This inversion allowed us to first

compare how the network connections were modulated between US and SS trials in a

baseline network without the effects of TFUS. Specifically, we wanted to examine how

changes in connectivity (DCM B matrix) distinguished between unsuccessful and

successful inhibition. Analysis of this general linear model (described below) focused on

the B matrix which describes how connections changed between conditions. This

comparison was done to mirror the standard comparison of SS and US trials that is

typically employed to reveal what inhibitory mechanisms were more potently activated

during SS trials (Aron et al, 2014). Furthermore, using this contrast of conditions

provides a baseline to ask whether the same connectivity mechanisms were altered

when comparing No-TFUS and Stop-TFUS SS trials. Therefore, the second DCM model

inversion was applied to the No-TFUS and Stop-TFUS SS trials. The No-TFUS trials

were used as a baseline for the fit. Again, this allowed a focus on the connectivity

modulation between the two conditions and treated the non-stimulation condition as the

baseline network.

To analyze the resultant DCM condition-specific changes and the modulation of

connectivity parameters by experiment relevant variables (e.g., SSRT), we tested

group-level effects using the Parametric Empirical Bayes (PEB) framework. We give a

brief overview of PEB (for in-depth discussion, see Friston et al., 2016) for hypothesis

testing and connectivity parameter extraction using a PEB. Building a PEB statistical

model involves creating a hierarchical model with, in our case, two levels. The lower

level is the subject level, which is the results of DCM fits to individual's ERP data. This

first level includes the posterior means and uncertainties for each subject's DCM

connectivity parameters. The PEB framework statistically models these parameters

using a Bayesian general linear model (GLM) at the group level. As is the case with

GLMs (and mixed models), the model attempts to explains the connectivity parameters

as between-subject and within-subject variability, while allowing for between-subject

differences in connectivity parameters to be treated as random effects. The PEB

Bayesian GLM allows using subject-based DCM parameters to be examined using a

linear model with respect to explanatory variables at the between-subject (group) level.

Because the PEB framework yields group-level estimates as empirical priors, this approach also allows changing parameters that were estimated at the subject level by allowing them to be estimates distributed around a group mean effect. This process stands in contrast to the typically used "summary statistic" approach, which often involves applying several t-tests or correlations per connection.

For clarity, we built two separate PEB models for the DCMs fit to (1) No-TFUS US and SS trials, and (2) No-TFUS and Stop-TFUS SS trials. The former PEB model was used to test hypotheses regarding mean changes in connectivity between unsuccessful and successful inhibition at baseline (No-TFUS). The latter PEB model was used to address (1) how the inhibition network connectivity changed on average between TFUS conditions (intercept in GLM), and (2) how individual differences in the change in SSRT between TFUS conditions was embedded in changes in connectivity parameters across subjects (covariate). Therefore, in the first model we only used the mean intercept as the explanatory variable to examine mean changes in connectivity between No-TFUS US and SS trials. The mean change parameters represent the gain change in connectivity going from US to SS trials. In the second model, which examined No-TFUS SS and Stop-TFUS SS trials, we used a design matrix of explanatory variables that included an (1) intercept representing mean changes in connectivity (mean TFUS effect), (2) the change in individual subject's SSRT between TFUS conditions, and (3) the TFUS change in individual subject SSRT variability. Before entering the SSRT covariates, they were transformed to a gain change by taking the log-ratio of Stop-TFUS SSRT (mean or variability) over the No-TFUS SSRT (mean or

variability). The covariates were then z-scored to have a zero mean and yield

standardized PEB model parameters.

When initially estimating each of the two PEBs, we always entered in the full

model including all extrinsic and intrinsic connection modulatory parameters. Because

our hypotheses centered around changes along prefrontal and deep areas, and ventral

pathway interactions, we first compared the PEB model with and without dorsal pathway

connection parameters. We show in the results that, across both PEBs, the model

without dorsal pathway connection changes yielded a better model with respect to the

explanatory variables. This allowed a substantial reduction in connectivity parameters

needing to be tested.

Once the group-level GLM parameters were estimated with respect to

modulations of extrinsic and intrinsic connectivity, we used this framework to test

several hypotheses regarding mean and SSRT driven changes in connectivity

parameters. Hypothesis testing proceeds by Bayesian model reduction of the GLM. This

involves turning off/on different connectivity parameters and comparing the free energy

of reduced models. Comparing models in this manner is similar to performing classical

hypothesis testing via model reduction in mixed models by employing likelihood ratio or

F-tests.

Hypotheses were tested by designing models with different combinations of

parameters on/off. The model space of these hypotheses was defined in a factorial

space that focused on 4 factors that could be modulated by mean connectivity changes

(PEBs 1 and 2) or connectivity modulation via TFUS induced changes in SSRT (PEB 2).

These factor/hypothesis spaces were driven by previous work. The first set of models

considered how inhibition is related to pathways the backward connections from rIFG

and pre-SMA to the deep (i.e., basal ganglia) nodes. The second factor, driven by work

indicating that pre-SMA interacts with rIFG before deep projections (Rae et al., 2009),

tested for modulation of their backwards and forwards connections. The third factor was

motivated by proposals that differences in inhibition (SS vs. US) might be mediated

changes in attentional orienting, which predict changes in intrinsic self-inhibitory gain in

either rIFG, rIOG, or both. Therefore, this factor examined for modulation of the

superficial pyramidal cell gain across the nodes. Finally, to examine how SS versus US

and different TFUS effects on inhibition depend on top-down vs bottom-up processing,

our final comparison tested for the inclusion of either backward, forward or both sets of

connections along the ventral pathway. Given that each of 4 factors had 3 levels each,

plus a null (all zero) model, the first PEB surmounted to testing 3^4+1 (82) models, and

PEB model 2 included $2*3^4+1$ (163) models.

Rather than summarize these effects as the free energy for each model, which

would surmount to a severe reduction in discerning the probability of a winning model,

the hypotheses were grouped into families, and model hypotheses were tested at the

family level. After using family model comparison on reduced GLMs, we used Bayesian

model averaging (BMA) to obtain GLM model estimates of connectivity parameters

having a posterior probability >95%. BMA was performed on all families within a factor,

weighting the summarized parameters by the probability of the family. For example,

BMA parameters for the second PEB mode that includes the mean SSRT as a

predictor, for example, should be interpreted as would a linear regression coefficient;

similarly, the mean term would represent the mean change in connectivity. These BMA

parameters were reported as the final changes in connectivity and should be considered

to have a 95% probability of being non-zero.

Before conducting our main analysis presented, and to reduce the model space,

we first considered whether the best PEB model would include ventral and dorsal

pathway projections after fitting to individual subjects. Specifically, we asked if

describing the baseline difference of no-TFUS SS and US inhibition involved modulation

of either or both the dorsal and ventral pathways. Using the family-based hypothesis

comparisons test described above for PEBs, we created factor spaces including the

dorsal, ventral, or both pathways. Both sets of pathways included top-down and bottom-

up connections that were grouped together. The dorsal pathway included the rParietal

and rDLPFC nodes and the ventral pathway included rIFG and rTemporal. Bayesian

model comparison (probability = 1) revealed strong evidence in favor of the model with

just ventral pathway connection modulations from US to SS trials. Based on this result,

we excluded dorsal pathway connections from the rest of our DCM analysis to reduce

the space of parameters.

Results

Human participants performed a Stop-Signal task with online, trial-by-trial TFUS.

Subjects were divided into groups based on receiving one of three stimulation type: (1)

active stimulation targeted to right pars opercularis, (2) an active stimulation control site

(ipsilateral somatosensory, S1) to account for non-site specific TFUS, and (3) sham

stimulation to account for TFUS auditory artifacts. Since TFUS has been demonstrated

to illicit immediate effects on ERPs (Lee et al., 2016), stimulation was applied online

(see Methods) either at the onset of the go or stop signal cue, during both Go and Stop

trials (Figure 1A). We hypothesized that if rIFG implemented motor inhibition, then

TFUS behavioral effects would be limited to alteration of stopping but not going. TFUS

in the pars opercularis group improved inhibition, while exerting no effects in control

groups. TFUS also altered inhibition related ERPs, which were quantified at electrode

and source-localized levels, while also assessing TFUS impact on effective network

connectivity assessed using DCM.

Figure 1 **A.** visual layout of Stop-Signal task and corresponding times when transcranial focused
ultrasound (*TFUS*) was delivered. **B.** The left plot shows the average neuronavigation location of TFUS
applied to all MRIs used in the rIFG group. The structural scans on the right show renderings of the
targeted TFUS point in pars opercularis in 4 subjects. **C.** Lateral maximum pressure profile obtained at
30-mm depth focus in both water and transcranial ultrasound simulation on a CT scan from one patient
(solid and dotted lines, respectively). Horizontal red and blue lines denote full-width half maximum of the
spatial profile of lateral pressure. **D.** Simulated transcranial pressure profile onto T1 MRI plot shown as a
color overlay.

Numerical simulation of TFUS to rIFG

To determine intensity and accuracy of TFUS after skull transmission in the rIFG

group, we used numerical simulations based on CT and MRI data from 3 preoperative

patients and validated the simulation results using a water tank test. At the focus,

modeling of transcranial simulations predicted an average maximum intensity of 2.8

W/cm^2. This is in the range of intensity of non-thermal neuromodulation (Legon et al.,

2014). Additionally, predicted shifts in peak pressure due to skull transmission was 1.25

mm laterally, relative to a water model simulation, and had a lateral full-width half

maximum of 5.1 mm (Fig. 1C). These simulations indicate a high spatial precision with

>95% of energy limited to pars opercularis (Fig. 1D).

Only TFUS to rIFG alters response inhibition

We first addressed how probability of failing to inhibit responses,

P(respond|signal) (Fig. 2A), changed across TFUS conditions within and across groups,

by fitting a 2-parameter logistic linear mixed-model to obtain a slope (**β)** of the response

inhibition curve across all subjects and TFUS conditions. Modeling indicated a good fit

of the logistic curve (mean R^2 = 84%). Analysis of **β** indicated only the rIFG group

exhibited a TFUS-altered P(respond|signal). Importantly, behavioral effects of TFUS

were not found for either control groups. Anova results indicated a significant Group x

TFUS interaction (F(2,50) = 3.8, p = 0.034, η_p^2 = 0.17), and an overall effect of TFUS

condition (F(2,50) = 11.74, p = 0.002, η_p^2 = 0.29). Follow-up one-way ANOVAs across

TFUS onsets (e.g., coincident with stop signal), but within-groups, showed only the rIFG

group exhibited differences across onsets. Follow-up t-tests in this group showed **β** for

Stop-TFUS was lower than No-TFUS and Go-TFUS conditions (both p < 0.01: mean **β**'s

indicating change in probability for approximately 25% change in normalized SSD: No-

TFUS = 0.35 (0.12), Stop-TFUS=0.27 (0.08), Go-TFUS=0.35 (0.11)). These results

indicate only the rIFG group was affected by TFUS, as predicted, and only during Stop-

TFUS.

Figure 2A shows improved inhibition performance during Stop-TFUS for the rIFG

group occurred at longer SSDs (65% and 95% SSD). A repeated-measures ANOVA on

P(respond|signal) for the rIFG group across all SSD levels and TFUS onsets revealed a

significant interaction (F(6,102) = 8.21, p < 0.0001,   = 0.33). Contrast t-tests between

Stop-TFUS and the average of No- and Go-TFUS across all SSDs indicated the

interaction resulted from a reduction in P(respond|signal) for Stop-TFUS in the highest

two SSDs (all p < 0.01; Bonferroni α = 0.0125). These results indicate Stop-TFUS

induced improvements of inhibition were more pronounced at later SSDs.

Figure 2. **A.** Probability of responding across stop signal delays in the rIFG TFUS group for all TFUS
onsets. **B.** Violin plots of rIFG group across-subject distribution of stop signal reaction times (SSRT).

Based on the prediction that rIFG implements an inhibitory process and our

finding that TFUS improved inhibition in this group, we hypothesized rIFG TFUS

changes to P(respond|signal) should result from a shortening of the stopping speed

(i.e., SSRT); notably, TFUS did not affect other behavioral variables (e.g., Go RTs, see

Supplementary Material). SSRT analysis in a mixed-design ANOVA indicated a

significant Group x TFUS interaction (F(4,100) = 10.2, p < 0.001, η_p^2 = 0.21). Follow-up

one-way ANOVAs within-groups indicated only rIFG group SSRTs (Fig. 2B) differed

between TFUS onsets (p < 0.05), with t-tests confirming that SSRTs were indeed

shortest and only altered during Stop-TFUS (Fig. 2B). This result, along with the above

P(respond|signal), indicate rIFG Stop-TFUS altered inhibition by shortening the stop

process (SSRT).

Neural components underlying inhibition

In the rest of the results we focus our analysis on the rIFG group because this

was the only group exhibiting behavioral effects of TFUS. Furthermore, we only analyze

No-TFUS and Stop-TFUS conditions because Go activity was subtracted from neural

data at Stop trials (see Methods).

Our first analysis examined sensor-level ERPs across three contrasts using

cluster-based permutation t-tests: (1) successful – unsuccessful stopping contrast over

both TFUS conditions, (2) successful (No-TFUS) – successful stopping (Stop-TFUS)

contrast, and (3) interaction comparing successful – unsuccessful stopping between the

No-TFUS and Stop-TFUS conditions. The cluster-based scalp maps show the

progression of clusters time-locked to the stop-signal onset (0 ms, Fig. 3A,B).

Figure 3. **A.** Scalp plots of cluster-corrected permutation paired t-tests (*p* < 0.01). Colored contours represent significant clusters. (1). Contrast of all SS and US trials. (2). Contrast of TFUS conditions SS. (3). Interaction contrast calculated as SS-US of No-TFUS trials minus SS-US Stop-TFUS trials. **B.** Average ERP time courses of three clusters identified by the permutation testing that were differentiated by statistical contrasts. From left to right, the first cluster (left column) is right-parietal electrodes (CP6, CP4, P6, P8), the second cluster (middle column) is fronto-central electrodes (C1, Cz, C2), and the third cluster (right column) is right-frontal electrodes (F8, F6, F4). The vertical dashed lines represent the stop-signal reaction times for the No-TFUS (magenta) and Stop-TFUS (maroon). Latencies in A and B are expressed relative to stop signal onset (0 ms).

We identified several ERPs that differentiated successful stopping (SS) and

unsuccessful stopping (US) trials. The first contrast of SS-US (Figure 3A, row 1)

indicated SS trials exhibited a larger ERP centralized over a right-parietal cluster around

the time-range typically found for the P100 peak (100-148 ms). The interaction of SS-

US and TFUS showed this effect occurred earlier and during the peak onset (80-100

ms) of Stop-TFUS trials. Consideration of the SS-SS contrast (TFUS effect; Figure 3A-

2nd row) indicated the largest effect was attributable to the SS TFUS trials, supporting

previous conjectures that inhibition and stopping speed is directly related to larger P100

parietal responses. This result also aligns with previous studies reporting an enhanced

P100 during successful stopping (Boehler et al., 2009).

We also examined the N100 in the right-frontal cluster in the 80-120 ms window

(Figure 3B). We did not find a difference in the frontal N100 with respect to the overall

SS – US contrast (Figure 3A). This result is in line with several other studies noting a

non-significant effect of this contrast (Kenemans, 2015). However, we did find the ERP

was substantially larger in SS TFUS trials compared to SS No-TFUS trials indicating a

direct contribution to stopping efficacy. This increase for SS TFUS suggests this ERP

may stem from rIFG and provide an index of stopping speed, rather than success.

The ERPs most commonly associated with response is the N200/P300 complex.

Notably, the N200 often appears in both right-frontal and fronto-central clusters, while

the P300 is more aligned with the fronto-central (Huster et al., 2013; Kenemans, 2015).

When examining the N200, which typically only appears during US (Liotti et al., 2010;

Wessel et al., 2015), we found an ERP peaking around 200 ms in both clusters that only

appeared in US trials (Figure 3B). This N200 emerged during both No- and Stop-TFUS

(Figure 3, bottom right), with a larger amplitude during US Stop-TFUS.

Of all possible ERPs, the fronto-central P300 has been regarded as the most

robust marker of response inhibition and stopping speed (Wessel and Aron, 2015).

Accordingly, we found P300 amplitude differed between SS and US trials, with US trials

exhibiting a larger amplitude around the peak (290-320 ms) (Figs. 3A-B). However,

because P300 peaks occur after the SSRT, this implies it is too temporally protracted to

reflect inhibition (Huster et al., 2013). Alternatively, others have indicated the P300

onset latency is related to inhibition because it scales with individuals SSRTs (Wessel

and Aron, 2015). Based on this notion, we considered whether the P300 onset was

causally related to the SSRT shortening induced by TFUS. We tested the specific

prediction that Stop-TFUS SS trials should exhibit an earlier onset that correlates with

individual changes in SSRT. Visually, contrasting waveforms of SS trials across TFUS

conditions (Fig. 3B, upper-middle) supports this intuition that P300 onset (zero-crossing

in Figure 3B) shifted earlier in alignment with Stop-TFUS induced SSRT shifts. This was

quantitatively supported by a significant, across-subject correlation between the TFUS

induced in change P300 onset and SSRT (0.61, $p < 0.05$), providing direct support that

P300 latencies reflect the timing of inhibition speed.

Whole-brain source SPM and source-based regression analysis

We also examined source-based activity to localize TFUS-induced changes in

evoked activity and generating source location priors for DCM. We hypothesized that, if

differential rIFG activation indexed SS versus US, then conjunction analysis should

reveal an overlap between SS – US and No-TFUS – Stop-TFUS conditions if rIFG

activity is related to successful stopping (Aron et al., 2014). As expected, whole-brain

SPMs (Fig. 4A) revealed the only area exhibiting an overlap was a pars opercularis-

centered rIFG peak during the 20-100 ms time window (results of analysis of other time

windows are reported in Supplementary Material).

To understand how TFUS altered stopping efficacy, we compared changes in

SSRT between Stop-TFUS and No-TFUS SS trials using whole-brain SPM linear

regression. In the regression, positive contrasts indicate ΔSSRT (No-TFUS – TFUS) is

associated with larger activity in No-TFUS trials. Negative contrasts indicate TFUS SS

trials exhibit larger evoked activity predicted by larger changes in SSRT. The negative

contrast in Figure 4B indicates that the first site to exhibit increased TFUS-related

predictions is rIFG. This occurred both in the -40:20 and 20:100 time-windows. The

positive contrast found in early time windows indicates Stop-TFUS exerted effects on

stopping by also decreasing early activity in both bilateral parietal and right temporal

sites. We also found pre-SMA was only predictive of SSRT after rIFG, with the pre-SMA

modulation peaking at 100:180 ms. These results show that areas typically associated

with successful inhibition were predictive of TFUS-induced changes in behavior while

occurring before the SSRT itself.

FWE P	Cluster Extent	Uncorr. P	Peak-F	X	Y	Z	Cluster Peak
F-contrast: Successful Stop – Unsuccessful Stop (SS-US)							
<0.005	893	0.0001	16.35	50	-40	-26	R Inferior Temporal
<0.05	586	0.0001	15.16	54	26	18	R Inferior Frontal
F-contrast: non-tFUS – stop-tFUS (tFUS)							
<0.01	620	0.0001	16.65	-42	-72	-8	L Inferior Occipital
<0.005	899	0.0001	16.09	52	18	22	R Inferior Frontal:
<0.01	699	0.0001	15.52	48	-74	-10	R Inferior Occipital
Conjunction of F-contrasts							
<0.05	580	0.005	14.13	49	18	20	R Inferior Frontal:

Figure 4. **A.** Whole-brain SPM F-contrasts of evoked activity in the 20:100 ms window. Figure shows the surface mesh projections of the overall SS – US (left) TFUS contrast (middle), and conjunction (right). Below the surface meshes is a table listing the statistics. **B.** Whole-brain SPM linear regression of activity No-TFUS and Stop-TFUS SS trials against SSRT changes in No-TFUS and Stop-TFUS SS trials.

TFUS effects in an inhibition network: Dynamic Causal Modeling (DCM)

The above TFUS results provide evidence that rIFG activity is causally related to inhibition. Principally, differences in local activation can result from both local and inter-areal connectivity (David et al., 2006). We used DCM to quantify network effects. Bayesian model selection established a winning model as a hierarchical network in

which rDLPFC and rIFG sat at the top of the hierarchy and pre-SMA being below these

areas. The model also included nodes for right temporal (rTemporal), right inferior

occipital gyrus (rIOG), right parietal (rParietal), a hidden subcortical node (Deep) and

left motor cortex (M1). These results accord with networks proposed by previous studies

implicating both motor inhibition and attentional orienting (Weicki and Frank, 2013;

Munkata et al., 2011). The DCM (Fig. 5) was fit to individual subject's data to determine

how connectivity parameters differentiated (1) No-TFUS baseline SS and US inhibition

differences, and (2) between No-TFUS and Stop-TFUS SS trials and accompanying

changes in SSRT. The above approach revealed the models agreeably captured the

spatiotemporal properties of the ERP scalp data across both sets of model fits (Fig. 5).

Figure 5. **A**. Connections used to implement the final dynamic causal model structure. Exogenous inputs
entered through rIFG and rIOG represented as a green box. All nodes except Left M1 and Thalamus

were all located in the right hemisphere. **B**. Left panel shows the mean observed and predicted scalp
ERP data derived from the dynamic causal model fit to the No-TFUS US and SS trials. Right panel shows
the mean observed and predicted scalp ERP data derived from the dynamic causal model fit to the No-
TFUS SS and Stop-TFUS SS trials. These results show that the final model provided a good fit to the
data. Data are plotted relative to stop signal onset (0 ms on x-axis).

The first set of analyses examined what connectivity parameters were altered

during No-TFUS successful inhibition. The first comparison examined whether

differential rIFG and pre-SMA interactions were related to baseline successful stopping,

as suggested by several functional and anatomical studies (Duann et al., 2009; Rae et

al., 2015; Swaan et al., 2012; Fig. 6A). Model comparisons revealed a winning family

included interactions between both areas, but with a moderate posterior probability

(0.78). BMA across families revealed both the connection from rIFG to pre-SMA (136%)

and pre-SMA to rIFG (193%) were altered during SS trials. Changes in this connection

suggest that SS trials were supported by prefrontal interactions. Considering previous

data indicating increases in pre-SMA projections (Forstmann et al., 2008) to striatum

render increased RTs, this result might reflect an effect of blocking the impetus pre-

SMA provides towards responding to the Go signal (Verbruggen and Logan, 2009).

The next comparison tested the hypothesis that both pre-SMA and rIFG

projections to deep areas are necessary for successful inhibition (Fig. 6B). Only a family

including changes from rIFG to deep was predictive of successful inhibition. BMA

indicated rIFG exhibited a mean reduction in the backwards connection of 65% during

successful stopping. This result agrees with previous fMRI studies (Aron et al., 2006;

Jahfari et al., 2011) indicating inhibitory processes are driven by cortical to basal ganglia

interactions.

Figure 6. Family-based model comparison for different hypothesized interactions. Plots contain modulatory parameter of connections with strong positive evidence as being different between US and SS trials. Parameter estimates (>95% posterior probability) are in parentheses next to modulated connection in exponential percentage change. Anything above 100% reflects an increase in SS trials compared to US (opposite for below 100%). **A.** Test of rIFG and pre-Sma interactions. **B.** Tests of rIFG and pre-SMA backward projections to deep node. **C.** Tests for the ventral pathway connections. **D.** Comparison of intrinsic superficial pyramidal cell gains.

Another hypothesis that has been put forth regarding successful inhibition is that

it is mainly mediated by attentional orienting in ventral pathways (Hampshire and Sharp,

2015). DCM implementations of attentional process can be cast in terms of hierarchical

predictive coding models (Feldman and Friston, 2010). Previous work suggests

increased attention in sensory areas emerges as increased recurrent intrinsic gain

(increased disinhibition) of superficial pyramidal cells thought to report prediction errors

through forward connections (Auksztulewicz et al., 2015). In DCM, recurrent gains

would weight prediction error signals driven changes in top-down (backward)

connections (Bastos et al, 2012). We designed the next two comparisons to examine

whether SS trials exhibited these network changes. Extrinsic connection analysis in No-

TFUS SS indicated a winning family including only changes in top-down backward

connections predicted successful inhibition (Pp =1). BMA indicated, however, the

inhibitory connection from rIFG to rTemporal was reduced while backward connections

from rTemporal to rIOG increased in connectivity (Fig. 6C).

926   The final SS and US modulation comparison analyzed changes in recurrent

gains of rIFG, rTemporal and rIOG (Fig. 6D). The winning model (Pp = 0.99) included

an increase in gain for both rIOG (176%) and rIFG (153%). Importantly, the increased

rIOG gain is predicted by attentional orienting models of response inhibition (Hampshire

and Sharp, 2015) and accords with previous DCM studies that have manipulated

attentional cueing (Auksztulewicz et al., 2015). Mechanistically, the increased rIOG gain

in SS trials results in ascending signal that has a larger effect on decreasing rIFG top-

down expectation signals in backward connections.

Stopping efficiency (SSRT) is driven by lower and prefrontal interactions

935   Another primary goal of understanding inhibitory control is quantifying how the

efficiency of stopping, evaluated through the SSRT, is implemented via network

pathways. Previous work has employed between-subjects' correlations of SSRT and

connectivity parameters to isolate the pathways involved in this process (e.g., Jahfari et

al., 2011). Because previous studies using fMRI and a variety of connectivity methods

revealed pre-SMA and rIFG to basal ganglia connections are correlated in different

directions with SSRT (Jahfari et al., 2011), we adopted this approach when analyzing

the baseline No-TFUS SS versus US contrast. This correlational analysis indicated both

connections were correlated in opposite directions with SSRT. Figure 7 shows that

increasing connectivity from rIFG to deep targets predicted shorter SSRTs. The

opposite pattern was found for pre-SMA, wherein increasing pre-SMA backward activity

predicted longer SSRTs.

Figure 7. Left and right plots show linear regression fits predicting individual subjects' No-TFUS
SSRT from the backward connection to the deep area projecting from rIFG and pre-SMA,
respectively. Larger values on the x-axis denote decreasing backward inhibitory connectivity.

952 TFUS to rIFG causally dissociates inhibitory mechanisms

Building on TFUS's effect of increasing inhibition performance, a primary

question was whether the changes in connectivity strength between No-TFUS SS and

Stop-TFUS SS trials would reflect stepwise changes in the connections modulated in

the SS and US No-TFUS comparison. This comparison is in line with the idea of failed

stop trials resulting partially from less active inhibition mechanisms. By applying TFUS

to rIFG, we were able to causally dissociate rIFG's role in implementing inhibitory

mechanisms. Below, mean changes represent the average modulation of connectivity

going from No-TFUS SS to Stop-TFUS SS trials. SSRT-based effects represent how

changes in SSRT between TFUS conditions predict the change in connectivity between

TFUS conditions.

In first analyzing the family of models examining the rIFG and pre-SMA

interaction, there was no clear winning family of models for both the mean and SSRT

effect (Fig. 8A). For the mean, the model with projections from rIFG to pre-SMA (122%)

and pre-SMA to rIFG (75%) was highest in probability (Pp = 0.79). Of these

connections, SSRT change was only negatively related to the rIFG to pre-SMA pathway

(r^2 = 74%). indicating that an increase in inhibitory connectivity from rIFG to pre-SMA

during Stop-TFUS trials predicted a larger change in SSRT. Therefore, changes in

SSRT were not related to forward connections between these nodes, but instead were

modulated by a top-down inhibitory modulation from rIFG to pre-SMA.

When comparing models testing the role of rIFG or pre-SMA to deep backward

connections, family analysis revealed a strong effect of the mean change and SSRT

change for both backward connections (Pp > 0.95 both effects). BMA in the winning

family indicated both effects were above threshold for both connections. Both rIFG

(143%) and pre-SMA (139%) exhibited increased connectivity during TFUS. Concerning

changes in TFUS related changes in SSRT, the direction of effects was opposite for

rIFG (154%) and pre-SMA (67%): for rIFG, decreased inhibition yielded increased

SSRT change, and the opposite for pre-SMA. Together, these results point to

differential interactions of both pre-frontal areas with a deep node in responding to

TFUS. These results are further in line with the directionally opposite linear correlations

of the No-TFUS SSRT to each of these backward connections (Fig. 8B).

Finally, we examined ventral pathway parameters for TFUS modulation of top-

down, bottom-up, and intrinsic gain changes. Model averaging revealed the rIFG to

rTemporal top-down connections (66%) decreased with TFUS, suggesting a direct

effect from rIFG TFUS. This decrease in top-down rIFG connection agrees as well with

the results found during the SS versus US comparison, suggesting a causal and

overlapping pathway driving inhibition performance. All bottom-up connections

increased as well (Fig. 8C). With respect to SSRT change (Figure 8D), only the bottom-

up connections were related to changes in the SSRT. However, the forward connectivity

relations to SSRT were in opposite directions for modulations of the rIOG to rTemporal

(110%) and rTemporal to rIFG (78%). Finally, we found rTemporal recurrent gain

increased in mean (150%) and was positively related to changes in SSRT (163%). This

relation between the SSRT effect and rTemporal to rIFG forward connection was strong

enough such that a leave-one-out cross validation prediction of SSRT change using this

connection exhibited a large correlation with actual SSRT change (0.84, $p < 0.01$).

These results indicate neural implementation of stopping speed involves processing

efficacy of bottom-up, temporal cortex (prediction error) signals.

Figure 8. Results of the family-based model comparison for different hypotheses tested when comparing modulation from No-TFUS SS to Stop-TFUS SS trials. The family-based posterior probability (F-Pp) for the winning model is listed below each model. The outcomes of these plots can be interpreted of as revealing the modulatory parameters connections with very strong, positive evidence of US and SS trials being different. Parameters estimates with a greater than 95% posterior probability in these families are presented in parentheses next to the modulated connection. The parameters are presented in exponential form of percentage change. Values above 100% equates to a parameter increase in SS trials compared to US (and the opposite for values below 100%). Parameters in-active in each model are in a gray color. **A**. hypothesis test of rIFG and pre-Sma interactions. **B.** Hypothesis tests of rIFG and pre-SMA backward projections to deep node. **C.** comparison of intrinsic superficial pyramidal cell gains. **D.** tests for the ventral pathway connections.

Discussion

The present study examined whether pars opercularis sector of rIFG is explicitly

involved in motor response inhibition. We used a stop-signal task, online TFUS, source-

localized EEG, and dynamic causal modeling of ERPs to examine this hypothesis and

examine underlying response inhibition network mechanisms. Behaviorally, TFUS

applied to pars opercularis and coincident with the stop signal increased the likelihood

of successful inhibition. Because TFUS enhancements of inhibition resulted from faster

stopping processes (i.e., SSRT), without altering Go RTs, this supports pars opercularis'

role as directly triggering action stopping. Examination of scalp ERP analysis indicated

TFUS rendered a shift in the fronto-central P300 onset, which has been previously

hypothesized as neural marker of stopping speed (Wessel and Aron, 2015).

Imperatively, the shifted onsets were directly linked to and correlated with TFUS-

induced changes in individuals SSRT. Spatial accuracy of TFUS was supported by

whole-brain evoked activity indicating only pars opercularis exhibited a conjunction

effect of TFUS with successful compared to failed inhibition (Figure 3). The hypothesis

that pars opercularis activity is directly related to stopping efficacy was confirmed by

TFUS-driven activity differences predicting between-subjects SSRT change (Figure 4B).

Despite these results, recent work suggested TFUS effects result from auditory artifacts

(Sato et al., 2018). Auditory effects, however, cannot explain our findings because

control groups exhibited no behavioral effects, and no group exhibited evoked auditory

cortex activity (Figure S7). We interpret the above results as indicating TFUS selectively

modulated pars opercularis activity, and pars opercularis explicitly implements response

inhibition.

Generally, response inhibition involves several processes embedded in

connections across a neural network, ranging from sensory cue detection, attention,

performance monitoring, and presumably explicit motor inhibition (Munkata et al., 2011;

Weicki and Frank, 2013; Wessel and Aron, 2017). Many have proposed that motor

inhibition is implemented directly in rIFG connectivity to basal ganglia (STN and

striatum), either in parallel or routed through pre-SMA (Aron et al., 2014), or both. In

contrast, others have proposed motor inhibition is better understood as an emergent

outcome of attentional orienting and biased competition between neural processing of

response and inhibition cues (Hampshire and Sharp, 2015). Extant evidence indicates,

however, that motor inhibition and attentional processing are likely all involved in

different processing stages of inhibitory control tasks (Wessel and Aron, 2017).

Given the evidence for response inhibition as a multi-process phenomenon, a

key question is what network connections and biophysical mechanisms support action

stopping. A core component of models positing a direct rIFG motor inhibitory process is

that its connectivity with subcortical nodes should change with stopping success or

efficacy. The present study's DCM analysis of No-TFUS successful and failed inhibition

is consistent with the hypothesis that rIFG and pre-SMA subcortical connections are

relevant for motor inhibition. DCM results of No-TFUS successful stopping also revealed

an anti-correlation of SSRT to pre-SMA and rIFG deep connection strength (Figure 7).

Our findings agree with another fMRI connectivity study (Jahfari et al., 2011) that found

successful stopping accompanied increasing pre-SMA to striatum connectivity and

predicted longer SSRTs, with the opposite correlation for rIFG to striatum. Pivotal

support for rIFG's role is the result that TFUS effects on SSRT were directly related to

changes in rIFG and pre-SMA to deep connectivity with a similar anti-correlation pattern

as the baseline. We take these results to indicate pars opercularis subcortical

connectivity is directly involved in driving motor inhibition through feedback pathways.

This indicate a potential mechanistic effect of TFUS, wherein inhibition improved by

altering the connectivity of layer V rIFG pyramidal neurons by increasing the excitability

of these cells. Notably, this conjecture of an excitatory TFUS effect on pyramidal

neurons and our TFUS parameterization are in accordance with the neuronal

intramembrane cavitation excitation (NICE) model that has been used recently to

explain the acoustical stimulation effects on the biophysics of neuronal firing (see

Plaksin et al., 2016).

An unresolved issue regarding the inhibition network is how rIFG and pre-SMA

interactions areas relate to inhibition in general (Duann et al., 2009; Swaan et al., 2012).

Our DCM optimization indicated rIFG projected to pre-SMA through backwards

inhibitory connections and pre-SMA to rIFG via excitatory forward connections. This

bidirectional connectivity agrees with previous fMRI stop-signal and DTI studies (Duann

et al., 2009; Rae et al., 2015). Functionally, No-TFUS DCM results revealed successful

stopping was accompanied by increases in both connections. However, when

comparing baseline and TFUS DCMs, our results indicate that only the rIFG to pre-SMA

connections in this subset of connections were effectively related to inhibition in terms of

the mean gain change in backwards connectivity and its covariation with TFUS induced

changes in SSRT (Figure 6A-B). This result is directly relevant to several studies that

have either concluded pre-SMA drives rIFG (Swann et al., 2011) during inhibition or the

opposite (Duann et al., 2009). Combining DCM and TFUS indicated the inhibitory effect

of rIFG onto pre-SMA is causally responsible for driving inhibition at a cortical level.

This raises the question of why the pre-SMA to rIFG connection was only relevant

during baseline successful inhibition. An alternative interpretation is found in

neuroimaging (Crone et al., 2006), ECoG (Swaan et al., 2012), and TMS studies of

proactive inhibition and response switching (Rushworth et al., 2002). These studies

have indicated pre-SMA encodes a set of potential actions. During response inhibition,

this predicts successful stopping involves pre-SMA signaling the action(stopping)-rule to

rIFG. Therefore, the action rule and connectivity conveying it should not differ for

successful No-TFUS and Stop-TFUS trials because it should have been similarly

communicated under both conditions.

The above results provide causal evidence that pars opercularis and its

connectivity are directly involved in motor inhibition. Nonetheless, neuroimaging studies

have shown rIFG-related activation predicts attentional orienting and stimulus

expectancy processes during response inhibition (Erika-Florence et al., 2014;

Hampshire and Sharp, 2015; Xu et al., 2017). Although attention was not manipulated in

our study, we derived predicted network mechanisms from predictive coding models of

attention (Feldman and Friston, 2010) and DCM-EEG studies directly manipulating

attention and stimulus expectancy (Auksztulewicz et al., 2015). Importantly, the

microcircuits implemented in the DCM presently used are directly related to predictive

coding models and have explicit mechanisms supported by previous DCM studies. For

example, these studies have shown increased attention is linked to increased recurrent

gain on ascending (forward) sensory (prediction error) signals, while violations of

stimulus expectations were linked to decreased top-down and increased in bottom-up

connectivity, respectively (Auksztulewicz et al., 2015; Fardo et al., 2017). Along these

lines, our DCM during baseline stopping featured increased rIOG recurrent gain as

expected if attention increased the precision afforded to the sensory processing of stop

cues (Moran et al., 2013; Figure 7C). Concerning top-down changes, rIFG to rTemporal

decreased and rTemporal to rIOG increased (Figure 7D). Successful stopping at

baseline therefore may rely on an increased rIOG gain weighted sensory signals that

drive top-down changes in rIFG. However, increased connectivity from rTemporal to

rIOG seems at odds with this interpretation because it implies a larger reliance on top-

down information during successful stopping. This is explained by the finding that

optimized inputs to rIOG were negative and therefore inhibitory (rather than excitatory)

across subjects. Therefore, increasing rTemporal backward connectivity rendered an

enhanced negative rIOG signals.

Although the preceding results seem to accord with attention-based formulations

of inhibition, examination of TFUS effects on connectivity indicate inhibition success

was generally unrelated to these mechanisms. The only overlap between baseline and

TFUS conditions was a decrease in top-down rIFG to rTemporal backwards

connectivity. Still, this effect was not directly predictive of a change in SSRT. TFUS

effects also involved an increase in bottom-up connectivity that was predictive of SSRT

change (Figure 8D) in agreement with expectation violation effects found in other EEG-

DCM studies (Auksztulewicz et al., 2015). The most interesting effect was recurrent

gains only increased in rTemporal, indicating SSRT changes were not due to increased

sensory weighting. We propose that cortically-related SSRT changes were driven by

TFUS altering the effects the rIFG to rTemporal connection had on the rTemporal gain.

A partial correlation supported this hypothesis by showing covariation of rTemporal gain

and SSRT change was rendered null when accounting for the correlation of rIFG to

rTemporal and rTemporal gain. A potential explanation for this result is that rIFG

engages in a proactive inhibitory function whereby it biases bottom-up processing of the

temporal cortex, which itself may encode the expected probability of stop signal

occurring. This interpretation is consistent with fMRI stop-signal studies demonstrating

temporal cortex encodes a prediction error of stop-signal probability (Hu et al., 2015; Ide

et al., 2013).

In summary, TFUS can induced enhanced response inhibition performance,

allowing the underlying mechanisms to be linked to direct and source-resolved

electrophysiological neural processes in humans. By pairing TFUS with DCM, we found

a network model of response inhibition suggesting pars opercularis explicitly invokes

motor inhibition through deep pyramidal connections directly synapsing onto subcortical

nodes, as well as pre-SMA and temporal cortex. These results also significantly extend

the possible applications by showing TFUS combined with network modeling has the

potential to alter and infer the effective connection between biophysical network

mechanisms and behavior.

1145

1146

1147

1148

1149

1150

1151

1152

1153

1154

1155

1156

1157

References

Aron, A. R., Fletcher, P. C., Bullmore, E. T., Sahakian, B. J., & Robbins, T. W. (2003). Stop-signal inhibition disrupted by damage to right inferior frontal gyrus in humans. *Nature Neuroscience*. https://doi.org/10.1038/nn1203-1329a

Aron, A. R., Robbins, T. W., & Poldrack, R. A. (2014). Inhibition and the right inferior frontal cortex: one decade on. https://doi.org/10.1016/j.tics.2013.12.003

Aron, A. R. (2006). Cortical and Subcortical Contributions to Stop Signal Response Inhibition: Role of the Subthalamic Nucleus. *Journal of Neuroscience*. https://doi.org/10.1523/JNEUROSCI.4682-05.2006

Aron, A. R., Herz, D. M., Brown, P., Forstmann, B. U., & Zaghloul, K. (2016). Frontosubthalamic Circuits for Control of Action and Cognition. *The Journal of Neuroscience*. https://doi.org/10.1523/jneurosci.2348-16.2016

Aubry, J.-F., Tanter, M., Pernot, M., Thomas, J.-L., & Fink, M. (2003). Experimental demonstration of noninvasive transskull adaptive focusing based on prior computed tomography scans. *The Journal of the Acoustical Society of America*. https://doi.org/10.1121/1.1529663

Auksztulewicz, R., & Friston, K. (2015). Attentional enhancement of auditory mismatch responses: A DCM/MEG study. *Cerebral Cortex*. https://doi.org/10.1093/cercor/bhu323

Baddeley, A. (1996). Exploring the Central Executive. *Quarterly Journal of Experimental Psychology Section A: Human Experimental Psychology*. https://doi.org/10.1080/713755608

Bari, A., & Robbins, T. W. (2013). Inhibition and impulsivity: Behavioral and neural basis of response control. *Progress in Neurobiology*. https://doi.org/10.1016/j.pneurobio.2013.06.005

Bastos, A. M., Usrey, W. M., Adams, R. A., Mangun, G. R., Fries, P., & Friston, K. J. (2012). Canonical Microcircuits for Predictive Coding. *Neuron*. https://doi.org/10.1016/j.neuron.2012.10.038

Bates, D., Maechler, M., & Bolker, B. (2013). lme4: Linear mixed-effects models using S4 classes. *R Package Version 1.1-7*. https://doi.org/citeulike-article-id:1080437

Bhatt, M. B., Bowen, S., Rossiter, H. E., Dupont-Hadwen, J., Moran, R. J., Friston, K. J., & Ward, N. S. (2016). Computational modelling of movement-related beta-oscillatory dynamics in human motor cortex. *NeuroImage*. https://doi.org/10.1016/j.neuroimage.2016.02.078

Boehler, C. N., Münte, T. F., Krebs, R. M., Heinze, H. J., Schoenfeld, M. A., & Hopf, J. M. (2009). Sensory MEG responses predict successful and failed inhibition in a stop-signal task. *Cerebral Cortex*. https://doi.org/10.1093/cercor/bhn063

Boehler, C., Author, C., & Nicolas Boehler, C. (2010). Pinning down response inhibition in the
brain-conjunction analyses of the Stop-signal task4). *Neuroimage, 52*(1), 1621–1632.
https://doi.org/10.1016/j.neuroimage.2010.04.276

Cai, W., George, J. S., Verbruggen, F., Chambers, C. D., & Aron, A. R. (2012). The role of the
right presupplementary motor area in stopping action: two studies with event-related
transcranial magnetic stimulation. *Journal of Neurophysiology*.
https://doi.org/10.1152/jn.00132.2012

Chambers, C. D., Bellgrove, M. A., Stokes, M. G., Henderson, T. R., Garavan, H., Robertson, I.
H., … Mattingley, J. B. (2006). Executive "brake failure" following deactivation of human
frontal lobe. *Journal of Cognitive Neuroscience*.
https://doi.org/10.1162/089892906775990606

Chambers, C. D., Garavan, H., & Bellgrove, M. A. (2009). Insights into the neural basis of
response inhibition from cognitive and clinical neuroscience. *Neuroscience and
Biobehavioral Reviews*. https://doi.org/10.1016/j.neubiorev.2008.08.016

Chatham, C. H., Claus, E. D., Kim, A., Curran, T., Banich, M. T., & Munakata, Y. (2012).
Cognitive control reflects context monitoring, not motoric stopping, in response inhibition.
PLoS ONE. https://doi.org/10.1371/journal.pone.0031546

Chikazoe, J., Jimura, K., Asari, T., Yamashita, K. I., Morimoto, H., Hirose, S., … Konishi, S.
(2009). Functional dissociation in right inferior frontal cortex during performance of go/no-
go task. *Cerebral Cortex*. https://doi.org/10.1093/cercor/bhn065

Corbetta, M., & Shulman, G. L. (2002). Control of goal-directed and stimulus-driven attention in
the brain. *Nature Reviews. Neuroscience*. https://doi.org/10.1038/nrn755

Crone, E. A., Wendelken, C., Donohue, S. E., & Bunge, S. A. (2006). Neural evidence for
dissociable components of task-switching. *Cerebral Cortex*.
https://doi.org/10.1093/cercor/bhi127

David, O., Kiebel, S. J., Harrison, L. M., Mattout, J., Kilner, J. M., & Friston, K. J. (2006).
Dynamic causal modeling of evoked responses in EEG and MEG. *NeuroImage*.
https://doi.org/10.1016/j.neuroimage.2005.10.045

David, O., Maess, B., Eckstein, K., & Friederici, A. D. (2011). Dynamic Causal Modeling of
Subcortical Connectivity of Language. *Journal of Neuroscience*.
https://doi.org/10.1523/jneurosci.3433-10.2011

Delorme, A., & Makeig, S. (2004). EEGLAB: an open source toolbox for analysis of single-trial
EEG dynamics including independent component analysis. *Journal of Neuroscience
Methods*. https://doi.org/10.1016/j.jneumeth.2003.10.009

Duann, J.-R., Ide, J. S., Luo, X., & Li, C. R. (2009). Functional connectivity delineates distinct
roles of the inferior frontal cortex and presupplementary motor area in stop signal inhibition.
*The Journal of Neuroscience : The Official Journal of the Society for Neuroscience,
29*(32), 10171–10179. https://doi.org/10.1523/JNEUROSCI.1300-09.2009

Evans, A. C., Kamber, M., Collins, D. L., & MacDonald, D. (1994). An MRI-Based Probabilistic
Atlas of Neuroanatomy. In *Magnetic Resonance Scanning and Epilepsy*.
https://doi.org/10.1007/978-1-4615-2546-2_48

Erika-Florence, M., Leech, R., & Hampshire, A. (2014). A functional network perspective on
response inhibition and attentional control. *Nature Communications*.
https://doi.org/10.1038/ncomms5073

Fardo, F., Auksztulewicz, R., Allen, M., Dietz, M. J., Roepstorff, A., & Friston, K. J. (2017).
Expectation violation and attention to pain jointly modulate neural gain in somatosensory
cortex. *NeuroImage*. https://doi.org/10.1016/j.neuroimage.2017.03.041

Feldman, H., & Friston, K. J. (2010). Attention, Uncertainty, and Free-Energy. *Frontiers in
Human Neuroscience*. https://doi.org/10.3389/fnhum.2010.00215

Fini, M., & Tyler, W. J. (2017). Transcranial focused ultrasound: a new tool for non-invasive
neuromodulation. *International Review of Psychiatry*.
https://doi.org/10.1080/09540261.2017.1302924

Friston, K. J., Litvak, V., Oswal, A., Razi, A., Stephan, K. E., Van Wijk, B. C. M., … Zeidman, P.
(2016). Bayesian model reduction and empirical Bayes for group (DCM) studies.
NeuroImage. https://doi.org/10.1016/j.neuroimage.2015.11.015

Forstmann, B. U., Dutilh, G., Brown, S., Neumann, J., von Cramon, D. Y., Ridderinkhof, K. R., &
Wagenmakers, E.-J. (2008). Striatum and pre-SMA facilitate decision-making under time
pressure. *Proceedings of the National Academy of Sciences*.
https://doi.org/10.1073/pnas.0805903105

Hampshire, A., Chamberlain, S. R., Monti, M. M., Duncan, J., & Owen, A. M. (2010). The role of
the right inferior frontal gyrus: inhibition and attentional control. *NeuroImage*.
https://doi.org/10.1016/j.neuroimage.2009.12.109

Hampshire, A., & Sharp, D. J. (2015). Contrasting network and modular perspectives on
inhibitory control. *Trends in Cognitive Sciences*. https://doi.org/10.1016/j.tics.2015.06.006

Hayner, M., & Hynynen, K. (2002). Numerical analysis of ultrasonic transmission and absorption
of oblique plane waves through the human skull. *The Journal of the Acoustical Society of
America*. https://doi.org/10.1121/1.1410964

Haynes, W. I. A., & Haber, S. N. (2013). The Organization of Prefrontal-Subthalamic Inputs in
Primates Provides an Anatomical Substrate for Both Functional Specificity and Integration:
Implications for Basal Ganglia Models and Deep Brain Stimulation. *Journal of
Neuroscience*. https://doi.org/10.1523/jneurosci.4674-12.2013

Hu, S., Ide, J. S., Zhang, S., & Li, C. shan R. (2015). Anticipating conflict: Neural correlates of a
Bayesian belief and its motor consequence. *NeuroImage*.
https://doi.org/10.1016/j.neuroimage.2015.06.032

Huster, R. J., Enriquez-Geppert, S., Lavallee, C. F., Falkenstein, M., & Herrmann, C. S. (2013).
Electroencephalography of response inhibition tasks: Functional networks and cognitive
contributions. *International Journal of Psychophysiology*.
https://doi.org/10.1016/j.ijpsycho.2012.08.001

Ide, J. S., Shenoy, P., Yu, A. J., & Li, C. -s. R. (2013). Bayesian Prediction and Evaluation in the
Anterior Cingulate Cortex. *Journal of Neuroscience*. https://doi.org/10.1523/jneurosci.2201-
12.2013

Jahfari, S., Waldorp, L., van den Wildenberg, W. P. M., Scholte, H. S., Ridderinkhof, K. R., &
Forstmann, B. U. (2011). Effective Connectivity Reveals Important Roles for Both the
Hyperdirect (Fronto-Subthalamic) and the Indirect (Fronto-Striatal-Pallidal) Fronto-Basal
Ganglia Pathways during Response Inhibition. *Journal of Neuroscience*.
https://doi.org/10.1523/JNEUROSCI.5253-10.2011

Kenemans, J. L. (2015). Specific proactive and generic reactive inhibition. *Neuroscience and
Biobehavioral Reviews*. https://doi.org/10.1016/j.neubiorev.2015.06.011

Levy, B. J., & Wagner, A. D. (2011). Cognitive control and right ventrolateral prefrontal cortex:
Reflexive reorienting, motor inhibition, and action updating. *Annals of the New York
Academy of Sciences*. https://doi.org/10.1111/j.1749-6632.2011.05958.x

Lee, W., Kim, H. C., Jung, Y., Chung, Y. A., Song, I. U., Lee, J. H., & Yoo, S. S. (2016).
Transcranial focused ultrasound stimulation of human primary visual cortex. *Scientific
Reports*. https://doi.org/10.1038/srep34026

Legon, W., Sato, T. F., Opitz, A., Mueller, J., Barbour, A., Williams, A., & Tyler, W. J. (2014).
Transcranial focused ultrasound modulates the activity of primary somatosensory cortex in
humans. *Nature Neuroscience*. https://doi.org/10.1038/nn.3620

Legon, W., Bansal, P., Tyshynsky, R., Ai, L., & Mueller, J. K. (2018). Transcranial focused
ultrasound neuromodulation of the human primary motor cortex. *Scientific Reports*.
https://doi.org/10.1038/s41598-018-28320-1

Litvak, V., Garrido, M., Zeidman, P., & Friston, K. (2015). Empirical Bayes for Group (DCM)
Studies: A Reproducibility Study. *Frontiers in Human Neuroscience*.
https://doi.org/10.3389/fnhum.2015.00670

Logan, G. D., & Cowan, W. B. (1984). On the ability to inhibit thought and action: A theory of an
act of control. *Psychological Review*. https://doi.org/10.1037/0033-295X.91.3.295

Mallet, N., Schmidt, R., Leventhal, D., Chen, F., Amer, N., Boraud, T., & Berke, J. D. (2016).
Arkypallidal Cells Send a Stop Signal to Striatum. *Neuron*.
https://doi.org/10.1016/j.neuron.2015.12.017

Mathôt, S., Schreij, D., & Theeuwes, J. (2012). OpenSesame: An open-source, graphical
experiment builder for the social sciences. *Behavior Research Methods*.
https://doi.org/10.3758/s13428-011-0168-7

Mattia, M., Spadacenta, S., Pavone, L., Quarato, P., Esposito, V., Sparano, A., … Mirabella, G. (2012). Stop-event-related potentials from intracranial electrodes reveal a key role of premotor and motor cortices in stopping ongoing movements. *Frontiers in Neuroengineering*. https://doi.org/10.3389/fneng.2012.00012

Matzke, D., Love, J., Wiecki, T. V., Brown, S. D., Logan, G. D., & Wagenmakers, E. J. (2013). Release the BEESTS: Bayesian Estimation of Ex-Gaussian STop-Signal reaction time distributions. *Frontiers in Psychology*. https://doi.org/10.3389/fpsyg.2013.00918

Moran, R. J., Campo, P., Symmonds, M., Stephan, K. E., Dolan, R. J., & Friston, K. J. (2013). Free Energy, Precision and Learning: The Role of Cholinergic Neuromodulation. *Journal of Neuroscience*. https://doi.org/10.1523/JNEUROSCI.4255-12.2013

Morein-Zamir, S., Dodds, C., van Hartevelt, T. J., Schwarzkopf, W., Sahakian, B., Müller, U., & Robbins, T. (2014). Hypoactivation in right inferior frontal cortex is specifically associated with motor response inhibition in adult ADHD. *Human Brain Mapping*. https://doi.org/10.1002/hbm.22539

Munakata, Y., Herd, S. A., Chatham, C. H., Depue, B. E., Banich, M. T., & O'reilly, R. C. (2011). A unified framework for inhibitory control Opinion. *Trends in Cognitive Sciences*, *15*(10), 453–459. https://doi.org/10.1016/j.tics.2011.07.011

Obeso, I., Robles, N., Marrón, E. M., & Redolar-Ripoll, D. (2013). Dissociating the Role of the pre-SMA in Response Inhibition and Switching: A Combined Online and Offline TMS Approach. *Frontiers in Human Neuroscience*. https://doi.org/10.3389/fnhum.2013.00150

Opitz, A., Legon, W., Rowlands, A., Bickel, W. K., Paulus, W., & Tyler, W. J. (2013). Physiological observations validate finite element models for estimating subject-specific electric field distributions induced by transcranial magnetic stimulation of the human motor cortex. *NeuroImage*. https://doi.org/10.1016/j.neuroimage.2013.04.067

Penny, W. D., Stephan, K. E., Daunizeau, J., Rosa, M. J., Friston, K. J., Schofield, T. M., & Leff, A. P. (2010). Comparing families of dynamic causal models. *PLoS Computational Biology*. https://doi.org/10.1371/journal.pcbi.1000709

Plaksin, M., Kimmel, E., & Shoham, S. (2016). Cell-Type-Selective Effects of Intramembrane Cavitation as a Unifying Theoretical Framework for Ultrasonic Neuromodulation. *ENeuro*. https://doi.org/10.1523/eneuro.0136-15.2016

Rae, C. L., Hughes, L. E., Anderson, M. C., & Rowe, J. B. (2015). The prefrontal cortex achieves inhibitory control by facilitating subcortical motor pathway connectivity. *The Journal of Neuroscience : The Official Journal of the Society for Neuroscience*. https://doi.org/10.1523/JNEUROSCI.3093-13.2015

Ray Li, C. -s. (2006). Imaging Response Inhibition in a Stop-Signal Task: Neural Correlates Independent of Signal Monitoring and Post-Response Processing. *Journal of Neuroscience*. https://doi.org/10.1523/jneurosci.3741-05.2006

Sato, T., Shapiro, M. G., & Tsao, D. Y. (2018). Ultrasonic Neuromodulation Causes Widespread
Cortical Activation via an Indirect Auditory Mechanism. *Neuron*.
https://doi.org/10.1016/j.neuron.2018.05.009

Schmajuk, M., Liotti, M., Busse, L., & Woldorff, M. G. (2006). Electrophysiological activity
underlying inhibitory control processes in normal adults. *Neuropsychologia*.
https://doi.org/10.1016/j.neuropsychologia.2005.06.005

Sharp, D. J., Bonnelle, V., De Boissezon, X., Beckmann, C. F., James, S. G., Patel, M. C., &
Mehta, M. A. (2010). Distinct frontal systems for response inhibition, attentional capture,
and error processing. *Proceedings of the National Academy of Sciences*.
https://doi.org/10.1073/pnas.1000175107

Shulman, G. L., Astafiev, S. V., Franke, D., Pope, D. L. W., Snyder, A. Z., McAvoy, M. P., &
Corbetta, M. (2009). Interaction of Stimulus-Driven Reorienting and Expectation in Ventral
and Dorsal Frontoparietal and Basal Ganglia-Cortical Networks. *Journal of Neuroscience*.
https://doi.org/10.1523/jneurosci.5609-08.2009

Swann, N. C., Cai, W., Conner, C. R., Pieters, T. A., Claffey, M. P., George, J. S., … Tandon, N.
(2012). Roles for the pre-supplementary motor area and the right inferior frontal gyrus in
stopping action: Electrophysiological responses and functional and structural connectivity.
NeuroImage. https://doi.org/10.1016/j.neuroimage.2011.09.049

Tomaiuolo, F., MacDonald, J. D., Caramanos, Z., Posner, G., Chiavaras, M., Evans, A. C., &
Petrides, M. (1999). Morphology, morphometry and probability mapping of the pars
opercularis of the inferior frontal gyrus: An in vivo MRI analysis. *European Journal of
Neuroscience*. https://doi.org/10.1046/j.1460-9568.1999.00718.x

Treeby, B. E., & Cox, B. T. (2010). k-Wave: MATLAB toolbox for the simulation and
reconstruction of photoacoustic wave fields. *Journal of Biomedical Optics*.
https://doi.org/10.1117/1.3360308

Verbruggen, F., & Logan, G. D. (2009). Models of response inhibition in the stop-signal and
stop-change paradigms. *Neuroscience and Biobehavioral Reviews*.
https://doi.org/10.1016/j.neubiorev.2008.08.014

Verbruggen, F., Aron, A. R., Stevens, M. A., & Chambers, C. D. (2010). Theta burst stimulation
dissociates attention and action updating in human inferior frontal cortex. *Proceedings of
the National Academy of Sciences*. https://doi.org/10.1073/pnas.1001957107

Vossel, S., Thiel, C. M., & Fink, G. R. (2006). Cue validity modulates the neural correlates of
covert endogenous orienting of attention in parietal and frontal cortex. *NeuroImage*.
https://doi.org/10.1016/j.neuroimage.2006.05.019

Wessel, J. R., & Aron, A. R. (2015). It's not too late: The onset of the frontocentral P3 indexes
successful response inhibition in the stop-signal paradigm. *Psychophysiology*.
https://doi.org/10.1111/psyp.12374

Wessel, J. R., & Aron, A. R. (2017). On the Globality of Motor Suppression: Unexpected Events
and Their Influence on Behavior and Cognition. *Neuron*.
https://doi.org/10.1016/j.neuron.2016.12.013

White, P. J., Clement, G. T., & Hynynen, K. (2006). Local frequency dependence in transcranial
ultrasound transmission. In *AIP Conference Proceedings*.
https://doi.org/10.1063/1.2205477

Wiecki, T. V., & Frank, M. J. (2013). A computational model of inhibitory control in frontal cortex
and basal ganglia. *Psychological Review*. https://doi.org/10.1037/a0031542

Xu, K. Z., Anderson, B. A., Emeric, E. E., Sali, A. W., Stuphorn, V., Yantis, S., & Courtney
Correspondence, S. M. (2017). Neural Basis of Cognitive Control over Movement
Inhibition: Human fMRI and Primate Electrophysiology Evidence. *Neuron, 96*.
https://doi.org/10.1016/j.neuron.2017.11.010

Yamawaki, N., Borges, K., Suter, B. A., Harris, K. D., & Shepherd, G. M. G. (2014). A genuine
layer 4 in motor cortex with prototypical synaptic circuit connectivity. *ELife*.
https://doi.org/10.7554/eLife.05422

1394

1395

1396

1397

1398

1399

1400

1401

1402

1403

1404

1405

1406

1407

1408

Supplementary Materials

S1. Dynamic causal modeling structure optimization

Below we show the structures used for building and optimizing the dynamic causal

model based on the no-TFUS successful stopping event related potentials. The full steps of the

process are described in the Methods section. Figure S1 shows the tested structures and

possible connections. Figure S2 shows outputs for comparing these models using Bayesian

model comparisons, as well as model comparisons for which nodes receiving thalamic input,

and tests for the type of connection projecting from cortical to deep areas (forward, backward, or

both).

1418

1419 Figure S1. Differences in dynamic causal model spaces that were compared for the lower network
1420 structure and prefrontal network structure (top and bottom panel, respectively). The legend on the bottom
1421 shows how different connections are coded in these putative model spaces.

Model structure optimization on mean no-tFUS SS trials

1422

Figure S2. Model probabilities for the different model tests. Each of these tests was examined on the
mean ERP data for No-TFUS SS trials using family-wise fixed effect Bayesian model selection. Above
each plot shows the model component being optimized. For example, the first bar plot shows the different
prefrontal hierarchy arrangements. Above each plot is the difference in log-free energy between the best
family and the 2nd best. The log difference of free energies approximates a log-Bayes factor (Penny,
2012). It is considered positive evidence in favor of a model if this value is > 5. Evidence in favor of the
winning models (bars enclosed in red) was very strong. The plots on the right show the different structural
arrangements used to examine the prefrontal and lower-level hierarchical structure.

1431

1432

S2. Go Reaction Times

We addressed whether TFUS (real or sham) exerted any effects on simply responding to

the Go signal. We analyzed this by extracting the ex-gaussian-based mean RT using a

maximum likelihood approach (Lacoutoure and Cousineau, 2008). The means were calculated

separately for all subjects, and then analyzed using a 3 × 2 mixed-design ANOVA with factors of

the 3 groups, and TFUS state: No-TFUS and Go-TFUS trials. This analysis allowed us to

assess if "going", independent of "stopping", was altered by potential TFUS auditory artifacts,

stimulation to unrelated areas (S1 group), or whether TFUS to rIFG also influenced Go RT

processes independent of a stopping context signal. Table 1 shows the per group mean

difference of the RT between No-TFUS and Go-TFUS per group. Across groups, there was a

consistent negative difference indicating that RTs during TFUS may have been slightly shorter.

These differences in means seemed marginally larger for the rIFG group. However, the ANOVA

did not reveal any significant effect of TFUS, group, or their interaction (all $p > 0.05$). These

results suggest that neither TFUS (rIFG and S1 groups) nor auditory factors alone (sham rIFG

group) altered Go RTs independent of a stop signal.

S3. Signal-respond RTs and context independence

Calculating a measure of stopping latency (SSRT) based on the independent race-

model (Logan and Cowan, 1984; Bissett and Logan, 2014) assumes that signal-respond RTs

are Go processes resulting from a censoring of the Go RT distribution. Testing this assumption

predicts that (a) mean signal-respond RT should be faster than mean Go RT, and (b) during

fixed-SSD paradigms like the one employed here, signal-respond RT should increase with

longer SSDs because there are more failed inhibitory responses. Before testing (a) and (b), we

wanted to discern whether our TFUS manipulation exerted any effects on the signal-respond

RTs. Based on the context-independence assumption of the race model that signal-respond

RTs are Go processes escaping inhibition, and (2) if the TFUS-driven changes in inhibition

(P(respond|signal) (Fig. 3) are due to changes in inhibition, we should expect no differences in

signal-respond RTs between conditions or groups. We used a mixed-design ANOVA to examine

the subject level signal-respond RT means with Group (3 levels) and TFUS (3 levels: no-TFUS,

Go-TFUS, Stop-TFUS). We found no significant interactions or effects. Importantly, follow-up t-

tests revealed no differences across TFUS conditions within the rIFG group (mean signal-

respond RT: no-TFUS: 375 ±18 ms; Go-TFUS: 368 ±13 ms; Stop TFUS: 353 ±13 ms). This

supports the conclusion that TFUS did not alter the mean of Go processes that escaped

inhibition.

Given the lack of difference in mean signal-respond RTs (across all SSDs) across TFUS

conditions, we collapsed these RT means across TFUS conditions and compared them to each

subject's mean Go RT with a mixed-design ANOVA. There was no interaction or Group

differences ($p > 0.05$). As expected, the mean signal-respond RT (all groups: 365 ±15) was

significantly shorter (F(1,50) = 89.77, $p < 0.0001$) than the mean Go RT (all groups: 436 ±22).

Next, we examined whether the signal-respond RT increased with increasing SSD by

regressing all subjects' signal-respond RTs (collapsed across TFUS conditions) onto their SSDs

to obtain a single-slope parameter. This revealed a significant regression slope of 0.44 ($p <$

0.001), indicating signal-respond RTs did increase with increasing SSD.

Having confirmed that TFUS did not alter signal-respond RTs or the Go RTs, we sought

to test the context-independence assumption of the race model. This has been done in several

ways (see Bissett, 2014 for a non-parametric approach). The standard approach for examining

this assumption is comparing predicted signal-respond RTs from fitting the independent race

model to the observed signal-respond RTs (Verbruggen and Logan, 2009). The independence

assumption is typically assessed by showing that observed signal-respond RTs and those

predicted by the independent race model are not different. Because we used a parametric (ex-

gaussian) based approach to estimate the SSRT (Matzke et al, 2013), we verified these derived

fits by using a posterior predictive model comparison to the observed data. The models were

used to simulate 500 predictive distributions of signal-respond RTs to estimate the absolute

goodness of fit (Gelman & Hill, 2007) to each individual subject's signal-respond RT. This

approach generates p-values that test for the difference in the predicted and observed signal-

respond RTs at each SSD level. The typical metric for assuming goodness of fit is that the p-

value is close to 0.5, while being below and above 0.05 and 0.95, respectively, is considered a

poorly predictive model. As pointed out by Matzke et al. (2013), these estimates are most stable

for SSDs in which several signal-respond RTs are observed. Therefore, we analyzed each

subject's p-values averaged across the two highest SSDs.

The mixed-design ANOVA across groups and TFUS conditions did not reveal any

effects or interaction on the *p*-values. This result also agrees with the analysis showing no

differences in signal-respond RTs between TFUS conditions, indicating the Bayesian procedure

for estimating SSRT provided accurate predictions of signal-respond RTs. For all participants,

the *p*-values were in the range of 0.1 to 0.9 with a mean of 0.48 and standard deviation of 0.2.

1497

S4. SSRT Variability was not altered by TFUS.

One possible driver of increased response inhibition performance in the rIFG TFUS

group is that it may have reduced within-subject SSRT variability. Therefore, as TFUS did not

alter mean or variance of Go RTs, if it reduced SSRT variability, this would increase the

likelihood of successful inhibition. Using the same statistical approach for the mean SSRT in the

main text, we found no significant effects of SSRT variability between any of the conditions (all *p*

> 0.05).

1505

S5. SS (No-TFUS) – SS (Stop-TFUS) whole-brain regression table

Table S1 lists the peak coordinates of clusters of the difference in evoked activity of No-

TFUS and Stop-TFUS SS with the between-subjects changes in SSRT. These coordinates were

used to identify which areas exhibited differential activation with respect to the speed of

stopping across subjects. Coordinates for relevant regions of interest were used as prior

locations in subsequent dynamic causal modeling in the main text.

1512

Regression of ΔSSRT and SS (No-TFUS) – SS (Stop-TFUS)					
Time-Window (ms)	Regression Direction	Region	Peak Coordinates (X,Y,Z MNI)	Z-value	Extent (Voxels)
-40:20	Positive	R Supramarginal	56, -40, 16	2.05	316
		L Supramarginal	-56, -50, 18	1.94	123
		R Inferior Temporal	40, -4, -40	1.92	45
	Negative	R Inferior Frontal	40, 30, 10	1.94	55
20:100	Negative	R Inferior Frontal	56, 28, 12	2.64	455
100:180	Positive	R Superior Frontal	26, 50, 18	3.18	568
		R Paracentral Lobule	6, -20, 68	2.98	389
		L Paracentral Lobule	-6, -20, 68	2.94	563
		L Superior Frontal	22, 52, 28	2.74	574
	Negative	L Supramarginal	-60, -28, 38	1.76	52
180:260	Negative	R Inferior Occipital	46, -82, -6	4.12	41
		R Inferior Temporal	58, -40, -24	3.62	552
		L Inferior Temporal	-58, -38, -18	3.55	342
		R Middle Temporal	62, -2, -24	3.16	22
260:340	Positive	L inferior Frontal	-46, 14, 32	4.25	39
		L Middle Frontal	-30, 22, 34	2.84	236
		R Middle Frontal	26, 14, 42	2.61	442
	Negative	R Angular	44, -62, 38	3.62	62
		L Postcentral	-42, -34, 52	2.91	92
		R Middle Temporal	50, 8, -30	2.81	73

1513   Table S1. Significant regions of regression for the change in TFUS induced SSRT.

1514

S6. Whole-brain SPM analysis

We performed a whole-brain SPM analysis for (1) positive t-contrast for SS (No-TFUS) –

SS (Stop-TFUS) trials to examine where activity was larger for TFUS compared to No-TFUS

trials, (2) SS-US main effects, and (3) TFUS main effect. In all of the analysis below, we used a

cluster-forming voxel threshold of p< 0.005

SS-SS t-contrast. In the stop-signal locked window (-40:20 ms), we found differences in

left supramarginal gyrus. In the N100 window (20:100 ms), the contrast was significant, as

anticipated, in rIFG. This result supports our scalp ERP measures and previous results

indicating that an N100 occurs in the rIFG that is predictive of stop success. In the N200 window

(100:180 ms), we found increased source activity in TFUS trials in bilateral clusters with

locations indicative of pre-SMA (Mayka et al., 2006). Consideration of the fourth time window

(180:260 ms), which overlapped with the SSRT, revealed that only the right inferior occipital

area produced a larger response in TFUS trials. In the last window (260:340 ms), we found

clusters of activity larger in TFUS trials for left superior occipital, right cuneus, and anterior

cingulate. Because this window always occurred after the SSRT for both No-TFUS and TFUS

conditions, it is likely these changes in activity represent a component of performance

monitoring rather than inhibition *per se.* Examination of the contrast for larger No-TFUS SS trial

activity revealed this contrast was only significant in the latest time-window (260:340 ms). This

contrast indicated source ERP activity was larger in No-TFUS trials in bilateral postcentral gyrus

clusters. An important result from these contrasts is the primary areas in which activity was

larger during TFUS trials. We found that both rIFG and pre-SMA activity coincided with

increased TFUS-related stopping. Interestingly, this analysis confirmed that rIFG differences

occurred before those in pre-SMA. However, a temporal precedent of change in ERP does not

necessitate that stopping-related changes occurred in rIFG before pre-SMA. The source cluster

MNI locations and cluster sizes are presented in Table S2.

1540

1541

SS (No-TFUS) – SS (Stop-locked TFUS)					
Time-Window (ms)	t-contrast Direction	Region	Peak Coordinates (X,Y,Z MNI)	Z-value	Extent (Voxels)
20:100	TFUS > No-TFUS	Right Inferior Frontal	44, 28, 5	2.8	979
100:180	TFUS > No-TFUS	L pre-SMA	-10, -10, 50	2.7	1108
		R pre-SMA	6, 22, 50	2.67	1069
		R DLPFC	30, 28 40	2.91	850
180:260	TFUS > No-TFUS	Right Inferior Occipital	50, -70, 8	2.62	47
260:340	TFUS > No-TFUS	Left Superior Occipital	-18, -82, 34	2.34	594
		Right Cuneus	18, -70, 32	2.31	259
		Anterior Cingulate	8, 6, 24	2.01	253
	TFUS < No-TFUS	Left Postcentral	-18, -38, 68	2.02	578
		Right Postcentral	26, -24, 70	1.99	362
P. Uncorrected Voxel < 0.005 and P. Cluster Uncorrected < 0.05					

Table S2. MNI locations for SS-SS contrasts

SS-US and TFUS F-contrast. Here we show brain areas that exhibited differential

evoked activation according to (1) successful compared to unsuccessful stopping (SS – US F-

contrast), and (2) which areas were modulated by TFUS (TFUS F-contrast). We computed

these whole-brain SPM contrasts using a flexible factorial model to implement a repeated-

measures ANOVA. We examined these contrasts over three time windows that covered the

time from stop-signal onset till after the range of subject SSRTs. The main result from all of

these analyses is that the only brain area exhibiting overlap between a significant SS-US and

TFUS contrast was a right inferior frontal gyrus cluster centered on pars opercularis, during the

20:100 ms time window. This result supports the accuracy of our TFUS manipulation and its

effects on inhibition performance. These data are shown in Figures S3-5.

1553

FWE P	Cluster Extant	Uncorr. P	Peak-F	X	Y	Z	Cluster Peak
F-contrast: Successful Stop – Unsuccessful Stop (SS-US)							
<0.005	893	0.0001	16.35	50	-40	-26	R Inferior Temporal
<0.05	586	0.0001	15.16	54	26	18	R Inferior Frontal
F-contrast: non-tFUS – stop-tFUS (tFUS)							
<0.01	620	0.0001	16.65	-42	-72	-8	L Inferior Occipital
<0.005	899	0.0001	16.09	52	18	22	R Inferior Frontal:
<0.01	699	0.0001	15.52	48	-74	-10	R Inferior Occipital

1554

1555   Figure S3. SPM F-contrasts of source evoked power during the 20:100 ms post stop signal
1556   window. F-contrasts and their cluster corrected statistics are shown for the SS-US contrast and
1557   the contrast of SS trials during non-TFUS and stop-TFUS.

FWE P	Cluster Extant	Uncorr. P	Peak-F	X	Y	Z	Cluster Peak
F-contrast: Successful Stop – Unsuccessful Stop (SS-US)							
<0.001	1147	<0.0001	24.87	24	-34	68	R Postcentral
<0.0005	1279	<0.0001	21.85	-22	-28	72	L Pre/Postcentral
<0.05	479	<0.005	19.23	-36	-52	42	L Inferior Parietal
F-contrast: non-tFUS – stop-tFUS (tFUS)							
<0.0001	1337	<0.0001	18.41	22	8	56	R pre-SMA
=0.2	343	<0.01	16.15	52	18	22	R Middle Temporal
<0.05	590	<0.001	13.18	48	-74	-10	L Middle Temporal
<0.001	1113	<0.0001	11.99	-6	-28	56	L Precentral

1558

1559   Figure S4. SPM F-contrasts of source evoked power during the 100:180 ms post stop signal
1560   window. F-contrasts and their cluster corrected statistics are shown for the SS-US contrast and
1561   the contrast of SS trials during non-TFUS and stop-TFUS.

1562

1563

FWE P	Cluster Extant	Uncorr. P	Peak-F	X	Y	Z	Cluster Peak
F-contrast: Successful Stop – Unsuccessful Stop (SS-US)							
<0.05	497	<0.005	16.83	46	-20	-14	R Middle Temporal
F-contrast: non-tFUS – stop-tFUS (tFUS)							
<0.0001	1702	<0.00001	19.64	42	-58	-10	R Inferior Temporal
<0.001	1270	<0.0001	17.92	-58	-40	-14	L Middle Temporal
<0.05	564	<0.005	13.49	-4	60	0	L Medial Superior Frontal

1564

1565 Figure S5. SPM F-contrasts of source evoked power during the 180:260 ms post stop signal
1566 window. F-contrasts and their cluster corrected statistics are shown for the SS-US contrast and
1567 the contrast of SS trials during non-TFUS and stop-TFUS.

1568

Interaction t-contrasts. The most common contrast in analyzing stop-signal neural data

involves comparing successful to unsuccessful stop activation (SS – US) as we did for the

above whole-brain analysis. The rationale is based on the assumption that areas directly related

to stopping/inhibition are more 'potently' active in successful trials, and that failed stop trials

(US) also reflect go activity according to the independent race model of Logan and Cowan

(1984). This choice of contrast also stems from the fact that typical stop-signal tasks do not offer

a second set of SS trials for comparison (which our experiment does). Therefore, this raises the

following question: Did TFUS yield changes in stopping by merely altering what would have

been US Stop-TFUS trials? If TFUS merely raised the overall level of activity across all stop

trials, (i.e., SS and US), then we should expect no interactions. We examined the whole-brain

SPM interaction across the 4 time windows used in the previous analysis. In each window, we

examined the t-contrast interaction that compared for a bigger SS-US difference in either Stop-

TFUS or No-TFUS trials. We only found these effects for the 100:180 ms and 180:260 ms

window. The interaction t-contrast SPMs and corresponding tables are shown in Figures S6-7.

1583

Window: 100:180 ms

FWE P	Cluster Extent	Uncorr. P	Peak-T	X	Y	Z	Cluster Peak
F-Contrast: Interaction SS-US (stop-tfUS) > SS-US (non-tFUS)							
<0.001	1364	<0.0001	4.42	52	-18	-12	L Postcentral
<0.005	1153	<0.00001	4.41	42	-90	4	R Middle Occipital
<0.005	988	<0.00001	4.30	32	-4	38	R Inferior Temporal
<0.05	670	<0.005	3.70	44	14	10	R Frontal Operculum /Inferior Frontal /Pars Opercularis

1584

1585   Figure S6. SPM interaction t-contrasts of source evoked power during the 100:180 ms post stop
1586   signal window. The t-contrast tested for areas in which the SS-US contrast was higher during
1587   stop-TFUS compared to non-TFUS. The cluster corrected statistics are shown are shown below
1588   the plot.

1589

1590

Window: 180:260 ms

FWE P	Cluster Extent	Uncorr. P	Peak-T	X	Y	Z	Cluster Peak
F-Contrast: Interaction SS-US (stop-tfUS) > SS-US (non-tFUS)							
<0.01	706	<0.001	4.21	52	-18	-12	R Middle Temporal
<0.005	984	<0.00001	3.90	42	-90	4	R Middle Occipital
<0.05	567	<0.0001	3.73	46	16	4	R Inferior Frontal Gyrus / Pars Opercularis
<0.005	822	<0.001	3.61	-2	14	64	pre-SMA/SMA
<0.005	780	<0.005	3.25	-34	-68	28	L Middle Occipital

1591

1592 Figure S7. SPM interaction t-contrasts of source evoked power during the 180:260 ms post stop
1593 signal window. The t-contrast tested for areas in which the SS-US contrast was higher during
1594 stop-TFUS compared to non-TFUS. The cluster corrected statistics are shown are shown below
1595 the plot.

1596

We note that we only found significant contrasts in the latter two time windows, i.e., 100-

180 and 180-260 ms from the Go signal. In both contrasts, the SS-US difference was larger in

the Stop-TFUS conditions. Generally, these results agree with that of the main text examining

just the SS-SS contrast and other studies (Aron and Poldrack, 2006; Boehler et al., 2010).

However, as Boehler et al. (2010) point out, this contrast is conservative which lends itself to

identifying areas primarily involved in successful stopping. It is also unlikely that this

conservative contrast can identify areas involved in the broader stopping network. It is therefore

not surprising that we did not find difference in parietal cortices, for example. Importantly,

though, these results demonstrating mainly activation differences in inferior frontal and occipital

cortices is almost identical to that found in Boehler et al. (2010). This indicates that our results

generally agree with previous studies using the SS − US contrast.

1608

S7. ROI Interaction analysis

In the main text we presented a time-based source analysis of the main ROIs of interest,

including Right IFG, Right pre-SMA, Right DLPFC, Left M1, and Right Inferior Temporal

cortices. In the main text's analysis, we compared SS trials across TFUS conditions. To address

the change in time-course activation at the ROI level, we computed the interaction of SS-US

(no-TFUS) and SS-US (Stop-TFUS) trials using the same MNI locations as used for analyses

reported in the main text (Fig. S6).

1616

Figure S8. Source-base evoked power time-series for different regions-of-interest listed in the
figure. Each ROI was extracted with a radius of 6 mm.

Only three of these ROIs are different before the minimum SSRT across TFUS conditions (185

ms). These include pars opercularis, Right Inferior Occipital, and Right inferior temporal ROIs.

This result is mostly in line fMRI (Boehler et al., 2010; Li et al., 2006) and EEG/MEG (Boehler et

al., 2009; Wessel and Aron, 2015) studies, although previous work has also identified Right

temporal cortex (Rae et al., 2015; Xu et al., 2017). Nevertheless, previous findings of differential

temporal cortex activity have been based on comparing SS to Go trials rather than the SS − US

(and interaction) comparison we used here. Given the DCM and SSRT regression results

indicating changes in temporal cortex activity – as well as the temporal cortex's role in visual

signal in both the top-down and bottom-up processing directions – these results suggest the

success of stopping may rely on information transfer of node between sensory and prefrontal

ventral areas, e.g., rIFG. Finally, an interesting result is the lack of difference in r pre-SMA

before SSRT. Notably, there was differential activation but only after SSRT. This result speaks

to the broader debate of whether pre-SMA or rIFG lead to stopping in a serial process fashion

(Aron et al., 2016; Obeso et al., 2013). For example, some studies have suggested that

information for stopping passes through pre-SMA onto rIFG and vice versa in other accounts.

These serial accounts, though, forego three factors. First, visual information regarding stop

contexts are passed up both the ventral and dorsal pathways which inherently project

separately to rIFG and r pre-SMA, respectively. Second, the prefrontal cortex is likely arranged

hierarchically with both areas connected with basal ganglia structures, such as the STN and

striatum. The third comparison is based on considering the current EEG-DCM (main test)

results to those from other fMRI-DCM and other connectivity results (Jahfari et al., 2017) on

inhibition. Specifically, the previous bilinear DCM models used in fMRI accounts of response

inhibition are unlikely to capture the fast-timescale processes that underlies stopping processes.

Along this same line, though, when considering connectivity as a property of inhibition

processes, caution should also be used in interpreting temporal precedence of control between

brain areas based solely on activation. Thus, the above results indicate that the set of expected

ROIs were differentially modulated by TFUS during SS stopping.

S8. Assessment of possible auditory effects of TFUS

Recent work examining the effects of TFUS on cortical activity has employed animal models to consider the possibility that TFUS may alter activity in auditory pathways. Using a single-element transducer and optical imaging on a mouse model, Guo et al. (2018) showed that, regardless of the transducer placement and target, TFUS caused activation of the auditory pathway. They suggested that this activity may spread cortically and induce artifactual effects of TFUS in cortical areas not directly targeted by TFUS. It remains unknown how auditory pathway activation via TFUS would yield our behavioral and neural effects. Nevertheless, it is critically important to quantify the extent to which auditory pathway activation might have affected our results. To address this question, we analyzed source-localized evoked results across the rIFG group, as well as S1 and sham rIFG control groups. Because our goal was to determine whether TFUS altered the evoked activity with respect to No-TFUS, we compared the time courses of source power for stop-locked data by using a source-based ROI of right auditory cortex. We used the source location of X: 46, Y: -14, Z: 8 for right auditory cortex, which was obtained from Rademacher et al. (2001). These locations were used to extract the eigenvariate time-course after source localization in a sphere with radius of 8 mm to be conservative. These source-time courses were converted to a pseudo activation using the (exp(SOURCE)+exp(-SOURCE))/2 transform. We used this procedure to ensure we could properly detect differences regardless of the ERP activity sign (Fig. S9).

If the behavioral results in the rIFG group were merely the result of changes in auditory pathway activity, we should at least see a difference when contrasting the SS-US trials for No-TFUS and Stop-TFUS conditions in the rIFG group, or at least across groups. Paired samples t-tests did not reveal any significant differences (after false discovery rate correction of $p < 0.05$) when comparing the time courses of the No-TFUS and Stop-TFUS conditions (Fig. S9).

1670 Therefore, given the spatiotemporal fidelity of EEG, we conclude that our results were not

1671 contaminated by TFUS artifacts activating the auditory pathway.

1684

Figure S9. Group source-evoked power time series from right auditory cortex.

Overall, these results suggest that, although TFUS may exert an auditory pathway

effects measurable in mouse single neurons and LFPs, (1) this activity may not be measurable

at the macroscopic level, and (2) support the notion that changes in auditory cortical activity

cannot account for our TFUS-related neural and behavioral effects. Additionally, we note that

Guo et al. (2018) found that the auditory pathway activation was accompanied by startle-like

reflexes. Our behavioral results (Fig. 2) are not compatible with a startle release reflex. If that

were the case, TFUS auditory-related startle activity would likely predict shorter RTs during Go

trials and failed inhibition Stop trials. The effects of TFUS on our behavioral responses are not

compatible with the involvement of startle reflexes, as we found no TFUS effects on the Go

RTs. Therefore, we conclude that the effects of TFUS on stopping behavior, nor processing by

the cortical and subcortical nodes of the inhibition network, were not induced by artifactual

stimulation of the auditory pathway.

1698

S9. DCM family model hypothesis spaces and results

1699

Below we show the resultant family model posterior probabilities for the family comparisons over

the 4 factors (3 levels each) for both parametric empirical Bayesian (PEB) GLMs (Figure S10-

11). PEB GLM model 1 examined the changes in mean connectivity from No-TFUS US to SS

trials. Changes in mean connectivity represent the gain on connectivity to represent the No-

TFUS SS trials. PEB GLM model 2 examined the effects of TFUS-induced changes in mean

connectivity and changes in connectivity that accompanied change in SSRT across subjects.

Figures S8 and S9 show the changes in connections and family model probabilities for PEB

model 1. Figure S12 shows the results of tests of the exact same factor space displayed as bar

plots to represent the family probability of each model separately for the TFUS-induced mean

and SSRT change in successful inhibition connectivity.

1710

Figure S10. Hypothesis space and results of the Family based PEB Bayesian model
comparison for different hypotheses. Family-based posterior probability (F-Pp) and the log-

bayes factor with respect to the winning model are listed below each model and were computed
as the difference in free energy between families (F-Pp/Log-Bayes). The winning model family is
enclosed in a red box. The outcomes of these plots can be interpreted of as revealing the
modulatory parameters connections with very strong, positive evidence being different between
US and SS trials. Parameters estimates with a greater than 95% posterior probability in these
families are presented in parentheses next to the modulated connection. The parameters are
presented in exponential form of percentage gain. Values above 100% equates to a parameter
increase in SS trials compared to US (and the opposite for values below 100%). Parameters in-
active in each model are in a gray color. The top Panel shows the hypothesis test of rIFG and
pre-SMA interactions. The bottom panel shows the hypothesis test of intrinsic gains modulation.

1723

1724

Figure S11. Hypothesis space and results of the Family based PEB Bayesian model
comparison for 3 different families comparing hypothetical different interactions between rIFG
and the deep node, pre-SMA and the deep node, or both interacting with the deep pathway. The
plot is in the same format as Figure S8. Both of these nodes had backward, inhibitory
connections with the deep pathway. The top panel compares families comparing the rIFG and
pre-SMA to deep backwards connection. The bottom panel compares families testing for
differences in top-down v bottom-up connections along the ventral pathway.

1732

1733

1734 Figure S12. Resultant family posterior probabilities for the comparisons of each factor. Each bar
1735 plot shows the marginal probability of each family marginalized separately for changes in the
1736 mean connectivity (blue bars) and connectivity changes predicted by TFUS induced changes in
1737 SSRT (orange bars). The Bayesian model averaged parameters are presented in the main text.