# Fast sequence-based microsatellite genotyping development workflow for any non-model species.

Olivier Lepais[1,2*], Emilie Chancerel[1], Christophe Boury[1], Franck Salin[1], Aurélie Manicki[2], Laura Taillebois[2], Cyril Dutech[1], Abdeldjalil Aissi[3], Cecile F. E. Bacles[2], Françoise Daverat[4], Sophie Launey[5], Erwan Guichoux[1].

[1] BIOGECO, INRA, Univ. Bordeaux, 33610 Cestas, France

[2] ECOBIOP, INRA, Université de Pau et Pays de l'Adour, 64310 Saint-Pée-sur-Nivelle, France

[3] LAPAPEZA, University of Batna 1 Hadj Lakhdar, 0500 Batna, Algeria

[4] EABX, IRSTEA, Cestas Cedex 33612, France

[5] ESE, Ecology and Ecosystem Health, Agrocampus Ouest, INRA, 35000 Rennes, France

[*]Corresponding author

**Abstract:** Application of high-throughput sequencing technologies to microsatellite genotyping (SSRseq) has been shown to remove many of the limitations of electrophoresis-based methods and to refine inference of population genetic diversity and structure. However, early proof of concept and species specific development studies resulted in dispersed information making it cumbersome for prospective users to identify a clear path to SSRseq approach set up in species of new interest. To overcome these difficulties, we present here a streamlined SSRseq development workflow that includes microsatellite development, multiplexed marker amplification and sequencing, and automated bioinformatics data analysis. We demonstrate its application to five groups of species across kingdoms (fungi, plant, insect and fish) with different levels of polymorphism and genomic resource availability. We found that relying on previously developed microsatellite assay is not optimal and leads to a resulting low number of reliable locus being genotyped. In contrast, *de novo ad hoc* primer designs gives highly multiplexed microsatellite assays that can be sequenced to produce high quality genotypes for 20 to 40 loci. We highlight critical upfront development factors to consider for effective SSRseq setup in a wide range of situations. The automated sequence analysis pipeline, which accounts for all linked polymorphisms along the sequence, quickly generates a powerful multi-allelic haplotype-based genotypic dataset. Cost and time effective application of SSRseq approaches are within reach for any species, calling to new theoretical and analytical frameworks to extract more information from multi-nucleotide polymorphism marker systems.

**Keywords**: sequence-based microsatellite genotyping, GBS-SSR, SSRseq, haplotype sequence, SNPSTR, HapSTR.

## Introduction

Development of high-throughput sequencing (HTS) technologies since the mid 2000's has greatly widened the scope of genetic applications in research fields such as agronomy, ecology and evolutionary biology where the use of molecular markers is common place. The easy collection of millions of DNA sequences has also facilitated the development of molecular markers such as single nucleotide polymorphism (SNP; (Garvin *et al.* 2010; Delord *et al.* 2018)) or microsatellites (or single sequence repeats SSR; (Guichoux et al., 2011; Malausa et al., 2011)) for any species of interest. Beyond marker development, HTS has been largely adopted for genotyping-by-sequencing (GBS), thus taking advantage of restriction enzymes to target sequencing efforts in specific parts of the genome that are shared by conspecific individuals (Miller *et al.* 2007; Baird *et al.* 2008). Restriction site-associated DNA polymorphism sequencing allows simultaneous identification and genotyping of SNP in non-model species and interrogation of thousands to hundreds of thousands of loci genome-wide but remains expensive when population samples comprise hundreds of individuals. Recent development of amplicon sequencing which targets previously identified SNP makes it possible to genotype hundreds to thousands of individuals at a few hundreds of SNP at a very reasonable cost (Nguyen-Dumont *et al.* 2013; Campbell *et al.* 2015; Aykanat *et al.* 2016). If sequence-based genotyping of SNP is gaining momentum, application of HTS to microsatellite genotyping was until recently lagging behind. Updating microsatellite genotyping to modern technologies remains important for several reasons. Firstly, some current scientific questions in ecology or evolutionary biology can be answered using a moderate number (e.g. a dozen) of highly polymorphic multi-allelic loci such as microsatellites (Harrison *et al.* 2013). Secondly, variation in the number of repeated oligonucleotide motif is a unique kind of polymorphism with specific mutation mechanism and rate which in itself provides a complementary picture of genetic variation to nucleotide substitutions across populations (Haasl and Payseur 2011) and genomes (Willems *et al.* 2014). Thirdly, it becomes more and more obvious that microsatellite polymorphism is involved in numerous biological processes such as gene expression regulation and epigenetic mechanisms (Bagshaw 2017; Sadd *et al.* 2018), and more generally in phenotypic variation (Xie *et al.* 2019) including human diseases (Gymrek 2017; Hannan 2018). Thus, while marker preference evolves through time with specific markers dominating the genotyping field over a period of time following technological advances (Schlötterer 2004; Seeb *et al.* 2011), maintaining our capability to interrogate any kind of polymorphism in the context of rapid HTS technological advances is paramount and should be prioritized.

In the age of HTS, and in comparison with SNP genotyping, microsatellite genotyping suffers from several drawbacks: homoplasy (alleles of identical size having different underlying sequence (Viard *et al.* 1998; Estoup *et al.* 2002)), time and cost consuming development and genotyping, low

throughput, lack of automation and data standardization (Moran *et al.* 2006; Ellis *et al.* 2011). Yet, all of these limitations are linked to the fact that currently microsatellite genotyping relies on allele discrimination based on amplicon size assessed by capillary electrophoresis (De Barba *et al.* 2016) and do not hold true if microsatellite genotyping transitions to sequence-based genotyping. Previous direct comparisons of capillary electrophoresis and sequence-based microsatellite genotyping (called SSRseq thereafter) validated SSRseq as a reliable method (Darby *et al.* 2016; Vartia *et al.* 2016). Advantages of sequence-based over capillary electrophoresis-based microsatellite genotyping are significant. Direct access to allele sequence reveals additional polymorphisms that remain hidden when using only allele size to identify variation (Darby *et al.* 2016; Vartia *et al.* 2016; Barbian *et al.* 2018). Sequence data thus reduces allele homoplasy because alleles of the same size may contain molecular variation that does not translate into size variation such as SNP, indels masking variation in repeat number, or presence of two adjacent SSR motifs with complementary size variation (Darby *et al.* 2016). As a result, SSRseq offers refined genetic diversity estimation and population structure inference (Darby *et al.* 2016; Bradbury *et al.* 2018; Neophytou *et al.* 2018; Viruel *et al.* 2018).

Studies published so far have explored specific technical or analytical aspects of SSRseq. Several bioinformatics approaches have been developed (Hoogenboom *et al.* 2016; Suez *et al.* 2016; Zhan *et al.* 2017; Barbian *et al.* 2018), different laboratory protocols tested (Vartia *et al.* 2016; Pimentel *et al.* 2018) and means to account for molecular variation on population genetic inference compared (Neophytou *et al.* 2018; Viruel *et al.* 2018). Together, these studies explored numerous issues surrounding the technical and analytical advantages of SSRseq over traditional methods, each of them focusing on a given model species making it difficult to assess finding generalization to other biological models. In addition, laboratory methods for microsatellite marker amplification as well as analytical approaches to convert raw sequences to codominant genotypes markedly differ between studies making it cumbersome for prospective users to identify a clear path to setup a new SSRseq approach for a species of new interest.

Here, we propose an integrative workflow for the development of a SSRseq analysis for application to any species. We apply this workflow to five species groups from families taken across Phylums (Basidiomycota: *Armillaria ostoyae*; Angiosperms: *Quercus faginea* and *Q. canariensis*; Euarthropoda: *Melipona variegatipes*; Chordata: *Alosa alosa, A. fallax* and *Salmo salar*) that markedly differ in the amount of genomic data already available for them. We compare a broad range of possible development scenarios including sequencing of already optimized microsatellite assay traditionally genotyped on capillary sequencers, optimizing primers around already developed microsatellites, and developing microsatellite *de novo* from a range of genomic resources when available, or from newly generated low coverage random genome sequences for species without existing genomic

resources. Building on our previous experience in development of highly multiplex microsatellite genotyping protocols (Guichoux et al., 2011; Lepais & Bacles, 2011), we propose a streamlined, simple and cost effective approach with demonstrated application to groups of species which a wide range of genetic and evolutionary characteristics. We offer an automated microsatellite sequence data analysis pipeline to produce haplotypic data, validated by extensive blind-repeat genotyping to estimate SSRseq error rates. We explore two different data analysis approaches that either account for the whole range of polymorphism detected within the repeated motif and the flanking regions, or restrict the analysis to the repeated motif itself. We find that relying on previously developed microsatellite assay is far from optimal and result in a low number of reliable locus genotyped. In contrast, primer redesign based on strict criteria around known locus or *de novo* microsatellite development gives successful single highly multiplexed PCR amplification and sequencing, producing high quality genotypes for 20 to 40 loci. Overall, we highlight critical upfront developments needed to generalize SSRseq setup effectively. We emphasize that efficient and powerful multi-polymorphism haplotype-based genotyping approaches are time and cost effective and achievable in virtually any species, calling for new theoretical and analytical development to extract more information from this new kind of molecular markers.

## Materials and Methods

### Studied species, SSRseq development strategies and DNA isolation

We selected a range of species from different Kingdoms among biological models studied in our laboratories (Table 1). For *S. salar* we took the most straightforward route by amplifying and sequencing microsatellites using previously developed primers. To develop a refined workflow, we chose species with different level of genomic resource availability to test alternative *de novo* microsatellite development strategies that are likely to cover a wide range of situations (Table 1). Protocols for DNA isolation and final DNA quality differ between species (Table 1).

Table 1: SSRseq development strategy and DNA characteristics of species used in this study.

| Kingdom | Class | Species | SSRseq development strategy | Number of DNA extracted individuals | DNA extraction protocol | DNA quality | Reference |
|---------|-------|---------|------------------------------|--------------------------------------|-------------------------|-------------|-----------|
| Animalia | Actinopterygii | *Salmo salar* | Previously developed loci | 1152 | Salt-chloroform (Gauthey *et al.* 2015) | High | (Bacles et al., 2018; Lepais et al., 2017) this study |
| Plantae | Eudicots | *Quercus faginea*, *Q.canariensis* | Re-designed primers around already developed loci using reference genome sequence of closely related species | 380 | Invisorb DNA Plant HTS 96 kit | High | |
| Animalia | Actinopterygii | *Alosa alosa*, *A.fallax* | *De novo* loci development based on available repeat-enriched library sequencing | 382 | Invitrogen PureLink Genomic DNA Mini kit | Highly degraded | (Taillebois et al., submitted) |
| Fungi | Agaricomycetes | *Armillaria ostoyae* | *De novo* loci development based on reference genome sequence | 384 | CTAB (Prospero *et al.* 2008) | Heterogeneous | this study |
| Animalia | Insecta | *Melipona variegatipes* | *De novo* loci development based on newly generated low coverage whole genome shotgun sequencing | 91 | Qiagen DNeasy 96 Blood & Tissue Kit | High | this study |

**SSRseq using previously-developed primers**

The most straightforward approach to SSRseq microsatellite genotyping, i.e. based on sequence information from existing primers, was applied to *S. salar* using two marker selection strategies. In the first strategy, we selected 23 primers from a list of 81 microsatellites available for *S. salar* (O'Reilly and Wright 1995; Slettan *et al.* 1996; Ozaki *et al.* 2001; Rexroad *et al.* 2001; Gilbey *et al.* 2004; Paterson *et al.* 2004; King *et al.* 2005; Vasemägi *et al.* 2005; Thorsen *et al.* 2005; Yano *et al.* 2013, see details in Supporting Table S1) and mapped (Vasemägi *et al.* 2010). Selection criteria included allele size smaller than 300 bp to ensure that sequencing reads can span the entire allele length including library construction and absence of sequences complementarity between primers tested using Multiplex Manager (Holleley and Geerts 2009). In the second strategy, we chose to sequence a set of 15 microsatellites (Supporting Table S1) that are routinely amplified in a single multiplexed PCR and genotyped using electrophoresis-based capillary sequencers (Bacles et al., 2018; Lepais et al., 2017).

**SSRseq with microsatellite (re)development**

*Genomic resources for microsatellite (re)development*

For the other species, microsatellites primers were either re-designed or developed *de novo* from various genomic resources (Figure 1 top panel).

For *Quercus* sp., primers were re-design primers in flanking regions of existing microsatellite markers to optimize multiplex amplification and sequence interpretability while taking advantage of already validated microsatellite markers. We extracted primer sequences from 259 polymorphic and mapped EST-derived (Durand *et al.* 2010) and 35 genomic microsatellites (Steinkellner *et al.* 1997; Kampfer *et al.* 1998)). The primer sequences were mapped on the *Q. robur* reference genome (Plomion *et al.* 2018, GenBank accession GCA_003013145.1) using bowtie 2 v2.3.4.1 (Langmead and Salzberg 2012) and genomic sequence spanning from 200 bp downstream of the forward primer to 200 bp upstream of the reverse primer position were extracted using bedtools v2.25.0 (Quinlan 2014) resulting in 294 sequences used as genomic resource to design new primers.

For *Alosa* sp., we used sequences obtained from a Roche 454 GS-FLX sequencing run on a microsatellite-enriched DNA library (Rougemont *et al.* 2015) following the method described in Malausa *et al.* (2011).

For *A. ostoyae*, we used the reference genome sequence as genomic resource to identify microsatellites loci (Sipos *et al.* 2017, GenBank accession GCA_900157425.1).

Finally, no genomic resources were available for *M. variegatipes*. We therefore used DNA from one individual to construct a whole-genome sequencing library using Illumina TruSeq DNA kit. The resulting library was sequenced on an Illumina MiSeq flowcell using v3 2x300 pb paired-end sequencing kit. Mothur software v1.39.5 (Schloss *et al.* 2009) was used to assemble paired reads and keep paired reads with a minimum overlap of 100 bp without mismatch. We randomly subsampled 500,000 reads from the resulting 6.74 million paired reads for subsequent microsatellite identification, because a few hundreds of thousands of random sequences are sufficient to identify thousands of microsatellites (Castoe et al., 2010; Lepais & Bacles, 2011).

Figure 1: Workflow for SSRseq markers optimization or development depending on genomic resource availability, from selection to multiplexed amplification and library preparation to bioinformatics analysis (refined approach).

De novo *microsatellite development or primer re-design*

The command line version of QDD pipeline v 3.1 (Meglécz *et al.* 2010, 2014) was run on either i) a reference genome sequence, ii) a set of low coverage random sequences or iii) sequence extracted around already characterised microsatellite loci, to detect sequences containing microsatellites, identify good quality sequence (singletons and consensus) from problematic sequences (sequences showing low complexity, minisatellites or multiple BLAST hits with other sequences) and design primer pairs flanking the identified microsatellites (Figure 1). QDD pipeline was run with default parameters, except for the primer design step (pipe3) where parameters were stringently defined in order to improve their capacity to be amplified jointly in a single multiplexed PCR (Qiagen Multiplex

kit handbook; Lepais & Bacles 2011): primer optimal size was set to 25 nucleotides (min: 21, max: 26), optimal annealing temperature to 68°C (min: 60°C, max: 75°C) with a maximal difference of 10°C between primers of a same pair and optimal percentage of cytosine and guanine of 50% (min: 40%, max: 60%). In addition, PCR product size was set between 120 and 200 bp to be compatible with a wide range of sequencing platforms and to produce robust genotyping assays that can be used to analyse degraded or non-invasive samples. QDD analysis results in a large number of candidate loci with designed primer pairs from which a restricted number of loci can be selected (Figure 1). At the exception of *Quercus* sp. where a restricted set of input sequences necessarily limited choice among resulting candidate microsatellites, several quality criteria were used to select 60 microsatellites among hundreds to thousands candidates for further testing. We followed recommendations of Meglécz *et al.* (2014) to prioritize primer pairs with increased likelihood of amplification success by selecting microsatellite from singletons and not from consensus sequence, with pure repeat instead of compound motifs, showing at least 20 bp between the primers and the repeat motif, and with flanking region showing high complexity (e.g. primer pairs from the design group A following QDD terminology: no minisatellite, no other microsatellite in the flanking region, no homopolymer in the flanking region or the primer). In addition, we further selected microsatellites with the highest number of repeats to increase the probability of selecting polymorphic loci, avoided motif that can form hairpin such as AT repeats and when possible included a variety of di-, tri- and tetra nucleotide repeats.

**Primer modification and simplex amplification tests**

For Ion Torrent sequencing (Table 2), tag A 5'-GCCTTGCCAGCCCGCTCAG-3' was add to the 5' end of each forward primer and tag B sequence 5'-GCCTCCCTCGCGCCA-3' (Margulies *et al.* 2005; Blacket *et al.* 2012) was added to the 5' end of each reverse primer. For Illumina sequencing (Table 2), specific tags 5'-TCGTCGGCAGCGTCAGATGTGTATAAGAGACAG-3' and 5'-GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAG-3' were added to the 5' end of the forward and reverse primer sequences respectively. Designed primers were tested for potential primer dimer formation using Primer Pooler (Brown *et al.* 2017). Primer pairs showing a deltaG lower than -6 kcal/mol are likely to form dimer and result in poor amplification in a multiplexed PCR. For locus involved in significant interactions, alternative primers were selected or in absence of alternative, another locus was selected from the candidate list. Oligonucleotides were ordered in a plate format from Integrated DNA Technologies with standard desalt purification at a final concentration of 100 µM. Primer pairs were tested using simplex amplification performed on one individuals per species using Qiagen Multiplex kit in a final volume of 10µL and with a final concentration of each primer of 0.2µM. Amplification conditions consisted of an initial denaturation step at 95°C for 15 minutes,

followed by 35 cycles consisting of denaturation at 95°C for 20s, annealing at 59°C for 60s and extension at 72°C for 30s, and a final extension step for 10 minutes at 72°C. Amplicons and 1 Kb size standard were then loaded on a 3% agarose gel containing GelRed or SyberSafe dye and migrated at 100v for 15 minutes. Each locus was screened under UV light for positive amplification with a clear band at the expected size.

Table 2: Summary of the tested scenarios for SSRseq genotyping

| Species | SSRseq development strategy | Candidate loci | Screened loci | Number of loci in a single multiplexed PCR | Sequencing Plateform | Total number of individuals sequenced | Number of individual analyzed for this study | Number of repeated individuals | Sequenced loci[†] | Mean (cv) sequences / loci / individual | Reliable loci genotyped[‡] | Overall success rate |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| *S. salar* | Previously developed loci | 81 | - | 23 | Ion Torrent PGM i316 | 960 | 66 | 66 | 20 | 161 (61%) | 9 | 39% |
| | | | | 23 | Illumina MiSeq - 1/2 nano PE | 192 | 96 | 96 | 20 | 70 (41%) | 10 | 43% |
| *S. salar* | Previously developed loci | 15[§] | - | 15 | Illumina MiSeq - 1/2 nano PE | 192 | 96 | 96 | 13[§] | 99 (65%) | 7 | 47% |
| *Quercus* sp. | Primer redesign around previously developped loci | 462 | 60 | 60 | Illumina MiSeq - 1/3 V2 PE | 380 | 46 | 46 | 53 | 260 (32%) | 40 | 67% |
| *Alosa* sp. | *De novo* | 2872 | 60 | 28 | Illumina MiSeq - 1 nano SE | 382 | 156 | 156 | 25 | 95 (58%) | 21 | 75% |
| | | | | 28 | Illumina MiSeq - 2 nano SE | 382 | 156 | 156 | 26 | 198 (58%) | 24 | 86% |
| | | | | 28 | Illumina MiSeq - 3 nano SE | 382 | 156 | 156 | 26 | 267 (58%) | 24 | 86% |
| *A. ostoyae* | *De novo* | 1806 | 60 | 51 | Illumina MiSeq - 1/2 V2 PE | 384 | 384 | 96 | 48 | 243 (83%) | 38 | 75% |
| *M. variegatipes* | *De novo* | 8937 | 60 | 54 | Illumina MiSeq - 1/4 V2 PE | 182 | 91 | 91 | 49 | 176 (45%) | 39 | 72% |

[†]: loci showing substantial evidence for minimum sequencing success (at least 20 sequences in at least 50% of the individuals); [‡]: reliable loci (less than 50% of missing data among individuals and less than 6% of genotyping error based on comparison of repeated genotyping); [§] routinely genotyped using optimized multiplexed PCR and capillary-based sequencer. *FDSTools* analysis using two parameter sets: *stutterfinder* -s:-1:50,+1:10 *allelefinder* -m 15 -n 20; and *stuttermark* -s:-1:70,+1:10 *allelefinder* -m 10 -n 20. For each marker, four parameter combination were used (two strategies and two parameters set) and for each strategy, the best parameter set was used for a given locus.

**Multiplex microsatellite amplification and sequencing library construction**

From 192 to 960 individuals were analyzed depending on the taxa considered including from 46 to 156 repeated individuals to check the reproducibility of the method (Table 2). For each of the taxonomic groups, a three-round multiplex PCR approach was used to amplify all loci simultaneously and improve amplification homogeneity and thus coverage of sequence between loci (Chen *et al.* 2016). In the first round, a multiplexed PCR including all selected locus primers was performed (Figure 1, Supporting Table S1 for locus characteristics including primer sequences). PCR amplification were carried out in 96-well plates in a final volume of 5µL or 10µL using Qiagen Multiplex kit, 0.05 µM of each forward and reverse tailed primers and about 40 ng of template DNA (depending on the species, 1µL of undiluted or diluted isolated DNA). PCR cycles were performed on Applied Biosystems 2720 or Verity thermocyclers and consisted of a denaturing step of 5 min at 95°C followed by 20 cycles of 95°C for 30 s, 59°C for 180 s and 72°C for 30 s. In the second round, additional Taq polymerase added with the aim to use remaining primers completely. The PCR mixture of a final volume of 5 or 10 µL consisted of 2.5 or 5 µL of Qiagen Multiplex kit, 1.5 or 3 µL of undiluted amplicon and 1 or 2 µL of water. The PCR cycles were identical as in the first round. The third round is the indexing PCR (Figure 1) that add Ion Torrent or Illumina sequencing adaptors and barcodes used to assign each sequence to an individual. For Ion Torrent sequencing, we used 106 different barcodes resulting in a total of 960 barcode combinations. For Illumina sequencing, we used the Nextera XT index set allowing for 384 barcode combinations. The PCR mixture of a final volume of 10 µL consisted of Qiagen Multiplex kit Master Mix, 0.5 µM of sequencing platform-specific adaptor and 5 µL of undiluted amplicon resulting from the second PCR round. PCR cycles consisted of a denaturing step of 5 min at 95°C followed by 15 cycles of 95°C for 30 s, 59°C for 90 s and 72°C for 30 s and a final extension step of 68°C for 10 minutes. Amplicons from the 96 wells within a plate were pooled together in an Eppendorf tube, and purified with 1.8X Agencourt AMPure XP beads (Beckman Coulter, UK). Quality check and quantification were done using Agilent Tapestation D1000 kit and Qubit fluorometric system (Thermo Fisher Scientific), and quantified using Kapa libraryquantification kit in a Roche LightCycler 480 quantitative PCR. The resulting two to ten pools were pooled in equimolar concentration and sequenced using an Ion Torrent PGM i316 chip or Illumina MiSeq flowcell using nano or v2 2x250 bp paired-end sequencing kit (Table 2) at the Genome Transcriptome Facility of Bordeaux.

**Bioinformatics data analysis**

*Sequence preparation*

After sequence demultiplexing and adaptor trimming using a sequencer platform built-in software, reads shorter than 70 bp were removed using *cutadapt* (Martin 2011). When paired-end sequencing was used, paired reads were assembled into contigs using *pear* (Zhang *et al.* 2014) with a minimum overlap of 50 bp and a maximum assembled sequence length of 450 bp. For *Alosa* sp., reverse reads quality was generally poor, therefore, only the forward read was used as its length (250 bp) encompasses the whole length of the sequenced loci (max. 200 bp). In some cases, microsatellite amplicons from several species were pooled prior to PCR indexing so that two individuals from two species shared an identical barcode combination and were sequenced in a single run. Forward primer sequences of species-specific loci were used to sort sequences belonging to different species into different fastq files using *fqgrep* tool (https://github.com/indraniel/fqgrep) allowing for one mismatch.

*Converting microsatellite sequences to genotypes*

We used the first three steps of the *FDSTools* v1.1.1 pipeline (Hoogenboom *et al.* 2016) to identify sequences corresponding to the microsatellite alleles and call genotypes for each individual (Figure 1). This analytical tool was chosen because it accounts for any kind of polymorphism detected across the analysed sequences (including variation in the number of repeated motifs, SNP or indels) while integrating specific tools to detect true allele from stutter mutation introduced during amplification that are typical of microsatellite markers.

In the first step (Figure 1), *tssv* (Anvar *et al.* 2014) matches primer sequences, allowing for 8% of mismatch, to identify sequences originating from each locus and count the occurrence of each unique sequence found for each locus for each individual. In the second step (Figure 1), *Stuttermark* uses the number of repeats of the microsatellite motif and the coverage of each unique sequence to flag unique sequences as potential allele, stutter resulting from slippage mutation during PCR and erroneous sequences (noise). In the third step (Figure 1), *Allelefinder* calls one or two alleles among the most abundant sequences flagged as potential alleles by *Stuttermark*. Following the FDSTools analysis, several custom-made *bash* routines were used to format the tabulated genotypic table, compare genotypes from repeated individuals to estimate locus specific allelic error rate based on the number of mismatches. In addition, allele level information was extracted including allele sequences, three-digit code used for genotype annotation, number of occurrence across individuals and allele length. Locus characteristics such as missing data rate and number of alleles are also summarized across the analysed individuals (Figure 1). All these bioinformatics steps have been embedded into a single bash script (*SSRseq_DataAnalysis_ParametersComparison.sh* available at https://doi.org/10.15454/HBXKVA) that allow to modify key analytical parameters (described below)

to evaluate their effect on the quality of the genotypic call (number of detected alleles, error and missing data rates).

Preliminary analyses showed that no single set of pipeline parameters resulted in good quality genotypic call across all locus. Instead, specific sets of parameters must be used to optimise genotypic call quality for each locus. However, testing numerous parameter combinations would rapidly become cumbersome and impractical. We thus restricted our analysis to two sets of parameters, called PS1 and PS2 thereafter. PS1 and PS2 were selected on the basis of high quality selection for locus showing moderate to high stutters and unbalanced allele coverage for heterozygote genotypes, because such patterns were typical of the loci developed here in non-model species (see Supporting Materials & Methods for full description). A single execution of the custom-made bash script (*SSRseq_DataAnalysis_ParametersComparison.sh* available at https://doi.org/10.15454/HBXKVA) run the analysis using the two parameter sets PS1 and PS2 for direct comparison of locus-specific genotypic call quality and selection of the most appropriate settings for each locus.

In addition, two analytical strategies were compared. In the first strategy (called *FullLength* thereafter), all variation identified between primers was considered as a haplotype irrespective of the nature of the polymorphism because all polymorphisms are physically-linked to each other in reads that encompass the whole locus. The *FullLenghth* strategy may be too complex for some loci or species showing high levels of polymorphism.  In the second strategy, the analysis was therefore restricted to the repeat motif only (strategy called *RepeatFocused* thereafter). In this case, primer and flanking sequences surrounding the repeated motif are indicated in the primer sequence field of the *FDSTools* input file. The *tssv* step then extracts the sequence corresponding to the repeat motif region (still allowing for 8% of mismatch which will accommodate flanking sequence polymorphism) to perform subsequent genotypic call with *Stuttermark* and *Allelefinder* as described above. As the analytical approach used by *FDSTools* is based on counting coverage of unique sequences, any variation identified within the repeated motif region, including variation in repeated motif number, SNP and indel within the motif region, will still be accounted for when defining alleles. While this *RepeatFocused* strategy may be more robust due to the shorter length of sequence analysed, it should identify a smaller number of alleles compared to the *FullLength* strategy.

*Comparing analytical approaches*

We compared genotypic call qualities between the four generated datasets (i.e. *FullLength_PS1*, *FullLength_PS2*, *RepeatFocused_PS1* and *RepeatFocused_PS2*). For each locus, we determined the best parameter set for each strategy and then the best overall analysis approach using the following

criteria by order of importance: estimated allelic error, amount of missing genotypes and number of detected alleles. Loci that showed more than 6% of allelic error or more than 50% of missing data across individuals were flagged as failed and removed from further inspection. For each species and analytical strategy, we recorded the number of genotyped loci, mean allelic error and missing data rate and the total number of alleles (haplotypes) across loci. We then determined the best overall approach for each locus and used it to genotype each locus generating a final *Combined* genotypic dataset for each species using a specific bash script (*SSRseq_DataAnalysis_FinalGenotyping.sh* available https://doi.org/10.15454/HBXKVA). Finally, the overall development success rate was computed for each species by dividing the number of reliable loci by the number of loci included in the multiplexed PCR.

**Gain from sequence information**

The number of identified alleles based on sequence information (haplotypes) was compared to the number of alleles differing in amplicon length only for all analysed loci to assess the gain of information obtained by using sequence data and estimate size homoplasy. We investigated further the nature of the detected polymorphism by counting, for each locus, the number of variations in the number of repeats, SNP and indels in the repeated motif and the flanking regions that differ between haplotypes.

# Results

**Sequence-based genotyping of previously developed microsatellites**

Attempts to genotype previously developed microsatellites in *S. salar* either from a new combination of 23 loci or a routinely-used multiplex of 15 loci resulted in a low number of reliable loci genotyped (Table 2). The overall success rate (the percentage of reliable loci over the number of loci amplified in the multiplexed PCR) ranged from 39% to 47% and the number of reliable loci from 7 to 10 (Table 2). Moreover, the quality of the generated genotypic datasets is relatively low with a rate of missing data and allelic error above 10% and 1% respectively (Figure 2). The sequences produced on the PGM Ion Torrent platform resulted in the lowest genotypic data quality (Figure 2). The same genotyping protocol sequenced on the Illumina MiSeq platform produced higher genotypic quality with lower missing data and allelic error rates (Figure 2) in spite of a 2.3 times lower mean coverage per sequenced locus per individual (Table 2). The Illumina MiSeq sequencing platform was thus used for subsequent analyses in other species.

Figure 2: Results of SSRsq development from previously developed microsatellites in *S. salar* for (a) a new multiplex of 23 microsatellite sequenced with Ion Torrent PGM and (b) Illumina MiSeq sequencing platforms, and (c) a routinely-used multiplex of 15 microsatellites sequenced with Illumina MiSeq sequencing platform. Number of reliable loci, total number of alleles, missing data and allelic error rates are indicated for three bioinformatics analysis strategies that focused either on all polymorphism across the sequence, on polymorphism within the repeated motif only, or a combination of the best strategy for each locus.

## Sequence-based genotyping of *de novo* developed microsatellites

In contrast, overall success rate of *de novo* microsatellite development ranged from 67% to 86% (Table 2). Given the high number of candidate microsatellites typically identified from high-throughput sequencing or reference genome sequence, we could afford to screen as much as 60 new loci, and combined most of them (from 28 to 60) in a single multiplexed PCR for amplification (Table 2). As a result, the final number of reliable loci was consistently high amounting to 24 for *Alosa* sp., 38 for *A. ostoyae*, 39 for *M. variegatipes* and 40 for *Quercus* sp. (Table 2). All these protocols produced high quality genotypic dataset, with low missing data (2.6% for *Quercus sp.*, 6.8% for *Alosa sp.*, 5.4% for *M. variegatipes*) at the exception of *A. ostoyae* (20.7%) and low allelic error rates as estimated based on blind-repeat genotyping (0.4% for *Quercus* sp., 0.6% for *Alosa* sp., 0.9% for *M. variegatipes* and 0.7% for *A. ostoyae*).

We explored the effect of sequence coverage on genotypic data quality on *Alosa* sp. by sequencing the same set of 28 microsatellites amplified in 156 individuals using one, two or three Illumina MiSeq nano flowcells (Table 2). The resulting increased coverage, from 95 to 198 and 267 sequences respectively per locus per individual (Table 2), recovers more data for those highly degraded DNA samples. First, increasing the coverage from 95 to 198 sequences by locus by individual detected three additional loci, while a further increase in coverage failed to recover additional locus (Table 2). Second, the missing data rate linearly decreases with the increase in coverage, from 16.4% to 9.0% and 6.8% with 95, 198 and 267 sequences per locus per individual respectively (Supporting Material Figure S1). It is worth noting that the allelic error rate is not affected by genome coverage, as it

varied only slightly between 0.5% and 0.7% without any correlation to coverage (Supporting Material Figure S1). A significant result is that even in conditions when only highly degraded DNA templates are available, reliable genotypic data can be obtained with moderate coverage, while increasing coverage will reduce missing data but not genotyping error rate.

**Whole sequence VS repeated motif polymorphism analysis**

Focusing the analysis on the repeated motif slightly increased the number of reliable loci and tends to produce fewer missing data and allelic errors (Figure 3). However, numerous polymorphisms that may be present in the flanking sequences are not accounted for. Indeed, analysing all polymorphism detected between the PCR primers resulted in a higher mean number of allele per locus, at the expense of slightly higher missing data and allelic error rates (Figure 3). Interestingly, 17% of the loci can be analysed reliably using either the *FullLength* or the *RepeatFocused* analytical approach. Thus combining analytical strategies by selecting the best approach for each locus resulted into an optimized dataset (Figure 3). Even for loci with reliable genotypes irrespectively of the analytical approach chosen, selecting the one that produces the best quality data (in terms of number of alleles, missing data and error rate) leads to an improved dataset quality. This combined strategy results in recovering the highest number of loci and alleles while keeping missing data and allelic error rates at the lowest (Figure 3).

Figure 3: Results of SSRseq development based on newly optimized microsatellites in (a) *Quercus* sp., (b) *Alosa* sp., (c) *A. ostoyae* and (d) *M. variegatipes* sequenced with Illumina MiSeq sequencing platform. Number of reliable loci, total number of alleles, missing data and allelic error rates are indicated for three bioinformatics analysis strategies that focused either on all polymorphism across the sequence, on polymorphism within the repeated motif only, or a combination of the best strategy for each locus.

**Types of polymorphism detected across species**

While most common population genetics applications do not necessitate to characterize the nature of the polymorphism differentiating alleles, the main advantage of sequence data (in addition to analysing a much higher number of loci) is to be able to identify allelic variation that does not translate into size variation, i.e. the only variation that is detected when using classical electrophoretic approaches. Across species, the proportion of alleles would have remained undetected by capillary electrophoresis (size homoplasy) ranging from 6% for *M. variegatipes*, 11% for *S. salar*, 14% for *Alosa* sp., 35% for *Quercus* sp. and 53% for *A. ostoyae* (Table 3). Conversely, the increase in the proportion of allele detected by accessing sequence data ranges from 6% for *M. variegatipes*, 13% for *S. salar*, 16% for *Alosa* sp., 56% for *Quercus* sp. and 113% for *A. ostoyae* (Table 3). Indeed, beside variation in repeat number, we identified numerous SNP and indel either in the flanking sequence or in the repeat motif itself (Figure 4, Table 3). In fact, additional polymorphism beyond variation in repeat number was the rule rather than the exception (Table 3). However, differences in the proportions of the type of the detected polymorphism were found between species (Figure 4). While repeat number variation represented more than 80% of the polymorphism detected in *S. salar*, *Quercus* sp., *Alosa* sp. and *M. variegatipes*, SNP in the flanking sequence was the most frequent polymorphism for *A. ostoyae* representing 49.8% of the variation, much high than variation of repeat number estimated at 29.8% (Figure 4). For *A. ostoyae*, SNP in the repeat motif and indel in the flanking sequence also represent a significant proportion of the polymorphism detected (12.6% and 6.5% respectively). *Quercus* sp. were characterised by SNP both within the repeat motif (7.5%) and the flanking region (9.3%). The second most comment polymorphism for *Alosa* sp. was SNP in the repeat motif (8.5%) and for *M. variegatipes* SNP in the flanking region (5.5%). For *S. salar*, variation in the number of motif was much more frequent than for other species (93.0%) compared to other polymorphisms that represent a marginal proportion of the variation (less than 3% each).

Figure 4: Proportion of detected polymorphism types within the repeat motif or in the flanking sequence for each sample per species group.

Table 3: Detected polymorphism.

| Species | Number of loci | Number of alleles with sequence difference[†] | Number of alleles differing by amplicon size[‡] | % of size homoplasy | % of increase in alleles due to sequence | Polymorphism in the repeat motif[§] | | | Polymorphism in the flanking sequences[§] | |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | | | | Repeat number variation | SNP | Indel | SNP | Indel |
| *S. salar*[¶] | 14 | 122 | 108 | 11% | 13% | 107 (14) | 3 (3) | - | 2 (2) | 3 (3) |
| *Quercus* sp. | 40 | 537 | 346 | 35% | 55% | 406 (40) | 38 (25) | 1 (1) | 47 (18) | 13 (10) |
| *Alosa* sp. | 24 | 174 | 150 | 14% | 16% | 130 (23) | 13 (15) | 2 (2) | 4 (4) | 3 (3) |
| *A. ostoyae* | 38 | 398 | 187 | 53% | 113% | 187 (33) | 79 (26) | 8 (7) | 312 (23) | 41 (16) |
| *M. variegatipes* | 39 | 166 | 156 | 6% | 6% | 147 (39) | 3 (3) | 1 (1) | 9 (7) | 5 (5) |

[†]: irrespectively of polymorphism type, computed based on the Combined analysis strategy; [‡]: simulating the number of alleles that would have been identified using traditional capillary electrophoresis, computed based on the FullLength analysis strategy and accounting for allele size only on the same locus as the Combined approach.
[§]: total number of alleles (and number of loci in brackets) for each polymorphism type. [¶]: combination of two sequence based microsatellite genotyping protocols.

## Discussion

While relying on previously developed microsatellite assay is far from optimal, our study demonstrates that primer redesign around known locus or *de novo* microsatellite development based on strict criteria gives successful single highly multiplexed PCR amplification and sequencing for 20 to 40 loci. However, critical upfront factors need to be considered for efficient SSRseq setup. The development workflow and analysis pipeline provided in this study are able to produce powerful multilocus haplotypic data in a time and cost effective manner in virtually any species.

**Not all roads lead to Rome: navigating pitfalls when adopting HTS for microsatellite genotyping**

Our preliminary attempts to apply SSRseq using previously developed microsatellite primers or capillary-based multiplexed microsatellite genotyping protocols clearly failed to produce reliable genotypic data for a sufficient number of loci. In the best case scenario, sequencing of 15 microsatellites in *Salmo* led to the reliable genotyping of 7 loci with 15.6% of missing data and 1.19% of allelic error, a result far worse than the high quality dataset obtained for the same multiplex using traditional capillary-electrophoresis of 14 loci with 0.5% missing data and 0.35% allelic error (Bacles *et al.* 2018). Similar trends were observed in previous studies that validated the use of HTS to genotype microsatellites and relied on previously developed primers: the generated datasets were characterised by high levels of missing data (up to 45%, (Vartia *et al.* 2016)) or low number of genotyped loci (7 to 8, (Suez *et al.* 2016; Barbian *et al.* 2018)). Two characteristics are problematic for SSRseq from microsatellites initially developed for capillary electrophoresis-based genotyping. First, high variability in locus length and primer characteristics leads to heterogeneous amplification intensity and sequencing coverage across loci. Indeed, efficient multiplexed PCR necessitates careful primer design using strict criteria (Guichoux et al., 2011; Lepais & Bacles, 2011). In addition, the need for variable locus length for optimal multiplexing without allele size range overlap in capillary-based electrophoresis becomes unnecessary for sequencing, because same size loci can be reliably identified simply based on primer sequences. Secondly, as sequencing errors accumulate proportionally to sequence length, lengthy sequences are more difficult to analyse, while complex or interrupted microsatellite motifs produce difficult to interpret stutter patterns. In that case, starting from a limited number of loci results in a handful of reliable loci that may be too low for downstream applications. This conclusion agrees well with a previous study developing SSRseq in *S. salar*, where only one out of six (17%) of previously developed loci was successfully integrated into the final panel, compared with a 26% success rate for newly developed primers (Bradbury *et al.* 2018). Adapting previously developed loci in the same species resulted in a success rate of about 45% in our case (7

out of 15 and 10 out of 23 reliable loci), but *de novo* development in other species was much more successful with a success rate of 75% on average. Not relying of previous primer design is thus of primary importance for the success of reliable SSRseq protocol development.

**Workflow for efficient SSRseq development in non-model species**

We propose here a workflow for developing new SSRseq approaches and we demonstrated its efficiency for a range of species with marked differences in polymorphism level. Starting from a reasonable number of candidate loci that were identified using readily available or newly generated genomic resources, resulted in an average of 75% of the loci included in the PCR multiplex generating reliable genotypic data. This result compares favourably with previous studies where success rates were lower (47% (Farrell *et al.* 2016), 53% (Neophytou *et al.* 2018)) or similar (78% (Tibihika *et al.* 2018)) when developing about 20 to 40 loci from a moderate number of candidate loci. However, extensive screening of numerous markers resulted in a much lower success rate: 101 validated loci from 385 tested (26% success rate (Bradbury *et al.* 2018)) or 43 validated loci from 448 locus tested (10% (Zhan *et al.* 2017)). In this study, we propose a balanced approach for cost-effective development that relies on simple amplification tests of a limited number of carefully selected candidate loci. Locus validation is then made when sequencing the final set of loci in a way that the validation step, based on the analyse of blind-repeat of at least 48 individuals, jointly generates useful genotypic data (for additional individuals included in the sequencing run). In addition, most of the ordered primers will be validated and integrated in the final set of reliable loci which greatly minimise development costs. The workflow is also flexible in terms of number of loci chosen for analysis: if a higher number of loci is required; additional sets of 60 loci could be selected from the candidate locus list and amplified in separated PCR multiplexes that can be pooled together before PCR indexing leading to a time and cost effective way to genotype additional loci (Bradbury *et al.* 2018). We did not test multiplexing more than 60 loci in a single PCR, but we did not see difficulties in doing so (Campbell *et al.* 2015). In such a situation, careful primer design with strict criteria and control for primer interactions will be key to increase multiplexed amplification success.

The proposed workflow for SSRseq development minimized laboratory steps and analytical optimizations. We chose to start from a moderate number of loci, designed to maximize their compatibility and sequence interpretability, and remove any locus that failed to amplify, produce interpretable or repeatable genotypes. This strategy avoids tedious optimization and reduce the number and complexity of laboratory steps necessary to produce genotypes. Admittedly, additional DNA or amplicon clean up or normalisation might improve sequence quality and coverage across individuals and loci. However, we chose to keep the laboratory procedure as simple as possible and

compensate the increase in amplification heterogeneity by additional sequence output resulting in sufficient coverage (220 reads/loci/individuals on average) to obtain a nearly complete genotypic dataset (generally 5% of missing data excluding the atypical case of *Armilaria* species). Moreover, increasing the number of laboratory steps inflates the risk of handling error and subsequently of genotyping error (Vartia *et al.* 2016). In this respect, highly multiplexed PCR are a useful technique to reduce laboratory steps and potential associated errors in addition to saving time and cutting costs. Additional tests not presented here showed that the two-stage multiplex amplification prior to PCR indexing  that was used to increase amplification homogeneity across loci is not necessary: high coverage is still efficient to compensate the increase in amplification heterogeneity when using a single multiplex PCR step for locus amplification as performed in previous studies (De Barba *et al.* 2016; Zhan *et al.* 2017; Bradbury *et al.* 2018; Tibihika *et al.* 2018).

We chose to rely on the set of 384 Illumina barcodes combinations because we found it to fit well with the output of the MiSeq sequencing platform when analysing from 20 loci to 300 loci depending on the type of flow cell used. However, studying more than 384 individuals from a single species necessitates either several MiSeq runs (Bradbury *et al.* 2018) or custom made dual-indexing strategies, as was successfully performed for the Ion Torrent PGM run (960 barcode combinations used) or in previous studies using the MiSeq platform (960 and 1024 barcode combinations (Farrell *et al.* 2016; Zhan *et al.* 2017)).

Finally, including repeated individuals is of paramount importance to assess the reliability of the produced genotypic data for each locus. This procedure aims to be best practice in capillary electrophoresis-based microsatellite genotyping (Hoffman and Amos 2005; Pompanon *et al.* 2005; Guichoux *et al.* 2011) but is even more important in SSRseq. Indeed, not all sequenced loci produced reliable genotypic data.  It is thus necessary to be able to identify and exclude loci producing high genotyping errors. At the bioinformatics analysis stage, we settled on testing a limited number of parameter combinations (four parameter sets) that were found to provide satisfactory results for most loci across the different species analysed. Selecting the best parameter set for each locus among the four parameter combinations is an easy task that greatly improve the number of reliable loci and decrease the genotyping error rate. However, a few loci will still show high genotyping error rate due to low coverage or complex polymorphism patterns and should be excluded from the final genotypic dataset. Exploratory analyses testing a much wider number of parameter combinations resulted in limited success in increasing the final number of reliable locus. Extensive parameter testing is a tedious task requiring high computation time with only minor improvement for the final genotypic dataset quality and is not worth the extra effort as long as a sufficient number of loci have been included in the genotyping panel. Furthermore, additional analyses using alternative

microsatellite sequences analysis tools such as Megasat (Zhan *et al.* 2017) and MicNeSs (Suez *et al.* 2016) also lead to the conclusion that all loci cannot be analysed reliably using a single set of parameters. Thus, we stress the importance to include a significant number of repeated individuals (at least 48) in the first analysis of a new SSRseq panel to (1) coarsely optimize analysis parameters for reliable loci, (2) quantify genotyping error rate and (3) identify and exclude unreliable loci. Such procedure has been implemented only in a limited number of previous studies describing SSRseq methods (5 out of 12 published studies thus far, (De Barba *et al.* 2016; Zhan *et al.* 2017; Bradbury *et al.* 2018; Barbian *et al.* 2018; Viruel *et al.* 2018)) but should be generalized.

**Implications of haplotype based genotyping**

We took advantage of the fact that reads span across entire loci to analyse all linked and phased polymorphisms encountered using the FDSTools pipeline (Hoogenboom *et al.* 2016). This haplotype approach differs from the methods implemented in other sequence-based microsatellite genotyping software such as Megasat (Zhan *et al.* 2017) which focuses on amplicon length or Micness (Suez *et al.* 2016) which estimates the number of repeated motifs while accounting for up to one substitution within the microsatellites motif. The haplotype approach has several advantages. Firstly, it is relatively insensitive to sequencing error because the analysis focused on unique sequence with high coverage and thus does not consider the noise generated by sequencing error which produce numerous unique low coverage sequences that are removed. Secondly, by analysing unique sequences, it accounts for any kind of polymorphisms, while at the same time includes an algorithm to identify stutters resulting from slippage mutation due to microsatellite instability during PCR amplification. Thirdly, it is flexible enough to be parameterized for analysis of the whole sequence or of the repeated motif only which is very useful to adapt the analysis to recalcitrant loci genotyping or to specific downstream applications.

Previous studies have demonstrated the added benefit of analysing different types of polymorphism within sequences compared to using amplicon size only (as in traditional capillary electrophoresis-based genotyping) to differentiate alleles (Darby *et al.* 2016; Neophytou *et al.* 2018; Barbian *et al.* 2018; Tibihika *et al.* 2018). Size homoplasy, due to alleles identical by size but not by sequence, ranges from to 32% and 44% (Darby *et al.* 2016; Vartia *et al.* 2016; Barbian *et al.* 2018). Here, we found high variability in size homoplasy ranging from 6% to 53% between the studied species in direct correlation to species polymorphism level and type of variation. SNP in the repeat motif or in the flanking region was the main source of size homoplasy which showed high prevalence in *A. ostoyae* (SNP represented 62.4% of the detected variation and 53% of size homoplasy) and to a lesser extent in *Quercus* sp. (SNP represented 16.8% of the detected variation and 35% of apparent

homoplasy). Even for species with lower apparent polymorphism levels, such as *M. variegatipes* (6%), *S. salar* (11%) and *Alosa* sp. (16%), the increase in the observed number of alleles will substantially improve genotypic resolution power (Darby *et al.* 2016; De Barba *et al.* 2016). It is important to note that not all SSRseq approaches maximize the information gained by the sequence information at the same level. If only allele size length is taken into account at the bioinformatics analysis step, conditions and thus results will be very similar to capillary electrophoresis-based genotyping (Zhan *et al.* 2017) with minimal information gain. Additional information will be obtained from analysing polymorphism within the repeat microsatellite motif, either by counting the number of repeated motifs including one substitution (Suez *et al.* 2016) or better still using the haplotype approach which is not constrained by the number of substitutions or the complexity of interrupted motifs. However, haplotype-based analysis that accounts for all linked variations across the whole sequence will make the most of the information available from sequence data (Barthe *et al.* 2012). Previous application of such an analytical approach used custom-made scripts implementing generalist bioinformatics tools and manual scoring of final genotypes (Darby *et al.* 2016; Neophytou *et al.* 2018) making them difficult to automate and transfer to other non- model systems (but see Barbian *et al.* 2018). Here, we integrated FDSTools algorithms into an automated analysis that includes parameters comparison and final genotyping into two pipelines based on bash script (https://doi.org/10.15454/HBXKVA), thus making it easy to estimate genotyping error based on repeat genotyping, to optimize parameters and identify reliable loci and produce the final multilocus genotypic table.

Finally, the ability to detect different sources of variation originating from several mutation mechanisms occurring at different rates provides renewed opportunities to study ecological and evolutionary events that occur at different timescales (Ramakrishnan and Mountain 2004; Barthe *et al.* 2012). Combining information on linked microsatellites and SNP (into a system called SNPSTR (Mountain *et al.* 2002) or HapSTR (Hey *et al.* 2004; Sorenson and Dacosta 2011)) was demonstrated to be a promising approach thanks to the increased phylogenetic resolution offered by explicitly considering complementary mutation properties of the markers. While theoretical and analytical implications of these approaches have been derived (Ramakrishnan and Mountain 2004; Hey *et al.* 2004; Payseur and Cutter 2006), empirical applications remain scarce and restricted to a very small number of systems due to the previous difficulties encountered to generate such empirical data (Mountain *et al.* 2002; Hey *et al.* 2004). This early limitation does not hold anymore with the generalisation of sequence-based microsatellite genotyping, as proposed herein, and the new ability to analyse linked microsatellites and SNP as haplotype. In addition, the flexibility of coalescent programs to simulate linked loci of different types (e.g. fastsimcoal2 (Excoffier *et al.* 2013)) will authorize far more realistic simulation of mutation mechanisms specifically tailored to each marker

system. Such improvement, especially when applied to tens of marker systems, would make simulation-based inference (Beaumont *et al.* 2002) more accurate over an extended timescale range even for complex evolutionary history scenarios (Mountain *et al.* 2002).

In conclusion, this study proposes an integrated approach to expedite the development of SSRseq protocol for any species. We found it to be time efficient and cost effective for a range of species across Kingdoms. Building on our experience, we provided several recommendations to improve development efficiency. The two most important advices are to optimize marker selection and primer design for effective multiplex PCR amplification and sequence interpretability, and to use repeated individuals to assess the quality of the generated genotypic data. The automated genotyping pipeline developed in this study is flexible enough to optimize analysis parameters and improve genotypic data quality, and rapidly process sequences to estimate locus-level statistics (including missing data and genotyping error rates). Furthermore, it generates formatted genotypic tables ready for downstream data analysis from sequence information ensuring easy data interoperability (correspondence between allele code and allele sequence). As our method is based on simple bioinformatics algorithms, hundreds of individuals can be genotyped at tens of markers within a couple of hours on any standard laptop running on a Linux operating system. The ability of SSRseq to characterize SNP and indel present along the sequences, in addition to the targeted microsatellite, represents a new opportunity to produce empirical data to apply existing theoretical and statistical frameworks that integrate linked polymorphism with different mutation characteristics (Payseur and Cutter 2006). Finally, the ease and cost effectiveness of parallel development for multiple species make these approach convenient to develop powerful multilocus datasets for comparative population and community genetics studies (Crutsinger 2016), and to further investigate the functional implications (Bagshaw 2017) and adaptive potential of microsatellite variation among natural populations (Xie *et al.* 2019).

**Acknowledgments**

This research was supported by grants to OL of the Agence de l'Eau Adour-Garonne (project 2017/3273), the Région Nouvelle-Aquitaine (project 2016-1R20602-00007239), the Agence Française pour la Biodiversité (project 2016–18 A13) and INRA. A. A. benefited from a travel grant from the Department of Agronomy ISVSA of the University Batna 1 Hadja Lakhdar. We thank François Meurgey for the sampling of *Melipona sp.* individuals. Technical developments and sequencing were performed at the Genome Transcriptome Facility of Bordeaux (Grants from Investissements d'Avenir, Convention attributive d'aide EquipEx Xyloforest ANR-10-EQPX-16-01).

**References:**

Anvar S. Y., K. J. Van Der Gaag, J. W. F. Van Der Heijden, M. H. A. M. Veltrop, R. H. A. M. Vossen, *et al.*, 2014 TSSV: A tool for characterization of complex allelic variants in pure and mixed genomes. Bioinformatics 30: 1651–1659. https://doi.org/10.1093/bioinformatics/btu068

Aykanat T., M. Lindqvist, V. L. Pritchard, and C. R. Primmer, 2016 From population genomics to conservation and management: A workflow for targeted analysis of markers identified using genome-wide approaches in Atlantic salmon Salmo salar. J. Fish Biol. 1–22. https://doi.org/10.1111/jfb.13149

Bacles C. F. E., C. Bouchard, F. Lange, A. Manicki, C. Tentelier, *et al.*, 2018 Estimating the effective number of breeders from single parr samples for conservation monitoring of wild populations of Atlantic salmon Salmo salar. J. Fish Biol. 92: 699–726. https://doi.org/10.1111/jfb.13537

Bagshaw A. T. M., 2017 Functional mechanisms of microsatellite DNA in eukaryotic genomes. Genome Biol. Evol. 9: 2428–2443. https://doi.org/10.1093/gbe/evx164

Baird N. A., P. D. Etter, T. S. Atwood, M. C. Currey, A. L. Shiver, *et al.*, 2008 Rapid SNP discovery and genetic mapping using sequenced RAD markers. PLoS One 3: e3376. https://doi.org/10.1371/journal.pone.0003376

Barba M. De, C. Miquel, S. Lobréaux, P. Y. Quenette, J. E. Swenson, *et al.*, 2016 High-throughput microsatellite genotyping in ecology: Improved accuracy, efficiency, standardization and success with low-quantity and degraded DNA. Mol. Ecol. Resour. 17: 492–507. https://doi.org/10.1111/1755-0998.12594

Barbian H. J., A. J. Connell, A. N. Avitto, R. M. Russell, A. G. Smith, *et al.*, 2018 CHIIMP: An automated high-throughput microsatellite genotyping platform reveals greater allelic diversity in wild chimpanzees. Ecol. Evol. 8: 7946–7963. https://doi.org/10.1002/ece3.4302

Barthe S., F. Gugerli, N. A. Barkley, L. Maggia, C. Cardi, *et al.*, 2012 Always look on both sides: phylogenetic information conveyed by simple sequence repeat allele sequences. PLoS One 7: e40699. https://doi.org/10.1371/journal.pone.0040699

Beaumont M. A., W. Zhang, and D. J. Balding, 2002 Approximate bayesian computation in population genetics. Genetics 162: 2025–2035.

Blacket M. J., C. Robin, R. T. Good, S. F. Lee, and A. D. Miller, 2012 Universal primers for fluorescent labelling of PCR fragments – an efficient and cost-effective approach to genotyping by fluorescence. Mol. Ecol. 12: 456–463. https://doi.org/10.1111/j.1755-0998.2011.03104.x

Bradbury I. R., B. F. Wringe, B. Watson, I. Paterson, J. Horne, *et al.*, 2018 Genotyping-by-sequencing of genome-wide microsatellite loci reveals fine-scale harvest composition in a coastal Atlantic salmon fishery. Evol. Appl. 11: 918–930. https://doi.org/10.1111/eva.12606

Brown S. S., Y.-W. Chen, M. Wang, A. Clipson, E. Ochoa, *et al.*, 2017 PrimerPooler: automated primer pooling to prepare library for targeted sequencing. Biol. Methods Protoc. 2: 1–10. https://doi.org/10.1093/biomethods/bpx006

Campbell N. R., S. Harmon, and S. R. Narum, 2015 Genotyping-in-Thousands by sequencing (GT-seq): A cost effective SNP genotyping method based on custom amplicon sequencing. Mol. Ecol. Resour. 15: 855–867. https://doi.org/10.1111/1755-0998.12357

Castoe T. A., A. W. Poole, W. Gu, A. P. Jason de Koning, J. M. Daza, *et al.*, 2010 Rapid identification of

thousands of copperhead snake (Agkistrodon contortrix) microsatellite loci from modest amounts of 454 shotgun genome sequence. Mol. Ecol. Resour. 10: 341–347. https://doi.org/10.1111/j.1755-0998.2009.02750.x

Chen K., Y. xun Zhou, K. Li, L. xin Qi, Q. fei Zhang, *et al.*, 2016 A novel three-round multiplex PCR for SNP genotyping with next generation sequencing. Anal. Bioanal. Chem. 1–7. https://doi.org/10.1007/s00216-016-9536-6

Crutsinger G. M., 2016 A community genetics perspective: Opportunities for the coming decade. New Phytol. 210: 65–70. https://doi.org/10.1111/nph.13537

Darby B. J., S. F. Erickson, S. D. Hervey, and S. N. Ellis-Felege, 2016 Digital fragment analysis of short tandem repeats by high-throughput amplicon sequencing. Ecol. Evol. 6: 4502–4512. https://doi.org/10.1002/ece3.2221

Delord C., G. Lassalle, A. Oger, D. Barloy, M.-A. Coutellec, *et al.*, 2018 A cost-and-time effective procedure to develop SNP markers for multiple species: A support for community genetics, (M. Bunce, Ed.). Methods Ecol. Evol. 9: 1959–1974. https://doi.org/10.1111/2041-210X.13034

Durand J., C. Bodenes, E. Chancerel, J. M. Frigerio, G. Vendramin, *et al.*, 2010 A fast and cost-effective approach to develop and map EST-SSR markers: oak as a case study. BMC Genomics 11. https://doi.org/10.1186/1471-2164-11-570

Ellis J. S., J. Gilbey, A. Armstrong, T. Balstad, E. Cauwelier, *et al.*, 2011 Microsatellite standardization and evaluation of genotyping error in a large multi-partner research programme for conservation of Atlantic salmon (Salmo salar L.). Genetica 139: 353–367. https://doi.org/10.1007/s10709-011-9554-4

Estoup A., P. Jarne, and J.-M. Cornuet, 2002 Homoplasy and mutation model at microsatellite loci and their consequences for population genetics analysis. Mol. Ecol. 11: 1591–1604. https://doi.org/10.1046/j.1365-294X.2002.01576.x

Excoffier L., I. Dupanloup, E. Huerta-Sánchez, V. C. Sousa, and M. Foll, 2013 Robust demographic inference from genomic and SNP data. PLoS Genet. 9: e1003905. https://doi.org/10.1371/journal.pgen.1003905

Farrell E. D., J. E. L. Carlsson, J. Carlsson, and E. D. Farrell, 2016 Next Gen Pop Gen : implementing a high-throughput approach to population genetics in boarfish (Capros aper). R. Soc. Open Sci. 3: 160651. https://doi.org/http://dx.doi.org/10.1098/rsos.160651

Garvin M. R., K. Saitoh, and A. J. Gharrett, 2010 Application of single nucleotide polymorphisms to non-model species: a technical review. Mol. Ecol. Resour. 10: 915–934. https://doi.org/10.1111/j.1755-0998.2010.02891.x

Gauthey Z., M. Freychet, A. Manicki, A. Herman, O. Lepais, *et al.*, 2015 The concentration of plasma metabolites varies throughout reproduction and affects offspring number in wild brown trout (Salmo trutta). Comp. Biochem. Physiol. A. Mol. Integr. Physiol. 184: 90–96. https://doi.org/10.1016/j.cbpa.2015.01.025

Gilbey J., E. Verspoor, A. McLay, and D. Houlihan, 2004 A microsatellite linkage map for Atlantic salmon ( *Salmo salar* ). Anim. Genet. 35: 98–105. https://doi.org/10.1111/j.1365-2052.2004.01091.x

Guichoux E., L. Lagache, S. Wagner, P. Chaumeil, P. Léger, *et al.*, 2011 Current trends in microsatellite genotyping. Mol. Ecol. Resour. 11: 591–611. https://doi.org/10.1111/j.1755-0998.2011.03014.x

Gymrek M., 2017 A genomic view of short tandem repeats. Curr. Opin. Genet. Dev. 44: 9–16.

https://doi.org/10.1016/j.gde.2017.01.012

Haasl R. J., and B. A. Payseur, 2011 Multi-locus inference of population structure: a comparison between single nucleotide polymorphisms and microsatellites. Heredity (Edinb). 106: 158–171. https://doi.org/10.1038/hdy.2010.21

Hannan A. J., 2018 Tandem repeats mediating genetic plasticity in health and disease. Nat. Rev. Genet. 19: 286–298. https://doi.org/10.1038/nrg.2017.115

Harrison H. B., P. Saenz-Agudelo, S. Planes, G. P. Jones, and M. L. Berumen, 2013 Relative accuracy of three common methods of parentage analysis in natural populations. Mol. Ecol. 22: 1158–1170. https://doi.org/10.1111/mec.12138

Hey J., Y.-J. Won, A. Sivasundar, R. Nielsen, and J. A. Markert, 2004 Using nuclear haplotypes with microsatellites to study gene flow between recently separated Cichlid species. Mol. Ecol. 13: 909–919. https://doi.org/10.1046/j.1365-294X.2003.02031.x

Hoffman J. I., and W. Amos, 2005 Microsatellite genotyping errors: detection approaches, common sources and consequences for paternal exclusion. Mol. Ecol. 14: 599–612. https://doi.org/10.1111/j.1365-294X.2004.02419.x

Holleley C. E., and P. G. Geerts, 2009 Multiplex Manager 1.0: a cross-platform computer program that plans and optimizes multiplex PCR. Biotechniques 46: 511–517. https://doi.org/10.2144/000113156

Hoogenboom J., P. de Knijff, J. F. J. Laros, R. H. de Leeuw, K. J. van der Gaag, *et al.*, 2016 FDSTools: A software package for analysis of massively parallel sequencing data with the ability to recognise and correct STR stutter and other PCR or sequencing noise. Forensic Sci. Int. Genet. 27: 27–40. https://doi.org/10.1016/j.fsigen.2016.11.007

Kampfer S., C. Lexer, J. Glössl, and H. Steinkellner, 1998 Characterization of (GA)n microsatellite loci from Quercus robur. Hereditas 129: 183–186. https://doi.org/10.1111/j.1601-5223.1998.00183.x

King T. L., M. S. Eackles, and B. H. Letcher, 2005 Microsatellite DNA markers for the study of Atlantic salmon (Salmo salar) kinship, population structure, and mixed-fishery analyses. Mol. Ecol. Notes 5: 130–132. https://doi.org/10.1111/j.1471-8286.2005.00860.x

Langmead B., and S. L. Salzberg, 2012 Fast gapped-read alignment with Bowtie 2. Nat. Methods 9: 357–359. https://doi.org/10.1038/nmeth.1923

Lepais O., and C. F. E. Bacles, 2011 Comparison of random and SSR-enriched shotgun pyrosequencing for microsatellite discovery and single multiplex PCR optimization in Acacia harpophylla F. Muell. Ex Benth. Mol. Ecol. Resour. 711–724. https://doi.org/10.1111/j.1755-0998.2011.03002.x

Lepais O., A. Manicki, S. Glise, M. Buoro, and A. Bardonnet, 2017 Genetic architecture of threshold reaction norms for male alternative reproductive tactics in Atlantic salmon (Salmo salar L.). Sci. Rep. 7: 43552. https://doi.org/10.1038/srep43552

Malausa T., A. Gilles, E. Meglécz, H. Blanquart, S. Duthoy, *et al.*, 2011 High-throughput microsatellite isolation through 454 GS-FLX Titanium pyrosequencing of enriched DNA libraries. Mol. Ecol. Resour. 11: 638–644. https://doi.org/10.1111/j.1755-0998.2011.02992.x

Margulies M., M. Egholm, W. E. Altman, S. Attiya, J. S. Bader, *et al.*, 2005 Genome sequencing in microfabricated high-density picolitre reactors. Nature 437: 376–380. https://doi.org/10.1038/nature03959

Martin M., 2011 Cutadapt removes adapter sequences from high-throughput sequencing reads. EMBnet.journal 17: 10. https://doi.org/10.14806/ej.17.1.200

Meglécz E., C. Costedoat, V. Dubut, A. Gilles, T. Malausa, *et al.*, 2010 QDD: a user-friendly program to select microsatellite markers and design primers from large sequencing projects. Bioinformatics 26: 403–4. https://doi.org/10.1093/bioinformatics/btp670

Meglécz E., N. Pech, A. Gilles, V. Dubut, P. Hingamp, *et al.*, 2014 QDD version 3.1: a user-friendly computer program for microsatellite selection and primer design revisited: experimental validation of variables determining genotyping success rate. Mol. Ecol. Resour. 14: 1302–1313. https://doi.org/10.1111/1755-0998.12271

Miller M. R., J. P. Dunham, A. Amores, W. A. Cresko, and E. A. Johnson, 2007 Rapid and cost-effective polymorphism identification and genotyping using restriction site associated DNA (RAD) markers. Genome Res. 17: 240–8. https://doi.org/10.1101/gr.5681207

Moran P., D. J. Teel, E. S. LaHood, J. Drake, and S. Kalinowski, 2006 Standardising multi-laboratory microsatellite data in Pacific salmon: an historical view of the future. Ecol. Freshw. Fish 15: 597–605. https://doi.org/10.1111/j.1600-0633.2006.00201.x

Mountain J. L., A. Knight, M. Jobin, C. Gignoux, A. Miller, *et al.*, 2002 SNPSTRs: Empirically derived, rapidly typed, autosomal haplotypes for inference of population history and mutational processes. Genome Res. 12: 1766–1772. https://doi.org/10.1101/gr.238602

Neophytou C., E. Torutaeva, S. Winter, H. Meimberg, H. Hasenauer, *et al.*, 2018 Analysis of microsatellite loci in tree of heaven (Ailanthus altissima (Mill.) Swingle) using SSR-GBS. Tree Genet. Genomes 14. https://doi.org/10.1007/s11295-018-1295-4

Nguyen-Dumont T., B. J. Pope, F. Hammet, M. C. Southey, and D. J. Park, 2013 A high-plex PCR approach for massively parallel sequencing. Biotechniques 55: 69–74. https://doi.org/10.2144/000114052

O'Reilly P., and J. M. Wright, 1995 The evolving technology of DNA fingerprinting and its application to fisheries and aquaculture. J. Fish Biol. 47: 29–55. https://doi.org/10.1111/j.1095-8649.1995.tb06042.x

Ozaki A., T. Sakamoto, S. Khoo, K. Nakamura, M. R. M. Coimbra, *et al.*, 2001 Quantitative trait loci (QTLs) associated with resistance/susceptibility to infectious pancreatic necrosis virus (IPNV) in rainbow trout (Oncorhynchus mykiss). Mol. Genet. Genomics 265: 23–31. https://doi.org/10.1007/s004380000392

Paterson S., S. B. Piertney, D. Knox, J. Gilbey, and E. Verspoor, 2004 Characterization and PCR multiplexing of novel highly variable tetranucleotide Atlantic salmon (Salmo salar L.) microsatellites. Mol. Ecol. Notes 4: 160–162. https://doi.org/10.1111/j.1471-8286.2004.00598.x

Payseur B. A., and A. D. Cutter, 2006 Integrating patterns of polymorphism at SNPs and STRs. Trends Genet. 22: 424–429. https://doi.org/10.1016/j.tig.2006.06.009

Pimentel J. S. M., A. O. Carmo, I. C. Rosse, A. P. V. Martins, S. Ludwig, *et al.*, 2018 High-throughput sequencing strategy for microsatellite genotyping using neotropical fish as a model. Front. Genet. 9: 73. https://doi.org/10.3389/fgene.2018.00073

Plomion C., J. M. Aury, J. Amselem, T. Leroy, F. Murat, *et al.*, 2018 Oak genome reveals facets of long lifespan. Nat. Plants 4: 440–452. https://doi.org/10.1038/s41477-018-0172-3

Pompanon F., A. Bonin, E. Bellemain, and P. Taberlet, 2005 Genotyping errors: causes, consequences and solutions. Nat. Rev. Genet. 6: 847–59. https://doi.org/10.1038/nrg1707

Prospero S., B. Lung-Escarmant, and C. Dutech, 2008 Genetic structure of an expanding Armillaria root rot fungus (Armillaria ostoyae) population in a managed pine forest in southwestern France. Mol. Ecol. 17: 3366–3378. https://doi.org/10.1111/j.1365-294X.2007.03829.x

Quinlan A. R., 2014 BEDTools: The Swiss-Army Tool for Genome Feature Analysis. Curr. Protoc. Bioinforma. 47: 11.12.1-11.12.34. https://doi.org/10.1002/0471250953.bi1112s47

Ramakrishnan U., and J. L. Mountain, 2004 Precision and accuracy of divergence time estimates from STR and SNPSTR variation. Mol. Biol. Evol. 21: 1960–1971. https://doi.org/10.1093/molbev/msh212

Rexroad C. E., R. L. Coleman, A. M. Martin, W. K. Hershberger, and J. Killefer, 2001 Thirty-five polymorphic microsatellite markers for rainbow trout (Oncorhynchus mykiss). Anim. Genet. 32: 317–319. https://doi.org/10.1046/j.1365-2052.2001.0730b.x

Rougemont Q., A. L. Besnard, J. L. Baglinière, and S. Launey, 2015 Characterization of thirteen new microsatellite markers for allis shad (Alosa alosa) and twaite shad (Alosa fallax). Conserv. Genet. Resour. 7: 259–261. https://doi.org/10.1007/s12686-014-0352-z

Sadd B. M., S. Schaack, X. Zhao, C. Sun, and L. Su, 2018 Tandem repeats contribute to coding sequence variation in bumblebees (Hymenoptera: Apidae). Genome Biol. Evol. 10: 3176–3187. https://doi.org/10.1093/gbe/evy244

Schloss P. D., S. L. Westcott, T. Ryabin, J. R. Hall, M. Hartmann, *et al.*, 2009 Introducing mothur: open-source, platform-independent, community-supported software for describing and comparing microbial communities. Appl. Environ. Microbiol. 75: 7537–41. https://doi.org/10.1128/AEM.01541-09

Schlötterer C., 2004 The evolution of molecular markers — just a matter of fashion? Nat. Rev. Genet. 5: 63–69. https://doi.org/10.1038/nrg1249

Seeb J. E., G. Carvalho, L. Hauser, K. Naish, S. Roberts, *et al.*, 2011 Single-nucleotide polymorphism (SNP) discovery and applications of SNP genotyping in nonmodel organisms. Mol. Ecol. Resour. 11: 1–8. https://doi.org/10.1111/j.1755-0998.2010.02979.x

Sipos G., A. N. Prasanna, M. C. Walter, E. O'Connor, B. Bálint, *et al.*, 2017 Genome expansion and lineage-specific genetic innovations in the forest pathogenic fungi Armillaria. Nat. Ecol. Evol. 1: 1931–1941. https://doi.org/10.1038/s41559-017-0347-8

Slettan A., I. Olsaker, and Ø. Lie, 1996 Polymorphic Atlantic salmon, Salmo salar L., microsatellites at the SSOSL438, SSOSL439 and SSOSL444 loci. Anim. Genet. 27: 57–58. https://doi.org/10.1111/j.1365-2052.1996.tb01180.x

Sorenson M. D., and J. M. Dacosta, 2011 Genotyping HapSTR loci: phase determination from direct sequencing of PCR products. Mol. Ecol. Resour. 11: 1068–1075. https://doi.org/10.1111/j.1755-0998.2011.03036.x

Steinkellner H., S. Fluch, E. Turetschek, C. Lexer, R. Streiff, *et al.*, 1997 Identification and characterization of (GA/CT)n microsatellite loci from Quercus petraea. Plant Mol. Biol. 33: 1093–1096. https://doi.org/10.1023/A:1005736722794

Suez M., A. Behdenna, S. Brouillet, P. Graça, D. Higuet, *et al.*, 2016 MicNeSs: Genotyping microsatellite loci from a collection of (NGS) reads. Mol. Ecol. Resour. 16: 524–533. https://doi.org/10.1111/1755-0998.12467

Thorsen J., B. Zhu, E. Frengen, K. Osoegawa, P. J. de Jong, *et al.*, 2005 A highly redundant BAC library of Atlantic salmon (Salmo salar): an important tool for salmon projects. BMC Genomics 6: 50.

https://doi.org/10.1186/1471-2164-6-50

Tibihika P. D., M. Curto, E. Dornstauder-Schrammel, S. Winter, E. Alemayehu, *et al.*, 2018 Application of microsatellite genotyping by sequencing (SSR-GBS) to measure genetic diversity of the East African Oreochromis niloticus. Conserv. Genet. 1–16. https://doi.org/10.1007/s10592-018-1136-x

Vartia S., J. L. Villanueva-Cañas, J. Finarelli, E. D. Farrell, P. C. Collins, *et al.*, 2016 A novel method of microsatellite genotyping-by-sequencing using individual combinatorial barcoding. R. Soc. Open Sci. 3: 150565. https://doi.org/10.1098/rsos.150565

Vasemägi A., J. Nilsson, and C. R. Primmer, 2005 Expressed sequence tag-linked microsatellites as a source of gene-associated polymorphisms for detecting signatures of divergent selection in Atlantic salmon (Salmo salar L.). Mol. Biol. Evol. 22: 1067–76. https://doi.org/10.1093/molbev/msi093

Vasemägi A., R. Gross, D. Palm, T. Paaver, C. R. Primmer, *et al.*, 2010 Discovery and application of insertion-deletion (INDEL) polymorphisms for QTL mapping of early life-history traits in Atlantic salmon. BMC Genomics 11: 156. https://doi.org/10.1186/1471-2164-11-156

Viard F., P. Franck, M. P. Dubois, A. Estoup, and P. Jarne, 1998 Variation of microsatellite size homoplasy across electromorphs, loci, and populations in three invertebrate species. J. Mol. Evol. 47: 42–51.

Viruel J., A. Haguenauer, M. Juin, F. Mirleau, D. Bouteiller, *et al.*, 2018 Advances in genotyping microsatellite markers through sequencing and consequences of scoring methods for Ceratonia siliqua (Leguminosae). Appl. Plant Sci. 6: e01201. https://doi.org/10.1002/aps3.1201

Willems T., M. Gymrek, G. Highnam, T. 1000 G. P. 1000 Genomes Project Consortium, D. Mittelman, *et al.*, 2014 The landscape of human STR variation. Genome Res. 24: 1894–904. https://doi.org/10.1101/gr.177774.114

Xie K. T., G. Wang, A. C. Thompson, J. I. Wucherpfennig, T. E. Reimchen, *et al.*, 2019 DNA fragility in the parallel evolution of pelvic reduction in stickleback fish. Science (80-. ). 363: 81–84. https://doi.org/10.1126/science.aan1425

Yano A., B. Nicol, E. Jouanno, E. Quillet, A. Fostier, *et al.*, 2013 The sexually dimorphic on the Y-chromosome gene (sdY) is a conserved male-specific Y-chromosome sequence in many salmonids. Evol. Appl. 6: 486–496. https://doi.org/10.1111/eva.12032

Zhan L., I. G. Paterson, B. A. Fraser, B. Watson, I. R. Bradbury, *et al.*, 2017 megasat: Automated inference of microsatellite genotypes from sequence data. Mol. Ecol. Resour. 17: 247–256. https://doi.org/10.1111/1755-0998.12561

Zhang J., K. Kobert, T. Flouri, and A. Stamatakis, 2014 PEAR: A fast and accurate Illumina Paired-End reAd mergeR. Bioinformatics 30: 614–620. https://doi.org/10.1093/bioinformatics/btt593

**Data Accessibility**

- DNA sequences: sequence reads from SSRseq run were submitted to the European Nucleotide Archive SRA under the accession number PRJEB31908 (*Alosa* sp.), PRJEB31909 (*A. ostoyae*), PRJEB31910 (*M. variegatipes*), PRJEB31913 (*Quercus* sp.) and PRJEB31914 (*S. salar*). Random shotgun sequencing reads from one *M. variegatipes* individual for microsatellite discovery was submitted under the accession number ERR3255838.

- Pipeline scripts for automated sequence-based microsatellite genotyping with documentation and example files are available at Data Inra https://doi.org/10.15454/HBXKVA.

**Authors Contributions**

OL designed the study and developed the data analysis pipeline. OL, EC, CB, FS and EG developed the laboratory protocols. AM, LT, AA and CFEB contributed to laboratory analyses. EC, LT and CFEB contributed to data analysis. CD, FD and SL contributed to sampling and preliminary data collection. OL wrote the manuscript which was revised critically and approved by all authors.

**Supporting / Supplemental information**

Table S1: Full details about the loci used or developed in this study (including species, locus name, primer sequences, characteristics, and reference)

Figure S1: Effect of sequence coverage on SSRseq data quality for *Alosa* species.

Supporting Materials and Methods: detailed explanation of FDSTools parameters used in this study.