The advantages and disadvantages of short- and long-read metagenomics to infer bacterial and eukaryotic community composition

Keywords: metagenomics, Nanopore, Illumina, long read, community composition

William S. Pearman[1], Nikki E. Freed[1], Olin K. Silander[1]

1 School of Natural and Computational Sciences, Massey University, Auckland, New Zealand

Corresponding Authors:

William S Pearman

Olin K Silander

Private Bag 102904, North Shore, 0745 Auckland, New Zealand

Email addresses: wpearman1996@gmail.com; olinsilander@gmail.com

Abstract

Background

The first step in understanding ecological community diversity and dynamics is quantifying community membership. An increasingly common method for doing so is through metagenomics. Because of the rapidly increasing popularity of this approach, a large number of computational tools and pipelines are available for analysing metagenomic data. However, the majority of these tools have been designed and benchmarked using highly accurate short read data (i.e. illumina), with few studies benchmarking classification accuracy for long error-prone reads (PacBio or Oxford Nanopore). In addition, few tools have been benchmarked for non-microbial communities.

Results

Here we use simulated error prone Oxford Nanopore and high accuracy Illumina read sets to systematically investigate the effects of sequence length and taxon type on classification accuracy for metagenomic data from both microbial and non-microbial communities. We show that very generally, classification accuracy is far lower for non-microbial communities, even at low taxonomic resolution (e.g. family rather than genus).

Conclusions

We then show that for two popular taxonomic classifiers, long error-prone reads can significantly increase classification accuracy, and this is most pronounced for non-microbial communities. This work provides insight on the expected accuracy for metagenomic analyses for different taxonomic groups, and establishes the point at which read length becomes more important than error rate for assigning the correct taxon.

Introduction

Applying Metagenomic Methods to Quantify Community Composition

To understand ecological community diversity, it is essential to quantify taxon frequency. The most common method of quantifying taxa frequencies is through metabarcoding (Ji et al. 2013). In this method, conserved genomic regions (often 16S rRNA in the case of bacterial and archaeal species; 18S rRNA or Cytochrome c oxidase I for eukaryotic species) are amplified from the sample of interest, sequenced (most often using high-throughput methods such as Illumina), and then classified using one of several available pipelines (e.g. QIIME, MEGAN, Mothur) (Caporaso et al. 2010; Huson et al. 2016; Schloss et al. 2009). Many of these pipelines have been designed around the analysis of bacterial datasets.

In contrast to metabarcoding, metagenomic approaches do not rely on the amplification of specific genomic sequences, which can introduce bias. Instead, they aim to quantify community composition based on the recovery and sequencing of all DNA from community samples. Not only do metagenomic methods profile taxon composition in a less biased way than metabarcoding, but they can also yield insight into the functional diversity present in ecosystems (Schloss and Handelsman 2005; Keeling et al. 2014).

While metabarcoding approaches have been widely applied to both microbial and eukaryotic taxa, the vast majority of metagenomic studies have focused only on microbial communities. Unsurprisingly, the various advantages and disadvantages of using metagenomic analyses for microbial communities are well-documented (Roumpeka et al. 2017; Thomas, Gilbert, and Meyer 2012; Temperton and Giovannoni 2012). There are likely several factors driving this microbe-centric application of metagenomics, including (1) the greater level of diversity of microbial taxa; (2) the considerable number of microbial taxa that are "unculturable," making it difficult to collect the requisite amount of DNA for genomic sequencing; (3) the availability of a multitude of non-molecular methods for quantifying multicellular taxa; and (4) the relative paucity of genomic sequence for multicellular organisms in databases (Escobar-Zepeda, Vera-Ponce de León, and Sanchez-Flores 2015) (Supp. Fig.1). This latter factor is perhaps the single largest factor in driving the bias toward microbial metagenomics.

However, the amount and diversity of eukaryotic genomic sequence data is rapidly

increasing. Although multicellular metabarcode databases are currently far more complete

relative to genomic databases, this gap is closing quickly. For example, the Earth

BioGenome project aims to sequence the genomes of upwards of one million eukaryotic

species within the next decade (Lewin et al. 2018). Regardless of the success of this effort,

there are a host of ongoing eukaryotic sequencing projects, including Bat 1K (Teeling et al.

2018), Bird 10K (10,000 bird genomes (OBrien, Haussler, and Ryder 2014)), G10K (10,000

vertebrate genomes (10K Community of Scientists 2009)), and i5K (5000 arthropod

genomes (Robinson et al. 2011)), among others. This suggests that within the next five

years, most multicellular organisms will have at least one member of their family present in

genomic databases, with some groups of multicellular organisms being completely

represented at the genus level.

This would increase the utility of metagenomics for assessing membership in plant and

animal communities, especially for cases in which organisms are difficult to observe or

degraded. This is frequently the case for diet studies (Pearman et al. 2018), many

invertebrate communities such as in treeholes (Gossner et al. 2016) or algal holdfasts

(Ojeda and Santelices 1984).

81 Analysis of Short-read Metagenomic Data

Many metagenomic classification analyses rely on first pass classifiers to assign reads to

one or more taxa, followed by second pass classifiers that can improve on the initial

classification by taking into account the number and relationship of taxa identified in the first

pass. This second step often relies on a lowest common ancestor algorithm (Wood and

Salzberg 2014; Kim et al. 2016; Huson et al. 2016), or by refining taxonomic representation

by examining the results from the first pass classifier (Lu et al. 2016).

The most widely used first pass classifier is BLAST, and it is considered gold standard

(McIntyre et al. 2017). However, BLAST is not computationally efficient enough to deal with

tens or hundreds of millions of reads. Thus, algorithms for fast metagenomic classification

have been the subject of intense research over the last few years, and include k-mer based

approaches such as CLARK (Ounit et al. 2015), Kraken and related tools (Kraken, Kraken2,

and KrakenUniq) (Wood and Salzberg 2014), Centrifuge (Kim et al. 2016), EnSVMB (Jiang

et al. 2017), and Kaiju (Menzel, Ng, and Krogh 2016), as well as reduced alphabet amino

acid based approaches such as DIAMOND (Buchfink, Xie, and Huson 2015). In almost all

cases these have been designed and benchmarked using short read data (McIntyre et al.

2017).

Analysis of Long-Read Data Metagenomic data

The advent of "third generation" single molecule long read technologies (PacBio and Oxford

Nanopore) has significant implications for metagenomic analyses, most notably for genome

assembly (Frank et al. 2016; Nicholls et al. 2019). These technologies allow read lengths of

10 kilobase pairs (Kbp) and beyond, in strong contrast with the approximately 300 base pairs

(bp) limit of Illumina. However, both PacBio and Nanopore technologies have far higher error

rates (88-94% accuracy for Nanopore (Wick, Judd, and Holt 2018) and 85-87% for PacBio

(Ardui et al. 2018)). The lower accuracy of Nanopore and PacBio (non-circular consensus)

sequence reads may affect the success of current classification methods, and there are few

algorithms designed to exploit long-read data.

As a first approach toward determining the use of long-read technologies for metagenomic

applications, we would like to understand the relative advantages and disadvantages of

using short accurate reads versus long error-prone reads. Recent work has shown that

relatively high genus level classifications of approximately 93% have been achieved using

Nanopore-based metagenomic analyses of a mock bacterial community (Brown et al. 2017).

Here we expand this analysis to allow direct comparison between short and long read

approaches. In addition, we compare metagenomic classification success in microbial

communities as compared to communities of multicellular organisms. We find that longer

reads, despite their higher error rate, can considerably improve classification accuracy

compared to shorter reads, and that this is especially true for specific taxa.

Methods

Genomic data

120   For each of four major taxonomic divisions (bacteria, fungi, animals, and plants), we

121   downloaded 20 genomes from GenBank (Benson et al. 2013). Within each of these

122   divisions, we included genomes from a total of 22 classes, 46 orders, and 58 families (Figure

123   1).

Read simulation

We simulated Nanopore reads using NanoSim 2.0.0 (Yang et al. 2017) with the default error

parameters for *E. coli* R9 1D data. This method uses a mixture model to produce simulated

reads with indel and error rates similar to real datasets. The error model is applied equally to

all parts of a read, and the read lengths are drawn from a distribution approximating real

data. To create simulated read data of specific lengths, we truncated the simulated reads

after the relevant number of basepairs using a custom perl script (i.e. to simulate 100bp

Nanopore reads, we truncated all reads in a simulated dataset to 100bp). We did this for

read lengths varying from 100 bp to 4,000 bp at 100 bp intervals, simulating 1,000 reads per

interval for all taxa (a total of 40,000 reads for each taxon, and 3.2 million reads for all taxa

and read lengths).

Figure 1. Cladogram of species included in the simulated mock community.

We simulated Illumina data using dwgsim 0.1.12 (Homer 2017) with the following options:

dwgsim -e 0.0001 -E 0.0001 -N 2000 -1 100 -2 100 -r 0.0001 -R 0.01 -y 0.000 -c 0

This implements errors to mirror those in Illumina data, with constant error rates of 1e-4 and

no indels (which are extremely rare in Illumina data). We generated 1,000 reads for each

genome, at three read lengths: 100 bp, 150 bp, and 300 bp (a total of 240,000 reads across

all taxa and lengths), and used only single end reads for all analyses.

Sequence Classification

We used BLAST 2.7.1 (Madden 2013) and Kraken2 (Wood and Salzberg 2014) for

sequence classification. We created a local custom database consisting of the NCBI nt

database (downloaded on Feb 8 2019) and the genomes of the 80 taxa that we used to test

classification success. We used the default alignment parameters for BLAST, except for

implementing a maximum e-value of 0.1. We used the match with the highest bit score for all

downstream analyses. For Kraken2 analyses we used the default parameters (in which the

k-mer length is 35 bp and default minimiser length is 31 bp). For Kraken2 we used the taxon

assigned by the lowest common ancestor (LCA) algorithm employed in Kraken2.

Accuracy metrics

To assess the effects of read length on classification accuracy we focus our analysis only on

how often a read is assigned to the correct taxon. For our simulated reads there are three

possible outcomes when querying a database (**Table 1**).

We expect that taxa that are well represented in the database, and which have few closely

related taxa, will have high rates of true matches. Taxa with many close relatives in the

database will have many false matches. Taxa that are poorly represented in the database

will have high rates of failed queries. Both of these latter results are in a class usually

referred to as false negatives: we falsely infer taxon A is absent. However, they largely arise

from different mechanisms. Importantly, as genomic databases become more complete, we

expect the fraction of failed queries will decrease. At the same time we expect that the

fraction of false matches may increase, as more and more closely related taxa become

present in the database. The exact nature of this tradeoff is not well explored. Novel

statistical approaches, such as Bayesian re-estimation of species frequencies, may mitigate

167   the problem (Lu et al. 2016); however, improved methods are required address this problem

168   (Nasko et al. 2018).

170   **Table 1. Description of outcomes for database queries.**

Description of outcome	Metric	Notation
A read query from taxon A returns a match from taxon A	True match (we correctly infer taxon A is present)	M_{true}
A read query from taxon A returns a match from a taxon that is not A	False match (we infer taxon A is absent due to a secondary match)	M_{false}
A read query from taxon A returns no hit at all	Failed query (we infer taxon A is absent due to database paucity).	M_{fail}

There are other aspects of classification success that we do not focus on here. The first of

these is the notion of a true negative: a sequence that is known to *not* arise from any taxa,

should not return a match to any taxa. This is not a biologically realistic situation (all

sequences arise from a taxon), although this aspect is useful when trying to assess the

performance of different classifiers [ref Gardner] and presenting the full truth table. The

second aspect we do not consider here are false positives: if a read query matches taxon A,

but does not arise from taxon A. We would thus falsely interpret taxon A as being present in

a community. This metric is intrinsic to the composition of the community rather than just

each taxon and the database. For example, if taxon A dominates the community, then it

cannot have high rates of false positives relative to true positives simply because the vast

majority of read queries from the community will be from taxon A and thus true positives.

Conversely if taxon B is extremely rare, there will be a large number of false positives

relative to true positives, as very few read queries will be from taxon B, resulting in a very

small number of true positives.

Thus, we use a simplified set of metrics (see **Table 1**) that are not intrinsically related to

community composition: true matches, false matches, and failed queries. We used our

simulated genomic sequence reads from 80 taxa to quantify these three outcomes at both

the genus and family level. To assign genus and family from species, we used the NCBI

taxonomy database (Federhen 2012) (which is used by BLAST as the default taxon

classifier).

We calculate two ratios from the three metrics in **Table 1**. The first is the fraction of true

positives classified correctly (i.e. recall):

$Recall = M_{true}/(M_{true} + M_{false} + M_{fail})$

The second is the ratio of true matches to false matches. This simply excludes failed queries

from the equation. We term this second metric classification success.

$Classification\ Success = M_{true}/(M_{true} + M_{false})$

The critical difference between these metrics is that taxa which are poorly represented in the

database may nevertheless have high rates of classification success, although recall will

necessarily be low. However, as the fraction of failed queries approaches zero (which we

expect as genomic databases grow), these two metrics become equivalent.

Results

We first looked only at short read lengths to quantify the effects of sequencing technology

and classifier (BLAST or Kraken2) on recall at the level of genus. For both bacteria and

fungi, we found that recall was at or above 99.9% for Illumina reads of any length (100bp,

150bp, or 300bp), for both BLAST and Kraken2 (**Fig. 2**). In strong contrast, for Nanopore

data, recall was far lower; approximately 25% for 100bp reads and increasing to 75% at

300bp. In general, Kraken2 had slightly lower recall than BLAST.

Figure 2. Recall is consistently higher in bacteria and fungi than plants or animals for both short Illumina and Nanopore reads. Each panel shows recall for the different kingdoms. Recall for individual taxa is indicated in grey, with median recall shown by dashed (Nanopore) or solid (Illumina). Blue lines indicate recall rates for reads classified using Kraken2; red for reads classified using BLAST. Illumina reads exhibit consistently higher recall; bacteria and fungi exhibit higher recall than plants or animals.

However, for plants and animals, average recall was low regardless of sequencing

technology. Average recall for Illumina reads peaked at approximately 55% and 75% for

animals and plants, respectively (**Fig 2,** light blue lines). Nanopore recall rates peaked at just

over 20% and 35% for animals and plants, respectively. However, this was highly taxon-

dependent, with some taxa consistently having recall near 100%, while others remained

close to 0% regardless of sequencing technology or read length (**Fig. 2,** grey lines). Perhaps

surprisingly, on average Kraken2 outperformed BLAST for Illumina reads for both plant and

animal taxa.

We next quantified differences in classification success (the proportion of all classified reads

that were correctly classified), again considering only short read lengths. For bacteria and

fungi, both Illumina and Nanopore reads exhibited high classification success, with the

exception of Kraken2 classification of Nanopore reads (**Fig. 3**). For each sequencing

method and classifier, classification success for plants and animals was low relative to

bacteria and fungi. For both Illumina and Nanopore, BLAST resulted in approximately 87%

and 97% of reads being correctly classified, for animals and plants respectively. However,

Kraken2 success was far lower, especially for Nanopore reads, peaking at 54% in animals

(**Fig. 3**). Over this range of read lengths, we found only a weak relationship between read

length and classification success, in contrast to the results for recall.

It is perhaps expected that highly accurate Illumina reads would result in more accurate

taxonomic classification than long error-prone Nanopore reads. However, it is possible to

obtain Nanopore reads far in excess of 300bp (reads up to 2 megabase pairs have been

sequenced), so we next quantified recall and classification success for reads with lengths up

to 4,000 bp. Because such read lengths are not currently possible to obtain using Illumina

technology, we did not measure recall and classification success for Illumina reads of similar

lengths.

Figure 3. Classification success for short reads is weakly related to read length and strongly dependent on classification method. Each panel shows recall for the different kingdoms. Classification success for individual taxa is indicated in grey, with median classification success shown by solid lines (Illumina) or dashed lines (Nanopore). Blue lines indicate recall rates for reads classified using Kraken2; red for reads classified using BLAST. For bacteria and fungi, median classification rates of Illumina-BLAST, Illumina-Kraken2, and Nanopore-BLAST are almost exactly 100% for all read lengths.

We observed similar relationship between read length and recall for both BLAST and

Kraken2. For bacteria and fungi, recall increased from ~20% using 100 bp reads to almost

100% when using 1500 bp reads. For animals and plants we observed similar trends,

although at no point did recall approach 100%. However, long Nanopore reads surpassed

the recall of even the longest Illumina reads (300 bp) classified with Kraken, with crossover

points at approximately 3000 bp for animals and 2500 bp for plants **(Fig. 4**, red and blue

solid lines).

**Figure 4. Long Nanopore reads equal or surpass the recall of the longest Illumina
reads for both BLAST and Kraken2.** Each panel shows recall for the different kingdoms.
Recall for Nanopore reads for individual taxa is indicated in grey, with median recall
indicated by dashed lines, either blue (Kraken2) or red (BLAST). The recall rates for 300 bp
Illumina reads are shown as thin solid lines, again either blue (Kraken2) or red (BLAST).
Coloured points show the recall for all Illumina reads of all lengths (100 bp, 150 bp, and 300
bp).

We also considered this metric at the level of family. In this case found that for animals,

Nanopore reads surpassed Illumina reads only at lengths close to 4000 bp, reaching

approximately 70% recall at this point (**Supp Fig. 2**). However, for plants Nanopore recall

surpassed Illumina recall at 2500 bp, with 4000 bp reads yielding a recall of approximately

90%. We found again that for both animals and plants, Kraken2 recall surpassed BLAST

when relying on Illumina reads.

Figure 5. Classification success for long reads is dependent on read length only for
Kraken2 classification. Each panel shows classification success for the different kingdoms.
Classification success for individual taxa is indicated in grey, with median classification
success shown by solid lines (Illumina) or dashed lines (Nanopore). Blue indicates
classification success rates for reads classified using Kraken2, while red indicates those
classified using BLAST. For animal and plants, the classification success of Kraken2
depends strongly on read length, and never surpasses BLAST or Illumina at any length.

We next examined classification success at longer read lengths. For BLAST we observed no

relationship between classification success and read length for any taxon (**Fig 5**.). Bacteria

and fungi both had consistently high classification success (median 100%), while animals

and plants had lower classification success (median 82% and 96%, respectively). However,

for Kraken2 we observed a consistent increase in classification success as read length

increased. However, this never exceeded the classification success we observed for BLAST,

nor did it succeed the classification success we observed for short accurate Illumina reads.

Finally, we tested classification success at the level of Family. In this case, we observed that

for BLAST, the classification success for plants was approximately 99% overall read lengths,

while for Kraken2 only 4000 bp reads reached this level. For animals, BLAST classification

success was approximately 95% over all read lengths, but for Kraken2 reached a maximum

of 85% at the longest read lengths (**Supp. Fig. 3**).

294   Discussion

Here we have compared the relative accuracy of taxon classification using simulated short

accurate reads (Illumina) and long, error-prone reads (Nanopore) with known ground truth.

We have used two simple metrics of success: recall (the ratio of correctly classified reads to

all reads) and classification success (the ratio of correctly classified reads to all classified

reads). We have tested taxon classification using a broad range of taxa, including bacteria,

fungi, animals, and plants.

Recall for both BLAST and Kraken2 was improved by the use of long reads, especially in the

case of animals and plants, for which recall improved almost three-fold as read length

increased from 300 bp to 4,000 bp. Generally both Kraken2 and BLAST achieved similar

levels of recall. The exception was for short reads for animals and plants, for which Kraken2

was more accurate than BLAST.

We found no relationship between classification success and read length for BLAST. This

implies that the ratio of correctly classified reads to all classified reads remains relatively

constant over different read lengths. However, the number of reads that are classified *at all*

increases with read length (causing an increase in recall). These observations are in line

with what has been observed by others (McHardy et al. 2007). The exception to this lack of

relationship between classification success and read length was for Kraken2, for which the

proportion of correctly classified reads increases with read length by more than 50% for both

plants and animals.

Our results also indicate that recall for long Nanopore reads was equal to or higher than

short Illumina reads. This was true regardless of kingdom, or classification method, with

Nanopore surpassing 300 bp Illumina reads at approximately 1500 bp for plants and

animals, and surpassing 150 bp Illumina reads at between 1500 bp and 3000 bp for bacteria

and fungi, depending on the methodology (**Fig. 4**). Even the longest Illumina reads, at 300

bp, were outclassed by Nanopore at between 3500 and 4000 bp, depending on

methodology.   These results do suggest that one approach to improve Nanopore

classification accuracy is to impose minimum read lengths. This can be achieved by

performing size selection during library preparation or during computational analyses.

At first glance, then, there appears to be a clear trade-off between short read Illumina and

long read Nanopore sequencing for metagenomic analyses. While Nanopore allows higher

recall at long read lengths, this advantage is offset by the fact that Illlumina generally

provides more reads per run. At most, recall for Nanopore improves 50% beyond 300 bp

Illumina reads, while classification success is similar (using BLAST). Thus, if the read

capacity of Illumina runs is 50% or more than Nanopore, the number of classified reads will

be maximised using Illumina technology - on a per sequencing run basis. However, for many

researchers the more relevant metric is cost per read. In this case, MinION read yields are

approximately equal to MiSeq, and only HiSeq or NovaSeq provides a clear cost advantage

over Nanopore MinION. On the other hand, cost per read for PromethION are not far from

NovaSeq. Thus, we find no clear advantage in using Illumina over Nanopore given the

observed classification accuracy for long inaccurate Nanopore reads.

Differences in accuracy between bacteria, fungi, animals, and plants

We find very large differences in classification accuracy (mostly in terms of recall) for

bacteria and fungi versus plants and animals. The discrepancy between taxonomic groups

likely arises from a variety of factors. Among these are the higher degree of divergence

between bacterial species relative to animal and plant species, and the complexity of

bacterial genomes compared to eukaryotic genomes. We discuss these factors below.

Bacterial taxa are often considered separate species once they have diverged by 6% ANI

(Average Nucleotide Identity) on a genomic level (Stackebrandt and Goebel 1994;

Konstantinidis and Tiedje 2005). The degree of nucleotide divergence between eukaryotic

species is not standardised (Cognato 2006), and species are generally designated as such

based on the biological species concept put forward by Mayr (Mayr 1999). Although

divergence levels differ substantially between loci (as for bacteria), for some loci general

ranges for eukaryotic species have emerged. For example, for mitochondrial COI, between-

species divergence is usually greater than 3% (Song et al. 2008; Lefébure et al. 2006).

These loci are among the fastest diverging loci in plant and animal genomes, and many

other loci may differ by far less than 1% between species. Due to this low level of

divergence, metagenomic classifiers may frequently classify animal and plant genera with

lower accuracy than bacterial genera.

A second explanation for the increased classification success in bacteria and fungi is that

these genomes contain fewer repetitive elements than animals or plants (Treangen et al.

2009). Although such repetitive regions are usually masked from classifiers (including

BLAST and Kraken2), this masking may not be complete.

A third reason is that the genomic databases for plants and animals are far less complete

than for bacteria and fungi. There is a large difference in the number of genomes and

sequences available for different Kingdoms, with bacteria having significantly more species

present than the next closest kingdom (See Supp. Fig.1). However, we expect this factor will

be mitigated in the future as genomic databases continue to expand and computational

search methods continue to improve.

Differences in accuracy between Kraken2 and BLAST

We observed similar levels of recall for BLAST and Kraken2 over most reads lengths. However, there were strong differences in classification success. For short reads, Kraken2 classification success was far lower than BLAST. As read lengths increased, Kraken2 classification success approached BLAST. Part of this is likely due to longer reads allowing multiple k-mer matches, decreasing the probability of a false positive classification. One perhaps underappreciated advantage of Kraken2 over BLAST is that Kraken2 has reduced sensitivity to structural variation within reads. As Kraken2 allows multiple k-mers to match within a read, structural changes (e.g. inversions) are less likely to influence the outcome of Kraken2 matching. Such structural changes may influence BLAST due to the matching and extend algorithm. Thus for long reads, classifiers that are insensitive to synteny may be more successful, especially for taxa in which structural rearrangements are common.

Conclusions

Here we have shown despite being error-prone, Nanopore reads are useful for metagenomic classification due to their increased length, and that for plant and animal communities, the classification accuracy of long Nanopore reads exceeds that of Illumina. We found that classification accuracy is more dependent on the set of taxa being considered than on the metagenomic classifier being used (Kraken2 or BLAST), and that this was true for both short accurate (Illumina) and long error-prone (Nanopore) sequence data. Together these data suggest that one consideration in selecting a metagenomic sequencing method (i.e. long or short read) is the taxonomic group of interest.

Declarations

Ethics approval and consent to participate – not applicable

Consent for publication – approved by all authors

Availability of data and material – <u>data for this manuscript will be uploaded to DataDryad if</u>

<u>manuscript is accepted</u>

Competing interests – the authors have no competing interests to declare

Funding – Some of this work was supported by a Massey University Strategic Research

Excellence Fund awarded to NF.

Authors' contributions - WP, NF, and OS conceived the project, WP and OS simulated

and generated the data. WP and OS analysed the data. WP, NF, and OS wrote the paper.

Acknowledgements - Thanks to Paul Gardner for his helpful and insightful comments on

the manuscript.

396 References

10K Community of Scientists, Genome. 2009. "Genome 10K: A Proposal to Obtain Whole-

Genome Sequence for 10 000 Vertebrate Species." *The Journal of Heredity*.

https://academic.oup.com/jhered/article-abstract/100/6/659/839176.

Ardui, Simon, Adam Ameur, Joris R. Vermeesch, and Matthew S. Hestand. 2018. "Single

Molecule Real-Time (SMRT) Sequencing Comes of Age: Applications and Utilities for

Medical Diagnostics." *Nucleic Acids Research* 46 (5): 2159–68.

Benson, Dennis A., Mark Cavanaugh, Karen Clark, Ilene Karsch-Mizrachi, David J. Lipman,

James Ostell, and Eric W. Sayers. 2013. "GenBank." *Nucleic Acids Research* 41

(Database issue): D36–42.

Brown, Bonnie L., Mick Watson, Samuel S. Minot, Maria C. Rivera, and Rima B. Franklin.

2017. "MinION nanopore Sequencing of Environmental Metagenomes: A Synthetic

Approach." *GigaScience* 6 (3): 1–10.

Buchfink, Benjamin, Chao Xie, and Daniel H. Huson. 2015. "Fast and Sensitive Protein

Alignment Using DIAMOND." *Nature Methods* 12 (1): 59–60.

Caporaso, J. Gregory, Justin Kuczynski, Jesse Stombaugh, Kyle Bittinger, Frederic D.

Bushman, Elizabeth K. Costello, Noah Fierer, et al. 2010. "QIIME Allows Analysis of

High-Throughput Community Sequencing Data." *Nature Methods* 7 (5): 335–36.

Cognato, Anthony I. 2006. "Standard Percent DNA Sequence Difference for Insects Does

Not Predict Species Boundaries." *Journal of Economic Entomology* 99 (4): 1037–45.

Escobar-Zepeda, Alejandra, Arturo Vera-Ponce de León, and Alejandro Sanchez-Flores.

2015. "The Road to Metagenomics: From Microbiology to DNA Sequencing

Technologies and Bioinformatics." *Frontiers in Genetics* 6 (December): 348.

Federhen, Scott. 2012. "The NCBI Taxonomy Database." *Nucleic Acids Research* 40

(Database issue): D136–43.

Frank, J. A., Y. Pan, A. Tooming-Klunderud, V. G. H. Eijsink, A. C. McHardy, A. J.

Nederbragt, and P. B. Pope. 2016. "Improved Metagenome Assemblies and Taxonomic

Binning Using Long-Read Circular Consensus Sequence Data." *Scientific Reports* 6

(May): 25373.

Homer, Nils. 2017. *DWGSIM* (version 0.1.12). Github. https://github.com/nh13/DWGSIM.

Huson, Daniel H., Sina Beier, Isabell Flade, Anna Górska, Mohamed El-Hadidi, Suparna

Mitra, Hans-Joachim Ruscheweyh, and Rewati Tappu. 2016. "MEGAN Community

Edition - Interactive Exploration and Analysis of Large-Scale Microbiome Sequencing

Data." *PLoS Computational Biology* 12 (6): e1004957.

Jiang, Yuan, Jun Wang, Dawen Xia, and Guoxian Yu. 2017. "EnSVMB: Metagenomics

Fragments Classification Using Ensemble SVM and BLAST." *Scientific Reports* 7 (1):

9440.

Ji, Yinqiu, Louise Ashton, Scott M. Pedley, David P. Edwards, Yong Tang, Akihiro

Nakamura, Roger Kitching, et al. 2013. "Reliable, Verifiable and Efficient Monitoring of

Biodiversity via Metabarcoding." *Ecology Letters* 16 (10): 1245–57.

Keeling, Patrick J., Fabien Burki, Heather M. Wilcox, Bassem Allam, Eric E. Allen, Linda A.

Amaral-Zettler, E. Virginia Armbrust, et al. 2014. "The Marine Microbial Eukaryote

Transcriptome Sequencing Project (MMETSP): Illuminating the Functional Diversity of

Eukaryotic Life in the Oceans through Transcriptome Sequencing." *PLoS Biology* 12 (6):

e1001889.

Kim, Daehwan, Li Song, Florian P. Breitwieser, and Steven L. Salzberg. 2016. "Centrifuge:

Rapid and Sensitive Classification of Metagenomic Sequences." *Genome Research* 26

(12): 1721–29.

Konstantinidis, Konstantinos T., and James M. Tiedje. 2005. "Genomic Insights That

Advance the Species Definition for Prokaryotes." *Proceedings of the National Academy*

of Sciences of the United States of America 102 (7): 2567–72.

Lefébure, T., C. J. Douady, M. Gouy, and J. Gibert. 2006. "Relationship between

Morphological Taxonomy and Molecular Divergence within Crustacea: Proposal of a

Molecular Threshold to Help Species Delimitation." *Molecular Phylogenetics and*

Evolution 40 (2): 435–47.

Lewin, Harris A., Gene E. Robinson, W. John Kress, William J. Baker, Jonathan Coddington,

Keith A. Crandall, Richard Durbin, et al. 2018. "Earth BioGenome Project: Sequencing

Life for the Future of Life." *Proceedings of the National Academy of Sciences of the*

United States of America 115 (17): 4325–33.

Lu, Jennifer, Florian P. Breitwieser, Peter Thielen, and Steven L. Salzberg. 2016. "Bracken:

Estimating Species Abundance in Metagenomics Data." https://doi.org/10.1101/051813.

Madden, Thomas. 2013. *The BLAST Sequence Analysis Tool*. National Center for

Biotechnology Information (US).

Mayr, Ernst. 1999. *Systematics and the Origin of Species, from the Viewpoint of a Zoologist*.

Harvard University Press.

McHardy, Alice Carolyn, Héctor García Martín, Aristotelis Tsirigos, Philip Hugenholtz, and

Isidore Rigoutsos. 2007. "Accurate Phylogenetic Classification of Variable-Length DNA

Fragments." *Nature Methods* 4 (1): 63–72.

McIntyre, Alexa B. R., Rachid Ounit, Ebrahim Afshinnekoo, Robert J. Prill, Elizabeth Hénaff,

Noah Alexander, Samuel S. Minot, et al. 2017. "Comprehensive Benchmarking and

Ensemble Approaches for Metagenomic Classifiers." *Genome Biology* 18 (1): 182.

Menzel, Peter, Kim Lee Ng, and Anders Krogh. 2016. "Fast and Sensitive Taxonomic

Classification for Metagenomics with Kaiju." *Nature Communications* 7 (April): 11257.

Nasko, Daniel J., Sergey Koren, Adam M. Phillippy, and Todd J. Treangen. 2018. "RefSeq

Database Growth Influences the Accuracy of K-Mer-Based Lowest Common Ancestor

Species Identification." *Genome Biology* 19 (1): 165.

Nicholls, Samuel M., Joshua C. Quick, Shuiquan Tang, and Nicholas J. Loman. 2019. "Ultra-

Deep, Long-Read Nanopore Sequencing of Mock Microbial Community Standards."

GigaScience 8 (5). https://doi.org/10.1093/gigascience/giz043.

OBrien, Stephen J., David Haussler, and Oliver Ryder. 2014. "The Birds of Genome10K."

GigaScience 3 (1): 32.

O'Leary, Nuala A., Mathew W. Wright, J. Rodney Brister, Stacy Ciufo, Diana Haddad, Rich

McVeigh, Bhanu Rajput, et al. 2016. "Reference Sequence (RefSeq) Database at NCBI:

Current Status, Taxonomic Expansion, and Functional Annotation." *Nucleic Acids

Research* 44 (D1): D733–45.

Ounit, Rachid, Steve Wanamaker, Timothy J. Close, and Stefano Lonardi. 2015. "CLARK:

Fast and Accurate Classification of Metagenomic and Genomic Sequences Using

Discriminative K-Mers." *BMC Genomics* 16 (March): 236.

Robinson, Gene E., Kevin J. Hackett, Mary Purcell-Miramontes, Susan J. Brown, Jay D.

Evans, Marian R. Goldsmith, Daniel Lawson, Jack Okamuro, Hugh M. Robertson, and

David J. Schneider. 2011. "Creating a Buzz about Insect Genomes." *Science* 331

(6023): 1386.

Roumpeka, Despoina D., R. John Wallace, Frank Escalettes, Ian Fotheringham, and Mick

Watson. 2017. "A Review of Bioinformatics Tools for Bio-Prospecting from Metagenomic

Sequence Data." *Frontiers in Genetics* 8 (March): 23.

Schloss, Patrick D., and Jo Handelsman. 2005. "Metagenomics for Studying Unculturable

Microorganisms: Cutting the Gordian Knot." *Genome Biology* 6 (8): 229.

Schloss, Patrick D., Sarah L. Westcott, Thomas Ryabin, Justine R. Hall, Martin Hartmann,

Emily B. Hollister, Ryan A. Lesniewski, et al. 2009. "Introducing Mothur: Open-Source,

Platform-Independent, Community-Supported Software for Describing and Comparing

Microbial Communities." *Applied and Environmental Microbiology* 75 (23): 7537–41.

Song, Hojun, Jennifer E. Buhay, Michael F. Whiting, and Keith A. Crandall. 2008. "Many

Species in One: DNA Barcoding Overestimates the Number of Species When Nuclear

Mitochondrial Pseudogenes Are Coamplified." *Proceedings of the National Academy of*

Sciences of the United States of America 105 (36): 13486–91.

Stackebrandt, E., and B. M. Goebel. 1994. "Taxonomic Note: A Place for DNA-DNA

Reassociation and 16S rRNA Sequence Analysis in the Present Species Definition in

Bacteriology." *International Journal of Systematic and Evolutionary Microbiology* 44 (4):

846–49.

Teeling, Emma C., Sonja C. Vernes, Liliana M. Dávalos, David A. Ray, M. Thomas P.

Gilbert, Eugene Myers, and Bat1K Consortium. 2018. "Bat Biology, Genomes, and the

Bat1K Project: To Generate Chromosome-Level Genomes for All Living Bat Species."

Annual Review of Animal Biosciences 6 (February): 23–46.

Temperton, Ben, and Stephen J. Giovannoni. 2012. "Metagenomics: Microbial Diversity

through a Scratched Lens." *Current Opinion in Microbiology* 15 (5): 605–12.

Thomas, Torsten, Jack Gilbert, and Folker Meyer. 2012. "Metagenomics - a Guide from

Sampling to Data Analysis." *Microbial Informatics and Experimentation* 2 (1): 3.

Treangen, Todd J., Anne-Laure Abraham, Marie Touchon, and Eduardo P. C. Rocha. 2009.

"Genesis, Effects and Fates of Repeats in Prokaryotic Genomes." *FEMS Microbiology*

Reviews 33 (3): 539–71.

Wick, Ryan, Louise M. Judd, and Kathryn E. Holt. 2018. *Comparison of Oxford Nanopore*

Basecalling Tools. https://doi.org/10.5281/zenodo.1188469.

Wood, Derrick E., and Steven L. Salzberg. 2014. "Kraken: Ultrafast Metagenomic Sequence

Classification Using Exact Alignments." *Genome Biology* 15 (3): R46.

Yang, Chen, Justin Chu, René L. Warren, and Inanç Birol. 2017. "NanoSim: Nanopore

Sequence Read Simulator Based on Statistical Characterization." *GigaScience* 6 (4): 1–

6.

Supplementary Materials

Supplementary Figure 1. The number of species present in the NCBI RefSeq database has
grown roughly exponentially over time. Note that the y-axis is plotted on a log scale. Data were
retrieved from the RefSeq database (O'Leary et al. 2016):
https://ftp.ncbi.nlm.nih.gov/genomes/GENOME_REPORTS/eukaryotes.txt and
https://ftp.ncbi.nlm.nih.gov/genomes/GENOME_REPORTS/prokaryotes.txt

Supplementary Figure 2 Recall at the family level. Each panel shows recall for the different kingdoms. Recall for Nanopore reads for individual taxa is indicated in grey, with median recall indicated by dashed lines, either blue (Kraken2) or red (BLAST). The recall rates for 300 bp Illumina reads are shown as thin solid lines, again either blue (Kraken2) or red (BLAST). Coloured points show the recall for all Illumina reads of all lengths (100 bp, 150 bp, and 300 bp).

Supplementary Figure 3 Classification success at the family level. Each panel shows classification success for the different kingdoms. Classification success for individual taxa is indicated in grey, with median classification success shown by solid lines (Illumina) or dashed lines (Nanopore). Blue indicates classification success rates for reads classified using Kraken2, while red indicates those classified using BLAST. For animal and plants, the classification success of Kraken2 depends strongly on read length, and never surpasses BLAST or Illumina at any length.

Supplementary Table 1. List of species include in the *in silico* mock community, with associated Kingdom and NCBI

Species	Kingdom	NCBI Accession
Actinidia chinensis	Plantae	CM009654.1
Ananas comosus	Plantae	CM003813.1
Arabidopsis thaliana	Plantae	CP002684.1
Brassica nigra	Plantae	CM004491.1
Camelina sativa	Plantae	CM002729.1

Citrus sinensis	Plantae	CM001701.1
Dioscorea rotundata	Plantae	BDMI01000001.1
Eutrema salsugineum	Plantae	CM001778.1
Gossypioides kirkii	Plantae	CM008980.1
Leersia perrieri	Plantae	CM002476.1
Malus domestica	Plantae	CM007867.1
Micromonas sp.	Plantae	CP001574.1
Panicum hallii	Plantae	CM008046.2
Raphanus sativus	Plantae	CM007999.1
Rosa chinensis	Plantae	CM009582.1
Setaria italica	Plantae	CM004364.1
Solanum lycopersicum	Plantae	CM001064.3
Sorghum bicolor	Plantae	CM000760.3
Theobroma cacao	Plantae	LT594788.1
Trifolium pratense	Plantae	LT555306.1
Amphiprion percula	Animalia	CM009708.1
Bos indicus	Animalia	CM003021.1
Capra hircus	Animalia	CM001710.2
Chrysemys picta	Animalia	CM002655.1
Columba livia	Animalia	CM007525.1
Cyprinus carpio	Animalia	LN590701.1
Drosophila busckii	Animalia	CP012523.1
Equus caballus	Animalia	CM000377.2
Falco peregrinus	Animalia	CM007505.1
Homo sapiens	Animalia	CM004593.1
Lycaon pictus	Animalia	CM007565.1
Macaca mulatta	Animalia	CM000308.1
Microcebus murinus	Animalia	CM007661.1

Mus musculus	Animalia	CM004154.1
Oncorhynchus tshawytscha	Animalia	CM009202.1
Oryctolagus cuniculus	Animalia	CM000790.1
Ovis aries	Animalia	CM008472.1
Takifugu rubripes	Animalia	HE602535.1
Timema cristinae	Animalia	CM007794.2
Xiphophorus maculatus	Animalia	CM008938.1
Agaricus bisporus	Fungi	CP015470.1
Alternaria solani	Fungi	CP022024.1
Colletotrichum higginsianum	Fungi	CM004455.1
Cryptococcus gattii	Fungi	CP025759.1
Debaryomyces hansenii	Fungi	CR382133.2
Eremothecium sinecaudum	Fungi	CP014242.1
Flammulina velutipes	Fungi	CM002695.1
Fusarium verticillioides	Fungi	CM000578.1
Kluyveromyces lactis	Fungi	CR382121.1
Komagataella phaffii	Fungi	LT962476.1
Lachancea nothofagi	Fungi	LT598449.1
Malassezia sympodialis	Fungi	LT671813.1
Millerozyma farinosa	Fungi	FO082059.1
Ogataea parapolymorpha	Fungi	CM002300.1
Saccharomyces cerevisiae	Fungi	BK006935.2
Sporisorium scitamineum	Fungi	CP010913.1
Trichoderma reesei	Fungi	CP016232.1
Valsa mali	Fungi	CM003098.1
Yarrowia lipolytica	Fungi	HG934059.1
Zygosaccharomyces rouxii	Fungi	CU928173.1
Acidithiobacillus ferrivorans	Bacteria	LT841305.1

Bacillus thuringiensis	Bacteria	CP015250.1
Bacillus velezensis	Bacteria	CP025939.1
Bifidobacterium longum	Bacteria	CP013673.1
Bordetella bronchiseptica	Bacteria	CM002881.1
Brucella melitensis	Bacteria	CP018494.1
Campylobacter jejuni	Bacteria	CP012689.1
Caulobacter crescentus	Bacteria	AE005673.1
Cellvibrio japonicus	Bacteria	CP000934.1
Escherichia albertii	Bacteria	AP014855.1
Gordonibacter sp.	Bacteria	LT827128.1
Klebsiella pneumoniae	Bacteria	CP025088.1
Mycobacterium tuberculosis	Bacteria	CP023640.1
Ornithobacterium rhinotracheale	Bacteria	CP006828.1
Pseudomonas arsenicoxydans	Bacteria	LT629705.1
Salmonella enterica	Bacteria	CP007400.2
Serratia symbiotica	Bacteria	LN890288.1
Staphylococcus aureus	Bacteria	CP012974.1
Treponema pallidum	Bacteria	CP020366.1
Vibrio cholerae	Bacteria	LT907989.1