Systematic exploration of protein conformational space using a Distance Geometry approach

Thérèse E. Malliavin,[*,†] Antonio Mucherino,[‡] Carlile Lavor,[¶] and Leo Liberti[*,§]

†*Unité de Bioinformatique Structurale, Institut Pasteur, UMR 3528, CNRS, Centre de Bioinformatique, Biostatistique et Biologie Intégrative, Institut Pasteur, USR 3756, CNRS, Paris, (75015) France*

‡*IRISA, Rennes, France*

¶*IMECC-UNICAMP, University of Campinas, Campinas, SP, Brazil*

§*CNRS LIX, Ecole Polytechnique, 91128 Palaiseau, France*

E-mail: therese.malliavin@pasteur.fr; leo.liberti@lix.polytechnique.fr

Short title

Systematic exploration of protein conformations by TAiBP

Corresponding authors

Thérèse E Malliavin

Institut Pasteur and CNRS UMR 3528, Unité de Bioinformatique Structurale, 25 rue du Dr Roux, 75015 Paris, France E-mail address: therese.malliavin@pasteur.fr

Leo Liberti

CNRS LIX (UMR 7161), Ecole Polytechnique, 91128 Palaiseau, France. E-mail address: leo.liberti@lix.polytechnique.fr

Abstract

The optimization problem that arises for protein structure determination is undergoing a change of perspective due to the larger importance in biology taken by the disordered regions of biomolecules and intrinsically disordered proteins. Indeed, in such cases, the algorithm convergence criterion is more difficult to set up; moreover, the enormous size of the space makes it difficult to achieve a complete exploration. The interval Branch-and-Prune (iBP) approach, based on a reformulating of the Distance Geometry Problem (DGP) and proposed few years ago, provides a theoretical frame for the fast generation of protein conformations, by systematically sampling the conformational space. When an appropriate subset of inter-atomic distances is known exactly, this worst-case exponential-time algorithm is provably complete and fixed-parameter tractable. These guarantees, however, quickly disappear as distance measurement errors are introduced. Here we propose a variant of this approach: the threading-augmented interval Branch-and-Prune (TAiBP), where the combinatorial explosion of the original iBP approach arising from its exponential complexity is alleviated by partitioning the input instances into consecutive peptide fragments and by using Self-Organizing Maps (SOMs) to obtain clusters of similar solutions. A validation of the TAiBP approach is presented here on a set of proteins of various sizes and structures. The calculation inputs are: a uniform covalent geometry extracted from force field covalent terms, the backbone dihedral angles with error intervals, and some long-range distances. For most of the protein smaller than 50 residues and interval widthes of $20°$, the TAiBP approach yielded solutions with RMSD values smaller than 3 Å with respect to the initial protein conformation. The efficiency of TAiBP approach for proteins larger than 50 residues will require the use of non-uniform covalent geometry, and may benefit from the recent development of residue-specific force-field.

Introduction

Since the early days of structural biology, optimization techniques have been at the heart of biomolecular structure calculation. Indeed, most of the experimental information is only indirectly related to protein structure. In addition, this information is noisy. Furthermore, the sparsity of data is made even bigger as most of biophysical techniques concentrates on time-average or space-average data in order to obtain large enough signal-to-noise ratio.

Several optimisation schemes have been used for Nuclear Magnetic Resonance (NMR) structure determination, such as simulated annealing[1] and genetic algorithms.[2] More recently, a Bayesian scheme,[3] using a Markov chain Monte Carlo (MCMC) scheme for the conformational space sampling,[4,5] allowed the increase of the convergence radius for problems of protein structure determination by NMR. In addition, the use of a logharmonic shape for distance restraint potential,[6] along with a Bayesian approach for the restraint weighting,[7] allowed an improvement of the quality of NMR protein structures.[8–10]

The optimization schemes used up to now come without a completeness/exactness guarantee. At convergence they can at most ensure local optimality, although they are commonly used in the hope of obtaining the global minimum or several global minima of the optimisation problem. This, however, depends on the choice of a starting point for the computation, which is why calculations of protein conformations under NMR restraints are repeated several times during the procedure of structure determination.[11] The convergence of these algorithms is generally required in order to accept a set of conformations as a solution. But, this iterative frame[12,13] encounters difficulties when the problem has many local minima that are far apart. Such cases started to occur more frequently in the field of structural biology with the growing interest to disordered regions of biomolecules.[14–16] Monte Carlo approaches have been proposed for intrinsically disordered proteins.[17] Molecular dynamics simulations[18] are also used on all kind of biomolecular polymers, but they do not provide a definitive answer to the problem of finding all minima.

Since NMR studies biomolecules in solution, and due to the large number of various

parameters it can measure, it is particularly sensitive to the effect of internal mobility. NMR measures are inter-atomic distances and angles, which are closely related parameters. The problem of protein structure determination by NMR can be thus considered as a Distance Geometry Problem (DGP).[19,20]

The interval Branch-and-Prune (iBP) approach has been developed[21] for solving the DGP in the frame of the calculation of protein conformations. This approach, based on reordering the atoms on the backbone, has been used to solve the classical DGP[22,23] defined on the studied system. A reordering list ensures that there is a restricted and manageable locus for the spatial position of any atom. This is achieved by using a relaxed form of trilateration with respect to the three preceding atoms in the order. More precisely, two out of three of the distances involved in trilateration must be known (suffcently) exactly, and one is allowed to have error represented by an interval. Any atom, together with its three *reference predecessors*, form a 4-clique. In that way, the iBP approach mimics the approach of exploring protein conformation in torsion angle space.[24–27] The exact distances are given by covalent bond lengths and covalent angle values, using the cosine law. Note that applying this framework using generic information from a force field instead of measured distances makes the implicit assumption of a uniform covalent geometry within the protein structure. Analyses of high-resolution crystallographic structures,[28,29] however, have shown that this assumption is not necessarily verified. Independent parallel work has conducted to the development of residue-specfic force field.[30–33]

Basing on the atom reordering, it is possible to describe a tree exploration algorithm in order to find all solutions of a DGP instance. Each tree node represents a spatial position for an atom. The level of a node in the tree is the index of the atom in the reordering. This means that a whole level represents all of the possible spatial positions for the atom indexed by the level. The tree width increases exponentially in the worst case, but certain reorderings make it possible to bound the width,[34] which yields a fixed-parameter tractable behavior (at least with exact distances). We note that the exploration of this tree is implicitly incomplete, in

the sense that certain subtrees are pruned because the atomic positions at their root nodes are not consistent with long range distances to preceding atoms. Naturally, each pruned node induces the pruning of the subtree rooted at that node. It was demonstrated[20,21] that, starting from a set of exact distances measured in a given PDB structure, the search tree can be completely (but implicitly) explored in a relatively small amount of CPU time.

In this paper, we employ the iBP algorithm in a setting which is considerably closer to the protocols of protein structure determination than the mathematical setting in which it was initially conceived. Instead of exact distances measured on a given PDB structure, this requires the use of a mixed set of distance intervals and of exact distances arising from a covalent geometry defined through a force field. Several attempts have been made in this direction in the recent past. A significant exploration of the conformational space of some α-helical 15 to 51-residues proteins was performed in,[35] and more recently, the iBP approach was re-implemented[36] in order to make it closer to an application to real-life case of protein structure determination. First, the number of tree branches was reduced[37] by taking into account the information from improper angles. Second, a parser and a grammar have been defined to convert the topology, parameter and atom type information used in molecular modeling to the distance information which is the main input of iBP. Third, a syntax has been defined to make the atom reordering information a user-defined input of the calculation. This new implementation makes it possible to perform tree branching on intervals determined on ϕ and ψ backbone angles, which may be obtained through chemical shift measurements.[38]

In the present work, we use this newer implementation to extensively test the iBP approach on a set of various protein structures. The expected combinatorial explosion is prevented by several ingredients: (i) the division of the protein into fragments which are sampled independently and then assembled, (ii) the use of signed improper angle values to reduce the tree size of each fragment, (iii) the use of self-organizing maps to cluster intermediate fragments. The input restraints are: (i) ϕ and ψ values with error intervals of 20°,

40° and 60°, (ii) distance values connecting fragment extremities with error intervals of 6 Å, (iii) long-range distances connecting middle residues of fragments with error intervals of 10 Å. The method is called threading-augmented iBP (TAiBP) approach, as it intends on one hand, to generate conformation of peptide fragments using iBP and on the other hand, to place/thread these fragments in the 3D space to build protein conformations. We point out that the idea to separate iBP instances in sub-instances is not completely new, but it was explored so far only in the context of parallel[39] and distributed[40] computing. Also, a building of protein conformations from fragment assembly was proposed initially[41–43] in the Rosetta approach for protein structure modeling.

The proposed methodology is innovative with respect to the state of the art because it is designed to find all possible configurations compatible with a given set of angle and distance restraints on a given protein. This is in contrast to classical methods for structure determination,[1] which might at best produce different protein conformations. The approach is different from the more recently proposed methods aiming at determining the global minimum configuration of the system:[44–49] or at determining all few relative positions of monomers within a protein homo-oligomer.[50] On the contrary, the exhaustive list of conformations generated by TAiBP provides a solution for the structural analysis of highly flexible or disordered regions of biomolecules.

It is important to note that our purpose is beyond finding a conformation close to the target one, since we aim instead to the much more ambitious goal of finding many (and hopefully all) incongruent but geometrical consistent conformations. Moreover,because our algorithm approach is not iterative but based on branching, we have no need for considering "convergence to a local optimum" a requirement for accepting a conformation. The results of our computational experiments, however, have been validated by detecting whether conformations close to the target PDB structure have been sampled during the tree exploration, by Root Mean Square Distance (RMSD) of atomic coordinates to the target structure. The proposed approach allows us to explore the tree for proteins up to 50 residues. The error on

the backbone angles, as well as the non-uniform covalent geometry, are both factors which (by now) prevent our method from being successful on proteins larger than 50 residues.

Materials and Methods

Test case calculations

The database of protein structures was built in the following way. The protein structures contained in `kinemage.biochem.duke.edu/databases/top100.php`[51] have been downloaded: these structures are X-ray crystallographic structures with resolution in range 1.4-1.7 Å, on which hydrogens have been added with rotational optimization of OH, SH and NH^{3+} positions.[51] This database was chosen as it contains high resolution X-ray crystallographic structure inclusing carefully positioned hydrogens, thus producing objects corresponding exactly to those iBP is designed to calculate.

Among these structures, 30 structures with number of residues between 21 and 107 have been selected, containing only trans peptidic bonds and corresponding to the following list of 24 proteins: 1aacH, 1benABH, 1bkfH, 1bpiH, 1ckaH, 1cnrH, 1ctjH, 1difH, 1edmBH, 1fxdH, 1igdH, 1iroH, 1isuAH, 1mctIH, 1ptfH, 1ptxH, 1rroH, 256bAH, 2bopAH, 3b5c, 3ebxH, 451cH, bio1rpoH and bio2wrpH. In 3b5c, the N terminal residue THR88 was removed because of missing backbone atoms. On each structure, the conformation of chain A was selected for preparing the iBP input, and in the case multiple conformations have been observed for a residue, the A conformation was selected.

Input values for the iBP calculation

The parameters defining the covalent and improper geometries were taken from the geometric force field PARALLHDG (version 5.3)[52] (Table 1). The atom re-ordering is the same proposed in the most recent implementation of iBP[36] (Table 2).

Two sets of values were used for the ϕ and ψ backbone angles: (i) the ϕ_{angl}, ψ_{angl} angles

measured on the X-ray crystallographic structures using VMD,[53] (ii) the ϕ_{dist}, ψ_{dist} angles calculated from the distances dNN and dCC between N and C atoms of successive residues, assuming the covalent geometry uniform and described in Table 1. The dihedral or pseudo-dihedral angle Ω between ordered atoms i-3,i-2,i-1 and i, is determined using the cosine law from a trihedron:[37]

$$\cos\Omega = \frac{\cos\gamma - \cos\alpha\cos\beta}{\sin\alpha\sin\beta}, \tag{1}$$

where α is the angle between atoms (i-3,i-2,i-1), β is the angle between atoms (i-1,i-2,i), and γ is the angle between atoms (i-3,i-2,i). For Ω angles being ϕ or ψ, the angles α, β and γ are calculated from the bond lengths and bond angles among heavy backbone atoms, as well as the distances dNN and dCC between successive residues along the protein sequence.

The input restraints for iBP processing of sub-peptides are: (i) the restraints corresponding to the bond length and bond angles of the force field PARALLHDG (version 5.3);[52] (ii) the backbone angles ϕ and ψ, determined as described previously; (iii) the distances between Cα atoms located at the two extremities residues of each sub-peptides; (iv) the long-range distances between Cα atoms of the residues located at the middle of each fragment. Several error bounds have been tested: error of $\pm$ 10°, 20° and 30° for the angles ϕ and ψ, error of $\pm$ 3 Å for the Cα-Cα distance between extremities of peptide fragments, and error of $\pm$ 5 Å for the long-range Cα-Cα distance between sub-peptides.

Interval Branch-and-Prune calculation of peptide fragments

As the TAiBP approach intends to explore the conformation of protein backbone, the processed protein is initially converted to a poly-Alanine chain. The protein is then divided in 15-residues peptide fragments, two successive fragments having a sequence of 5 superimposed residues. The peptides are then assembled together to produce protein conformations, as it will be described in the next subsection.

For each fragment, the iBP tree of possible conformations is systematically explored. We employ the most recent implementation of IBP[36] (in the C programming language), which is tuned for the calculation of protein conformations based on the force field knowledge for the covalent geometry. The tree branching is performed on the ϕ and ψ backbone angles.

For any vertex i in the order, we seek the embedded coordinates $\mathbf{x}_i$, given distances between the three preceding vertices $i-1$, $i-2$ and $i-3$. As described above, the distances $d_{i,i-1}$, $d_{i,i-2}$ and $d_{i,i-3}$ are known, where $d_{i,i-3}$ is potentially an interval. The remaining distances in the clique formed by four vertices are given by the coordinates $\mathbf{x}_{i-1}$, $\mathbf{x}_{i-2}$ and $\mathbf{x}_{i-3}$. We shall introduce the variables d_i, θ_i and τ_i and σ_i for embedding vertex i, where d_i denotes $d_{i,i-1}$.

Given d_i, θ_i, τ_i, and σ_i, the embedded coordinates of vertex i are given by the following equation:

$$\mathbf{x}_i = \mathbf{p}_1 + \tau_i\,\mathbf{p}_2 + \sigma_i\sqrt{1 - \tau_i^2}\,\mathbf{p}_3$$

where $\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3 \in \mathbb{R}^3$ depend only on $\mathbf{x}_{i-1}$, $\mathbf{x}_{i-2}$, $\mathbf{x}_{i-3}$, d_i and θ_i,

$$\mathbf{p}_1 = -\left(\frac{d_i}{\|\mathbf{r}_{12}\|}\right)\left(\left(\cos(\theta_i) - \frac{\|\mathbf{r}_{12}\|}{d_i}\right)\mathbf{x}_{i-1} - \cos(\theta_i)\,\mathbf{x}_{i-2}\right)$$

$$\mathbf{p}_2 = -\left(\frac{d_i}{\|\mathbf{r}_{12}\|}\right)\left(\frac{\sin(\theta_i)}{\|\mathbf{r}_{12} \times \mathbf{r}_{23}\|}\right)\left(\|\mathbf{r}_{12}\|^2\,\mathbf{r}_{23} - (\mathbf{r}_{12} \cdot \mathbf{r}_{23})\,\mathbf{r}_{12}\right)$$

$$\mathbf{p}_3 = -\left(\frac{d_i}{\|\mathbf{r}_{12}\|}\right)\left(\frac{\sin(\theta_i)}{\|\mathbf{r}_{12} \times \mathbf{r}_{23}\|}\right)\|\mathbf{r}_{12}\|\,(\mathbf{r}_{12} \times \mathbf{r}_{23})$$

and we have introduced $\mathbf{r}_{12}, \mathbf{r}_{23} \in \mathbb{R}^3$ for notational simplicity,

$$\mathbf{r}_{12} = \mathbf{x}_{i-1} - \mathbf{x}_{i-2}$$

$$\mathbf{r}_{23} = \mathbf{x}_{i-2} - \mathbf{x}_{i-3}$$

The angle θ_i is obtained from the cosine law using the relevant distances,

$$\theta_i \triangleq \cos^{-1}\left(\frac{d_i^2 + d_{i-1,i-2}^2 - d_{i,i-2}^2}{2d_i\, d_{i-1,i-2}}\right) \tag{2}$$

The dihedral angle τ_i formed by the four vertex is determined from the cosine law for a trihedron:[37]

$$\tau_i = \frac{2d_{i-2,i-1}^2\left(d_{i-3,i-2}^2 + d_{i-2,i}^2 - d_{i-3,i}^2\right) - d_{i-3,i-2,i-1}d_{i-2,i-1,i}}{\sqrt{4d_{i-3,i-2}^2 d_{i-2,i-1}^2 - d_{i-3,i-2,i-1}^2}\sqrt{4d_{i-2,i-1}^2 d_{i-2,i}^2 - d_{i-2,i-1,i}^2}} \tag{3}$$

where

$$d_{i-3,i-2,i-1} \triangleq d_{i-3,i-2}^2 + d_{i-2,i-1}^2 - d_{i-3,i-1}^2$$

$$d_{i-2,i-1,i} \triangleq d_{i-2,i-1}^2 + d_{i-2,i}^2 - d_i^2$$

The variable $\sigma_i \in \{-1, +1\}$ is the sign of $\sin\omega_i$. When ω_i is known from either protein chemistry or measurement, we may directly compute τ_i, as well as the sign $\sigma_i \in \{-1, +1\}$.

The branching was reduced to one branch in the case when the dihedral angle between the four concerned atoms corresponds to an improper angle ω_i (Table 1). Indeed, in that case, the absolute value and the sign of improper angle being known, its cosine and sine values are unambigously defined, which generates a single branch in the tree.

The number of saved conformations is reduced by applying a RMSD filter of 3 Å between two successively saved conformations. In order to avoid pruning due to slight discrepancy between distance restraints, a tolerance of 0.05 Å has been added to the bounds of distance intervals. The minimum discretization factor, which is the minimum ratio between each distance interval to the number of tree branches generated within the interval, was set to 0.05 Å, in order that the branching does not oversample small intervals. The maximum number of branchs at each tree level was 4. No pruning due to the van der Waals radii of the force field protein-allhdg5-4 PARALLHDG (version 5.3)[52] was applied. A maximum number

of saved conformations of 10^9 was permitted for each iBP run. The solutions are stored in a multiframe dcd format.[54]

Assembling the peptide fragments and clustering

The generated conformations of neighbouring peptide fragments in the protein sequence are then assembled by superimposing the five last and initial residues of the fragments located first and second in the sequence. The conformations of fragments are assembled to each other by root-mean-square superimposition of backbone atoms located in the five superimposed residues. For each superimposition, the residue number for which the smallest distance was observed between corresponding atoms in the two peptides is used to decide where to stop with the first peptide and to continue with the second one. The assembled conformation is then submitted to two pruning devices: (i) a device checking whether there is no clash between the two fragments, i.e. no Cα atoms closer than 1 Å, (ii) a device checking that long-range Cα-Cα distance restraints between peptide middle residues are verified. The fragment assembly is implemented using a python script based on the MDAnalysis[55,56] and numpy[57] python packages.

To scale down the combinatorial of the calculation, a clustering approach, the **S**elf-**O**rganizing **M**aps (SOM),[58–61] which are unsupervised neural networks, were used to reduce the number of conformations. The method was implemented through a set of python scripts.[62] The SOM approach was used after a iBP calculation or after an assembly step as soon as the number of saved conformations was larger than 1000. The conformations sampled by iBP were encoded from the distances d_{ij} calculated between the n C$_\alpha$ atoms of the fragment, by diagonalizing the covariance matrix C:

$$C_{i,j} = \frac{1}{n} \sum_{k=1}^{n} \sum_{l=1}^{n} (d_{i,k} - \bar{d}_i)(d_{l,j} - \bar{d}_j) \tag{4}$$

where $\bar{d}_i = \frac{1}{n} \sum_{j=1}^{n} d_{i,j}$. The matrix C can be replaced by its four largest eigenvalues along

wih the corresponding eigenvectors. The eigenvalue and eigenvector descriptors were used to train a periodic Euclidean 2D self-organizing map (SOM), defined by a three-dimensional matrix. The first two dimensions were chosen to be 100×100 and defined the map size.

The self-organizing maps were initialized with a random uniform distribution covering the range of values of the input vectors. At each step, an input vector is presented to the map, and the neuron closest to this input is updated. The maps are trained in two phases. During the first phase, the input vectors are presented to the SOM in random order to avoid mapping bias with a learning parameter of 0.5, and a radius parameter of 36.[63] During the second phase, the learning and radius constants were decreased exponentially from starting values 0.5 and 36, respectively, during 10 cycles of presentation of all the data in random order. Once the caculation of the SOM has been realized, the conformations corresponding to local maxima of homogeneity, are detected and the total set of conformation is replaced by these representative conformations.

Once the full protein chain has been reconstructed by iterative assembly of growing fragments, the final protein conformations were compared to the initial structure, by calculating the root-mean-square (RMSD) between Cα atomic coordinates (Å).

Results

Probing the hypothesis of uniform covalent geometry

The 24 structures extracted from the database of Word et al[51] have been processed to analyze the geometry of covalent angles (Figure 1a). The distributions of covalent angles between C-N-Cα (blue curve), N-Cα-C (magenta curve) and Cα-C-N (green curve) (Figure 1a) are centered on about 121.3°, 110.6° and 116.8°, with standard deviations of about 2.2°, 3.0° and 2.2°. These distributions agrees with the ones observed by Hinsen et al:[64] C-N-Cα (121.4° $\pm$ 1.6°), N-Cα-C (111.1° $\pm$ 2.9°), Cα-C-N (116.6° $\pm$ 1.3°), and in agreement to this work, the largest width is observed for the bond angle N-Cα-C.

In order to verify whether the variations in bond angles could arise from variations in protein internal mobility, the bond angle values were compared to the B factor value averaged on the corresponding residues (Figure 1b). There are no correlation between the values of bond angles and the B factors, which shows that the variations of covalent geometry cannot be assigned to differences in protein internal mobility.

The variations in covalent geometry were then plotted (Figure 1c-e) along the positions of protein residues in the Ramachandran diagram, by coloring the point describing the (ϕ, ψ) angle values of a given residue, according to the values of the residus bond angles. The Ramachandran plots are multi-colored according to the values of bond angles C-N-Cα (Figure 1c), N-Cα-C (Figure 1d) and Cα-C-N (Figure 1e). All α-helix regions, around (-60°,-45°), display a quite monocolor pattern, with values mostly in the range 100°-105° for angle C-N-Cα (Figure 1c), in the range 125°-130° for angle N-Cα-C (Figure 1d) and in the range 120°-125° for angle Cα-C-N (Figure 1e). At contrary, the β-strand region and the loops region of each diagram display a larger heterogeneity in bond angle values than the α-helix region. This heterogeneity has certainly a strong influence on the overall tertiary structures. Indeed, the β strands are extended structures in which local variation can have strong influence on the orientation at long distance. On the other hand, the change of direction of protein backbone can be also very sensitive to local loop structure variation. In that way, both β strand orientations and loop directions have a strong impact on the protein tertiary structure.

In order to investigate the relevance of the uniform geometry hypothesis for the iBP calculation, the ϕ_{angl} and ψ_{angl} values measured on the top100 conformations have been compared to the ϕ_{dist} and ψ_{dist} values obtained in the following way. The bond angles and bond lengths defined in the force field PARALLHDG[52] have been used to determine the distances between atoms related by one of two covalent bonds. These distances along with distances between nitrogen and carbonyls located in successive residues along the protein sequence, have been used to determine the backbone angles ϕ_{dist} and ψ_{dist} using the spherical

cosine relationship (Eq. 1).

For each residue K, the cumulative sums of the differences between *angl* and *dist* backbone angles for residues i, i varying from 1 to K, were calculated:

$$\Phi_K = \sum_{i=1}^{K}(\phi_i^{\mathsf{angl}} - \phi_i^{\mathsf{dist}}) \tag{5}$$

$$\Psi_K = \sum_{i=1}^{K}(\psi_i^{\mathsf{angl}} - \psi_i^{\mathsf{dist}}). \tag{6}$$

In Figure 2, the variations of Φ_K (green curves) and of Ψ_K (magenta curves) have been plotted along K, for the 24 studied proteins. The most important observation from these curves is that Φ_K and Ψ_K display extraordinary large variations along protein primary sequence. These variations extend from about $100°$ for the proteins 1benABH, 2bopAH, 1okAaH, bio1rpoH, up to several hundreds of degrees. The drift of Φ_K and Ψ_K depends of course on the total number of residues in the protein. Another observation is that Φ_K and Ψ_K curves do not display the same features. Φ_K curves are positive and increase along K, whereas Ψ_K curves are mostly negative and decrease along K.

To summarize, the analysis of protein structures involved in the present validation reveals that the hypothesis of uniform covalent geometry is far from being verified even in high-resolution crystallographic structures as the ones selected in this database[51] with resolution in the range 1.0-1.5 Å. Due to this point, the differences between angles ϕ_{angl}, ψ_{angl} and ϕ_{dist}, ψ_{dist} display large cumulative drift along the protein sequence.

Exploring the conformational space of fragments using iBP

iBP calculations were performed on individual peptides spanning the analyzed proteins and the obtained results are presented in Figure 3. The run durations (Figure 3a) are in the range of 1 to 10^5s, which corresponds to a maximum duration of about one day for an individual calculation. Given that calculations of peptides spanning one individual protein

can be launched in parallel, this duration is not prohibitive. Distributions of run durations centered on on 10^2s, are observed in the case of error intervals on backbone angles of $20°$ (blue curve) and $40°$ (magenta curve), centered on 10^2s, whereas the durations are shifted to larger values in the 10^3-10^4s range for an error of $60°$ (green curve).

The tree sizes were reduced using the signed values of improper angles (Table 1), and are in the range 10^5-10^9. For each run, a maximal number of 10^9 conformations to generate was required as input. The number of conformations generated during each run (Figure 3b) is in the range of 1 to 10^5 which is several order of values smaller than this maximal number. All trees have thus been completely parsed during the iBP calculations.

The number of generated (Figure 3b) and saved (Figure 3c) conformations increase linearly along the run duration. The number of generated conformations is in the range 10^4-10^8, whereas the number of saved conformations is in the range 1-10^5. For the largest error intervals ($40°$ and $60°$, magenta and green dots), similar numbers of conformations are generated (Figure 3b) and these numbers depend mainly on the duration of the run. In the case of the smallest interval width (blue dots in Figure 3b), few runs display much smaller numbers of generated conformations. The number of saved conformations (Figure 3c) is of course smaller by two or three orders of magnitude from the number of generated conformations, but is also more dispersed. These numbers sample superimposed ranges for error intervals of $20°$ and $40°$ (blue and magenta dots), whereas they sample larger values for error interval of $60°$ (green dots). The result of tree parsing thus depends only slightly on the interval width, but a difference of behavior for the number of saved conformations is observed for error interval larger than $40°$ (Figure 3c), and this difference is also visible in the run duration (Figure 3a).

In NMR of biomolecules, the conversion of experimental information into interval restraints is a major problem.[6,7] In iBP, a discretisation of distance intervals transforms the DGP into a discrete problem. But, the quantity of lost information during interval discretisation is an important question to explore and was analyzed (Figure 3d) through the

discretisation factor, which is the ratio between each distance interval to the number of tree branches generated within the interval. The standard deviation of this factor is plotted along its average value, both being calculated for an individual iBP tree. Overall, one should notice that the largest discretisation factors are smaller than 0.25 Å. This number is of the order of the positional error in atomic coordinates for high-resolution X-ray crystallographic structures.[65,66] The discretisation if distance intervals thus remains with the X-ray crystallographic uncertainties on atomic positions, and does not induce major lost of information in the iBP calculations.

For the various error intervals, the couples of average and standard deviation values for the discretisation factor (Figure 3d) are clustered around different points: (0.11, 0.11) Å for the intervals of 20°, (0.18, 0.08) Å for the intervals of 40°, (0.23, 0.07) Å for the intervals of 60°. The average value increases with the interval width on ϕ and ψ angles. More surprisingly, the standard deviations decrease with the interval width: this is due to the calculation inputs. Indeed, the maximum number of branches is limited by 4 in all calculations, but the discretisation factor should be always larger than a threshold of 0.05 Å. These two parameters induce the saturation of the number of branchs for large widths as 60°. At the contrary, for smaller interval widths, the maximum number of branches is not attained due to the required threshold. This induces a larger variability between the number of tree branches and the resulting standard deviation is larger in the case of smaller errors.

During each iBP calculation, the conformations generated by branching on the ϕ and ψ intervals are then pruned or not according to the violation or to the verification of the distance interval between the Cα atoms located at the N and C terminal residues. Percentages of pruned conformations (Figure 3e) is observed up to 100%. In the case of interval widths of 20° and 40°, three and one runs do not provide any solutions. As all these runs were performed using as input ϕ_{dist} and ψ_{dist} backbone angles, the pruning of all solutions is due to the inconsistency between the ϕ and ψ angle restraints and the extremities distance restraint. This inconsistency arises directly from the non-uniform covalent geometry described in the

first section and is amplified by the use of a small error on backbone angle restraints.

Similarly to the number of runs without solutions, contrasted distributions are observed (Figure 3e) for the percentages of pruned conformations, depending on the width of intervals on backbone angles. For the smallest width (20°: blue curve), the percentage of pruned conformations displays a weak maximum at around 25%, but a non-negligible number of runs display percentages of pruned conformations in the 50-90%. Such high pruning percentages arise because in the case of narrow intervals on backbone angles, the hypothesis of uniform covalent geometry made by iBP has much more chances to induce solutions which do not verify the distance restraint between peptide extremities. For the larger interval widths on backbone angles (40°: magenta curve, 60°: green curve), the distribution is much more focused on respective ranges of 40-70% and 60-80%, which are larger than the 25% observed for the error of 20°. The global picture is that the increase of intervals on backbone restraints induces more pruning, but a better consistency between the obtained conformations and the distance restraint as the discrepancy arising from uniform covalent geometry hypothesis is counter-balanced by larger intervals for ϕ and ψ andgles. This behavior is promising for the application of iBP to cases with error on restraints similar to situations encountered with experimental data.

After generating peptide conformations using iBP, a procedure based on the self-organizing map[61] is used to cluster the conformations and to extract representative ones. The distributions of the number of representative conformations (Figure 3f) are centered on the 0-100 and 0-50 range for the widths of 20° and 40°. In the case of the larger width 60°, much larger numbers of representative conformations can be obtained, up to 250. The average number of representative conformations extracted from the SOM clustering of an iBP run on peptide fragment, is of the order of 10^2, which makes the number of maximum combinations of peptides during the step of fragments assembly to be about 10^4, and will permit to overcome the combinatorial explosion, as it will be shown in the following.

Efficiency of the TAiBP assembly strategy

Starting from the conformations of individual peptides generated by iBP, the individual peptide conformations were superimposed on the backbone atoms of their last and initial five residues, in order to grow step by step the protein structure from the N terminal to the C terminal extremity. The proposed fragment assembly is then conserved or pruned according to two successively applied criteria: (i) the clashing criterion tests whether $C\alpha$ atoms of each fragment are farther apart from a given threshold (1 Å), (ii) the pruning distance criterion tests whether distance between the central $C\alpha$ of all inserted peptides is within 5Å of the distances observed in the initial PDB structure. Several assembly strategies have been used: (a) the fragments are added one by one from the N terminal to the C terminal extremities of the protein, (b) all possible assemblies of two fragments are formed along the sequence, and then assembled together successively from N to C terminal, (c) all possible assemblies of three fragments are formed along the sequence, and then assembled together successively from N to C terminal. Depending on the protein target, one approach can be more efficient than the others, but no general trend of efficiency for one strategy was found during the analysis, so the results of the three strategies are presented together.

Some statistics on the assembly steps of TAiBP are presented in Figure 4. The numbers of distance pruning events (#DistPruning), of clash pruning events (#ClashPruning) and of processed conformations (#Processed) are plotted along the size of the processed peptide in Figures 4a,b,c. The numbers of distances and clash pruning events (Figure 4b) are mostly in the range 10^2-10^4 for all fragment sizes and interval of backbone restraints. In addition, these numbers increase of more than one order of value when the fragment size changes from 25 residues (assembly of two fragments) to larger values. Larger pruning experienced in the case of larger fragments is probably induced by the excluded volume effect arising from the construction of the protein fold.

For fragment sizes larger than 25 residues, the numbers of pruning events (#DistPruning and #ClashPruning) are mostly around 10^3-10^5 (Figures 4a,b), in a range similar to the

number of processed (#Processed) conformations (Figure 4c) which proves a large efficiency of pruning events for reducing the ensemble of solutions. Interestingly, distance pruning (Figure 4a) and clash pruning (Figure 4b) events are in similar range, displaying thus similar efficiency to reduce the number of solutions.

The parameters #DistPruning and #ClashPruning (Figures 4a,b) both shift toward larger values when increasing the interval widths on backbone angles. For widthes of 40° and 60° (magenta and green dots), the shift is steeper for the distance pruning events (Figure 4a) than for the clash pruning events (Figure 4b), whereas similar shifts are obtained for the interval width of 20° (blue dots). The widening of intervals has thus a stronger effect on the distance restraints between the peptide fragments than on the clash level. Finally, for fragments larger than 50 residues, the assembled fragments vanish except for the smaller intervals of 20° (blue dots), due to a pruning of all solutions. This pruning is the consequence of the discrepancy between non-uniform covalent geometry observed in the PDB structures and of the hypothesis of uniform covalent geometry made in the frame of iBP calculations.

The two by two comparison of the events of distance and clash pruning, and of the number of processed conformations reveal the following trends (Figure 4d,e,f). The numbers of pruning events by clashes or by distances do not display any correlation (Figure 4d). At the contrary, #ClashPruning displays a quite strong correlation with #Processed (Figure 4e), specially for the largest angle interval (60°: green points). A similar tendency is observed for #DistPruning with two superimposed behaviors (Figure 4f): a correlation similar to the one observed for #ClashPruning, and other points with relatively smaller numbers of distance pruning events. This second set of points corresponds mostly to the case of the fragments of 25 residues. Indeed, as these fragments are much smaller than the full protein, they have less chance to be rejected by pruning distance information.

Each assembled fragment has been compared to the corresponding region in the PDB target structure. This comparison was performed using RMSD (Å) between coordinates of heavy backbone atoms (Figure 5a,b) and not the TM score.[67,68] Indeed, the statistical

validation of TM score was performed on protein structures larger than 80 residues,[68] which do not correspond to the set of proteins studied here. For each fragment, maximum and minimum RMSD values are plotted in Figure 5a,b with respect to the fragment size.

In the case of narrow intervals (20°: blue points) on backbone angle restraints minimum RMSD values are mostly smaller than 3.0 Å for all fragments up to 65 residues (Figure 5a). The increase of width in backbone angle intervals induces a drift of RMSD toward larger values: the RMSD drift is limited to 2-4 Å up to 35 residues, but jumps up to 5-6 Å for larger fragments. The threading-augmented iBP procedure proposed here thus allows one to obtain fragment conformations close to the PDB conformations for fragment sizes smaller than 65 residues. In that case, the Φ_K and Ψ_K drifts previously described (Eq. 6 and Figure 2) have thus been overcome.

The maximum RMSD values (Figure 5b) are located in the 5-20 Å range. These maximum values were put in perspective with an analysis[69] in which protein structures were compared to a representative set of protein-like alternative structures generated using threading. Most of the RMSD R values for an N-residue protein fall in the interval: $3.333N^{1/3} - 2.0 \leq R \leq 3.333N^{1/3} + 2.0$, producing distributions of values smaller than 20 Å. This upper limit of 20 Å is similar to the one observed in the present calculation, which means that the TAiBP approach was able to mostly span the possible range of RMSD values.

At the end of TAiBP calculation, the poly-Ala sequence was replaced by the protein specific sequence and the residue sidechains have been added using the relax tool of the Rosetta suite.[43] The obtained conformations have been then analysed (Table 3) and compared (Figure 6) to the conformation of the protein present in the database top100. From the 24 top100 structures initially processed, the 29 calculations realized on the 7 proteins smaller than 50 residues (Figure 6) display conformations calculated with TAiBP close to top100 conformations. Indeed, 18 runs display RMSD to initial top100 conformation smaller than 3 Å and 23 runs display RMSD smaller than 3.7 Å. Negative Rosetta scores calculated with the model[70] were obtained for all calculations, except the calculations on 1mctIH with

an interval width of 60°.

For a given protein, the origin of ϕ and ψ restraints: angles or distances, introduced in the subsection "Probing the hypothesis of uniform covalent geometry" display various influences (Table 3) on the cordinate RMSD between the TAiBP and top100 conformations. For 1ben1BH and 1bpiH, the RMSD is smaller if the ϕ and ψ target values were extracted from the distances between successive nitrogen amide and carbonyl atoms, assuming uniform covalent geometry. At the contrary, 1fxdH, 1cnrH, 1edmBH display the opposite patterns. For the other proteins, the influence of ϕ and ψ origin on the coordinate RMSD displays a mixed pattern.

The distribution of RMSD values, calculated on the whole sets of conformations obtained for a given TAiBP calculation (Figure 7) span values up to 10 Å. This upper bound is smaller than the one observed for the maximum RMSD in Figure 5b. If the angles ϕ, ψ are measured on the initial top100 conformation (ϕ_{angl} and ψ_{angl}), the increase of interval width induces a drift of RMSD toward larger values (magenta, brown and orange curves). The pattern is less clear if ϕ, ψ restraints are of *distances* origin, ie. calculated from measured distances in top100 conformations: in that case, many RMSD curves (blue, green and cyan curves) are more or less superimposed whatever is the interval width. The different proteins display quite different RMSD distributions which some distributions quite centered to a narrow interval and other much wider. These contrasted features arise from the various efficiencies of pruning long-range distances in the frame of different 3D protein topologies.

Calculations with hypothesis of uniform covalent geometry

The covalent geometry of the 1benABH, 1cnrH, 1edmBH, 1fxdH, 1igdH, 1isuAH, 1mctlH, bio1rpoH, 1bpiH conformations obtained using TAiBP and then relaxed using Rosetta displays some characteristics (Figure 8) quite different from the one analyzed on the top100 conformations at the beginning of the present work (Figure 1). Indeed, the distribution of covalent angles (Figure 8a) is much thinner, although some invidual values can display quite

large drifts from the distribution centers (Figure 8b). These large drifts are observed in the protein regions in which two neighboring peptide fragments were superimposed. The ω dihedral values (Figure 8c) are distributed around 180° and -180°, in a similar way than in the initial structures (data not shown).

The comparison of the Ramachandran plots between Figures 1 and 8 reveal two differences. First, the TAiBP Ramachandran plots display a narrower range of colors than the top100 ones in agreement with the thinner distribution of covalent angles. Second, the (ϕ,ψ) distributions are fuzzier in Figures 8d-f than in Figures 1c-e. The convergence toward a uniform covalent geometry is thus accompanied by a expansion of the regions sampled in the Ramachandran diagram. This expansion of allowed Ramachandran regions has been also observed in the recent analysis of allowed Ramachandran regions.[71] In agreement with covalent geometry close to uniformity, the plots of cumulative sum of the differences: Φ_K and Ψ_K (Eq. 6) display much smaller drifts on the TAiBP conformations with a large majority of the values ranging between -50° and 50° (data not shown).

The protein conformations generated by the TAiBP approach and relaxed with Rosetta (Table 3 and Figure 6), have been used as target conformations for a new run of the TAiBP approach, in order to investigate whether, in the case of mostly uniform covalent geometry, different results could be obtained. In a way similar to the previous TAiBP run, the coordinate RMSD (Å) between the new target and TAiBP fragments display a drift toward larger values for increasing fragment size (data not shown). The minimal and maximal RMSD distributions calculated for the reconstructed full chains of protein targets (Figure 9) show that for all targets except 1fxdH, the distribution of minimal values are mostly in the 1-4 Å for the interval width of 20° (blue curves). For all targets except 1bpiH and on a lesser extend 1cnrH, the increase of interval width do not have strong impact on the minimal RMSD distribution (full lines). Using target conformations closer to the hypothesis of uniform covalent geometry thus reduces the impact of increasing the interval width for ϕ and ψ values. The distributions of maximal RMSD values (dashed lines) displays more variability

than minimum RMSD distributions. Unsurprisingly, these distributions shift toward larger values and/or become broader in the case of increased interval width for ϕ and ψ values.

Discussion-Conclusion

A method has been described to generate the protein structure by systematically exploring all possible conformations of the protein. This method is based on a threading-augmented interval Branch-and-Prune (TAiBP) approach in which the interval Branch-and-Prune (iBP) [21,37,72] is first used to systematically explore the conformations of 15-residues peptides fragments of the protein, followed by the construction of protein structure by a systematic assembly of the fragments conformations, and by pruning conformations displaying clashes and violations of few long-range distance restraints.

This two steps approach along with clustering using self-organized maps [61] allows to overcome the combinatorial explosion arising from the exponential complexity of the iBP algorithm. The duration of a total calculations is of the order of tenths of hours. This could be even speed up by using compiled language in place of python scripts used for fragments assembly and clustering.

The largest problem to which the proposed approach faces is the non-uniformity of covalent geometry among the PDB structures. This aspect of PDB structures looks very minor as it involves only few degrees variations among the covalent angles, but, as this non-uniformity induces biases in the direction of extended and loop parts of protein structures, it is obvious that it may have big consequences in fragment assembly. Also, as this non-uniformity is present since the first days of structural biology and is probably deeply related to amino-acid type and the Ramachandran plot, [73–79] it is quite difficult to sort it out. Nevertheless, it should be noticed that attemps have be made [80] to explore the relationship between backbone conformations and covalent geometry.

On the other hand, the iBP approach on which we based the conformational sampling of

protein fragments, was developed using the initial hypothesis of a uniform covalent geometry. This hypothesis permitted to set up an algorithm which displays good scaling properties in case of a sufficient number of exactly know inter-atomic distances.[21] Modifying the algorithm to take into account possible variations in the covalent geometry would increase enormously its complexity. Nevertheless, the very recent development of residue-specific force fields[30–33] opens new avenues for taking into account these aspects.

The calculations performed using the TAiBP approach were validated by detecting whether iBP provides at least one solution close to the target solution. This detection was performed using the coordinate RMSD between backbone atoms. For most of the proteins smaller than 50 residues and interval widthes of 20° for backbone angles, solutions were obtained with RMSD values smaller than 4 Å. Larger protein sizes and/or larger interval widths induce drift in the obtained conformations, which usually conduct to a pruning of all conformations because the long-range distance restraints are no more verified.

In order to disentangle the relative effects of interval widthes and of non-uniform geometry, additional calculations were performed using iBP outputs as target structures. For these calculations, the drift effect induced by larger interval widthes is mitigated.

Acknowledgments

Thérèse Malliavin thanks the Pasteur Foundation for postdoctoral support of Dr Bradley Worley and thanks Dr Bradley Worley for its implementation of iBP and Dr Guillaume Bouvier for his support in python scripting. The authors wish to thank Institut Pasteur, Ecole Polytechnique, CNRS, FAPESP and CNPq for financial support.

References

(1) Nilges, M.; Clore, G. M.; Gronenborn, A. M. Determination of three-dimensional structures of proteins from interproton distance data by dynamical simulated annealing from

a random array of atoms. Circumventing problems associated with folding. *FEBS Lett.* **1988**, *239*, 129–136.

(2) Pearlman, D. A. FINGAR: A new genetic algorithm-based method for fitting NMR data. *J. Biomol. NMR* **1996**, *8*, 49–66.

(3) Rieping, W.; Habeck, M.; Nilges, M. Inferential structure determination. *Science* **2005**, *309*, 303–306.

(4) Habeck, M.; Nilges, M.; Rieping, W. Bayesian inference applied to macromolecular structure determination. *Phys Rev E Stat Nonlin Soft Matter Phys* **2005**, *72*, 031912.

(5) Habeck, M.; Nilges, M.; Rieping, W. Replica-exchange Monte Carlo scheme for bayesian data analysis. *Phys. Rev. Lett.* **2005**, *94*, 018105.

(6) Rieping, W.; Habeck, M.; Nilges, M. Modeling errors in NOE data with a log-normal distribution improves the quality of NMR structures. *J. Am. Chem. Soc.* **2005**, *127*, 16026–16027.

(7) Habeck, M.; Rieping, W.; Nilges, M. Weighting of experimental evidence in macro-molecular structure determination. *Proc. Natl. Acad. Sci. U.S.A.* **2006**, *103*, 1756–1761.

(8) Nilges, M.; Bernard, A.; Bardiaux, B.; Malliavin, T.; Habeck, M.; Rieping, W. Accurate NMR structures through minimization of an extended hybrid energy. *Structure* **2008**, *16*, 1305–1312.

(9) Bernard, A.; Vranken, W. F.; Bardiaux, B.; Nilges, M.; Malliavin, T. E. Bayesian estimation of NMR restraint potential and weight: a validation on a representative set of protein structures. *Proteins* **2011**, *79*, 1525–1537.

(10) Rosato, A. et al. Blind testing of routine, fully automated determination of protein structures from NMR data. *Structure* **2012**, *20*, 227–236.

(11) Habeck, M.; Rieping, W.; Linge, J. P.; Nilges, M. NOE assignment with ARIA 2.0: the nuts and bolts. *Methods Mol. Biol.* **2004**, *278*, 379–402.

(12) Linge, J. P.; Habeck, M.; Rieping, W.; Nilges, M. ARIA: automated NOE assignment and NMR structure calculation. *Bioinformatics* **2003**, *19*, 315–316.

(13) Wurz, J. M.; Kazemi, S.; Schmidt, E.; Bagaria, A.; Guntert, P. NMR-based automated protein structure determination. *Arch. Biochem. Biophys.* **2017**, *628*, 24–32.

(14) Mohan, A.; Oldfield, C. J.; Radivojac, P.; Vacic, V.; Cortese, M. S.; Dunker, A. K.; Uversky, V. N. Analysis of molecular recognition features (MoRFs). *J. Mol. Biol.* **2006**, *362*, 1043–1059.

(15) Cumberworth, A.; Lamour, G.; Babu, M. M.; Gsponer, J. Promiscuity as a functional trait: intrinsically disordered regions as central players of interactomes. *Biochem. J.* **2013**, *454*, 361–369.

(16) Schneider, R.; Blackledge, M.; Jensen, M. R. Elucidating binding mechanisms and dynamics of intrinsically disordered protein complexes using NMR spectroscopy. *Curr. Opin. Struct. Biol.* **2018**, *54*, 10–18.

(17) Salmon, L.; Nodet, G.; Ozenne, V.; Yin, G.; Jensen, M. R.; Zweckstetter, M.; Blackledge, M. NMR characterization of long-range order in intrinsically disordered proteins. *J. Am. Chem. Soc.* **2010**, *132*, 8407–8418.

(18) Hollingsworth, S. A.; Dror, R. O. Molecular Dynamics Simulation for All. *Neuron* **2018**, *99*, 1129–1143.

(19) Havel, T. F. An evaluation of computational strategies for use in the determination of protein structure from distance constraints obtained by nuclear magnetic resonance. *Prog. Biophys. Mol. Biol.* **1991**, *56*, 43–78.

(20) Liberti, L.; Lavor, C.; Maculan, N.; Mucherino, A. Euclidean Distance Geometry and Applications. *SIAM Rev* **2014**, *56*, 3–69.

(21) Lavor, C.; Liberti, L.; Mucherino, A. The interval Branch-and-Prune algorithm for the discretizable molecular distance geometry problem with inexact distances. *J Glob Optim* **2013**, *56*, 855–871.

(22) C Lavor and J Lee and A Lee-St.John and L Liberti and A Mucherino and M Sviridenko, Discretization orders for distance geometry problems. *Optimization Letters* **2012**, *6*, 783–796.

(23) Lavor, C.; Liberti, L.; Donald, B.; Worley, B.; Bardiaux, B.; Nilges, M. Minimal NMR distance information for rigidity of protein graphs. *Discrete Applied Mathematics* **2018**, in press.

(24) Braun, W.; Go, N. Calculation of protein conformations by proton-proton distance constraints. A new efficient algorithm. *J. Mol. Biol.* **1985**, *186*, 611–626.

(25) Abagyan, R.; Totrov, M.; Kuznetsov, D. CMA new method for protein modeling and design: Applications to docking and structure prediction from the distorted native conformation. *J Comput Chem* **1994**, *15*, 488–506.

(26) Guntert, P. Automated NMR structure calculation with CYANA. *Methods Mol. Biol.* **2004**, *278*, 353–378.

(27) Bardiaux, B.; Malliavin, T. E.; Nilges, M.; Mazur, A. K. Comparison of different torsion angle approaches for NMR structure determination. *J. Biomol. NMR* **2006**, *34*, 153–166.

(28) Berkholz, D. S.; Shapovalov, M. V.; Dunbrack, R. L.; Karplus, P. A. Conformation dependence of backbone geometry in proteins. *Structure* **2009**, *17*, 1316–1325.

(29) Moriarty, N. W.; Tronrud, D. E.; Adams, P. D.; Karplus, P. A. A new default restraint library for the protein backbone in Phenix: a conformation-dependent geometry goes mainstream. *Acta Crystallogr D Struct Biol* **2016**, *72*, 176–179.

(30) Zhou, C.; Jiang, F.; Wu, Y. Residue-specific force field based on protein coil library. RSFF2: modification of AMBER ff99SB. *J Phys Chem B* **2015**, *119*, 1035–1047.

(31) Kang, W.; Jiang, F.; Wu, Y. Universal Implementation of a Residue-Specific Force Field Based on CMAP Potentials and Free Energy Decomposition. *J Chem Theory Comput* **2018**, *14*, 4474–4486.

(32) Liu, H.; Song, D.; Lu, H.; Luo, R. Intrinsically disordered protein-specific force field CHARMM36IDPSFF. *Chem Biol Drug Des* **2018**, *92*, 1722–1735.

(33) Choi, J.; Pappu, R. Improvements to the ABSINTH Force Field for Proteins Based on Experimentally Derived Amino Acid Specific Backbone Conformational Statistics. *J Chem Theory Comput* **2019**, *15*, 1367–1382.

(34) L. Liberti and C. Lavor and A. Mucherino, The discretizable molecular distance geometry problem seems easier on proteins. *Distance Geometry: Theory, Methods and Applications. Mucherino, Lavor, Liberti, Maculan (eds.)* **2014**, 47–60.

(35) Cassioli, A.; Bardiaux, B.; Bouvier, G.; Mucherino, A.; Alves, R.; Liberti, L.; Nilges, M.; Lavor, C.; Malliavin, T. An algorithm to enumerate all possible protein conformations verifying a set of distance constraints. *BMC Bioinformatics* **2015**, *16*, 23–37.

(36) Worley, B.; Delhommel, F.; Cordier, F.; Malliavin, T.; Bardiaux, B.; Wolff, N.; Nilges, M.; Lavor, C.; Liberti, L. Tuning interval Branch-and-Prune for protein structure determination. *Journal of Global Optimization* **2018**, *72*, 109–127.

(37) Lavor, C.; Alves, R.; Figueiredo, W.; Petraglia, A.; Maculan, N. Clifford Algebra and

the Discretizable Molecular Distance Geometry Problem. *J Glob Optim* **2015**, *25*, 925–942.

(38) Shen, Y.; Bax, A. Protein backbone and sidechain torsion angles predicted from NMR chemical shifts using artificial neural networks. *J. Biomol. NMR* **2013**, *56*, 227–241.

(39) A. Mucherino and C. Lavor and L. Liberti and E-G. Talbi, A Parallel Version of the Branch & Prune Algorithm for the Molecular Distance Geometry Problem. *IEEE Conference Proceedings, ACS/IEEE International Conference on Computer Systems and Applications (AICCSA10)* **2010**, 1–6.

(40) F. Fidalgo and D.S. Gonçalves and C. Lavor and L. Liberti and A. Mucherino, A Symmetry-based Splitting Strategy for Discretizable Distance Geometry Problems. *Journal of Global Optimization* **2018**, *71*, 717–733.

(41) Das, R.; Baker, D. Macromolecular modeling with rosetta. *Annu. Rev. Biochem.* **2008**, *77*, 363–382.

(42) Ovchinnikov, S.; Park, H.; Kim, D. E.; Liu, Y.; Wang, R. Y.; Baker, D. Structure prediction using sparse simulated NOE restraints with Rosetta in CASP11. *Proteins* **2016**, *84 Suppl 1*, 181–188.

(43) Ovchinnikov, S.; Park, H.; Kim, D. E.; DiMaio, F.; Baker, D. Protein structure prediction using Rosetta in CASP12. *Proteins* **2018**, *86 Suppl 1*, 113–121.

(44) Alipanahi, B.; Krislock, N.; Ghodsi, A.; Wolkowicz, H.; Donaldson, L.; Li, M. Determining protein structures from NOESY distance constraints by semidefinite programming. *J. Comput. Biol.* **2013**, *20*, 296–310.

(45) Huang, Y. J.; Mao, B.; Xu, F.; Montelione, G. T. Guiding automated NMR structure determination using a global optimization metric, the NMR DP score. *J. Biomol. NMR* **2015**, *62*, 439–451.

(46) Carr, J. M.; Whittleston, C. S.; Wade, D. C.; Wales, D. J. Energy landscapes of a hairpin peptide including NMR chemical shift restraints. *Phys Chem Chem Phys* **2015**, *17*, 20250–20258.

(47) Joo, K.; Joung, I.; Lee, J.; Lee, J.; Lee, W.; Brooks, B.; Lee, S. J.; Lee, J. Protein structure determination by conformational space annealing using NMR geometric restraints. *Proteins* **2015**, *83*, 2251–2262.

(48) Khoo, Y.; Singer, A.; Cowburn, D. Integrating NOE and RDC using sum-of-squares relaxation for protein structure determination. *J. Biomol. NMR* **2017**, *68*, 163–185.

(49) Hong, S. H.; Joung, I.; Flores-Canales, J. C.; Manavalan, B.; Cheng, Q.; Heo, S.; Kim, J. Y.; Lee, S. Y.; Nam, M.; Joo, K.; Lee, I. H.; Lee, S. J.; Lee, J. Protein structure modeling and refinement by global optimization in CASP12. *Proteins* **2018**, *86 Suppl 1*, 122–135.

(50) Martin, J. W.; Zhou, P.; Donald, B. R. Systematic solution to homo-oligomeric structures determined by NMR. *Proteins* **2015**, *83*, 651–661.

(51) Word, J.; Lowell, S.; LaBean, T.; Taylor, H.; Zalis, M.; Presley, B.; Richardson, J.; Richardson, D. Visualizing and Quantifying Molecular Goodness-of-Fit: Small-probe Contact Dots with Explicit Hydrogen Atoms. *J Mol Biol* **1999**, *285*, 1713–1733.

(52) R Engh and R Huber, Accurate bond and angle parameters for X-ray protein structure refinement. *Acta Crystallogr A* **1991**, *47*, 392–400.

(53) Humphrey, W.; Dalke, A.; Schulten, K. VMD: visual molecular dynamics. *J. Mol. Graph.* **1996**, *14*, 33–38,27–28.

(54) Humphrey, W.; Dalke, A.; Schulten, K. VMD – Visual Molecular Dynamics. *Journal of Molecular Graphics* **1996**, *14*, 33–38.

(55) Michaud-Agrawal, N.; Denning, E. J.; Woolf, T. B.; Beckstein, O. MDAnalysis: a toolkit for the analysis of molecular dynamics simulations. *J Comput Chem* **2011**, *32*, 2319–2327.

(56) Richard J. Gowers,; Max Linke,; Jonathan Barnoud,; Tyler J. E. Reddy,; Manuel N. Melo,; Sean L. Seyler,; Jan Domaski,; David L. Dotson,; Sbastien Buchoux,; Ian M. Kenney,; Oliver Beckstein, MDAnalysis: A Python Package for the Rapid Analysis of Molecular Dynamics Simulations. Proceedings of the 15th Python in Science Conference. 2016; pp 98–105.

(57) Oliphant TE, *A guide to NumPy*; Trelgol Publishing, USA, 2006.

(58) Kohonen, T. Self-Organized formation of topologically correct feature maps. **1982**, *43*, 59–69.

(59) Kohonen, T. Self-Organizing Maps. **2001**,

(60) Miri, L.; Bouvier, G.; Kettani, A.; Mikou, A.; Wakrim, L.; Nilges, M.; Malliavin, T. E. Stabilization of the integrase-DNA complex by Mg2+ ions and prediction of key residues for binding HIV-1 integrase inhibitors. *Proteins* **2014**, *82*, 466–478.

(61) Bouvier, G.; Duclert-Savatier, N.; Desdouits, N.; Meziane-Cherif, D.; Blondel, A.; Courvalin, P.; Nilges, M.; Malliavin, T. Functional motions modulating VanA ligand binding unraveled by self-organizing maps. *J Chem Inf Model* **2014**, *54*, 289–301.

(62) Bouvier, G.; Desdouits, N.; Ferber, M.; Blondel, A.; Nilges, M. An automatic tool to analyze and cluster macromolecular conformations based on Self-Organizing Maps. *Bioinformatics* **2014**, *31*, 1–3.

(63) Spill, Y.; Bouvier, G.; Nilges, M. A convective replica-exchange method for sampling new energy basins. *J Comput Chem* **2013**, *34*, 132–140.

(64) Hinsen, K.; Hu, S.; Kneller, G. R.; Niemi, A. J. A comparison of reduced coordinate sets for describing protein structure. *J Chem Phys* **2013**, *139*, 124115.

(65) BC Finzel and TL Poulos and J Kraut, Crystal structure of yeast cytochrome c peroxidase refined at 1.7-Å resolution. *The Journal of Biological Chemistry* **1984**, *259*, 13027–13036.

(66) Umena, Y.; Kawakami, K.; Shen, J. R.; Kamiya, N. Crystal structure of oxygen-evolving photosystem II at a resolution of 1.9 . *Nature* **2011**, *473*, 55–60.

(67) Zhang, Y.; Skolnick, J. Scoring function for automated assessment of protein structure template quality. *Proteins* **2004**, *57*, 702–710.

(68) Xu, J.; Zhang, Y. How significant is a protein structure similarity with TM-score = 0.5? *Bioinformatics* **2010**, *26*, 889–895.

(69) Reva, B. A.; Finkelstein, A. V.; Skolnick, J. What is the probability of a chance prediction of a protein structure with an rmsd of 6 A? *Fold Des* **1998**, *3*, 141–147.

(70) Alford, R. F. et al. The Rosetta All-Atom Energy Function for Macromolecular Modeling and Design. *J Chem Theory Comput* **2017**, *13*, 3031–3048.

(71) Hollingsworth, S. A.; Karplus, P. A. A fresh look at the Ramachandran plot and the occurrence of standard structures in proteins. *Biomol Concepts* **2010**, *1*, 271–283.

(72) C Lavor and L Liberti and N Maculan and A Mucherino, The Discretizable Molecular Distance Geometry Problem. *Computational Optimization and Applications* **2012**, *52*, 115–146.

(73) Hollingsworth, S. A.; Lewis, M. C.; Karplus, P. A. Beyond basins: ϕ,ψ preferences of a residue depend heavily on the ϕ,ψ values of its neighbors. *Protein Sci.* **2016**, *25*, 1757–1762.

(74) Moriarty, N. W.; Tronrud, D. E.; Adams, P. D.; Karplus, P. A. A new default restraint library for the protein backbone in Phenix: a conformation-dependent geometry goes mainstream. *Acta Crystallogr D Struct Biol* **2016**, *72*, 176–179.

(75) Brereton, A. E.; Karplus, P. A. On the reliability of peptide nonplanarity seen in ultra-high resolution crystal structures. *Protein Sci.* **2016**, *25*, 926–932.

(76) Moriarty, N. W.; Tronrud, D. E.; Adams, P. D.; Karplus, P. A. Conformation-dependent backbone geometry restraints set a new standard for protein crystallographic refinement. *FEBS J.* **2014**, *281*, 4061–4071.

(77) Hollingsworth, S. A.; Lewis, M. C.; Berkholz, D. S.; Wong, W. K.; Karplus, P. A. $(\phi,\psi)_2$ motifs: a purely conformation-based fine-grained enumeration of protein parts at the two-residue level. *J. Mol. Biol.* **2012**, *416*, 78–93.

(78) Tronrud, D. E.; Karplus, P. A. A conformation-dependent stereochemical library improves crystallographic refinement even at atomic resolution. *Acta Crystallogr. D Biol. Crystallogr.* **2011**, *67*, 699–706.

(79) Berkholz, D. S.; Shapovalov, M. V.; Dunbrack, R. L.; Karplus, P. A. Conformation dependence of backbone geometry in proteins. *Structure* **2009**, *17*, 1316–1325.

(80) Berkholz, D. S.; Krenesky, P. B.; Davidson, J. R.; Karplus, P. A. Protein Geometry Database: a flexible engine to explore backbone conformations and their relationships to covalent geometry. *Nucleic Acids Res.* **2010**, *38*, D320–325.

Table 1

Table 1: Geometric parameters for covalent and improper bonds and angles taken from the force field PARALLHDG (version 5.3).[52]

Bond atoms	Bond length (Å)	Bond angle atoms	Bond angle value (°)
C CH1E	1.525	C CH1E HA	108.9914
CH1E HA	1.080	C CH1E NH1	111.1396
CH1E NH1	1.458	C CH1E NH2	106.9610
CH1E NH2	1.486	C NH1 CH1E	121.6541
C NH1	1.329	CH1E C NH1	116.1998
C O	1.231	CH1E C O	120.8258
C OC	1.249	CH1E NH1 H	119.2367
H NH1	0.980	C NH1 H	119.2489
H NH2	0.980	C NH2 H	118.1853
		HA CH1E NH1	108.0508
		H NH2 CH1E	109.5000
		H NH2 H	107.3000
		NH1 C O	122.9907
		NH2 CH1E HA	108.4800
		NH2 C O	122.6277
		CH1E C OC	118.0611
		OC C OC	123.3548
Improper atoms	Improper angle value (°)	Atom name	Definition of atom type
C CH1E HA HA	-70.4072	CH1E	α carbon
CH1E C NH1 HA	66.2535	C	carbonyl carbon
C NH1 HA HA	-70.8745	O	carbonyl oxygen
HA CH1E HA HA	-66.5692	OC	C-terminal carbonyl oxygen
CH1E C NH1 CH1E	178.0	HA	Hα hydrogen
O C NH1 H	178.0	NH1	amide nitrogen
C CH1E NH1 O	0.0	NH2	N-terminal nitrogen
NH1 CH1E C HA	119.0	H	amide hydrogen
NH2 CH1E C HA	119.0		
NH1 HA CH1E C	121.0		
NH2 HA CH1E C	116.0		
C NH1 CH1E H	180.0		
CH1E C NH1 O	180.0		
NH2 H H CH1E	41.0		
CH1E OC C OC	178.0		

Table 2

Table 2: Atom re-ordering used during the iBP calculation step within the first, the last and the inner residues of the peptide fragment.

Residue type	order
first	N, H1, H2, CA, N, HA, CA, C
inner	N, -O, -CA, -C, N, CA, C, +N,
	-C, N, CA, H1, N, CA, C, HA, C, CA
last	N, -O, -CA, -C, N, CA, C,
	-C, N, CA, H1, N, CA, C, HA,
	C, CA, O, C, O2

Table 3

Table 3: Analysis of the TAiBP conformations further relaxed using the Rosetta suite.[43] The Rosetta energy terms are defined from the ref2015 score function.[70] Origin of ϕ and ψ angles describes whether their target values have been directly measured from the initial top100 conformation (angles) or whether they have been determined from distances measured on the top100 conformation using the hypothesis of a uniform covalent geometry (distances).

Protein	origine ϕ,ψ	interval width on ϕ,ψ	Number of residues	Minimal RMSD value (Å)	Rosetta fa_atr (ref2015)	Rosetta total (ref2015)
1benABH	angles	20°	21	1.1	-55.0	-9.49
1benABH	distances	20°	21	1.0	-62.0	-12.8
1benABH	angles	40°	21	2.2	-59.6	-13.6
1benABH	distances	40°	21	3.1	-58.5	-7.0
1benABH	angles	60°	21	3.7	-50.0	-4.8
1benABH	distances	60°	21	3.0	-52.7	-4.7
1bpiH	angles	40°	55	6.5	-142.2	-16.0
1bpiH	distances	40°	55	5.0	-151.4	-22.4
1bpiH	distances	60°	55	4.7	-168.6	-16.3
bio1rpoH	angles	20°	61	1.4	-207.7	-70.6
1fxdH	angles	20°	55	1.7	-168.2	-39.0
1fxdH	distances	40°	55	5.5	-139.1	-27.8
1fxdH	angles	40°	55	4.5	-150.8	-26.6
1mctIH	angles	20°	25	0.7	-52.7	-6.8
1mctIH	distances	20°	25	2.9	-50.8	-1.9
1mctIH	angles	40°	25	0.8	-53.4	-7.6
1mctIH	distances	40°	25	2.6	-45.9	-1.9
1mctIH	angles	60°	25	2.7	-49.5	9.1
1mctIH	distances	60°	25	2.2	-53.4	0.4
1cnrH	angles	20°	45	2.0	-133.5	-37.6
1cnrH	distances	20°	45	2.0	-123.4	-30.8
1cnrH	angles	40°	45	3.4	-121.9	-24.8
1cnrH	distances	40°	45	5.0	-123.2	-23.6
1cnrH	distances	60°	45	3.4	-115.7	-23.7
1edmBH	angles	20°	35	1.2	-93.9	-25.9
1edmBH	distances	20°	35	2.0	-80.2	-16.6
1edmBH	angles	40°	35	2.4	-78.2	-19.4
1edmBH	distances	40°	35	3.3	-82.6	-18.1
1edmBH	distances	60°	35	2.7	-93.2	-10.6

Figures

Figure 1: Variability of the covalent geometry within the 24 protein structures analyzed from the database top100.[51] (a) Distribution of bond angles (°) between atoms C-N-Cα (blue), N-Cα-C (magenta) and Cα-C-N (green). (b) Comparison between bond angles and B factors averaged on the residue of the atom Cα. The color code is the same in (a) and (b). (c-e) Ramachandran plots of the protein structures colored according to the value of bond angles (°) between atoms C-N-Cα (c), N-Cα-C (d) and Cα-C-N (e). On each plot (c-e), the color scale is defined in the inset legend.

Figure 2: Variations of the Phi_K (green curves) and Psi_K (magenta curves) (°) plotted along the primary sequences of target proteins (see Eq. 6). The top100 ids of the analyzed proteins are labelled on each plot.

Figure 3: Analysis of the iBP runs on peptide fragments. The colors of lines/points correspond to the error interval on backbone angles: 20° (blue), 40° (magenta) and 60° (green). (a) Distribution of the run duration (s) plotted on log scale. Number of generated (b) and saved (c) conformations plotted along the run duration (s). Both axes are in decimal log scale. (d) Standard deviation (Å) of discretization factor along the average discretization factor (Å). (e) Distribution of the percentage of pruned conformations. (f) Distribution of the number of representative conformations obtained by clustering through self-organizing maps.

Figure 4: Analysis of the assembly of protein fragments to build the protein folds. All parameters except the number of residues are expressed in decimal logarithmic scale. The colors of points correspond to the error interval on backbone angles: 20° (blue), 40° (magenta) and 60° (green). Plots of: (a) number of distance pruning versus number of residues, (b) number of clash prunings versus number of residues, (c) number of processed conformations versus number of residues, (d) number of distance prunings versus number of clashe prunings, (e) number of processed conformations versus number of clashe prunings, (f) number of processed conformations versus number of distance prunings.

Figure 5: Comparison of the assembled fragment conformations to the corresponding region in the PDB target structure. For each processed fragment, the minimum (a) and maximum (b) values of the coordinates RMSD (Å) between heavy backbone atoms are plotted along the number of residues. Dashed lines have been added at the levels of 2, 3 and 4 Å in minimum RMSD plot.

Figure 6: Comparison of the TAiBP conformations to the initial top100 conformations. The TAiBP conformations proceed using the relax tool from Rosetta[43] are drawn in read cartoon and superimposed to the target top100 conformations, drawn in green cartoon. Some residues of the protein core are drawn in licorice. Close to each superimposed structures, the origin of the backbone angles restraints ϕ, ψ is given as *distances* or *angles*. If the angles ϕ, ψ are measured on the initial top100 conformation, the restraints are of *angles* origin, whereas if the angles are calculated from measured distances in top100 conformations, assuming a uniform covalent geometry, the restraints are of *distances* origin. The interval widths of ϕ, ψ restraints are also plotted.

Figure 7: Distribution of RMSD values (Å) of each conformation obtained at the end of TaiBP approach with respect to the conformation of the corresponding protein in the database top100.[51] The curves are colored according to the origin of the backbone angle restraints (*distances* or *angle*) introduced in the subsection "Probing the hypothesis of uniform covalent geometry", and to the interval of ϕ, ψ restraints. The color used are: blue (*distances*, 20°), magenta (*angles*, 20°), green (*distances*, 30°), brown (*angles*, 30°), cyan (*distances*, 40°), orange (*angles*, 40°).

Figure 8: Variability of the covalent geometry within the set of the conformations obtained on 1benABH, 1cnrH, 1edmBH, 1fxdH, 1igdH, 1isuAH, 1mctlH, bio1rpoH, 1bpiH using TAiBP and then relaxed using Rosetta. (a) Distribution of bond angles (°) between atoms C-N-Cα (blue), N-Cα-C (magenta) and Cα-C-N (green). (b) Distribution of bond angles (°) between atoms C-N-Cα (blue), N-Cα-C (magenta) and Cα-C-N (green) with increased scale to detect the minority peaks, (c) Distribution of the ω dihedral angle (°). (d-f) Ramachandran plots of the protein conformations colored according to the value of bond angles (°) between atoms C-N-Cα (c), N-Cα-C (d) and Cα-C-N (e). On each plot, the color scale is defined in the inset legend.

Figure 9: Distribution of minimum (full line) and maximum (dotted line) RMSD (Å) values for the TAiBP run using as targets protein conformations generated by the TAiBP approach and relaxed with Rosetta (Table 3 and Figure 6). The calculations were realized with interval of 20° (blue), 40° (green), and 60° (magenta) widths on the ϕ and ψ values.