LETTER

Castration-mediated IL-8 Promotes Myeloid Infiltration and Prostate Cancer Progression

Keywords: androgen receptor, CXCL8, CXCR2, immunotherapy, PMN-MDSCs, myeloid-derived suppresor cells.

Zoila A. Lopez-Bujanda[1,2,3], Michael C. Haffner[1], Matthew G. Chaimowitz[3], Nivedita Chowdhury[1,2,3], Nicholas J. Venturini[3], Aleksandar Obradovic[3], Corey S. Hansen[4], Joanna Jacków[4], Karen S. Sfanos[1,5], Charles J. Bieberich[6,7], Paula J. Hurley[5], Mark J. Selby[8], Alan J. Korman[8,9], Angela M. Christiano[4,10], Angelo M. De Marzo[1,5], and Charles G. Drake[3,11,12].

[1]Department of Pathology, Johns Hopkins University School of Medicine, Baltimore, MD, USA; [2]Bloomberg-Kimmel Institute for Cancer Immunotherapy, Johns Hopkins University, Baltimore, MD, USA; [3]Columbia Center for Translational Immunology, Columbia University Medical Center, New York, NY, USA; [4]Department of Dermatology, Columbia University, New York, NY, USA; [5]Departments of Urology and Oncology, Johns Hopkins University School of Medicine, Baltimore, MD, USA; [6]Department of Biological Sciences, University of Maryland Baltimore County, Baltimore, Maryland; [7]University of Maryland Marlene and Stewart Greenebaum Cancer Center, Baltimore, Maryland; [8]Bristol-Myers Squibb, Redwood City, CA, USA; [9]Current affiliation: Vir Biotechnology, San Francisco, CA; [10]Department of Genetics & Development, Columbia University, New York, NY, USA;

[11]Herbert Irving Comprehensive Cancer Center, Columbia University, New York, NY,

USA; [12]Department of Medicine, Division of Hematology/Oncology, Columbia University,

New York, NY, USA.

Summary

Immunotherapy is a treatment for many types of cancer, primarily due to deep and durable

clinical responses mediated by immune checkpoint blockade (ICB)[1,2]. Prostate cancer is

a notable exception in that it is generally unresponsive to ICB. The standard treatment

for advanced prostate cancer is androgen-deprivation therapy (ADT), a form of castration

(CTX). ADT is initially effective, but over time patients eventually develop castration-

resistant prostate cancer (CRPC). Here, we focused on defining tumor-cell intrinsic

factors that contribute to prostate cancer progression and resistance to immunotherapy.

We analyzed cancer cells isolated from castration-sensitive and castration-resistant

prostate tumors, and discovered that castration resulted in significant secretion of

Interleukin-8 (IL-8) and it's likely murine homolog Cxcl15. These chemokines drove

subsequent intra-tumoral infiltration with polymorphonuclear myeloid-derived suppressor

cells (PMN-MDSCs), promoting tumor progression.   PMN-MDSC infiltration was

abrogated when IL-8 was deleted from prostate cancer epithelial cells using

CRISPR/Cas9, or when PMN-MDSC migration was blocked with antibodies against the

IL-8 receptor CXCR2. Blocking PMN-MDSC infiltration in combination with anti-CTLA-4

delayed the onset of castration resistance and increased the density of polyfunctional

CD8 T cells in tumors. Taken together, our findings establish castration-mediated IL-8

secretion and subsequent PMN-MDSC infiltration as a key suppressive mechanism in the

progression of prostate cancer. Targeting of the IL-8/CXCR2 axis around the time of

ADT, in combination with ICB, represents a novel therapeutic approach to delay prostate

cancer progression to advanced disease.

Main

After primary therapy with surgery or radiation, approximately 40% of prostate cancer

patients develop progressive disease. The standard treatment for recurrent prostate

cancer is androgen-deprivation therapy (ADT), but the majority of these patients

eventually develop castration-resistance (CR). Although some patients with metastatic

castration-resistant prostate cancer (mCRPC) benefit from the cancer vaccine sipuleucel-

T[3], neither CTLA-4 blockade[4,5] nor PD-1 blockade[6] has reliably produced meaningful

clinical responses. Potential reasons for this include a low total mutational burden (TMB)

as well as poor infiltration by CD8 T cells[7].

We and others have shown that ADT initially increases CD8 T cell infiltration into prostate

tumors[8-10], and this response is augmented pre-clinically with anti-CTLA-4[11]. Emerging

data suggest that immune-resistance in prostate cancer involves dysfunctional myeloid

cells known as myeloid-derived suppressor cells (MDSCs) in the tumor microenvironment

(TME)[12,13]. MDSCs secrete IL-23, which acts directly on prostate cancer epithelial cells

to drive castration-resistance[14]. Importantly, the mechanism(s) by which suppressive

MDSCs are recruited to the prostate TME are largely unknown.

To identify immune-related tumor-cell intrinsic factors involved in prostate cancer

progression, we performed expression analyses on murine prostate cancer cells pre- and

post- castration. We used the MCRedAL prostate cancer cell line; an RFP expressing version of the Myc-Cap cell line characterized by *MYC* overexpression[15]. Like human prostate cancer, MCRedAL tumors are initially castration-sensitive (CS), but castration-resistance (CR) develops approximately 30 days after castration (Extended Data Fig. 1a). Pre- and post- ADT tumor cells were sorted to > 96% purity (Extended Data Fig. 1b) and analyzed (Fig. 1a-b and Extended Data Fig. 1c). A number of cytokine and chemokine transcripts were significantly up-regulated post-ADT (Fig. 1b right), including *Cxcl15*, a CXC chemokine with a conserved ELR motif (Extended Data Table 1), which is the likely murine homolog of human *IL-8* (*CXCL8*)[16-19]. qRT-PCR and ELISA assays confirmed the upregulation of Cxcl15 post-ADT at the protein level (Extended Data Fig. 1d). In addition to the chemokines above, GSEA revealed the upregulation of several pro-inflammatory pathways post-ADT (Fig. 1c). *In vitro* experiments using the human androgen-responsive LNCaP cell line corroborated a role for these pro-inflammatory signals, showing that in the absence of androgen, TNFα upregulated IL-8 expression in a dose-dependent manner (Fig. 1d left); while AR signaling in the absence of inflammation did not affect IL-8 expression (Fig. 1d right). These data led to the hypothesis that AR signaling directly suppresses *IL-8* expression in prostate cancer cells. We performed *in silico* ChIP-Seq analyses using human LNCaP cells (GSE83860) and found AR binding at the *IL-8* promoter in the presence of the potent androgen dihydrotestosterone (DHT; Fig. 1e top). This androgen dependent binding was verified by ChIP-qRT-PCR (Fig. 1f).

To further explore the role of AR in IL-8 regulation, we interrogated RNA polymerase binding and transcription marks found at sites of active promoters[20]. In the presence of DHT, binding of RNA polymerase II (pol II), phosphorylated serine 2 RNA polymerase II

(pSer2 pol II) and histone H3 tri-methyl Lys4 (H3K4me3) to the IL-8 locus were

substantially reduced, consistent with reduced transcriptional activity (Fig. 1f).

Conversely, pSer2 pol II binding to the promoter of the well-established AR-regulated

gene *PSA* (*KLK3*), was significantly increased in the presence of DHT as expected

(Extended Data Fig. 1e). Consistent with a role for inflammation, TNFα significantly

increased p65 binding at the *IL-8* (*CXCL8*) promoter in LNCaP cells (Fig. 1e bottom). No

significant binding of AR was detected at the promoters of the chemokines *CXCL1*,

CXCL2, *CXCL5* or *CXCL12* (Extended Data Fig. 1f). These data suggest that AR directly

suppresses *IL-8* expression through repressive AR binding to the *IL-8* promoter. Taken

together, we found that IL-8 transcription is up-regulated by pro-inflammatory signaling,

and down-regulated by AR signaling (Fig. 1g).

We next investigated the effects of ADT on the expression of *Cxcl15 in vivo*, using RNA

in situ hybridization (RISH) to study Myc-Cap tumors.   We found that CR tumors

expressed increased *Cxcl15* as compared to CS tumors, particularly in epithelial

(PanCK[+]) tumor cells (Fig. 2a, Extended Data Fig. 2a). These findings were confirmed *in*

vitro, both at the mRNA and protein level (Fig. 2b). To investigate these findings in the

context of human prostate cancer, we used three paired cell lines in which isogenic CR

lines were derived from CS progenitors. For each pair, the CR line expressed significantly

increased IL-8 as compared to the CS counterpart, both at the mRNA and protein level

(Fig. 2c-d). This observation held across a panel of AR expressing prostate cancer cell

lines; with higher levels of *IL-8* expression in cell lines from castration-resistant disease

(Extended Data Fig. 2b). To test whether AR modulates *Cxcl15* expression in benign

prostate epithelium, we used RISH to study WT mice treated with ADT, and WT mice

treated with ADT followed by testosterone (T) repletion (Extended Data Fig. 2c). These data (Fig. 2e-f) showed increased epithelial *Cxcl15* expression in ADT samples with expression significantly decreased by testosterone repletion (Fig. 2f). This observation was further corroborated by interrogating a dataset (GSE8466) profiling human prostate epithelial cells isolated by laser-capture microdissection (LCM) from men undergoing ADT and ADT with testosterone supplementation. Testosterone repletion significantly reduced *IL-8* mRNA expression (Fig. 2g), supporting the hypothesis that AR signaling down-regulates IL-8 expression. In agreement with these data from benign prostate tissues, we LCM-enriched tumor prostate epithelium from high-risk PCa patients treated with ADT on a neo-adjuvant trial (NCT01696877) and found increased *IL-8* expression as compared to tumors from age and stage-matched untreated controls (Fig. 2h). Taken together, analyses using human tissues strongly support the notion that castration increases *IL-8* expression in prostate epithelial cells.

We next quantified castration-mediated immune infiltration in Myc-Cap allografts (Fig. 3a). Consistent with prior data[11], ADT promoted a transient T cell influx, without significant changes in tumor associated macrophage (TAM) populations (Fig. 3b). By contrast, PMN-MDSC infiltration was significantly increased in CR tumors (Fig. 3b), as verified by IHC (Fig. 3c). We found similar results in human prostate cancer xenografts (Extended Data Fig. 3a-b). PMN-MDSC infiltration also increased in WT mice treated with ADT, but not in WT mice treated with ADT then repleted with testosterone (Extended Data Fig. 3c), supporting a causal relationship between ADT and PMN-MDSC infiltration. Molecular profiling of the infiltrating myeloid cells revealed a signature consistent with functional

PMN-MDSCs, including up-regulation of *IL-1b*, *Arg2* and *IL-23a*[14] (Fig. 3d; Extended Data

Table 2). In particular, increased expression of *IL-23a* and *Cxcr2* was verified by qRT-

PCR (Fig. 3e) and flow cytometry (Extended Data Fig. 3d). To test whether blocking the

IL-8/CXCR2 axis was sufficient to attenuate post-ADT PMN-MDSC infiltration, we treated

prostate-tumor bearing mice with anti-CXCR2 and found that blocking CXCR2

significantly diminished tumor infiltration with PMN-MDSCs in both human (PC3) and

murine (Myc-Cap) immunodeficient and immunocompetent models (Fig. 3f and Extended

Data Fig. 3e-f). To confirm this observation at the genetic level, we used CRISPR/Cas9

to generate human (PC3) and mouse (Myc-Cap) lines that were knocked out for human

IL-8 or the murine IL-8 homolog Cxcl15, respectively. We observed a clear decrease in

PMN-MDSC infiltration in both settings (Fig. 3g and Extended Data Fig. 3e-f).

We next asked whether the supernatants from castration-resistant MCRedAL (CR-

MCRedAL) cells were sufficient to drive PMN-MDSC migration *in vitro*. In line with *in vivo*

results (Fig. 3f-g and Extended Data Fig. 4a-c), we found that PMN-MDSC migrated

towards the supernatant of CR tumors and migration was significantly attenuated by

CXCR2 blockade (Extended Data Fig. 4d). Human prostate cancer (PC3) showed an

identical pattern. To confirm a role for IL-8 in PMN-MDSC migration, we generated IL-8

KO CR-LNCaP (LNCaP-abl) using CRISPR/Cas9. Supernatants from IL-8 KO cells were

significantly attenuated in their ability to promote PMN-MDSC migration (Extended Data

Fig. 4e). These PMN-MDSCs were functional and suppressed CD8 T cell proliferation in

a dose-dependent manner (Extended Data Fig. 4f-i). Although CXCR2 blockade

decreased PMN-MDSC migration, it did not significantly alter their suppressor function

(Extended Data Fig. 4j). Similarly, Cxcl15 loss did not diminish the suppressive function

of PMN-MDSCs (Extended Data Fig. 4k). Taken together these findings reinforce a

functional role for castration-mediated IL-8 secretion in PMN-MDSC migration.

Finally, we investigated the pre-clinical activity of blocking the IL-8/CXCR2 axis at the time

of androgen-deprivation in the Myc-Cap model. Notably, in the absence of

immunotherapy the combination of ADT and CXCR2 blockade was not effective

(Extended Data Fig. 5a). In contrast, combining CXCR2 blockade with ICB (anti-CTLA-

4; Fig. 4a) resulted in significantly increased survival (Fig. 4b). This triple combination

(ADT + anti-CXCR2 + anti-CTLA-4) was effective even when tumors were relatively

advanced (400 mm^3) at the time of treatment (Extended Data Fig. 5b&d). Macrophage

modulation with anti-CSF1R was not effective therapeutically in this setting (Extended

Data Fig. 5c&e). Mechanistically, the increased anti-tumor effects mediated by the

addition of anti-CXCR2 to ADT + anti-CTLA-4 did not appear to be due to increased T cell

infiltration (Fig. 4c and Extended Data Fig. 5f-h), nor due to decreased Treg infiltration

(Fig. 4d), but rather correlated with an increase in polyfunctional effector CD8 T cells in

tumor-draining lymph nodes (TDLN) and spleens (Fig. 4e&f).

In summary, these studies showed that castration mediates increased IL-8 secretion by

prostate cancer epithelial cells by releasing AR-mediated transcriptional repression. IL-8

(and Cxcl15) up-regulation then drives prostate tumor infiltration with PMN-MDSCs. We

found that blocking CXCR2 at the time of androgen-deprivation therapy attenuates PMN-

MDSC infiltration, rendering prostate tumors more responsive to ICB. It is noteworthy

that in other murine models the recruitment of PMN-MDSC and neutrophils may be driven

by other chemokines, including Cxcl1[21] and Cxcl12[22]. Our findings are corroborated by

clinical data showing that PMN-MDSCs accumulate in the blood of patients with advanced

prostate cancer[23-25], and that an intratumoral PMN signature is associated with poor

outcome[26]. Our data are also supported by pre-clinical studies showing that blocking

MDSC function increases the efficacy of ICB in animal models of CRPC[12]. Consistent

with recent data, we found that the PMN-MDSCs infiltrating prostate tumors express IL-

23[14]. We further showed that inhibiting the recruitment of these cells peri-castration

augmented the CD8 T cell effector function initiated by ICB. Based on these findings, we

have initiated a phase 1b/2 trial (NCT03689699) to test whether adding ICB and anti-IL-

8 to a short course of ADT can prevent PMN-MDSC infiltration and delay progression in

men with castration-sensitive prostate cancer. In summary, targeting the IL-8/CXCR2

pathway following ADT in combination with immune checkpoint blockade may represent

a novel treatment paradigm to improve responses to immunotherapy and delay the onset

of castration-resistance.

Acknowledgments

We thank members of the Drake Lab for discussion and insightful comments; F. Veglia

for advice with *in vitro* suppression assays; K.C. Smith, A. Floratos, and the Center for

Computational Biology and Bioinformatics at Columbia University for ChIP-Seq analysis;

S. Coley, T. Swayne, E. Munteanu, and the Confocal and Specialized Microscopy Shared

Resource at Columbia University for help with microscopy; L. Dasko-Vincent from the

Sidney Kimmel Comprehensive Cancer Center Imaging Facility at Johns Hopkins for

support with LCM, J. Pevsner for assistance on protein homology analyses, and B.

Johnson for help with statistical analyses. This study was supported by U.S. Department
of Defense (W81XWH-13-1-0369), U.S. National Institutes of Health National Cancer
Institute (R01: CA127153), the Patrick C. Walsh Fund, the OneInSix Foundation, and the
Prostate Cancer Foundation. Research reported in this publication was performed in the
CCTI Flow Cytometry Core, supported in part by the Office of the Director, National
Institutes of Health under awards S10OD020056. H&E/IHC staining and image collection
for this work was performed in the Molecular Pathology Shared Resource and the
Confocal and Specialized Microscopy Shared Resource of the Herbert Irving
Comprehensive Cancer Center at Columbia University, supported by NIH grant #P30
CA013696 (National Cancer Institute). The content is solely the responsibility of the
authors and does not necessarily represent the official views of the National Institutes of
Health.

Author contributions

215 Z.A.L.B., M.C.H., M.G.C., N.C., N.J.V., and A.O. performed experiments; C.H., J.J.,
216 C.J.B., P.J.H., M.J.S., and A.J.K. contributed essential reagents; Z.A.L.B., M.C.H., A.M.C.
217 and C.G.D. designed and supervised experiments; M.C.H., K.S.S., and A.D.M.
218 coordinated the study on human samples; C.G.D. supervised the study. Z.A.L.B. and
219 C.G.D. wrote the manuscript, which was edited by all authors.

Author information

222 C.G.D. has stock or ownership interests in Compugen, Harpoon, Kleo, Potenza, and
223 Tizona Therapeutics, and has served as a consultant for Agenus, Dendreon, Janssen

Data Availability

235 The data that support the findings of this study are available from the corresponding

236 author upon reasonable request.

Biological Materials

239 Biological materials used in this study may be requested from the corresponding

240 author, with the exception of anti-CTLA-4 and anti-CXCR2 antibodies which were

241 obtained through an MTA with A.K and M.S.

Figures

Figure 1 | Androgen-Deprivation Therapy (ADT) Increases IL-8 Expression in Prostate Cancer Cells. a, Differential expression profile of tumor epithelial cells isolated from castration-sensitive (CS) and ADT-treated MCRedAL tumor bearing mice. Heatmap showing transcripts 3 standard deviations away from the mean (n = 3 per group). **b,** Differential chemokine expression of tumor epithelial cells isolated from CS and ADT tumor bearing mice (replicate numbers as in **a**). Left, volcano plot showing gene

expression among all MTA 1.0 microarray transcripts. Right, heatmap of normalized

chemokine transcripts. **c,** Hallmarks gene sets pathway analysis post-ADT shows NF-κB

up-regulation post-ADT. **d,** qRT-PCR quantification of *IL-8* in LNCaP cells cultured at

indicated concentrations of TNFα and DHT, cells cultured in androgen-free media as

described in materials and methods (n = 3 per condition, repeated x 2). Expression levels

normalized to mean ΔCT level in samples cultured in androgen free media without TNFα

or DHT. **e,** ChIP-Seq analysis of AR at the *IL-8* (*CXCL8*) promoter in LNCaP cells cultured

in the presence of either vehicle (DMSO), DHT (100 nM), or TNFα (1000 U/ml) (n = 2 per

group; GSE83860). **f,** ChIP quantitative RT-PCR (qRT-PCR) analysis of AR, pSer2 Pol

II, pol II, and H3K4me3 at the *IL-8* (*CXCL8*) promoter (n = 3 per group). Transfected

LNCaP cells treated for 24 hours with or without DHT (100 nM). **g,** Schematic model of

the interplay between AR and NFkB in the regulation of IL-8 transcription. For **e,** loci with

significant differential binding (black bar) were identified as described in materials and

methods. Error bars represent standard error. Unpaired t-tests were performed, *p*-values

≤ 0.05 (*), 0.01 (**), 0.001 (***) and 0.0001 (****); *p*-values ≥ 0.05 (ns).

Figure 2 | IL-8 is Up-Regulated in Post-Castration and Castration-Resistant Prostate

Cancer Cells. **a,** Representative images of *Cxcl15* fluorescent detection (murine

homologue of IL-8) in Myc-Cap tumors. Tumors were harvested when volumes reached

~500mm³ (CS group), 7 days after androgen-deprivation (ADT), or at the time of

castration-resistance (CR) and hybridized with CF568-labeled probe sets (white) to

Cxcl15, CF640-labeled anti-PanCK antibody (red), and CF488-labeled anti-CD45

antibody (green). Nuclei counterstained with DAPI (blue). Repeated x 3. **b,** Gene and

protein expression of Cxcl15 in MCRedAL cells of indicated tumor samples by qRT-PCR

and ELISA, respectively (n = 3 per group, repeated x 2). **c,** qRT-PCR quantification of *IL-*

8 in human AR positive castration-sensitive cells (CS: LNCaP, LAPC4, and VCaP) and

their castration-resistant counterparts (CR: LNCaP-abl, LAPC4-CR, and VCaP-CR),

replicate numbers as in **b**. **d,** IL-8 protein expression in the isogenic cell pairs from **c**

quantified by ELISA, replicate numbers as in **c**. **e,** Representative images of *Cxcl15*

fluorescent detection in benign murine prostate tissue samples from castration-sensitive

(CS), androgen-deprivation treated (ADT), and ADT-treated mice that received

testosterone repletion (ADT + T). Tissue sections hybridized with CF568-labeled probe

sets (white) to *Cxcl15*, and CF640-labeled anti-PanCK antibody (red). Nuclei were

counterstained with DAPI (blue). Repeated x 3. **f,** qRT-PCR analysis of *Cxcl15*

expression in prostate luminal epithelial cells from indicated treatment groups (n = 3 per

group). Prostate luminal epithelial cells were isolated based on their

$GFP^+CD49f^{int}CD24^+CD45^-F4/80^-CD11b^-$ expression by flow sorting into Trizol LS. **g,**

Expression of *IL-8* in human prostate epithelial cells micro-dissected from patients in a

clinical trial (NCT00161486) receiving placebo, androgen-deprivation treatment (ADT), or

ADT plus testosterone repletion (ADT + T). Z-score values of microarray transcripts from

benign prostate biopsies were normalized to placebo samples (n = 4 per group;

GSE8466).   **h,** Expression of *IL-8* in human prostate cancer epithelial cells micro-

dissected from untreated or ADT-treated (NCT01696877; n = 8 per group) patients as

determined by qRT-PCR.   RISH images are at 60X magnification; scale bar = 100 μm.

Gene expression levels were normalized to the mean ΔCT level in samples from CS,

untreated or placebo groups.   For **b-g**, unpaired t-tests were performed; for **h** a Mann-

Whitney U test was used due to the non-normal data distribution observed. p-values ≤

0.05 (*) and 0.01 (**); p-values ≥ 0.05 (ns) shown. The range in box and whiskers plots

shows min and max values such that all data are included.

Figure 3 | Castration-mediated IL-8 Up-Regulation Promotes PMN-MDSC

Infiltration. a, Gating strategy used to profile the immune compartment of the TME by

flow cytometry. Tumor associated macrophages (TAMs) gated based on CD45$^+$Ly6G$^-$F4/80$^+$CD11b$^+$, Inflammatory (Inf.) TAMs as CD45$^+$CD11b$^+$F4/80$^+$Ly6C$^+$MHCII$^-$, immature (Imm.) TAMs as CD45$^+$CD11b$^+$F4/80$^+$Ly6C$^+$MHCII$^+$, MHCII^{hi} TAMs as CD45$^+$CD11b$^+$F4/80$^+$Ly6C$^-$MHCII$^+$, MHCII^{low} TAMs as CD45$^+$CD11b$^+$F4/80$^+$Ly6C$^-$MHCII$^-$, tumor Infiltrating Lymphocytes (TILs) CD45$^+$CD4$^+$ or CD45$^+$CD8$^+$, tumor infiltrating polymorphonuclear myeloid-derived suppressor cells (PMN-MDSCs) as CD45$^+$CD11b$^+$Ly6C$^+$Ly6G$^+$. **b,** TAM, TIL, and PMN-MDSC density normalized to mg of tumor weight (cells/mg; n ≥ 3 per group, repeated x 2). **c,** Representative H&E and immunohistochemistry (F4/80 and Ly6G) of indicated murine allografts (repeated x 3). **d,** Normalized expression of selected genes determined by NanoString nCounter gene analysis in sorted myeloid fractions defined as in **a** (n = 3 per group). **e,** qRT-PCR quantification of *Cxcr2* and *Il-23* in indicated populations of Myc-Cap tumors (n = 3 per group). **f** and **g,** Density of PMN-MDSCs normalized to mg of tumor weight (cells/mg) in Myc-Cap and PC3 tumors (n ≥ 4 per group, repeated x 2). Cells quantified by flow cytometry as in **a**, tumors implanted and harvested as in materials and methods. H&E and IHC images at 40X magnification; scale bar = 50 μm. Gene expression levels normalized to the mean ΔCT level in samples from the Immature TAMs (Imm.) group. Unpaired t-tests performed, *p*-values ≤ 0.05 (*), 0.01 (**), 0.001 (***) and 0.0001 (****); *p*-values ≥ 0.05 (ns).

Figure 4 | CXCR2 Blockade Improves Response to Immune Checkpoint Blockade

Following Androgen-Deprivation Therapy. **a,** Treatment scheme, scale = weeks.

Animals sacrificed for immune phenotyping 1 week post-ADT. **b,** Tumor growth and

survival curves of mice from isotype vs. anti-CTLA-4 vs. anti-CTLA-4 + anti-CXCR2

groups treated as described in **a** (black line vs. orange line vs. purple line, respectively; n

≥ 8 per group, repeated x 2). **c,** Tumor infiltrating lymphocyte (TILs) density in indicated

treatment groups (n ≥ 5 per group, repeated x 2). **d,** Treg percentages (as fraction of

CD4) in indicated tissues (n ≥ 5 per group, repeated x 2). **e,** Polyfunctional CD8 T cells,

left panel = density, center/right panels = percentage of total CD8, animals numbers as in

d. **f,** Representative histograms and dot plots of polyfunctional $CD8^+$ $IFN\gamma^+Gz\beta^+TNF\alpha^+$

from tumor draining lymph nodes (TDLN). Repeated x 2. For **a-f**, treatment was initiated

when tumor volumes reached $200mm^3$. Average tumor volume (±s.e.m.) for each

experimental group. Wilcoxon test used for survival analysis. Flow cytometry as in

materials and methods. Unpaired t-tests performed, *p*-values ≤ 0.05 (*), 0.01 (**), 0.001

(***) and 0.0001 (****); *p*-values ≥ 0.05 (ns).

References

1   Topalian, S. L., Drake, C. G. & Pardoll, D. M. Immune checkpoint blockade: a common denominator approach to cancer therapy. *Cancer Cell* **27**, 450-461, doi:10.1016/j.ccell.2015.03.001 (2015).

2   Drake, C. G. Prostate cancer as a model for tumour immunotherapy. *Nat Rev Immunol* **10**, 580-593, doi:10.1038/nri2817 (2010).

3   Kantoff, P. W. *et al.* Sipuleucel-T immunotherapy for castration-resistant prostate cancer. *N Engl J Med* **363**, 411-422, doi:10.1056/NEJMoa1001294 (2010).

4   Kwon, E. D. *et al.* Ipilimumab versus placebo after radiotherapy in patients with metastatic castration-resistant prostate cancer that had progressed after docetaxel chemotherapy (CA184-043): a multicentre, randomised, double-blind, phase 3 trial. *Lancet Oncol* **15**, 700-712, doi:10.1016/S1470-2045(14)70189-5 (2014).

5   Beer, T. M. *et al.* Randomized, Double-Blind, Phase III Trial of Ipilimumab Versus Placebo in Asymptomatic or Minimally Symptomatic Patients With Metastatic Chemotherapy-Naive Castration-Resistant Prostate Cancer. *J Clin Oncol* **35**, 40-47, doi:10.1200/JCO.2016.69.1584 [pii]
10.1200/JCO.2016.69.1584 (2017).

6   Topalian, S. L. *et al.* Safety, activity, and immune correlates of anti-PD-1 antibody in cancer. *N Engl J Med* **366**, 2443-2454, doi:10.1056/NEJMoa1200690 (2012).

7   Gao, J. *et al.* VISTA is an inhibitory immune checkpoint that is increased after ipilimumab therapy in patients with prostate cancer. *Nat Med* **23**, 551-555, doi:10.1038/nm.4308 (2017).

8   Mercader, M. *et al.* T cell infiltration of the prostate induced by androgen withdrawal

in patients with prostate cancer. *Proc Natl Acad Sci U S A* **98**, 14565-14570,

doi:10.1073/pnas.251140998 (2001).

9   Gannon, P. O. *et al.* Characterization of the intra-prostatic immune cell infiltration

in androgen-deprived prostate cancer patients. *J Immunol Methods* **348**, 9-17,

doi:10.1016/j.jim.2009.06.004 (2009).

10   Drake, C. G. *et al.* Androgen ablation mitigates tolerance to a prostate/prostate

cancer-restricted antigen. *Cancer Cell* **7**, 239-249, doi:10.1016/j.ccr.2005.01.027

(2005).

11   Shen, Y. C. *et al.* Combining intratumoral Treg depletion with androgen deprivation

therapy (ADT): preclinical activity in the Myc-CaP model. *Prostate Cancer Prostatic*

Dis **21**, 113-125, doi:10.1038/s41391-017-0013-x (2018).

12   Lu, X. *et al.* Effective combinatorial immunotherapy for castration-resistant

prostate cancer. *Nature* **543**, 728-732, doi:10.1038/nature21676 (2017).

13   Lopez-Bujanda, Z. & Drake, C. G. Myeloid-derived cells in prostate cancer

progression: phenotype and prospective therapies. *J Leukoc Biol* **102**, 393-406,

doi:10.1189/jlb.5VMR1116-491RR (2017).

14   Calcinotto, A. *et al.* IL-23 secreted by myeloid cells drives castration-resistant

prostate cancer. *Nature* **559**, 363-369, doi:10.1038/s41586-018-0266-0 (2018).

15   Watson, P. A. *et al.* Context-dependent hormone-refractory progression revealed

through characterization of a novel murine prostate cancer cell line. *Cancer Res*

65, 11565-11571, doi:10.1158/0008-5472.CAN-05-3441 (2005).

16   Hol, J., Wilhelmsen, L. & Haraldsen, G. The murine IL-8 homologues KC, MIP-2,

and LIX are found in endothelial cytoplasmic granules but not in Weibel-Palade

bodies. *J Leukoc Biol* **87**, 501-508, doi:10.1189/jlb.0809532 (2010).

17   Rossi, D. L. *et al.* Lungkine, a novel CXC chemokine, specifically expressed by

lung bronchoepithelial cells. *J Immunol* **162**, 5490-5497 (1999).

18   Schmitz, J. M., McCracken, V. J., Dimmitt, R. A. & Lorenz, R. G. Expression of

CXCL15 (Lungkine) in murine gastrointestinal, urogenital, and endocrine organs.

J Histochem Cytochem **55**, 515-524, doi:10.1369/jhc.6A7121.2007 (2007).

19   Chen, R. *et al.* Telomerase Deficiency Causes Alveolar Stem Cell Senescence-

associated Low-grade Inflammation in Lungs. *J Biol Chem* **290**, 30813-30829,

doi:10.1074/jbc.M115.681619 (2015).

20   Karlic, R., Chung, H. R., Lasserre, J., Vlahovicek, K. & Vingron, M. Histone

modification levels are predictive for gene expression. *Proc Natl Acad Sci U S A*

107, 2926-2931, doi:10.1073/pnas.0909344107 (2010).

21   Kumar, V. *et al.* Cancer-Associated Fibroblasts Neutralize the Anti-tumor Effect of

CSF1 Receptor Blockade by Inducing PMN-MDSC Infiltration of Tumors. *Cancer*

Cell **32**, 654-668 e655, doi:10.1016/j.ccell.2017.10.005 (2017).

22   Patnaik, A. *et al.* Cabozantinib Eradicates Advanced Murine Prostate Cancer by

Activating Antitumor Innate Immunity. *Cancer Discov* **7**, 750-765,

doi:10.1158/2159-8290.CD-16-0778 (2017).

23   Kawahara, T. *et al.* Neutrophil-to-lymphocyte ratio predicts prostatic carcinoma in

men undergoing needle biopsy. *Oncotarget* **6**, 32169-32176,

doi:10.18632/oncotarget.5081 (2015).

24   Yin, X. *et al.* Prognostic Role of Neutrophil-to-Lymphocyte Ratio in Prostate

Cancer: A Systematic Review and Meta-analysis. *Medicine (Baltimore)* **95**, e2544,

doi:10.1097/MD.0000000000002544 (2016).

25   Alfaro, C. *et al.* Tumor-Produced Interleukin-8 Attracts Human Myeloid-Derived

Suppressor Cells and Elicits Extrusion of Neutrophil Extracellular Traps (NETs).

Clin Cancer Res **22**, 3924-3936, doi:10.1158/1078-0432.CCR-15-2463 (2016).

26   Gentles, A. J. *et al.* The prognostic landscape of genes and infiltrating immune

cells across human cancers. *Nat Med* **21**, 938-945, doi:10.1038/nm.3909 (2015).

Materials and Methods

Patient Samples

Formalin fixed, paraffin embedded (FFPE) human prostate cancer samples were obtained from consented patients treated with ADT (degarelix; 240 mg SQ) in a neo-adjuvant trial (NCT01696877)[1] and matched control radical prostatectomies were obtained from patients treated at the Johns Hopkins Sidney Kimmel Comprehensive Cancer Center (Baltimore, MD) under IRB-approved clinical protocol J1265. All patients provided written, informed consent.

Cell Lines

Myc-Cap, derived from spontaneous prostate cancer in c-Myc transgenic mice [2,3], was a generous gift from Dr. C. Sawyers. To generate MCRedAL, Myc-Cap cells were transfected with pRetroQ-mCherry-C1 (Clontech) using lipofectamine 2000 (Invitrogen) and isolated by FACS sorting based on mCherry expression (Extended Data Fig. 1a). Myc-Cap and MCRedAL cells were cultured in DMEM as previously described[2]. LNCaP, VCaP, E006AA, CWR22Rv1, DU145, and PC3 cell lines were obtained and cultured as recommended by the ATCC. LAPC4 (a gift from Dr. S. Yegnasubramanian) were maintained in RPMI-1640 (Corning) supplemented with 10% fetal bovine serum (FBS; Gemini Bio-Products). Androgen independent LNCaP-abl cells were a gift from Dr. Z. Culig and cultured as descrived previously[4]. LAPC4-CR and VCaP-CR (a gift from S. Yegnasubramanian) were derived by passaging LAPC4 and VCaP cells through castrated animals and further subculturing in RPMI-1640 supplemented with 10% charcoal stripped serum (CSS; Gemini Bio-Products) supplemented with 1X B-27

Neuronal Supplement (Gibco). For experiments when cells were grown in androgen-free

conditions, 10% FBS was substituted for 10% CSS in complete media. For

migration/chemotaxis assays, prostate cancer cell lines were cultured in complete media

containing either 0.5% or 2.5% FBS for human and murine cells, respectively. All cell

lines were cultured in 1% penicillin/streptomycin media at 37°C, 5% CO_2.

Mouse Strains

Seven-week-old FVB/NJ, J:NU, C57BL/6-Tg(TcraTcrb)1100Mjb/J (OT-I), and B6.SJL-

PtprcaPepcb/BoyJ (CD45.1) male mice were purchased from The Jackson Laboratory.

A breeding pair of *Hoxb13-rtTA|TetO-H2BGFP* (HOXB13-GFP) mice[5] was received from

UMBC and experimental animals were bred in-house. Animals were kept in a specific

pathogen-free facility at either Johns Hopkins University School of Medicine or Columbia

University Medical Center. All animal experiments were performed in accordance with

protocols approved by the Institutional Animal Care and Use Committee (IACUC) at the

respective institutions.

Tumor Allografts and Xenografts

Eight-week-old male FVB/NJ and J:NU mice were subcutaneously inoculated with either

Myc-Cap or MCRedAL (1×10^6 cells/mouse), and LNCaP or PC3 (3×10^6 cells/mouse) in

the right flank, respectively. Tumor diameters were measured with electronic calipers

every 3 days as indicated and the tumor volume was calculated using the formula:

[longest diameter × (shortest diameter)^2]/2. Myc-Cap tumor bearing mice received

androgen-deprivation therapy (ADT) 4 weeks after tumor implantation when tumor

volume reached ~500mm^3, as indicated in figure legends. ADT was administered via

subcutaneous (sc) injection of degarelix acetate (a GnRH receptor antagonist; Ferring

Pharmaceuticals Inc.) at a dosage of 0.625 mg/100 μl H$_2$O/25 g body weight every 30

days, unless otherwise indicated. Onset of castration-resistance was defined as the time

to tumor size increased by 30% (~650 mm^3) after ADT. Chemical castration by ADT was

compared to bilateral orchiectomy as described in Extended Data Fig. 1a.

Luminal Epithelial Regression/Regeneration

Eight-week-old male HOXB13-GFP mice carrying the *Hoxb13-rtTA* transgene and a

Tetracycline operator–Histone 2B-Green Fluorescent Protein (TetO-H2BGFP), which

results on GFP expression being restricted to luminal epithelial Hoxb13$^+$ cells (described

previously[5]), were castrated via bilateral orchiectomy. A cycle of prostate

regression/regeneration was induced as described previously[6]. Briefly, mice were

allowed to regress for six weeks to reach the fully involuted state. Mice were randomized

to ADT or ADT + testosterone (T) treatment groups. Testosterone was administered for

four weeks for prostate regeneration by subcutaneous pellets; this regimen yields

physiological levels of serum testosterone. All mice received 2mg/ml of Doxycycline

(Sigma) in the drinking water to induce GFP expression[5] under the control of the luminal

epithelial promoter, HoxB13, one week prior euthanizing them for their analysis.

Antibody Blockade

Anti-CXCR2 (murine IgG1-D265A, clone: 11C8; a non-FcγR-binding mutant with

deficient FcγR-mediated depletion), anti-CSF1R (rat IgG2a, clone: AFS98; with

competent FcγR-mediated depletion), and anti-CTLA-4 (murine IgG2a, clone: 12C11; with competent FcγR-mediated depletion)[7] were used. Antibody treatment was administered via intraperitoneal (ip) injection at a dose of 50 mg/kg body weight for 3 doses every 4 days for CXCR2, 50 mg/kg body weight every 3 days for the duration of the experiment for CSF1R, and/or10 mg/kg body weight for 3 doses every 3 days for CTLA-4. Mouse IgG1 (clone: 4F7), rat IgG2a (clone: 2A3), and mouse IgG2a (clone: 4C6) were used as isotype controls. Anti-CXCR2 and anti-CSF1R treatments started 7 days before ADT; while anti-CTLA-4 treatment was started either 3 or 12 days before ADT ($400mm^3$ vs. $200mm^3$, respectively).

Flow cytometry

Single-cell suspensions from prostate tumor and tissues were prepared using the mouse tumor dissociation kit according to the manufacturer's recommendations (Miltenyi). Single-cell suspensions of tumor-draining lymph nodes (TDLNs) and spleens were homogenized mechanically with the back of a syringe. Cells were Fc-blocked with purified rat anti-mouse CD16/CD32 (Clone: 2.4 G2, Becton Dickinson BD) for 15 minutes at RT. Dead cells were discriminated using the LIVE/DEAD (L/D) fixable viability dye eFluor 506 or near-IR dead cell stain kit (Thermo Fisher) and samples were stained for extracellular and intracellular markers. The following antibodies were used: CD45 (30F-11), CD45.2 (104), CD24 (M1/69), CD49f (GOH3), Ly6C (HK1.4), Ly6G (1A8), Gr1 (RB6-8C5), CD11b (M1/70), F4/80 (BM8), MHCII (2G9), PD-L1 (10F.9G2), CD4 (RM4-5), CD8 (53-6.7), CD44 (IM7), CD62L (MEL-14), CD25 (PC61), Ki67 (16A8), IFN-γ (XMG1.2), TNF-α (MP6-XT22), IL-2 (JES6-5H4), GZβ (GB11), CXCR2 (242216), and IL-23

(FC23CPG). For intracellular staining, cells were fixed and permeabilized using BD

Perm/Wash (BD Biosciences) at room temperature for 45 minutes. For intracellular

cytokine staining, cells were stimulated with PMA (50 ng/ml) and ionomycin (500 ng/ml)

for 4 hours in the presence of protein transport inhibitor cocktail (eBiosciences). Gates

of cytokines were determined by fluorescence minus one (FMO) controls. Staining was

visualized by fluorescence activated cell sorting (FACS) analysis using a BD

FACSCelesta™ (BD Biosciences) and analyzed using FlowJo® (Flowjo LLC). Prostate

luminal epithelial cells are defined as $CD45^-CD11b^-F4/80^-CD24^+CD49f^{int}GFP^+$, and

prostate epithelial tumor cells are defined as $CD45^-CD11b^-F4/80^-mCherry^+$. Tumor

associated macrophages (TAMs) are referred to as $CD45^+CD11b^+F4/80^+$, inflammatory

TAMs   as $CD45^+CD11b^+F4/80^+Ly6C^+MHCII^-$,   immature   TAMs   as

$CD45^+CD11b^+F4/80^+Ly6C^+MHCII^+$, MHCII^{hi} TAMs as $CD45^+CD11b^+F4/80^+Ly6C^-MHCII^+$,

MHCII^{low} TAMs as $CD45^+CD11b^+F4/80^+Ly6C^-MHCII^-$. PMN-MDSCs are defined as

$CD45^+CD11b^+Ly6C^+Ly6G^+$.   CD4 T cells as $CD45^+CD4^+$, regulatory T cells as

$CD45^+CD4^+CD25^+$, CD8 T cells as $CD45^+CD8^+$, polyfunctional CD8 T Cells as

$CD45^+CD8^+IFN\gamma^+TNF\alpha^+Gz\beta^+$, and memory CD8 T cells as $CD45^+ CD8^+CD44^+CD62L^-$.

123Count eBeads counting beads (Thermo Fisher) were used to normalize the numbers

of PMN-MDSCs in migration/chemotaxis experiments.

Protein Quantification

Tumors collected at different treatment time points were minced, lysed in CelLytic MT

(Sigma) containing halt protease and phosphatase inhibitor (Thermo Fisher) in a 1:100

ratio, and incubated on ice for 30 minutes with intermittent vortexing. Tumor lysates were

assayed for raw protein concentration with Coomassie assay (Bio-Rad). IL-8 and Cxcl15

were analyzed by ELISA kits following the manufacturer's instructions (BD Bioscience

and R&D Systems, respectively).

Immunohistochemical staining (IHC)

Tumor and tissue samples were fixed with either 10% formalin (Fisher Scientific,

Pittsburgh, PA) or zinc fixative (BD) for 24 hours before paraffin embedding and

sectioning. Sections were stained with hematoxylin and eosin (H&E), and antibodies

against mouse Ly6G (1A8; BD Pharmingen) and F4/80 (BM8; eBioscience). Staining

was performed by the Molecular Pathology core of the Herbert Irving Comprehensive

Cancer Center at Columbia University. All images were acquired on a Leica SCN 400

system with high throughput 384 slide autoloader (SL801) and a 40X objective; files were

processed with Aperio ImageScope v12.3.1.6002.

RNA In Situ Hybridization (RISH) and Immunohistochemistry

Manual fluorescent RISH was performed on formalin-fixed and zinc-fixed paraffin

embedded sections using company protocols. Briefly, 5μm sections were cut, baked at

60 ℃ for 1 hour, dewaxed, and air-dried before pre-treatments. RISH *Cxcl15* probe, 3-

plex positive control probes (*Polr2a*, *Ppib*, *Ubc*) and 3-plex negative control probes (*DapB*

of Bacillus subtilis strain) from Advanced Cell Diagnostics (ACD) were used in this study.

Detection of specific probe binding sites was performed with RISH Multiplex Fluorescent

Reagent Kit v2 Reagent kit from ACD following the manufacturer's instructions. Tyramide

CF568 (Biotium) was used to visualize RISH signal.

For a more precise identification of cells expressing *Cxcl15*, RISH was coupled to

immunohistochemistry of PanCK (Poly; Dako) and CD45 (30-F11; BD Biosciences).

Immediately after RISH detection, samples were permeabilized with 0.2% TBS-Tween 20

for 10 minutes at RT, and then blocked with 2.5% of normal goat serum (Vector) for 30

minutes at RT. Primary antibody for PanCK was diluted 1/400 in renaissance background

reducing diluent (Biocare Medical) and incubated overnight at 4 °C. After washing off the

primary antibody, the slides were incubated 15 minutes at RT horseradish peroxidase

(HRP) secondary antibody (Vector). Tyramide CF640R (Biotium) was used to visualize

PanCK staining. In some cases, CD45 staining was also performed. For this, HRP signal

was abolished by a 30 minute incubation at RT with PeroxAbolish (Biocare Medical) and

then blocked with 2.5% of normal goat serum (Vector) for 30 minutes at RT. Primary

antibody for CD45 was diluted 1/50 in renaissance background reducing diluent (Biocare

Medical) and incubated 90 minutes at RT. After washing off the primary antibody, the

slides were incubated 15 minutes at RT HRP-secondary antibody (Vector). Tyramide

CF488A (Biotium) was used to visualize CD45 staining. All images were acquired on a

Nikon A1RMP confocal microscope using a 60X objective.   Comparisons of ISH-IHC

results were performed using ImageJ.

Whole Genome Expression Profiling and Analysis

MCRedAL tumor were harvested when their tumor volume reached ~500mm^3 (CS group),

and 7 days after chemical castration (ADT). MCRedAL cells were isolated based on their

mCherry$^+$ CD45$^-$ F4/80$^-$ CD11b$^-$ expression by flow sorting on a DakoCytomation MoFlo.

RNA was extracted using Trizol LS (Invitrogen) and treated with DNAse-I using RNA

clean & Concentrator (Zymo Research). The analysis was performed using Affymetrix

Mouse Clariom D (MTA 1.0) array according to the manufacturer's instructions. Resulting

CEL files were analyzed in Affymetrix Expression Console (v. 1.4) using the SST-RMA

method, and all samples passed the quality control. Log2 probe intensities were

extracted from CEL (signal intensity) files and normalized using RMA quantile

normalization, then further analyzed using Partek Genomics Suite v6.6. Illustrations

(volcano plots, heatmaps, and histograms) were generated using TIBCO Spotfire

DecisionSite with Functional Genomics. Gene set enrichment analysis (GSEA) of

differently expressed genes was performed using the hallmark gene sets Molecular

Signature Database (MSigDB).

Nanostring

RNA extraction was performed using the Trizol LS reagent (Thermo Fisher) as per

manufacturer's instructions. For NanoString analysis, the nCounter mouse PanCancer

Immune Profiling panel was employed using the nCounter Analysis System (NanoString,

Seattle, WA). Analysis was conducted using nSolver software (NanoString). Heatmap

analyses were performed using The R Project for Statistical Computing (https://www.r-

project.org/).

Pairwise Alignment

The homology of the murine chemokines Cxcl1, Cxcl2, Cxcl5, Cxcl15, Cxcl12, and Cxcl17

to   human   IL-8   was   evaluated   using   BLASTP   2.9.0+

(https://blast.ncbi.nlm.nih.gov/Blast.cgi?PAGE=Proteins)[8].   Proteins were consider

homologous if they shared > 30% amino acid identity[9]. Expected values of <0.05 were

consider statistically significant. The expected value includes an inherent Bonferroni

correction.

Chromatin immunoprecipitation assay (ChIP)-Seq

ChIP-Seq   data   was   obtained   from

https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE83860 which contains ChIP-

Seq data acquired with androgen receptor (AR) and nuclear factor NF-kappa-B p65

subunit (p65) specific antibodies on cell lysates from LNCaP cells cultured under the

following treatments: DMSO, DHT, and TNFα. For each treatment the dataset contains

two ChIP-Seq replicates pulled down using the AR and p65 antibodies[10]. ChIP-Seq data

were aligned to the hg38 reference version using the subread package, and then the BAM

files were sorted and indexed using SAMtools. Loci with significant differential binding

(FDR = 0.05) of pulled-down proteins to DNA were identified using the csaw package for

ChIP-Seq analysis, closely following Lun and Smyth's script[7]. ChIP-Seq visualization

was performed using the Integrative Genomics Viewer (IGV) from the Broad Institute

(http://software.broadinstitute.org/software/igv/).

ChIP-qRT-PCR

Chromatin immunoprecipitation was performed as described[11]. In brief, LNCaP cells

were washed with serum-free media and then grown in media containing 10% charcoal

stripped FBS for 48 hours. Cells were treated with 100nM DHT or vehicle for 8 hours.

DNA was cross-linked with 1% formaldehyde in PBS for 10 minutes and crosslinking was

quenched by addition of 0.125 M glycine. Fixed cells were then lysed in lysis buffer (1%

SDS, 5mM EDTA, 50mM Tris HCl, pH8.1) and sonicated to a fragment size of 200-600

bp using a Covaris water bath sonicator (Woburn, MA). Sheared chromatin was then

incubated with primary antibodies (AR [06-680, Millipore], H3K4me3 [ab8580, Abcam],

phospho-Ser5 RNA polymerase 2 [ab5131, Abcam], RNA polymerase 2 [4H8, Cell

Signaling Technologies] or control IgG [Cell Signaling Technologies]) overnight at 4ºC.

Complexes were immobilized on Dynabeads (Thermo Fisher) by incubating for 4 hours

at 4ºC. Beads were sequentially washed with TSEI (0.1% SDS, 1% Triton X-100, 2mM

EDTA, 20mM Tris HCl, pH 8.1, 150mM NaCl), TSEII (0.1% SDS, 1% Triton X-100, 2mM

EDTA, 20mM Tris HCl, pH 8.1, 500mM NaCl) and TSEIII (0.25 M LiCl, 1% NP-40, 1%

deoxycholate, 1mM EDTA, 10mM Tris HCl, pH 8.1). DNA was eluted with IP Elution buffer

(1% SDS, 0.1M NaHCO$_3$, proteinase K) and incubated at 56ºC for 15 minutes. Enriched

DNA libraries were analyzed using primers specific to *IL-8* locus: Forward: 5'

AGCTGCAGAAATCAGGAAGG 3' and Reverse: 5' TATAAAAGCCACCGGAGCA 3'

using quantitative (q) RT-PCR. Data is shown as relative enrichment normalized to input

DNA.

Quantitative (q) RT-PCR

Total RNA was extracted using Trizol (Ambion). cDNA was prepared from total RNA

preps using the RNA to cDNA EcoDry Premix (Clontech). Real-time assays were

conducted using TaqMan real-time probes (Applied Biosystems). ΔΔ CT method was

used for relative gene expression. Expression of the target gene was normalized to the

reference gene (18S) and the mean expression level of the control group. LCM samples

were normalized to 18S, TBP, and GAPDH reference genes.

Laser Capture Microscopy (LCM)

Formalin fixed-paraffin embedded radical prostatectomy specimens, from patients

enrolled in a neoadjuvant clinical trial (NCT01696877)[1] who received 240 mg (SQ) of

degarelix and matched control cases (patients that did not receive any hormone therapy),

were sectioned at a thickness of 8 µm and transferred onto PEN membrane glass slides

(Leica). Sections were deparaffinized, hydrated and stained with hematoxylin prior to

microdissection. Individual cancer cells and cancer cell clusters were microdissected by

a trained pathologist using a LMD 7000 laser capture microscope (Leica). RNA was

recovered from the microdisseceted material using the RNeasy FFPE kit (Qiagen).

Quantitative RT-PCR was performed as described above. For the analysis, a Mann-

Whitney U test was performed.

IL-8 and Cxcl15 CRISPR/Cas9 Knock Outs

The 20 bp long gRNA, designed using Deskgen online software, for targeting *IL-8* and

Cxcl15 in exon 3 (5'- *TTCAGTGTAAAGCTTTCTGA* -3' and 5'-

ACAGAGCAGTCCCAAAAAT -3', respectively) were incorporated into two

complementary 100-mer oligonucleotides and cloned into a gRNA containing plasmid

containing the (NeoR/KanR) cassette (Addgene # 41824). The human codon optimized

pCAGGS-Cas9-mCherry was used for gene-editing experiments (a gift from Stem Cell

Core Facility at Columbia University).   gRNA and Cas9 containing plasmids were

introduced to prostate epithelial cells using the basic nucleofector kit (Amaxa, Lonza)

following the manufacturer's instructions for primary mammalian epithelial cells (program

W001). Successfully transfected cells were selected by culturing in the presence of

400µg/ml of neomycin sulfate analog (G418; Sigma), and isolated based on their mCherry

expression 24 hours after transfection. Knock out clones were screened for IL-8 and

Cxcl15 expression by ELISA and gene-editing confirmed by PCR amplification and

Sanger sequencing (GENEWIZ) using primers ~200bp away from the cut site (IL-8

Forward: 5'- TTTGGACTTAGACTTTATGCCTGAC -3; IL-8 Reverse: 5'-

TCCTGGGCAAACTATGTATGG -3; Cxcl15 Forward: 5'-

GCTAGGCACACTGATATGTGTTAAA -3; Cxcl15 Reverse: 5'-

ACATTTGGGGATGCTACTGG -3).

Migration/Chemotaxis Assay

Cells and supernatants used in this assay were resuspended in culture media containing

0.5% or 2.5% FBS. Transwell plates of 3-mm pore size were coated with Fibronectin

(Corning Costar) and loaded with 500 ml of medium or with different cell supernatants in

triplicates (lower chamber). Cells were resuspended at $2x10^7$ cells/ml, and 200 ml of this

suspension was placed in each of the inserts (upper chamber). After 2.5 hours of

incubation at 37°C and 5% CO_2, inserts were removed and 10,000 beads (Thermo Fisher)

were added to each well. In some cases, either isotype or anti-CXCR2 (200 µg/ml) were

added at the beginning of the experiment. The cells in the lower chamber were collected

along with the starting cell population, stained with L/D, CD11b, Ly6C, and Ly6G and

evaluated by flow cytometry in a BD FACSCelesta™ (BD Biosciences). The ratio of beads

to cells was determined, allowing calculation of the number of cells that had migrated to

the bottom well. *In vivo*, LD-PMN-MDSCs were collected as described below from

splenocytes of CR-Myc-Cap tumor bearing mice and labeled with DiD (DilC18(5) or 1,1'-

Dioctadecyl-3,3,3',3'-

Tetramethylindodicarbocyanine, 4-Chlorobenzenesulfonate Salt; Invitrogen), a lipophilic

membrane dye, as described previously[12]. DiD$^+$ LD-PMN-MDSCs were adoptively

transferred into FVB/NJ recipient 8-week male mice and their ability to migrate in

response to 200ng of recombinant Cxcl15 was evaluated 4 hours after injection. Beads

were also used to calculate absolute numbers of Ly6G$^+$ PMNs and DiD$^+$ LD-PMN-MDSCs

in vivo.

PMN-MDSC Enrichment

Animals were sacrificed and spleens were collected. After dissociating cell clumps, the

cell suspension was centrifuged (740 g, 10 minutes, RT) and resuspended in 1 ml HBSS–

EDTA containing 0.5% BSA. Cells were then resuspended in 50% Percoll solution and

treated on a three-layer Percoll gradient (55%, 72%, and 81%) at (1500 g, 30 minutes,

10°C without break). LD-PMN-MDSCs were collected from the 50-55% and 55-72%

interfaces. Red blood cells (RBCs) were eliminated with RBC lysis solution (Miltenyi).

In vitro Suppression Assays

PMN-MDSCs were isolated from the spleen of CR-Myc-Cap-tumor bearing mice using

the neutrophil isolation kit (Miltenyi) according to the manufacturer's instructions; greater

than 95% enrichment was confirmed by flow cytometry. Unless otherwise indicated, a

density gradient separation was performed prior to column purification. OT-I (CD45.2)

transgenic splenocytes were mixed at a 1:10 ratio with sex-matched CD45.1 splenocytes.

Splenocytes containing CD8 T responder cells were stained with CellTrace Violet (5µM

CTV; Thermo Fisher) and plated on a 96-well round-bottom plate at a density of 2×10^5

cells per well. PMN-MDSCs cells were added at 2-fold dilutions starting from 2×10^5 cells,

in the presence of their cognate peptides (5pM OVA) and incubated for 60 hours.

Proliferation of CD8 T responder cells (gated as L/D⁻CD8⁺CTV⁺) was quantified by flow

cytometry based on the dilution of Cell Trace Violet (CTV). Percent suppression (%

Suppression) was calculated by the following formula: % Suppression = [1-(% divided

cells of the condition/ the average of % divided cells of T responder only conditions)] x

100.

Z-score Analysis

IL-8 expression was evaluated in a publicly available data set (GSE8466)[13] using z-score

values of quantile-normalized microarray transcripts from benign prostate biopsies. Z-

score values were obtained by scaling the data for each gene in each patient to:

(expression - mean expression across all genes) / (standard deviation of expression

across all genes).

Statistical Analysis

Statistical analysis was performed using Prism 7 (GraphPad). Unpaired two-tailed t-tests,

Mann-Whitney U test, Tukey's multiple comparisons tests, or Wilcoxon rank sum tests

were conducted and considered statistically significant at *p*-values ≤0.05 (*), 0.01 (**),

0.001 (***) and 0.0001 (****).

Methods References

1 Antonarakis, E. S. *et al.* Neoadjuvant randomized trial of degarelix (Deg) ±

cyclophosphamide/GVAX (Cy/GVAX) in men with high-risk prostate cancer (PCa)

undergoing radical prostatectomy (RP). *Journal of Clinical Oncology* **35**, 5077-

5077, doi:10.1200/JCO.2017.35.15_suppl.5077 (2017).

2 Watson, P. A. *et al.* Context-dependent hormone-refractory progression revealed

through characterization of a novel murine prostate cancer cell line. *Cancer Res*

65, 11565-11571, doi:10.1158/0008-5472.CAN-05-3441 (2005).

3 Ellwood-Yen, K. *et al.* Myc-driven murine prostate cancer shares molecular

features with human prostate tumors. *Cancer cell* **4**, 223-238 (2003).

4 Culig, Z. *et al.* Switch from antagonist to agonist of the androgen receptor

bicalutamide is associated with prostate tumour progression in a new model

system. *Br J Cancer* **81**, 242-251, doi:10.1038/sj.bjc.6690684 (1999).

5 Rao, V. *et al.* A Hoxb13-driven reverse tetracycline transactivator system for

conditional gene expression in the prostate. *Prostate* **72**, 1045-1051,

doi:10.1002/pros.22490 (2012).

6 Wang, X. *et al.* A luminal epithelial stem cell that is a cell of origin for prostate

cancer. *Nature* **461**, 495-500, doi:10.1038/nature08361 (2009).

7 Shen, Y. C. *et al.* Combining intratumoral Treg depletion with androgen deprivation

therapy (ADT): preclinical activity in the Myc-CaP model. *Prostate Cancer Prostatic*

Dis **21**, 113-125, doi:10.1038/s41391-017-0013-x (2018).

8   Altschul, S. F. *et al.* Protein database searches using compositionally adjusted

substitution matrices. *Febs J* **272**, 5101-5109, doi:10.1111/j.1742-

4658.2005.04945.x (2005).

9   Pevsner, J. *Bioinformatics and functional genomics*. (John Wiley & Sons, 2015).

10   Malinen, M., Niskanen, E. A., Kaikkonen, M. U. & Palvimo, J. J. Crosstalk between

androgen and pro-inflammatory signaling remodels androgen receptor and NF-

kappaB cistrome to reprogram the prostate cancer cell transcriptome. *Nucleic*

Acids Res **45**, 619-630, doi:10.1093/nar/gkw855 (2017).

11   Haffner, M. C. *et al.* Androgen-induced TOP2B-mediated double-strand breaks

and prostate cancer gene rearrangements. *Nat Genet* **42**, 668-675,

doi:10.1038/ng.613 (2010).

12   Carlson, A. L. *et al.* Tracking Single Cells in Live Animals Using a Photoconvertible

Near-Infrared Cell Membrane Label. *Plos One* **8**, doi:ARTN e69257

10.1371/journal.pone.0069257 (2013).

13   Mostaghel, E. A. *et al.* Intraprostatic androgens and androgen-regulated gene

expression persist after testosterone suppression: therapeutic implications for

castration-resistant prostate cancer. *Cancer Res* **67**, 5033-5041,

doi:10.1158/0008-5472.CAN-06-3332 (2007).