1 **Biomass, abundances, and abundance and**

2 **geographical range size relationship of birds along a**

3 **rainforest elevational gradient in Papua New Guinea**

4

5 Katerina Sam[1,2], Bonny Koane[3]

6

7 [1] Biology Centre of the Czech Academy of Sciences, Entomology Institute, Ceske

8 Budejovice, Czech Republic

9 [2] University of South Bohemia, Faculty of Science, Ceske Budejovice, Czech Republic

10 [3] The New Guinea Binatang Research Centre, Madang, Papua New Guinea

11
12
13 Corresponding Author:

14 Katerina Sam[1,2]

15 Branisovska 31, 370 05, Ceske Budejovice, Czech Republic

16 Email address: Katerina.sam.cz@gmail.com

17
18 ## Abstract

19 The usually positive inter-specific relationship between geographical range size and the
20 abundance of local bird populations comes with exceptions. On continents, the majority of
21 these exceptions have been described from tropical montane areas in Africa, where
22 geographically-restricted bird species are unusually abundant. We asked how the local
23 abundances of passerine and non-passerine bird species along an elevational gradient on Mt.
24 Wilhelm, Papua New Guinea relate to their geographical range size. We collected data on
25 bird assemblages at eight elevations (200 – 3,700 m, at 500 m elevational increments). We
26 used a standardized point-counts at 16 points at each elevational study site. We partitioned
27 the birds into feeding guilds, and we obtained data on geographical range sizes from Bird-
28 Life International data zone. We observed a positive relationship between abundance and
29 geographical range size in the lowlands. This trend changed to a negative one towards higher
30 elevations. The total abundances of the assemblage showed a hump-shaped pattern along the
31 elevational gradient, with passerine birds, namely passerine insectivores, driving the observed

32 pattern. In contrast to abundances, the mean biomass of the bird assemblages decreased with
33 increasing elevation. Our results show that montane bird species maintain dense populations
34 which compensate for the decreased available area near the top of the mountain.
35

Introduction

Many previous studies have found a positive inter-specific relationship between geographical
range size and the abundance of local populations (Brown 1984; Gaston & Blackburn 2000;
Gaston et al. 2000). The authors hypothesized that (1) species utilizing a wider range or more
abundant resources become more abundant and widely distributed (Brown 1984), (2) that
high population growth leads to higher abundances and to more occupied sites or that (3)
intensive dispersal produces a positive inter-specific abundance-range size relationship
(Borregaard & Rahbek 2010; Gaston et al. 2000). While there is extensive literature devoted
to the patterns of species diversity along elevational gradients (McCain 2009; Rahbek 1995),
these studies rarely combine species richness with the study of bird abundance and biomass,
arguably more important parameters when it comes to the impact of birds on other trophic
levels (but see e.g., (Romdal 2001; Terborgh 1977). Even fewer studies have combined these
attributes of bird communities with an estimate of available resources (Ding et al. 2005;
Ghosh-Harihar 2013; Price et al. 2014) and/or available area along mountain ranges (e.g.,
Ferenc et al. 2016; Price et al. 2014).

Furthermore, many studies do not address the potential differences between
passerines and non-passerines or, they completely exclude non-passerine species. Klopfer &
MacArthur (1960) suggested that phylogenetically younger passerines should be relatively
more abundant than non-passerines in unstable environments. They assumed that younger
passerines have a less limited central nervous capacity than non-passerines, making them
more capable of fitting changing environmental stimuli. In this study, we aimed to test the
analogous hypothesis that non-passerines will be more abundant in favorable tropical
lowlands with stable climatic conditions than in the more variable environments at higher
elevations. In the Himalayas, the ratio of passerines to non-passerines increased very slowly
between 160 and 2,600 m a.s.l., and abruptly between ca. 3,000 – 4,000 m a.s.l. (Price et al.
2014) (but note that not all non-passerines were surveyed). Similarly, passerine abundance
increased relative to non-passerines with increasing elevation in the Andes (Terborgh 1977).
Finally, bird studies focusing on the patterns of abundance or biomass in different feeding
guilds along elevational gradients are rare, however, they are essential for improving our
understanding of ecosystem dynamics and function.

Macroecological studies have often revealed positive interspecific correlations
between geographical range sizes and the abundance of local populations (Brown 1984;
Gaston & Blackburn 2000; Gaston et al. 2000). The negative correlation between abundance
and range-size was showed on temperate datasets (but see Blackburn et al. 2006) and in
montane Africa, where the geographically restricted species are generally more abundant than
species with large geographical ranges (Fjeldså et al. 2012, Ferenc et al. 2016, Reif et al.
2006). Several other recent studies of tropical montane taxa report that abundance is
uncorrelated with (or negatively correlated to) geographical range size (Nana et al. 2014;
Reeve et al. 2016) but see (Theuerkauf et al. 2017). The only existing study on this topic from

Papua New Guinea showed that abundance was not related to range size (Freeman 2018).
However, in contrast to other studies, this one was based solely on mist-netting data and
range size was calculated as elevational breath instead of area (Freeman 2018).
The drivers behind the high abundances of montane forest species are unknown.
However, several mutually non-exclusive hypotheses have been considered (Ferenc et al.
2016). These are: (1) Long-term climatic stability allows specialization of new ecological
forms, which then leads to high local abundances of species at mountain tops (Fjeldså et al.
2012). (2) Species-poor communities compensate with increased density at high altitudes
which then leads to high abundances of montane bird species (MacArthur 1972). (3) Locally
abundant tropical montane species have a higher chance of surviving despite their small
range sizes while insufficiently abundant species go extinct (Johnson 1998).
To investigate the relationship between abundance and area in different regions, we
focused on bird assemblages along the elevational gradient of Mt. Wilhelm in Papua New
Guinea. Our goals were to investigate (1) the trends in abundances of birds along the
elevational gradient, (2) the changes in relative abundances of different groups of birds
(passerines and non-passerines, various feeding guilds), and (3) the effects of geographical
range sizes on the abundance of individual species.
92

Materials & Methods

The study was performed along Mt Wilhelm (4,509 m a.s.l.) in the Central Range of Papua
New Guinea (Figure 1a, b). The complete rainforest gradient spanned from the lowland
floodplains of the Ramu river (200 m a.s.l., 5° 44'S 145° 20'E) to the treeline (3700 m a.s.l.,
5° 47'S 145° 03'E; Fig. 1). We completed the study along a 30 km long transect, where eight
sites were evenly spaced at 500 m elevational increments. Because of the steep terrain,
elevation could deviate by 50 m within each study site. Survey tracks and study sites at each
elevation were directed through representative and diverse microhabitats (e.g., ridges,
valleys, rivulets; ≥ 250 m from forest edge). In the lowlands, average annual precipitation is
3,288 mm, rising to 4,400 mm at 3,700 m a.s.l. There is a distinct condensation zone at
around 2,500 – 2,700 m a.s.l. (McAlpine et al. 1983). Mean annual temperature typically
decreases at a constant rate of 0.54°C per 100 elevational meters; from 27.4°C at the lowland
site (200 m a.s.l.) to 8.37°C at the tree line (3700 m a.s.l.). The habitats of the elevational
gradient is described as lowland alluvial forest (200 m a.s.l.), foothill forest (700 and 1,200 m
a.s.l.), lower montane forest (1,700 – 2,700 m a.s.l.), and upper montane forest (3,200 and
3,700 m a.s.l.; according to Paijmans (1976). Plant species composition of the forest
(Paijmans 1976), general climatic conditions (McAlpine et al. 1983) and habitats at
individual study sites (Sam & Koane 2014) are described elsewhere.
Data on bird communities were collected in 2010, 2011 and 2012 during the wet and
dry seasons, using a standardized point-counts at 16 points per elevation (Sam & Koane
2014; Sam et al. 2019). Both visual observations and identifications based on calls were used.
The surveys were conducted in the mornings between 0545 and 1100. Each of the 16 sample
points had a radius of 50 m (area 0.785 ha per point, which makes 12.56 ha per elevational
study site). Points were located 150 m apart to lower the risk of multiple encounters of the
same individuals. We visited each point 14 times (8 times during the dry season and 6 times

during the wet season). The order of the points was altered during each re-survey, to avoid
biases due to time of day. Birds were detected for 15 minutes during each visit at each point.
This resulted in 240 minutes of daily surveys. During the point-counts, we used a distance
sampling protocol. The birds were recorded in five 10-m-wide radial distance bands
(Buckland et al. 2001). Detection adjustments, however, proved to come with significant
problems in the tropics (Banks☐Leite et al. 2014). Therefore, we used the observed
abundance only estimates instead of the distance sampling-based estimates in the analyses
(see similar reasons and discussion by Ferenc et al. 2016). To evaluate the consistency in our
data, we (1) compared the abundances of birds observed during point-counts (reported here)
and from mist-netting conducted at the same sites during the same surveys (Sam et al. 2019),
(2) we ran all the analyses reported here also with mist-netting data, and we (3) we compared
the abundances of the birds recorded during point-counts done in wet and dry season (Figure
S1-S3). The data showed that abundances obtained by mist-netting and by point-counts and
by point-counts in wet and dry season are well correlated, and that the trends remain
unchanged, when only mist-netting data are used (Figure S1-S3).

We recorded the number of individuals of each species at any of the 15-min intervals
and summed them across all 16 points of each survey day at the certain elevation. Then we
averaged these daily abundances across the 14 days (or 6 days of wet season and 8 days of
dry season respectively at each elevation). Hereafter we call this measure "*mean elevational*
abundance" of a given species at a certain elevational site. After that, we averaged the *mean*
elevational abundance across the elevations where the bird species was present to calculate
"*mean abundance*" of a given species along the elevational gradient. Species at their
elevational range limits usually have low abundances which might be difficult to detect
correctly. Therefore, potential errors, which would lead to an erroneous *mean abundance*,
might occur if rare individuals at the sites close to their range limits are missed during census.
To ensure that our observations are valid, we also repeated the analyses with the "*maximal*
mean elevational abundance" of each species (Figure S4).

To summarize the abundances of bird assemblages at a given elevation (hereafter
"*total abundance*") we calculated the sum of the *mean elevational abundance* of all species
present at each site (i.e. at 16 points within a 4-hour long survey). Elevations between
minimal and maximal range where birds were missing were not considered, i.e., data were
not extrapolated, and the birds were given zero abundance at this elevation. The taxonomy
used followed the International Ornithological Congress World Bird List version 6.1.

The elevational *weighted mean point* was calculated as the elevation, where the
species had potentially the highest abundances. *Weighted mean point* was calculated as a sum
of elevations weighed by an abundance of the given species at this elevation which was
divided by the sum of the abundances (e.g. Elevation 1 * abundance at elevation 1 + elevation
2 * abundance at elevation 2)/(abundance at elevation 1 + abundance at elevation 2) (Figure
S4). Based on the weighted mean point, we divided the species into three groups as follows:
(a) "lowland" group - species with their elevational weighted mean point in the lower part of
the elevational gradient (up to 800 m a.s.l.), (b) "middle" group - species with a weighted
mean point between 800 and 1600 m a.s.l., and (c) "montane" group - species with their
weighted mean point in the upper third of the gradient (above 1,600 m a.s.l.). Note that a
single species (Great cuckoo-dove - *Reinwardtoena reinwardti*), occurring at all sites along

the complete gradient between 200 and 3,200 m, thus falls into the group of montane species.
To confirm the validity of our data, we also repeated our analyses with the maximal mean
elevational abundance point, i.e. the elevational site where we recorded the *maximal mean*
elevational abundance (Figure S4).
All recorded bird species were classified into five trophic guilds: insectivores,
frugivores, frugivores-insectivores, insectivores-nectarivores and nectarivores based on
dietary information from standard references (Hoyo et al. 1992-2011; Pratt & Beehler 2015)
and our data (Sam et al. 2019; Sam et al. 2017). Abundances of passerines and non-passerines
and individual feeding guilds were compared by non-parametric Kruskal-Wallis tests. We
report the mean ± SE and abundances per 12.56 ha recorded in a 15-minute-long census
unless we state otherwise. Geographical range sizes of all birds were obtained from Bird-Life
International data zone web pages accessed in July 2016. Bodyweight (mean for males) of the
birds were obtained from Hoyo et al. (1992-2011). Bird metabolism was calculated from
bodyweight according to available equations (McNab 2009).
We conducted the field work under the Institutional Animal Care and Use Committee
approval permit No. 118 000 561 19 and 999 020 778 29 awarded by the PNG National
Research Institute permit. Research was also permitted by Australian Bird and Bat Banding
permit No. 3173. The data were collected at the land of several rainforest dwelling
communities. The customary landowners (Peter Sai, Family Mundo, Alois Koane, Simon
Yamah, Samson Yamah, Joe Black) gave fully informed verbal and prior consent to the study
of bird communities on their land along Mt. Wilhelm gradient. The negotiations with
landowners were organized via The New Guinea Binatang Research Centre, an NGO in
Papua New Guinea.
185

Results
In total, we recorded 25,715 birds belonging to 249 (Table S1) species from the point-counts
along the elevational gradient of Mt. Wilhelm. This represents 87% of bird species recorded
along the gradient so far (Marki et al. 2016; Sam & Koane 2014; Sam et al. 2019). Total bird
species richness seemed to show a plateau at lower elevations (up to 1700 m a.s.l.) and
decreased with increasing elevation afterward (Figure 2a). In contrast, *total abundance* of
birds showed a humped shaped pattern, peaking between 1,700 and 2,700 m a.s.l. with ca.
420-450 individuals of all birds per 16 sampling points (i.e., 12.86 ha) (Figure 2c).
194

Passerines and non-passerines
Passerines were overall more species rich along the elevational gradient, represented by 161
species in comparison to non-passerines represented by 88 species (Figure 2b). We observed
a linearly decreasing pattern in species richness of non-passerine birds (N = 8, y = -5.9167x +
60.056, R^2 = 0.96) along the elevational gradient and a hump-shaped pattern (N = 8, y = -
$2.1012x^2$ + 18.982x + 27.315, R^2 = 0.92) in species richness of passerine birds (Figure 2b).
The species richness of passerines (r = 0.52, P = 0.19, N = 8) and non-passerines (r = 0.91, P
= 0.001, N = 8) correlated with their *total abundance* (Figure 2b, c).
The *mean elevational abundance* of passerine birds was overall significantly higher
(mean ± SD = 3.90 ± 4.8) than the *mean elevational abundance* of non-passerines (mean ±

SD = 2.46 ± 3.1; W = 21438; P < 0.001). The *mean elevational abundance* of assemblages
increased with increasing elevation, with approximately 2.5 times as many individuals per
non-passerine species and nearly twice as many individuals per passerine species at the
highest elevation than in the lowlands (Figure 3). The pattern was similar in both wet and dry
seasons (Figure S5). This pattern remains to be valid even when only the *maximal mean*
elevational abundance were considered. Birds having their *maximal mean elevational*
abundance at higher elevations had abundances higher than birds with maxima in lowlands
(Figure S6)
Passerine birds with an elevational weighted mean point in the montane forest (above
1600 m a.sl.) had a higher *mean abundance* than those with a middle and lowland distribution
(Figure 4a, Table S1). However, with their increasing elevational weighted mean point, the
geographical ranges of the species decreased (Figure 4b). We found no significant change in
the *mean elevational abundance* of non-passerine birds with an elevational weighted mean
point (Figure 4c) but similarly to passerines, non-passerines with a higher elevational
weighted mean point had smaller ranges (Figure 4c). The abundance range-size relationships
for all bird species of the complete forestal gradient of Mt. Wilhelm showed a significantly
negative relationship ($F_{1,248}$ = 8.22, P = 0.004, Figure S7). The trends remained negative,
albeit nonsignificant, for passerines ($F_{1,159}$ = 1.17, P = 0.28) and non-passerines ($F_{1,86}$ = 2.6, P
= 0.10) separately (Figure S7). However, the relationship of the three bird groups with
different elevational weighted mean points showed a variable pattern, as the trend changed
from a positive relationship in the lowland group of species, to no trend for middle species,
and a negative trend for montane species (Figure S8). The pattern remained similar, when we
split the data into abundances in the wet and dry season (Figure S9). Furthermore, the pattern
remains unchanged even when the *maximal mean elevational abundance* is considered in
analyses, as the *maximal mean elevational abundance point* and weighted mean point
correlated closely (Figure S10). Finally, more abundant passerine montane birds had not only
larger geographical ranges, but also longer elevational ranges (Figure S11).
232
Feeding guilds
Without respect to which feeding guild they belong, species occurring at low elevations
usually had a lower *mean elevational abundance* than species occurring at high elevations
(Figure 4a) i.e., their *mean elevational abundance* increased with increasing elevation.
Nectarivorous and insectivore-nectarivorous species had the highest *mean elevational*
abundance which increased towards higher elevations (Figure 5a). Within insectivore-
nectarivores, the pattern was driven purely by the presence of flocks of nectar-feedings
lorikeets at high elevations (i.e. the pattern disappeared when we removed lorikeets from the
dataset).
Total abundances of bird assemblages belonging to different feeding guilds however
showed different patterns (Figure 5b). While *total abundances* of insectivores followed a
mid-elevational peak (Figure 5b), *total abundances* of other feeding guilds showed no trend
(Figure 5b).
Within passerine birds, the *mean elevational abundance* of birds belonging to
different feeding guilds (except frugivores) increased with their elevational weighted mean

point (Figure 5c). In contrast, the *mean elevational abundance* of non-passerines birds
belonging to various feeding guilds showed various patterns (Figure 5d).

Mean biomass of bird communities (Figure 6) recorded at each elevational study site
decreased with increasing elevation, thus showing a different pattern from *mean elevational*
abundance and *total abundance*. At the two highest elevations (3,200 and 3,700 m) mean
biomass of passerines was relatively larger than biomass of non-passerines. The pattern of
decreasing biomass was observed both with passerines and non-passerines (Figure 6a), as
well as in all feeding guilds (Figure 6b). Because large species may have larger ranges
(Gaston 1996), we tested how strong the relationship was between body size and
geographical range. We found a weakly positive correlation between body size and range size
in non-passerines, and no correlation in passerine birds (Figure S12).

Discussion

In this study we focused on the patterns and relationships in species richness, abundance and
range size in assemblages of passerine and non-passerine birds along a tropical, elevational
gradient. After a detectable, initial plateau at low elevations, overall species richness declined
with increasing elevation on Mt. Wilhelm (Sam et al. 2019), a pattern that is typical for
mountains with a humid base (McCain 2009). Further, when considered separately the
species richness patterns of passerines (hump-shaped) and non-passerines (steeply
decreasing) differed. The findings are, in part, similar to those from Himalaya (Price et al.
2014), where species richness of non-passerines decreased with increasing elevation but
passerines followed a hump-shaped pattern and their richness peaked at ca. 2000 m.

Total abundance of bird assemblages at individual elevations also follows a different
hump-shaped pattern. This mismatch in patterns between observed species richness and
abundance is an interesting observation. Previous studies show that unimodal or linearly
decreasing patterns in density are usually paralleled by the patterns of total species richness
along the same gradients (e.g., Romdal 2001; Terborgh 1977). Our findings are similar to
patterns in abundances of birds observed along an elevational gradient in Cameroon (Ferenc
et al. 2016). In this study, declining species richness was associated with minimal changes in
the *total abundance* (i.e. the number of individuals per species) of birds because the average
number of individuals per species increased with increasing elevation.

Overall, the trends exhibited by *total abundance* and species richness do not
correspond to one another. However, when *total abundance* is partitioned into the hump-
shaped trend for passerines and decreasing trend for non-passerine birds, the trends become
complementary with their respective species richness. To our knowledge, there is not a single
study focusing separately on the abundance patterns of passerine and non-passerine birds
along an elevational gradient. Our data further show that the species richness and abundance
of passerines increase relative to non-passerines with increasing elevation. This might be in
concordance with previous suggestions that phylogenetically younger passerines should be
relatively more abundant in less favorable and unstable environments. Klopfer & MacArthur
(1960) showed that the proportions of non-passerines to passerines changes from north to
south. A study comparable to ours by Price et al. (2014) indicated that the ratio between the

abundances of passerines and non-passerines increased very slowly between 160 and 2,600 m
a.s.l., and then increased abruptly between ca. 3,000 – 4,000 m a.s.l..

The widespread pattern that abundance is positively correlated with geographic range
size (Gaston & Blackburn 2000) does not seem to apply to New Guinean birds distributed
along elevational gradients. Contrary to this widely accepted pattern, we described a negative
correlation between the local abundance of birds and the complete range size of the given
species. The deviation from a positive abundance-area relationship is caused by the
combination of the decreasing range size and increasing abundance of birds towards high
elevations. This observation is also consistent with the idea of taxon cycles whereby endemic
species are confined to mountain tops. This observation also fits the predictions of the density
compensation hypothesis in which individual species may increase their abundances to fill
the available ecological space in species-poor assemblages (MacArthur et al. 1972). The
hypothesis thus assumes that small-range species that have insufficiently sparse local
populations become extinct.

We showed that New Guinean bird species with small ranges are associated with high
local abundances, as has been suggested for marsupials in Australia (Johnson 1998), birds of
the Australian wet tropics (Williams et al. 2009) and Afromontane birds (Ferenc et al. 2016).
There are only a few previous examples of datasets that report either nonsignificant or
negative abundance–range-size relationships from birds in temperate zones (Gaston 1996;
Päivinen et al. 2005). However, several studies have reported nonsignificant or negative
abundance–range-size relationships from the birds in the tropics (Ferenc et al. 2016; Nana et
al. 2014; Reeve et al. 2016; Reif et al. 2006). Although, studies reporting a positive trend
(Theuerkauf et al. 2017) or no trend (Freeman 2018) in the tropics also exist.

The species richness of birds declined (with a lowland plateau) with increasing
elevation on Mt Wilhelm (Sam et al. 2019). This is a typical pattern for mountains with a
humid base (McCain 2009). However, we found that the number of individuals per bird
species increased with increasing elevation and decreasing area. Further investigations of our
data and its partitioning into feeding guilds showed that patterns of abundances for passerine
birds are driven by insectivorous birds, while frugivores drive the decreasing pattern in non-
passerines. This, in turn, is driven solely by the species richness of the feeding guild within
the two groups of birds. A high proportion of the non-passerine birds of Mt. Wilhelm are
identified as frugivorous (44%) and insectivorous (29%), whereas, most of the passerines
(59%) are insectivorous.

The contrasting pattern for the *total abundance* of passerine and non-passerine bird
assemblages is an interesting observation considering the decreasing trend in overall
environmental productivity (McCain 2009) and food availability (estimated by the abundance
of insects and fruits) along the elevational gradient (e.g., Janzen et al. 1976; Loiselle & Blake
1991), especially along wet mountains like Mt. Wilhelm (McCain 2009). Along Mt.
Wilhelm, abundances of arthropods followed a humped-shaped pattern with a peak at ca.
1700 m (Sam et al. 2020, Supriya et al. 2019, Volf et al. 2020), providing the best food
resources for insectivorous passerines in the middle of the gradient. In contrast, abundance as
well as biomass of fruits decreased steeply with increasing elevation (Hazel 2019, Segar et al.
2013). Considering that 71% of non-passerines feed on fruits and/or nectar while at least 68%
of passerines feed solely on insects, the observed patterns in total abundance might be shaped

by the availability of resources. Additionally, abundance patterns in both groups of birds are parallel to the species richness of these groups along our gradient. This corresponds to previously reported results on the relationships between abundance and species richness along elevational gradients (Terborgh 1977).

Mean biomass of bird communities recorded at each elevational study site decreased quite steeply with increasing elevation, showing a different pattern than the *total abundance* of birds at given sites. At the upper most two elevations (3,200 and 3,700 m) mean biomass of passerines was relatively larger than biomass of non-passerines which corresponds, in part, with their *mean elevational abundance* at these elevations. The decrease in bird biomass suggests a decrease in energy flux at given elevations, very likely because of reduction of primary productivity (Dolton & de L. Brooke 1999).

Conclusions

In direct contrast to the abundance-geographical range size relationship hypothesis investigated here, we found that montane species which associated with small geographical ranges have locally higher abundances than lowland species which are associated with large geographical ranges. The *mean abundances* of passerine and non-passerine birds follow a similar trend (significant for passerines, but nonsignificant for non-passerines), with montane birds having higher abundances then lowland birds. Abundances of passerines seem to be driven by insectivores, while non-passerines seem to be driven by frugivores. Our data further show that passerines and non-passerines have different patterns of species richness and *total abundance* along the same elevational gradient.

Acknowledgments

We wish to thank numerous field assistants from Kausi, Numba, Bundi, Bruno Sawmill, Sinopass, and Kegesugl for help in the field and hospitality.

References

Banks☐Leite C, Pardini R, Boscolo D, Cassano CR, Püttker T, Barros CS, and Barlow J. 2014. Assessing the utility of statistical adjustments for imperfect detection in tropical conservation science. *Journal of Applied Ecology* 51:849-859.

Blackburn TM, Cassey P, and Gaston KJ. 2006. Variations on a theme: sources of heterogeneity in the form of the interspecific relationship between abundance and distribution. *Journal of Animal Ecology* 75:1426-1439.

Borregaard MK, and Rahbek C. 2010. Causality of the relationship between geographic distribution and species abundance. *The quarterly review of biology* 85:3-25.

Brown JH. 1984. On the relationship between abundance and distribution of species. *The American Naturalist* 124:255-279.

Buckland S, Anderson D, Burnham K, Laake J, Borchers D, and Thomas L. 2001. Introduction to distance sampling: Oxford University Press. *Oxford Considerations and Management Recommendations: Wildlife Society Bulletin* 34:1393-1395.

Ding TS, Yuan HW, Geng S, Lin YS, and Lee PF. 2005. Energy flux, body size and density in relation to bird species richness along an elevational gradient in Taiwan. *Global Ecology and Biogeography* 14:299-306.

Ferenc M, Fjeldså J, Sedláček O, Motombi FN, Nana ED, Mudrová K, and Hořák D. 2016. Abundance-area relationships in bird assemblages along an Afrotropical elevational gradient:

380   space limitation in montane forest selects for higher population densities. *Oecologia* 181:225-
381   233.
382 Fjeldså J, Bowie RC, and Rahbek C. 2012. The role of mountain ranges in the diversification of birds.
383   *Annual Review of Ecology, Evolution, and Systematics* 43:249-265.
384 Freeman BG. 2018. No evidence for a positive correlation between abundance and range size in birds
385   along a New Guinean elevational gradient. *Emu-Austral Ornithology*:1-9.
386 Gaston KJ. 1996. The multiple forms of the interspecific abundance-distribution relationship.
387   *Oikos*:211-220.
388 Gaston KJ. 1996. Species-range-size distributions: patterns, mechanisms and implications. *Trends in*
389   *Ecology & Evolution*. 11:197-201.
390 Gaston KJ, and Blackburn TM. 2000. *Pattern and Process in Macroecology*. Oxford: Blackwell
391   Publishing.
392 Gaston KJ, Blackburn TM, Greenwood JJ, Gregory RD, Quinn RM, and Lawton JH. 2000.
393   Abundance–occupancy relationships. *Journal of Applied Ecology* 37:39-59.
394 Ghosh-Harihar MJ. 2013. Distribution and abundance of foliage-arthropods across elevational
395   gradients in the east and west Himalayas. *Journal of animal ecology* 28:125-130.
396 Hazel RJ. 2019. Functional alpha- and beta-diversity of birds and fruits on an elevational gradient in
397   Papua New Guinea. PhD thesis. University of Sussex.
398 Hoyo dJ, Elliott J, Sargatal J, and Christie DA. 1992-2011. Handbook of the Birds of the World (vol.
399   1-16). Barcelona, Spain: Lynx Editions.
400 Janzen DH, Ataroff M, Fariñas M, Reyes S, Rincon N, Soler A, Soriano P, and Vera M. 1976.
401   Changes in the arthropod community along an elevational transect in the Venezuelan Andes.
402   *Biotropica* 8:193-203.
403 Johnson C. 1998. Species extinction and the relationship between distribution and abundance. *Nature*
404   394:272-274.
405 Klopfer PH, and MacArthur RH. 1960. Niche size and faunal diversity. *American Naturalist* :293-
406   300.
407 Loiselle BA, and Blake JG. 1991. Temporal variation in birds and fruits along an elevational gradient
408   in Costa Rica. *Ecology* :180-193.
409 MacArthur RH. 1972. *Geographical ecology: patterns in the distribution of species*: Princeton
410   University Press.
411 MacArthur RH, Diamond JM, and Karr JR. 1972. Density compensation in island faunas. *Ecology*
412   53:330-342.
413 Marki PZ, Sam K, Koane B, Kristensen JB, Kennedy JD, and Jønsson KA. 2016. New and
414   noteworthy bird records from the Mt. Wilhelm elevational gradient, Papua New Guinea.
415   *Bulletin of Brittish Ornithology Club* 137:263-271.
416 McAlpine JR, Keig R, and Falls R. 1983. Climate of Papua New Guinea. *CSIRO and Australian*
417   *National University Press, Canberra*.
418 McCain CM. 2009. Global analysis of bird elevational diversity. *Global Ecology and Biogeography*
419   18:346-360.
420 Nana ED, Sedláček O, Bayly N, Ferenc M, Albrecht T, Reif J, Motombi FN, and Hořák D. 2014.
421   Comparison of avian assemblage structures in two upper montane forests of the Cameroon
422   volcanic line: lessons for bird conservation. *Biodiversity and Conservation* 23:1469-1484.
423 Paijmans K. 1976. New Guinea Vegetation. In: Paijmans K, editor. Canberra: National University
424   Press. p 212 pp.
425 Päivinen J, Grapputo A, Kaitala V, Komonen A, Kotiaho JS, Saarinen K, and Wahlberg N. 2005.
426   Negative density-distribution relationship in butterflies. *Bmc Biology* 3:5.
427 Pratt TK, and Beehler BM. 2015. *Birds of New Guinea*: Princeton University Press.
428 Price TD, Hooper DM, Buchanan CD, Johansson US, Tietze DT, Alström P, Olsson U, Ghosh-
429   Harihar M, Ishtiaq F, and Gupta SK. 2014. Niche filling slows the diversification of
430   Himalayan songbirds. *Nature* 509:222-225.
431 Rahbek C. 1995. The elevational gradient of species richness: a uniform pattern? *Ecography* 18:200-
432   205.
433 Reeve AH, Borregaard MK, and Fjeldså J. 2016. Negative range size–abundance relationships in
434   Indo▯Pacific bird communities. *Ecography* 39:990-997.

Reif J, Hořák D, Sedláček O, Riegert J, Pešata M, Hrázský Z, Janeček Š, and Storch D. 2006. Unusual
abundance–range size relationship in an Afromontane bird community: the effect of
geographical isolation? *Journal of Biogeography* 33:1959-1968.
Romdal TS. 2001. Altitudinal distribution and abundance patterns of bird species in the Eastern Arc
Mountains, Tanzania. *Scopus* 21:35-54.
Sam K, and Koane B. 2014. New avian records along the elevational gradient of Mt. Wilhelm, Papua
New Guinea. *Bulletin of the British Ornithologists' Club* 134:116-133.
Sam K, Koane B, Bardos DC, Jeppy S, and Novotny V. 2019. Species richness of birds along a
complete rain forest elevational gradient in the tropics: Habitat complexity and food resources
matter. *Journal of Biogeography* 46:279-290.
Sam K, Koane B, Jeppy S, Sykorova J, and Novotny V. 2017. Diet of land birds along an elevational
gradient in Papua New Guinea. *Scientific Reports* 7:44018.
Sam K, Koane B, Sam L, Mrazova A, Segar S, Volf M, Moos M, Simek P, Sisol M, and Novotny V.
2020. Insect herbivory and herbivores of *Ficus* species along a rain forest elevational gradient
in Papua New Guinea. *Biotropica*. 52:263-276.
Segar ST, Volf M, Zima Jnr J, Isua B, Sisol M, Sam L, Sam K, Souto□Vilarós D, and Novotny V.
2017. Speciation in a keystone plant genus is driven by elevation: a case study in New
Guinean Ficus. *Journal of Evolutionary Biology*. 30:512-523.
Supriya K, Moreau CS, Sam K, and Price TD. 2019. Analysis of tropical and temperate elevational
gradients in arthropod abundance. *Frontiers of Biogeography*, 11:e4310.
Terborgh J. 1977. Bird species diversity on an Andean elevational gradient. *Ecology* 58:1007-1019.
Theuerkauf J, Chartendrault V, Desmoulins F, Barré N, and Gula R. 2017. Positive range–abundance
relationships in Indo□Pacific bird communities. *Journal of Biogeography* 44:2161-2163.
Volf M, Laitila JE, Kim J, Sam L, Sam K, Isua B, Sisol M, Wardhaugh CW, Vejmelka F, Miller SE,
and Weiblen GD. 2020. Compound specific trends of chemical defences in *Ficus* along an
elevational gradient reflect a complex selective landscape. *Journal of Chemical Ecology*.
46:442–454
Williams S, Williams YM, VanDerWal J, Isaac JL, Shoo LP, and Johnson CN. 2009. Ecological
specialization and population size in a biodiversity hotspot: how rare species avoid extinction.
Proceedings of the National Academy of Sciences 106:19737-19741.

465
466

Figure 1. Location of the elevational gradient of Mt. Wilhelm in Papua New Guinea (a) and the study
sites along the gradient (b).

Figure 2. Species richness (fitted with exponential function: $y = -2.4107x^2 + 11.756x + 93.946$, $R^2 =$
0.95) of all birds recorded during point-counts from along the elevational gradient of Mt. Wilhelm (a);
species richness of passerine and non-passerine birds separately (b). Total (i.e. summed) abundances
of passerine (grey) and non-passerine (black) bird assemblages at respective elevational sites (c).

Figure 3. *Mean elevational abundance* of a passerine and non-passerine bird species (±SE) (i.e. mean
number of individuals of a given species at a given elevational site) occurring in the assemblages
along the elevational gradient of Mt Wilhelm (fitted with loess smooth function). The patterns remain
to be valid both in wet and dry season (Figure S5) and when maximal mean elevational abundances
are considered (Figure S6).

Figure 4. Passerine (a ,b) and non-passerine (c, d) birds divided into three groups based on the
position of their weighted mean point of elevational distribution on Mt. Wilhelm, and their *mean*
abundances (a, c) and geographical range sizes in km^2 (b, d). Kruskal-Wallis - passerines (a) $\chi^2 = 16.3$,
df = 2, N = 161, P < 0.001; (b) $\chi^2 = 67.3$, df = 2, N = 161, P < 0.001; non-passerines (c) $\chi^2 = 1.2$, df =
2, N = 88, P = 0.549; (d) $\chi^2 = 19.5$, df = 2, N = 88, P < 0.001. Lowland group = elevational weighted
mean point up to 800m a.s.l., mid group = elevational weighted mean point between 801 and 1600m
a.s.l., and montane group = elevational weighted mean point above 1600 m a.s.l. The patterns remain

valid both in wet and dry season (Figure S9) and when maximal mean elevational abundance point is
considered instead of weighted mean point in analyses (Figure S10).
490
Figure 5. *Mean elevational abundances* of birds partitioned into feeding guilds (a) and *total*
abundance of bird assemblages partitioned into feeding guilds (b). *Mean abundances* of birds
partitioned into feeding guilds and into passerines (c) and non-passerines (d). *Mean elevational*
abundance refers to mean number of individuals of a given species at a given elevation. Subsequently,
mean abundance refers to averaged *mean elevational abundances* of a species across all elevations
where it was present. *Total abundance* refers to aggregated abundances of bird assemblage at a given
elevations. Ne – Nectarivores, In – Insectivores, In-Ne – Insectivore-nectarivores, Fr – Frugivores, Fr-
In – Frugivore-insectivores. Standard errors of the mean are not shown for the clarity of the graph.
Lowland group = elevational weighted mean point up to 800m a.s.l., mid group = elevational
weighted mean point between 801 and 1600m a.s.l., and montane group = elevational weighted mean
point above 1600 m a.s.l.
502
503 Figure 6. Mean biomass (across the re-surveys of all point-counts) of passerine and non-passerine
504 birds (a) and birds partitioned into feeding guilds (b) of Mt. Wilhelm (total biomass in kg/12.86 ha).

532   Figure 1.

551   Figure 2.

554   Figure 3.

571   Figure 4.

587   Figure 5.

590
591

603   Figure 6.