Biomass, abundances, and abundance and range size relationship of birds along a rainforest

elevational gradient in Papua New Guinea

Katerina Sam[1,2], Bonny Koane[3]

[1] Biology Centre CAS, Entomology Institute, Ceske Budejovice, Czech Republic

[2] University of South Bohemia, Faculty of Science, Ceske Budejovice, Czech Republic

[3] The New Guinea Binatang Research Centre, Madang, Papua New Guinea

Keywords: Feeding guilds, Mt. Wilhelm, non-passerines, passerines

Author Contributions: KS conceived and designed the experiments. KS and BK conducted the field

work. KS analysed the data and wrote the manuscript. BK commented on the manuscript.

Abstract

The usual positive inter-specific relationship between range size and abundance of local populations

was shown to have notable exceptions in Afrotropical montane areas, where range-restricted bird

species are unusually abundant. We tested how the local abundances of passerines and non-passerine

of Mt Wilhelm elevational gradient in Papua New Guinea relate to their geographic range size. We

collected the data on bird assemblages at eight elevations (200 – 3,700 m, 500 m elevational

increment) using a standardised point count at 16 locations per elevation. We partitioned the birds into

feeding guilds and we obtained data on range sizes from Bird-Life International data zone. We

observed positive trends of the abundance and range size relationship in lowland changing to negative

trends towards higher elevations. The total assemblage abundances showed hump-shaped pattern,

with passerine birds, namely then passerine insectivores, driving the observed pattern. In contrast to

abundances, mean biomass of the bird assemblages decreased with increasing elevation (i.e. showed

opposite pattern than mean abundances). Our data suggest that montane bird species have abilities to

maintain dense populations which compensate for lower area available near mountain tops.

Introduction

An extensive literature has been devoted to the patterns of species diversity along elevational gradients (McCain 2009; Rahbek 1995). In 2009, McCain (2009) review 150 datasets along elevational gradients and found that birds display four distinct diversity patterns in nearly equal frequency on mountains: decreasing diversity, low-elevation plateaus, low-elevation plateaus with mid-peaks, and unimodal mid-elevational peaks. Biodiversity richness of elevational gradients thus triggered further attention of many scientists, while many aspects of bird assembly remain unknown. Specifically, these studies of species richness have been rarely combined with the study of bird abundance and biomass, arguably more important parameters when it comes to the impact of birds on other trophic levels (but see e.g. Romdal 2001; Terborgh 1977). Even fewer studies have combined these attributes of bird communities with an estimate of available resources (Ding et al. 2005; Ghosh-Harirar and Price 2013; Price et al. 2014) and/or available area along mountain ranges (e.g. Ferenc et al. 2016).

In many studies, attention was not paid to potential differences between passerine and non-passerine species, or passerine species were considered only. Klopfer and MacArthur (1960) suggested that phylogenetically younger passerines should be relatively more abundant in unstable environments. They also showed that the proportions on non-passerines increased from north towards the tropics, where the conditions are more favourable for the existence of non-passerines. In our work, we aimed to test an analogous hypothesis that the non-passerines will be more abundant in favourable tropical lowlands with stable climatic conditions compared to the higher elevations with less favourable environments. In Himalayas, the ratio of passerines to non-passerines increased very slowly between 160 and 2,600 m a.s.l., and abruptly between ca. 3,000 – 4,000 m a.s.l. (Price et al. 2014) (but note that not all non-passerines were surveyed). Similarly, passerine abundance increased relative to non-passerines with increasing elevation in the Andes (Terborgh 1977).

The patterns of abundance or biomass in different feeding guilds with elevation have been also rarely investigated in birds. However, they are important for our understanding of ecosystem dynamics and function; arguably, birds as such do not share many ecological functions (Sekercioglu

2006). Along many elevational gradients, insectivorous birds are the most abundant of all feeding guilds (Ghosh-Harirar and Price 2013; Price et al. 2014; Terborgh 1977).

Ability of the species to occupy large ranges, might also affect their abundances within the range. Macroecological studies have revealed a pervasive positive interspecific correlation between range sizes and abundance in local populations (Gaston 2000; Gaston and Blackburn 2000). Recently, it has been shown that majority of positive abundance range-size relationships have been demonstrated on datasets from temperate regions (but see Blackburn et al. 2006)). Ferenc et al. (2016) discussed that the rule is systematically violated in avian assemblages in African montane forest environments (Reif et al. 2006), where range restricted species are generally more abundant than species with large ranges (Fjeldså et al. 2012). A number of other recent studies of tropical montane taxa report that abundance is uncorrelated with (or negatively correlated to) range size (Nana et al. 2014; Reeve et al. 2016) but see (Theuerkauf et al. 2017). The only existing study on this topic from Papua New Guinea showed that abundance (capture rates) was not related to range size (measured elevational breadth) (Freeman 2018).

Factors underlying high abundances of montane forest species remain unknown, but several mutually non-exclusive hypotheses have been discussed (Ferenc et al. 2016). These are: (1) Long-term eco-climatic stability facilitates ecological specialization, which consequently results in high local abundances of montane species (Fjeldså et al. 2012). (2) High abundances of montane forest species are a result of density compensation in species-poor communities at high altitudes (MacArthur et al. 1972). (3) Locally abundant tropical montane species survive despite their small range sizes, whereas insufficiently abundant species are systematically removed by an extinction filter (Johnson 1998).

In this study, we focused on bird assemblages along the elevational gradient of Mt. Wilhelm in Papua New Guinea. The specific goals were to investigate (1) trends in abundances of birds along the elevational gradient, (2) changes in relative abundances of different groups of birds (passerines and non-passerines, various feeding guilds), and (3) effects of range sizes on the abundance of individual species.

Methods

The study was performed along of Mt Wilhelm (4,509 m a.s.l.) in the Central Range of Papua New

Guinea (Figure 1a, b). The complete rainforest gradient spanned from the lowland floodplains of the

Ramu river (200 m a.s.l., 5° 44'S 145° 20'E) to the timberline (3700 m a.s.l., 5° 47'S 145° 03'E; Fig.

1). The study was completed along a 30 km long transect with eight sites, evenly spaced at 500 m

elevational increments. Because of the steep terrain, elevation could deviate by 50 m within each

study site. Survey tracks and study sites at each elevation were directed through representative and

diverse microhabitats (e.g., ridges, valleys, rivulets; ≥ 250 m from forest edge). Average annual

precipitation is 3,288 mm in the lowlands, rising to 4,400 mm at 3,700 m a.s.l., with a distinct

condensation zone around 2,500 – 2,700 m a.s.l. Mean annual temperature decreases from 27.4°C at

the lowland site to 8.37°C at the tree line at a constant rate of 0.54°C per 100 elevational metres. The

habitats within surveyed transect could be described as lowland alluvial forest (200 m a.s.l.), foothill

forest (700 and 1,200 m a.s.l.), lower montane forest (1,700 – 2,700 m a.s.l.), and upper montane

forest (3,200 and 3,700 m a.s.l.; according to Paijmans, 1976). The typical plant species composition

of forest (Paijmans 1976), general climatic conditions (McAlpine et al. 1983) and habitats at

individuals study sites (Sam and Koane 2014) are described elsewhere.

Data on bird communities were collected in 2010, 2011 and 2012 during the wet and dry

seasons, using a standardised point-count at 16 locations per elevation (Sam and Koane 2014; Sam et

al. 2019) where other survey methods and complete species list are presented). The surveys were

conducted in the mornings between 5:45 and 11:00 am. Each of the 16 sample points had a radius of

50 m (area 0.785 ha per point, which makes 12.56 ha per elevational study site), and were located at

least 150 m apart to diminish the risk of multiple counting of individuals. Each point was visited on

14 different days and the order of the points was changed during each visit to avoid biases due to

daytime. Birds were detected for 15 minutes during each visit at each point. This resulted into 240

minutes of daily surveys. During the point-counts, a distance sampling protocol was used, with birds

recorded in five 10-m-wide radial distance bands (Buckland et al. 2001). Application of detection

adjustments is, however, generally problematic in tropical environments (Banks☐Leite et al. 2014)

and we used the observed abundance estimates instead of the distance sampling-based estimates in

further analyses (see similar reasons and discussion by (Ferenc et al. 2016).

The mean number of individuals of a species recorded at 15-min interval (for all 16 points

together) was recorded as "mean elevational abundance". Elevations between minimal and maximal

range where birds were missing were not considered, i.e. data were not extrapolated. The aggregate

abundances of birds in assemblages at a given elevation (hereafter "total abundance") were calculated

as the sum of mean elevational abundances of all species present. The taxonomy used in this study

followed the International Ornithological Congress World Bird List version 6.1.

The elevational mean-point (i.e. optimal elevational distribution of the species) was calculated

from the lower and upper elevational limit of a species distribution on Mt. Wilhelm. Species were

divided into three exclusive groups according to the position of their elevational mean-point as

follows: (a) species having their elevational mean-point in the lower third of the elevational gradient

(up to 800 m a.s.l.) were included in the "lowland" group, (b) species with mean-point between 800

and 1600 m a.s.l. comprised the "middle" group, and (c) species with their mean-point in the upper

third of the gradient (above 1,600 m a.s.l.) represented the "montane" species group. Note that species

(N = 1) occurring from along the complete gradient (200-3,200 m) thus fall into the group of montane

species. The mean-point and mid-point of elevational range differed slightly (max 640 m) for 22

species. Mean-point moved 13 species from middle to montane group and 9 from lowland to middle

group.

Recorded birds were partitioned into five trophic guilds: insectivores, frugivores, frugivores-

insectivores, insectivores-nectarivores and nectarivores based on dietary information in standard

references (Hoyo et al. 1992-2011; Pratt and Beehler 2015) and our data (Sam et al. 2019; Sam et al.

2017). Abundances of passerines and non-passerines and individual feeding guilds were compared by

non-parametric Kruskal-Wallis tests. We report mean ± SE and abundances per 12.56 ha recorded in

15-minute-long census unless we state otherwise. Range sizes of all birds were obtained from Bird-

Life International data zone web pages accessed in July 2016. Body weight (mean for males) of the

birds were obtained from Hoyo et al. (1992-2011). Bird metabolism was calculated from body weight

according to available equations (McNab 2009).

Results

All together, we recorded 25,715 birds belonging to 249 bird (Supplementary Table S1) species during the point-counts along the elevational gradient of Mt. Wilhelm during this project. Total bird species richness seemed to show a plateau at lower elevations (up to 1700 m a.s.l.) and decreased with increasing elevation afterwards (Figure 2a) (see also Marki et al. 2016; Sam and Koane 2014; Sam et al. 2019 for further results). In contrast, *total abundance* of birds showed a humped shaped pattern, peaking between 1,700 and 2,700 m a.s.l. with ca. 420-450 individuals of all birds per 16 sampling points (i.e. 12.86 ha) (Figure 2c).

Passerines and non-passerines

Passerines were overall more species rich along the elevational gradient, represented by 161 species in comparison to non-passerines represented by 88 species (Figure 2b). We observed a linearly decreasing pattern in species richness of non-passerine birds (y = -5.9167x + 60.056, R² = 0.96) along the elevational gradient and hump-shaped pattern (y = -2.1012x^2 + 18.982x + 27.315, R² = 0.92) in species richness of passerine birds (Figure 2b). The species richness of both passerines and non-passerines correlated significantly with their total abundances (r = 0.82 and 0.84, P = 0.013 and 0.008 respectively) (Figure 2 b, c).

Mean elevational abundances of passerine birds were overall significantly higher (4.16 ± 0.25) than mean elevational abundances of non-passerines (3.12 ± 0.3; U = 5128; $Z_{88, 161}$ = - 3.59, P < 0.001). The number of individuals per bird species increased continually with elevation, with approximately 2.5 times as many individuals per non-passerine species and nearly twice as many individuals per passerine species at the highest elevation as in the lowlands (Supplementary Figure S1). The mean elevational abundances of passerine birds (calculated as the average of elevational abundances across occupied elevations) of individual species increased with their elevational mean-point (Kruskal-Wallis test: χ^2 = 6.16, df = 2, 161, P = 0.046, Figure 2d). Passerine birds having their elevational mean-point in the montane forest (above 1600 m a.sl.) had the highest mean elevational

abundances (Figure 2d), but the smallest range-sizes (Figure 3a). We found no significant change

(Kruskal-Wallis test: $\chi^2 = 0.95$, df = 2, 88, P = 0.62) in mean elevational abundances of non-passerine

birds with elevational mean-point (Figure 2d) but similarly to passerines, non-passerines with higher

elevational midpoint had smaller ranges (Figure 3b). The relationship between local mean abundance

and range size across all bird species of the entire forested gradient of Mt. Wilhelm showed a

significantly negative trend ($F_{1,248} = 8.22$, P = 0.004). The trends remained negative, albeit non-

significant, for passerines ($F_{1,159} = 1.17$, P = 0.28) and non-passerines ($F_{1,86} = 2.6$, P = 0.10) separately

(Supplementary Figure S2).

Feeding guilds

With no respect to which feeding guild they belong, species occurring at low elevations had usually

lower mean elevational abundances than species occurring at high elevations (Figure 4a; i.e. their

mean elevational abundance increased with increasing elevation). The highest mean abundances had

nectarivorous and insectivore-nectarivorous species (Figure 4a) and the pattern was likely driven by

presence of flocks of nectar-feedings lorikeets.

Total abundances of birds belonging to different feeding guilds however showed different

patterns (Figure 4b). While total abundances of frugivores and frugivore-nectarivores decreased with

increasing elevation, total abundance of insectivore-nectarivores and nectarivores showed no trend,

and insectivores showed a mid-elevational peak (Figure 4b).

Within passerine birds, the mean elevational abundances of birds belonging to different

feeding guilds increased with their elevational mean-point (Figure 4c). In contrast, the mean

elevational abundances of non-passerines birds belonging to various feeding guilds showed various

patterns (Figure 4d).

Mean biomass of bird communities (Figure 5) recoded at each elevational study site

decreased with increasing elevation, showing thus opposite pattern than mean elevational abundances.

At the upper most two elevation (3,200 and 3,700 m) mean biomass of passerines was relatively

larger than biomass of non-passerines which corresponds partly also with their mean elevational

abundances at these elevations (Figure 5a). The decreasing pattern in decreasing biomass was

observed both with passerines and non-passerines (Figure 5a), as well as in all feeding guilds (Figure

5b).

Discussion

In this study, we studied the relationships between species richness, abundances, and range sizes

in passerine and non-passerine assemblages along a tropical elevational gradient. Avian species

richness declines monotonically with elevation on Mt. Wilhelm (Sam et al. 2019), which is a typical

pattern for wet-based mountains (McCain 2009). However, total abundances of bird assemblages at

the individual elevations show a different, hump-shaped pattern. This is an interesting observation, as

previous studies showed that unimodal or linearly decreasing patterns on density paralleled the

patterns of total species richness along the same gradients (e.g. (Romdal 2001; Terborgh 1977). Our

findings are also different from patterns in abundances of birds observed along elevational gradient in

Cameroon (Ferenc et al. 2016), where a decline in species richness and uniform abundance of birds

were observed with increasing elevation.

The overall pattern in abundance of bird assemblages we observed can be partitioned into

hump-shaped pattern for passerine birds and decreasing trend for non-passerine birds. Such

partitioned patterns correspond better with respective species richness than overall species richness

and overall abundance. However, note it might be difficult to interpret patterns non-passerine birds,

which represent paraphyletic group of birds. To our knowledge, there is not a single study focusing

separately on abundance pattern in passerine and non-passerine birds along an elevational gradient.

Our data further show that species richness and abundance of passerines increase relative to non-

passerines with increasing elevation. This might be in concordance with previous suggestions that

phylogenetically younger passerines should be relatively more abundant in less favourable and stable

environments. Klopfer and MacArthur (1960) showed that the proportions of non-passerines towards

passerines change from north to south. A study focusing on a similar pattern along an elevational

gradient in Himalaya indicated that ratio between abundances of passerines/non-passerines increases

only very slowly between 160 and 2,600 m a.s.l., and then increased abruptly between ca. 3,000 –

4,000 m a.s.l.

(Price et al. 2014). Unfortunately, this study did not survey all non-passerines (Price et al. 2014).

The widespread pattern that abundance is positively correlated with geographic range size

does not seem to apply to New Guinean birds distributed along elevational gradients (Gaston and

Blackburn 2000). Contrary to this widely accepted pattern, we found a negative relationship between

the local abundance and complete range size of the species. Deviation from positive abundance-area

relationship is caused by a combination of decreasing range sizes and increasing abundances of birds

towards high elevations. Such an observation fits to predictions of the density compensation

hypothesis, which suggest that in species-poor assemblages, individual species may increase their

abundances to fill the available ecological space (MacArthur et al. 1972). Such a hypothesis

ultimately assumes that extinction filters remove small-range species that have scarce local

populations.

Basically, we showed that New Guinean bird species with small ranges select for high local

abundances, as has been suggested for marsupials in Australia (Johnson 1998), birds of the Australian

wet tropics (Williams et al. 2009) or Afromontane birds (Ferenc et al. 2016). There are only few

previous examples of datasets that report either nonsignificant or negative abundance–range-size

relationships from the temperate zone birds (Gaston 1996; Päivinen et al. 2005), but several studies

have reported nonsignificant or negative abundance–range-size relationships from the tropics, both in

birds (Ferenc et al. 2016; Nana et al. 2014; Reeve et al. 2016; Reif et al. 2006). However, studies

reporting a positive trend (Theuerkauf et al. 2017) or no trend (Freeman 2018) in the tropics also

exist.

Avian species richness declines monotonically with elevation on Mt Wilhelm (Sam et al.

2019), which is a typical pattern for wet-based mountains (McCain 2009), and we show here that the

number of individuals per species increases with elevation. Such an observation fits to predictions of

the density compensation hypothesis, which suggests that in species-poor assemblages, individual

species

may increase their abundances to fill the available free ecological space (MacArthur 1972). Further

investigations of our data and its partitioning into feeding guilds showed that patterns of abundances

for passerine birds are driven by insectivorous birds, while frugivores drive the decreasing pattern in

non-passerines. This seems to be given solely by species richness of the feeding guild within the two

groups of birds. While high proportion of the non-passerine birds of Mt. Wilhelm is identified as

frugivores (44%), followed by insectivores (29%), most of the passerines (59%) are insectivorous.

The contrasting pattern for total abundance of passerine and non-passerine bird assemblages

and is an interesting fact considering that overall environmental productivity (McCain 2009) and food

availability as estimated by abundance of insects or fruits seem to decrease with increasing elevation

(e.g. (Janzen et al. 1976; Loiselle and Blake 1991), especially along wet mountains similar to Mt.

Wilhelm (McCain 2009). On the other hand, observed patterns in abundances of both groups of birds

are parallel to the species richness of these groups along our gradient which corresponds with

previously reported results on relationship on abundance and species richness along elevational

gradients (Terborgh 1977).

Mean biomass of bird communities recoded at each elevational study site decreased quite

steeply with increasing elevation, showing thus opposite pattern than mean elevational abundances.

At the upper most two elevation (3,200 and 3,700 m) mean biomass of passerines was relatively

larger than biomass of non-passerines which corresponds partly also with their mean elevational

abundances at these elevations. The decreasing pattern in decreasing biomass was observed both with

passerines and non-passerines, as well as in all feeding guilds. The decrease in biomass suggest

decrease in energy flux into the birds at given elevation, very likely as a result of reduction f primary

productivity (Dolton and de L. Brooke 1999).

In conclusion, our data show that passerines and non-passerines have different patterns of

species richness and total abundance along the same elevational gradient. However, the mean

elevational abundances of passerine and non-passerine birds follow a similar trend (significant for

passerines, but non-significant for non-passerines), with montane birds having higher abundances then

lowland birds. This observation is in direct contrast to abundance-area relationship hypothesis.

Abundances of passerines driven seems to be driven by insectivores, while non-passerines seem to be

driven by frugivores.

Acknowledgement

We wish to thank numerous field assistants from Kausi, Numba, Bundi, Bruno Sawmill, Sinopass,

and Kegesugl for help in the field and hospitality. The work of K. Sam was supported by Czech

Science Foundation Grants 18□23794Y and the infrastructure and logistics of the project was

financially

supported by the European Science Foundation 669609 grant, the Darwin Initiative for the Survival of

Species grant 22□002 and the Christensen Foundation grant 2016□8734.

Conflict of Interest: The authors declare that they have no conflict of interest.

References

Banks□Leite C et al. (2014) Assessing the utility of statistical adjustments for imperfect detection in

tropical conservation science. Journal of Applied Ecology 51:849-859

Blackburn TM, Cassey P, Gaston KJ (2006) Variations on a theme: sources of heterogeneity in the

form of the interspecific relationship between abundance and distribution. Journal of Animal

Ecology 75:1426-1439

Buckland S, Anderson D, Burnham K, Laake J, Borchers D, Thomas L (2001) Introduction to

distance sampling: Oxford University Press. Oxford. Considerations and Management

Recommendations: Wildlife Society Bulletin 34:1393-1395

Ding TS, Yuan HW, Geng S, Lin YS, Lee PF (2005) Energy flux, body size and density in relation to

bird species richness along an elevational gradient in Taiwan. Global Ecology and Biogeography

14:299-306

Dolton CS, de L. Brooke M (1999) Changes in the biomass of birds breeding in Great Britain, 1968–

88. Bird Study 46:274-278

Ferenc M et al. (2016) Abundance-area relationships in bird assemblages along an Afrotropical

elevational gradient: space limitation in montane forest selects for higher population densities.

Oecologia 181:225-233

Fjeldså J, Bowie RC, Rahbek C (2012) The role of mountain ranges in the diversification of birds.

Annual Review of Ecology, Evolution, and Systematics 43:249-265

Freeman BG (2018) No evidence for a positive correlation between abundance and range size in birds

along a New Guinean elevational gradient. Emu-Austral Ornithology:1-9

Gaston KJ (1996) The multiple forms of the interspecific abundance-distribution relationship.

Oikos:211-220

Gaston KJ (2000) Global patterns in biodiversity. Nature 405:220-227

Gaston KJ, Blackburn TM (2000) Pattern and process in macroecology. Blackwell Publishing, Oxford

Ghosh-Harirar M, Price TD (2013) A test for community saturation along the Himalayan bird

diversity gradient, based on within-species geographical variation. Journal of Animal Ecology.

83:628-38

Hoyo dJ, Elliott J, Sargatal J, Christie DA (eds) (1992-2011) Handbook of the Birds of the World

(vol. 1-16). Lynx Editions, Barcelona, Spain

Janzen DH et al. (1976) Changes in the Arthropod Community along an Elevational Transect in the

Venezuelan Andes. Biotropica 8:193-203

Johnson C (1998) Species extinction and the relationship between distribution and abundance. Nature

394:272-274

Klopfer PH, MacArthur RH (1960) Niche size and faunal diversity. American Naturalist 94:293-300

Loiselle BA, Blake JG (1991) Temporal variation in birds and fruits along an elevational gradient in

Costa Rica. Ecology 72:180-193

MacArthur RH (1972) Geographical ecology: patterns in the distribution of species. Princeton

University Press

MacArthur RH, Diamond JM, Karr JR (1972) Density compensation in island faunas. Ecology

53:330-342

Marki PZ, Sam K, Koane B, Kristensen JB, Kennedy JD, Jønsson KA (2016) New and noteworthy

bird records from the Mt. Wilhelm elevational gradient, Papua New Guinea. Bulleting of Brittish

Ornithological Club 137:263-271

McAlpine JR, Keig R, Falls R (1983) Climate of Papua New Guinea. CSIRO and Australian National

University Press, Canberra

McCain CM (2009) Global analysis of bird elevational diversity. Global Ecology and Biogeography

18:346-360

McNab BK (2009) Ecological factors affect the level and scaling of avian BMR. Comparative

Biochemistry and Physiology Part A: Molecular & Integrative Physiology 152:22-45

Nana ED et al. (2014) Comparison of avian assemblage structures in two upper montane forests of the

Cameroon volcanic line: lessons for bird conservation. Biodiversity and conservation 23:1469-

1484

Paijmans K (ed) (1976) New Guinea Vegetation. National University Press, Canberra

Päivinen J et al. (2005) Negative density-distribution relationship in butterflies. Bmc Biology 3:5

Pratt TK, Beehler BM (2015) Birds of New Guinea. Princeton University Press

Price TD et al. (2014) Niche filling slows the diversification of Himalayan songbirds. Nature

509:222-225

Rahbek C (1995) The elevational gradient of species richness: a uniform pattern? Ecography 18:200-

205

Reeve AH, Borregaard MK, Fjeldså J (2016) Negative range size–abundance relationships in

Indo☐Pacific bird communities. Ecography 39:990-997

Reif J et al. (2006) Unusual abundance–range size relationship in an Afromontane bird community:

the effect of geographical isolation? Journal of Biogeography 33:1959-1968

Romdal TS (2001) Altitudinal distribution and abundance patterns of bird species in the Eastern Arc

Mountains, Tanzania. Scopus 21:35-54

Sam K, Koane B (2014) New avian records along the elevational gradient of Mt. Wilhelm, Papua

New Guinea. Bulletin of the British Ornithologists' Club 134:116-133

Sam K, Koane B, Bardos DC, Jeppy S, Novotny V (2019) Species richness of birds along a complete

rain forest elevational gradient in the tropics: Habitat complexity and food resources matter.

Journal of Biogeography 46:279-290

Sam K, Koane B, Jeppy S, Sykorova J, Novotny V (2017) Diet of land birds along an elevational

gradient in Papua New Guinea. Scientific reports 7: 44018

Sekercioglu CH (2006) Increasing awareness of avian ecological function. Trends in ecology &

evolution 21:464-471

Terborgh J (1977) Bird species diversity on an Andean elevational gradient. Ecology 58:1007-1019.

doi: doi:10.2307/1936921

Theuerkauf J, Chartendrault V, Desmoulins F, Barré N, Gula R (2017) Positive range–abundance

relationships in Indo Pacific bird communities. Journal of Biogeography 44:2161-2163

Figure 1. Location of the elevational gradient of Mt. Wilhelm in Papua New Guinea (a) and the study

sites along the gradient (b).

Figure 2. Bird species richness (**c**, fitted with exponential function: $y = -2.4107x^2 + 11.756x +$

93.946, $R^2 = 0.95$) of all birds recorded during point-counts from along the elevational gradient of Mt.

Wilhelm; species richness of passerine and non-passerine birds separately (**b**). Total abundance of

passerine and non-passerine birds (left axis, lines) and all birds together (right axis, grey bars) (**b**) and

mean elevational (±S.E.) abundance of all passerine and non-passerine species (**c**). Mean elevational

abundance of passerine and non-passerine species divided into the three groups based on the position

of their mid-point of elevational distribution on Mt. Wilhelm (**d**; Kruskal-Wallis test: passerines $\chi^2 =$

6.16, $df = 2$, 161, $P = 0.046$ and non-passerines $\chi^2 = 0.95$, $df = 2$, 88, $P = 0.62$). Part d) reports mean,

conf. int., non-outlier range. Lowland group = elevational mid-point up to 800m a.s.l., middle group =

elevational mid-point between 801 and 1600m a.s.l., and montane group = elevational mid-point

above 1600 m a.s.l.

Figure 3. Passerine and non-passerine birds divided into three groups based on the position of their

midpoint of elevational distribution on Mt. Wilhelm and their range sizes in km^2. Note log scale used

on y-axis. Passerines: Kruskal-Wallis $\chi^2 = 88.45$, df = 2, N = 161, P < 0.001, non-passerines: Kruskal-

Wallis $\chi^2 = 23.14$, df = 2, N = 88, P < 0.001. Lowland group = elevational mid-point up to 800m a.s.l.,

mid group = elevational mid-point between 801 and 1600m a.s.l., and montane group = elevational

mid-point above 1600 m a.s.l.

Figure 4. Mean elevational abundances of birds partitioned into feeding guilds (a) and total

abundance of bird assemblages partitioned into feeding guilds (b). Mean elevational abundances of

birds partitioned into feeding guilds and into passerines (c) and non-passerines (d). Ne – Nectarivores,

In – Insectivores, In-Ne – Insectivore-nectarivores, Fr – Frugivores, Fr-In – Frugivore-insectivores.

Standard errors of the mean are not shown for the clarity of the graph. Lowland group = elevational

mid-point up to 800m a.s.l., mid group = elevational mid-point between 801 and 1600m a.s.l., and

montane group = elevational mid-point above 1600 m a.s.l.

Figure 5. Mean biomass (across the re-surveys of all point-counts) of passerine and non-passerine

birds (a) and birds partitioned into feeding guilds (b) of Mt. Wilhelm (total biomass in kg/12.86 ha).

441   Figure 1.

460   Figure 2.

476   Figure 3.

497   Figure 4.

512   Figure 5.