Title: Environmental (e)DNA detection of the invasive pink salmon *Oncorhynchus gorbuscha* during

the 2017 Norwegian invasion

Authors: Laura M. Gargan[1], Frode Fossøy[2], Tor A. Mo[2], Jeanette E. L. Carlsson[1], Bernard Ball[1], Jens

Carlsson[1]

[1]Area 52 Research Group, Earth Institute/School of Biology and Environmental Science, University

College Dublin, Ireland

[2]Norwegian Institute for Nature Research (NINA), Trondheim, Norway

Corresponding author: Laura M. Gargan, E-mail: laura.gargan@ucd.ie

Keywords: *Oncorhynchus gorbuscha*; invasive species; digital droplet PCR; environmental DNA;

Norway

ABSTRACT

The pink salmon *Oncorhynchus gorbuscha* was introduced from its native range in the Pacific to

Northwest Russia several times since the 1950's. While this species has been regularly observed in

rivers in Northern Norway since that time, there has been an upsurge in the numbers of odd-year *O.*

gorbuscha individuals observed in rivers in southern Norway in recent years, and particularly in 2017.

Although the wide-scale effects of this species presence are currently uncertain, there are concerns

regarding potential competition between *O. gorbuscha* and native species – most notably the Atlantic

salmon *Salmo salar.* Environmental (e)DNA is becoming a widely used tool to monitor rare and

invasive species in aquatic environments. In the present pilot study, primers and a probe were

developed to detect *O. gorbuscha* from eDNA samples taken from a Norwegian river system where

the species was observed. Water samples were taken at both upstream and downstream locations of

the Lysakerelva river during Autumn 2017 (to coincide with spawning) and during late Spring 2018.

Autumn samples were positive for *O. gorbuscha* at both sampling locations, whereas Spring samples

showed positive detection of this species in the upstream region of the river, when smolt should have

left, or be in the process of leaving the river. These findings reveal that eDNA-based methods can be

used detect the presence of *O. gorbuscha* during their spawning season. This suggests that odd-year

populations have the potential to become established in the studied river system. We recommend that

eDNA sampling is repeated to determine whether individuals of this odd-year population have

survived at sea and return to spawn. Our assay specificity tests indicate that the tools developed in the

present study can be used for detection of *O. gorbuscha* in both Norwegian and other European river

systems where presence/absence data is required. We also suggest some modifications to our

methodology that may improve upon the detection capabilities of *O. gorbuscha* using eDNA.

INTRODUCTION

Pink salmon *Oncorhynchus gorbuscha* (Walbaum, 1792) are native to the Pacific Ocean,

where they typically spawn in the freshwater ecosystems of bordering countries between the latitudes

of 40° and 70°. This species follows a strict two-year life cycle (Figure 1). Adult *O. gorbuscha*

migrate from the open sea and up-river to spawn in the autumn, after which all spawning individuals

die. The juveniles emerge the following spring, ready to migrate down-river and out into the open sea

to mature for one winter. They return as adults to freshwater in the next autumn to spawn, thus

completing the life cycle (Heard 1991).

O. gorbuscha was originally introduced from their native Pacific range to North Western

Russia several times since the 1950's, when fry were stocked in several rivers that drain into the

White Sea and the Barents Sea (Bakshtansky 1980). While *O. gorbuscha* has been regularly found in

rivers in Northern Norway since 1960 (Berg 1961), there has been an upsurge in the observations of

odd-year *O. gorbuscha* in Norwegian rivers in recent years (Mo *et al.,* 2018), as well as in rivers in

the UK and Ireland (Armstrong *et al.,* 2018, Whelan 2017, Millane *et al*., 2019). However, self-

reproducing populations have not yet appeared to become established, most likely because they have

not adapted time of spawning to match local conditions outside of their native range (Mo *et al.,* 2018).

55   Established populations of *O. gorbuscha* in rivers in North Western Russia are dominated by

odd-year individuals (Gordeeva & Salmenkova, 2011). Thus far, odd-year individuals have been

noticeably more abundant throughout their invasive range, with a particularly large number of

spawning *O. gorbuscha* recorded in 2017 (Armstrong *et al.,* 2018, Mo *et al.,* 2018). Therefore, it is

these odd-year stocks that are most likely to become established as populations and the next spawning

season (which should take place during Autumn of 2019) will be an important time for research

scientists to determine the extent of this invasion and its potential ecological effects.

62   Presently, the spawning time of *O. gorbuscha* in Norway does not appear to overlap with

native salmonids (e.g. Atlantic salmon *Salmo salar* and brown trout *S. trutta*). However, *O. gorbuscha*

and *S. salar* have similar preferences for spawning habitats and so there is a risk of competition for

optimal spawning sites (Sandlund *et al.,* 2018). While *O. gorbuscha* juveniles emerge ready to

migrate to sea, observations in Norwegian rivers suggest that they spend some time feeding in

freshwater (from weeks to months) and during this time there may be interactions between juveniles

of native salmonids (Sandlund *et al.,* 2018). However, it is also possible that the eggs and fry of *O.*

gorbuscha can provide a source of food for other native salmonid species (Rasputina *et al.,* 2016). In

order to fully assess the impacts of the presence of *O. gorbuscha* in Norwegian (and other European)

rivers, it is first necessary to determine the spatial, as well as the temporal (e.g. time of spawning and

migration) distribution of this species.

73   Environmental DNA (eDNA) is a genetic survey method that relies on the detection of taxa

from extracellular and intracellular material that is deposited into the environment. Subsequently, this

material can be isolated from the environmental sample (such as water, air or soil; Taberlet *et al.,*

2012) and interrogated using genetic markers for multi-species (Thomsen *et al.,* 2012, Hänfling *et al.,*

2016) or targeted species (Ficetola *et al.,* 2008, Jerde *et al.,* 2011, Gustavson *et al.,* 2015) detection.

While many studies have used quantitative PCR (qPCR) for targeted detection of species in

an eDNA sample (Thomsen *et al.,* 2012, Wilcox *et al.,* 2013, Atkinson *et al.,* 2018), more recently the

approach of using digital droplet PCR (ddPCR) has been adopted by some researchers (Doi *et al.,*

2015, Nathan *et al.,* 2015, Evans *et al.,* 2017, Baker *et al.,* 2018), with either similar (Nathan *et al.,*

2015) or increased (Doi *et al.,* 2015, Hunter *et al.,* 2017) sensitivity reported for ddPCR platforms

compared to qPCR for the analysis of eDNA. ddPCR is a relatively recent technological

advancement, allowing for accurate estimation of low copy DNA number (Hindson *et al.,* 2011). It is

an absolute quantification method which, unlike qPCR, does not rely on standard curves to estimate

target DNA concentration. For DNA samples extracted from environmental water and containing

potentially very low copy number of target DNA, the ability of ddPCR to detect rare events in a

reaction is of particular interest, especially for invasive species in aquatic ecosystems where they may

exist in low abundance and may be difficult to detect using conventional survey methods (e.g. netting

and electrofishing).

Whether analysis is being carried out using qPCR or ddPCR, eDNA-based detection is

becoming increasingly used in aquatic freshwater environments for a range of low abundance or

invasive taxa, such as amphibians (Pilliod *et al.,* 2013, Spear *et al.,* 2014), molluscs (Goldberg *et al.,*

2013, Peñarrubia *et al.,* 2016, Carlsson *et al.,* 2017) and crustaceans (Tréguier *et al.,* 2014, Harper *et*

al., 2018), as well as fish (Takahara *et al.,* 2013, Klymus *et al.,* 2015, Davison *et al.,* 2016) –

including salmonid species (Gustavson *et al.,* 2015, Atkinson *et al.,* 2018, Rusch *et al.,* 2018). To

date, there has been no published studies for implementing eDNA for the detection of *O. gorbuscha.*

We hypothesise that eDNA can be used for detection of non-native *O. gorbuscha* in running water. In

addition, should any eradication measures be employed in the future, eDNA methods could be used to

determine the efficacy of such efforts (Banks *et al.,* 2015).

This study aimed to address questions relating to the presence and distribution of *O.*

gorbuscha in a Norwegian river, as part of a pilot study employing the non-intrusive and genetic

survey method of eDNA collection and analysis. A species-specific probe-based assay was developed

for this species and deployed in an urban river system where *O. gorbuscha* had been previously

observed. We also developed this assay with the intention that it can be deployed in other European

freshwater or marine ecosystems where this species may currently, or potentially, invade. This is of

particular relevance for detection of the species during spawning season of this year (Autumn 2019),

where it is unknown if the putative increased mortality of *O. gorbuscha* at sea (resulting from a

mismatch between emigration time and food availability; Armstrong *et al.,* 2018) will result in a less

significant invasion.

MATERIALS AND METHODS

Water sampling and eDNA extraction

114   For this study, water samples were collected from the Lysakerelva river in the south-east

region of Norway (Table 1 and Figure 2) during Autumn of 2017 and Spring/early Summer of 2018.

This river system runs through a large urban area outside of Oslo and is characterised by a number of

natural migration barriers, including waterfalls. The smallest of these waterfalls (Møllefoss; Figure 2)

is a few metres in height and is fitted with fish ladders to enable upstream migration. The next

waterfall, Granfoss (Figure 2), is sufficiently high (~12-15 metres) as to completely prevent upstream

fish migration. In terms of the diversity of fish found in the study area, the Lysakerelva river fish

fauna is dominated by *S. salar*, *S. trutta* and the European minnow *Phoxinus phoxinus* (Saltveit *et al.,*

2013). Several *O. gorbuscha* have previously been caught in the Lysakerelva river (>20 fish;

Sandlund *et al.,* 2018).

124   During September 2017, two water samples were taken below Møllefoss and just above upper

high tide close to the mouth of the Lysakerelva river, and two water samples were taken immediately

below the Granfoss waterfall (~800 metres upstream from the river mouth; Table 1 and Figure 2).

These samples were taken to coincide with the spawning season of *O. gorbuscha.* This sampling was

repeated the following May, the time period when it is expected that juvenile *O. gorbuscha* would be

undertaking seaward migration (Table 1). In June 2018, two water samples were also taken from the

estuary (Figure 2) as it was considered possible that *O. gorbuscha* may also be found in the coastal

area after migrating downriver (Heard 1991; Moore *et al.,* 2016). Additional samples (*n*=4; Table 1)

were taken from above the Granfoss waterfall in June 2018 (Figure 2) as negative eDNA field control

samples, as it was expected that no *O. gorbuscha* would be present at this location due to the barrier

of the Granfoss waterfall.

For each locality, we filtered two replicate samples of either *c.* 1 L water on a 0.45 µm

cellulose filter (Pall MicroFunnel 300 ST; Pall Corporation, New York, USA; Table 1) or two

replicate samples of *c.* 10 L water on a 2.0 µm glassfiber filter (Merck Millipore, Burlington,

Massachusetts, USA; Table 1) using a peristaltic pump (Vampire sampler, Bürkle, Bad Bellingen,

Germany). The 0.45 µm cellulose filters were immediately placed in 2 mL tubes with 1440 µL ATL-

buffer (Qiagen, Hilden, Germany), whereas the 2.0 µm glassfiber filters were placed in 5 mL tubes

with 4050 µL ATL-buffer. All samples were stored at room temperature until further processing in the

genetics laboratory. All field equipment (e.g. filtering tubes and collection bottles) was sterilised

between collection of each sample using 10% bleach solution for approximately 60 minutes.

In the laboratory, 160 µL or 450 µL Proteinase-K (Qiagen) was added to the 2 mL and 5 mL

sampling tubes, respectively. All samples were incubated overnight at 56°C. DNA was isolated from

0.45 µm cellulose filters using DNeasy DNA Blood & Tissue kit (Qiagen), and from the 2.0 µm

glassfiber filters using NucleoSpin Plant II Midi kit (Macherey-Nagel, Düren, Germany), following

the manufacturers protocol except that Qiagen buffers were used instead of those supplied with the

kit. DNA extracted from the 0.45 µm cellulose filters was eluted in 100 µL AE buffer, whereas DNA

extracted from the 2.0 µm glassfiber filters was eluted in 200 µL AE buffer. All samples were re-

eluted in order to maximise the output of DNA. Final concentrations of the eluted DNA samples

(Supplementary Table 1) were determined using a Nanodrop 1000 Spectrophotometer (Thermo Fisher

Scientific, Waltham, Massachusetts, USA).

 Molecular assay development and specificity testing

An assay was designed to amplify a 98 bp region of the mitochondrial *COI* gene of *O.*

gorbuscha (Table 2). This assay consisted of primers and a 5' VIC labelled TaqMan® minor groove

binding (MGB) probe. The primers and probes were designed using Primer3 software (Rozen and

Skaletsky 2000). Specificity of the assay was checked *in-silico*, by aligning the primers and probe

with the consensus sequence generated from publicly available *O. gorbuscha* sequences as well as

those from other salmonids commonly occurring in the study area (e.g. *S. salar* and *S. trutta*). The

primer and probe sequences were also checked against the NCBI database to ensure specificity to the

target organism. Furthermore, specificity of the assay was checked using qPCR, with tissue-extracted

DNA from *O. gorbuscha*, as well as from *S. salar* and *S. trutta*. Tissue-extracted DNA was also

acquired and tested by qPCR for rainbow trout (*O. mykiss*) which, while not a native salmonid, has

been widely introduced throughout Europe (Stanković *et al.,* 2015). In addition, DNA extracted from

the closely-related chum salmon *O. keta* tested to check the specificity of the assay. The latter is not

currently found in Europe but this species overlaps with *O. gorbuscha* in its native range.

All qPCR reactions took place in a 20 µl reaction volume, containing 10 µl of TaqMan™

Environmental Master Mix 2.0 (Thermofisher), 2 µl of each primer (2 µM), 2 µl of probe (2 µM;

Applied Biosystems) and 2 µl of template DNA (where extracts were normalised to 34 ng/µl). The

PCR program consisted of 50°C for 2 min, 95°C for 10 min, followed by 40 cycles of 95°C for 15 s

and 60°C for 1 min. All qPCR reactions were carried out using QuantStudio™ 7 Flex Real-Time PCR

System (Applied Biosystems). All qPCR analysis took place in University College Dublin.

Potential cross-amplification of other salmonids (*O. mykiss, S. salar, S. trutta, Salvelinus*

alpinus, Salvelinus fontinalis, Salvelinus namaycush and Thymallus thymallus) using the *O.*

gorbuscha assay was also tested using ddPCR at NINA using the same PCR conditions as those

detailed in the next section.

 Digital droplet (dd)PCR analysis of eDNA samples

Sample collection and eDNA extraction resulted in a total of 14 samples originating from the

Lysakerelva river (Table 1). Detection and concentration of target-DNA was assessed using droplet-

digital-PCR (QX200 Droplet Digital PCR system with AutoDG, Bio-Rad Laboratories, Hercules,

USA). A tissue-extracted DNA sample of *O. gorbuscha* was included in the analysis, as a positive

control. A no-template control was also included in the analysis. The eDNA samples, along with

positive, negative and no-template controls, were analysed in triplicate with the exception of the

samples from 2017 (Table 1), which were analysedsingularly. The ddPCR reactions consisted of 0.9

µM forward and reverse primers, 0.25 µM of the probe, ddPCR™ Supermix for Probes (No dUTP)

(Bio-Rad Laboratories), dH$_2$O and 5 µL template. In order to assess potential problems with inhibition

in our samples we also reran the samples with only 1 µL template. However, we found no indications

of inhibition among our samples. To generate droplets, an AutoDG Instrument (Bio-Rad

Laboratories) was used, with subsequent PCR amplification in a Veriti96-Well Thermal Cycler

(Applied Biosystems). The following thermal cycling conditions were used: an initial denaturation

step at 95°C for 10 min, 40 cycles of denaturation at 95°C for 30 sec, annealing and extension at 60°C

for 1 min, a final step of denaturation at 98°C for 10 min, and a final hold at 4°C. PCR plates were

transferred to a QX200 Droplet Reader (Bio-Rad Laboratories) to automatically detect the fluorescent

signal in the droplets. QuantaSoft software v.1.7.4 (Bio-Rad Laboratories) was used to separate

positive from negative droplets according to manufacturer's instructions. A threshold of minimum 3

positive droplets were used as a criterion for positive detection Wacker et al. 2019). All ddPCR-

analyses took place at the genetic lab at NINA in Trondheim, Norway.

RESULTS

Molecular assay development

Testing of the *O. gorbuscha* assay with non-target salmonids using tissue-extracted DNA and

qPCR, revealed that the assay did not cross-amplify *S. salar*, *S. trutta,* or *O. mykiss.* ddPCR testing of

non-target salmonid species (*O. mykiss, S. salar, S. trutta, S. alpinus, S. fontinalis, S. namaycush, T*

thymallus) did not produce any detectable amplification. However, amplification was observed for *O.*

keta DNA analysed by qPCR (data not shown). This species is closely related to *O. gorbuscha* and

there are very few nucleotide differences between these species for the *COI* region targeted by our

assay (Supplementary Figure 3). However, this should not present a major concern for researchers

wishing to apply this assay for *O. gorbusha* detection in Norwegian and other European locations, as

neither *O. keta* nor other species of Pacific salmon (*O. nerka*, *O. kisutch*, *O. tshawytscha* and *O.*

masoudo) currently occur in these regions.

ddPCR analysis of eDNA samples

An average number of 15,299 droplets were analysed for each sample included in the ddPCR

analyses. Using the *O. gorbuscha* assay in a ddPCR analysis, there was target DNA detected in six out

of 14 eDNA samples (Table 3).

In Autumn 2017, *O. gorbuscha* DNA was detected at Granfoss but not at Møllefoss (Table 3,

Figure 3). In Spring, target DNA was detected in both samples taken below the Granfoss waterfall,

which indicates that there were still young *O. gorbuscha* in the sampling area that had not yet left the

river. No target DNA was found in samples taken near to the mouth of the river in June 2018, which

is perhaps unsurprising, considering that upstream and downstream samples were taken 20 days apart

and young *O. gorbuscha* may have already migrated to sea.

No detectable target DNA was found in samples taken from the estuary. It is possible that,

similar to the samples taken in the downstream region of the Lysakerelva river, young *O. gorbuscha*

may have already migrated out to sea when this sampling took place in the Summer. Alternatively, a

lack of detection of *O. gorbuscha* at this location may be a false negative, as a result of the relatively

low number of samples taken over a large sampling area (Table 1 and Figure 2).

Interestingly, target DNA was also detected in low copy number from one out of four of the

control samples above the migration barrier (Figure 3) indicating alternative means of target DNA

transport to this site, or contamination of the sample in the field (see Discussion). All no-template

controls were negative for target DNA.

DISCUSSION

The year 2017 was unprecedented in terms of the number of *O. gorbuscha* that were observed

in Norwegian rivers (Mo *et al.,* 2018), as well as in Ireland (Whelan 2017, Millane *et al*., 2019) and

Scotland (Armstrong *et al.,* 2018,). In the present study, we were successfully able to detect *O.*

gorbuscha from environmental water samples in Norway during this invasion. Target DNA was

amplified from samples taken in both Autumn (during adult spawning) and the following Spring

(during the migration of juveniles), indicating that spawning in the focal river system was successful.

However, the survival of juvenile *O. gorbuscha* at sea is poorly understood and it is unknown whether

mortality will limit the ability of this species to return to the river and establish self-reproducing

populations.

The assay developed and validated in this study not only has applications for monitoring

presence of *O. gorbuscha* spatially and temporally in Norwegian rivers, but also in countries where it

currently exists outside of its native range, as well as those where it has not been observed but may

potentially occur in the future. Further, this assay can be used to monitor *O. gorbuscha* presence

following any future eradication efforts that may be implemented. While the utility of the assay

developed in this study is limited to those areas where *O. gorbuscha* does not overlap with *O. keta*,

this should not be of concern for researchers using this assay to detect *O. gorbuscha* within its

invasive range as *O. keta* is not currently found outside the North Pacific.

The finding of a positive detection of *O. gorbuscha* above the impassable barrier of Granfoss

waterfall demonstrates the potential drawbacks of using eDNA to infer species presence. There was

no indication of contamination in any of the negative controls included in this study. Specificity

testing by qPCR and ddPCR revealed that this assay does not amplify *COI* from other commonly co-

occurring native or introduced salmonids. The sample location was the only site visited on the day of

sampling, yet it is impossible to rule out contamination of field equipment from previous sampling

events downstream of the barrier. It is also possible that an alternative vector, such as predation (by

birds for example; Merkes *et al.,* 2014), can explain the detection of target DNA in a location where

the organism is not present.

The results of our pilot study show that this assay can be used to detect the presence of *O.*

gorbuscha in running water. However, we can make some suggestions to increase the efficacy of our

approach for future studies. The lack of detection of *O. gorbuscha* in water samples taken from the

estuary and at the mouth of the Lysakerelva river in early Summer indicated that the juveniles had

already migrated to sea. We would recommend an increase in temporal sampling, as data derived

from these samples may have been able to reveal the timing of juvenile migration to sea, increasing

our knowledge about *O. gorbuscha* behaviour in Norwegian rivers and indicating the extent of

potential interactions with local fauna.

We detected a mismatch in the forward primer sequence (Supplementary Figure 1), based on

the *O. gorbuscha* consensus sequence generated from publicly available *COI* records on GenBank.

The source of this mismatch is a sequence from an odd-year individual (Accession No. MG951587.1)

from the White Sea that was submitted to the NCBI database after the assay was developed and

tested. It is therefore possible that this haplotype is found in Norwegian rivers. This mismatch is

found on the 5' end of the forward primer, therefore it is unlikely that our assay efficiency was

severely compromised in the present study. However, to ensure optimal efficiency in the assay (and

particularly to ensure sensitive detection of the low copy numbers frequently found in eDNA

samples), we recommend that the forward primer incorporate a degenerate base at this position should

the assay be deployed in other studies. Further, we concur with other eDNA researchers (e.g.

Goldberg *et al.,* 2016) that concerted efforts are made to minimise any contamination in the field,

through the implementation of careful sterilisation and sampling techniques, as well as the use of

strict controls (e.g. field blanks, filter blanks) to monitor for exogenous sources of DNA during the

entire eDNA sampling and analysis workflow.

ACKNOWLEDGMENTS

The authors would like to thank Hege Brandsegg for running the ddPCR genetic analysis. We also

gratefully acknowledge Sigrid Skoglund for drawing the pink salmon life cycle.

COMPLIANCE WITH ETHICAL STANDARDS

Conflict of interest The authors declare that they have no conflict of interest.

Ethical approval All applicable international, national, and/or institutional guidelines for the care and

use of animals were followed.

REFERENCES

Atkinson, S., Carlsson, J. E. L., Ball, B., Egan, D., Kelly-Quinn, M., Whelan, K., and Carlsson, J.
(2018) 'A quantitative PCR-based environmental DNA assay for detecting Atlantic salmon (*Salmo*
salar L.)', *Aquatic Conservation: Marine and Freshwater Ecosystems*, 28(5), pp:1238–1243.
https://doi.org/10.1002/aqc.2931

Baker, C. S., Steel, D., Nieukirk, S., and Klinck, H. (2018) 'Environmental DNA (eDNA) From the
Wake of the Whales: Droplet Digital PCR for Detection and Species Identification', *Frontiers in*
Marine Science, 5. https://doi.org/10.3389/fmars.2018.00133

Bakshtansky, E. L. (1980) 'Pink salmon in the Kola Peninsula', In: Thorpe JE (ed) *Salmon ranching*.
Academic Press, New York, pp:245–259

Banks, J. C., Demetras, N. J., Hogg, I. D., Knox, M. A., & West, D. W. (2016) 'Monitoring brown
trout (*Salmo trutta*) eradication in a wildlife sanctuary using environmental DNA', *New Zealand*
Natural Sciences, 41, pp:1–13. https://doi.org/10289/10302

Berg, M. (1961) 'Pink salmon (*Oncorhynchus gorbuscha*)in northern Norway in the year 1960', *Acta*
Borealia, A Scientia, Tromsø, 17, pp:1–24

Carlsson, J. E. L., Egan, D., Collins, P. C., Farrell, E. D., Igoe, F. and Carlsson, J. (2017) 'A qPCR
MGB probe based eDNA assay for European freshwater pearl mussel (*Margaritifera margaritifera*
L.)', *Aquatic Conservation: Marine and Freshwater Ecosystems*, 27(6), pp:1341–1344. doi:
10.1002/aqc.2788

Davison, P. I., Creach, V., Liang, W.-J., Andreou, D., Britton, J. R. and Copp, G. H. (2016)
'Laboratory and field validation of a simple method for detecting four species of non-native
freshwater fish using eDNA.', *Journal of Fish Biology*, 89(3), pp. 1782–1793. doi: 10.1111/jfb.13086.

Doi, H., Uchii, K., Takahara, T., Matsuhashi, S., Yamanaka, H. and Minamoto, T. (2015) 'Use of
Droplet Digital PCR for Estimation of Fish Abundance and Biomass in Environmental DNA
Surveys', *Plos One*, 10(3). doi: 10.1371/journal.pone.0122763.

Ficetola, G. F., Miaud, C., Pompanon, F. and Taberlet, P. (2008) 'Species detection using
environmental DNA from water samples', *Biology letters*, 4(4), pp. 423–425. doi:

10.1098/rsbl.2008.0118.

Goldberg, C. S., Sepulveda, A., Ray, A., Baumgardt, J. and Waits, L. P. (2013) 'Environmental DNA as a new method for early detection of New Zealand mudsnails (*Potamopyrgus antipodarum*)', *Freshwater Science*, 32(3), pp. 792–800. doi: 10.1899/13-046.1.

Goldberg, C. S., Turner, C. R., Deiner, K., Klymus, K. E., Thomsen, P. F., Murphy, M. A., Spear, S. F., McKee, A., Oyler-McCance, S. J., Cornman, R. S., Laramie, M. B., Mahon, A. R., Lance, R. F., Pilliod, D. S., Strickler, K. M., Waits, L. P., Fremier, A. K., Takahara, T., Herder, J. E. and Taberlet, P. (2016) 'Critical considerations for the application of environmental DNA methods to detect aquatic species', *Methods in Ecology and Evolution*, 7(11), pp. 1299–1307. doi: 10.1111/2041-210X.12595.

Gordeeva, N. V. and Salmenkova, E. A. (2011) 'Experimental microevolution: Transplantation of pink salmon into the European North', *Evolutionary Ecology*, 25(3), pp. 657–679. doi: 10.1007/s10682-011-9466-x.

Gustavson, M. S., Collins, P. C., Finarelli, J. A., Egan, D., Conchúir, R., Wightman, G. D., King, J., Gauthier, D., Whelan, K., Carlsson, J., and Carlsson, J. (2015) 'An eDNA assay for Irish *Petromyzon marinus* and *Salmo trutta* and field validation in running water', *Journal of Fish Biology*, 87(5), pp:1254–1262. https://doi.org/10.1111/jfb.12781

Hänfling, B., Handley, L. L., Read, D. S., Hahn, C. and Li, J. (2016) 'Environmental DNA metabarcoding of lake fish communities reflects long-term data from established survey methods', *Molecular Ecology*, 25(13), 3101-3119. doi: 10.1111/mec.13660.

Harper, K. J., Anucha, N. P., Turnbull, J. F., Bean, C. W. and Leaver, M. J. (2018) 'Searching for a signal: Environmental DNA (eDNA) for the detection of invasive signal crayfish, *Pacifastacus leniusculus* (Dana, 1852)', *Management of Biological Invasions*, 9(9). doi: 10.3391/mbi.2018.9.2.07

Hindson, B. J., Ness, K. D., Masquelier, D. a., Belgrader, P., Heredia, N. J., Makarewicz, A. J., Bright, I. J., Lucero, M. Y., Hiddessen, A. L., Legler, T. C., Kitano, T. K., Hodel, M. R., Petersen, J. F., Wyatt, P. W., Steenblock, E. R., Shah, P. H., Bousse, L. J., Troup, C. B., Mellen, J. C., Wittmann, D. K., Erndt, N. G., Cauley, T. H., Koehler, R. T., So, A. P., Dube, S., Rose, K. A., Montesclaros, L., Wang, S., Stumbo, D. P., Hodges, S. P., Romine, S., Milanovich, F.P., White, H. E., Regan, J. F., Karlin-Neumann, G. A., Hindson, C. M., Saxonov, S. and Colston, B. W. (2011) 'High-throughput droplet digital PCR system for absolute quantitation of DNA copy number', *Analytical Chemistry*, 83(22), pp. 8604–8610. doi: 10.1021/ac202028g.

Heard, W. R. (1991) 'Life history of pink salmon (*Oncorhynchus gorbuscha*)', In: Groot C, Margolis L (eds) *Pacific salmon life histories*. UBC Press, Vancouver, pp:121–123

Hunter, M. E., Dorazio, R. M., Butterfield, J. S. S., Meigs-Friend, G., Nico, L. G. and Ferrante, J. A. (2017) 'Detection limits of quantitative and digital PCR assays and their influence in presence–absence surveys of environmental DNA', *Molecular Ecology Resources*, 17(2), pp. 221–229. doi: 10.1111/1755-0998.12619.

Jerde, C. L., Mahon, A. R., Chadderton, W. L. and Lodge, D. M. (2011) '"Sight-unseen" detection of rare aquatic species using environmental DNA', *Conservation Letters*, 4(2), pp. 150–157. doi: 10.1111/j.1755-263X.2010.00158.x.

Klymus, K. E., Richter, C. A., Chapman, D. C. and Paukert, C. (2015) 'Quantification of eDNA shedding rates from invasive bighead carp *Hypophthalmichthys nobilis* and silver carp *Hypophthalmichthys molitrix*', *Biological Conservation*, 183, pp. 77–84. doi: 10.1016/j.biocon.2014.11.020.

Millane, M., Walsh, L., Roche, W. K., and Gargan, P. G. (2019) 'Unprecedented widespread occurrence of Pink Salmon *Oncorhynchus gorbuscha* in Ireland in 2017', *Journal of Fish Biology*. doi:10.1111/jfb.13994

Mo, T. A., Thorstad, E. B., Sandlund, O. T., Berntsen, H. H., Fiske, P., and Uglem, I. (2018) 'The pink salmon invasion: a Norwegian perspective', *Journal of Fish Biology*, 93, pp. 5–7. doi:10.1111/jfb.13682

Moore, J. W., Gordon, J., Carr-Harris, C., Gottesfeld, A. S., Wilson, S. M. and Russell, J. H. (2016) 'Assessing estuaries as stopover habitats for juvenile Pacific salmon', *Marine Ecology Progress Series*, 559, pp:201–2015. doi:10.3354/meps11933

Nakano, M., Komatsu, J., Matsuura, S. I., Takashima, K., Katsura, S. and Mizuno, A. (2003) 'Single-molecule PCR using water-in-oil emulsion', *Journal of Biotechnology*, 102(2), pp. 117–124. doi: 10.1016/S0168-1656(03)00023-3.

Nathan, L. M., Simmons, M., Wegleitner, B. J., Jerde, C. L. and Mahon, A. R. (2014) 'Quantifying Environmental DNA Signals for Aquatic Invasive Species Across Multiple Detection Platforms', *Environmental Science & Technology*, 48(21), pp. 12800–12806. doi: 10.1021/es5034052.

Penãrrubia, L., Alcaraz, C., Vaate, A. B. D., Sanz, N., Pla, C., Vidal, O., and Vinãs, J. (2016) 'Validated methodology for quantifying infestation levels of dreissenid mussels in environmental DNA (eDNA) samples', *Scientific Reports*, 6. https://doi.org/10.1038/srep3906

Pilliod, D. S., Goldberg, C. S., Arkle, R. S., Waits, L. P. and Richardson, J. (2013) 'Estimating occupancy and abundance of stream amphibians using environmental DNA from filtered water samples', *Canadian Journal of Fisheries and Aquatic Sciences*, 70(8), pp. 1123–1130. doi: 10.1139/cjfas-2013-0047.

Pinheiro, L. B., Coleman, V. A., Hindson, C. M., Herrmann, J., Hindson, B. J., Bhat, S. and Emslie, K. R. (2012) 'Evaluation of a droplet digital polymerase chain reaction format for DNA copy number quantification', *Analytical Chemistry*, 84(2), pp. 1003–1011. doi: 10.1021/ac202578x.

Rasputina, E. N., Shustov, Y. A. and Tyrkin, I. A. (2016) 'Eggs of pink salmon *Oncorhynchus gorbuscha* as additional nontraditional food of juvenile Atlantic salmon *Salmo salar* in rivers of the Kola Peninsula', *Russian Journal of Biological Invasions*, 7, pp: 294–296.

Rusch, J. C., Hansen, H., Strand, D. A., Markussen, T., Hytterød, S., and Vrålstad, T. (2018) 'Catching the fish with the worm: A case study on eDNA detection of the monogenean parasite *Gyrodactylus salaris* and two of its hosts, Atlantic salmon (*Salmo salar*) and rainbow trout (*Oncorhynchus mykiss*)', *Parasites and Vectors*, 11(1). https://doi.org/10.1186/s13071-018-2916-3

Sandlund, T., Berntsen, H. H., Fiske, P., Kuusela J., Muladal, R., Niemelä, E., Uglem, I., Forseth, T., Mo, T. A., Thorstad, E., Veselov, A. E., Vollset, K. W. and Zubchenko, A. V. (2018) 'Pink salmon in Norway: the reluctant invader', *Biological Invasions*, doi: 10.1007/s10530-018-1904-z.

Saltveit, S. J., Brittain, J. E. and Bremnes, T. (2013) 'The return of Atlantic Salmon (*Salmo salar* L.) and improved water quality in urban rivers in Oslo, Norway', *River Res. Applic*. 30, pp:571–577. doi: 10.1002/rra.2670.

Spear, S. F., Groves, J. D., Williams, L. a and Waits, L. P. (2014) 'Using environmental DNA methods to improve detectability in a hellbender (*Cryptobranchus alleganiensis*) monitoring program', *Biological Conservation*, 183, pp. 38-45. doi: 10.1016/j.biocon.2014.11.016.

Taberlet, P. Coissac, E., Hajibabaei, M. and Rieseberg, L. (2012) 'Environmental DNA', *Molecular Ecology*, 21(8), pp. 1789–1793. doi: 10.1111/j.1365-294X.2012.05542.x.

Takahara, T., Minamoto, T. and Doi, H. (2013) 'Using Environmental DNA to Estimate the Distribution of an Invasive Fish Species in Ponds', *PLoS ONE*, 8(2). doi: 10.1371/journal.pone.0056584.

Thomsen, P. F., Kielgast, J., Iversen, L. L., Wiuf, C., Rasmussen, M., Gilbert, M. T. P., Orlando, L.

and Willerslev, E. (2012) 'Monitoring endangered freshwater biodiversity using environmental
DNA', *Molecular Ecology*, 21(11), pp. 2565–2573. doi: 10.1111/j.1365-294X.2011.05418.x.

Tréguier, A., Paillisson, J. M., Dejean, T., Valentini, A., Schlaepfer, M. A. and Roussel, J. M. (2014)
'Environmental DNA surveillance for invertebrate species: Advantages and technical limitations to
detect invasive crayfish Procambarus clarkii in freshwater ponds', *Journal of Applied Ecology*, 51(4),
pp. 871–879. doi: 10.1111/1365-2664.12262

Wacker, S., Fossøy, F., Larsen, B. M., Brandsegg, H., Sivertsgård, R., Karlsson, S. (2019)
'Downstream transport and seasonal variation in freshwater pearl mussel (*Margaritifera*
margaritifera) eDNA concentration', *Environmental DNA*, doi: 10.1002/edn3.10.

Whelan K (2017) 'Pink invaders', *Off the Scale*, issue 18, Sept– Oct 2017, pp. 14–21.
http://www.offthescaleangling.ie/issue18/offthescale_issue18.pdf. Accessed 22 Mar 2019

Wilcox, T. M., McKelvey, K. S., Young, M. K., Jane, S. F., Lowe, W. H., Whiteley, A. R. and
Schwartz, M. K. (2013) 'Robust Detection of Rare Species Using Environmental DNA: The
Importance of Primer Specificity', *PLoS ONE*, 8(3). doi: 10.1371/journal.pone.0059520.

TABLES AND FIGURES

Figure 1: Life cycle of the pink salmon *Oncorhynchus gorbuscha, ill.: Sigrid Skoglund, NINA.*

Figure 2: Map showing the sampling area in Norway, with water sampling locations for detection of
O. gorbuscha along the Lysakerelva river and in the estuary indicated by orange dots. Barriers are
indicated by blue bars.

Table 1: Details of water sampling in Lysakerelva river.

Date	Locality	No. samples	Water volume	Filter size	Latitude, Longitude
04.09.2017	Granfoss below waterfall	2	1	0.45µm	59.917894, 10.631348
04.09.2017	Møllefoss below waterfall	2	1	0.45µm	59.914629, 10.636455
31.05.2018	Granfoss below waterfall	2	0.8	0.45µm	59.917894, 10.631348
20.06.2018	Estuary	2	10	2.0µm	59.911005, 10.643129
20.06.2018	Møllefoss below waterfall	2	10	2.0µm	59.914629, 10.636455
22.06.2018	Granfoss above waterfall	2	10	2.0µm	59.918166, 10.630236
22.06.2018	Granfoss above waterfall	2	0.8	0.45µm	59.918166, 10.630236

441 **Table 2:** Details of the assay that was designed, tested and deployed for detection of a 98bp region of
442 the mitochondrial *COI* gene of *O. gorbuscha* in this study.

Primer Type	Name	Sequence 5' -3'	Length (bp)
Forward Primer	*PinkF*	CACCGCCCTAAGCCTACTAA	20
Reverse Primer	*PinkR*	AGGCATGGGCTGTAACGATT	20
Probe *	*PinkPr*	CGCTCTTCTAGGGAATGACCA	21

443 * 5' VIC labelled reporter dye and 3' non-fluorescent quencher

445 **Table 3:** Detection of *O. gorbuscha* using eDNA shown as the number of samples that were positive
446 for detection out of the total number of samples analysed by ddPCR.

Locality	Date	eDNA detection
Granfoss below waterfall	04.09.2017	1/2
Granfoss below waterfall	31.05.2018	2/2
Granfoss above waterfall	22.06.2018	1/4
Møllefoss below waterfall	04.09.2017	2/2
Møllefoss below waterfall	20.06.2018	0/2
Estuary	20.06.2018	0/2

450 **Figure 3:** Boxplot showing the number of positive droplets from ddPCR detection of *O. gorbuscha* in
451 relation to date and locality. The horizontal dashed line indicates the lower threshold of three droplets
452 for assessing a sample as positive. See Table 1 for details of each sample.

17