Dissecting the eQTL micro-architecture in *Caenorhabditis elegans*

Mark G. Sterken[*,1], Roel P. J. Bevers[*], Rita. J. M. Volkers[*], Joost A. G. Riksen[*], Jan E. Kammenga[*], L. Basten Snoek[*,†,1]

[*]Laboratory of Nematology, Wageningen University, Droevendaalsesteeg 1, 6708 PB Wageningen, The Netherlands

[†]Theoretical Biology and Bioinformatics, Utrecht University, Padualaan 8, 3584 CH Utrecht, The Netherlands

[1]Corresponding authors

Running title: *C. elegans* eQTL micro-architecture

Keywords: genetic architecture, eQTL, *Caenorhabditis elegans*, introgression line,

recombinant inbred line

Corresponding authors:

Mark Sterken

Laboratory of Nematology, Wageningen University and Research

Postbus 8123

6700ES, Wageningen

+31 317 482 998

mark.sterken@wur.nl

Basten Snoek

Theoretical Biology and Bioinformatics, Utrecht University

Padualaan 8

3584CH, Utrecht

+31 30 253 0000

l.b.snoek@uu.nl

Abstract

The study of expression quantitative trait loci (eQTL) using natural variation in inbred populations has yielded detailed information about the transcriptional regulation of complex traits. Studies on eQTL using recombinant inbred lines (RILs) led to insights on local and distant regulatory loci of transcript abundance. However, determining the underlying causal polymorphic genes or variants is difficult, but ultimately essential for the understanding of regulatory networks of complex traits. This requires insight into whether associated loci are single eQTL or a combination of closely linked eQTL, and how this QTL micro-architecture depends on the environment. We addressed these questions by mapping eQTL in N2 x CB4856 *C. elegans* RIL populations across three different environments (control, heat-stress, and recovery). To test for independent replication of the RIL eQTL, we used introgression lines (ILs). Both populations indicate that the overall heritability, number, and position of eQTL differed among environments. Across environments we were able to replicate 70% of the local- and 40% of the distant-eQTL using the ILs. Simulation models revealed that additive effects explain up to 60-93% of RIL/IL heritability across environments. Closely linked eQTL explained up to 40% of RIL/IL heritability in the control environment whereas only 7% in the heat-stress and recovery environments. In conclusion, we show that reproducibility of eQTL was higher for local vs. distant eQTL and that the environment affects the eQTL micro-architecture.

Introduction

The genetic architecture of quantitative traits in genetically segregated populations such as recombinant inbred lines (RILs) differs within and across species and strongly depends on the environment and type of trait (FOURNIER AND SCHACHERER 2017). In some cases, quantitative traits are (nearly) monogenic and display Mendelian characteristics. For instance the Kallman syndrome in humans is caused by a knockout of FGFR1 (which encodes fibroblast growth factor receptor 1) (MUENKE *et al.* 1994), industrial melanism in British peppered moths is caused by a single gene mutation (VAN'T HOF *et al.* 2011), and natural variation in the *npr-1* gene in the nematode *Caenorhabditis elegans* is a major determinant of multiple life-history traits (ANDERSEN *et al.* 2014; STERKEN *et al.* 2015). These monogenic traits are relatively easy to detect but rather exceptions than the rule. Most quantitative traits display a complex polygenic genetic architecture, regulated by multiple QTLs, where each locus explains a proportion of the trait variation (ALBERT AND KRUGLYAK 2015).

Studying quantitative genetics of the transcriptome has proven useful in understanding the architecture of complex traits. Expression quantitative trait loci (eQTL) can be measured reliably in a high-throughput fashion in many organisms like: yeast, *Arabidopsis thaliana*, tomato, maize, *C. elegans*, mice, and humans (SCHADT *et al.* 2003; BREM AND KRUGLYAK 2005; LI *et al.* 2006; KEURENTJES *et al.* 2007; RANJAN *et al.* 2016). The advantage of the eQTL-approach lies in the assessment of thousands of traits simultaneously for which the genetic architectures can be compared (JANSEN AND NAP 2001; GILAD *et al.* 2008). First, the variance explained per underlying QTL will cover the whole range from essentially monogenic (Mendelian) to nearly undetectable. Second, correlating gene expression to loci can reveal the biological processes and phenotypes that lay downstream of (e)QTL (*e.g.* (JIMENEZ-GOMEZ *et al.* 2010; TERPSTRA *et al.* 2010; JIMENEZ-GOMEZ *et al.* 2011; ANDERSEN *et al.* 2014; SCHMID *et al.* 2015; STERKEN *et al.* 2017; ALBERT *et al.* 2018)). In general, two

patterns stand out in eQTL studies: i) the occurrence of regulatory hotspots (*trans*-bands) where a single locus affects the expression of many genes across the genome and ii) an abundance of locally regulated eQTL (*cis*-eQTL) where the locus is located on or near the gene of which it affects the expression (NICA AND DERMITZAKIS 2013; ALBERT *et al.* 2018). *Cis*-eQTL can occur due to polymorphisms affecting the transcription of the gene under study, *e.g.* deletions or non-functional promotor regions. The occurrence of *trans*-bands was found to be specific for environmental conditions linked to genetic effects. Together, the *cis*-eQTL and *trans*-bands typically cover ~75% of the mapped eQTL in any study (e.g. (SNOEK *et al.* 2017; ALBERT *et al.* 2018)).

The detection of (e)QTL depends on many different factors like developmental stage, background mutations, and the ambient environment, to name a few (LI *et al.* 2006; VINUELA *et al.* 2010; DUVEAU AND FELIX 2012; SNOEK *et al.* 2012; ANDERSEN *et al.* 2014; FRANCESCONI AND LEHNER 2014). Importantly, QTL detection is mainly determined by the number of genetically segregating RILs and the density and structure of the genetic marker map. Further characterization of QTLs often involves the identification of the underlying causal polymorphic genes in order to understand molecular mechanisms affecting complex traits and/or to facilitate breeding for selected traits. An important hallmark of QTL mapping has been the fine mapping and identification of polymorphic regulators or quantitative trait nucleotides (QTN) (RADWAN AND BABIK 2012; ROCKMAN 2012).

Fine mapping of causal variants requires determining if associated loci are single QTL or combinations of closely linked QTLs, here coined the QTL micro-architecture, and how this micro-architecture depends on the environment. Apparent single QTLs may be truly single QTL or a combination of closely linked QTLs not segregating in the studied population. These are often hard to characterize because of a lack of resolution of the genetic map or shortage of a sufficient number of recombination events. Here, we set out to

investigate the QTL micro-architecture of eQTLs in the nematode *C. elegans* in a segregated population of RILs derived from a cross between wild-types Bristol N2 and CB4856 (LI *et al.* 2006; THOMPSON *et al.* 2015). We use a set of previously detected eQTL for their replication in a separate, independent, population of introgression lines (ILs) in this study (DOROSZUK *et al.* 2009; THOMPSON *et al.* 2015; SNOEK *et al.* 2017). Compared to the RILs, which are genetic mosaics of loci derived from both parents, ILs contain a single genomic segment of one parent in a genetic background of another parent. Typically, introgression lines are used to verify the existence of QTL, or to narrow down the location of a QTL in order to find the underlying causal polymorphism (*e.g.* see (GAO *et al.* 2018)). The RIL and IL populations are exposed to three environments: heat-shock, recovery from heat-shock, and a control environment at a standard rearing temperature. We investigate three aspects of eQTL mapped in RILs: i) we address how well eQTL are replicated in ILs, ii) we test if the predicted numbers of eQTL agree with the observed number of differentially expressed genes in ILs, and iii) we simulate QTL architectures explaining the relative heritability in RILs and ILs and match our data to these models. Finally, we show that the eQTL microarchitecture under ambient conditions consists of closely linked eQTL.

Material and methods

Strains used

The wild-type strains N2 and CB4856 were used and 57 introgression lines (ILs) with segments of CB4856 in an N2 genetic background (DOROSZUK *et al.* 2009). Most of the ILs have been sequenced, and the genetic map was based on the sequenced genotypes (THOMPSON *et al.* 2015), a file with the strains and the map has been included in **Table S1**. The data used for the recombinant inbred lines (RILs) is accessible in (SNOEK *et al.* 2017).

Nematode culturing

The strains were cultured as described previously (SNOEK *et al.* 2017). In short, strains were kept in maintenance at 12 °C on 6-cm Nematode Growth Medium (NGM) Petri dishes seeded with the *Escherichia coli* strain OP50 as food source (BRENNER 1974). Before starting the experiment, single hermaphrodites of each strain in the L2 stage were sub-cultured in 12-wells plates and grown at 20 °C. The offspring was screened for the occurrence of males by microscopy and only populations consisting solely of hermaphrodites were transferred to 9-cm NGM Petri dishes containing *E. coli* OP50 and grown until the bacterial food was depleted.

Treatments for the transcriptomic experiment

The experiment described in our previous paper was repeated on the IL population and the parental strains (SNOEK *et al.* 2017). We started the experiment by transferring a starved population to a new 9-cm NGM Petri dish seeded with *E. coli* OP50. After 60 h at 20 °C, the populations consisting of egg-laying adults were bleached to obtain the eggs, which were transferred to a new 9-cm NGM Petri dish (BRENNER 1974). Around 46 h the developmental stage of the population was assessed by microscopy, verifying that the population consisted

mainly of L4 larvae. Populations not consisting of L4 larvae were not used. The strains were

exposed to one of three environments: (i) a control environment; grown for 48 h at 20 °C, (ii)

a heat-stress environment; grown for 46 h at 20 °C and subsequently exposed to 35 °C for 2 h,

and (iii) a recovery environment; grown for 46 h at 20 °C, exposed to 35 °C for 2 h, and

thereafter returned to 20 °C for 2 h. Directly after the treatment the animals were washed off

the Petri dish using M9 buffer and collected in a 1.5 mL Eppendorf tube, which was

centrifuged and the pellet was flash frozen in liquid nitrogen. After quality control of the

RNA isolation, we obtained 56 ILs for the control and heat-stress environment, and 55 ILs for

the recovery environment. For the parental strain N2: four control samples, four heat stress

samples and three recovery samples passed quality control. For the parental strain CB4856:

three control samples, three heat stress samples and five recovery samples passed quality

control.

RNA isolation

The procedure was followed as described before (SNOEK *et al.* 2017). The RNA of the

samples was isolated using the RNEasy Micro Kit from Qiagen (Hilden, Germany) following

the 'Purification of Total RNA from Animal and Human Tissues'. The lysis step was

modified, pellets were lysed in 150 µl RLT buffer, 295 µl RNAse-free water, 800 µg/ml

proteinase K and 1% ß-mercaptoethanol, which was incubated at 55 °C and 1000 rpm in a

Thermomixer (Eppendorf, Hamburg, Germany) for 30 minutes, where after the

manufacturer's protocol was followed.

cDNA synthesis, labelling and hybridization

Before starting cDNA synthesis, the quality and quantity of the RNA was measured by

NanoDrop-1000 spectrophotometer (Thermo Scientific, Wilmington DE, USA). The integrity

of the RNA was assessed by loading 3 μL per sample on a 1% agarose gel. Samples with

good quality RNA were processed following the 'Two-Color Microarray-Based Gene

Expression Analysis; Low Input Quick Amp Labeling' -protocol, version 6.0 from Agilent

(Agilent Technologies, Santa Clara, CA, USA). The microarrays used were Agilent *C.*

elegans (V2) Gene Expression Microarray 4X44K slides.

Array scanning, data extraction, and normalization

After washing, the microarrays were scanned using an Agilent High Resolution C Scanner

following the recommended settings. For data extraction, Agilent Feature Extraction Software

(version 10.7.1.1) was used, following manufacturer's guidelines. The extracted data was

normalized using the limma package in "R" (version 3.4.2, x64) (RITCHIE *et al.* 2015). As

recommended, the data was not background corrected before normalization (ZAHURAK *et al.*

2007), the within-array normalization used the 'Loess' method and between-array

normalization used the Quantile method (SMYTH AND SPEED 2003).

Data analysis

Data was analyzed using "R" (version 3.4.2, x64) with custom written scripts (TEAM 2017),

accessible via https://git.wur.nl/published_papers/sterken_2019_closely_linked_qtl. For

analysis, the dplyr and tidyr packages were used for data organization (WICKHAM 2018a;

WICKHAM 2018b), and plots were generated using ggplot2 (WICKHAM 2009), except for plots

displaying simulation results. The transcriptome data analyzed was deposited at ArrayExpress

under E-MTAB-5779 (RIL data) (SNOEK *et al.* 2017), and E-MTAB-7424 (IL and parental

data, described in this paper).

eQTL data

The eQTL data from the RILs were obtained from a previous publication (SNOEK *et al.* 2017)

and can be explored at www.bioinformatics.nl/wormqtl2 (SNOEK *et al.* 2019). In short, eQTL

were mapped using a single marker model, which was conducted separately for each

environment. The threshold for each of the three environments was set at $-\log_{10}(p) > 3.9$

(false discovery rate, FDR = 0.05) (see (SNOEK *et al.* 2017) for details).

Data transformation

The expression data was transformed to a z-score based on the standard deviation and mean of

the N2-replicates per environment, using

$$Z_{N2,i,j} = \frac{I_{i,j} - \mu_{i,N2}}{\sigma_{i,N2}}$$

where Z_{N2} is the Z-score of spot *i* (1, 2, ..., 45220) of strain *j* (one of the RILs or ILs) and *I* is

the normalized intensity. The μ is the mean of the intensity over the N2 strains for spot *i* and

the σ is the standard deviation over the N2 strains for spot *i*. This transformation shows the

effect of the CB4856 introgressions in the ILs.

Additionally, the expression data was transformed to a z-score based on the standard

deviation and mean of the CB4856-replicates per environment, using

$$Z_{CB4856,i,j} = \frac{I_{i,j} - \mu_{i,CB4856}}{\sigma_{i,CB4856}}$$

where Z_{CB4856} is the Z-score of spot *i* (1, 2, ..., 45220) of strain *j* (one of the RILs or ILs) and *I*

is the normalized intensity. The μ is the mean of the intensity over the CB4856 strains for spot

i and the σ is the standard deviation over the CB4856 strains for spot *i*. This transformation

shows the effect of the genetic background in the ILs.

In order to compare IL gene expression with eQTL effect sizes, another data

transformation was used, as there the number of standard deviations difference with the mean

is less informative than the effect in log$_2$-ratio with the mean. Hence, we transformed the data

by

$$R_{N2,i,j} = \log_2 \frac{I_{i,j}}{\mu_{i,N2}}$$

where R_{N2} is the log$_2$-ratio with the mean of spot i of strain j, I is the normalized intensity, and

μ is the mean intensity over the N2 samples.

Confirmation of IL genotype

Before mapping, we assured that samples were labelled correctly by applying a *cis*-eQTL

based analysis of gene expression compared to the genetic map of the IL population, for

details and explanation, see (ZYCH *et al.* 2017). Samples of three strains were removed

(WN263, WN264, and WN284), as the strains did not match the genotypes in the IL

population (DOROSZUK *et al.* 2009; THOMPSON *et al.* 2015).

Confirmation of eQTL in introgression lines

The differential expression per IL was calculated based on the Z-score calculated using the

N2 parental strains (Z_{N2}). As each IL was measured once, we estimated the significance

assuming a normal distribution based on the genetic-background parent. These significances

were corrected for multiple testing by the Benjamini-Hochberg method, as applied by the

p.adjust function in R (BENJAMINI AND HOCHBERG 1995). As a control and benchmark, the

same procedure was applied to the individual RILs. The number of differentially expressed

genes with an expected eQTL per CB4856 locus were counted for both the ILs and the RILs.

These were expressed as a percentage of the expected eQTL in that locus (separately for *cis*-

and *trans*-eQTL).

Another method of confirming eQTL is by correlating the differential expression per

inbred line compared to the genetic background-parent (R_{N2}) with the QTL effect. By

calculating the Pearson correlation of R_{N2} with the eQTL effect per IL or RIL separately for

the *cis-* and *trans-*eQTL, it can be assessed how well the overall QTL patterns were

recapitulated per strain. These correlations were calculated using the *cor* function and the

significances of correlation were calculated using the *cor.test* function.

Confirmation of trans-bands using ILs

In the eQTL study on the RILs (SNOEK *et al.* 2017), 19 *trans-*bands were identified (**Table**

S3). For each *trans-*band 2 to 6 ILs covering or flanking the locus were tested by correlating

the *trans-*eQTL effect with the R_{N2} in the ILs. Significance was determined based on the

correlation values in the N2 samples. Only if the correlation was stronger than the highest-

correlating N2 sample, it was called as significant. Only significant positive correlations

(effect direction matches the eQTL model) were scored as confirming the *trans-*band.

To determine the overall false-positive rate of this analysis, we determined the

correlation of the ILs that were not in or near the *trans-*band region. These correlations were

compared to the correlation value at which the *trans-*band was called confirmed. This resulted

in an overall false positive rate of 0.11. It should be noted that this approach assumes that

*trans-*bands stem from a single location and that these locations were accurately pinpointed in

the RIL mapping. Hence, the derived threshold can be considered strict.

Expectation of differential expressed genes based on eQTL

First, the number of differentially expressed genes in each RIL and IL was calculated using

the contrast with either of the ancestral strains, Z_{N2} and Z_{CB4856}, at an estimated significance of

$p < 1*10^{-5}$. Second, the expected number of differentially expressed genes was estimated for

each RIL and IL based on the eQTL architecture in the RIL population (eQTL threshold, -

$\log_{10}(p) > 3.9$; FDR = 0.05) (SNOEK *et al.* 2017). These values were normalized to the RIL

population (the RIL population average was set to 1), to compare the relative number of

differentially expressed genes expected versus observed in the ILs.

Differential gene-expression in N2 and CB4856

Differential gene expression between the two parental strains was independently calculated

for each environment using a linear model

$$\log_2(I_{i,j}) \sim G_j + e_i$$

where *I* is the normalized intensity of spot *i* (1, 2, ..., 45220) of strain *j* (CB4956 or N2) is

explained over genotype *G* and residual variance *e*. For the parental strain N2: n = 4 in control

and heat-stress environments, n = 3 in recovery environment. For the parental strain CB4856:

n = 3 in control and heat stress environments and n = 5 recovery in recovery environment.

The obtained significances were corrected for multiple testing using the Benjamini-Hochberg

method implemented in the *p.adjust* function in R (BENJAMINI AND HOCHBERG 1995). A

threshold of FDR = 0.1 was taken as requirement for differential expression.

Estimation of trait genetic architecture

Two approaches were taken to determine trait architectures using both RIL and IL population:

heritability calculations and directly testing the amount of variance between both populations.

Both approaches are based on the notion that the trait variance in the populations scale with

trait complexity, where

$$\sigma^2_{trait} \propto \sigma^2_{QTL\ 1} + \sigma^2_{QTL\ 2} + \cdots + \sigma^2_{QTL\ n} + \sigma^2_e$$

here σ^2 is the trait variance, proportional to the variance captured by *n* QTL (which can be

multiple loci interacting; epistasis) and the measurement error. The contrast between the RIL

and the IL population is that the QTL segregating in the IL population will affect fewer strains

per QTL, whereas in the RIL population each QTL will segregate on average in 50% of the

strains. In other words, the IL population shares most QTL effects because the genetic

background is the same, whereas in the RIL population most QTL segregate.

Heritability was calculated as in (BREM AND KRUGLYAK 2005; KEURENTJES *et al.*

2007), where

$$H_{i,F}^2 = \frac{\sigma_{i,F}^2 - \sigma_{i,P}^2}{\sigma_{i,F}^2}$$

where H^2 was the heritability of spot i (1, 2, ..., 45220) of population F (either RIL or IL)

within one of the three environments. The σ^2 indicated the trait variance of spot i. Here the

pooled trait variance of the parental lines N2 and CB4856 (denoted with P) was used as an

estimate of the measurement error, which was subtracted from the trait variance of either the

RIL or IL population. We used a permutation approach to determine the threshold for

significant heritability (BREM AND KRUGLYAK 2005; VINUELA *et al.* 2012). Thereto, the gene

expression values per spot per environment were randomized over the strains (RIL and IL

separately, but together with the same parental lines) and recalculated the heritability as

described above. This was repeated 1000 times, and the 5% highest value for each spot was

taken as the FDR = 0.05 threshold. For the RILs, 4478 spots (2910 genes) displayed

significant heritability (FDR = 0.05) in control environment ($H^2 \geq 0.68$), 9133 spots (5072

genes) in heat-shock environment ($H^2 \geq 0.70$), and 13683 spots (7537 genes) in recovery

environment ($H^2 \geq 0.62$). For the ILs, 1361 spots (1012 genes) displayed significant

heritability (FDR = 0.05) in control environment ($H^2 \geq 0.70$), 1647 spots (1123 genes) in heat-

shock environment ($H^2 \geq 0.71$), and 1578 spots (1092 genes) in recovery environment ($H^2 \geq$

0.64).

The second measure was by determining the ratio of trait variance between the IL and

the RIL population. The assumption was that the variance contributed by random sources

(σ^2_e) was equal in both populations. Hence, this is a measure of relative heritability and was

calculated as

$$R_{H,i} = \log_2(\frac{\sigma_{i,IL}^2}{\sigma_{i,RIL}^2})$$

where R_H was the heritability ratio of spot i (1, 2, ..., 45220), and σ^2 was the trait variance in

either the IL or the RIL population within one of the tree environments. We took the $\log_2$

ratio, meaning that traits with a higher heritability in the IL population resulted in a positive

number, and traits with a higher heritability in the RIL population with a negative number.

We performed two tests to place the relative heritability in perspective. First, we took the

spots showing significant heritability (FDR = 0.05). Second, to test whether variance was

significantly different between the two populations, we used the non-parametric Fligner-

Killeen test for homogeneity of variances, this we corrected for multiple testing using the

Benjamini-Hochberg method (FDR = 0.05) (CONOVER *et al.* 1981; BENJAMINI AND

HOCHBERG 1995).

Simulating trait architecture impact on relative heritability

To determine how the trait variance behaves proportionally in each population, we simulated

different trait architectures in both the IL and RIL population. First, all QTL effect sizes were

simulated by random drawing from a standard normal distribution (μ = 0, σ = 1). These QTL

were simulated in trait architectures containing 1 to 100 QTL (or 1 to 100 clusters of closely

linked QTL, each 33331 times. To recapitulate the observed QTL distribution in *C. elegans*,

we distributed these QTL along the informative markers in our IL and RIL population. This

assured an enrichment of QTL on the chromosome arms compared to the chromosome tips

and centers (ROCKMAN *et al.* 2010; SNOEK *et al.* 2017). To approach the residual variance

(σ_e^2) in our data, we added random variation based on a normal distribution (*rnorm*) such that

the average heritability of each simulated trait was 0.85

The following eight architectures were simulated: (i) randomly distributed additive

QTL, (ii) randomly distributed additive QTL that only show their effect in one genotype, (iii)

clusters of closely linked QTL with random effects, (iv) clusters of closely linked QTL that

only show their effect in one genotype, (v) clusters of closely linked QTL that are balanced

within the ancestral genotypes, (vi) clusters of closely linked QTL that are balanced within the

ancestral genotypes and only show their effect in one genotype, (vii) randomly distributed

interacting QTL, (viii) clusters of closely linked interacting QTL. The number of closely

linked QTL per cluster was simulated from two to five. The additive and closely-linked model

were similar in essence, but differ in the architecture, where we modelled the closely-linked

QTL such that these were typically not segregated in the population used.

The distribution of traits over a balanced closely-linked architecture and additive

architecture was done by measuring the number of heritable traits with a heritability ratio

between -3 and -2 (peak) and between 0.5 and 1.5 (intersect). These intervals were chosen as

they represent areas where the distributions of these models differ. The number of expected

traits per interval were counted as the median of all simulated architectures (so 1 to 100 QTL,

and 2 to 5 QTL per closely-linked locus) in order to get an estimation that is less dependent

on the chosen simulation parameters. The predicted number of traits at the peak and the

intersect for the additive model (a) and the balanced closely-linked model (b) were used to

solve the true distribution based on the real number of heritable traits in these heritability ratio

intervals (c). This was done by solving the equations

350
$$a_{peak} * x_1 + b_{peak} * x_2 = c_{peak} \text{ and}$$

$$a_{intercept} * x_1 + b_{intercept} * x_2 = c_{intercept}$$

351   for x_1 and x_2 using the *solve* function in R. This led to the estimated number of additive traits

352   (x_1) and balanced closely linked traits (x_2).

Results

Genome-wide confirmation of eQTL effects and location using introgression lines

Analysis of differential expression showed that introgression lines (ILs) could confirm both *cis-* and *trans*-eQTL mapped in the recombinant inbred line (RIL) population (SNOEK *et al.* 2017). A population of 48 RILs, 56 ILs ((DOROSZUK *et al.* 2009); **Table S1**), and at least three replicates of the ancestral strains N2 and CB4856 were exposed to the three environments (**Figure 1A**). Based on the number of eQTL expected in the CB4856 locus of each IL, we found that on average 69.5% (FDR = 0.05) of the *cis*-eQTL found using the RILs could be confirmed using the ILs over all three environments (FDR = 0.05). The *trans*-eQTL found in the RILs, were confirmed on average 42.2% (FDR = 0.05) in the ILs (**Figure 1A and 1B; Table S2**). Note that the tiling structure of the ILs meant that each QTL location was covered by 0-5 (5-95% quantiles) ILs (**Figure S1**). The percentage of confirmed eQTL was positively correlated with the RIL significance threshold (**Figure S2A**). Furthermore, correlation analysis of the $\log_2$ ratio with the N2 ancestor in the ILs versus the eQTL effect measured by RILs showed that eQTL effects in the CB4856 locus of each strain were well approximated in the ILs (average Pearson correlation, rho = 0.661; **Figure S2B**; **Table S2**). Together, over all three environments, the analyses confirmed eQTL occurrence and effect sizes in the RIL population in independent experiments using an IL population.

Figure 1: Experimental setup and eQTL confirmation (**A**) The four types of populations used in the experiment: recombinant inbred lines (RILs), the ancestral strain CB4856, the ancestral strain N2, and introgression lines (ILs). These were exposed to three separate treatments: control (48 hours at 20^{o}C), heat-stress (46 hours at 20^{o}C followed by 2 hours at 35^{o}C), and recovery (as heat-stress, followed by an additional 2 hours at 20^{o}C). After the experiment, populations of the strains were harvested and used for transcriptome measurement by microarray. (**B**) the percentage of eQTL confirmed based on differentially expressed genes per IL and per RIL (FDR = 0.05). The percentages are split-out over the three environments, and the two eQTL type (*cis* or *trans*). The eQTL were mapped using the RIL population, hence that population is expected to confirm the eQTL to a high degree. (**B**) An example of one *cis*-eQTL (for gene F26E4.1) and one *trans*-eQTL (for gene K06H6.1) in the IL population. On the x-axis the genotype at the QTL locus is plotted, and on the y-axis the expression (R$_{N2}$)

Next, we investigated the *trans*-bands identified in the RIL population more closely.

The *trans*-bands were described in a previous study (SNOEK *et al.* 2017). For each *trans*-band,

we took 2 to 6 ILs covering and flanking the locus and tested if the *trans*-eQTL effects

correlated significantly with differentially expressed genes in the ILs compared to N2. We

found supporting evidence for 14 out of 19 *trans*-bands at an overall false-positive rate of

0.11; see materials and methods. Furthermore, we narrowed down the location for some of

these *trans*-bands (**File S1**; **Table S3**). In particular the *trans*-bands in the control and heat-

stress environment were well supported (5/5 and 6/7, respectively), whereas those in the

recovery environment were less (3/7). The poor support in the recovery environment is likely

due to the few *trans*-eQTL per *trans*-band in that environment (median of 25 genes in

recovery versus 51 in control and 125 in heat stress). Altogether, we conclude that *trans*-

bands were highly replicable, to a higher extent than individual *trans*-eQTL.

Next, we asked if the number of differentially expressed genes in the IL population

could be predicted from the RIL population. Therefore we compared the gene expression in

both populations to the CB4856 ancestor to measure the effect of the N2 loci and to the N2

ancestor to measure the effect of the CB4856 loci. The effect of CB4856 introgressions in an

N2 genetic background on gene expression was negatively correlated with the expectations

based on the RIL population. In other words, in the RIL population the number of genes

differentially expressed due to the introgression was under-estimated and the number of genes

differentially expressed due to the genetic background was over-estimated (**Figure 2**). In the

ILs we found that the relative number of differentially expressed genes compared to N2 was

6.7-fold higher than expected based on the differentially expressed genes found in the RILs

(paired t-test, $p < 1*10^{-54}$). Furthermore, when compared to CB4856, the number of

differentially expressed genes in the ILs was 3.3-fold lower than expected from the RILs

(paired t-test, $p < 1*10^{-128}$). The differences between RILs and ILs were consistent when

changing both the threshold for differential expression and the threshold for eQTL-based expectations (**Figure S3**). These results show that the expectations from a single marker model are a poor predictor for differentially expressed genes in ILs. Importantly, a single marker model under-estimates the impact of small introgressions and over-estimates the impact of the genetic background.

Figure 2: Number of differentially expressed genes (DEG) expected (exp.) from eQTL mapping and observed experimentally (obs.). On the x-axis, the population type is shown and on the y-axis, the expected and observed ratio of differentially expressed genes is shown, both normalized for the expectation in the RILs (eQTL threshold, $-\log_{10}(p) > 3.9$; DEG threshold, $-\log_{10}(p) > 5$). In the left panel the effect compared to N2 is shown (effect of the CB4856 loci; introgressions in the IL panel) and in the right panel the effect compared to CB4856 is shown (effect of the N2 loci; genetic background in the IL panel).

Micro-architecture of eQTL

The distribution and differences of genetic recombination over ILs and RILs allowed studying the architecture of eQTL in *C. elegans*. We hypothesized that different underlying trait architectures would leave different trait-variance distributions in the IL and RIL population. For example, a simple trait architecture dominated by one major QTL would only lead to an effect on trait levels in a single or a few ILs, but in ~50% of the RILs. Hence, resulting in a low trait variance over the genome-wide IL population compared to the RIL population. As

heritability was a straightforward way to interpret trait variance, we estimated the broad-sense

heritability (H^2) in both populations (**Table S4**). In general, heritability was higher in the RIL

population. In total, over all conditions, we found 10673 unique genes that showed significant

heritability in the RIL population, whereas we found 2952 unique genes that showed

significant heritability in the IL population (permutation, FDR = 0.05; **Figure S4A**). Only 929

genes in the RILs and 8 genes in the ILs showed heritable gene expression variation over all

three environments (**Figure S4B** and **C**). The reason for the difference between populations

was partly due to 534 out of 929 genes having an eQTL in the RILs (of which 354 had a *cis-*

eQTL), which captured little heritable variation in the whole IL population. From the

heritability analysis, we conclude that, as expected, heritability was higher in the RIL

population, and that heritability is highly environment dependent.

To gain insight in the impact of different trait architectures, we simulated eight distinct

architectures, including additive and epistatic architectures (see Materials and methods,

Figure S5, and **Figure S6**) and measured trait variance ratios between the IL and RIL

population. Subsequently, we assumed that the measurement error (σ_e^2) was similar for both

populations and calculated the heritability ratio between both populations (**Figure 3A**). Based

on the additive simulation, we expected 95% of the variance ratios to fall between –3.6 and –

1.7, whereas under a model with closely-linked QTL that cancel each other out in the same

genetic background (balanced), the 95% interval falls between –24.9 and 4.9. The balanced

closely-linked QTL architecture was modelled such that QTL within such a cluster would

only separate occasionally by a recombination in the RIL and IL population. The shape of the

distribution of genes with significant heritability seems to form a combination of additive and

balanced closely linked QTL (**Figure 3A**). To get an estimate of the contribution of the two

models, we used the difference in the additive and balanced closely linked distributions from

the simulation to estimate how the traits are divided over the two QTL models (**Figure 3B**).

Analysis of the two distributions revealed that in all three environments additivity was the

most likely QTL model (60 – 93% of genes with significant H^2). Balanced closely linked

QTL explained up to 40% of relative heritability in the control environment, but only 7% in

the heat stress and recovery environments (**Table S5**). Therefore, we hypothesize that the

balanced closely linked QTL model is an important eQTL architecture in the nematode *C.*

elegans.

Figure 3: Balanced closely linked QTL can explain heritability ratios between the RIL and IL populations. (**A**) Histogram of the heritability ratio in the IL versus the RIL population of genes with significant gene expression heritability (permutation, FDR = 0.05). The light orange surface indicates microarray spots of genes with heritable variation in gene expression only in the RIL population (3262 in control, 7545 in heat-stress, and 12176 in recovery), the pink surface microarray spots of genes with only heritable variation in gene expression in the IL population (145, 59, and 71 respectively) and the orange color microarray spots of genes with heritable variation in gene expression in both populations (1216, 1588, and 1507 respectively). (**B**) The density distribution of the heritability ratios of genes with heritable gene gene expression variation over the balanced closely linked (green) and the additive (purple) QTL model. Trait models consist only of additive or balanced closely linked QTL. (**C**) Overlap in differentially expressed genes between ILs (threshold $-\log_{10}(p) > 5$). The

comparisons are grouped per chromosome (x-axis): the intra-chromosomal comparisons and all other chromosomes versus chromosome I, II, III, IV, V, or X. On the y-axis the fraction overlap is shown. The dots represent one IL versus IL comparison the size corresponds to the number of overlapping differentially expressed genes.

Two predictions from the balanced closely linked eQTL microarchitecture

The balanced closely linked QTL model comes with two testable predictions: (i) the genes with heritable variation in gene expression were not differentially expressed in the parental strains, (ii) differential expression of genes was not only linked to one location in ILs but to clusters of ILs perturbing distant loci (multiple balanced clusters). Indeed, only 9.1 to 15.4% of the genes with heritable variation in gene expression were differentially expressed when comparing the two parental strains (FDR = 0.1; **Table S6**). Also, the second prediction was found to be correct, on average 21.9% of the differentially expressed genes were shared between IL strains covering loci on different chromosomes, which was only slightly less than the average of 25.1% between ILs covering the same chromosome (**Figure 3C** and **Figure S7**, and for examples **Figure 4**). Therefore, we concluded that localized genetic complexity was a major part of the eQTL architecture.

Furthermore, differences in heritability between the RIL and IL populations over environments indicated that the control environment was different from heat-shock and recovery. The balanced closely linked architecture, which predicted a broad range of relative heritability, seemed to be most prominent in the control environment. It could be that these complex architectures were more important for the developmental process. This environmental effect could be illustrated by the expression variation of the genes: *mtl-2*, *npp-1*, and *clec-62*. Of which *mtl-2* and *npp-1* showed an interaction with the environment. The gene *npp-1* only showed this interaction in the control environments, environment where ILs on multiple chromosomes showed differential expression of this gene, yet for *mtl-2* ILs with differential expression on different chromosomes were found during recovery conditions,

whereas *clec-62* shows ILs with differential expression on different chromosomes in control

and heat stress conditions. (**Figure 4**). The nuclear core complex protein NPP-1 is involved

in spindle formation and important for oogenesis, a process that was interrupted during heat-

shock (SCHETTER *et al.* 2006; JOVIC *et al.* 2017). MTL-2, one of two metallothioneins in *C.*

elegans, was expressed in the intestine and was induced after heat-shock (FREEDMAN *et al.*

1993). The pattern observed for *clec-62* was exceptional as it showed a consistent response

over all three environments where IL strains with an introgression on chromosome I,

chromosome IV, and chromosome X showed much lower expression compared to the N2

ancestor. Furthermore, also an IL strain with an introgression on chromosome II and one with

an introgression on chromosome V displayed this phenotype. C-type lectin 62 is a gene

belonging to the extensive *clec*-family, which is thought to be involved in *C. elegans*

immunity (SCHULENBURG *et al.* 2008) although its exact function is not known.

Figure 4: Three genes with high heritability in gene expression variation in the IL population compared to the RIL population. (**A**) Comparison in expression between IL and RIL population for *clec-62*, *mtl-2*, and *npp-1*. On the x-axis, the population type and on the y-axis, the z-score with the N2 parental lines. The dashed horizontal lines indicate the significance threshold (z-score, -$\log_{10}$(p) > 5). (**B**) The expression per introgression. On the x-axis, the introgression location is shown per chromosome. On the y-axis, the z-score with the N2 parental line is shown. The dashed horizontal lines indicate the significance threshold (z-score, -$\log_{10}$(p) > 5). The colors indicate the environments (blue for control, red for heat-stress, and green for recovery).

Discussion

Most eQTL mapped in RILs are replicable in ILs

Here, we show that eQTL mapped in a RIL population can be confirmed in a different type of inbred line population. We found that the majority of *cis*-eQTL are replicable in the IL population (on average 69.5% per IL), whereas *trans*-eQTL are less replicable in ILs (on average 42.2% per IL). This is likely to reflect the mainly monogenic architecture of *cis*-eQTL, versus the more polygenic and environment-dependent *trans*-eQTL (LI *et al.* 2006; KEURENTJES *et al.* 2007; ROCKMAN *et al.* 2010; SNOEK *et al.* 2017; ALBERT *et al.* 2018). Especially the confirmation of *trans*-bands by ILs shows that these regulatory hot-spots are robust regardless of genetic background. It should be noted that this study does have limited power per introgression line as each line was only measured once per condition. Hence we avoid conclusions based on a single ILs throughout this paper. However, the tiling nature of the population does mean that each locus is covered by multiple ILs (DOROSZUK *et al.* 2009).

This study helps the search for candidate genes underlying these *trans*-bands, by narrowing-down the regions. However, additional experiments and replicates are recommendable and currently pursued for the heat-stress *trans*-band on chromosome IV. Furthermore, it shows how ILs can be used to narrow-down these eQTL hotspots (SNOEK *et al.* 2012), using correlation analysis previously used to link *trans*-bands to genes and biological processes (ANDERSEN *et al.* 2014; STERKEN *et al.* 2017). These findings show on a large scale that QTL mapped using a single marker model in a moderately sized RIL population are reliably replicable in a population with a different genetic structure, which confirms findings for single traits reported in *C. elegans* and beyond (for example, see (SNOEK *et al.* 2012; ANDERSEN *et al.* 2014; GAO *et al.* 2018)).

Introgression lines reveal the presence of parental-balanced, polygenic, traits

The difference in genetic complexity between RILs and ILs can be leveraged to understand

trait architectures. Despite confirming many QTL, it has been noted that ILs often tell a

different story than RILs, implying abundant genetic interactions, not uncovered by RILs (as

reviewed by (MACKAY 2014)). In this study, we also show that the number of differentially

expressed genes in the N2 genetic background of the ILs is lower than expected compared to

RIL-based estimations. Furthermore, eQTL mapped in the RIL population also led to an

underestimation of the number of differentially expressed genes due to the introgressions in

the ILs. To summarize, the introgressions display more-than-additive effects, whereas the

genetic background shows less-than-additive effects.

Our findings are in line with observations in introgression lines in *C. elegans*, showing

that for some traits more and different QTL than expected from RILs can be found (*e.g.*

(GAERTNER *et al.* 2012; GLATER *et al.* 2014; SNOEK *et al.* 2014a)). Additionally, our findings

are also in line with findings in other organisms, where such effects have been reported for

different types of traits (SHAO *et al.* 2008; GALE *et al.* 2009; SPIEZIO *et al.* 2012). In this

study, we show this effect in general, over many gene expression traits. Moreover, a direct

comparison of trait mapping in a genome-wide IL population versus a RIL population has

only been conducted in a few studies, as far as we are aware (GLATER *et al.* 2014; SNOEK *et*

al. 2014a). It has been argued that ILs showing more- or less-than-additive effects compared

to the parental strains is a hallmark of epistasis (as reviewed by (MACKAY 2014)). However,

recent studies in yeast show that additivity underlies most of the heritable trait variation

among inbred lines, where epistasis accounts for phenotypic extremes (BLOOM *et al.* 2015;

FORSBERG *et al.* 2017; ALBERT *et al.* 2018). Thus remains the question, what kind of trait

architecture drives our observations: pervasive epistasis or widespread additivity?

The developmental trait architecture consists of balanced closely linked QTL

Trait architectures underlying natural variation in *C. elegans* differ strongly over traits. For example, currently 25 quantitative trait nucleotides (QTNs) are known in *C. elegans*, capturing a majority of the heritable variation for particular traits (reviewed by (ROCKMAN 2012) and (GAERTNER AND PHILLIPS 2010) and studies by (ANDERSEN *et al.* 2014; NOBLE *et al.* 2015; SCHMID *et al.* 2015; COOK *et al.* 2016; GREENE *et al.* 2016a; GREENE *et al.* 2016b; LARGE *et al.* 2016; BEN-DAVID *et al.* 2017; STERKEN *et al.* 2017; ZDRALJEVIC *et al.* 2017; HAHNEL *et al.* 2018; ZDRALJEVIC *et al.* 2019). However, there are also examples of traits that are highly heritable but have only yielded complex or few QTL (let alone QTNs). For example, a study on bacterial preference of *C. elegans* noted a relatively high heritability (0.46) for *Serratia marcescens* over *E. coli*, yet uncovered only a single QTL in the RILs used for mapping whereas multiple were expected (GLATER *et al.* 2014). Furthermore, recent work from our group, studying metabolite abundances showed high heritability (from 0.32 up to 0.82) corresponding to a few uncovered QTL (GAO *et al.* 2018). Intriguingly, several studies imply trait architectures consisting of closely linked QTL that are balanced in the parental strains (GAERTNER *et al.* 2012; GLATER *et al.* 2014). By simulation, we show that some trait architectures can be differentiated by relative heritability between IL and RIL populations which is a property we can reliably measure in our two populations, as it only relies on determining trait variance. In this way, we found that for genes without an eQTL, but showing high heritability, the most parsimonious explanation lies in an architecture comprised of (multiple) closely linked QTL clusters. This type of architecture seems especially prominent during normal development where it could affect approximately 40% of genes with heritable expression variation. It should be noted that only strictly balanced closely linked and strictly additive QTL distributions were modelled. We think the observed distribution suggests that a combination of (both or multiple) types of QTL make up the complete genetic architecture of a trait.

In recent years, the number of parameters known to affect gene expression variation in

the context of natural variation has steadily grown. It is currently clear that environment (LI *et*

al. 2006; SNOEK *et al.* 2017), age (VINUELA *et al.* 2010), development (FRANCESCONI AND

LEHNER 2014), and background mutations (STERKEN *et al.* 2017) affect gene expression

variation and eQTL distribution in *C. elegans*. These factors could also contribute to the

observed heritability differences between ILs and RILs. For environment, we can confirm that

it plays an important role: the heritability ratio is strongly dependent on environment. This is

in line with effects observed for *trans*-eQTL, which are also strongly environment dependent,

implying heritability is as well (LI *et al.* 2006; SNOEK *et al.* 2017). Developmental differences

between the strains in each population could also drive some of the observed heritability

ratios. For example, ILs could be developmentally more homogenous than RILs, which could

result in lower levels of trait variance in ILs. It is clear that age affects *trans*-eQTL and

heritability (VINUELA *et al.* 2010; VINUELA *et al.* 2012). However, it is unclear how that

would affect the heritability ratios we measure in general; it seems unlikely that

developmental differences result in a specific variance signature as changes occur in both

populations. More specific, in this experiment strains are tightly synchronized and

developmental stage is observed to be L4 before isolation, therefore we expect that these

potential sources of increased variance are controlled. Hence, the relative heritabilities under

normal development imply clusters of closely linked QTL.

The role of cryptic genetic variation on trait complexity

We used the contrast in genetic complexity of the RIL and IL population to investigate trait

complexity. The relative heritabilities between these populations differed over the three

environments. In our previous study on the RILs, we show that the *trans*-eQTL architecture is

different over the three environments, showing that *trans*-eQTL contribute most to cryptic

genetic variation. Here, we find that the micro-architectures is more complex in an ambient environment.

Development of an organism is a tightly regulated process and completion of development into a reproducing adult is essential for reproduction, hence fitness. The nematode *C. elegans* experiences outbreeding depression (DOLGIN *et al.* 2007), given the strains N2 and CB4856 are genetically distinct, it is likely that inbred lines constructed from these two strains disrupt some linked loci involved in this process (SEIDEL *et al.* 2008; SNOEK *et al.* 2014a). We hypothesize that especially the ambient environment stimulating normal development is prone to accentuate the effects of genetic 'mismatches' resulting in trait variation. First, local sites are expected to co-segregate over many generations in *C. elegans*, making it likely that they are filled with compensatory mutations (ROCKMAN *et al.* 2010). Second, development is a polygenic, tightly regulated, process. On the level of gene expression, there are strong changes over the entire development from egg to adult, and during the last juvenile stage (L4) especially (FRANCESCONI AND LEHNER 2014; HENDRIKS *et al.* 2014; SNOEK *et al.* 2014b). Together, these two effects may cause the *C. elegans* populations in ambient environments to show micro-architectures that are more complex.

On the other hand, stress responses are also tightly regulated, but initially aimed at reaching a stress-survival mode, which is a process that involves fewer genes than development. The heat-shock response in *C. elegans* starts by de-phosphorylation of HSF-1 and DAF-16 and entry of these transcription factors into the nucleus (as reviewed by (RODRIGUEZ *et al.* 2013)). Once there, they regulate the expression of chaperones that function to minimize cellular damage. Our results show that populations in both the heat-shock and recovery from heat-shock environments show fewer QTL with complex micro-architectures. We hypothesize that this is due to the effect of the strong transcriptional response induced by heat shock (BRUNQUELL *et al.* 2016; JOVIC *et al.* 2017). It is possible that

this abrupt disturbance emphasizes regulatory variation in only a few key response pathways

compared to the more subtle and complex process of development.

Conclusions

We present an eQTL experiment conducted with ILs in three environments covering the same

conditions as a RIL experiment published previously (SNOEK *et al.* 2017). We show that ILs

can replicate both *cis-* and *trans*-eQTL; furthermore, we present evidence supporting 14

eQTL hot spots (*trans*-bands). Yet, eQTL mapped in the RIL population systematically

under-estimate the impact of a single introgression on the transcriptome and over-estimate the

impact of the genetic background, suggesting additional genetic complexity. We show that

during normal growth multiple clusters of closely linked QTL are underlying additional

genetic complexity.

Further understanding of this phenomenon requires systematic dissection of many

QTL – a process that in this study was only undertaken for *trans*-bands – using introgression

lines. It remains to be determined what kind of role epistasis plays in trait variation. Currently,

RIL populations in most species are typically of insufficient size to detect any epistasis

beyond two loci interactions. Careful crosses with ILs to generate double-loci ILs might be a

way forward to further our understanding of trait regulation in the context of natural genetic

variation.

Acknowledgements

The authors thank the GRAPPLE project partners: Olga Valba, Sreenivas Chavali, Benjamin Lang, Mirko Francesconi, Rachel Brenchley, Arjen van 't Hof, Sergei Nechaev, Olga Vasieva, Madan Babu, Andrew Cossins, and Ben Lehner. We thank Harm Nijveen for assistance with WormQTL2.

Funding

LBS was funded by ERASysbio-plus ZonMW project GRAPPLE - Iterative modelling of gene regulatory interactions underlying stress, disease and ageing in *C. elegans* (project 90.201.066) and The Netherlands Organization for Scientific Research (project no. 823.01.001). JEK was funded by NIH grant 1R01AA 026658-01. The funding bodies had no role in the design nor the collection, analysis, and interpretation of the data, nor the writing of the manuscript.

Availability of data and materials

The strains used in this study can be requested from the authors. The transcriptome datasets used in the analysis are deposited at ArrayExpress E-MTAB-5779 and E-MTAB-7424.

Author contributions

MGS, LBS and JEK conceived and designed the experiments. MGS wrote the manuscript, with input from JEK and LBS. MGS, RPJB, RJMV, and JAGR performed the experiments. MGS analyzed the data, with input from LBS. All authors commented on the manuscript.

Supplementary information

Figure S1: Coverage per locus and per QTL. (**A**) the coverage in CB4856 loci per location on

the genome, split out for ILs and RILs. The 56 ILs together have a higher coverage over the

chromosome arms, where also most QTL map. The 48 RILs have a more homogenous

distribution, only at the *peel-1/zeel-1* locus on chromosome I there is low coverage (SEIDEL *et*

al. 2008). (**B**). A histogram of the number of CB4856 loci covering an eQTL. Typically, an

eQTL is covered by CB4856 loci of 2 ILs and 23 RILs (median).

Figure S2: Confirmation of eQTL by significance thresholds and correlation analysis. (**A**)

The percentage of confirmed eQTL per population per eQTL-type. On the x-axis, the $-\log_{10}(p)$

significance threshold in the RILs is shown (binned per 0.5 in the analysis) and on the y-axis

the percentage of confirmed eQTL per bin per strain is shown (at least five confirmable eQTL

required per bin). For example, for the RIL WN98 under recovery conditions there were 115

cis-eQTL with a significance between 4.0 and 4.5 in its CB4856 regions, 91 of these were

confirmed based on the z-score (79.1%). This example results in a dot where the size

corresponds to the number of eQTL and the color indicates the environments. The line shown

is a linear regression curve, where the surrounding grey area is the confidence interval. (**B**)

The Pearson correlation of eQTL effects with gene expression in the ILs and the RILs. On the

x-axis the population type is plotted and on the y-axis the Pearson correlation. The

correlations are split-out over the three environments (control, heat stress, and recovery) and

over eQTL type (*cis* or *trans*).

Figure S3: Differentially expressed genes in RILs and ILs. (**A**) Number of differentially

expressed genes (DEG) expected from eQTL mapping and observed experimentally, split out

per environment. On the x-axis the population and on the y-axis the expected and observed

ratio of differentially expressed genes is shown, both normalized for the expectation in the

RILs (eQTL threshold, $-\log_{10}(p) > 3.9$; DEG threshold, $-\log_{10}(p) > 5$). In the top panel the

effect compared to N2 is shown (effect of the CB4856 loci; introgressions in the IL panel) and

in the bottom panel the effect compared to CB4856 is shown (effect of the N2 loci; genetic

background in the IL panel). (**B**) The effect of different thresholds for eQTL significance (x-

axis) and DEG calling (y-axis) on the ratio between observed and expected.

Figure S4: Genes with heritable variation in gene expression in the RIL and IL populations.

(**A**) A histogram of the measured heritabilities for the IL and RIL populations, negative values

have been discarded. Colours indicate whether the heritability was significant (FDR = 0.05) in

the control (blue), heat-stress (red), or recovery (green) environment. (**B**) The overlap in

genes with significant heritability in gene expression variation over the environments for the

RIL population (FDR = 0.05). (**C**) The overlap in genes with significant heritable variation in

gene expression over the environments for the IL population (FDR = 0.05).

Figure S5: A graphical overview of the modelled trait architectures. The arrows indicate QTL

effect sizes as found in the N2 genotype (orange) and the CB4856 genotype (blue). Perfectly

opposed arrows, one of which orange and the other blue, indicate that the same QTL has an

opposite effect in the two genotypes (e.g. as seen for Additive random distribution). When an

arrow lacks a perfectly opposed arrow of another colour, it means that the QTL is only found

in that particular genotype. Epistatic interactions are indicated as a line connecting two

rectangles.

Figure S6: Variance ratios between IL and RIL population for the simulated trait

architectures (**Figure S5**). The density of occurrence of variance ratios per simulation is

given. The colour scale indicates how many QTL for a trait were simulated.

Figure S7: Overlap in differentially expressed genes between ILs (threshold $-\log_{10}(p) > 5$.

The ILs are ordered based on introgression on the x- and y-axis. The location of the CB4856

segment is indicated behind the IL strain-code. The fraction overlap shown is calculated as the

percentage of unique differentially expressed genes in the two compared ILs. The overlap is

shown for the control environment (**A**), the heat-stress environment (**B**), and the recovery

environment (**C**).

File S1: Comparison of gene expression in ILs with eQTL in *trans*-bands identified in the RIL

population. Per *trans*-band, a figure was constructed. In (**A**) the name of the *trans*-band is

given at the top of the panel (e.g. chromosome I, 3.5-4 million bases) the correlation of the

relative gene expression (R_{N2}) was correlated with the eQTL effect for the ILs covering or

nearby the *trans*-band. Significance was determined based on the highest correlation in the N2

strains, only strains with a stronger correlation than N2 were scored as significant. In (**B**) the

genetic map of the ILs of the *trans*-band chromosome is shown. On the x-axis the genomic

location (in million bases) is shown and on the y-axis the ILs. In (**C**) the actual correlation

between expression in the ILs and eQTL effect in the RILs is shown. With on the x-axis the

eQTL effect and on the y-axis R_{N2}. Each dot represents a spot on the microarray with an

eQTL in the *trans*-band investigated. The line indicates the estimated slope and the pale blue

area around the line indicates the confidence interval of the fit.

Table S1: The genotypes of the strains used in this study, -1 denotes a CB4856 genotype, a 1

denotes an N2 genotype. The genotypes are given per chromosome and per position (based on

WormBase version WS258)

Table S2: Table of the eQTL with gene expression comparisons per strain per environment.

The table lists the strain name, strain type and the comparisons are split out for the *cis-* and

trans-eQTL. The correlation reported is the Pearson correlation and its significance. The

number of eQTL present in the CB4856 loci is given, as is the percentage that is also

differentially expressed (FDR = 0.05).

Table S3: Summary of the comparison of gene expression in the ILs with the *trans*-bands

identified previously (Snoek *et al.* 2017). The location of the *trans*-band and the number of

genes and microarray spots affected are listed. The genotypes of the ILs with an introgression

adjacent to – or covering the *trans*-band are listed. The genotypes of the ILs that confirm the

trans-band and the *trans*-band locus that these ILs imply are also given.

Table S4: Table with the heritabilities per spot per environment per population. The estimated

genetic variance (V_g) and the technical variance (measurement error, V_e) are given, as is the

heritability (H^2). The FDR0.05 column indicates the threshold of significance based on

permutation.

Table S5: A table with the estimated fraction and amount of traits for an additive QTL

architecture and a balanced closely-linked QTL architecture. The numbers are calculated

based on only a single type of QTL occurring for a particular architecture. For example,

additive means *n* QTL that display an additive effect, without any other type of effect for that

simulated trait.

Table S6: differentially expressed genes between N2 and CB4856 for the three environments:

control, heat-stress, and recovery from heat stress. The columns in the table: SpotID, the

Agilent spot identifier, the information on the gene expression detected by that spot is given

(chromosome, location (start/stop), strand, and three identifiers). Two columns show if genes

are significantly heritable; general (in any population) and specific (which population, or

both). The significance and effect columns give the output of the linear model when

comparing CB4856 versus N2 (positive effect indicates the gene is higher expressed in N2).

The last column gives the significance as corrected for multiple testing (FDR).

References
751
Albert, F. W., J. S. Bloom, J. Siegel, L. Day and L. Kruglyak, 2018 Genetics of trans-
regulatory variation in gene expression. Elife 7.
Albert, F. W., and L. Kruglyak, 2015 The role of regulatory variation in complex traits and
disease. Nat Rev Genet 16: 197-212.
Andersen, E. C., J. S. Bloom, J. P. Gerke and L. Kruglyak, 2014 A variant in the neuropeptide
receptor npr-1 is a major determinant of Caenorhabditis elegans growth and
physiology. PLoS Genet 10: e1004156.
Ben-David, E., A. Burga and L. Kruglyak, 2017 A maternal-effect selfish genetic element in
Caenorhabditis elegans. Science 356: 1051-1055.
Benjamini, Y., and Y. Hochberg, 1995 Controlling the False Discovery Rate - a Practical and
Powerful Approach to Multiple Testing. Journal of the Royal Statistical Society Series
B-Statistical Methodology 57: 289-300.
Bloom, J. S., I. Kotenko, M. J. Sadhu, S. Treusch, F. W. Albert et al., 2015 Genetic
interactions contribute less than additive effects to quantitative trait variation in yeast.
Nature Communications 6.
Brem, R. B., and L. Kruglyak, 2005 The landscape of genetic complexity across 5,700 gene
expression traits in yeast. Proc Natl Acad Sci U S A 102: 1572-1577.
Brenner, S., 1974 The genetics of Caenorhabditis elegans. Genetics 77: 71-94.
Brunquell, J., S. Morris, Y. Lu, F. Cheng and S. D. Westerheide, 2016 The genome-wide role
of HSF-1 in the regulation of gene expression in Caenorhabditis elegans. BMC
Genomics 17: 559.
Conover, W. J., M. E. Johnson and M. M. Johnson, 1981 A Comparative-Study of Tests for
Homogeneity of Variances, with Applications to the Outer Continental-Shelf Bidding
Data. Technometrics 23: 351-361.
Cook, D. E., S. Zdraljevic, R. E. Tanny, B. Seo, D. D. Riccardi et al., 2016 The Genetic Basis
of Natural Variation in Caenorhabditis elegans Telomere Length. Genetics 204: 371-
383.
Dolgin, E. S., B. Charlesworth, S. E. Baird and A. D. Cutter, 2007 Inbreeding and
outbreeding depression in Caenorhabditis nematodes. Evolution 61: 1339-1352.
Doroszuk, A., L. B. Snoek, E. Fradin, J. Riksen and J. Kammenga, 2009 A genome-wide
library of CB4856/N2 introgression lines of Caenorhabditis elegans. Nucleic Acids
Res 37: e110.
Duveau, F., and M. A. Felix, 2012 Role of pleiotropy in the evolution of a cryptic
developmental variation in Caenorhabditis elegans. PLoS Biol 10: e1001230.
Forsberg, S. K., J. S. Bloom, M. J. Sadhu, L. Kruglyak and O. Carlborg, 2017 Accounting for
genetic interactions improves modeling of individual quantitative trait phenotypes in
yeast. Nat Genet 49: 497-503.
Fournier, T., and J. Schacherer, 2017 Genetic backgrounds and hidden trait complexity in
natural populations. Curr Opin Genet Dev 47: 48-53.
Francesconi, M., and B. Lehner, 2014 The effects of genetic variation on gene expression
dynamics during development. Nature 505: 208-211.
Freedman, J. H., L. W. Slice, D. Dixon, A. Fire and C. S. Rubin, 1993 The novel
metallothionein genes of Caenorhabditis elegans. Structural organization and
inducible, cell-specific expression. J Biol Chem 268: 2554-2564.
Gaertner, B. E., M. D. Parmenter, M. V. Rockman, L. Kruglyak and P. C. Phillips, 2012 More
than the sum of its parts: a complex epistatic network underlies natural variation in
thermal preference behavior in Caenorhabditis elegans. Genetics 192: 1533-1542.

Gaertner, B. E., and P. C. Phillips, 2010 Caenorhabditis elegans as a platform for molecular
quantitative genetics and the systems biology of natural variation. Genetics Research
92: 331-348.
Gale, G. D., R. D. Yazdi, A. H. Khan, A. J. Lusis, R. C. Davis *et al.*, 2009 A genome-wide
panel of congenic mice reveals widespread epistasis of behavior quantitative trait loci.
Mol Psychiatry 14: 631-645.
Gao, A. W. W., M. G. Sterken, J. U. de Bos, J. van Creij, R. Kamble *et al.*, 2018 Natural
genetic variation in C. elegans identified genomic loci controlling metabolite levels.
Genome Research 28: 1296-1308.
Gilad, Y., S. A. Rifkin and J. K. Pritchard, 2008 Revealing the architecture of gene regulation:
the promise of eQTL studies. Trends Genet 24: 408-415.
Glater, E. E., M. V. Rockman and C. I. Bargmann, 2014 Multigenic natural variation
underlies Caenorhabditis elegans olfactory preference for the bacterial pathogen
Serratia marcescens. G3 (Bethesda) 4: 265-276.
Greene, J. S., M. Brown, M. Dobosiewicz, I. G. Ishida, E. Z. Macosko *et al.*, 2016a Balancing
selection shapes density-dependent foraging behaviour. Nature 539: 254-258.
Greene, J. S., M. Dobosiewicz, R. A. Butcher, P. T. McGrath and C. I. Bargmann, 2016b
Regulatory changes in two chemoreceptor genes contribute to a Caenorhabditis
elegans QTL for foraging behavior. Elife 5.
Hahnel, S. R., S. Zdraljevic, B. C. Rodriguez, Y. Zhao, P. T. McGrath *et al.*, 2018 Extreme
allelic heterogeneity at a Caenorhabditis elegans beta-tubulin locus explains natural
resistance to benzimidazoles. PLoS Pathog 14: e1007226.
Hendriks, G. J., D. Gaidatzis, F. Aeschimann and H. Grosshans, 2014 Extensive oscillatory
gene expression during C. elegans larval development. Mol Cell 53: 380-392.
Jansen, R. C., and J. P. Nap, 2001 Genetical genomics: the added value from segregation.
Trends Genet 17: 388-391.
Jimenez-Gomez, J. M., J. A. Corwin, B. Joseph, J. N. Maloof and D. J. Kliebenstein, 2011
Genomic Analysis of QTLs and Genes Altering Natural Variation in Stochastic Noise.
Plos Genetics 7.
Jimenez-Gomez, J. M., A. D. Wallace and J. N. Maloof, 2010 Network Analysis Identifies
ELF3 as a QTL for the Shade Avoidance Response in Arabidopsis. Plos Genetics 6.
Jovic, K., M. G. Sterken, J. Grilli, R. P. J. Bevers, M. Rodriguez *et al.*, 2017 Temporal
dynamics of gene expression in heat-stressed Caenorhabditis elegans. PLoS One 12:
e0189445.
Keurentjes, J. J. B., J. Y. Fu, I. R. Terpstra, J. M. Garcia, G. van den Ackerveken *et al.*, 2007
Regulatory network construction in Arabidopsis by using genome-wide gene
expression quantitative trait loci. Proceedings of the National Academy of Sciences of
the United States of America 104: 1708-1713.
Large, E. E., W. Xu, Y. Zhao, S. C. Brady, L. Long *et al.*, 2016 Selection on a Subunit of the
NURF Chromatin Remodeler Modifies Life History Traits in a Domesticated Strain of
Caenorhabditis elegans. PLoS Genet 12: e1006219.
Li, Y., O. A. A. Lvarez, E. W. Gutteling, M. Tijsterman, J. J. Fu *et al.*, 2006 Mapping
determinants of gene expression plasticity by genetical genomics in C-elegans. Plos
Genetics 2: 2155-2161.
Mackay, T. F. C., 2014 Epistasis and quantitative traits: using model organisms to study gene-
gene interactions. Nature Reviews Genetics 15: 22-33.
Muenke, M., U. Schell, A. Hehr, N. H. Robin, H. W. Losken *et al.*, 1994 A common mutation
in the fibroblast growth factor receptor 1 gene in Pfeiffer syndrome. Nat Genet 8: 269-
274.

Nica, A. C., and E. T. Dermitzakis, 2013 Expression quantitative trait loci: present and future.
Philos Trans R Soc Lond B Biol Sci 368: 20120362.
Noble, L. M., A. S. Chang, D. McNelis, M. Kramer, M. Yen *et al.*, 2015 Natural Variation in
plep-1 Causes Male-Male Copulatory Behavior in C. elegans. Curr Biol 25: 2730-
2737.
Radwan, J., and W. Babik, 2012 The genomics of adaptation Introduction. Proceedings of the
Royal Society B-Biological Sciences 279: 5024-5028.
Ranjan, A., J. M. Budke, S. D. Rowland, D. H. Chitwood, R. Kumar *et al.*, 2016 eQTL
Regulating Transcript Levels Associated with Diverse Biological Processes in
Tomato. Plant Physiology 172: 328-340.
Ritchie, M. E., B. Phipson, D. Wu, Y. Hu, C. W. Law *et al.*, 2015 limma powers differential
expression analyses for RNA-sequencing and microarray studies. Nucleic Acids Res
43: e47.
Rockman, M. V., 2012 The Qtn Program and the Alleles That Matter for Evolution: All That's
Gold Does Not Glitter. Evolution 66: 1-17.
Rockman, M. V., S. S. Skrovanek and L. Kruglyak, 2010 Selection at Linked Sites Shapes
Heritable Phenotypic Variation in C. elegans. Science 330: 372-376.
Rodriguez, M., L. B. Snoek, M. De Bono and J. E. Kammenga, 2013 Worms under stress: C.
elegans stress response and its relevance to complex human disease and aging. Trends
Genet 29: 367-374.
Schadt, E. E., S. A. Monks, T. A. Drake, A. J. Lusis, N. Che *et al.*, 2003 Genetics of gene
expression surveyed in maize, mouse and man. Nature 422: 297-302.
Schetter, A., P. Askjaer, F. Piano, I. Mattaj and K. Kemphues, 2006 Nucleoporins NPP-1,
NPP-3, NPP-4, NPP-11 and NPP-13 are required for proper spindle orientation in C.
elegans. Dev Biol 289: 360-371.
Schmid, T., L. B. Snoek, E. Frohli, M. L. van der Bent, J. Kammenga *et al.*, 2015 Systemic
Regulation of RAS/MAPK Signaling by the Serotonin Metabolite 5-HIAA. PLoS
Genet 11: e1005236.
Schulenburg, H., M. P. Hoeppner, J. Weiner, 3rd and E. Bornberg-Bauer, 2008 Specificity of
the innate immune system and diversity of C-type lectin domain (CTLD) proteins in
the nematode Caenorhabditis elegans. Immunobiology 213: 237-250.
Seidel, H. S., M. V. Rockman and L. Kruglyak, 2008 Widespread genetic incompatibility in
C. elegans maintained by balancing selection. Science 319: 589-594.
Shao, H., L. C. Burrage, D. S. Sinasac, A. E. Hill, S. R. Ernest *et al.*, 2008 Genetic
architecture of complex traits: large phenotypic effects and pervasive epistasis. Proc
Natl Acad Sci U S A 105: 19910-19914.
Smyth, G. K., and T. Speed, 2003 Normalization of cDNA microarray data. Methods 31: 265-
273.
Snoek, B. L., M. G. Sterken, R. P. J. Bevers, R. J. M. Volkers, A. Van't Hof *et al.*, 2017
Contribution of trans regulatory eQTL to cryptic genetic variation in C. elegans. BMC
Genomics 18: 500.
Snoek, B. L., M. G. Sterken, M. Hartanto, A.-J. van Zuilichem, J. E. Kammenga *et al.*, 2019
WormQTL2: an interactive platform for systems genetics in Caenorhabditis elegans.
bioRxiv: 766386.
Snoek, L. B., H. E. Orbidans, J. J. Stastna, A. Aartse, M. Rodriguez *et al.*, 2014a Widespread
genomic incompatibilities in Caenorhabditis elegans. G3 (Bethesda) 4: 1813-1823.
Snoek, L. B., M. G. Sterken, R. J. Volkers, M. Klatter, K. J. Bosman *et al.*, 2014b A rapid and
massive gene expression shift marking adolescent transition in C. elegans. Sci Rep 4:
3912.

Snoek, L. B., I. R. Terpstra, R. Dekter, G. Van den Ackerveken and A. J. Peeters, 2012
Genetical Genomics Reveals Large Scale Genotype-By-Environment Interactions in
Arabidopsis thaliana. Front Genet 3**:** 317.
Spiezio, S. H., T. Takada, T. Shiroishi and J. H. Nadeau, 2012 Genetic divergence and the
genetic architecture of complex traits in chromosome substitution strains of mice.
BMC Genet 13**:** 38.
Sterken, M. G., L. B. Snoek, J. E. Kammenga and E. C. Andersen, 2015 The laboratory
domestication of Caenorhabditis elegans. Trends Genet 31**:** 224-231.
Sterken, M. G., L. van Bemmelen van der Plaat, J. A. G. Riksen, M. Rodriguez, T. Schmid *et*
al., 2017 Ras/MAPK Modifier Loci Revealed by eQTL in Caenorhabditis elegans. G3
(Bethesda) 7**:** 3185-3193.
Team, R. C., 2017 R: A language and environment for statistical computing, pp. in
Vienna, Austria.
Terpstra, I. R., L. B. Snoek, J. J. Keurentjes, A. J. Peeters and G. van den Ackerveken, 2010
Regulatory network identification by genetical genomics: signaling downstream of the
Arabidopsis receptor-like kinase ERECTA. Plant Physiol 154**:** 1067-1078.
Thompson, O. A., L. B. Snoek, H. Nijveen, M. G. Sterken, R. J. Volkers *et al.*, 2015
Remarkably Divergent Regions Punctuate the Genome Assembly of the
Caenorhabditis elegans Hawaiian Strain CB4856. Genetics 200**:** 975-989.
van't Hof, A. E., N. Edmonds, M. Dalikova, F. Marec and I. J. Saccheri, 2011 Industrial
Melanism in British Peppered Moths Has a Singular and Recent Mutational Origin.
Science 332**:** 958-960.
Vinuela, A., L. B. Snoek, J. A. Riksen and J. E. Kammenga, 2010 Genome-wide gene
expression regulation as a function of genotype and age in C. elegans. Genome Res
20**:** 929-937.
Vinuela, A., L. B. Snoek, J. A. Riksen and J. E. Kammenga, 2012 Aging Uncouples
Heritability and Expression-QTL in Caenorhabditis elegans. G3 (Bethesda) 2**:** 597-
605.
Wickham, H., 2009 *Ggplot2 : elegant graphics for data analysis*. Springer, New York.
Wickham, H. F., R.; Henry, L; Müller, K., 2018a dplyr: A Grammar of Data Manipulation,
pp.
Wickham, H. H., L, 2018b tidyr: Easily Tidy Data with 'spread()' and 'gather()' Functions, pp.
Zahurak, M., G. Parmigiani, W. Yu, R. B. Scharpf, D. Berman *et al.*, 2007 Pre-processing
Agilent microarray data. BMC Bioinformatics 8**:** 142.
Zdraljevic, S., B. W. Fox, C. Strand, O. Panda, F. J. Tenjo *et al.*, 2019 Natural variation in C.
elegans arsenic toxicity is explained by differences in branched chain amino acid
metabolism. Elife 8.
Zdraljevic, S., C. Strand, H. S. Seidel, D. E. Cook, J. G. Doench *et al.*, 2017 Natural variation
in a single amino acid substitution underlies physiological responses to topoisomerase
II poisons. PLoS Genet 13**:** e1006891.
Zych, K., B. L. Snoek, M. Elvin, M. Rodriguez, K. J. Van der Velde *et al.*, 2017 reGenotyper:
Detecting mislabeled samples in genetic data. PLoS One 12**:** e0171324.
939