Development and External Validation of the Acute COPD Exacerbation Prediction Tool (ACCEPT)

Amin Adibi, MSc[1]; Don D Sin, MD[2]; Abdollah Safari, PhD[1]; Kate M Johnson, MSc[1]; Shawn D Aaron, MD[5], J Mark FitzGerald, MD[3,4]; Mohsen Sadatsafavi, MD, PhD[1,3,4]

Affiliations:

[1] Respiratory Evaluation Sciences Program, Faculty of Pharmaceutical Sciences, University of British Columbia, 2405 Wesbrook Mall, Vancouver, British Columbia, Canada V6T 1Z3.

[2] Division of Respiratory Medicine, Department of Medicine, The UBC Centre for Heart Lung Innovation, St. Paul's Hospital, University of British Columbia, 1081 Burrard Street, Vancouver, British Columbia, Canada V6Z 1Y6

[3] Institute for Heart and Lung Health, Division of Respiratory Medicine, Faculty of Medicine, University of British Columbia, 7th Floor, 2775 Laurel Street, Vancouver, British Columbia, Canada V5Z 1M9

[4] Centre for Clinical Epidemiology and Evaluation, University of British Columbia, 828 West 10th Avenue, Vancouver, BC, Canada, V5Z 1M9

[5] Ottawa Hospital Research Institute, University of Ottawa, Ontario, Canada K1H 8L6

Corresponding author: Don Sin, MD[2]

Division of Respiratory Medicine, Department of Medicine, The UBC Centre for Heart Lung Innovation, St. Paul's Hospital, University of British Columbia, 1081 Burrard Street, Vancouver, British Columbia, Canada V6Z 1Y6

Phone: (604) 806-8346 x68395

E-mail: Don.Sin@hli.ubc.ca

Conflict of interest:

Funding source: CIHR Catalyst Grant (#155554)

Abstract word count (Max 250): 257 words

Main text word count (excluding title page, abstract, references, tables, figures, statements): 3450 words

Number of tables: 4

Number of figures: 4

Number of references: 26

Number of appendices: 4

Keywords: COPD, exacerbation, clinical prediction model, prognosis, precision medicine

Research in context

Evidence before this study: Preventing future exacerbations is a major goal in COPD care. Because of

adverse effects, preventative treatments should be reserved for those at a higher risk of future

exacerbations. Predicting exacerbation risk in individual patients can guide these clinical decisions. A

2017 systematic review reported that of the 27 identified COPD exacerbation prediction tools, only two

had reported external validation and none was ready for clinical implementation. To find the studies that

were published afterwards, we searched PubMed for articles on development and validation of COPD

exacerbation prediction since 2015, using the search terms "COPD", "exacerbation", "model", and

"validation". We included studies that reported prediction of either the risk or the rate of exacerbations

and excluded studies that did not report external validation.

Added value of this study: ACCEPT is the first COPD exacerbation prediction tool that jointly estimates

the individualised rate and severity of exacerbations. ACCEPT is externally validated, generalisable, and

readily accessible.

Implications of all the available evidence: The availability of an individualised exacerbation risk can

enable patients and care providers to decide on preventive therapies based on patient-specific, objectively-

established thresholds for treatment benefit.

Abstract

Background: Accurate prediction of exacerbation risk enables personalised chronic obstructive pulmonary disease (COPD) care. We developed and validated a generalisable model to predict the individualised rate and severity of COPD exacerbations.

Methods: We pooled data from three COPD trials on patients with a history of exacerbations. We developed a mixed-effect model to predict exacerbations over one-year. Severe exacerbations were those requiring inpatient care. Predictors were a history of exacerbations, age, sex, body mass index, smoking status, domiciliary oxygen therapy, lung function, symptom burden, and current medication use. ECLIPSE, a multicentre cohort study, was used for external validation.

Results: The development dataset included 2,380 patients (mean 64·7 years, 57·7% men, mean exacerbation rate 1·42/year, 20·5% severe). When validated against all COPD patients in ECLIPSE (n=1819, mean 63·3 years, 65·2% men, mean exacerbation rate 1·20/year, 22·2% severe), the area-under-curve was 0·81 (95%CI 0·79–0·83) for ≥2 exacerbations and 0·77 (95%CI 0·74–0·80) for ≥1 severe exacerbation. Predicted rates were 0·25/year for severe and 1·31/year for all exacerbations, close to the observed rates (0·27/year and 1·20/year, respectively). In ECLIPSE patients with a prior exacerbation history (n=996, mean 63·6 years, 61·3% men, mean exacerbation rate 1·82/year, 22·0% severe), the area-under-curve was 0·73 (95%CI 0·70–0·76) for ≥2 exacerbations and 0·74 (95%CI 0·70–0·78) for ≥1 severe exacerbation. Calibration was accurate for severe exacerbations (predicted=0·37/year, observed=0·40/year) and all exacerbations (predicted=1·80/year, observed=1·82/year). The model is accessible at http://resp.core.ubc.ca/ipress/accept.

Interpretation: This model can be used as a decision tool to personalise COPD treatment and prevent exacerbations.

Funding: Canadian Institutes of Health Research (Grant #155554).

1.   Background and Objectives

Chronic Obstructive Pulmonary Disease (COPD) is characterised by symptoms of breathlessness and cough, which worsen acutely during exacerbations.[1] COPD is known to be a heterogeneous disorder with large variations in the risk of exacerbation across patients.[2] In clinical practice, a history of two or more exacerbations and one severe exacerbation per year is used to guide therapeutic choices for exacerbation prevention.[3] However, this approach is clinically limited owing to significant heterogeneity in risk even within those who frequently exacerbate.[4]

Prognostic clinical prediction tools enable personalised approaches to disease management. Despite potential benefits, no such tool is routinely used in the clinical management of COPD. This is unlike COPD-related mortality for which clinical scoring schemes such as the BODE index are available and frequently used.[5] A 2017 systematic review by Guerra and colleagues identified 27 prediction tools for COPD exacerbations.[6] Among these, only two reported on the validation of the model and none were deemed ready for personalised COPD management in the clinic.[6]

Here, we describe a new model, the Acute COPD Exacerbation Prediction Tool (ACCEPT), to predict, at an individual level, the rate and severity of COPD exacerbation and report on its performance in an independent external cohort, and explain, using case studies, its potential clinical application. As a decision tool, ACCEPT provides a personalised risk profile that allows clinicians to tailor treatment regimens to the individual needs of the patients.

2.   Methods

In reporting our prediction model, we have followed recommendations set forth by the Transparent Reporting of a Multivariable Prediction Model for Individual Prognosis or Diagnosis (TRIPOD) Statement.[7]

Target population and source of data

We developed the model using data from COPD patients, without a prior or current history of asthma, and who had experienced at least one exacerbation over the previous 12 months. We then externally validated the model: 1) in COPD patients regardless of their exacerbation history, and 2) in a subset of COPD patients with at least one exacerbation over the previous 12 months.

For discovery, we pooled data across all arms of three randomised controlled trials: Macrolide Azithromycin to Prevent Rapid Worsening of Symptoms in COPD (MACRO)[8], Simvastatin for the Prevention of Exacerbations in Moderate-to-Severe COPD (STATCOPE)[9], and the Optimal Therapy of

COPD to Prevent Exacerbations and Improve Quality of Life (OPTIMAL)[10]. We used Evaluation of

COPD Longitudinally to Identify Predictive Surrogate End-Points (ECLIPSE)[11] – an independent

longitudinal COPD cohort study – for external validation. The details of each of these studies have been

previously published. Briefly, the MACRO study[8] evaluated the effect of daily low-dose azithromycin

therapy on the rate of exacerbations in COPD patients; the STATCOPE study evaluated the effects of

daily simvastatin therapy on the rate of exacerbation[9], and the OPTIMAL study evaluated the effects of

tiotropium-fluticasone-salmeterol on the rate of exacerbation compared with tiotropium-fluticasone and

tiotropium alone.[10] In all three trials, which comprised the development dataset, patients who had a

history of at least one exacerbation over the previous 12 months were recruited. ECLIPSE, on the other

hand, was a multicentre three-year, non-interventional observational study whose primary aim was to

characterise COPD phenotypes and identify novel markers of disease progression.[11] The ECLIPSE study

included patients irrespective of their prior history of an exacerbation. *Table 1* provides a detailed

summary of these studies.

113   <<Table 1>>

Outcomes

The outcomes of interest were the rates of exacerbations and severe exacerbations over one year.

Exacerbations were the primary outcome of all three trials and a major outcome measure of the ECLIPSE

study. All studies used a similar definition of exacerbations, which was based on the criteria endorsed by

the Global Initiative for Chronic Obstructive Lung Disease (GOLD) scientific committee.[3] Briefly, an

exacerbation was defined as an acute episode of intensified symptoms that required additional therapy.[3]

Mild exacerbations were defined as those that were treated with short-acting bronchodilators. Moderate

exacerbations were those that required the institution of systemic corticosteroids and/or antibiotics and

severe exacerbations were those that required an emergency department visit or a hospitalisation.[3,8–10]

Predictors

To minimize the risk of bias, optimism, and overfitting, no data-driven variable selection was performed.

We pre-specified predictors based on clinical relevance and the availability of predictors in all the

datasets. Predictors included non-severe as well as severe exacerbations over the previous year, baseline

age, sex, smoking status, percent predicted post-bronchodilator forced expiratory volume in one second

(FEV_1 % predicted[12]), St. George's Respiratory Questionnaire (SGRQ) score, body mass index (BMI), and

the use of COPD and non-COPD medications as well as domiciliary oxygen therapy during the previous

12 months. COPD medications were defined as long-acting muscarinic receptor antagonists (LAMAs),

long-acting β2 agonists (LABAs), and inhaled corticosteroids (ICS). In addition to baseline medications,

the model adjusted for treatment assignment in the therapeutic trials (azithromycin in MACRO; statins in

STATCOPE; LABA/LAMA and ICS in OPTIMAL). To facilitate clinical implementation, a web

application was created (based on conversion factors that have been previously published), which enables

the use of a COPD Assessment Test (CAT) score in lieu of SGRQ.[13]

Follow up

We applied administrative censoring at the one-year follow-up time for patients who had data beyond this

threshold. The decision to limit predictions to one year was made *a priori* based on the assumption that

predicting exacerbations over this time frame was relevant for clinical COPD management and that

prediction accuracy of the model would decrease substantially beyond one year.

Statistical analysis methods

We used a joint accelerated failure time and logistic model to characterise the rate and severity of

exacerbations. We have previously published the details of this approach elsewhere.[14] In summary, this

framework assigns two random-effect terms to each individual, quantifying each individual's specific rate

of exacerbation and the probability that once an exacerbation occurs, it will be severe (Supplementary

Appendix I). For each patient, this framework fully specifies the hazard of all exacerbations (including

their severity) at any given point in time during follow-up, enabling different predictions such as the

probability of having a specific number of total and severe exacerbations during the next twelve months.

Two forms of uncertainty in predictions were quantified: the uncertainty due to the finite sample of the

validation set (represented by the 95% confidence interval [95%CI] around the mean of projected values),

and uncertainty due to the differences in patients' specific exacerbation frequency and severity

(represented by the 95% prediction interval around the mean, the interval which has a 95% probability to

contain a future observation of a patient with the same predictors). Shrinkage methods were not applied

given the low risk of bias due to complete pre-specification of the model and the relatively high events

per predictor in the development dataset.[15] Because in this framework, the correlation between the

previous and future exacerbations rates is modelled through random-effect terms, a history of

exacerbation did not enter the model as a predictor. Instead, a Bayesian approach was employed to model

the distribution of future exacerbation rate and severity, given the exacerbation history of an individual

subject (Supplementary Appendix II). The availability of full exacerbation history in the external

validation cohort enabled validation of this approach.

External validation

We used the first year of follow-up data in ECLIPSE to establish an accurate 1-year history of

exacerbation for each patient. Next, we used the second year of follow-up to validate the model. The

model was validated first in the entire COPD cohort of ECLIPSE (n=1,819), and then in a subset of

COPD patients who had at least one exacerbation in the first year of follow-up (n=996). This subset was

similar to the population characteristics of the development dataset, while the full ECLIPSE cohort

enabled assessment of the generalisability of the model beyond patients with an exacerbation history.

We examined model calibration (the degree to which the predicted and actual risks or rates of

exacerbations aligned) and discrimination (the extent to which the model separated individuals with

different risks).[16] Calibration was assessed by comparing the predicted and observed exacerbation rates

across subgroups with differential risks, evaluating the calibration plots, and calculating Brier scores (i.e.

the mean squared error of forecast). Discrimination was assessed by calculating receiver operating

characteristic (ROC) curves, and the area-under-the-curve (AUC), and then comparing them using a

DeLong's test.[17] ROC and AUC calculations were based on the occurrence of two or more exacerbations

of any type, or one or more severe exacerbations.[3]

Ethics Approval

The study was approved by the University of British Columbia/Providence Health Research Ethics Board

(H11-00786).

3. Results

Participants

Figure 1 presents the flowchart of sample selection. We excluded 96 patients who were either lost to

follow-up (n=33) or had missing values (n=63). The final development dataset included 2,380 patients

(1,107 from MACRO, 847 from STATCOPE, and 426 from OPTIMAL; overall mean age 64.7, 57.7%

males). Patients experienced a total of 3,056 exacerbations, 628 of which were severe. In the external

validation dataset, ECLIPSE, 109 patients had missing values. Thus the final sample included 1,819

COPD patients (mean age 63·3, 65·2% male). Among these 996 COPD patients had at least one

exacerbation in the first year (mean age 63·6, 61·3% male). *Table 2* provides a detailed comparison of the

development and validation datasets in terms of demographics, predictors, and outcome variables. The

average exacerbation rates in the development dataset, the validation set with all patients, and the

validation subset containing only those with a prior history of an exacerbation were 1·42, 1·20, and 1·80

events/year, respectively. For severe exacerbations, the average rate was 0·29, 0·27, and 0·40 events/year,

respectively.

<<Figure 1 and Table 2>>

Supplementary Appendix III – Table S1 shows the distribution of baseline predictors among different

studies that were included in the development dataset. Notably, none of the STATCOPE participants had

a history of statin use because patients with cardiovascular comorbidities were excluded from this trial.

We assumed that missing values were missing at random and opted for a complete case analysis given

that only 2·58% of patients in the combined development dataset and 5·65% of patients in the validation

dataset had missing data (*Supplementary Appendix II – Tables S2 and S3).*

Model specification and performance

Table 3 provides coefficient estimates for predictors. Regression coefficients are shown as log-hazard

ratios (ln-HR) for the rate component and log-odds ratios (ln-OR) for the severity component.

<<Table 3>>

When validated against all patients in ECLIPSE (including those without an exacerbation history),

ACCEPT slightly overestimated their actual overall exacerbation rates (observed 1·20 events/year,

predicted 1·31 events/year) but was accurate for severe exacerbation rates (observed 0·27 events/year,

predicted 0·25 events/year, *Figure 2A*). The same trend was observed in all major risk-factor subgroups

(Figure 2*A)* and in both men and women (Figure 3*A)*. The Brier score was 0·20 for all exacerbations and

0·12 for severe exacerbations. In patients with an exacerbation history, ACCEPT showed robust overall

calibration: predicted annual exacerbation rate closely matched the observed rate for all exacerbations

(observed 1·82 events/year, predicted 1·80 events/year), severe exacerbations (observed 0·40 events/year,

predicted 0·37 events/year), and risk-factor subgroups (*Figure 2B).* Calibration plots comparing per decile

average rate of exacerbations showed good agreement between observed and predicted rates for both

female and male patients (Figure 3*B*). The Brier score was 0·17 for all exacerbations and 0·16 for severe

exacerbations. Similar results for the development dataset are provided in *Supplementary Appendix IV.*

In all COPD patients, the model had an AUC of 0·81 (95%CI 0·79–0·83) for ≥2 exacerbations and 0·77

(95%CI 0·74–0·80) for ≥1 severe exacerbation (Figure 4*A)*. The corresponding AUCs for COPD patients

with an exacerbation history were 0·73 (95%CI 0·70–0·76) for two or more exacerbations and 0·74

(95%CI 0·70–0·78) for at least one severe exacerbation (Figure 4*B)*.

Compared to the current practice, which relies exclusively on a prior history of exacerbation to predict

future risk of a exacerbation, ACCEPT demonstrated higher performance in predicting severe

exacerbations in all COPD patients (AUC$_{ACCEPT}$=0·77 vs. AUC$_{Event History}$=0·66, p<0·0001) and in the

subset who had a prior history of an exacerbation (AUC_{ACCEPT}=0·74 vs. $AUC_{Event\ History}$=0·67, p<0·0001).

Similarly, ACCEPT showed better performance for all exacerbations regardless of severity (Figure 4).

225   <<Figure 2, Figure 3, Figure 4>>

4. Discussion

While preventing exacerbations is a major goal in COPD care, there are no tools in practice that can

accurately predict the risk or rate of exacerbations in a given individual. Prior studies suggest that patients

with a previous history of an exacerbation are more likely to exacerbate in the future than those without.[2]

However, this approach is hampered by a relatively poor resolution, leading to large variations in risk

across subjects even among those who have the same history of exacerbations. Our framework builds

upon this well-accepted approach and extends its use by incorporating other clinical features that enable a

more accurate prediction. We used pooled data from three clinical trials to show that a model containing

simple clinical and demographic variables in aggregate can be used to predict the risk of exacerbation

with improved accuracy over 12 months of follow-up.

A 2017 systematic review of clinical prediction models for COPD exacerbations found that only two of

the 27 reviewed models – CODEX[18] and Bertens' model[19] – reported on any external validation. When

the availability of predictors and practical applicability were also taken into account, none of the models

were deemed ready for clinical implementation.[6] We are aware of only two additional prediction models

published after this review – by Kerkhof[20] and Annavarapu[21] – that have reported external validation.

ACCEPT has several notable advantages compared with these models. Importantly, it is externally

validated in an independent cohort extending its generalisability beyond therapeutic clinical trials.

ACCEPT is also geographically generalisable because the external validation cohort contained data from

12 different countries across North America, Europe, and Oceania. In contrast, previous externally

validated models used geographically limited datasets: CODEX was Spanish[18], Bertens' model was

Dutch[19], Kerkhof's model was British[20], and Annavarapu's model was based on cross-sectional

administrative data from a non-single-payer context in the United States.[21] Bertens' model, CODEX, and

models by Kerkhof and Annavarapu reported validation AUCs of 0·66 and 0·59, 0·74, and 0·77,

respectively. However, the independence of the validation dataset in Kerkhof's model was questioned as

it was selected from the same database as the developmental population. Annavarapu did not report

calibration at all, and overall, both models suffered from a lack of generalisability given the local nature

of the data that were available to the investigators.

ACCEPT predicts the rate and severity of exacerbations jointly. This is crucial to appropriately tailoring

treatments to an individual, as the more granular nature of the output in ACCEPT provides more detailed

prediction to assist clinicians in their decision making. For example, ACCEPT can predict the number of

exacerbations at a given time period, time to next exacerbation, and the probability of experiencing a

specific number of non-severe or severe exacerbations within a given follow-up time (up to one year).

This is in contrast to the logistic regression models used in a majority of previous clinical prediction

models, which allow prediction probabilities of having at least one exacerbation in a single time window.[6]

Further, this framework can potentially be used for prognostic enrichment of randomised trials by

identifying patients who are more likely to exacerbate. Similar to asthma trials, the required sample size

and consequently the cost of large trials can be substantially reduced by using prediction models to recruit

patients above a certain threshold of expected exacerbation rate.[22,23]

Examples of Application in Clinic

Example 1

ACCEPT can combine predicted risk with effect estimates from randomised trials to enable personalised

treatment. For example, a benefit-harm analysis for roflumilast as preventive therapy for COPD

exacerbations reported that the benefits of roflumilast outweighed its potential harm when patients have a

severe exacerbation risk of at least 22% over a year.[24] Using data from this benefit-harm analysis, the

accompanying web app of ACCEPT can be used to inform therapeutic decisions on the use of roflumilast

for a given patient. Another example is in the potential use of preventative daily azithromycin therapy in

COPD. Azithromycin reduces exacerbation rate by 27%.[8] However, it is associated with increased risk of

hearing impairment and antimicrobial resistance and thus should be reserved for those at a high risk of

future exacerbations.[8] The accompanying web app illustrates this application by showing the risk of

exacerbations with and without daily azithromycin therapy in a given patient. Once care providers discuss

the risks of harm and benefits of the therapy and establish patient preference thresholds for the

benefit/harm trade-off, ACCEPT can be used to determine whether the preventive azithromycin therapy

for that individual reaches or surpasses this threshold.

279   <<Table 4>>

Example 2

ACCEPT generates nuanced predictions that allow clinicians to more accurately risk-stratify two patients,

who have an identical exacerbation history. For example, consider Patient A who has the following

characteristics: 57-years of age, male ex-smoker, with a chronic bronchitis phenotype, an FEV_1 of 51% of

the predicted value (GOLD grade 2), BMI of 18 kg/m^2, and a CAT score of 29. Now consider Patient B

who is 57 years of age, female ex-smoker with a chronic bronchitis phenotype, an FEV1 of 75% (GOLD

2), a CAT score of 10, and a BMI of 30 kg/m^2. Both patients experienced two exacerbations within the

past year, one of which was severe, requiring a hospitalisation. Both are using LAMA, LABA, and ICS,

but not statins. The complete baseline characteristics of these two patients, as well as their ACCEPT

predictions, are shown in Table 4. ACCEPT indicates that Patient A's 1-year probability of experiencing

a severe exacerbation is 29% (versus 18% for Patient B). His predicted severe exacerbation rate is 0.6

events/year (versus 0.3 events/year for Patient B). Based on the benefit-harm analysis for roflumilast[24],

the web app for ACCEPT predicts that roflumilast is likely to provide a net benefit to Patient A (net

benefit probability>80%) but not for Patient B (net benefit probability<20%).

Limitations

The pooled trial data we used to develop the model lacked data on certain variables such as comorbidities,

vaccination, blood markers (e.g. eosinophil count), and socio-economic status. As such, these predictors

could not be incorporated into the model. Moreover, the developmental dataset did not contain individuals

without exacerbations in the previous year; however, the model performed robustly in an external

validation dataset that included such patients. Neither the developmental nor the validation datasets

included patients with mild (GOLD I) severity and as such, we could not establish the accuracy of

predictions for this subgroup. Additionally, our model may not be generalisable to COPD patients with a

history of asthma, lifetime non-smokers, patients younger than 40 or older than 80 years of age, or

populations outside North America, Europe, and Oceania. Model updating and re-examination of its

external validity will be necessary when new sources of data become available.[25]

Compared to simple scoring systems such as the BODE index that can be manually calculated, ACCEPT

requires relatively sophisticated computational analysis. While parsimonious models are useful at the

bedside, given the complexity of processes involved in the pathogenesis of COPD exacerbations, we

believe such tools will have limited resolution. Given the proliferation of hand-held computational

devices in clinical practice and the wide availability of clinical parameters that are contained in the model,

ACCEPT is usable clinically. This is facilitated through its availability as a web app, a spreadsheet, or the

R package 'accept'.[26]

We emphasise that estimates in our model are predictive and should not be interpreted as "causal". The

observed association between being a smoker and a having a lower exacerbation rate (hazard ratio 0·82,

95%CI 0·73–0·93) is one such example. Smoking is likely a marker of disease severity with sicker

patients less likely to smoke than those with milder disease. As such, the information in the smoking

status variable has high predictive value for the tendency towards exacerbation but is not causally

interpretable.

How to Access the Model

An easy-to-use web application for ACCEPT is available at http://resp.core.ubc.ca/ipress/accept. For any

individual patient, the web app predicts 1) the exacerbation risk within the next 12 months, 2) the annual

rate of exacerbations, and 3) the probability of experiencing any given number of exacerbations. All

outcomes are reported separately for overall and severe exacerbations.

Additionally, we provide an R package[26], a spreadsheet template, public application programming

interfaces (APIs), and additional code at http://resp.core.ubc.ca/research/Specific_Projects/accept.

5.   Conclusions

ACCEPT is an externally validated and generalisable prediction model that enables nuanced prediction of

the rate and severity of exacerbations and provides individualised estimates of risks and uncertainty in

predictions. ACCEPT has good to excellent discriminatory power in predicting the rate and severity of

COPD exacerbations in all COPD patients and showed robust calibration in individuals with a history of

such exacerbations in the past year. Objective prediction of outcomes given each patient's unique

characteristics can help clinicians to tailor treatment of COPD patients based on their individualised

prognosis.

Acknowledgment

We would like to thank Ainsleigh Hill for her contribution to development and documentation of the R package and co-investigators Janet Sutherland, Kelly Ablog-Morrant, Drs. Larry Lynd, Teresa To, Annalijn Conklin, Wenjia Chen, Laura Nimmon, Iraj Poureslami, and Hui Xie for their input and feedback.

Contributors

MS, DDS, JMF, and SDA conceived the study. AA, AS, and MS developed and validated the model. DDS and SDA contributed to data acquisition. AA, KMJ, AS, JMF, DDS, SDA, and MS contributed to the interpretation of the data. AA wrote the first draft of the manuscript and created data visualizations. JMF, DDS, SDA, and MS provided clinical input and oversight. AA developed the Web Application with critical input from KMJ, SDA, DDS, and MS. MS and AA developed the interactive spreadsheet and the R package. All authors revised the manuscript critically and approved the final version to be published.

Declaration of Interest

Authors declare no conflicts of interest.

Table 1 Available datasets with data on the rate, time, and severity of COPD exacerbations.

Data Source	Design	Intervention	Study Period (Follow-up)	Centres	Inclusion Criteria	Exclusion Criteria
Development						
MACRO	Randomized trial	Azithromycin	03-2006 to 07-2010 (1 year)	17 sites in US	>40 yr. Clinical diagnosis of COPD. >10 pack-years of smoking. O_2 or systemic glucocorticoids therapy in the last yr. Hospitalization or ER visit.	Asthma. Exacerbation in the last month. HR above 100 /min. QT_C >450ms. QT_C prolonging or TdP-related medication except for amlodipine. Hearing impairment.
STATCOPE	Randomized trial	Simvastatin	03-2010 to 01-2014 (~2 years)	45 sites (29 in US and 16 in Canada)	40-80 yrs. Clinical diagnosis of COPD. >10 pack-years of smoking. O_2 OR systemic glucocorticoids OR antibiotics therapy OR hospitalization OR ER visit in the last yr.	Asthma. Receiving statins, or should have received statins. On drugs that contradicted with statins. Unable to take statins. Active liver disease, alcoholism, or allergy.
OPTIMAL	Randomized trial	Tiotropium in combination with Salmeterol or Fluticasone-Salmeterol	10-2003 to 01-2006 (1 year)	27 sites in Canada	>35 yr. Clinical diagnosis of COPD. >10 pack-years of smoking. Exacerbation requiring systemic glucocorticoids OR antibiotics therapy in the last yr.	Asthma <40 yrs. CHF with persistent severe LVD. Oral prednisone. Intolerance to tiotropium, salmeterol, or fluticasone-salmeterol. Glaucoma. Urinary tract obstruction. Lung transplant or volume reduction. Diffuse bilateral bronchiectasis. Pregnancy or breastfeeding.
Validation						
ECLIPSE	Cohort		12-2005 to 02-2010 (3 years)	46 sites in 12 countries	40-75 yrs. Clinical diagnosis of COPD. >10 pack-years of smoking.	Respiratory disorders other than COPD. Reported exacerbation in the past month. Significant inflammatory disease.

Table 2 Comparison of the Final Development and Validation Datasets

Predictor	Development (n=2380)		Validation (n=1819)		All COPD Patients		Validation (n=996)		COPD Patients with Event History
	%	Distribution	%	Distribution		%	Distribution		
Male Sex	57·69		65·20			61·28			
Current Smokers	25·80		27·49			25·48			
O$_2$ therapy previous year	46·85		6·87			10·23			
On statins	22·65		23·58			23·07			
On LAMA	65·04		70·97			80·54			
On LABA	52·06		68·78			78·54			
On ICS	57·23		71·69			81·34			
	Mean (SD)	*Distribution*				*Mean (SD)*	*Distribution*		
Age, years	64·68 (8·75)		63·30 (6·99)			63·53 (6·90)			
Follow–up time, years	0·90 (0·23)		0·97 (0·12)			0·97 (0·13)			
FEV$_1$, % predicted	40·60 (15·93)		48·40 (16·39)			44·60 (15·88)			
SGRQ Score[b]	49·95 (16·72)		47·41 (18·22)			51·43 (17·03)			
BMI	27·53 (6·44)		26·55 (5·80)			26·21 (5·77)			

Rate of Exacerbations [c]	Rate	Distribution	Rate	Distribution	Rate	Distribution
Total	1·42	0123456	1·20	0123456	1·82	0123456
Severe/very severe	0·29	0123	0·27	0123	0·40	0123

Abbreviations: SD, standard deviation; COPD, chronic obstructive pulmonary disease; LAMA, long-acting muscarinic antagonist; LABA, long-acting beta antagonist; ICS, inhaled corticosteroids; FEV1, % predicted forced expiratory volume in 1 second using Hankinson's method[12]; SGRQ, St. George's Respiratory Questionnaire; BMI, body mass index.

[a] Data are presented as mean (SD) for continuous predictors and number of subjects (% of column total) for dichotomous predictors, except where noted.

[b] Between 0 and 100, with a higher score indicating worse status.

[c] Numbers in the % column show the annual rate of exacerbations (episodes/patient).

[d] Number of participants with the specified number of COPD exacerbations during follow–up.

Table 3 Model coefficients for the Joint Rate–Severity Prediction Model of COPD Exacerbations [a]

Predictor	Rate Component			Severity Component		
	Estimate ln(HR)	95%CI	p	Estimate ln(OR)	95%CI	p
Intercept	-0·009	-0·58, 0·56	0·974	-3·849	-5·54, -2·16	<·0001
Male	-0·152	-0·25, -0·05	0·003	0·377	0·08, 0·67	0·012
Age at baseline (per 10–year)	-0·018	-0·08, 0·05	0·576	0·109	-0·07, 0·29	0·240
Current smoker [b] at Baseline	-0·195	-0·32, -0·07	0·003	0·390	0·03, 0·75	0·034
Oxygen therapy [b] last year	0·085	-0·03, 0·2	0·162	0·538	0·2, 0·88	0·002
Baseline FEV_1 (% predicted)	-0·428	-0·79, -0·07	0·020	-1·119	-2·24, 0·01	0·051
SGRQ Score [c] (per 10–unit)	0·100	0·07, 0·13	<·0001	0·199	0·11, 0·29	<·0001
BMI (per 10–unit)	-0·123	-0·21, -0·04	0·004	-0·103	-0·36, 0·15	0·434
CV-indicated statins [b]	0·095	-0·03, 0·22	0·127	0·315	-0·03, 0·67	0·078
LAMA [b]	0·144	0·03, 0·25	0·011	-0·134	-0·45, 0·18	0·402
LABA [b]	0·118	-0·01, 0·24	0·066	0·012	-0·34, 0·36	0·946
ICS [b]	0·216	0·09, 0·34	0·001	0·376	0·03, 0·72	0·034
Random Effect Variance	0·60	0·51, 0·69	<·0001	2·385	1·63, 3·14	<·0001
Random Effect Covariance	0·147		0·172			

Abbreviations: CI, confidence interval; FEV1, % predicted forced expiratory volume in 1 second using Hankinson's method[12];; SGRQ, St. George's Respiratory Questionnaire; LAMA, long-acting muscarinic antagonist; LABA, long-acting beta antagonist; ICS, inhaled corticosteroids; BMI, body mass index.
[a] All p–values and confidence limits were computed from the final Hessian matrix
based on a t distribution with default degrees of freedom (number of subjects minus number of random effects) in SAS NLMIXED.
[b] Binary predictor for medication use in the previous 12 months.
[c] Between 0 and 100, with a higher score indicating worse status.

Table 4 Case study: Despite having the same exacerbation history, predicted severe exacerbation rate and risk for patients A is almost twice as patient B leading to different recommendation for roflumilast therapy.

Predictor	Patient A	Patient B
Sex	Male	Female
Age	57	57
Current smoker	No	No
Oxygen therapy last year	No	No
FEV_1 (% predicted)	51 (GOLD 2)	75 (GOLD 2)
CAT Score	29	10
BMI	18	30
CV-indicated statins	No	No
LAMA	Yes	Yes
LABA	Yes	Yes
ICS	Yes	Yes
Exacerbations in the last year	2	2
Severe Exacerbations in the last year	1	1
Predictions	**Patient A**	**Patient B**
Predicted Exacerbation Risk	%82	%79
Predicted Severe Exacerbation Risk	%28	%18
Predicted Exacerbation Rate	2.1	1.9
Predicted Severe Exacerbation Rate	0.6	0.3
Chance of Net Benefit from Roflumilast	80%-90%	10%-20%
Roflumilast Prescription	Recommended	Not Recommended

Figure 1 Flow diagram

Subgroup Calibration performance of ACCEPT

A) In all COPD patients in ECLIPSE Study

B) In COPD patients with a history of exacerbation in ECLIPSE Study

Figure 2 Calibration of exacerbation and severe exacerbation rate in subgroups of external dataset

Calibration performance of ACCEPT

A) In all COPD patients in ECLIPSE Study

B) In COPD patients with a history of exacerbation in ECLIPSE Study

Figure 3 Calibration plot comparing per decile predicted vs. observed rate of exacerbations in female (right) and male (left) participants in the external validation dataset. Perfect agreement is shown by the dashed line.

Discrimination performance of ACCEPT vs. event history

A) In all COPD patients in ECLIPSE Study

B) In COPD patients with a history of exacerbation in ECLIPSE Study

Figure 4 Receiver operating characteristic (ROC) curves showing discriminative performance of ACCEPT compared with last year's history of exacerbation among patients with a prior history of exacerbation and all patients in the

external validation dataset. In line with GOLD recommendations, Area-under-the-curve (AUC) is shown for predicting at least two exacerbations and at least of severe exacerbation. DeLong's test for two correlated ROC curves was used to produce p-values.

References:

Aaron SD. Management and prevention of exacerbations of COPD. *BMJ* 2014; **349**: g5237.

Hurst JR, Vestbo J, Anzueto A, *et al.* Susceptibility to Exacerbation in Chronic Obstructive Pulmonary Disease. *N Engl J Med* 2010; **363**: 1128–38.

Global Initiative for Chronic Obstructive Lung Disease. POCKET GUIDE TO COPD DIAGNOSIS, MANAGEMENT, AND PREVENTION: A Guide for Health Care Professionals. *Glob Initiat Chronic Obstr Lung Dis* 2019. https://goldcopd.org/wp-content/uploads/2018/11/GOLD-2019-POCKET-GUIDE-DRAFT-v1.7-14Nov2018-WMS.pdf.

Obeidat M, Sadatsafavi M, Sin DD. Precision health: treating the individual patient with chronic obstructive pulmonary disease. *Med J Aust*; **0**. DOI:10.5694/mja2.50138.

Celli BR, Cote CG, Marin JM, *et al.* The Body-Mass Index, Airflow Obstruction, Dyspnea, and Exercise Capacity Index in Chronic Obstructive Pulmonary Disease. *N Engl J Med* 2004; **350**: 1005–12.

Guerra B, Gaveikaite V, Bianchi C, Puhan MA. Prediction models for exacerbations in patients with COPD. *Eur Respir Rev* 2017; **26**: 160061.

Collins GS, Reitsma JB, Altman DG, Moons K. Transparent reporting of a multivariable prediction model for individual prognosis or diagnosis (TRIPOD): the TRIPOD Statement. *BMC Med* 2015; **13**: 1.

Albert RK, Connett J, Bailey WC, *et al.* Azithromycin for prevention of exacerbations of COPD. *N Engl J Med* 2011; **365**: 689–98.

Criner GJ, Connett JE, Aaron SD, *et al.* Simvastatin for the Prevention of Exacerbations in Moderate-to-Severe COPD. *N Engl J Med* 2014; **370**: 2201–10.

Aaron SD, Vandemheen KL, Fergusson D, *et al.* Tiotropium in Combination with Placebo, Salmeterol, or Fluticasone–Salmeterol for Treatment of Chronic Obstructive Pulmonary Disease: A Randomized Trial. *Ann Intern Med* 2007; **146**: 545.

Agusti A, Calverley PM, Celli B, *et al.* Characterisation of COPD heterogeneity in the ECLIPSE cohort. *Respir Res* 2010; **11**: 122.

Hankinson JL, Odencrantz JR, Fedan KB. Spirometric reference values from a sample of the general U.S. population. *Am J Respir Crit Care Med* 1999; **159**: 179–87.

Morishita-Katsu M, Nishimura K, Taniguchi H, *et al.* The COPD assessment test and St George's Respiratory Questionnaire: are they equivalent in subjects with COPD? *Int J Chron Obstruct Pulmon Dis* 2016; **11**: 1543–51.

Sadatsafavi M, Sin DD, Zafari Z, *et al.* The Association Between Rate and Severity of Exacerbations in Chronic Obstructive Pulmonary Disease: An Application of a Joint Frailty-Logistic Model. *Am J Epidemiol* 2016; **184**: 681–9.

Ewout Steyerberg. Clinical Prediction Models - A Practical Approach to Development, Validation, and Updating. Springer New York, 2009.

Steyerberg EW, Vickers AJ, Cook NR, *et al.* Assessing the Performance of Prediction Models: A Framework for Traditional and Novel Measures. *Epidemiology* 2010; **21**: 128–38.

DeLong ER, DeLong DM, Clarke-Pearson DL. Comparing the areas under two or more correlated receiver operating characteristic curves: a nonparametric approach. *Biometrics* 1988; **44**: 837–45.

Almagro P, Soriano JB, Cabrera FJ, *et al.* Short- and Medium-term Prognosis in Patients Hospitalized for COPD Exacerbation: The CODEX Index. *Chest* 2014; **145**: 972–80.

Bertens L, Reitsma, Moons, *et al.* Development and validation of a model to predict the risk of exacerbations in chronic obstructive pulmonary disease. *Int J Chron Obstruct Pulmon Dis* 2013; : 493.

Kerkhof M, Freeman D, Jones R, Chisholm A, Price D. Predicting frequent COPD exacerbations using primary care data. *Int J Chron Obstruct Pulmon Dis* 2015; : 2439.

Annavarapu S, Goldfarb S, Gelb M, Moretz C, Renda A, Kaila S. Development and validation of a predictive model to identify patients at risk of severe COPD exacerbations using administrative claims data. *Int J Chron Obstruct Pulmon Dis* 2018; **Volume 13**: 2121–30.

FitzGerald JM, Sadatsafavi M. Improving precision in the prediction of asthma exacerbations. *Lancet Respir Med* 2017; **5**: 539–40.

Fuhlbrigge AL, Bengtsson T, Peterson S, *et al.* A novel endpoint for exacerbations in asthma to accelerate clinical development: a post-hoc analysis of randomised controlled trials. *Lancet Respir Med* 2017; **5**: 577–90.

Yu T, Fain K, Boyd CM, *et al.* Benefits and harms of roflumilast in moderate to severe COPD. *Thorax* 2014; **69**: 616–22.

Moons KGM, Kengne AP, Grobbee DE, *et al.* Risk prediction models: II. External validation, model updating, and impact assessment. *Heart Br Card Soc* 2012; **98**: 691–8.

Adibi A, Sadatsafavi M, Hill A. accept: The Acute COPD Exacerbation Prediction Tool (ACCEPT). 2019 https://CRAN.R-project.org/package=accept.