Strain-level diversity impacts cheese rind microbiome assembly and function

Brittany A. Niccum[1], Erik K. Kastman[1], Nicole Kfoury[2,3], Albert Robbat[2,3], Benjamin E. Wolfe[1,3]

[1]Tufts University, Department of Biology, Medford, MA, USA
[2]Tufts University, Department of Chemistry, Medford, MA, USA
[3]Tufts University Sensory and Science Center, Medford, MA, USA

*Correspondence: benjamin.wolfe@tufts.edu

ABSTRACT Taxa that are consistently found across microbial communities are often considered members of a core microbiome. One common assumption is that taxonomically identical core microbiomes will have similar dynamics and functions across communities. However, strain-level genomic and phenotypic variation of core taxa could lead to differences in how core microbiomes assemble and function. Using cheese rinds, we tested whether taxonomically identical core microbiomes isolated from distinct locations have similar assembly dynamics and functional outputs. We first isolated the same three bacterial species (*Staphylococcus equorum, Brevibacterium auranticum*, and *Brachybacterium alimentarium*) from nine cheeses produced in different regions of the United States and Europe. Comparative genomics identified distinct phylogenetic clusters and significant variation in genome content across the nine core microbiomes. When we assembled each community in initially identical compositions, community structure diverged over time resulting in communities with different dominant taxa. The core microbiomes had variable responses to abiotic (high salt) and biotic (the fungus *Penicillium*) perturbations, with some communities showing no response and others substantially shifting in community composition. Functional differences were also observed across the nine core communities, with considerable

variation in community pigment production (light yellow to orange) and composition of

volatile organic compound profiles emitted from the rind communities (nutty to sulfury).

Our work demonstrates that core microbiomes isolated from independent communities

may not function in the same manner due to strain-level variation of core taxa. Strain-

level diversity across core cheese rind microbiomes may contribute to variability in the

aesthetics and quality of surface-ripened cheeses.

INTRODUCTION

Metagenomic surveys of microbial communities often describe the existence of

core microbiomes. Although many definitions currently exist (1), core microbiomes are

generally considered to be the set of microbial taxa that are commonly found across all

(or many) sampled microbial communities. Many microbial communities, from plant

roots to wastewater treatment plants, contain a set of core microbiome components that

are common, highly abundant, and functionally significant (1–4). These core

microbiomes can range from just a few species to tens or hundreds of species. For

example, skin microbiomes of most humans are dominated by very similar

Corynebacterium, *Propionibacterium*, and *Staphylococcus* species (3, 5, 6).

One common and largely untested assumption about core microbiomes is that

taxonomically identical core microbiomes will have similar community assembly patterns

and functions. More specifically, when comparing 16S rRNA gene sequencing surveys,

samples that have very similar compositions of 16S sequences are often assumed to

have similar functional potentials. This assumption underlies the development of

taxonomy-based microbiome diagnostics and tools to predict function from taxonomic

sequences (7, 8). But independent evolution or coevolution of microbial species within

communities may generate previously underappreciated functional diversity across core

microbiomes. It is widely accepted that microbial genomes are highly variable within

species due to rapid rates of evolution and potential for lateral gene transfer (9–11).

Moreover, we know from decades of work in microbial ecology, physiology, and

genomics that there is considerable within species trait variation in microbes (12–14).

For example, a set of 11 strains of *Brevundimonas alba* isolated from the same

freshwater habitat had identical 16S rRNA sequences, but highly divergent carbon

utilization profiles and growth rates (15). This intraspecific trait diversity could be

ecologically significant, but the impact of strain-level diversity on core microbiome

assembly and function is poorly understood (16).

Cheese rinds provide an ideal opportunity to test whether taxonomically identical

core microbiomes have similar assembly dynamics and functions and more generally

the causes and consequences of core microbiome diversification. Rinds form on the

surfaces of cheeses aged in an aerobic environment and are composed of bacteria,

yeasts, and filamentous fungi (17–19). Our previous work used amplicon and shotgun

metagenomics to describe the bacterial and fungal diversity of 137 cheese rinds from

the United States and Europe (17). Three bacterial genera - *Staphylococcus*,

Brevibacterium, and *Brachybacterium* - were the most frequently detected across

cheese rinds, and can be considered a core microbiome for cheese rinds. Many

cheeses are made for decades in a single location and re-circulating populations of

microbes in cheese facilities could evolve during this time. Through variation in abiotic

and biotic selection pressures applied during cheese production and aging, including

abiotic (salinity, pH, resource availability) and biotic (presence of bacterial and fungal

neighbors), core cheese microbiomes have the potential to evolve new genotypes and

phenotypes with divergent functions.

Here we characterize core microbiome members across cheese rind

communities and determine the consequences of core microbiome diversification for

community assembly and function. We isolated the same three species of bacteria -

Staphylococcus equorum (hereafter *Staphylococcus*)*, Brevibacterium auranticum*

(hereafter *Brevibacterium*), and *Brachybacterium alimentarium* (hereafter

Brachybacterium) - from nine different cheeses made across the United States and

Europe (**Fig. 1A-B**). We refer to these nine sets of three co-isolated bacterial species as

taxonomically identical core microbiomes throughout the rest of the paper (**Fig. 1C**). We

chose these species because they represent the most common species of the three

most abundant bacterial genera in cheese rinds (17, 20). These species also commonly

enter the dairy environment from the raw milk used for cheese production and therefore

have the potential to co-occur and adapt to abiotic and biotic conditions within local

cheese production facilities (21–23). Each species has a distinct colony morphology

(**Fig. 1B**) making it easy to track composition in experimental communities. We

predicted that intraspecific variation of core microbiome members across cheese rind

communities would cause differences in community structure over time. We also

predicted that strain-level differences across core microbiomes would result in

differences in community functions relevant for cheese aging, including pigmentation of

the cheese rind biofilm and the production of aroma compounds.

RESULTS

Variation in genome content across taxonomically identical core microbiomes

To determine genomic variation across the nine taxonomically identical core

microbiomes, we constructed draft genomes of each strain (**Table S1**). We used single-

nucleotide polymorphisms (SNPs) in the core genes shared across all nine communities

to determine phylogenomic divergence of each of the core communities (24). We then

determined variation in functional gene content across the nine core communities using

PGAP (25). For functional gene content analysis, we focused on accessory genes that

were uniquely present in only one community as these genomic traits may help drive

divergence in core microbiome functions.

Across the nine communities, 8,069 gene clusters were shared among all three

species, making up the core metagenome of these communities. Using SNPs identified

in this core metagenome with PanSeq, clear phylogenomic divergence across the nine

cheese communities was apparent (**Fig. 2**). C1 was distant from the other eight core

microbiomes, driven by the highly divergent *Staphylococcus* genome in this community.

The eight other core microbiomes clustered into two broad phylogenomic groups: one

containing C6 and C2, and the other containing the remaining six communities (**Fig. 2**).

We next assessed the total number of unique accessory gene clusters across the

nine communities. Distribution of unique accessory gene content was highly variable,

ranging from 246 (C5) to 630 genes (C3) (**Fig. 2**, **Table S2**). Variability in the

abundance of accessory gene clusters was most prominent in *Staphylococcus* (ranging

from 36-280 unique gene clusters across strains) and *Brevibacterium* (ranging from 72-

213 unique gene clusters), suggesting that these taxa have the most dynamic

accessory gene content in the cheese rind core metagenome.

Several biological processes were significantly enriched in core communities

(**Table S3**). C3 had the most diverse enrichment of SEED categories, with

overrepresentation of genes in potassium metabolism, carbohydrates, and DNA

metabolism.   Protein   metabolism   and   phages/prophages/transposable

elements/plasmids were overrepresented in C4. In C2, the accessory genome was

significantly enriched with stress response genes. Carbohydrate-related genes were

enriched in the C6 core microbiome. Some of these unique accessory genes could be

functionally significant in the cheese rind environment. For example, *Brevibacterium* of

C3 has a unique potassium transport system with high similarity to the *kdfABCF* operon

(**Table S2**) which is known to play roles in salt stress in bacteria (26).

Collectively these genomic data demonstrate that taxonomically identical core

microbiomes isolated from distinct cheeses are phylogenomically diverse and have

variable genome content. Although the presence/absence of genes does not indicate

actual functional potential of microbes, these comparative genomic data suggested to

us that there could be divergence in how each taxa functioned within each community

and how they responded to perturbations.

Community assembly dynamics vary across taxonomically identical core

microbiomes

We next determined whether strain-level differences impacted how the cheese

rind communities assemble. A typical succession in our lab model involves early

colonization of *Staphylococcus* which can tolerate the low pH (5.0-5.2) of the cheese

curd, followed by growth of *Brachybacterium* and ending with dominance by

Brevibacterium (17, 27). We predicted two different potential impacts of strain-level

variation on community assembly. In one scenario, distinct strains of *Staphylococcus*,

Brachybacterium, and *Brevibacterium* across the nine communities may vary in genome

content or growth rates in isolation, but these differences may be too minor to impact

the dynamics of assembly of the three-member community. In this case, we expected

nearly identical community composition across the different core microbiomes as strains

of each species behaved similarly. Alternatively, strain-level differences may translate

into differences in interactions with other community members or rates of growth within

the community succession. In this scenario, we expected to observe reproducible

changes in the composition of the communities as they assembled and differences in

functional outputs.

To determine how strain-level differences across communities impact assembly

dynamics, we used *in vitro* community assembly assays to measure total colony forming

units (CFU) and community composition (relative abundance of each species) (**Fig. 3A**).

Communities were quantified at three and ten days after inoculating equal amounts of

each of the three bacterial species on the surface of cheese curd agar. Our previous

work has demonstrated that this assay mimics *in situ* community dynamics (17, 27). We

acknowledge that real cheese rind communities would develop over much longer time

scales (weeks to months). In the context of this work, we are using the community

assembly assay in a standardized environment to demonstrate the *potential* for

divergence in community assembly.

At both three and ten days of community assembly, there were nearly no

differences in total community abundance as measured by combined CFU of all three

species (**Fig 3B**, Day 3 ANOVA $F_{8,81}$ = 2.07, p = 0.05; Day 10 ANOVA $F_{8,79}$ = 0.46, p =

0.88). However, there were substantial differences in community composition across the

nine core communities (Day 3 permutational multivariate analysis of variance

[PERMANOVA] F = 4.005, p = 0.0001; Day 10 PERMANOVA F = 5.57, p = 0.0001).

Many communities (C1, C2, C6, C7) were dominated by *Brevibacterium* at the end of

succession (**Fig 3C**). Some communities had a relatively even mix of all three species

(C5, C3, C8, and C9). Community C4 had a highly dissimilar structure with a high

abundance of *Brachybacterium* at the end of succession and a low abundance of

Brevibacterium.

A simple explanation for differences in community composition across the nine

core communities is that individual bacterial strains have different growth abilities alone

and in the community. Those taxa and strains that grow best alone and with the

community present should be the most abundant members of the community. To test

this, we determined total growth of each of the 27 strains alone on cheese curd agar

and compared growth alone after ten days to growth in the community. All

Staphylococcus species grew well alone and had limited responses to growth in the

community (**Fig. 3D**). Two strains were slightly stimulated by growth in the community

(C5 and C7) and one was slightly inhibited (C6). In contrast to the relatively even growth

of the *Staphylococcus*, the *Brevibacterium* strains had variable growth alone across the

nine core communities. Four of the *Brevibacterium* strains grew poorly by themselves

on cheese curd agar (C2, C5, C8, and C9) and were strongly stimulated by growth in

the community. One *Brevibacterium* strain (C4) was inhibited by growth in the

community. All *Brachybacterium* strains grew well on cheese curd by themselves and

were generally inhibited when grown in the community.

For all three taxa, mean growth alone was a very poor predictor of mean relative

abundance in the community for each of the three species (*Staphylococcus* r^2 = 0.166,

p= 0.276; *Brevibacterium* r^2 = 0.001, p = 0.923; *Brachybacterium* r^2 = 0.020, p = 0.716).

A somewhat better predictor of mean relative abundance was how growth of each strain

was impacted by the community (*Staphylococcus* r^2 = 0.672, p<0.01; *Brevibacterium* r^2

= 0.013, p = 0.773; *Brachybacterium* r^2 = 0.319, p = 0.113). This suggests that

interactions between each of the strains and their communities may contribute to

differences in community composition across the nine core microbiomes. For example,

the inhibition of *Brevibacterium* and lack of inhibition of *Brachybacterium* in C4 may

partly explain why this community was the only to be dominated by *Brachybacterium*.

Variation in responses to abiotic and biotic perturbation across core

microbiomes

Core microbiomes may experience abiotic or biotic perturbations that could alter

community assembly and function. We predicted that if individual core members have

evolved different responses to stress or if the communities have coevolved stress-

response mechanisms, taxonomically identical core microbiomes may have divergent

responses to perturbations. Two major perturbations in cheese rind core microbiomes

are salt and interactions with fungi (17, 20, 28). Salt concentrations are initially high on

the surface of fresh cheese because salt is applied to the cheese surface or via a brine

(29). The salt diffuses into the cheese and eventually equilibrates to around 3% salt in

the rind environment of many cheeses. Core cheese rind microbiomes also experience

interactions with fungi, ranging from yeasts (e.g. *Debaryomyces* and *Galactomyces*

species) to molds (*Fusarium*, *Scopulariopsis*, and *Penicillium* species) (17, 20, 30).

Penicillium species are widespread in cheese rinds and can strongly inhibit diverse

cheese rind bacteria (17, 27, 31) potentially through the production of secondary

metabolites or other mechanisms.

To determine how the nine core microbiomes respond to salt and fungal

perturbations, we used the same community assembly assay described above with the

addition of two treatments: a 6% NaCl treatment and a +*Penicillium* treatment. We used

a strain of *Penicillium* that was isolated from a natural rind cheese and was previously

demonstrated to inhibit cheese rind bacterial growth (17). Across isolates of all three

taxa, both the 6% NaCl and +*Penicillium* treatments caused a general decrease in total

growth across all nine core microbiomes with +*Penicillium* causing stronger growth

inhibition (**Fig. 4A**). Communities had variable responses to the two perturbations. The

Penicillium perturbation caused the most significant shifts in community composition

with six out of nine core communities showing significant changes in community

composition (**Fig. 4B**-**C**). In some communities, *Penicillium* caused a major increase in

Brachybacterium relative abundance (C2 and C3). In other communities, *Penicillium*

caused an increase in the relative abundance of *Staphylococcus* (C1, C8, and C9). The

6% salt treatment caused fewer shifts in community composition with only two

communities (C5 and C6) responding to the higher salt environment. In both cases,

Brevibacterium increased in relative abundance.

Strain-level diversity of cheese rind core microbiomes drives divergent pigment and aroma production

Our experiments above demonstrate that strain-level diversity of the core cheese rind taxa drives divergence in community composition across the nine core microbiomes. Does this divergence lead to cheeses with different properties that could be perceived by consumers? Differences in community composition may not necessarily translate into differences in functional outputs. Many studies of the microbiome have suggested that communities with different compositions may have similar functions due to functional redundancy across community members (32–34). While our comparative genomic analysis above suggested potential functional differences across the cheese communities, many of the core community functions were conserved in the core genome and variation in accessory genes may have little impact on community functions. To determine whether divergence in composition of the core microbiomes can also translate into differences in functional outputs, we measured two important traits of cheese rind microbiomes: rind color and volatile organic compound (VOC) production.

Cheese rind bacteria define how the cheese appears to customers through the production of cellular pigments such as carotenoids or the secretion of pigmented extracellular metabolites into the curd (35–39). The three bacteria in our model community produce distinct pigments (**Fig. 1B**) and shifts in their relative abundance could translate into changes in rind color. Using a colorimeter, we measured the surface color of our model cheese communities aged for 10 days. Communities had significantly different color development (ANOVA $F_{9,39} = 524.9$, p <0.0001), with C3, C4, C6, C7, and

C9 having significantly greater a* values compared to the control, indicating more red

pigmentation (**Fig. 5A**). All communities had significantly greater b* values compared to

the control (ANOVA $F_{9,39}$ = 139.6, p <0.0001), with C3 and C4 having the greatest

values and appearing the most orange (**Fig. 5A**).

As the rind biofilm decomposes fats, proteins, and other components of the

cheese substrate, a diversity of VOCs are produced that are perceived by consumers

as aromas (40–42). Using headspace sorptive extraction followed by analysis with gas

chromatography-mass spectrometry (GC-MS)(43, 44), we quantified VOCs produced by

each community after 10 days of cheese rind development. Across all nine communities

248 unique VOCs were detected with significant differences in the mean VOCs per

community (**Fig. 5B,** ANOVA $F_{8,35}$ = 28.9, p <0.0001). The composition of VOCs across

the nine cheese communities was significantly different (**Fig. 5C**, PERMANOVA F =

62.38, p < 0.001, **Table S4**). Using a SIMPER analysis, nine compounds contributed

greater than 1% to the average overall Bray-Curtis dissimilarity: benzyl methyl ketone

(27% contribution; odor = floral/fruity odor), tetramethylpyrazine (19%; odor =

nutty/musty/chocolate/coffee),   2,5-dimethylpyrazine   (13%;   odor   =

nutty/musty/chocolate/coffee),   trimethylpyrazine   (12%;   odor   =

nutty/musty/chocolate/coffee), dimethyl disulfide (9%; odor = sulfurous/cabbage/onion),

dimethyl trisulfide (2%; odor = sulfurous/cabbage/onion), 2,6-diethylpyrazine (2%; odor

= nutty/musty/chocolate/coffee), unknown compound 520 (1%; odor unknown), and 3-

hydroxy-2-butanone (1%; odor = sweet/buttery/creamy). C5 had the most distinct VOC

profile of all communities with high amounts of tetramethylpyrazine, trimethylpyrazine,

and 3-hydroxy-2-butanone and low amounts of the major sulfur compounds, suggesting

a more nutty and buttery aroma profile.

DISCUSSION

Using taxonomically identical three-member communities isolated from nine

distinct cheeses, our work demonstrates the significance of strain-level variation for

microbiome community assembly and function. Studies of plant and animal

communities have demonstrated that intraspecific genetic and phenotypic diversity can

impact community assembly and function (45–47). Here we demonstrate that

intraspecific diversity of taxonomically identical core microbiome members can impact

the relative abundance of community members as well as functional outputs of the

communities. Many communities did converge on a similar composition despite having

substantial variation in accessory gene content. But several communities had

substantially different structures and functions despite starting off with identical

compositions. Some communities had relatively even coexistence of the three

community members, while others were dominated by either *Brevibacterium* or

Brachybacterium. The divergence was not due to stochastic community assembly

across replicates as we observed highly reproducible community structures across

replicate experiments.

The goal of this work was to determine whether taxonomically identical core

microbiomes have similar community dynamics and functions. The limited number of

core communities (nine) makes it difficult to pinpoint specific ecological or genetic

mechanisms that may be underlying the observed differences across the core

communities. One simple explanation for the dominance of different taxa across the

core microbiomes is differences in growth of individual strains. Our experiments

comparing growth alone versus in the community demonstrates variable growth rates

and interactions with the community for each of the three taxa. However it does not fully

explain community structure. For example, in C4 where *Brachybacterium* dominated,

the *Brachybacterium* strain had a similar levels of growth alone and interactions with

community members as other communities where *Brevibacterium* dominated (e.g. C5

and C6). Future work exploring the roles of inhibitory and cooperative interactions will

pinpoint specific mechanisms explaining the variable community assembly dynamics of

cheese rind core microbiomes.

The evolutionary processes that have generated the divergent species and

community-level responses of our core cheese microbiomes are currently unknown. It is

possible that each core microbiome has experienced different evolutionary histories in

each cheese production environment. As new batches of cheese are introduced to a

cave environment, communities may be repeatedly transferred to these new cheeses.

This repeated colonization of the cheese substrate could allow each of the core

microbiomes to evolve collectively as a community in the individual production

environments (48). Each environment may have unique abiotic selection pressures,

including salt concentrations, milk composition, temperature, that could shape the

evolutionary trajectories of these communities. The core microbiomes could also

experience highly divergent biotic environments. For example, these core communities

were isolated from cheeses with variable fungal environments, ranging from yeast to

filamentous fungi (17). Future work using experimental evolution to attempt to create

divergent communities from an ancestral core microbiome could begin to understand

the drivers of core microbiome diversification.

Our model communities represent the widespread bacterial taxa found in cheese

rinds. We acknowledge that these communities have several constraints that may

impact translation of our results to other systems. First, our communities only had three

bacterial species. While some widespread microbiomes have low species diversity (5,

49), many microbiomes have much higher levels of diversity. Would taxonomically

identical core microbiomes with higher taxonomic diversity also demonstrate divergence

in assembly and functions? With increasing potential for higher-order interactions and a

higher number of potential functions with increasing species diversity, we predict that

increasing diversity may lead to even more divergent communities. Our model

communities also used a single strain of each species within each core microbiome. In

constructing our communities, we chose to ignore potential intraspecific variation within

each of the nine core communities and assumed that the isolated taxa represented the

most common genomic type of the species within each of the core communities.

Metagenomic sequencing studies have identified multiple co-existing strains of the

same microbial species (3, 16, 56–60) and these strains may interact with each other

and other community members to impact community composition. It would be

fascinating to see how including intraspecific diversity within core microbiomes may

impact community assembly and function.

In a large amplicon-sequencing study of cheese rind microbiomes, we

demonstrated that taxonomically identical cheese rind communities could form in very

different cheesemaking regions (17). This was surprising given that these cheeses have

divergent sensory properties. Many of these differences could be driven by ingredients,

length of aging, or other cheese processes. Our current findings suggest that the

variability in the qualities of surface-ripened cheeses could also be driven by strain-level

differences across the cheese communities. We acknowledge that our lab cheese rinds

are not real cheeses and only represent potential patterns of cheese rind community

assembly. But it is very likely that the differences observed across the nine core

microbiomes would translate to actual cheese production. Previous studies of fermented

food microbes have pointed out strain-level differences of individual species used in

fermented foods (50–54), but studies demonstrating the functional significance of strain-

level variation at the community level are rare (55). To help preserve the unique

identities of cheeses made in specific regions, it may be helpful for cheese producers to

identify the unique genomic and functional properties of their core microbiomes and

maintain these communities.

More broadly, our work in these model microbiomes may have implications for

both the design and management of core microbiomes in other systems. First, our work

demonstrates that taxonomic profiling of microbiomes may not provide useful predictors

of assembly dynamics and functions. Amplicon based approaches of sequencing

microbiomes, such as using 16S rRNA gene sequencing, only capture high-level

taxonomic diversity. As we have demonstrated, taxonomically similar communities can

have very different dynamics. Fortunately, microbiome sequencing studies are moving

toward shotgun-metagenomic approaches that could capture the strain-level diversity

that we observed across our nine communities (3, 16, 56–60). Our work also suggests

that it might be hard to predict microbiome responses to disturbances using taxonomic

profiles alone. For example, across individuals that have similar skin core microbiomes,

responses to environmental stresses such as antibiotics may depend on the specific

strains and genomic content of the core communities. Finally, when designing synthetic

microbiomes, our work suggests that the individual 'parts' (strains of species) may alter

desired outcomes.

METHODS

Isolation and maintenance of core microbiome members

Frozen glycerol stocks of communities initially characterized using metagenomic

sequencing (Wolfe et al. 2014) were plated out on plate count agar with milk and salt

(PCAMS) to culture bacteria. Colonies with morphotypes that had the appearance of

one of the three target species were streaked from single colonies. *Staphylococcus*

equorum colonies are usually fast-growing, smooth, medium-sized, flat, and white or

light golden in color. *Brevibacterium auranticum* colonies are usually slow-growing,

medium-sized, and orange. *Brachybacterium alimentarium* colonies have medium

growth rates, are large and flat, and are yellow-green in color. Initial identification of the

isolates was done using the 16S rRNA region using primers 27f and 1492r.

Comparative genomics

The genome of each bacterial strain was sequenced, assembled, and annotated

as we previously described for *Staphylococcus* species (27). Briefly, DNA was extracted

using MoBio PowerSoil DNA extraction kits from streaks generated from a single bacterial

colony grown for one week on PCAMS. Approximately 1 µg of purified gDNA was sheared

using a Covaris S220 to approximately 450 base pair lengths and was used as the input for

a New England Biolabs (Ipswitch, MA) NEBNext Ultra DNA Library Prep Kit for Illumina.

Libraries were spread across multiple sequencing lanes with other projects and were

sequenced using 100 base-pair, paired-end reads on an Illumina HiSeq 2500.

Approximately 10 million reads were sequenced for each genome. Failed reads were

removed from libraries and reads were trimmed to remove low quality bases and were

assembled to create draft genomes using the *de novo* assembler in CLC Genomics

Workbench 8.0. Assembled genomes were annotated using RAST(61). All genome

assemblies have been deposited in NCBI (accession numbers in Table S1).

To identify phylogenomic relationships between each of the nine core

communities, we used PanSeq (24) to identify SNPs across the core genome of each of

the nine genomes for each of the three species. A SNP file for each species from each

community was then concatenated together to create a community SNP file. RAxML

8.2.11 (with GTR GAMMA nucleotide model and 100 bootstrap replicates) was used to

create a maximum likelihood phylogeny of the nine communities using the SNP file.

To compare the presence and absence of genes across strains and species,

core and accessory genes were identified by assigning protein-coding sequences to

functionally orthologous groups using the MultiParanoid method of the PanGenome

Analysis Pipeline (PGAP) (Zhao et al, 2012). Species-to-species orthologs were

identified by pairwise strain comparison using BLAST with PGAP defaults: a minimum

local coverage of 25% of the longer group and a global match of no less than 50% of

the longer group, a minimum score value of 50, and a maximum E value of 1E−8.

Multistrain orthologs were then found using MultiParanoid (80). Enrichment of SEED

subsystem categories in each of the nine core communities was determined using

Fisher's exact test with false-discovery rate correction.

Community assembly assays

To measure assembly of the distinct core communities, approximately 20,000

CFU of each species was inoculated on the surface of 150 µL of cheese curd agar (3%

salt) distributed into replicate wells of a 96-well plate, as previously described (17, 27).

Communities were incubated aerobically at 24°C in the dark, and harvested at 3 and 10

days after inoculation, which represent early and late community succession (17). To

determine community composition of individual replicate communities, the community

was pestled in 600 µL of 1X phosphate buffered saline, serially diluted, and plated onto

PCAMS. PCAMS plates were incubated for a week before counting the abundance of

each bacterial species. To measure growth alone, the same density of CFU of each

taxa alone was inoculated into wells. Five technical replicates of each community were

performed in each of two experimental replicates.

Salt (6%) and fungal (+*Penicillium*) perturbation experiments were conducted

using the same community assembly assay, but with 6% salt cheese curd agar or with

the addition of *Penicillium. Penicillium* strain #12, isolated from a natural rind cheese in

Vermont, was used in these experiments. We used this strain because it was isolated

from a cheese where the *Staphylococcus*, *Brachybacterium*, and *Brevibacterium* were

also found and it was used in previous experiments in our lab (27, 31). The exact

species identification of this mold is unknown, but it belongs to section *Fasciculata* with

other cheese *Penicillium* species. *Penicillium* was inoculated at an initial density of 2000

CFUs. Community composition in these experiments was determined as described

above except that cycloheximide was added to PCAMS plates used for bacterial

community isolation to eliminate fungal growth.

Color and VOC analyses

To measure rind color and VOC production, we constructed larger versions of

each of the nine core communities on cheese curd agar poured into Petri dishes (60mm

wide) to allow for a larger sampling area. To construct the rind communities, 600,000

CFU of each species was inoculated across the surface of the cheese curd agar.

Experimental cheeses were incubated for 10 days in the dark at 24°C before color and

VOC analyses.

To measure differences in color of the experimental cheeses, we used a CTI

A6CTI10 spectrocolorimeter. This handheld colorimeter uses the CIELAB color space to

quantify both lightness (L*) and two chromatic coordinates (a* and b*). Similar

colorimeters have been used to quantify cheese rind color (62). Higher values of a*

(a*+) indicate red colors while lower values (a*-) indicate green colors. Higher values of

b*(b*+) indicate yellow while lower values (b*-) indicate blue colors. Colorimeter

readings were taken by placing a 30mm Petri dish lid upside down on the middle of the

surface of the rind and then placing the colorimeter on the Petri dish surface. This was

done to protect the colorimeter from the sticky rind surface and to avoid cross-

contamination across replicates.

Cheese volatiles were collected from experimental cheese rinds by headspace

sorptive extraction (HSSE) using a polydimethylsiloxane (PDMS) coated magnetic stir-

bar. HSSE is an equilibrium-driven, enrichment technique in which 10mm long x 0.5 mm

thick stir-bars, Twister^{TM} (Gerstel), were suspended 1 cm above the sample by placing a

magnet on the top side of the collection vessel cover. Five replicates of each culture

were sampled for four hours. After collection, the stir-bar was removed and spiked with

10 ppm ethylbenzene-d$_{10}$, an internal standard obtained from RESTEK (Bellefonte, PA).

Organics were introduced into the gas chromatograph/mass spectrometer (GC/MS) by

thermal desorption. In addition to Twister blanks, analysis of the cheese curd agar

media was made to ensure background interferents were minimal. If present, these

compounds were subtracted from the data.

Analyses were performed using an Agilent (Santa Clara, CA) 7890A/5975C

GC/MS equipped with an automated multi-purpose sampler (Gerstel). The thermal

desorption unit (TDU, Gerstel) provided splitless transfer of the sample from the stir bar

into a programmable temperature vaporization inlet (CIS, Gerstel). The TDU was

heated from 40°C (0.70 min) to 275°C (3 min) at 600°C/min under 50ml/min of helium.

After 0.1 min the CIS, operating in solvent vent mode, was heated from -100°C to 275°C

(5 min) at 12°C/s. The GC column (30 m x 250 µm x 0.25 µm HP5-MS, Agilent) was

heated from 40°C (1 min) to 280 °C at 5°C/min with 1.2 mL/min of constant helium flow.

The MS was scanned from 40 to 350 *m/z*, with the EI source at 70 eV. A standard

mixture of C7 to C30 n-alkanes (Sigma–Aldrich, St. Louis, MO) was used to calculate

the retention index (RI) of each compound in the sample.

The Ion Analytics (Gerstel) spectral deconvolution software was used to analyze

the GC/MS data (63, 64).   A target/nontarget approach was employed where a

previously constructed database is used to deconvolve target compounds in the

sample. Once found, each compound's mass spectrum was subtracted from the total

ion current (TIC) signal. Each resulting peak scan was inspected to determine if the

residual ion signals were constant (±20%) or approximated background noise. If

constant, the software recorded the retention time, mass spectrum, 3-5 target ions and

their relative abundances into the database. Then the sample data is compared to

reference compound data in the database, viz., RI and MS (positive identification), or to

commercial libraries and literature (tentative identification). Once assigned, the

compound name, CAS#, and RI was added to the database. If neither positive nor

tentative identification could be made, a numerical identifier along with the same GC/MS

information was uploaded manually into the method. In contrast, if peak scans differed

(unresolved peak), the software searched for 3-5 invarient scans, averaged their

spectra, and then subtracted the average spectrum from the TIC signal. This process

was repeated until the residual signal at each scan approximated background noise. If

peak signals failed to meet the user-defined criterion below, no additional information

was obtained.

Four parameters were chosen as compound acceptance criterion. First, peak

scans must be constant for five or more consecutive scans (differences ≤ 20%) after

deconvolution. Second, the scan-to-scan variance (SSV or relative error) must be < 5.

The SSV calculates relative error by comparing the mass spectrum at each peak scan

against another. The smaller the difference, the closer the SSV is to zero, the better the

spectral agreement. Third, the Q-value must be ≥ 93. The Q-value measures the total

ratio deviation of the absolute value of the expected minus observed ion ratios divided

by the expected ion ratio times 100 for each ion across the peak. The closer the value is

to 100, the higher the certainty between sample and reference, library, and/or literature

spectra. Finally, the Q-ratio compares the ratio of the main ion intensity to confirming ion

intensities across the peak; it also must be ≤ 20%. These criteria form a single criterion

used for compound identification.

Statistical Analyses

To determine differences in community composition with all core microbiome

experiments, PERMANOVAs with Bray-Curtis dissimilarity were used. ANOVA on log-

transformed data was used to determine significant differences between total CFU

across experiments. In the cases of unequal variances (the individual taxa growth in

perturbations), Kruskal-Wallis tests were used. To determine relationships between

relative abundance and growth of individual strains, linear regressions were used. To

compare total growth alone to growth in the community, t-tests were used. Differences

in a* and b* values in the pigmentation assay were determined using ANOVA. To

determine differences in VOC composition across the nine communities,

PERMANOVAs on Bray-Curtis dissimilarity of relative peak area were used. A SIMPER

analysis of relative peak area of VOCs was used to identify the contributions of each

VOC to Bray-Curtis dissimilarity.

REFERENCES:

1. Shade A, Handelsman J. 2012. Beyond the Venn diagram: the hunt for a core microbiome.

Environ Microbiol 14:4–12.

2. Lundberg DS, Lebeis SL, Paredes SH, Yourstone S, Gehring J, Malfatti S, Tremblay J,

Engelbrektson A, Kunin V, del Rio TG, Edgar RC, Eickhorst T, Ley RE, Hugenholtz P,

Tringe SG, Dangl JL. 2012. Defining the core *Arabidopsis thaliana* root microbiome. Nature

488:86–90.

3. Oh J, Byrd AL, Park M, NISC Comparative Sequencing Program, Kong HH, Segre JA.

2016. Temporal stability of the human skin microbiome. Cell 165:854–866.

4. Albertsen M, Hansen LBS, Saunders AM, Nielsen PH, Nielsen KL. 2012. A metagenome of

a full-scale microbial community carrying out enhanced biological phosphorus removal.

ISME J 6:1094–1106.

5. Oh J, Byrd AL, Deming C, Conlan S, NISC Comparative Sequencing Program, Kong HH,

Segre JA. 2014. Biogeography and individuality shape function in the human skin

metagenome. Nature 514:59–64.

6. Findley K, Oh J, Yang J, Conlan S, Deming C, Meyer JA, Schoenfeld D, Nomicos E, Park

M, NIH Intramural Sequencing Center Comparative Sequencing Program, Kong HH, Segre

JA. 2013. Topographic diversity of fungal and bacterial communities in human skin. Nature

498:367–370.

7. Aßhauer KP, Wemheuer B, Daniel R, Meinicke P. 2015. Tax4Fun: predicting functional

profiles from metagenomic 16S rRNA data. Bioinformatics 31:2882–2884.

8. Nagpal S, Haque MM, Mande SS. 2016. Vikodak--A modular framework for inferring

functional potential of microbial communities from 16S metagenomic datasets. PLoS One

11:e0148347.

9. Toft C, Andersson SGE. 2010. Evolutionary microbial genomics: insights into bacterial host

adaptation. Nat Rev Genet 11:465–475.

10. McAdams HH, Srinivasan B, Arkin AP. 2004. The evolution of genetic regulatory systems in

bacteria. Nat Rev Genet 5:169–178.

11. Thomas CM, Nielsen KM. 2005. Mechanisms of, and barriers to, horizontal gene transfer

between bacteria. Nat Rev Microbiol 3:711–721.

12. Hahn MW, Jezberová J, Koll U, Saueressig-Beck T, Schmidt J. 2016. Complete ecological

isolation and cryptic diversity in *Polynucleobacter* bacteria not resolved by 16S rRNA gene

sequences. ISME J 10:1642–1655.

13. van der Woude MW, Bäumler AJ. 2004. Phase and antigenic variation in bacteria. Clin

Microbiol Rev 17:581–611, table of contents.

14. Cohan FM. 2002. What are bacterial species? Annu Rev Microbiol 56:457–487.

15. Jaspers E, Overmann J. 2004. Ecological significance of microdiversity: identical 16S rRNA

gene sequences can be found in bacteria with highly divergent genomes and

ecophysiologies. Appl Environ Microbiol 70:4831–4839.

16. Leventhal GE, Boix C, Kuechler U, Enke TN, Sliwerska E, Holliger C, Cordero OX. 2018.

Strain-level diversity drives alternative community types in millimetre-scale granular

biofilms. Nat Microbiol 3:1295–1303.

17. Wolfe BE, Button JE, Santarelli M, Dutton RJ. 2014. Cheese rind communities provide

tractable systems for in situ and in vitro studies of microbial diversity. Cell 158:422–433.

18. Wolfe BE, Dutton RJ. 2015. Fermented foods as experimentally tractable microbial

ecosystems. Cell 161:49–55.

19. Button JE, Dutton RJ. 2012. Cheese microbes. Curr Biol 22:R587–R589.

20. Irlinger F, Layec S, Hélinck S, Dugat-Bony E. 2015. Cheese rind microbial communities:

diversity, composition and origin. FEMS Microbiol Lett 362:1–11.

21. Quigley L, O'Sullivan O, Beresford TP, Paul Ross R, Fitzgerald GF, Cotter PD. 2012. High-

throughput sequencing detects subpopulations of bacteria not previously associated with

artisanal cheeses. Appl Environ Microbiol AEM.00918–12.

22. Quigley L, O'Sullivan O, Stanton C, Beresford TP, Ross RP, Fitzgerald GF, Cotter PD.

2013. The complex microbiota of raw milk. FEMS Microbiol Rev 37:664–698.

23. Verdier-Metz I, Gagne G, Bornes S, Monsallier F, Veisseire P, Delbès-Paus C, Montel M-C.

2012. Cow teat skin, a potential source of diverse microbial populations for cheese

production. Appl Environ Microbiol 78:326–333.

24. Laing C, Buchanan C, Taboada EN, Zhang Y, Kropinski A, Villegas A, Thomas JE, Gannon

VPJ. 2010. Pan-genome sequence analysis using Panseq: an online tool for the rapid

analysis of core and accessory genomic regions. BMC Bioinformatics 11:461.

25. Zhao Y, Wu J, Yang J, Sun S, Xiao J, Yu J. 2012. PGAP: pan-genomes analysis pipeline.

Bioinformatics 28:416–418.

26. Price-Whelan A, Poon CK, Benson MA, Eidem TT, Roux CM, Boyd JM, Dunman PM,

Torres VJ, Krulwich TA. 2013. Transcriptional profiling of *Staphylococcus aureus* during

growth in 2 M NaCl leads to clarification of physiological roles for Kdp and Ktr K+ uptake

systems. MBio 4.

27. Kastman EK, Kamelamela N, Norville JW, Cosetta CM, Dutton RJ, Wolfe BE. 2016. Biotic

interactions shape the ecological distributions of *Staphylococcus* species. MBio 7:e01157–

16.

28. Monnet C, Landaud S, Bonnarme P, Swennen D. 2015. Growth and adaptation of

microorganisms on the cheese surface. FEMS Microbiol Lett 362:1–9.

29. Fox PF, Uniacke-Lowe T, McSweeney PLH, O'Mahony JA. 2015. Chemistry and

Biochemistry of Cheese, p. 499–546. *In* Fox, PF, Uniacke-Lowe, T, McSweeney, PLH,

O'Mahony, JA (eds.), Dairy Chemistry and Biochemistry. Springer International Publishing,

Cham.

30. Marcellino N, Benson DR. 2013. The good, the bad, and the ugly: tales of mold-ripened

cheese. Microbiol Spectr 1:CM-0005-2012.

31. Zhang Y, Kastman EK, Guasto JS, Wolfe BE. 2018. Fungal networks shape dynamics of

bacterial dispersal and community assembly in cheese rind microbiomes. Nat Commun

9:336.

32. Louca S, Parfrey LW, Doebeli M. 2016. Decoupling function and taxonomy in the global

ocean microbiome. Science 353:1272–1277.

33. Vieira-Silva S, Falony G, Darzi Y, Lima-Mendez G, Garcia Yunta R, Okuda S, Vandeputte

D, Valles-Colomer M, Hildebrand F, Chaffron S, Raes J. 2016. Species–function

relationships shape ecological properties of the human gut microbiome. Nature

Microbiology 1:16088.

34. Allison SD, Martiny JBH. 2008. Colloquium paper: resistance, resilience, and redundancy in

microbial communities. Proc Natl Acad Sci U S A 105 Suppl 1:11512–11519.

35. Dufossé L, Mabon P, Binet A. 2001. Assessment of the coloring strength of brevibacterium

linens strains: spectrocolorimetry versus total carotenoid extraction/quantification. J Dairy

Sci 84:354–360.

36. Leclercq-Perlat M-N, Spinnler H-E. 2010. The type of cheese curds determined the

colouring capacity of *Brevibacterium* and *Arthrobacter* species. J Dairy Res 77:287–294.

37. Leclercq-Perlat MN, Corrieu G, Spinnler HE. 2004. The color of *Brevibacterium linens*

depends on the yeast used for cheese deacidification. J Dairy Sci 87:1536–1544.

38. Galaup P, Sutthiwong N, Leclercq-Perlat M-N, Valla A, Caro Y, Fouillaud M, Guérard F,

Dufossé L. 2015. First isolation of *Brevibacterium* sp. pigments in the rind of an industrial

red-smear-ripened soft cheese. Int J Dairy Technol 68:144–147.

39. Kamelamela N, Zalesne M, Morimoto J, Robbat A, Wolfe BE. 2018. Indigo- and indirubin-

producing strains of *Proteus* and *Psychrobacter* are associated with purple rind defect in a

surface-ripened cheese. Food Microbiol 76:543–552.

40. McSweeney PLH. 2004. Biochemistry of cheese ripening. Int J Dairy Technol 57:127–144.

41. Irlinger F, Mounier J. 2009. Microbial interactions in cheese: implications for cheese quality

and safety. Curr Opin Biotechnol 20:142–148.

42. Fox PF, Cogan TM. 2004. Factors that Affect the Quality of Cheese, p. 583–608. *In* Cheese

Chemistry, Physics and Microbiology. Elsevier.

43. Baltussen E, Sandra P, David F, Cramers C. 1999. Stir bar sorptive extraction (SBSE), a

novel extraction technique for aqueous samples: theory and principles. J Microcolumn Sep

11:737–747.

44.   Müller A, Faubert P, Hagen M, Zu Castell W, Polle A, Schnitzler J-P, Rosenkranz M. 2013.

Volatile profiles of fungi--chemotyping of species and ecological functions. Fungal Genet

Biol 54:25–33.

45.   Booth RE, Grime JP. 2003. Effects of genetic impoverishment on plant community diversity.

J Ecol.

46.   Violle C, Enquist BJ, McGill BJ, Jiang L, Albert CH, Hulshof C, Jung V, Messier J. 2012.

The return of the variance: intraspecific variability in community ecology. Trends Ecol Evol

27:244–252.

47.   Siefert A. 2012. Incorporating intraspecific variation in tests of trait-based community

assembly. Oecologia 170:767–775.

48.   Lawrence D, Fiegna F, Behrends V, Bundy JG, Phillimore AB, Bell T, Barraclough TG.

2012. Species interactions alter evolutionary responses to a novel environment. PLoS Biol

10:e1001330.

49.   Willger SD, Grim SL, Dolben EL, Shipunova A, Hampton TH, Morrison HG, Filkins LM,

O'Toole GA, Moulton LA, Ashare A, Sogin ML, Hogan DA. 2014. Characterization and

quantification of the fungal microbiome in serial samples from individuals with cystic

fibrosis. Microbiome.

50.   Bartowsky EJ, Borneman AR. 2011. Genomic variations of *Oenococcus oeni* strains and

the potential to impact on malolactic fermentation and aroma compounds in wine. Appl

Microbiol Biotechnol 92:441–447.

51.   Orlic S, Redzepovic S, Jeromel A, Herjavec S, Iacumin L. 2007. Influence of indigenous

Saccharomyces paradoxus strains on Chardonnay wine fermentation aroma. Int J Food Sci

Technol 42:95–101.

52.   Cano-García L, Flores M, Belloch C. 2013. Molecular characterization and aromatic

potential of *Debaryomyces hansenii* strains isolated from naturally fermented sausages.

Food Res Int 52:42–49.

53. Spus M, Li M, Alexeeva S, Wolkers-Rooijackers JCM, Zwietering MH, Abee T, Smid EJ.

2015. Strain diversity and phage resistance in complex dairy starter cultures. J Dairy Sci

98:5173–5182.

54. Walsh AM, Crispie F, Daari K, O'Sullivan O, Martin JC, Arthur CT, Claesson MJ, Scott KP,

Cotter PD. 2017. Strain-level metagenomic analysis of the fermented dairy beverage nunu

highlights potential food safety risks. Appl Environ Microbiol 83:e01144-17.

55. Erkus O, de Jager VCL, Spus M, van Alen-Boerrigter IJ, van Rijswijck IMH, Hazelwood L,

Janssen PWM, van Hijum SAFT, Kleerebezem M, Smid EJ. 2013. Multifactorial diversity

sustains microbial community stability. ISME J 7:2126–2136.

56. Li SS, Zhu A, Benes V, Costea PI, Hercog R, Hildebrand F, Huerta-Cepas J, Nieuwdorp M,

Salojärvi J, Voigt AY, Zeller G, Sunagawa S, de Vos WM, Bork P. 2016. Durable

coexistence of donor and recipient strains after fecal microbiota transplantation. Science

352:586–589.

57. Scholz M, Ward DV, Pasolli E, Tolio T, Zolfo M, Asnicar F, Truong DT, Tett A, Morrow AL,

Segata N. 2016. Strain-level microbial epidemiology and population genomics from shotgun

metagenomics. Nat Methods 13:435–438.

58. Asnicar F, Manara S, Zolfo M, Truong DT, Scholz M, Armanini F, Ferretti P, Gorfer V,

Pedrotti A, Tett A, Segata N. 2017. Studying vertical microbiome transmission from mothers

to infants by strain-level metagenomic profiling. mSystems 2:e00164-16.

59. Ercolini D. 2017. Exciting strain-level resolution studies of the food microbiome. Microb

Biotechnol 10:54–56.

60. Turaev D, Rattei T. 2016. High definition for systems biology of microbial communities:

metagenomics gets genome-centric and strain-resolved. Curr Opin Biotechnol 39:174–181.

61. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S,

Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK,

Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein

V, Wilke A, Zagnitko O. 2008. The RAST Server: rapid annotations using subsystems

technology. BMC Genomics 9:75.

62.　Dufossé L, Galaup P, Carlet E, Flamin C, Valla A. 2005. Spectrocolorimetry in the CIE

L*a*b* color space as useful tool for monitoring the ripening process and the quality of PDO

red-smear soft cheeses. Food Res Int 38:919–924.

63.　Kfoury N, Baydakov E, Gankin Y, Robbat A. 2018. Differentiation of key biomarkers in tea

infusions using a target/nontarget gas chromatography/mass spectrometry workflow. Food

Research International 113:414-423.

64.　Robbat A Jr, Kfoury N, Baydakov E, Gankin Y. 2017. Optimizing targeted/untargeted

metabolomics by automating gas chromatography/mass spectrometry workflows. J

Chromatogr A 1505:96–105.

ACKNOWLEDGEMENTS

Esther Miller, Grace Cox, Elizabeth Landis, Freddy Lee, and Megan Biango-Daniels provided very helpful feedback on earlier drafts of this manuscript. This work was supported by NSF grant 1715553 to B.E.W.

AUTHOR CONTRIBUTIONS

B.E.W. isolated the bacterial strains from the nine original cheeses. B.A.N. designed, conducted, and analyzed all *in vitro* cheese experiments. N.K. and A.R. designed and performed the VOC data collection and analysis. E.K.K. and B.E.W. performed bioinformatic analyses. B.A.N. and B.E.W. performed statistical analyses on the

709 community assembly and functional assays. B.A.N., E.E.K., N.K., A.R., and B.E.W.

710 wrote the manuscript.

712 **COMPETING INTERESTS STATEMENT**

713 The authors declare no competing interests.

715 **FIGURES and FIGURE LEGENDS:**

Figure 1: Isolation of nine taxonomically identical cheese rind core microbiomes. (A) The same three bacterial species - *Staphylococcus equorum*, *Brevibacterium auranticum,* and *Brachybacterium alimentarium* - were isolated from a set of 137 cheese rinds that were previously described using 16S rRNA gene amplicon sequencing (Wolfe et al. 2014). Each column represents average relative abundance data for one cheese rind microbiome. Data are clustered using an UPGMA tree based on Bray-Curtis dissimilarity. **(B)** The three core microbiome species have distinct colony morphologies. **(C)** Graphical representation of the nine core microbiomes as used throughout the manuscript.

725
726
727

728
Figure 2: Accessory genome of the cheese rind core microbiomes. Heatmap indicates
variation in the abundance of unique accessory gene clusters across the three individual taxa
(top) and across SEED functional categories (bottom). Phylogeny is a maximum likelihood
consensus tree constructed from SNPs identified across the nine core communities. Values are
bootstrap support.

734
735
736
737

Figure 3

739
Figure 3 (previous page): Divergent community assembly across the nine cheese rind
core microbiomes. (A) Experimental setup. Each set of three species from each core
microbiome was inoculated into wells of 96-well plates. Communities were harvested three and
ten days after inoculation. **(B)** Total community abundance as measured by CFUs of each of the
nine core microbiomes. **(C)** Relative abundance of each of the three bacterial species across
each of the nine core microbiomes. Each column represents a replicate. I1 and I2 indicate the
input compositions for the two independent experimental replicates. In the Day 3 and Day 10
datasets, the first five columns are from one experimental replicate and the second five are from
a second experimental replicate. Blank columns represent replicates that were lost due to
contamination. **(D)** Growth of each of the community members alone (open circles) and in the
presence of the community (closed black squares). Each point represents the mean CFUs of
the taxa and the error bars represent one standard deviation of the mean. Asterisks indicate
significant differences between growth alone and growth in the community (t-test, p < 0.05).
Figure 4 (next page): Response of the nine cheese rind core microbiomes to abiotic and
biotic perturbations. (A) Responses of each taxa to abiotic (6% salt) and biotic (*Penicillium*)
disturbance. Each point represents the mean CFUs of the taxa in that community at Day 10 and
the error bars represent one standard deviation of the mean. Asterisks indicate significant
difference in growth compared to control based on Kruskal-Wallis test (p<0.05). (B) Mean
community composition in the three treatments. Asterisk indicates significant difference in
community composition compared to control based on PERMANOVA. (C) Principal coordinates
analysis of replicate communities in the three treatments. PCoA is based on Bray-Curtis
dissimilarity of absolute abundances of each community member.

Figure 4

Figure 5

790
791
Figure 5: Functional diversity across nine cheese rind core microbiomes. (A) Color
profiles of experimental rind communities after ten days of rind development. Each dot
represents a replicate cheese rind community. Boxes in legend are representative photos of the
experimental cheese surface from each community. **(B)** Total volatile organic compound (VOC)
diversity across the nine cheese communities. Each point represents the mean number of
VOCs detected in each community and the error bars represent one standard deviation of the
mean. Core communities that share the same letter are not significantly different from one
another based on Kruskal-Wallis test (p<0.05)**. (C)** Non-metric multidimensional scaling of total
VOC profiles. Each dot represents a replicate cheese rind community. **(D)** Relative abundance
of VOCs that contributed the most to the Bray-Curtis dissimilarity across communities (as
determined by SIMPER analysis). Because total concentrations of VOCs are highly variable
across different compounds, visualization was simplified by relativizing the relative peak area
from GC-MS chromatograms within each VOC to the highest concentration detected for that
VOC. Data are clustered together by total VOC profiles using a UPGMA tree. Asterisks indicate
clusters with > 70% bootstrap support.

SUPPLEMENTARY TABLES:

Table S1: Overview of bacterial strains and genomes used in this study

Table S2: Distribution of gene clusters in the three taxa from each of the nine core
microbiomes. When a cell is filled, it indicates that a predicted gene belongs to a gene
cluster (row). In some communities, multiple genes belong to a single gene cluster. The
identifiers in the cells are the gene IDs of each of the genomes based on the RAST
annotation of that genome.

Table S3: Enrichment of SEED subsystem categories in core microbiomes based on
Fisher's exact test.

Table S4: Relative peak area of each volatile organic compound detected from the
experimental cheese communities. The "_1, _2, etc." indicates replicates within each of
the nine core communities.