Genomic potential for photoferrotrophy in a seasonally anoxic Boreal Shield lake

Tsuji JM[1], Tran N[1], Schiff SL[1], Venkiteswaran JJ[1,2], Molot LA[3], Tank M[4], Hanada S[4], Neufeld JD[1]*

[1]University of Waterloo, 200 University Avenue West, Waterloo, Ontario, Canada, N2L 3G1

[2]Wilfrid Laurier University, 75 University Avenue West, Waterloo, Ontario, Canada, N2L 3C5

[3]York University, 4700 Keele Street, Toronto, Ontario, Canada, M3J 1P3

[4]Tokyo Metropolitan University, 1-1 Minami-osawa, Hachioji, Tokyo, Japan 192-0397

*Corresponding author: jneufeld@uwaterloo.ca

Running title: Photoferrotrophy in a Boreal Shield lake

Keywords: photoferrotrophy; *Chlorobia*; *cyc2*; boreal lakes; Archaean Eon; metagenomics; anoxygenic photoautotrophy; iron cycling

Abstract

20   Photoferrotrophy, the light-induced oxidation of ferrous iron, is thought to have contributed to primary production within Earth's early anoxic oceans yet is presumed to be of little modern environmental relevance. Here we use genome-resolved metagenomics and enrichment cultivation to explore the potential for photoferrotrophy in the anoxic water columns of globally abundant Boreal Shield lakes. We recovered four high-completeness and low-contamination draft

25   genome bins assigned to the class *Chlorobia* (formerly phylum *Chlorobi*) from environmental metagenome data and enriched two novel sulfide-oxidizing species, also from the *Chlorobia*. The sequenced genomes of both enriched species, including the novel "*Candidatus* Chlorobium canadense", encoded the *cyc2* candidate gene marker for iron oxidation, suggesting the potential for photoferrotrophic growth. Surprisingly, one of the environmental genome bins encoded *cyc2*

30   and lacked sulfur oxidation gene pathways altogether. Despite the presence of *cyc2* in the corresponding draft genome, we were unable to induce photoferrotrophy in "*Ca.* Chlorobium canadense", suggesting that yet-unexplored mechanisms regulate expression of sulfide and ferrous iron oxidation gene systems, or that previously unrecognized functions for this outer membrane cytochrome exist. Doubling the known diversity of *Chlorobia*-associated *cyc2* genes,

35   metagenome data showed that putative photoferrotrophic populations occurred in one lake but that only sulfide-oxidizing populations were present in a neighboring lake, implying that strong ecological or geochemical controls govern the favourability of photoferrotrophy in aquatic environments. These results indicate that anoxygenic photoautotrophs in Boreal Shield lakes could have unexplored metabolic diversity that is controlled by ecological and biogeochemical

40   drivers pertinent to understanding Earth's early microbial communities.

Introduction

Anoxygenic photoautotrophs are common in oxygen-limited environments worldwide and serve as primary producers in sediments, hot springs, and anoxic water columns as deep as the Black Sea [1–3]. Microorganisms performing anoxygenic photosynthesis commonly use a reduced sulfur compound, such as sulfide or thiosulfate, as the photosynthetic electron donor [4, 5]. However, the discovery of alternative metabolic modes of photosynthesis challenge classical interpretations of how anoxic ecosystems function [6–8]. Photoferrotrophy, an anoxygenic photosynthetic process involving the oxidation of ferrous iron in place of reduced sulfur, has been described in several laboratory cultures but remains poorly understood with respect to its ecological role [9, 10]. Photoferrotrophs are of great interest for their potential involvement in early Earth microbial communities where they may have served as important primary producers in the ferruginous (i.e., iron-rich, sulfate-poor) and anoxic waters of Archaean Eon oceans (ca. 3.8-2.5 Ga) [11–13]. However, photoferrotrophy remains poorly explored in modern environments, and the process is assumed to have little ecological importance given the oxic nature of the Earth's atmosphere and higher levels of sulfate in most modern aquatic systems compared to Archaean oceans [10].

One of the key challenges in assessing the modern importance of photoferrotrophy has been a limited understanding of the biochemistry and genetics of iron oxidation. However, recent work has begun to unravel the molecular basis for iron oxidation in photoferrotrophs and other neutrophilic iron-oxidizing bacteria (reviewed in [14, 15]). Candidate gene systems are emerging that can be used to probe for selected types of iron oxidation environmentally. In particular, genes encoding for porin-cytochrome *c* protein complexes (PCC), such as *pioAB* and *mtoAB*, and genes encoding for outer-membrane monoheme c-type cytochromes, such as *cyc2*, have been identified as potentially useful markers for iron oxidation [15–19]. Recent environmental

metatranscriptomic work, for example, suggests tight links between zetaproteobacterial *cyc2* gene expression and iron oxidation for these microaerophilic iron-oxidizing bacteria [20]. The discovery of such genetic markers expands capabilities to explore the role of photoferrotrophy and other forms of iron cycling in nature.

70   Novel photoferrotrophic bacteria are also being isolated in the laboratory. For example, among the *Chlorobia* class (i.e., "green sulfur bacteria" and "*Candidatus* Thermochlorobacter aerophilum"), the only known photoferrotroph was *Chlorobium ferrooxidans* strain KoFox, which was enriched from freshwater lake sediments [21]. Recently, *Chlorobium phaeoferrooxidans* strain KB01, a bacteriochlorophyll *e*-containing member of the class

75 *Chlorobia*, was isolated from the anoxic water column of the meromictic and ferruginous Kabuno Bay [22]. In addition, *Chlorobium* sp. strain N1 was isolated from marine sediments [23]. Both *Chl. ferrooxidans* and *Chl. phaeoferrooxidans* oxidize ferrous iron as their sole photosynthetic electron donor, assimilating sulfur as sulfate [21, 22]. *Chl.* sp. N1 is capable of oxidizing either ferrous iron or sulfide as the photosynthetic electron donor and can also utilize organic

80 compounds [23]. A fourth recently isolated photoferrotroph, *Chlorobium* sp. BLA1, was shown to also be capable of oxidizing sulfide, although genome data is not yet available for this strain [24]. Phylogenetically, *Chl. ferrooxidans* and *Chl. phaeoferrooxidans* are sister groups to one another, whereas *Chl.* sp. N1 is closely related to *Chl. luteolum*, a sulfide-oxidizing species of *Chlorobia* hypothesized to be able to switch to photoferrotrophic growth [22, 23, 25]. Genome sequences

85 available for *Chl. ferrooxidans*, *Chl. phaeoferrooxidans*, *Chl.* sp. N1, and *Chl. luteolum* show that their genomes encode *cyc2* homologues, unlike all other sulfide-oxidizing *Chlorobia* [25–28]. These recent findings allow for first glimpses into the natural diversity of photoferrotrophs and provide some evidence that *cyc2* may serve as a genomic marker for photoferrotrophy within the class *Chlorobia*.

90   Current knowledge of photoferrotroph ecology derives from studies of a select few modern

environments. Among the best studied are meromictic ferruginous lakes that are used as

analogues of Archaean ocean conditions (reviewed in [29]; see also [30, 31]). Study of these

Archaean ocean analogues has led to varying conclusions about the environmental importance of

photoferrotrophy. In Lake Matano (Indonesia), bacteria classified to the *Chlorobium* genus were

95   detected beneath the oxic-anoxic zone interface [32]. However, only genes implicated in the

sulfide oxidation pathway among *Chlorobia* were detected in metagenome sequencing data, and

anoxygenic photoautotrophy rates were below detection limits, leaving the role of

photoferrotrophy unclear [33]. In Lake La Cruz (Spain), rates of photoferrotrophic iron oxidation

were estimated at 0.2-1.4 μM Fe^{2+} d^{-1}, yet microscopy data and photoautotrophy rates suggested

100   that photoferrotrophy was catalyzed by a small proportion of total phototrophic cells, and that

photoferrotrophs co-existed with other phototrophic bacteria, such as sulfide oxidizers [34]. In

Lake Kivu's Kabuno Bay (Democratic Republic of the Congo), anoxygenic phototrophs in the

upper anoxic zone used ferrous iron for most or all photosynthetic activity at rates of up to 100

μM Fe^{2+} d^{-1} [22]. Anoxygenic photoautotrophs were estimated as being capable of supplying up

105   to ~90% of total fixed carbon in the bay's anoxic zone [35]. Contrasting results from these lakes

suggest that unknown factors in ferruginous systems modulate the relative importance of various

anoxygenic photosynthetic modes. In addition, it appears as though sulfide-oxidizing

photoautotrophs can play a relatively important role in primary production even in sulfide-limited

systems (see also [36]). However, the limited number of such meromictic ferruginous lakes

110   makes further speculation about photoferrotroph ecology and *in situ* activity challenging.

        Recently, we combined 16S ribosomal RNA (16S rRNA) gene sequencing and iron isotope

data to propose photoferrotrophy by populations of *Chlorobia* as an important photosynthetic

process in the seasonally anoxic water columns of two ferruginous Boreal Shield lakes at the

International Institute for Sustainable Development Experimental Lakes Area (IISD-ELA; near

115 Kenora, Canada) [37]. We studied Lake 227, an experimentally eutrophied lake, and Lake 442, a

nearby and pristine reference lake [38, 39]. Boreal Shield lakes number in the tens of millions

across northern regions globally, and we proposed based on our preliminary data that Boreal

Shield lakes that develop seasonal anoxia commonly contain photoferrotrophic bacteria and could

serve as plentiful modern-day analogues of Archaean ocean ecosystems. Here, we use genome-

120 resolved environmental metagenomics to recover high-completeness, low-contamination draft

genomes of key populations of *Chlorobia* from the same two IISD-ELA lakes. We then apply

recent knowledge of iron functional gene markers to test whether these populations possess the

genetic potential for photoferrotrophy or other phototrophic modes. We also report the

enrichment of two novel species of *Chlorobia* from Lake 227 and nearby Lake 304 and analyze

125 their genetic and functional potential for phototrophic sulfide and iron oxidation. Through this

first usage of molecular tools to explore *Chlorobia*-driven photoferrotrophy in an environmental

context, we aim to clarify the diversity and ecology of anoxygenic photoautotrophic bacteria in

Boreal Shield lakes, with implications for understanding both Archaean microbial communities

and modern ferruginous environments.

Materials and Methods

Lake sampling, metagenome sequencing, assembly, and binning

The IISD-ELA sampling site, lake geochemistry, and sample collection methods were

described in detail previously [37]. Based on preliminary 16S rRNA gene data, we selected six

135 water column genomic DNA samples collected from Lakes 227 and 442 for re-sequencing via

shotgun metagenomics. All six samples were collected at or beneath the oxic-anoxic zone

boundary, at depths where low light penetration occurs, and had high relative abundances of anoxygenic phototrophs, dominated by populations of *Chlorobia*. Samples from Lake 227 were selected at 6 m and 8 m in 2013 and 2014, and samples from Lake 442 were selected at 16.5 m in 2011 and 13 m in 2014. The sequencing library was prepared using the NEBNext Ultra II DNA Library Prep Kit for Illumina (New England Biolabs; Ipswich, Massachusetts, USA) and was sequenced on a single lane of a HiSeq 2500 (Illumina; San Diego, California, USA) in Rapid Run Mode with 2x200 base paired-end reads, generating 26.3 to 39.5 million reads per sample. Library preparation and sequencing was performed by the McMaster Genome Facility (Hamilton, Ontario, Canada).

Raw metagenome reads were quality-trimmed, assembled, binned, and annotated using the ATLAS pipeline, version 1.0.22 [40]. The configuration file with all settings for the pipeline is available in Supplementary File 1. To improve genome bin completeness and reduce bin redundancy, QC processed reads from related samples (i.e., L227 2013 6 m and 8 m, L227 2014 6 m and 8 m, and L442 samples) were co-assembled and re-binned using differential abundance information via a simple wrapper around ATLAS, as described in the Supplementary Methods.

Identification of cyc2 genes and Chlorobia genome bins

Assembled contigs were assessed for the presence of *cyc2* gene homologues by building a custom profile Hidden Markov Model (HMM). The predicted primary sequences of *cyc2* were recovered from the four available genomes of *Chlorobia* known to possess the gene, *Chl. ferrooxidans*, *Chl. phaeoferrooxidans*, *Chl.* sp. N1, and *Chl. luteolum*, as well as from the genomes of reference microaerophilic iron-oxidizing bacteria as described by He and colleagues [14]. A cytochrome 572 gene from *Leptospirillum* sp. (EDZ39515.1) was omitted from the reference dataset, due to its high divergence from other sequences, to build a more robust

160   alignment for the *cyc2* clade relevant to the *Chlorobia* [41]. Collected sequences were aligned

using Clustal Omega, version 1.2.3 [42], and the alignment was used to build a profile HMM

using the *hmmbuild* function of HMMER3, version 3.2.1 [43]. Recovered *cyc2* genes over the

course of the study were added to the alignment and HMM; final HMMs are included in

Supplementary File 2.

165   Recovered *cyc2* genes were compared to *cyc2* reference sequences by building a maximum-

likelihood phylogeny. The alignment was masked using Gblocks, version 0.91b [44], as described

in the Supplementary Methods. The maximum likelihood phylogeny was then prepared from the

masked sequence alignment via IQ-TREE, version 1.6.10 [45], using the LG+F+I+G4 sequence

evolution model as determined by the ModelFinder module of IQ-TREE [46]. To build the

170   consensus tree, 1000 bootstrap replicates were performed, each requiring ~100-110 tree search

iterations for phylogeny optimization.

        Genome bins were dereplicated using dRep version 2.0.5 [47] with default settings. All

genome bins from individual assembles and co-assemblies were pooled for dereplication to help

ensure that the highest quality bins, whether from individual assemblies or co-assemblies, would

175   be retained for downstream analyses. Genome bins of *Chlorobia* with > 90% completeness and <

10% contamination based on CheckM statistics [48] were selected for further study. These

genome bins were imported into Anvi'o version 4 [49] and were manually examined for contigs

improperly binned based on read mapping, tetranucleotide frequencies, and contig taxonomic

classification. Import of ATLAS data into Anvi'o was performed using the *atlas-to-anvi.sh* script,

180   version 1.0.22-coassembly-r4, available in the *atlas-extensions* GitHub repository at

https://github.com/jmtsuji/atlas-extensions. Contigs in curated genome bins of *Chlorobia* were

then ordered via the Mauve Contig Mover development snapshot 2015-02-13 for Linux [50]

using the *Chl. luteolum* genome to guide ordering. To assess bin quality, tRNA genes were predicted using Prokka v1.13.3 [51].

Enrichment cultivation, sequencing, and assembly

Lake 227 and Lake 442 were sampled again in June 2016 and July 2017, along with the nearby Lake 304 in July 2017. Basic sampling information for each lake is summarized in Supplementary File 3. Lake 304 has been described previously and also has seasonally anoxic bottom waters [52, 53]. The lake was experimentally fertilized with phosphate in 1971-72 and 1975-76, with ammonium in 1971-74, and with nitrate in 1973-74, but the lake has not been manipulated since then and rapidly returned to its non-eutrophied state once additions ceased, having a water residence time of ~2.7 years [54–56]. Lake 304 was added to our sampling efforts for enrichment culturing to explore the broader distribution of photoferrotrophy among IISD-ELA lakes, because preliminary 16S rRNA gene sequencing data (not shown) indicated high relative abundances of *Chlorobia* populations in this lake. Anoxic water was collected from Lakes 227 and 304 at a depth of 6 m and Lake 442 at 15 m, where trace levels of light are generally detectable (i.e., in the range of 0.01-1 μmol photons m^{-2} s^{-1} between 400-700 nm wavelengths, as commonly measured when field sampling). Lake water was pumped to the surface using a gear pump and directly injected into N_2-filled 160 mL glass serum bottles that were sealed with blue butyl rubber stoppers (Bellco Glass; Vineland, New Jersey, USA), using a secondary needle to vent N_2 gas until the bottles were full. Water was kept cold (~4°C) and dark after collection and during shipping.

Enrichment cultures were grown using sulfide-containing Pfennig's medium, prepared as described by Imhoff [4, 5], or ferrous iron-containing freshwater medium, prepared as described by Hegler and colleagues [57]. The ferrous iron-containing medium contained 8 mM ferrous

chloride ($FeCl_2$), without filtration of precipitates, and used trace element solution SLA [4]. Initial enrichments contained 10-20% lake water at a total volume of 50 mL and were inoculated into 120-160 mL glass serum bottles sealed with blue butyl rubber stoppers (Bellco Glass) or black rubber stoppers (Geo-Microbial Technology Company; Ochelata, Oklahoma, USA). The

210   remaining headspace was flushed with a 90:10 N_2:CO_2 gas mix at left at a pressure of 1.5 atm. Several bottles additionally had anoxic DCMU (i.e., Diuron or 3-(3,4-dichlorophenyl)-1,1-dimethylurea; Sigma-Aldrich; St. Louis, Missouri, USA) added to a concentration of 50 µM to block activity of oxygenic phototrophs [58, 59]. Bottles were incubated at 22°C and ~50 cm away from far red PARSource PowerPAR LED Bulbs (LED Grow Lights Depot; Portland, Oregon,

215   USA) as the light source.

Two enrichment cultures survived repeated subculturing or dilution-to-extinction and contained green sulfur bacteria based on pigmentation and marker gene sequence analysis. Biomass from these cultures was collected by centrifugation, and genomic DNA was extracted using the DNeasy UltraClean Microbial Kit (Qiagen; Venlo, The Netherlands). Genomic DNA

220   was prepared into metagenome sequencing libraries using the Nextera DNA Flex Library Prep Kit (Illumina), and the library was sequenced on a fraction of a lane of a HiSeq 2500 (Illumina) in High Output Run Mode with 2x125 base paired-end reads. Library preparation and sequencing was performed by The Centre for Applied Genomics (TCAG; The Hospital for Sick Children, Toronto, Canada), generating ~6 million total reads per sample. Metagenomes were assembled

225   using ATLAS version 1.0.22 without co-assembly. Genome bins of *Chlorobia* were manually refined as described above, using Anvi'o version 5 [49] via *atlas-to-anvi.sh* commit 99b85ac; contigs of curated bins were subsequently ordered, and tRNA genes counted, as described above.

Comparative genomics of Chlorobia genomes

Refined genome bins of *Chlorobia*, which belonged to the *Chlorobium* genus, were
230   compared to genomes of reference strains from the *Chlorobiaceae* family. Genomes of all
available type strains and photoferrotrophs from the family were downloaded from the NCBI
(Supplementary File 4). Average nucleotide identity (ANI) between genomes was calculated
using FastANI version 1.1 [60]. The phylogenetic relationship between genomes was determined
by constructing a concatenated ribosomal protein alignment based on the rp1 set of 16 ribosomal
235   protein genes using GToTree version 1.1.10 [61, 62]. IQ-TREE version 1.6.9 [45] was used to
construct the maximum likelihood phylogeny from the alignment. The LG+F+R4 model of
sequence evolution was identified as optimal by the IQ-TREE ModelFinder module [33], and
phylogeny construction used 1000 bootstrap replicates, each requiring ~100-110 tree search
iterations for optimization.

240   The presence or absence of genes implicated in iron or sulfur cycling metabolism in the
collection of *Chlorobiaceae* genomes was assessed using reciprocal BLASTP [63]. Genes were
selected from the genomes of *Chl. ferrooxidans* and *Chl. clathratiforme* based on the genome
comparison of Frigaard and Bryant [25]. Predicted primary sequences of these genes were used to
identify putative homologues across other genomes of *Chlorobia* using the BackBLAST pipeline,
245   version 2.0.0-alpha2 (doi:10.5281/zenodo.3465955) [64]. The e-value cutoff for BLAST hits was
set at 10^{-40}, based on empirical testing of gene clusters known to be present or absent in reference
genomes of *Chlorobia*, and the identity cutoff was set to 20%. Selected genes (e.g., *qmoA*, *dsrJ*)
were omitted due to poor homology between reference genomes. Genes associated with
photosynthesis and carbon fixation were similarly assessed using BackBLAST, except that most
250   reference genes were selected from the genome of *Chl. tepidum* according to Bryant and
colleagues [65] and Tourova and colleagues [66].

Metagenome taxonomic and functional profiling

Environmental relative abundances of microbial populations were estimated by read

mapping to dereplicated genome bins (further details in Supplementary Methods). Taxonomy and

255   functional gene information were then overlaid onto relative abundance data. Genome bin

taxonomy was determined based on the Genome Taxonomy Database (GTDB) using GTDB-Tk,

version v0.2.2, relying on GTDB release 86, version 3 (April 2019) [67, 68]. As such, GTDB

taxonomy names are used throughout this manuscript. Genomes were tested for the presence of

the *cyc2* and *dsrA* functional genes using the HMM developed in this study and an HMM

260   available on FunGene [69], respectively. Gene amino acid translations were predicted for all

genome bins using prodigal version 2.6.3 [70], via the GTDB-Tk, and amino acid translations

were searched via HMMs using the MetAnnotate pipeline development release version 0.9.2

[71]. An e-value cutoff of 10^{-40} was used to filter low-quality gene hits. In case of bias due to

unassembled or incorrectly binned genes, genome bin-based environmental abundance metrics

265   were cross-compared to abundance metrics generated by directly scanning unassembled

metagenome reads, as described in the Supplementary Methods.

Assessment of ferrous iron oxidation potential of Chlorobia enrichments

After initial sequencing, cultures containing *Chlorobia* spp. continued to be purified in the

laboratory. A single culture, which was enriched on sulfide-containing medium, survived

270   continued cultivation and was provisionally named "*Candidatus* Chlorobium canadense strain

L304-6D" (ca.na.den'se N.L. neut. adj. *canadense* from or belonging to Canada). "*Ca*. Chl.

canadense" was purified through multiple rounds of incubation in deep agar dilution series and

picking of isolated green colonies [72]. For deep agar shakes, Pfennig's medium or modified

Chloracidobacterium thermophilum Midnight (CTM) medium [73] containing 1-3 mM buffered

275   sulfide feeding solution were used [4]. Eventually, in preparation for growth in ferrous iron-

containing medium, cultures of "*Ca.* Chl. canadense" were transferred to liquid freshwater

medium as described by Hegler and colleagues [57] (here termed, "Hegler freshwater medium")

but containing 0.5-1 mM buffered sulfide feeding solution, 10-20 μM $FeCl_2$, and 0.5 mg/L

resazurin to monitor the medium's redox status.

280   To test its ability to oxidize ferrous iron phototrophically, "*Ca.* Chl. canadense" was

inoculated into Hegler freshwater medium containing low levels of ferrous iron. Hegler

freshwater medium was made without ferrous iron nor sulfide and was aliquoted into 120 mL

glass serum bottles under flow of sterile 90:10 N_2:CO_2 gas, with 50 mL of medium per bottle.

Bottles were sealed with black rubber stoppers (Geo-Microbial Technology Company) and were

285   bubbled for at least nine minutes with additional 90:10 N_2:CO_2 gas to reduce trace oxygen. The

headspace of each bottle was left at 1.5 atm pressure. A sterile solution of $FeCl_2$ was added to

each bottle to reach a ferrous iron content of 100 μM, and to some bottles, a sterile and anoxic

solution of the chelator ethylenediaminetetraacetic acid (EDTA) was added to a concentration of

120 μM as done by Peng and colleagues [74] to explore whether this chelator could enhance

290   photoferrotrophic iron oxidation. Bottles were incubated at room temperature in the dark

overnight to allow for complexation of ferrous iron and EDTA. All media was confirmed to have

a pH of 6.5-7.

        "*Ca.* Chl. canadense" was grown in 100 mL Hegler freshwater medium using a total of ~2

mM sulfide, fed incrementally in 0.4 mM doses, until sulfide was completely oxidized and the

295   resazurin in the bottle went slightly pink. At the same time, a culture of *Chl. ferrooxidans* KoFox

was grown in the same freshwater medium containing ~8 mM $FeCl_2$ until ferrous iron was

completely oxidized. The entire 100 mL "*Ca.* Chl. canadense" culture was pelleted by

centrifugation at 7000 x g for 13 minutes, and the pellet was washed twice with unamended

Hegler freshwater medium. The ~200 mg of wet biomass recovered was suspended in 1 mL of

300   unamended freshwater medium, and 0.1 mL was inoculated into each relevant incubation bottle.

Similarly, 100 mL of the *Chl. ferrooxidans* reference culture was pelleted by centrifugation at

7000 x g for 5 minutes, washed twice, and suspended in 1 mL of unamended freshwater media,

and 0.1 mL was added to relevant bottles. Only the surfaces of *Chl. ferrooxidans* cell pellets,

which contained mostly green cells and not brown ferric iron precipitate, were suspended with

305   each wash, allowing for ~3 mg of nearly iron-free wet biomass to be recovered. Once cultures

were inoculated, they were transferred to a 22°C incubator without shaking where they received

30 μmol photons m^{-2} s^{-1} white light from a mix of fluorescent (F48T12/CW/VHO; Osram

Sylvania; Wilmington, Massachusetts, USA) and incandescent (60W) bulbs.

Cultures were sampled regularly over a 21-day incubation period to monitor iron

310   concentrations and iron oxidation states. At each sampling time point, an aliquot of culture was

removed from each bottle using a sterile 90:10 N_2:CO_2-flushed syringe and 25 G needle, and 330

μL of this culture aliquot was acidified immediately with 30 μL of 6 N hydrochloric acid (HCl).

Acidified samples were stored for no more than two days at 4°C (with the exception of samples

collected on day 14; see Supplementary Methods) before being assessed for iron species

315   concentrations via the ferrozine assay [75], with 5% ascorbic acid being used as the reducing

agent to determine total iron species [76].

Results

Recovery of Chlorobium genome bins

320   Binning of assembled contigs from the lake and enrichment culture metagenomes, followed

by dereplication of highly similar bins and manual curation, allowed for the recovery of six

highly complete, low-contamination genome bins that classified within the genus *Chlorobium* (Table 1). Three of the genome bins recovered from lake metagenomes had best representatives selected by dRep from the Lake 227 (2013 sample) co-assembly, and one had its best

325   representative from the Lake 442 (2011/2014 sample) co-assembly. In addition, a single *Chlorobium* genome bin was recovered from each enrichment culture. Recovered bins had at least 90.1% completeness and a maximum of 2.8% contamination based on analysis by CheckM [48]. Recovered genome bins were between 1.9 to 2.6 Mb in size, within the approximate length range of reference genomes associated with the family *Chlorobiaceae* (2.0-3.3 Mb; average 2.7

330   Mb), and were represented by 13 to 208 contigs. Other lower quality genome bins classified as belonging to the *Chlorobia* were also recovered (Supplementary File 5), but these were not considered for gene pathway analysis in order to minimize the risk of including false positives. In particular, L442 Bin 74 was excluded from the set of curated bins due to its high contig count (282) and elevated predicted contamination (4.12%) and strain heterogeneity (25%).

335   The constructed HMM for the *cyc2* gene enabled the recovery of five potential *cyc2* homologues associated with *Chlorobia* from assembled contigs (Fig 1). Three of these *cyc2* homologues were found in the set of six manually curated genome bins described above, and another homologue was detected in one of the lower quality genome bins of *Chlorobia* (L227 2014 Bin 92). The final homologue was detected on a short contig in the same lower quality

340   genome bin and was removed from subsequent analyses due to its lack of gene neighbours. Detected *cyc2* homologues were full-length genes, and their predicted primary sequences were detected in the HMM search with a maximum e-value of $\sim 10^{-140}$. All recovered *cyc2* homologues contained the heme-binding CXXCH motif near the N-terminus of their predicted primary sequence, and all homologues were predicted by I-TASSER [77] to contain a porin-like beta

345   barrel structure for membrane binding (Fig. 1A; one example shown) [13]. Aside from the closely

related genomes of *Chl. ferrooxidans* and *Chl. phaeoferrooxidans* (Supplementary Figure 1), all other genomes showed substantial genome re-arrangement around the *cyc2* homologue (Fig. 1B). However, detected *cyc2* homologues were always adjacent to a *c5* family cytochrome (Fig. 1B). These *c5* family cytochromes had low sequence identity to one another (i.e., as low as 30%) yet

350   also contained a CXXCH motif near the N-terminal end (Supplementary File 6). All recovered *cyc2* homologues grouped monophyletically with *cyc2* predicted primary sequences belonging to reference genomes of *Chlorobia* compared to the *cyc2* of other reference strains (Fig. 1C). Overall, the combination of HMM search specificity, genomic context, predicted gene motifs, and phylogenetic placement is strong evidence that the identified cytochrome genes are *cyc2*

355   homologues.

Enrichment cultivation of Chlorobia

Enrichment cultivation was attempted with anoxic lake water using both sulfide- and ferrous iron-containing anoxic media. For Lake 442, one enrichment in a ferrous iron-containing medium grew with evidence of light-driven iron oxidation, but 16S rRNA gene sequencing

360   showed that the culture was dominated by a member of the genus *Rhodopseudomonas*, which did not represent a detectable genus in environmental sequencing data. Given the known metabolic versatility of *Rhodopseudomonas* spp. and the negligible contribution of members of this genus to the studied lake microbial communities, this culture was not pursued further. No members of the *Chlorobia* were ever detected in any enrichment culture from Lake 442.

365   For Lakes 227 and 304, growth was observed for both media tested. Enrichments grown on sulfide-containing media developed green colour after ~12-15 weeks of incubation and contained populations of *Chlorobia* based on Sanger sequencing of both the V3-V4 region of the 16S rRNA gene and the partial *dsrAB* gene (using *Chlorobia*-targeted PCR primers 341f/GSB822r [78] and

PGdsrAF/PGdsrAR [79], respectively). However, the appearances of cultures grown on ferrous

370   iron-containing media for these lakes differed from those of reference photoferrotrophic members

of *Chlorobia*. Instead of forming a reddish-brown ferric iron precipitate, the enrichments

blackened, potentially indicative of sulfate reduction and metal sulfide formation. After

blackening, enrichments developed a green colour (Supplementary Figure 2). Suspecting that

sulfate reduction was occurring in the ferrous iron enrichment bottles to support sulfide-oxidizing

375   phototrophy, one Lake 227 enrichment was transferred into sulfide-containing media and still

developed green pigmentation, albeit slowly (Supplementary Figure 2). Amplification and Sanger

sequencing of the partial *dsrAB* gene cluster from this enrichment showed that the gene sequence

only differed by one ambiguous base from, and thus was essentially identical to, a corresponding

Lake 227 sulfide enrichment. Ferrous iron enrichments eventually stopped being followed due to

380   long growth times, low biomass, and instability of the cultures. Enrichments on sulfide-

containing media were continued instead.

Metagenome sequencing of the two successful enrichments of *Chlorobia* grown on sulfide

(L227 enrichment S-6D and L304 enrichment S-6D) showed that that genomes bins of the

Chlorobia from both enrichments encoded *cyc2* homologues. Although L227 enrichment S-6D

385   ceased to grow in laboratory culture, the L304 enrichment S-6D continued to be purified and was

named "*Candidatus* Chlorobium canadense strain L304-6D". At the time of running the ferrous

iron oxidation test (below), the "*Ca.* Chl. canadense" culture consisted of >90 % cells of

Chlorobia based on epifluorescence microscopy, and other contaminating cells were known to be

chemoorganoheterotrophs and sulfate reducing bacteria based on previous 16S rRNA gene

390   amplicon sequencing data (Supplementary File 7).

Metabolic diversity and phylogeny of Chlorobium genome bins

Recovered genome bins of *Chlorobia* were metabolically diverse and grouped into four general categories with respect to the presence of genes involved in iron and sulfur cycling (Fig. 2). One of the lake-recovered bins (L227 2013 Bin 56) lacked all genes involved in sulfide,

395 elemental sulfur, and thiosulfate oxidation but contained the *cyc2* gene homologue. Conversely, two other lake-recovered bins (L227 2013 Bin 55 and L442 Bin 64) contained all assessed genes in the sulfide oxidation pathway but lacked the *cyc2* gene. The final lake-recovered bin (L227 2013 Bin 22) lacked both *cyc2* and genes for sulfur oxidation. This lack of key metabolism genes in L227 2013 Bin 22 might be due to incomplete genome binning, although the presence a

400 bacterioferritin hypothesized by Frigaard and Bryant [25] to play a role in photoferrotrophy in this genome implies that photoferrotrophy might be possible. Lastly, the two enrichment culture bins contained genes involved in both sulfide and iron oxidation, similarly to *Chl.* sp. N1 and *Chl. luteolum*. Genes for hydrogen oxidation were detected in all of these genome bins (Fig. 2), along with marker genes for photosynthesis and bacteriochlorophyll biosynthesis (Supplementary

405 Figure 3). Marker genes for carbon fixation (via the reverse TCA cycle) were detected in all genome bins, with the exception of L227 2013 Bin 56, which was missing one of the two assessed gene clusters for the this process (*aclAB*; Supplementary Figure 3).

All recovered *Chlorobia* genome bins encoded the *cys* gene cluster for assimilatory sulfate uptake, which was noted previously to potentially be associated with photoferrotrophy due to its

410 absence from the genomes of most sulfur oxidizing strains [25]. The L227 2013 Bin 56 only contained half of the *cys* cluster, but the specific contig containing the cluster was checked and was found to have an assembly break point halfway through the gene cluster, making it likely that the complete *cys* cluster was present in the genome but failed to bin. However, the fact that *Chlorobia*-associated *cyc2* was completely undetectable in Lake 442 metagenome data (see

415   below) implies that, although Lake 442 Bin 64 contains the *cys* cluster, it does not encode *cyc2*.

Also, the genome of *Chl.* sp. N1 lacks the *cys* cluster, implying that the cluster is not necessary

for photoferrotrophic growth (Fig. 2). As such, the *cys* cluster is likely not a reliable indicator of

whether genomes of *Chlorobia* encode *cyc2*. Two of the genome bins also encoded putative

homologues to a bacterioferritin that is potentially involved in photoferrotrophy (see above).

420   Phylogenetic analysis of the recovered *Chlorobium* genome bins based on concatenated

ribosomal protein sequences showed that all four lake-recovered bins and the bin from the Lake

227 enrichment formed a monophyletic subgroup within the *Chlorobiaceae*, including *Chl.*

phaeoclathratiforme BU-1 (Fig. 2). Directly basal to this subgroup was a branch containing *Chl.*

ferrooxidans and *Chl. phaeoferrooxidans*. By comparison, the bin of "*Ca.* Chl. canadense"

425   grouped sister to *Chl. chlorochromatii*, separately from the above group and from the group

containing *Chl. luteolum* and *Chl.* sp. strain N1. Comparison of the concatenated ribosomal

protein phylogeny to the *cyc2* predicted primary sequence phylogeny reveals some congruency

between the two phylogenies for potentially photoferrotrophic members of the *Chlorobia*, with

the exception of L227-S-6D and "*Ca*. Chl. canadense" (Supplementary Figure 4). Overall, all six

430   recovered *Chlorobium* genome bins appear to represent novel species, having an ANI of < 79.6%

compared to the genomes of all available type strains (Supplementary Figure 1).

Functional and taxonomic profiling

Read mapping to the genome bins of *Chlorobia* showed that populations with varying

genomic potential for iron and sulfur-cycling were relevant to the lake environment (Fig. 3). Bins

435   containing *cyc2* (no *dsrA*) and bins containing *dsrA* (no *cyc2*) were both found at similar relative

abundances in Lake 227 samples in 2013 and 2014 (Fig. 3A). Generally, genome bins of

Chlorobia containing *cyc2* accounted for 0.6-2.8% of the Lake 227 microbial community data in

the assessed samples, whereas bins of *Chlorobia* containing *dsrA* accounted for 1.7-4.1% of the

data. Analyses of unassembled metagenome read data indicated similar relative abundances

440   (Supplementary Figure 5). In contrast, only genome bins encoding sulfide oxidation were

detected in Lake 442 metagenome assemblies. Even when unassembled metagenome data were

scanned, only a single HMM hit from the Lake 442 metagenomes matched *cyc2* classified as

belonging to a member of *Chlorobia*, compared to 27 000-37 000 total *rpoB* hits for these same

metagenomes, showing that the lack of *cyc2* genes detected in genome bins of *Chlorobia* from

445   L442 was not a result of incomplete assembly or genome binning. One genome bin of *Chlorobia*

overlapped between L227 and L442 (L442 2011 16.5m Bin 8); the rest were distinct between the

two lakes. In addition, the bins of *Chlorobia* recovered from the enrichment culture metagenomes

were not detected at high relative abundances in any of the six lake metagenomes.

Several non-*Chlorobia* genome bins were also found to contain the *cyc2* or *dsrA* gene

450   through MetAnnotate-based HMM searches (Fig. 3B). Genome bins containing potential *cyc2*

homologues grouped into the *Rhodoferax* or *Gallionella* genera known to include iron-cycling

bacteria [18, 80] or into families not associated with iron oxidation (i.e., *Methylomonadaceae* and

Steroidobacteraceae). The genome bins classified to the *Methylomonadaceae* family were at high

relative abundance in the lake metagenomes, suggesting an abundance potentially as high as

455   ~17% in Lake 227 (Fig. 3C). All *Methylomonadaceae*-affiliated bins, except for L442 2011

16.5m Bin 5, encoded particulate methane monooxygenase (*pmoA*) genes. Genome bins

containing *dsrA* included known chemolithotrophic sulfide-oxidizing bacteria of the *Sulfuritalea*

genus, heterotrophic sulfate-reducing bacteria of the *Desulfatirhabdiaceae* family, a bin classified

to the *Smithellaceae* family of the *Syntrophales* order, and a bin classified to the *Rhodoferax*

460   genus [81, 82]. These taxa, with the exception of *Rhodoferax* and *Smithellaceae*, also appeared in

unassembled read-based marker gene assessments of the metagenomes (Supplementary Figure 5) and could play a role in the broader iron and sulfur cycles of the lakes.

Chlorobia appeared to represent the dominant phototrophs in the lake anoxic zone samples. In total, genome bins classified to the *Chlorobia* represented as high as 8.3% of lake microbial

465   communities based on read mapping (Fig. 3C). Predicted relative abundances of populations of *Chlorobia* were 1.2-1.8 times higher when unassembled reads were assessed directly (Supplementary Figure 5), with a predicted maximum relative abundance of 12.3%, showing that high relative abundance is not a result of assembly bias (Fig. 3D). The only other potential chlorophototroph detected in the dataset at >1 % relative abundance was a single genome bin

470   classified to the *Chloroflexaceae* family (L442 Bin 82; 92.9/1.2% completeness/contamination), which made up 1.9% of the L442 microbial community in 2014 at 15 m depth (Fig. 3C) and contained the *bchL* and *pufM* photosynthesis genes (data not shown). This genome bin was undetected in all other samples.

Assessment of ferrous iron oxidation potential of "Ca. Chl. canadense"

475   Cultures of *Chl. ferrooxidans* showed the expected behaviour when incubated in ferrous iron containing medium. After a short lag phase, both *Chl. ferrooxidans* cultures containing 100 μM FeCl$_2$ (no EDTA) and exposed to light began oxidizing ferrous iron, and the cultures had nearly completely oxidized all ferrous iron within two days of the initial setup (Fig. 4). Similar bottles incubated in the dark showed consistent iron oxidation rates of ~5 μM d^{-1} that

480   corresponded to rates observed in uninoculated media controls; this iron oxidation was inferred to be abiotic, for example due to small amounts of oxygen contamination during sampling. No ferrous iron oxidation outside of this baseline effect was observed in the *Chl. ferrooxidans* cultures supplemented with 120 μM EDTA, in contrast to the stimulatory effect of EDTA

reported for *Rhodopseudomonas palustris* TIE-1 [74], which uses a different gene system

485 (*pioAB*) for extracellular electron transfer [19]. The EDTA-amended medium consistently

appeared to have ~50% ferrous iron oxidation at the beginning of the experiment, based on the

ferrozine assay, but this apparent oxidation has been observed previously with EDTA and could

potentially be due to ineffective binding of ferrozine to EDTA-chelated iron [76].

In contrast to *Chl. ferrooxidans*, "*Ca.* Chl. canadense" showed no photoferrotrophic ferrous

490 iron oxidation activity over the course of the experiment (Fig. 4; Supplementary Figure 6), and

the iron oxidation profiles of the cultures closely matched those of the uninoculated media

control bottles. A reference bottle of Hegler freshwater medium prepared in the same batch as

those in the main experiment but amended with 600 µM buffered sulfide feeding solution, in

place of ferrous iron, received the same inoculum of "*Ca.* Chl. canadense". Complete oxidation

495 of sulfide in the bottle was observed within five days of inoculation and incubation under light

(not shown), indicating that the media and inoculum were not the cause of the lack of observed

photoferrotrophic activity.

Discussion

500 Our study offers genome-resolved functional gene evidence and enrichment cultivation data

that provides a first glimpse into the ecology of photoferrotrophy in seasonally anoxic Boreal

Shield lakes. These results represent the first application of genome-resolved metagenomics to an

Archaean ocean analogue system and the first use of this approach to delineate potential

metabolic modes of coexisting, closely related photoautotrophs. We suggest the genomic

505 potential for photoferrotrophy in Lake 227 based on detection of the *cyc2* gene in multiple

genome bins of *Chlorobia*, and we argue for complex phototrophic ecology in Boreal Shield

lakes based on metabolic diversity of photoautotrophs within and between lake systems. Overall,

our findings provide genomic context to previous 16S rRNA gene data supporting the existence of photoferrotrophy in Boreal Shield lakes [37], adding nuance to the interpretation of these data

510   while expanding sparse knowledge of the diversity of *cyc2* within the class *Chlorobia*. Our research also reinforces the need for additional genetic work to understand the function and regulation of *cyc2* in photoferrotrophy.

Chlorobia-affliated *cyc2* genes were detected in high relative abundance genome bins in Lake 227 metagenomic data. The dominant *cyc2*-containing genome bin of *Chlorobia* in Lake

515   227 metagenomic data, L227 2013 Bin 56, was highly complete and represented as much as 2.8% of the upper anoxic zone microbial community based on read recruitment (Fig. 3A). Although populations of *Chlorobia* are commonly detected at high relative abundances in the anoxic zones of boreal lakes, these populations are typically assumed to oxidize sulfide (e.g., [83]). Because L227 2013 Bin 56 encoded no detectable genes for sulfide or thiosulfate oxidation (Fig. 2), the

520   corresponding population might rely solely on *cyc2* to supply electrons (e.g., from ferrous iron) needed for photosynthesis. Although the population represented by this genome bin might obtain energy chemotrophically via hydrogen oxidation (Fig. 2), as is possible for many members of the *Chlorobia* class [25], the higher relative abundance of *Chlorobia* in the upper compared to lower anoxic zone of Lake 227 (based on previous 16S rRNA gene sequencing data [37]) suggests that

525   light energy is being used in metabolism. The lack of detected sulfur oxidation genes in L227 2013 Bin 56 might be due to incomplete binning, given that the bin was found to encode only partial gene pathways for sulfate assimilation and the reverse TCA cycle (Fig. 2; Supplementary Figure 3), and additional work would be needed to confirm whether *cyc2* is actively expressed. However, our detection of *cyc2* affiliated with this high relative abundance genome bin of

530   *Chlorobia* in Lake 227 metagenomes provides important first evidence that the gene may bear ecological relevance for this Boreal Shield lake.

Our results reveal the coexistence of populations of *Chlorobia* with varying metabolic potential in Boreal Shield lakes. Genome bins of *Chlorobia* encoding *cyc2* co-occurred consistently with bins of *Chlorobia* encoding sulfide oxidation genes (and lacking *cyc2*) at similar relative abundances across all four analyzed Lake 227 metagenomes (Fig. 3A). Such coexistence aligns well with observations of co-occurring sulfide-oxidizing phototrophs and photoferrotrophs in Lake La Cruz [34] and agrees with existing evidence that phototrophic sulfide oxidation can occur in ferruginous systems [33]. Observations from enrichment culturing on ferrous iron-containing media suggest that sulfide oxidation among Boreal Shield lake *Chlorobia* is possible even when sulfide is at trace levels, generated directly from sulfate-reducing bacteria (Supplementary Figure 2). Such "cryptic sulfur cycling" has been reported in various other aquatic systems and is likely an ecologically common phenomenon beyond Boreal Shield lakes [84] and common among phototrophic processes even within low-sulfate environments. Cryptic iron cycling, which has been observed in other ferruginous lakes [36], might also be possible in Boreal Shield lakes but was not observed directly in this study.

The lack of detectable *Chlorobia*-affiliated *cyc2* genes in Lake 442 metagenomes has important implications for the ecology of *cyc2*-encoding *Chlorobia*. Although genome bins of *Chlorobia* were estimated to occur at high relative abundance in Lake 442 based on metagenome data (i.e., 6.5% at 15 m depth in 2014*)*, genome bins only encoded sulfide oxidation genes, not *cyc2* (Fig. 3A). Lake 442 has similar underlying geology to Lake 227 and is nearby (~13.5 km distance). This suggests that the favourability of *cyc2*-encoding *Chlorobia* may be strongly influenced by physicochemical drivers within the natural range possible among Boreal Shield lakes. Generally, levels of sulfate in the water column are higher in Lake 442 than Lake 227 (i.e., ~15 μM in the anoxic zone compared to ~7.5 μM [37]), and total dissolved iron concentrations in Lake 442 tend to be lower at the depths where populations of *Chlorobia* are present (i.e., ~5-15

µM compared to ~10-100 µM in Lake 227). Lake 304, from which the *cyc2*-encoding "*Ca.* Chl. canadense" was enriched, has sulfate and total dissolved iron concentrations in its upper anoxic zone (~10 uM and ~40-200 uM, respectively; unpublished data) that are closer to Lake 227 values than Lake 442 values. It is possible that these factors, or other factors such as hydrogen

560 and bicarbonate concentrations (observed to impact photoferrotrophy rates in *Rps. palustris* TIE-1 [85]), lake bathymetry, anoxia timing, pH, light quality, or dissolved organic carbon levels, could shift the geochemical balance in favour of *Chlorobia* encoding genes for phototrophic sulfide oxidation rather than *cyc2* in Lake 442. Although Lake 304 was historically fertilized until 1976, it is more likely, given Lake 304's rapid recovery after fertilization ceased and ~2.7 year

565 water residence time (see Methods), that other similarities between Lakes 227 and 304, such as lake bathymetry [86] or high sediment fluxes of phosphorous observed in Lake 304 even before artificial nutrient addition began [87], allow for the favourability of *cyc2*-encoding *Chlorobia* in these lakes more than legacy fertilization effects. Understanding key factors governing the favourability of *cyc2*-associated metabolism among a broader suite of Boreal Shield lakes could

570 prove highly valuable for predicting photoferrotrophic activity in both modern and Archaean ocean settings.

The lack of *Chlorobia*-affiliated *cyc2* genes in Lake 442 metagenomes also has implications for the interpretation of iron isotope data. The strong iron isotope fractionation observed between the oxic and anoxic zone boundaries of Lakes 227 and 442 previously by

575 Schiff and colleagues [37] appears to occur regardless of whether *Chlorobia*-affiliated *cyc2* genes are detectable, implying that this signature cannot be used as a sole indicator of photoferrotrophic metabolism. It is possible that microaerophilic iron oxidation could contribute to the observed iron isotope fractionation in Lake 442 given the detection of a *cyc2*-encoding genome bin within the *Gallionelles* genus in Lake 442 metagenomic data (Fig. 3B). However, other yet unknown

580   iron cycling processes might also be at work. More nuanced physico-chemical measurements or

knowledge of additional biological processes are likely needed to robustly predict the potential

for photoferrotrophy in modern and potentially ancient ferruginous environments.

We were unable to induce photoferrotrophy in "*Ca.* Chl. canadense" (Fig. 4), which was

enriched in the present study from Lake 304. This new species, along with the Lake 227

585   enrichment that was lost in early subculturing, encoded both the *cyc2* gene and the sulfide

oxidation gene pathway, suggesting an ability to switch between phototrophic sulfide oxidation

and photoferrotrophy, as observed for *Chl.* sp. N1. A lack of observed photoferrotrophic growth

in "*Ca.* Chl. canadense" might be a result of several factors. We noted that the c5 family

cytochrome encoded by "*Ca.* Chl. canadense" directly upstream of *cyc2* contains an additional

590   cysteine residue in the heme-binding motif in its primary sequence (i.e., CCXCH rather than the

typical CXXCH), unlike the primary sequences of the c5 family cytochromes in reference

cultures or environmental genome bins (Supplementary File 6). This additional cysteine might

not completely compromise the function of the protein but could have deleterious consequences

for iron oxidation [88]. If this gene neighbour to *cyc2* is involved in the same metabolic pathway

595   as *cyc2*, it could imply that *cyc2* in "*Ca.* Chl. canadense" has an impaired or alternative function.

Alternative functions could include electron exchange with humic substances as hypothesized by

He and colleagues [89] or even manganese oxidation as reported recently for *Chlorobia* in mixed

culture [90], although direct links between Cyc2 and oxidation of alternate substrates to ferrous

iron have not been demonstrated experimentally. Cyc2 might also allow for direct electron uptake

600   from solid-phase conductive substances, as has been observed in the photoferrotroph *Rps.*

palustris TIE-1 [91]. Some of these alternative potential functions of Cyc2 might be preserved or

even enhanced even if iron oxidation potential is impaired. Given that strong ecological controls

appear to exist on *cyc2*-encoding *Chlorobia* (see above), it is also possible that the *cyc2* gene

product in "*Ca*. Chl. canadense" is indeed involved in iron oxidation but requires carefully tuned

605   conditions to switch metabolic modes. This option would align with current research that

implicates *cyc2* in iron oxidation among neutrophilic iron-oxidizing bacteria (e.g., [20]). It would

also align with preliminary reports that the *cyc2*-encoding but sulfide-oxidizing *Chl. luteolum* is

capable of changing to photoferrotrophic growth (Kate Thompson and Sean Crowe, personal

communication), despite previous difficulties inducing this behaviour [23]. The "*Ca*. Chl.

610   canadense" culture provides a unique opportunity to probe the regulation and function of *cyc2*

within the class *Chlorobia* in future work.

       The addition of the genome bins of *Chlorobia* from this study to existing reference

genomes doubles the known diversity of *Chlorobia*-associated *cyc2* genes and allows for further

examination of the evolutionary history of *cyc2* in the class *Chlorobia*. Rather than forming a

615   single monophyletic grouping, the presence of *cyc2* in genomes of *Chlorobia* appears

polyphyletic based on the ribosomal protein phylogeny (Fig. 2), interspersed with genomes

encoding various forms of sulfur oxidation. The thiosulfate oxidation (*sox*) gene system among

Chlorobia is known to not follow an inheritance pattern matching taxonomic marker genes, and

lateral gene transfer within the *Chlorobia* may have been an important factor in its acquisition

620   [25, 92]. However, the *cyc2* predicted primary sequence phylogeny among *Chlorobia* (Fig. 1C) is

somewhat congruent to the *Chlorobia* ribosomal protein phylogeny, suggesting that loss from a

single common ancestor, rather than extensive lateral gene transfer within the class, may account

for the majority of how *cyc2* genes are distributed within the *Chlorobia* class (Supplementary

Figure 4). Such a scenario would imply that *cyc2* was acquired by a common ancestor of modern

625   *Chlorobia* but multiple clades of modern *Chlorobia* lost the *cyc2* gene over evolutionary time.

Recovery of additional *cyc2*-containing *Chlorobia* genomes is required to confirm this

hypothesis, but it is an intriguing example of the types of evolutionary analyses possible as *cyc2* among the *Chlorobia* is further studied at the genomic level.

Our research provides insight into the metabolic diversity and ecology of anoxygenic

630 photoautotrophs in seasonally anoxic Boreal Shield lakes. We show that high relative abundance genome bins of *Chlorobia* in lake metagenomes can have varying functional gene content with respect to their photosynthetic electron donor, including some genome bins from Lake 227 samples that encoded the *cyc2* gene implicated in photoferrotrophy. The absence of *Chlorobia*-affiliated *cyc2* genes in the neighbouring Lake 442 suggests that ecological or geochemical

635 controls strongly influence the favourability of *cyc2*-based metabolism in Boreal Shield lakes, with potential impacts on whole microbial community function in the anoxic zone. Although we were unable to induce photoferrotrophy in a laboratory setting, we enriched the novel *cyc2*-encoding species "*Ca.* Chl. canadense", which could be used in future research exploring the function and regulation of *cyc2* as genetic studies continue to progress in linking the *cyc2* gene

640 product to its cellular role [93]. Probing the metabolic diversity of anoxygenic phototrophs in Boreal Shield lakes could lead to novel inferences about photosynthesis in early Earth oceans, including how environmental variables in the range possible for Archaean oceans could modulate the dominant mode of photosynthesis. Altogether, our findings serve as an important basis for future work probing the rates, gene expression, biogeography, and enrichment culture activity of

645 *cyc2*-encoding *Chlorobia* in Boreal Shield lakes. Although further work is needed to demonstrate the role of the *cyc2* gene product in photoferrotrophy and the distribution of *cyc2* across a broader suite of Boreal Shield lakes, our data indicate that metabolic diversity among anoxygenic photoautotrophs may be an overlooked yet ecologically important phenomenon in Boreal Shield lakes globally. Knowledge of such metabolic diversity impacts our understanding of modern

650 ecosystems and predictions about the evolution of early life on Earth.

Data availability

Raw metagenome reads for the freshwater lake and enrichment culture metagenomes are available in the NCBI sequence read archive (SRA) under BioProjects PRJNA518727 and

655   PRJNA534305, respectively. Six curated genome bins of *Chlorobia* are available under the same BioProjects, along with all assembled contigs from freshwater lake metagenomes. The complete set of uncurated genome bins were uploaded to a Zenodo repository at doi:10.5281/zenodo.2720705. Code for downloading these data and performing the analyses presented in this paper is available at https://github.com/jmtsuji/Chlorobia-cyc2-genomics

660   (doi:10.5281/zenodo.3228523).

Acknowledgements

We thank R. Henderson and staff at the IISD-ELA for assistance in field sampling and interpretation of the dynamics of ELA lakes. In addition, we thank S. A. Crowe for valuable

665   feedback on the manuscript, K. J. Thompson for helpful discussion related to the work, L. H. Bergstrand for assistance with bioinformatics, and K. E. Engel, R. C. Beaver, and N. A. Shaw for assistance with laboratory tasks. JMT thanks M. S. M. Jetten, C. U. Welte, and staff at the Soehngen Institute of Anaerobic Microbiology (SIAM), for providing helpful training for anaerobic microbiology methods. This research was supported by Discovery and Strategic

670   Projects Grants from the National Sciences and Engineering Research Council of Canada (NSERC).

Author contributions

SLS, JJV, and LM were involved in the overall lake sampling project that led to this work

675   and provided input and assistance with sample collection and data interpretation. JMT and JDN

designed the experimental work in this study. JMT, NT, and MT performed enrichment

cultivation with support from SH. JMT analyzed the sequence data and drafted the manuscript

with the edits and feedback of all authors.

680   **Competing interests**

The authors declare no competing financial interests.

Supplementary information is available at The ISME Journal's website.

685 **Supplementary information**

Supplementary Methods

Supplementary Figure 1 – Average nucleotide identity between recovered genome bins and

reference genomes of *Chlorobia*. The same ribosomal protein tree as in Figure 2 is shown to the

690 left of the heatmap.

Supplementary Figure 2 – Photographs of enrichment cultures of *Chlorobia* in ferrous iron- and

sulfide-containing media.

695 Supplementary Figure 3 – Genomic potential for photosynthesis and carbon fixation in recovered

genome bins of *Chlorobia* compared to reference strains. The figure layout and phylogenetic tree

are identical to Figure 2. All query sequences for reciprocal BLASTP were derived from the

genome of *Chl. tepidum* TLS except for BciB, which was derived from the genome of *Chl.*

ferrooxidans KoFox.

Supplementary Figure 4 – Tanglegram comparing the concatenated ribosomal protein phylogeny

(Fig. 2) and Cyc2 phylogeny (Fig. 1C) among *Chlorobia*. The two tips whose placements differ

between the two phylogenies are highlighted in blue or green.

705 Supplementary Figure 5 – Bubble plot showing taxonomic and functional profiling of

unassembled metagenome data, in the same general layout as Figure 3. HMM hit counts were

normalized to the total hits of *rpoB*, a single-copy taxonomic marker gene, within each sample,

such that HMM hits are expressed as a percentage of *rpoB*. All families (NCBI taxonomy) with >

1% of normalized hits to the HMMs are shown. Although not shown, normalized *Chlorobiaceae*-

710 associated *cyc2* hits for Lake 227 at 8 m were 0.9% for both 2013 and 2014.

Supplementary Figure 6 – Iron oxidation activity of cultures of *Chlorobia* over an extended

incubation period of 21 days. The figure layout is identical to Figure 4.

715 Supplementary File 1 – ZIP file containing the YAML config files for the ATLAS metagenome

assemblies performed for this publication.

Supplementary File 2 – ZIP file of two alternative profile Hidden Markov Models for the *cyc2*

gene (general or *Chlorobia*-specific).

Supplementary File 3 – Excel file containing a summary of basic physical, sampling, and historic

parameters for Lakes 227, 442, and 304.

Supplementary File 4 – Excel file containing accessions of reference genomes of *Chlorobia* and

725 statistics from the reciprocal BLAST search of Fe/S gene pathways.

Supplementary File 5 – CSV file containing genome statistics for all dereplicated MAGs from

this study.

730 Supplementary File 6 – ZIP file of amino acid multiple sequence alignments: alignment of

Chlorobia-associated Cyc2 predicted primary sequences from this study with reference sequences

(unmasked and masked), alignment of c5 family cytochromes presented in this study, and alignment of additional low-homology Cyc2 hits from metagenomes.

735 Supplementary File 7 – ZIP file of purity data for the "*Ca.* Chl. canadense" culture, including an epifluorescence microscopy image of the culture and an amplicon sequence variant (ASV) table summarizing the percent relative abundances of members of the enrichment culture.

References

1.   Van Gemerden H, Mas J. Ecology of phototrophic sulfur bacteria. In: Blankenship RE, Madigan MT, Bauer CE (eds). *Anoxygenic Photosynthetic Bacteria*. 1995. Springer Netherlands, Dordrecht, pp 49–85.

2.   Tank M, Thiel V, Ward DM, Bryant DA. A panoply of phototrophs: an overview of the thermophilic chlorophototrophs of the microbial mats of alkaline siliceous hot springs in Yellowstone National Park, WY, USA. In: Hallenbeck PC (ed). *Modern Topics in the Phototrophic Prokaryotes: Environmental and Applied Aspects*. 2017. Springer International Publishing Switzerland, pp 87–137.

3.   Manske AK, Glaeser J, Kuypers MMM, Overmann J. Physiology and phylogeny of green sulfur bacteria forming a monospecific phototrophic assemblage at a depth of 100 meters in the Black Sea. *Appl Environ Microbiol* 2005; **71**: 8049–8060.

4.   Imhoff JF. The family *Chromatiaceae*. In: Rosenberg E, DeLong EF, Lory S, Stackebrandt E, Thompson F (eds). *The Prokaryotes*. 2014. Springer Berlin Heidelberg, pp 151–178.

5.   Imhoff JF. The family *Chlorobiaceae*. In: Rosenberg E, DeLong EF, Lory S, Stackebrandt E, Thompson F (eds). *The Prokaryotes*. 2014. Springer Berlin Heidelberg, pp 501–514.

6.   Griffin BM, Schott J, Schink B. Nitrite, an electron donor for anoxygenic photosynthesis. *Science* 2007; **316**: 1870–1870.

7.   Kulp TR, Hoeft SE, Asao M, Madigan MT, Hollibaugh JT, Fisher JC, et al. Arsenic(III) fuels anoxygenic photosynthesis in hot spring biofilms from Mono Lake, California. *Science* 2008; **321**: 967–970.

8.   Thiel V, Tank M, Bryant DA. Diversity of chlorophototrophic bacteria revealed in the omics era. *Annu Rev Plant Biol* 2018; **69**: 21–49.

9.   Ehrenreich A, Widdel F. Anaerobic oxidation of ferrous iron by purple bacteria, a new type of phototrophic metabolism. *Appl Environ Microbiol* 1994; **60**: 4517–4526.

10.   Camacho A, Walter XA, Picazo A, Zopfi J. Photoferrotrophy: remains of an ancient photosynthesis in modern environments. *Front Microbiol* 2017; **8**: 323.

11.   Poulton SW, Canfield DE. Ferruginous conditions: a dominant feature of the ocean through Earth's history. *Elements* 2011; **7**: 107–112.

12.   Kappler A, Pasquero C, Konhauser KO, Newman DK. Deposition of banded iron formations by anoxygenic phototrophic Fe(II)-oxidizing bacteria. *Geology* 2005; **33**: 865–868.

13.   Czaja AD, Johnson CM, Beard BL, Roden EE, Li W, Moorbath S. Biological Fe oxidation controlled deposition of banded iron formation in the ca. 3770Ma Isua Supracrustal Belt (West Greenland). *Earth Planet Sc Lett* 2013; **363**: 192–203.

14.   He S, Barco RA, Emerson D, Roden EE. Comparative genomic analysis of neutrophilic iron(II) oxidizer genomes for candidate genes in extracellular electron transfer. *Front Microbiol* 2017; **8**: 1584.

15.   Melton ED, Swanner ED, Behrens S, Schmidt C, Kappler A. The interplay of microbially mediated and abiotic reactions in the biogeochemical Fe cycle. *Nat Rev Micro* 2014; **12**: 797–808.

16.   Castelle C, Guiral M, Malarte G, Ledgham F, Leroy G, Brugna M, et al. A new iron-oxidizing/O_2-reducing supercomplex spanning both inner and outer membranes, isolated from the extreme acidophile *Acidithiobacillus ferrooxidans*. *J Biol Chem* 2008; **283**: 25803–25811.

17.   Barco RA, Emerson D, Sylvan JB, Orcutt BN, Meyers MEJ, Ramírez GA, et al. New insight into microbial iron oxidation as revealed by the proteomic profile of an obligate iron-oxidizing chemolithoautotroph. *Appl Environ Microbiol* 2015; **81**: 5927–5937.

785   18.   Kato S, Ohkuma M, Powell DH, Krepski ST, Oshima K, Hattori M, et al. Comparative genomic insights into ecophysiology of neutrophilic, microaerophilic iron oxidizing bacteria. *Front Microbiol* 2015; **6**: 1265.

19.   Gupta D, Sutherland MC, Rengasamy K, Meacham JM, Kranz RG, Bose A. Photoferrotrophs produce a PioAB electron conduit for extracellular electron uptake. *mBio* 2019;

790   **10**: e02668-19.

20.   McAllister SM, Polson SW, Butterfield DA, Glazer BT, Sylvan JB, Chan CS. Validating the Cyc2 neutrophilic Fe oxidation pathway using meta-omics of Zetaproteobacteria iron mats at marine hydrothermal vents. *bioRxiv* 2019; 722066.

21.   Heising S, Richter L, Ludwig W, Schink B. *Chlorobium ferrooxidans* sp. nov., a

795   phototrophic green sulfur bacterium that oxidizes ferrous iron in coculture with a "*Geospirillum*" sp. strain. *Arch Microbiol* 1999; **172**: 116–124.

22.   Llirós M, García–Armisen T, Darchambeau F, Morana C, Triadó–Margarit X, Inceoğlu Ö, et al. Pelagic photoferrotrophy and iron cycling in a modern ferruginous basin. *Sci Rep* 2015; **5**: 13803.

800   23.   Laufer K, Niemeyer A, Nikeleit V, Halama M, Byrne JM, Kappler A. Physiological characterization of a halotolerant anoxygenic phototrophic Fe(II)-oxidizing green-sulfur bacterium isolated from a marine sediment. *FEMS Microbiol Ecol* 2017; **93**: fix054.

24.   Lambrecht N, Link to external site this link will open in a new window. Insights into early Earth ocean biogeochemistry from intensive monitoring of two ferruginous meromictic

805   lakes. 2019. Ph.D., Iowa State University.

25.   Frigaard N-U, Bryant DA. Genomic insights into the sulfur metabolism of phototrophic green sulfur bacteria. In: Hell R, Dahl DC, Knaff D, Leustek T (eds). *Sulfur Metabolism in Phototrophic Organisms*. 2008. Springer Netherlands, pp 337–355.

26.   Crowe SA, Hahn AS, Morgan-Lang C, Thompson KJ, Simister RL, Llirós M, et al. Draft genome sequence of the pelagic photoferrotroph *Chlorobium phaeoferrooxidans*. *Genome Announc* 2017; **5**: e01584-16.

27.   Thompson KJ, Simister RL, Hahn AS, Hallam SJ, Crowe SA. Nutrient acquisition and the metabolic potential of photoferrotrophic *Chlorobi*. *Front Microbiol* 2017; **8**: 1212.

28.   Bryce C, Blackwell N, Straub D, Kleindienst S, Kappler A. Draft genome sequence of *Chlorobium* sp. strain N1, a marine Fe(II)-oxidizing green sulfur bacterium. *Microbiol Resour Announc* 2019; **8**: e00080-19.

29.   Koeksoy E, Halama M, Konhauser KO, Kappler A. Using modern ferruginous habitats to interpret Precambrian banded iron formation deposition. *Int J Astrobiol* 2016; **15**: 205–217.

30.   Savvichev AS, Kokryatskaya NM, Zabelina SA, Rusanov II, Zakharova EE, Veslopolova EF, et al. Microbial processes of the carbon and sulfur cycles in an ice-covered, iron-rich meromictic lake Svetloe (Arkhangelsk region, Russia). *Environ Microbiol* 2017; **19**: 659–672.

31.   Lambrecht N, Katsev S, Wittkop C, Hall SJ, Sheik CS, Picard A, et al. Biogeochemical and physical controls on methane fluxes from two ferruginous meromictic lakes. *Geobiology* 2019; **0**: 1–16.

32.   Crowe SA, Jones C, Katsev S, Magen C, O'Neill AH, Sturm A, et al. Photoferrotrophs thrive in an Archean Ocean analogue. *Proc Nat Acad Sci USA* 2008; **105**: 15938–15943.

33.   Crowe SA, Maresca JA, Jones C, Sturm A, Henny C, Fowle DA, et al. Deep-water anoxygenic photosythesis in a ferruginous chemocline. *Geobiology* 2014; **12**: 322–339.

34.   Walter XA, Picazo A, Miracle MR, Vicente E, Camacho A, Aragno M, et al. Phototrophic Fe(II)-oxidation in the chemocline of a ferruginous meromictic lake. *Front Microbiol* 2014; **5**: 713.

35.   Morana C, Roland FAE, Crowe SA, Llirós M, Borges AV, Darchambeau F, et al. Chemoautotrophy and anoxygenic photosynthesis within the water column of a large meromictic tropical lake (Lake Kivu, East Africa). *Limnol Oceanogr* 2016; **61**: 1424–1437.

835   36.   Berg JS, Jézéquel D, Duverger A, Lamy D, Laberty-Robert C, Miot J. Microbial diversity involved in iron and cryptic sulfur cycling in the ferruginous, low-sulfate waters of Lake Pavin. *PLOS ONE* 2019; **14**: e0212787.

37.   Schiff SL, Tsuji JM, Wu L, Venkiteswaran JJ, Molot LA, Elgood RJ, et al. Millions of Boreal Shield lakes can be used to probe Archaean Ocean biogeochemistry. *Sci Rep* 2017; **7**:

840   46708.

38.   Schindler DW, Armstrong FAJ, Holmgren SK, Brunskill GJ. Eutrophication of Lake 227, Experimental Lakes Area, northwestern Ontario, by addition of phosphate and nitrate. *J Fish Res Bd Can* 1971; **28**: 1763–1782.

39.   Campbell P. Phosphorus budgets and stoichiometry during the open-water season in two

845   unmanipulated lakes in the Experimental Lakes Area, northwestern Ontario. *Can J Fish Aquat Sci* 1994; **51**: 2739–2755.

40.   White III RA, Brown J, Colby S, Overall CC, Lee J-Y, Zucker J, et al. ATLAS (Automatic Tool for Local Assembly Structures) - a comprehensive infrastructure for assembly, annotation, and genomic binning of metagenomic and metatranscriptomic data. *PeerJ Preprints* 2017; **5**:

850   e2843v1.

41.   Jeans C, Singer SW, Chan CS, VerBerkmoes NC, Shah M, Hettich RL, et al. Cytochrome 572 is a conspicuous membrane protein with iron oxidation activity purified directly from a natural acidophilic microbial community. *ISME J* 2008; **2**: 542–550.

42.   Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, et al. Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Mol Syst Biol* 2011; **7**: 539.

43.   Eddy SR. Accelerated profile HMM searches. *PLOS Comput Biol* 2011; **7**: e1002195.

44.   Talavera G, Castresana J. Improvement of phylogenies after removing divergent and ambiguously aligned blocks from protein sequence alignments. *Syst Biol* 2007; **56**: 564–577.

45.   Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol* 2015; **32**: 268–274.

46.   Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat Methods* 2017; **14**: 587–589.

47.   Olm MR, Brown CT, Brooks B, Banfield JF. dRep: a tool for fast and accurate genomic comparisons that enables improved genome recovery from metagenomes through de-replication. *ISME J* 2017; **11**: 2864–2868.

48.   Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res* 2015; **25**: 1043–1055.

49.   Eren AM, Esen ÖC, Quince C, Vineis JH, Morrison HG, Sogin ML, et al. Anvi'o: an advanced analysis and visualization platform for 'omics data. *PeerJ* 2015; **3**: e1319.

50.   Rissman AI, Mau B, Biehl BS, Darling AE, Glasner JD, Perna NT. Reordering contigs of draft genomes using the Mauve Aligner. *Bioinformatics* 2009; **25**: 2071–2073.

51.   Seemann T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 2014; **30**: 2068–2069.

52.   Armstrong FAJ, Schindler DW. Preliminary chemical characterization of waters in the Experimental Lakes Area, northwestern Ontario. *J Fish Res Bd Can* 1971; **28**: 171–187.

53.   Schindler DW. Eutrophication and recovery in experimental lakes: implications for lake management. *Science* 1974; **184**: 897–899.

54.   Schindler D. The coupling of elemental cycles by organisms: evidence from whole-lake chemical perturbations. In: Stumm W (ed). *Chemical Processes in Lakes*. 1985. John Wiley and Sons, New York, NY, pp 225–250.

55.   Schindler DW. The dilemma of controlling cultural eutrophication of lakes. *Proceedings of the Royal Society B: Biological Sciences* 2012; **279**: 4322–4333.

56.   Curtis PJ, Schindler DW. Hydrologic control of dissolved organic matter in low-order Precambrian Shield lakes. *Biogeochemistry* 1997; **36**: 125–138.

57.   Hegler F, Posth NR, Jiang J, Kappler A. Physiology of phototrophic iron(II)-oxidizing bacteria: implications for modern and ancient environments. *FEMS Microbiol Ecol* 2008; **66**: 250–260.

58.   Milucka J, Kirf M, Lu L, Krupke A, Lam P, Littmann S, et al. Methane oxidation coupled to oxygenic photosynthesis in anoxic waters. *ISME J* 2015; **9**: 1991–2002.

59.   Clavier CGJ, Boucher G. The use of photosynthesis inhibitor (DCMU) for in situ metabolic and primary production studies on soft bottom benthos. *Hydrobiologia* 1992; **246**: 141–145.

60.   Jain C, Rodriguez-R LM, Phillippy AM, Konstantinidis KT, Aluru S. High throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nat Commun* 2018; **9**: 5114.

61.   Lee MD. GToTree: a user-friendly workflow for phylogenomics. *Bioinformatics* 2019; btz188.

62.   Hug LA, Baker BJ, Anantharaman K, Brown CT, Probst AJ, Castelle CJ, et al. A new view of the tree of life. *Nat Microbiol* 2016; **1**: 16048.

63.   Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. *Journal of molecular biology* 1990; **215**: 403–410.

905   64.   Bergstrand LH, Cardenas E, Holert J, Hamme JDV, Mohn WW. Delineation of steroid-degrading microorganisms through comparative genomic analysis. *mBio* 2016; **7**: e00166-16.

65.   Bryant DA, Liu Z, Li T, Zhao F, Costas AMG, Klatt CG, et al. Comparative and functional genomics of anoxygenic green bacteria from the taxa *Chlorobi*, *Chloroflexi*, and *Acidobacteria*. In: Burnap R, Vermaas W (eds). *Functional Genomics and Evolution of*

910   *Photosynthetic Systems*. 2012. Springer Netherlands, Dordrecht, pp 47–102.

66.   Tourova TP, Kovaleva OL, Gorlenko VM, Ivanovsky RN. Use of genes of carbon metabolism enzymes as molecular markers of *Chlorobi* phylum representatives. *Microbiology* 2013; **82**: 784–793.

67.   Parks DH, Chuvochina M, Waite DW, Rinke C, Skarshewski A, Chaumeil P-A, et al. A

915   standardized bacterial taxonomy based on genome phylogeny substantially revises the tree of life. *Nat Biotechnol* 2018.

68.   Chaumeil P-A, Mussig AJ, Hugenholtz P, Parks DH. GTDB-Tk: a toolkit to classify genomes with the Genome Taxonomy Database. *Bioinformatics* 2019; btz848.

69.   Fish JA, Chai B, Wang Q, Sun Y, Brown CT, Tiedje JM, et al. FunGene: the functional

920   gene pipeline and repository. *Front Microbiol* 2013; **4**: 291.

70.   Hyatt D, Chen G-L, LoCascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* 2010; **11**: 1–11.

71.	Petrenko P, Lobb B, Kurtz DA, Neufeld JD, Doxey AC. MetAnnotate: function-specific taxonomic profiling and comparison of metagenomes. *BMC Biol* 2015; **13**: 1–8.

72.	Pfennig N. *Rhodocyclus purpureus* gen. nov. and sp. nov., a ring-shaped, vitamin B12-requiring member of the family *Rhodospirillaceae*. *Int J Syst Evol Microbiol* 1978; **28**: 283–288.

73.	Tank M, Bryant DA. Nutrient requirements and growth physiology of the photoheterotrophic Acidobacterium, *Chloracidobacterium thermophilum*. *Front Microbiol* 2015; **6**.

74.	Peng C, Bryce C, Sundman A, Borch T, Kappler A. Organic matter complexation promotes Fe(II) oxidation by the photoautotrophic Fe(II)-oxidizer *Rhodopseudomonas palustris* TIE-1. *ACS Earth Space Chem* 2019.

75.	Stookey LL. Ferrozine - a new spectrophotometric reagent for iron. *Anal Chem* 1970; **42**: 779–781.

76.	Verschoor MJ, Molot LA. A comparison of three colorimetric methods of ferrous and total reactive iron measurement in freshwaters. *Limnol Oceanogr Methods* 2013; **11**: 113–125.

77.	Yang J, Yan R, Roy A, Xu D, Poisson J, Zhang Y. The I-TASSER Suite: protein structure and function prediction. *Nat Methods* 2015; **12**: 7–8.

78.	Overmann J, Coolen MJL, Tuschak C. Specific detection of different phylogenetic groups of chemocline bacteria based on PCR and denaturing gradient gel electrophoresis of 16S rRNA gene fragments. *Arch Microbiol* 1999; **172**: 83–94.

79.	Mori Y, Purdy KJ, Oakley BB, Kondo R. Comprehensive detection of phototrophic sulfur bacteria using PCR primers that target reverse dissimilatory sulfite reductase gene. *Microbes Environ* 2010; **25**: 190–196.

80.   Finneran KT, Johnsen CV, Lovley DR. *Rhodoferax ferrireducens* sp. nov., a psychrotolerant, facultatively anaerobic bacterium that oxidizes acetate with the reduction of Fe(III). *Int J Syst Evol Micr* 2003; **53**: 669–673.

81.   Watanabe T, Kojima H, Fukui M. Complete genomes of freshwater sulfur oxidizers *Sulfuricella denitrificans* skB26 and *Sulfuritalea hydrogenivorans* sk43H: genetic insights into the sulfur oxidation pathway of betaproteobacteria. *Syst Appl Microbiol* 2014; **37**: 387–395.

82.   Kuever J. The family *Desulfobacteraceae*. In: Rosenberg E, DeLong EF, Lory S, Stackebrandt E, Thompson F (eds). *The Prokaryotes: Deltaproteobacteria and Epsilonproteobacteria*. 2014. Springer Berlin Heidelberg, pp 45–73.

83.   Karhunen J, Arvola L, Peura S, Tiirola M. Green sulphur bacteria as a component of the photosynthetic plankton community in small dimictic humic lakes with an anoxic hypolimnion. *Aquat Microb Ecol* 2013; **68**: 267–272.

84.   Kappler A, Bryce C. Cryptic biogeochemical cycles: unravelling hidden redox reactions. *Environ Microbiol* 2017; **19**: 842–846.

85.   Croal LR, Jiao Y, Kappler A, Newman DK. Phototrophic Fe(II) oxidation in an atmosphere of H2: implications for Archean banded iron formations. *Geobiology* 2009; **7**: 21–24.

86.   Brunskill GJ, Schindler DW. Geography and bathymetry of selected lake basins, Experimental Lakes Area, northwestern Ontario. *J Fish Res Bd Can* 1971; **28**: 139–155.

87.   Mathias JA, Barica J. Factors controlling oxygen depletion in ice-covered lakes. *Can J Fish Aquat Sci* 1980; **37**: 185–194.

88.   Allen JWA, Sawyer EB, Ginger ML, Barker PD, Ferguson SJ. Variant c-type cytochromes as probes of the substrate specificity of the E. coli cytochrome c maturation (Ccm) apparatus. *Biochem J* 2009; **419**: 177–186.

89.   He S, Lau MP, Linz AM, Roden EE, McMahon KD. Extracellular electron transfer may be an overlooked contribution to pelagic respiration in humic-rich freshwater lakes. *mSphere* 2019; **4**: e00436-18.

90.   Daye M, Klepac-Ceraj V, Pajusalu M, Rowland S, Farrell-Sherman A, Beukes N, et al. Light-driven anaerobic microbial oxidation of manganese. *Nature* 2019; **576**: 311–314.

91.   Guzman MS, Rengasamy K, Binkley MM, Jones C, Ranaivoarisoa TO, Singh R, et al. Phototrophic extracellular electron uptake is linked to carbon dioxide fixation in the bacterium *Rhodopseudomonas palustris*. *Nat Commun* 2019; **10**: 1355.

92.   Meyer B, Imhoff JF, Kuever J. Molecular analysis of the distribution and phylogeny of the *soxB* gene among sulfur-oxidizing bacteria – evolution of the Sox sulfur oxidation enzyme system. *Environ Microbiol* 2007; **9**: 2957–2977.

93.   Chan C, McAllister SM, Garber A, Hallahan BJ, Rozovsky S. Fe oxidation by a fused cytochrome-porin common to diverse Fe-oxidizing bacteria. *bioRxiv* 2018; 228056.

94.   Petersen TN, Brunak S, von Heijne G, Nielsen H. SignalP 4.0: discriminating signal peptides from transmembrane regions. *Nat Methods* 2011; **8**: 785–786.

95.   Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, et al. UCSF Chimera—a visualization system for exploratory research and analysis. *J Comput Chem* 2004; **25**: 1605–1612.

Table and figure legends

Table 1. Quality statistics for genome bins of *Chlorobia* recovered in this study. Completeness and contamination were calculated using CheckM (see Methods). Note that no ribosomal RNA genes were recovered for the bins. A summary of bin statistics for lower-quality genome bins is

995   included in Supplementary File 5.

Figure 1. Recovered *cyc2* homologues of *Chlorobia* from metagenomes. **A**, Homology model of the Cyc2 protein from *Chl. phaeoferrooxidans* KB01, generated by I-TASSER based on sequence homology (with leading 18 signal peptides trimmed according to SignalP [94]) and visualized

1000   using UCSF Chimera [95]. The porin-like beta barrel structure is highlighted in dark blue, and the location of the single N-terminal heme-binding motif on the outer membrane portion of the structure is indicated by a black arrow. **B**, Genomic context of the recovered *cyc2* homologues. A ~10 kb region is shown of assembled contigs surrounding the homologues. The *cyc2* and adjacent c5 family cytochrome genes are highlighted in green and pink, respectively. Other shared genes

1005   between contigs are highlighted in yellow. Remaining genes are coloured grey if they lacked a functional annotation (e.g., no significant BLASTP hit, or hit to a domain of unknown function) or are coloured purple if their closest BLASTP hit corresponded to a gene associated with the *Chlorobiaceae* family. One gene in the 10 kb region had its closest BLASTP hit to a gene associated with the *Comamonadaceae* family (*Betaproteobacteria*; ~51% amino acid identity);

1010   this gene is highlighted in blue. **C**, Sequence comparison of recovered *cyc2* homologues to *cyc2* genes of known iron-oxidizing microorganisms. The phylogenetic tree was built from a 223 residue masked Cyc2 amino acid sequence alignment (see Methods). Bootstrap values over 50/100 are shown, and the scale bar represents the proportion of residue changes along the alignment. Due to the uncertain evolutionary history of *cyc2*, the phylogeny is midpoint-rooted. A

1015   green box highlights the monophyletic *Chlorobia* clade. Adjacent to the phylogeny, a subset of

the unmasked Cyc2 sequence alignment is shown (positions 40-80 of 609; N-terminus end).

Alignment positions having 100% sequence identity in the predicted heme-binding site (CXXCH

motif) are marked with red asterisks. Other positions with 100% sequence identity are marked

with black asterisks. The full sequence alignment is available in Supplementary File 6. Note that

1020   L227 2014 Bin 92 (name marked with an asterisk) was included in this figure for comparison of

its *cyc2* gene despite the bin quality being lower than the main *Chlorobia* genome bin set.

Abbreviations: *Chl. = Chlorobium*.

Figure 2. Iron- and sulfur-oxidizing genetic potential in recovered genome bins of *Chlorobia*

1025   compared to reference strains. The left side of the figure shows a maximum-likelihood

phylogenetic tree of the *Chlorobia* based on concatenated ribosomal protein amino acid

sequences (see methods). Bootstrap values over 50/100 are shown, and the scale bar represents

the percentage of residue changes across the 2243 amino acid alignment. Species of *Chlorobia*

are shaded based on their known or hypothesized metabolic potential. Genome bins of *Chlorobia*

1030   recovered from this study are bolded. On the right side, a heatmap is shown displaying the

presence/absence of genes implicated in iron and sulfur metabolism among *Chlorobia* based on

reciprocal BLASTP. Heatmap tiles are shaded based on the percent amino acid identity of an

identified gene compared to the reference sequence (*Chl. ferrooxidans* for iron-related genes, and

Chl. phaeoclathratiforme for sulfur-related genes). Although the cytochrome c5 family gene

1035   upstream of *cyc2* was not searched for directly due to poor sequence homology, the gene was

verified manually to be adjacent to all hits of *cyc2*. Abbreviations: *Chl. = Chlorobium*; *Cba. =*

Chlorobaculum; *Che. = Chloroherpeton*; *Ptc. = Prosthecochloris*.

Figure 3. Bubble plot showing predicted relative abundances of recovered genome bins in the lake and enrichment culture environments. The size of each bubble corresponds to the relative abundance of a genome bin or taxon within metagenome data based on mapping of assembled reads. Each bubble is also labelled with its percent relative abundance for clarity. **A**, Bins of *Chlorobia* coloured by their metabolic potential based on functional gene markers described in this study. Bolded names correspond to the higher-quality bins described in Table 1 and Fig. 2. In parentheses beside each name are bin quality statistics reported by CheckM – the predicted % completion, % contamination, and % strain heterogeneity, respectively. The displayed bins all classify to the *Chlorobium* genus based on GTDB taxonomy. **B**, Non-*Chlorobia* bins (≥ 0.01% relative abundance) found to have the same functional gene markers (*cyc2* or *dsrA*) in their genome sequences. Quality statistics are reported as in panel **A**. On the right side of the panel, the GTDB family and genus classifications of each bin are shown. **C**, Family-level taxonomic composition of the metagenomes based on GTDB classifications of genome bins with ≥ 1% relative abundance. The phylum of each family is displayed on the right side of the panel. For unresolved or non-Latin family names, the order (and class, if needed) is displayed in parentheses beside the family name for clarity. **D**, Assembly statistics for the metagenome data. The total number of quality-trimmed metagenome reads is represented by the total height of each bar. Reads that mapped to the filtered sequence assembly (i.e., excluding short contigs, see Methods) are highlighted in blue and are considered "assembled" reads. Assembled read totals were used to determine relative abundances of genome bins in panels A-C. Supplementary Figure 4 shows relative abundances based on predictions from unassembled metagenome reads for comparison.

Figure 4. Iron oxidation activity of two *Chlorobia* cultures under different growth conditions over time. The concentrations of ferrous iron, total iron, and ferric iron (determined by the

subtraction of ferrous iron from total iron) are shown over a 10 day incubation period for "*Ca.*

Chl. canadense" (enriched in this study) and *Chl. ferrooxidans*. The activity of the cultures in

1065   iron-amended media is shown when incubated in the dark with EDTA, in the light, and in the

light with EDTA. Each panel depicts the results of biological duplicates, with error bars

representing the standard deviation of technical duplicates in the ferrozine assay. The same results

are shown for the full 21-day incubation period in Supplementary Figure 6.

Figure 1

Figure 2

Figure 3

Figure 4