Evolution of Larval Segment Position across 12 *Drosophila* Species

Running title: Evolution of Segmentation across Drosophila

Gizem Kalay[1], Joel Atallah[1*], Noemie C. Sierra[2], Austin M. Tang[1], Amanda E. Crofton[1], Mohan K. Murugesan[1], Sherri Wykoff-Clary[1], Susan E. Lott[1]

[1]The Department of Evolution and Ecology, University of California, Davis, One Shields Avenue, Davis, CA 95616
[2]Earth and Planetary Sciences Department, University of California, Davis, One Shields Avenue, Davis, CA 95616

Corresponding authors: Email Gizem Kalay (gizemkly@gmail.com) or Susan E. Lott (selott@ucdavis.edu).

Author contributions: G.K. and S.E.L. designed the study. G.K., A.M.T., A.E.C., M.K.M. and S.W. collected the data. J. A. wrote the Python script that took measurements from samples in an automated manner. G.K and N.C.S. analyzed the data. G.K. and S.E.L. wrote the manuscript with input from J.A. and N.C.S.

Acknowledgements: We thank Lott Lab members Emily L. Cartwright, Anna A. Feitzinger, Charles S. Omura for comments on the manuscript; and Graham Coop for comments and statistical help. We also thank The National Drosophila Species Stock Center for the Drosophila species used in this project. This work was financially supported in part by National Institutes of Health grants, R01GM111362, R00GM098448, and start-up funds provided to S.E.L. by the University of California, Davis.

Data Archiving: All data and R code used for phylogenetic analyses for segment position are deposited in FigShare Repository at: https://doi.org/10.6084/m9.figshare.8085206. Data tables for segment position measurements are deposited in FigShare Repository at: https://doi.org/10.6084/m9.figshare.8170787. Python script for taking larval segment measurements is available at J.A.'s github: https://github.com/joelatallah/larval_imaging.

Competing interests: The authors declare no competing interests.

---

[*] current address: The Department of Biological Sciences, University of New Orleans, 2000 Lakeshore Drive, New Orleans, LA 70148

Abstract

Many developmental traits that are critical to the survival of the organism are also robust. These robust traits are resistant to phenotypic change in the face of variation. This presents a challenge to evolution. In this paper, we asked whether and how a well-established robust trait, *Drosophila* segment patterning, changed over the evolutionary history of the genus. We compared segment position scaled to body length at the first-instar larval stage among 12 *Drosophila* species. We found that relative segment position has changed many times across the phylogeny. Changes were frequent, but primarily small in magnitude. Phylogenetic analysis demonstrated that rates of change in segment position are variable along the *Drosophila* phylogenetic tree, and that these changes can occur in short evolutionary timescales. Correlation between position shifts of segments decreased as the distance between two segments increased, suggesting local control of segment position. The posterior-most abdominal segment showed the highest magnitude of change on average, had the highest rate of evolution between species, and appeared to be evolving more independently as compared to the rest of the segments. This segment was exceptionally elongated in the cactophilic species in our dataset, raising questions as to whether this change may be adaptive.

Keywords: segment patterning, *Drosophila*, evolution, robustness, larval stage

Introduction

Many developmental phenotypes are critical to the survival and fitness of the organism. These phenotypes are often observed to be robust, in that they produce a stereotyped outcome despite variation encountered during development (Wagner 2005; Félix and Wagner 2006; Masel and Siegal 2009; Siegal and Leu 2014). The variation experienced in ontogeny can come in a variety of forms, including stochastic, genetic, and environmental variation (Wagner 2005; Félix and Wagner 2006; Masel and Siegal 2009; Siegal and Leu 2014). Some robust phenotypes have specific mechanisms in place to ensure they are produced faithfully (Félix and Wagner 2006; Masel and Siegal 2009; Siegal and Leu 2014; Nijhout et al. 2017). However, these same robust phenotypes do evolve over periods of evolutionary time (Félix et al. 2000; Arthur and Chipman 2005; Lott et al. 2007; 2010; Fowlkes et al. 2011; Félix 2012; Combs and Fraser 2018). This poses a fundamental question: how do robust traits evolve if the phenotypic variation necessary to selection is suppressed?

One such developmental phenotype is segmentation along the head to tail (anterior-posterior) axis. Segmentation is the periodic repetition of anatomical structures. It is a shared feature of three big animal phyla, annelids, arthropods, and chordates (Davis and Patel 1999; Tautz 2004). In *Drosophila*, the foundation of segmentation is laid during the beginning of embryogenesis (St Johnston and Nüsslein-Volhard 1992), when development is under control of the maternal gene products, before the zygotic genome is activated. Maternal gene products that are located in the anterior and posterior parts of the fertilized egg set up a concentration gradient along

the length of the embryo (St Johnston and Nüsslein-Volhard 1992; Surkova et al. 2018).

These maternal gene products regulate one another, and also regulate genes

expressed later in development by the zygote (St Johnston and Nüsslein-Volhard 1992;

Surkova et al. 2018). This genetic network of regulators, which has been well-

established through decades of critical study (Nüsslein-Volhard and Wieschaus 1980;

Kornberg and Tabata 1993; Nasiadka et al. 2002; Nüsslein-Volhard et al. 2008; Clark

2017) precisely divide the embryo into progressively smaller subsections over

embryonic development, until the correct number of body segments is reached. At the

end of the embryonic stage, a first-instar larva with a highly organized segmented

pattern of differentiated structures is produced.

As long as segmentation has been investigated, its fundamental role in

Drosophila development has been clear, as defects in this process can be detrimental

(or lethal) to the organism (Wieschaus and Nuesslein-Volhard 2016). Indeed, the

lethality of homozygous mutations in segmentation genes was critical to their discovery

in mutant screens (Wieschaus and Nuesslein-Volhard 2016). Subsequent generations

of experiments using increasingly advanced methods have demonstrated that in

addition to being critical, segmentation is a very precise process, with measurements of

expression domains of segmentation genes being highly reproducible between embryos

(Houchmandzadeh et al. 2002; Gregor et al. 2007; Surkova et al. 2008; Jaeger 2010;

Petkova et al. 2014; Bentovim et al. 2017). Additionally, research has shown that

segmentation can proceed correctly, while keeping its precision, in the face of

substantial perturbations. For instance, in a seminal study launching the interface

between the gradient of the maternal gene *bicoid (bcd)* and its downstream target

hunchback (hb) as a model for understanding precision and scaling in developmental

signaling, Houchmandzadeh et al. (2002), also demonstrated that the relative position of

the Hb expression boundary (relative to embryo size) was robust to substantial genetic

and environmental perturbations. This study tested genetic perturbations in the form of

mutations in important maternal genes or absence of whole chromosomes, and found

very little or no variation in the relative position of Hb expression (Houchmandzadeh et

al. 2002). Moreover, the standard deviation of the Hb expression boundary did not show

a significant increase with these drastic perturbations, indicating that these genetic

changes did not decrease precision in the Hb boundary (Houchmandzadeh et al. 2002).

There is also substantial evidence that segmentation proceeds precisely in the face of

dramatic environmental perturbation. For example, when whole (Houchmandzadeh et

al. 2002) or two halves (Lucchetta et al. 2005) of *D. melanogaster* embryos were raised

at different temperature extremes after fertilization, inducing different developmental

rates, expression domains were formed in the same relative position and with the same

level of precision observed in control embryos, by the gap or pair rule stage of

embryonic development, respectively (Houchmandzadeh et al. 2002; Lucchetta et al.

2005).

The above studies, along with many others, show that *Drosophila* segmentation

is a complex trait that is robust to many forms of variation (Driever and Nüsslein-Volhard

1988; Houchmandzadeh et al. 2002; Lucchetta et al. 2005; Manu et al. 2009b). It is

perhaps unsurprising then that with the exception of some Hawaiian *Drosophila* species

(Spieth 1981), it is difficult to observe any gross differences in segment position, size

and shape, scaled to full body size, between adults of different *Drosophila* species.

However, over longer evolutionary time scales, segment patterning can vary

substantially between arthropod species (Regier et al. 2010). At the early embryonic

stage in *Drosophila*, previous studies have shown small quantitative differences in

relative position of segmentation gene expression domains between both closely related

(Lott et al. 2007; 2010; Fowlkes et al. 2011; Combs and Fraser 2018) and more

distantly related species (Fowlkes et al. 2011; Wunderlich et al. 2019). Within a species,

there is little evidence for variation in the relative position of segmentation gene

boundaries, with a few exceptions. For example, between *D. melanogaster* lines with

variable egg size, Lott et al. (2007) found no significant differences in relative position of

even-skipped (*eve*) stripe boundaries. Even when lines with extremes of egg size were

crossed to generate the full range of embryo lengths between the parental lines, relative

segment position was invariable between different genotypes. However, between *D.*

melanogaster lines that were subjected to strong artificial selection for egg size, Miles et

al. (2010) found few small quantitative differences in *eve* patterning using three-

dimensional imaging techniques. And, Jiang et al. (2015) was able to identify a line of *D.*

melanogaster from the *Drosophila* Genetic Reference Panel (Mackay et al. 2012) with

altered *even-skipped* pattern formation. It is not known, however, whether these

changes persist beyond the embryonic stage, or if these trends appear beyond the

limited number of species examined. So, while segment patterning is well documented

to be robust, it has also been demonstrated to have some small quantitative level of

variation between species, and in some circumstances, within species. Evolution of this

robust trait does occur, and the presence of some variation within species suggests that

evolution of small changes may be possible without causing catastrophic failure of

patterning. Thus, we hypothesized that the evolution of segmentation may occur by

relatively small shifts, rather than rare large leaps. This would require large sample

sizes to detect.

In order to investigate the evolution of segment patterning systematically across

a range of divergence times, we characterized segment position across 12 species of

Drosophila. These 12 species spanned the evolutionary history of the genus (40-60

million years; (Russo et al. 1995; Obbard et al. 2012; Russo et al. 2013) and included

three pairs of sister species. We measured position of each abdominal segment relative

to body length, referred to throughout the manuscript as relative segment position, in

the first-instar larvae (Fig. 1). Larval stage is the latest developmental stage where all

segments are visible and easy to measure, and the use of this stage facilitated the

measuring of more than a hundred larvae from each species. We found that relative

segment positions at the larval stage have changed many times in the evolutionary

history of *Drosophila*, and that these changes were mostly small in magnitude, with

some larger changes. Most species-pair comparisons showed differences in the relative

position of most abdominal segments. Most sister species were significantly different at

every segment, however some of the most diverged species in our dataset showed no

differences in their patterning. The magnitude of differences in relative position

increased towards the posterior segments in most species, most strikingly in *D.*

persimilis and *D. mojavensis*. Phylogenetic modelling showed that rate of segment

position change was highly variable among the branches along the *Drosophila*

phylogeny, and evolutionary rate changed even between closely related species.

Results of a correlation analysis for position change between segments within species

suggested local control of segment position, as correlations decrease with physical

distance. Correlations of changes in the rate of evolution of segment position between

branches of the phylogenetic tree also decreased with physical distance between

segments. Overall, these results demonstrate that this complex and robust

developmental trait does evolve, even over short timescales, and it does so primarily by

small frequent steps with occasional large leaps. This may permit the evolution of novel

patterns over long periods of evolutionary time, without compromising the integrity of

this critical developmental phenotype.

Materials and Methods

Species used

We took relative position measurements of abdominal segments from 12 *Drosophila*

species spanning the evolutionary history of the genus. These species were *D.*

melanogaster, *D. simulans*, *D. sechellia*, *D. yakuba*, *D. santomea*, *D. erecta*, *D.*

ananassae, *D. pseudoobscura*, *D. persimilis*, *D. willistoni*, *D. mojavensis wriglei* and *D.*

virilis (Fig. 2A). We used the sequenced lines from 11 of these species (Clark et al.

2007). For *D. santomea* we used stock #14021-0271.01 (The National *Drosophila*

Species Stock Center).

Fly husbandry and population size control

All flies were kept in plastic bottles in 20°C incubator with 60% humidity. Fly stocks were

raised on standard cornmeal food. The amount of time to sexual maturation varied

between species (Markow et al. 2008). Population control was conducted in a species-

specific manner through controlling the number of sexually mature adults added in a

given bottle, and how often the bottles were changed (Table S1). As a result, each

bottle ended up with 40-50 pupae. Adult flies were discarded 14 days after they became

fertile, to control for parental age.

Control experiment on the effect of the larval mounting procedure

In order to address whether the larval mounting procedure has species-specific effects

on measurements of relative segment position in first-instar *Drosophila* larvae, we

examined relative segment position in *D. melanogaster*, *D. sechellia* and *D. virilis* at

three different stages of the larval prep procedure: before heating, after heating, and

after mounting (Fig. S1). As these three species are variable in their egg (Markow et al.

2008) and larval size (Fig. 2C) and have varying phylogenetic distances (Fig. 2A), we

hypothesized that they might have a higher chance of being differentially affected by the

larval preparation protocol. First, first-instar larvae that hatched in distilled water were

temporarily immobilized using a 3-5 minute incubation on ice, put on a microscope slide

with water and then imaged using a Zeiss AxioImager microscope in brightfield. These

larvae were then put back in distilled water incubated at 60°C for 50 minutes and

imaged in water a second time using a brightfield microscope. The larvae were then

mounted in PVA Mounting Medium (BioQuip) (see next section). The larvae were then

imaged a third time using dark field microscopy (see "Imaging" section). In the images

of the iced (untreated control, the icing procedure slows larvae enough for imaging) and

then the heated larva, each of the 2nd and 3rd thoracic, plus eight abdominal segments

were marked by a node between bulges on the larval cuticle (Fig. S1). In the dark field

images, segment borders were determined by the anterior border of each denticle belt.

Denticle belts are rows of bristles on the ventral side of the larvae that are used for

traction while crawling (Bejsovec 2013). The anterior border of each denticle belt is a

proxy for the anterior border of each abdominal segment (Lohs-Schardin et al. 1979).

Using the Image J (version 1.47t) "Line tool", measurements were taken from the

anterior end of the larva to the border of each segment. Segment position was then

determined by dividing this value by the full length of the larvae (Fig. 1). Segment

position data was analyzed using the following linear models (using the "lm" function, all

terms were fixed), implemented in R (R Core Team 2019).

Relative segment position = μ + prep + species + prep x species + ε.

Relative segment position = μ + prep + species + segment + prep x species x segment

+ ε.

As a result, whether we included segment as a factor (p-value = 0.07 to 0.82) or not (p-

value = 0.42 to 0.83) in our linear model formula, we did not find any significant species-

specific effects of the preparation protocol on segment position measurements.

Preparing Drosophila first-instar larvae for imaging

For each species, 20-50 newly emerged adults were obtained from a population-

controlled bottle and put in an egg collection bottle with a cap containing glucose-agar

food. For *D. sechellia,* cornmeal fly food was used in an egg lay cap with yeast sprinkled

on it, as this prevents this species from withholding egg laying, a particular problem with

this species (Markow et al. 2008). The bottle was then placed in 20°C incubator upside

down. The next day eggs were collected from egg lay caps, put onto a mesh and

thoroughly washed using distilled water to remove residual yeast and egg lay cap food.

The eggs were then placed in a petri dish filled with distilled water. This petri dish was

placed in 20°C until the eggs developed into first-instar larvae. In addition to the

variability in the number of days necessary to reach sexual maturity, different species

also varied in the number of days necessary for a fertilized egg to develop into first-

instar larvae (Markow and O'Grady 2005). The petri dish with water and larvae was

then placed in 60°C oven for ~50 minutes (Table S1), which killed and straightened the

larvae. These larvae were then mounted on standard glass slides using PVA Mounting

Medium (BioQuip), standard coverslips and a dissection microscope. Larvae were

oriented such that their ventral side was facing up, their posterior spiracles were

protruding from their body and their left-right symmetry was protected. Once the slides

were ready, each coverslip was sealed with clear nail polish. The slides were then

incubated at 60°C overnight.

Imaging

The slides were imaged at 40X objective using a Zeiss AxioImager microscope and a

dark field filter. Using automated tiling, 64 high resolution images were taken for each

larva, which were later stitched using the ZEN 2012 (blue edition) software. Each image

was then exported to "tagged image format". These images will be made available prior

to publication.

Image processing

Measurements for the position of each abdominal denticle belt were made using a

custom Python script (the program will be made available prior to publication). This

program rotated and positioned each larva horizontally, anterior to the left and posterior

to the right, and cropped the image at the anterior, posterior and lateral borders. The

program then marked the anterior and posterior borders of the abdominal denticle belts.

It measured the distance from the anterior-most point of the larva to the anterior as well

as posterior borders of each denticle belt, and from the anterior-most point of the larva

to the posterior-most point of the larva (Fig. 1). Relative segment positions were

calculated as the distance of the anterior denticle boundary from the anterior of the

larva, divided by larval length (Fig. 1). The number of larval samples, from which

segment position measurements were taken and used for data analysis, varied from

105 to 145 for each species (Table S1).

Image editing

Some of the images were edited using ImageJ "Brush" tool to paint over bubbles around

the larvae and to adjust brightness and contrast when necessary. We found both of

these edits increased the number of successful runs by the custom image processing

program and did not change the measurements taken from these images.

Data analysis

The segment measurement data is available at

https://doi.org/10.6084/m9.figshare.8170787. A linear model ANOVA was fitted to the

data using R (R Core Team 2019) with the effects of species and segment and an

interaction term. "lm" and "aov" functions were used to implement the following

formulas. All terms were fixed.

Relative segment position. $= \mu +$ species $+$ segment $+$ species x segment $+ \varepsilon$

This was followed by Tukey's HSD (Honestly Significant Differences) (Steel et al. 1997)

test to conduct pairwise comparisons of the relative position of each segment between

species. Specifically, we used "HSD.test" function from the R package "agricolae" (De

Mendiburu 2019). We replicated these results using t-tests between each species pair

and then applying Bonferroni multiple test correction ("t.test" and "p.adjust" functions in

R, respectively) for both multiple species and multiple segments. These analyses were

done also for the posterior border and anterior-posterior width of denticle belts. For the

details of "posterior border" and "width" analyses of denticle belts, see File S1.

Principal component analysis was performed in R, using the "prcomp" function, and

mean centered positions of each segment as the data. To examine what PC1

represented in our data, we used Pearson correlation ("cor.test" function in R) to

correlate PC1 with the relative positions of each segment. Using the same method, we

also tested whether PC1 was correlated with larval length.

For correlation analysis between changes in relative segment position, we first

calculated deviation from the between-species mean of the relative position of each

segment in each species. We then used "cor" and "cor.test" functions in R to obtain

Pearson correlation coefficients and the associated p-values.

Phylogenetic Ancestral State Estimation

We performed phylogenetic analyses to infer the evolution of relative segment position

across the 12 *Drosophila* species here. For this, we first inferred an ultrametric

phylogeny published in Turelli et al. (2018) under three candidate relaxed molecular-

clock models. For each of the resulting phylogenies, we assessed the fit of four

candidate models that variously describe how rates of segment evolution vary across

branches of the phylogeny. Finally, we jointly inferred the phylogeny, model of segment

evolution, and the ancestral states for each segment under the preferred model of

segment evolution. We summarized various aspects of the evolution of relative segment

position from the resulting joint posterior probability distribution of ancestral states.

Complete details of these analyses are in File S2.

Results

Larval segment allometry is highly variable across *Drosophila* species

To elucidate whether relative segment position has changed over *Drosophila*

evolutionary history, we compared relative abdominal segment position (Fig. 1) at the

first-instar larval stage between 12 different species (Fig. 2A). We found that there are

many significant differences in relative segment position among species (Fig. 2B).

Overall, in 21 out of 66 species-pair comparisons, all eight segments had a significant

difference in their position. We note that species pairs do not represent independent

observations due to the underlying phylogeny (we perform a full phylogenetic analysis

below). For the majority of species-pair comparisons, the relative positions of five or

more segments differed (Fig. S2). In fact, there were only two pairs of species

compared (*D. simulans* - *D. ananassae* and *D. ananassae* - *D. willistoni*) where none of

the eight abdominal segments had a significant difference in their relative positions

(Table S2). Intriguingly, these species pairs are not closely related, with 15 and 32

million years of divergence, respectively. In all sister species comparisons, except for *D.*

yakuba – *D. santomea*, the relative position of the majority of abdominal segments were

different. On the other hand, between divergent species pairs, such as *D. melanogaster*

and *D. virilis*, relative positions of only two segments were different (Table S2).

Interestingly, when considering all species-pair comparisons together, some

species were responsible for a larger proportion of differences than others (Table S3).

Out of the 410 segment position differences observed over all species-pair

comparisons, differences with *D. persimilis* constituted 87 of these, the highest

proportion of any species (~21%). Differences with *D. sechellia*, *D. erecta* and *D.*

mojavensis followed with 78 each (~19%), whereas the number of differences with *D.*

ananassae was the lowest with 56 (~13.5%). This is consistent with the observation that

relative segment positions of *D. mojavensis* and *D. persimilis*, followed by *D. erecta* and

D. sechellia, have the highest total deviation from the species mean, whereas those of

D. ananassae are closest to the species mean making it the "average" species (Table

S3).

Next, we examined whether differences in segment position between species

were toward the head (anterior) or tail (posterior) of the larva. To do this, we calculated

mean positions of each segment across the 12 species and characterized whether a

particular segment in a particular species was more anterior or more posterior relative to

this mean (Fig. 3). For the majority of the species, the relative position of segments

physically closest to each other differed in the same direction. When the direction of the

difference changed, it was between segments that had the smallest (except for *D.*

willistoni) magnitude of difference in a given species (Fig. 3).

Notably, averaged over all species, deviation from the species mean in relative

segment position increased from anterior to posterior of the larva, with the posterior-

most segments having largest differences from the mean over all species. (Figs. 3,

S3A). Principal component analysis (Fig. S3B) also demonstrates this pattern, with PC1

correlating highly with the more posterior segment positions, explaining 88% of the

variance in the dataset. The trend of deviations in segment position increasing from

anterior to the posterior of the larva was strongest for *D. mojavensis* and *D. persimilis*,

(Figs. 3, S3C), but was nonetheless true in a number of other species as well (Fig. S3C,

Table S3). Indeed, when segment position data for *D. mojavensis* and *D. persimilis*

were excluded from the analysis, the trend still held (Fig. S3D).

In most species, magnitude of relative position change was highest for segment

A8 (Table S3), which extends from the anterior boundary of A8 to the tail of the larvae

(Fig. 1). We will refer to this segment as A8+tail. *D. mojavensis* and *D. persimilis* have

the largest differences for segment A8: *D. mojavensis* has the largest magnitude of

deviation from the species mean, with an anteriorly shifted segment A8 border and thus

a much longer A8+tail segment; while *D. persimilis* has a posteriorly shifted segment A8

border and thus an exceptionally short A8+tail segment. (Figs. 1, 2B and S5C). While

the magnitude of differences was highest for A8, the total number of differences are not

increased for this segment. Significant differences in relative position observed in

species-pair comparisons are highly similar in number for segments A3 through A8, with

segments A1 and A2 showing a slightly lower number of significant differences in

relative position (Fig. S4B, black line). Additionally, to determine if segment positioning

in the posterior of the embryo is less precise (noisier) as compared to the anterior, we

measured coefficient of variation for the position of each segment along the larva. The

coefficient of variation, averaged across all species, does not vary considerably and is

low across the length of the larva (~0.03) (Fig. S6A). This suggests that while segment

A8+tail has the largest magnitude differences in our dataset, its position is not more

variable between species, nor is it positioned any less precisely than the other

segments.

Changes in the size of A8+tail segment are responsible for a significant portion of the shifts in relative segment position between species

Given that the relative position of segment A8+tail, as compared to the other segments, had the largest deviations from the species mean for most of the 12 species (Fig. S3, Table S3), we asked how much of the total variation in relative segment position is driven by A8+tail. In order to address this, we recalculated relative segment position in the absence of A8+tail (see File S3). To determine whether any changes in segment position we see was due to the removal of A8+tail region specifically, and not simply due to the removal of a terminal segment, we made a separate recalculation of relative segment position after removing the head and thoracic region (h+t) (Fig. 1). In both cases we determined the total number of significant differences in relative segment position between pairs of species (t-test with Bonferroni correction), and corrected this number relative to the total number of comparisons. We found that the corrected total number of significant position differences was reduced from 51.25 to 38.42 (~25% decrease) when A8+tail was removed, but increased from 51.25 to 53.71 when h+t was removed (Fig. S4A, compare also Fig. 2B to Fig. S7A and B). This trend held even when segment position data for species with highest magnitude of posterior segment position differences, *D. mojavensis* and *D. persimilis*, were removed from calculations (from 31.63 down to 24, ~24% decrease, when A8+tail was removed, but up to 32.71 when h+t was removed). This suggests that changes in the size of A8+tail drove a substantial portion of the differences in relative segment position between species. On the other hand, changes in the size of h+t appear to have masked some of the inter-

species differences in relative position observed in the rest of the segments. These

results are consistent with the finding that the average difference in relative position of

A1, and hence the difference in the size of h+t, are the smallest in magnitude among all

segments (Figs. S3A, S5B), while relative position of A8, hence the size of A8+tail, has

the largest magnitude changes (Figs. S3A, S5C). They are also consistent with the

finding that total number of significant position changes are lower for segment A1 than

they are for A8 (Fig. S4B). For a more detailed analysis of "end removal" and its effects

as well as differences in the size of h+t and A8+tail, see File S3.

Changes in the length of larval body

As we had collected larval length measurements in order to calculate relative

segment positions, we also compared whole body length at the first-instar larval stage

among 12 *Drosophila* species (Fig. 2C), and tested whether length had an effect on

relative segment positioning. *D. santomea* was the largest at this stage, followed by *D.*

sechellia and *D. virilis*. *D. persimilis* and *D. pseudoobscura* were the smallest, followed

by *D. mojavensis*. Body length for the rest of the species showed few and smaller

differences (Fig. 2C). We had two comparisons between sister species with large size

differences, as *D. santomea* and *D. sechellia* were among the largest larvae, while their

sisters, *D. yakuba* and *D. simulans*, respectively, were of roughly average size. Sister

species *D. santomea* and *D. yakuba* had few differences in segment position, whereas

sister species *D. sechellia* and *D. simulans* were different for every segment, suggesting

that larval size is not predictive of segment position differences. Overall, larval length is

not correlated with number of segment position differences (Fig. S5A, R^2 = 0.0016, p-

value = 0.75). In other words, having a bigger difference in body length was not

correlated with having more differences in relative segment position between species.

Additionally, larval length had no effect on the direction (anterior vs. posterior) of

segment position differences, as we detected no relationship between body length and

the number of segments that are shifted to the anterior or posterior compared to the

species mean (Wilcoxon test, p-value = 1). Finally, we returned to our PCA (above), and

found that PC1, which was highly correlated with the more posterior segment positions,

was uncorrelated with larval length (Pearson correlation, p-value = 0.06).

Phylogenetic analysis shows variable rates of segment position evolution

To investigate how relative segment position has evolved during *Drosophila*

evolutionary history, we implemented phylogenetic methods to estimate the rates of

morphological evolution over the phylogeny. We employed a multivariate Brownian

motion model of evolution (Huelsenbeck and Rannala 2003; Lartillot and Poujol 2010),

and tested various morphological branch rate prior models that describe how rates of

morphological evolution vary across the branches of the tree. These analyses were

performed in RevBayes (v. 1.0.7; Höhna et al. 2016), and are outlined in detail in File

S2. Of the candidate models of segment evolution that we explored, the data

significantly preferred the uncorrelated lognormal relaxed molecular clock model, which

allowed rates to vary episodically along ancestor-descendant branches (*i.e.*, rates of

segment evolution are not correlated between ancestor-descendant branches). Results

presented here are for this model, but other morphological evolutionary models gave

similar results (File S2).

Our phylogenetic analyses indicate that evolutionary rates of relative segment

position are highly variable across branches of the *Drosophila* phylogeny (Figs. 4A,

S8A and B). Many of the highest rates of evolution are on branches for species with

sister species included in the analysis (i.e. *D. sechellia* and *D. simulans*; *D. santomea*

and *D. yakuba*; *D. persimilis* and *D. pseudoobscura*). This points to segment position

evolving quickly between these sibling species. The patterns vary among segments

(Fig. S9). Two lineages that are also evolving rapidly, particularly in the posterior half of

the larva (Figs. 4A, S9), those leading to *D. mojavensis* and *D. persimilis*, have the

longest and shortest posterior-most segment (A8+tail), respectively, in our dataset

(Figs. 2B, S5C). It is intriguing that the branch leading to *D. mojavensis* showed one of

the fastest rates of evolution, as it is also on a long branch. This high rate sustained

over a long branch is consistent with the A8+tail segment of *D. mojavensis* having the

largest magnitude change in our dataset (Figs. 2B, S3C).

Given that several species had large changes in segment position toward the

posterior of the larvae (Figs. 2B, 3, and S3C), we asked whether the overall rate of

evolution is also higher for some segments than it is for others. We examined the rate of

position evolution for each segment over all branches of the tree and found that rate of

evolution varies among segments, with posterior segments exhibiting elevated rates

(Fig. 4B). Segment A8 had the highest rate of change in segment position, with

segment A7 also having a slightly elevated rate than the rest of the segments (Fig. 4B).

The high rate of evolution for A8 is consistent with some of the largest magnitude

changes in our raw data (Figs. 2B, 3) being found for this segment.

To determine whether relative segment position has evolved through frequent

small changes or rare large changes, we examined the distribution of the amount of

change in relative segment position normalized across all branches of the *Drosophila*

evolutionary tree (i.e. amount of change per unit time; see File S2, S.3.4.5 and S.3.4.6

for details). This distribution of normalized magnitude of change in segment position

across all the branches of the phylogeny (Fig. 4C) showed that most segment positions

on most branches have experienced changes that are small in magnitude, whereas

some segment positions on some branches have experienced substantially larger

changes within a given amount of evolutionary time. This analysis shows that

throughout *Drosophila* evolutionary history, relative segment position has changed

predominantly through small quantitative steps, with occasional large leaps.

Is segment position along the anterior posterior axis controlled locally or

globally?

Within each species, we asked how the shift in the relative position of one

segment might be correlated to shifts in the relative position of other segments. This

would indicate whether there is a focal point along the anterior-posterior axis controlling

segment position or whether there is a more complex underlying regulatory mechanism.

We found that within all 12 species, the correlation between shifts in the relative

positions of any two segments is inversely proportional to the distance between the two

segments (Figs. 5A, S10, and S11). We did not observe a focal point governing the shift

in segments, where it might be expected that correlations would drop off as the distance

from that particular point increased. It appears that small adjustments in segment

position are made locally. These correlations are looking at the relationship between

segment positions across individuals within a species. As the variation within species

reflects developmental variation in a genetically identical line, these correlations show

that developmental variation in the positioning of one segment have stronger effects on

nearby segments than more distant segments. This is consistent with evidence from the

embryonic stage that showed correlation between deviation from mean in expression

boundary for various segmentation genes decreased with increasing distance between

the expression boundary (Lott et al. 2007). Interestingly, averaged over all species, shift

in the relative position of A8 was less well-correlated to shifts in the rest of the segments

(Figs. 5B, S12). This suggests that shifts in the relative position of A8 are more

independent than the shifts in the relative position of other segments within species

(See also File S4).

Given that the relative position of A8+tail changed more independently within a

species as compared to the other segments (Fig. 5B), we asked whether relative

position would be more tightly regulated across the other segments in the absence of

A8+tail or whether A8+tail was essential for proper segment positioning. To address this

question, we calculated correlation coefficients using relative segment positions

determined in the absence of A8+tail. These recalculated correlations resulted in lower

average correlation coefficients between position shifts of adjacent (as well as more

distant) segments as compared to when all segments were included in position

calculations. (Figs. S13, S14, red line, *D. sechellia* was exceptional, see Fig. S15). This

was also true when we recalculated correlations using relative segment positions

determined in the absence of h+t (Figs. S13, S14, blue line). These results suggest that

both ends of the larvae are needed for proper segment positioning, and removal of

either one decreases the level of coordination between segments, as demonstrated by

the reduction in correlation coefficients.

Coordination of segment position evolution across the larva

Within a species, segment position seems to be controlled locally in response to

developmental variation, as correlations between segments decrease with physical

distance in the larva. How then are evolutionary segment position changes correlated

across species? To address this question, we used the phylogenetic model to estimate

correlations in the rate of position change between segments across species. These

phylogenetic correlations may reflect underlying genetic correlations (in a quantitative

genetic sense) or be the result of constraints placed on the system by selection

pressures. Evolutionary rates of relative segment position were highly correlated across

the branches of the phylogeny, and correlation decreased as the anatomical distance

between segments increased (Fig. 5C). All correlations were positive, indicating that, for

example, when rapid evolution occurs in one segment, changes occur across all

segments. However, across species, the rate of evolution of segment A8 is not less

correlated than expected, given its distance from other segments. This suggests that

while this segment may evolve at a faster rate (Fig. 4B), on average it does not seem to

be evolving in a less coordinated fashion than the other segments, when the whole

phylogeny is considered.

Discussion

Many critical developmental processes and traits are known to be robust, in that

they are produced faithfully despite variation encountered in ontogeny. This tolerance of

variation does have limits, however, and exceeding these limits may disrupt the

development of these traits in such a way that development does not proceed. Hence,

robustness is desirable during development because it assures the precise production

of critical traits and processes within a range of developmental conditions.

Given the suppression of phenotypic variation in robust traits, their evolution has

been a subject of considerable interest to researchers over the years, and has produced

both theoretical and empirical work (Wagner 2005; Félix and Wagner 2006; Masel and

Siegal 2009; Siegal and Leu 2014; Payne and Wagner 2019). One proposal for the way

for a robust trait to evolve is for conditions to exceed the tolerance for variation in that

trait, and expose genetic variation that had previously been masked, i.e. cryptic genetic

variation, by the very robustness of the trait (Rutherford and Lindquist 1998; Gibson and

Dworkin 2004; Paaby and Rockman 2014). Exposing any amount of genetic variation of

unknown consequence to a critical trait at a time of great stress seems like a dangerous

proposition in a multicellular animal. Our data supports an alternative model, where

there is always some small amount of variation available, even in the most robust traits,

and that quantitative changes may evolve in robust traits without disrupting robustness.

Here, we focus on a well-studied robust process and trait, segmentation in

Drosophila. Previous studies have highlighted the ability of the segmentation network to

produce precisely localized segment markers in embryogenesis despite experimentally

produced perturbations in development (Félix and Wagner 2006; Masel and Siegal

2009). Our study extends this to measure the trait of segmentation in larvae, across 12

Drosophila species. Here, we demonstrate that segmentation varies considerably

across species, and that the rate of evolution of segment position varies considerably

across the phylogeny as well. The variation in segment position consists mostly of small

magnitude differences, suggesting that this trait evolves by many small changes, with

occasional larger changes. While it can be difficult to quantitatively compare earlier

stages of segmentation in the embryo to segments in the larva, this result is consistent

with patterning differences in the embryo observed between species (Lott et al. 2007;

2010). Our data shows that segment position differences between species exist also in

later developmental stages. Moreover, each line of each species measured here

precisely produces its characteristic segmentation pattern in the larva. This points to the

ability of segmentation to evolve without losing its ability to be precisely localized (Lott et

al. 2010). While we did not test the robustness of each species to genetic and

environmental variation specifically, this also suggests that segmentation can evolve

and remain robust to stochastic variation. If segmentation can evolve small quantitative

changes over evolutionary time without losing robustness, then this suggests that not all

genetic variation in this trait is cryptic (Miles et al. 2010; Jiang et al. 2015), and small-

scale differences (and the occasional larger-scale difference) in segmentation patterns

may be available to selection (Weber 1992). Or, perhaps there is a neutral space where

a range of segment positions are tolerated, and stabilizing selection keeps them in that

range.

The evolutionary patterns observed here may also be simply what might be

expected of complex traits generally, with changes in a number of genes producing

small quantitative changes in phenotype over evolutionary time. Here, we find that

segment position seems locally determined along the length of an individual, as position

shifts are highly correlated between adjacent segments, and correlations drop off over

distance, consistent with a previous result within species in embryos (Lott et al. 2007).

This is also consistent with the known regulatory interactions in this network, where

patterns are refined over developmental time (DiNardo and O'Farrell 1987; Surkova et

al. 2008) and regulation of segment patterning becomes more localized to a smaller

portion of the developing animal at each stage as compared to the previous one

(Nüsslein-Volhard and Wieschaus 1980; Kornberg and Tabata 1993; Nüsslein-Volhard

et al. 2008). Our results suggests that changes in many genes produced the observed

differences observed between species, and hence, it may be difficult to determine the

identity of individual genetic changes that may cause the differences in segment

position. However, the genetic network behind segment patterning has been fertile

ground for modeling in the embryo (Jaeger et al. 2004; Manu et al. 2009a; Wunderlich

and Depace 2011; Bentovim et al. 2017; Clark 2017; Verd et al. 2018; Petkova et al.

2019). Implementing a modeling framework developed for the embryonic stage, and

extending it past the embryonic segment polarity stage to the positions of segments in

the larva, is likely the best approach for identifying potential genetic causes of the

differences observed here.

Across the larva, the most striking pattern we observed was in the posterior-most

segment, from A8 to the tail of the animal. We found that this segment has the largest

magnitude of differences between species, and evolves faster between species than the

other segments. It is the most independently controlled segment within a species, with

correlations between its position and all other segments being the lowest found. As the

h+t region anterior of A1 does not share these properties, it is not simply a feature of

terminal segments, but specific to the A8+tail region. The A8+tail region contains the

posterior spiracles, important breathing structures in the larva (Hu and Castelli-Gair

1999), as well as the genital imaginal disc from which the genital structures in the adult

are produced (Sánchez and Guerrero 2001). While genital structures are known to

evolve rapidly between species (Eberhard 2013), it is unclear whether this would

produce differences in the entire segment in which the genital imaginal disc is found.

Alternatively, the differing conditions in which larvae find themselves may produce

differences in behavior that would require differences in posterior spiracle length. We

are currently exploring this possibility in relatives of *D. mojavensis*, as this species has

an exceptionally long A8+tail segment. In our care, *D. mojavensis* larvae burrow deeply

in food with only their long posterior spiracles visible above the surface of the food. As

D. mojavensis and its relatives are desert-dwelling cactus specialists (Oliveira et al.

2012), perhaps the larvae burrow deeply into cactus in their natural environment to find

a more hospitable microclimate (McKenzie and McKechnie 1979; Green et al. 1983).

Overall, our results show for the first time that segment position in *Drosophila* has

changed frequently throughout the evolutionary history of the genus. The changes were

mostly small in magnitude, presumably representing only small perturbations to the

development of the organism. Rate of evolution of this trait varies across the phylogeny,

with larger magnitude differences and a higher rate of evolution observed for the tail of

the larva. Changes in segment position within species, as well as rates of evolution

between species, are highly correlated between neighboring segments, indicating the

highly coordinated nature of the genetic network underlying this trait. Future studies are

needed to unravel the nature of the genetic changes underlying segment position

differences between species and whether any of the observed phenotypic differences

between species might be adaptive.

602
603   **References**
604

Arthur, W., and A. D. Chipman. 2005. The centipede Strigamia maritima: what it can tell
us about the development and evolution of segmentation. Bioessays 27:653–660.

Bejsovec, A. 2013. Wingless/Wnt signaling in *Drosophila*: The pattern and the pathway.
Mol. Reprod. Dev. 80:882–894.

Bentovim, L., T. T. Harden, and A. H. Depace. 2017. Transcriptional precision and
accuracy in development: from measurements to models and mechanisms.
Development 144:3855–3866.

Clark, A. G., M. B. Eisen, D. R. Smith, C. M. Bergman, B. Oliver, T. A. Markow, T. C.
Kaufman, M. Kellis, W. Gelbart, V. N. Iyer, D. A. Pollard, T. B. Sackton, A. M.
Larracuente, N. D. Singh, J. P. Abad, D. N. Abt, B. Adryan, M. Aguade, H. Akashi, W.
W. Anderson, C. F. Aquadro, D. H. Ardell, R. Arguello, C. G. Artieri, D. A. Barbash, D.
Barker, P. Barsanti, P. Batterham, S. Batzoglou, D. Begun, A. Bhutkar, E. Blanco, S. A.
Bosak, R. K. Bradley, A. D. Brand, M. R. Brent, A. N. Brooks, R. H. Brown, R. K. Butlin,

C. Caggese, B. R. Calvi, A. Bernardo de Carvalho, A. Caspi, S. Castrezana, S. E.
Celniker, J. L. Chang, C. Chapple, S. Chatterji, A. Chinwalla, A. Civetta, S. W. Clifton, J.
M. Comeron, J. C. Costello, J. A. Coyne, J. Daub, R. G. David, A. L. Delcher, K.
Delehaunty, C. B. Do, H. Ebling, K. Edwards, T. Eickbush, J. D. Evans, A. Filipski, S.
Findeiß, E. Freyhult, L. Fulton, R. Fulton, A. C. L. Garcia, A. Gardiner, D. A. Garfield, B.
E. Garvin, G. Gibson, D. Gilbert, S. Gnerre, J. Godfrey, R. Good, V. Gotea, B. Gravely,
A. J. Greenberg, S. Griffiths-Jones, S. Gross, R. Guigo, E. A. Gustafson, W. Haerty, M.
W. Hahn, D. L. Halligan, A. L. Halpern, G. M. Halter, M. V. Han, A. Heger, L. Hillier, A.
S. Hinrichs, I. Holmes, R. A. Hoskins, M. J. Hubisz, D. Hultmark, M. A. Huntley, D. B.
Jaffe, S. Jagadeeshan, W. R. Jeck, J. Johnson, C. D. Jones, W. C. Jordan, G. H.
Karpen, E. Kataoka, P. D. Keightley, P. Kheradpour, E. F. Kirkness, L. B. Koerich, K.
Kristiansen, D. Kudrna, R. J. Kulathinal, S. Kumar, R. Kwok, E. Lander, C. H. Langley,
R. Lapoint, B. P. Lazzaro, S.-J. Lee, L. Levesque, R. Li, C.-F. Lin, M. F. Lin, K.
Lindblad-Toh, A. Llopart, M. Long, L. Low, E. Lozovsky, J. Lu, M. Luo, C. A. Machado,
W. Makalowski, M. Marzo, M. Matsuda, L. Matzkin, B. McAllister, C. S. McBride, B.
McKernan, K. McKernan, M. Mendez-Lago, P. Minx, M. U. Mollenhauer, K. Montooth,
S. M. Mount, X. Mu, E. Myers, B. Negre, S. Newfeld, R. Nielsen, M. A. F. Noor, P.
O'Grady, L. Pachter, M. Papaceit, M. J. Parisi, M. Parisi, L. Parts, J. S. Pedersen, G.
Pesole, A. M. Phillippy, C. P. Ponting, M. Pop, D. Porcelli, J. R. Powell, S. Prohaska, K.
Pruitt, M. Puig, H. Quesneville, K. Ravi Ram, D. Rand, M. D. Rasmussen, L. K. Reed,
R. Reenan, A. Reily, K. A. Remington, T. T. Rieger, M. G. Ritchie, C. Robin, Y.-H.
Rogers, C. Rohde, J. Rozas, M. J. Rubenfield, A. Ruiz, S. Russo, S. L. Salzberg, A.
Sanchez-Gracia, D. J. Saranga, H. Sato, S. W. Schaeffer, M. C. Schatz, T. Schlenke, R.
Schwartz, C. Segarra, R. S. Singh, L. Sirot, M. Sirota, N. B. Sisneros, C. D. Smith, T. F.
Smith, J. Spieth, D. E. Stage, A. Stark, W. Stephan, R. L. Strausberg, S. Strempel, D.
Sturgill, G. Sutton, G. G. Sutton, W. Tao, S. Teichmann, Y. N. Tobari, Y. Tomimura, J.
M. Tsolas, V. L. S. Valente, E. Venter, J. Craig Venter, S. Vicario, F. G. Vieira, A. J.
Vilella, A. Villasante, B. Walenz, J. Wang, M. Wasserman, T. Watts, D. Wilson, R. K.
Wilson, R. A. Wing, M. F. Wolfner, A. Wong, G. Ka-Shu Wong, C.-I. Wu, G. Wu, D.
Yamamoto, H.-P. Yang, S.-P. Yang, J. A. Yorke, K. Yoshida, E. Zdobnov, P. Zhang, Y.
Zhang, A. V. Zimin, J. Baldwin, A. Abdouelleil, J. Abdulkadir, A. Abebe, B. Abera, J.
Abreu, S. Christophe Acer, L. Aftuck, A. Alexander, P. An, E. Anderson, S. Anderson,
H. Arachi, M. Azer, P. Bachantsang, A. Barry, T. Bayul, A. Berlin, D. Bessette, T.
Bloom, J. Blye, L. Boguslavskiy, C. Bonnet, B. Boukhgalter, I. Bourzgui, A. Brown, P.
Cahill, S. Channer, Y. Cheshatsang, L. Chuda, M. Citroen, A. Collymore, P. Cooke, M.
Costello, K. D'Aco, R. Daza, G. De Haan, S. DeGray, C. DeMaso, N. Dhargay, K.
Dooley, E. Dooley, M. Doricent, P. Dorje, K. Dorjee, A. Dupes, R. Elong, J. Falk, A.
Farina, S. Faro, D. Ferguson, S. Fisher, C. D. Foley, A. Franke, D. Friedrich, L.
Gadbois, G. Gearin, C. R. Gearin, G. Giannoukos, T. Goode, J. Graham, E. Grandbois,
S. Grewal, K. Gyaltsen, N. Hafez, B. Hagos, J. Hall, C. Henson, A. Hollinger, T. Honan,
M. D. Huard, L. Hughes, B. Hurhula, M. Erii Husby, A. Kamat, B. Kanga, S. Kashin, D.
Khazanovich, P. Kisner, K. Lance, M. Lara, W. Lee, N. Lennon, F. Letendre, R. LeVine,
A. Lipovsky, X. Liu, J. Liu, S. Liu, T. Lokyitsang, Y. Lokyitsang, R. Lubonja, A. Lui, P.
MacDonald, V. Magnisalis, K. Maru, C. Matthews, W. McCusker, S. McDonough, T.

Mehta, J. Meldrim, L. Meneus, O. Mihai, A. Mihalev, T. Mihova, R. Mittelman, V.
Mlenga, A. Montmayeur, L. Mulrain, A. Navidi, J. Naylor, T. Negash, T. Nguyen, N.
Nguyen, R. Nicol, C. Norbu, N. Norbu, N. Novod, B. O'Neill, S. Osman, E. Markiewicz,
O. L. Oyono, C. Patti, P. Phunkhang, F. Pierre, M. Priest, S. Raghuraman, F. Rege, R.
Reyes, C. Rise, P. Rogov, K. Ross, E. Ryan, S. Settipalli, T. Shea, N. Sherpa, L. Shi, D.
Shih, T. Sparrow, J. Spaulding, J. Stalker, N. Stange-Thomann, S. Stavropoulos, C.
Stone, C. Strader, S. Tesfaye, T. Thomson, Y. Thoulutsang, D. Thoulutsang, K.
Topham, I. Topping, T. Tsamla, H. Vassiliev, A. Vo, T. Wangchuk, T. Wangdi, M.
Weiand, J. Wilkinson, A. Wilson, S. Yadav, G. Young, Q. Yu, L. Zembek, D. Zhong, A.
Zimmer, Z. Zwirko, P. Alvarez, W. Brockman, J. Butler, C. Chin, M. Grabherr, M. Kleber,
E. Mauceli, and I. MacCallum. 2007. Evolution of genes and genomes on the
Drosophila phylogeny. Nature 450:203–218.

Clark, E. 2017. Dynamic patterning by the *Drosophila* pair-rule network reconciles long-
germ and short-germ segmentation. Plos Biol 15:e2002439.

Combs, P. A., and H. B. Fraser. 2018. Spatially varying cis-regulatory divergence in
Drosophila embryos elucidates cis-regulatory logic. PLoS Genet 14:e1007631–23.

Davis, G. K., and N. H. Patel. 1999. The origin and evolution of segmentation. Trends
Cell Biol. 9:M68–72.

Felipe De Mendiburu 2019. agricolae: Statistical Procedures for Agricultural Research.
R package version 1.3-1. https://CRAN.R-project.org/package=agricolae.

DiNardo, S., and P. H. O'Farrell. 1987. Establishment and refinement of segmental
pattern in the *Drosophila* embryo: spatial control of engrailed expression by pair-rule
genes. Genes & Development 1:1212–1225.

Driever, W., and C. Nüsslein-Volhard. 1988. The bicoid protein determines position in
the *Drosophila* embryo in a concentration-dependent manner. Cell 54:95–104.

Eberhard, W. G. 2013. Sexual Selection and Animal Genitalia. Harvard University
Press.

Félix, M.-A. 2012. *Caenorhabditis elegans* vulval cell fate patterning. Phys. Biol.
9:045001–20.

Félix, M.-A., and A. Wagner. 2006. Robustness and evolution: concepts, insights and
challenges from a developmental model system. Heredity 100:132–140.

Félix, M.-A., P. De Ley, R. J. Sommer, L. Frisse, S. A. Nadler, W. K. Thomas, J.
Vanfleteren, and P. W. Sternberg. 2000. Evolution of Vulva Development in the
Cephalobina (Nematoda). Developmental Biology 221:68–86.

Fowlkes, C. C., K. B. Eckenrode, M. D. Bragdon, M. Meyer, Z. Wunderlich, L.
Simirenko, C. L. Luengo Hendriks, S. V. E. Keränen, C. Henriquez, D. W. Knowles, M.
D. Biggin, M. B. Eisen, and A. H. Depace. 2011. A Conserved Developmental
Patterning Network Produces Quantitatively Different Output in Multiple Species of
Drosophila. PLoS Genet 7:e1002346.

Gibson, G., and I. Dworkin. 2004. Uncovering cryptic genetic variation. Nat Rev Genet
5:681–690.

Green, C. H., B. Burnet, K. J. Connolly, 1983. 1983. Organization and patterns of inter-
and intraspecific variation in the behaviour of *Drosophila* larvae. Animal Behaviour
31:282–291.

Gregor, T., D. W. Tank, E. F. Wieschaus, and W. Bialek. 2007. Probing the Limits to
Positional Information. Cell 130:153–164.

Houchmandzadeh, B., E. Wieschaus, and S. Leibler. 2002. Establishment of
developmental precision and proportions in the early *Drosophila* embryo. Nature
415:798–802.

Höhna, S., M. J. Landis, T. A. Heath, B. Boussau, N. Lartillot, B. R. Moore, J. P.
Huelsenbeck, and F. Ronquist. 2016. RevBayes: Bayesian Phylogenetic Inference
Using Graphical Models and an Interactive Model-Specification Language. Syst Biol
65:726–736.

Hu, N., and J. Castelli-Gair. 1999. Study of the posterior spiracles of *Drosophila* as a
model to understand the genetic and cellular mechanisms controlling morphogenesis.
Developmental Biology 214:197–210.

Huelsenbeck, J. P., and B. Rannala. 2003. Detecting correlation between characters in
a comparative analysis with uncertain phylogeny. Evolution 57:1237–1247.

Jaeger, J. 2010. The gap gene network. Cell. Mol. Life Sci. 68:243–274. Cellular and
molecular life sciences : CMLS.

Jaeger, J., M. Blagov, D. Kosman, K. N. Kozlov, Manu, E. Myasnikova, S. Surkova, C.
E. Vanario-Alonso, M. Samsonova, D. H. Sharp, and J. Reinitz. 2004. Dynamical
Analysis of Regulatory Interactions in the Gap Gene System of *Drosophila*
melanogaster. Genetics 167:1721–1737.

Jiang, P., M. Z. Ludwig, M. Kreitman, and J. Reinitz. 2015. Natural variation of the
expression pattern of the segmentation gene even-skipped in melanogaster.
Developmental Biology 405:173–181.

Kornberg, T. B., and T. Tabata. 1993. Segmentation of the *Drosophila* embryo. Current
Opinion in Genetics & Development 3:585–594.

Lartillot, N., and R. Poujol. 2010. A Phylogenetic Model for Investigating Correlated
Evolution of Substitution Rates and Continuous Phenotypic Characters. Molecular
Biology and Evolution 28:729–744.

Lohs-Schardin, M., C. Cremer, and C. Nüsslein-Volhard. 1979. A fate map for the larval
epidermis of *Drosophila melanogaster*: localized cuticle defects following irradiation of
the blastoderm with an ultraviolet laser microbeam. Developmental Biology 73:239–255.

Lott, S. E., M. Kreitman, A. Palsson, E. Alekseeva, and M. Z. Ludwig. 2007.
Canalization of segmentation and its evolution in *Drosophila*. Proceedings of the
National Academy of Sciences 104:10926–10931.

Lott, S. E., M. Z. Ludwig, and M. Kreitman. 2010. EVOLUTION AND INHERITANCE OF
EARLY EMBRYONIC PATTERNING IN *DROSOPHILA SIMULANS* AND *D.
SECHELLIA*. Evolution 65:1388–1399.

Lucchetta, E. M., J. H. Lee, L. A. Fu, N. H. Patel, and R. F. Ismagilov. 2005. Dynamics
of *Drosophila* embryonic patterning network perturbed in space and time using
microfluidics. Nature 434:1134–1138.

Mackay, T. F. C., S. Richards, E. A. Stone, A. Barbadilla, J. F. Ayroles, D. Zhu, S.
Casillas, Y. Han, M. M. Magwire, J. M. Cridland, M. F. Richardson, R. R. H. Anholt, M.
Barrón, C. Bess, K. P. Blankenburg, M. A. Carbone, D. Castellano, L. Chaboub, L.
Duncan, Z. Harris, M. Javaid, J. C. Jayaseelan, S. N. Jhangiani, K. W. Jordan, F. Lara,
F. Lawrence, S. L. Lee, P. Librado, R. S. Linheiro, R. F. Lyman, A. J. Mackey, M.
Munidasa, D. M. Muzny, L. Nazareth, I. Newsham, L. Perales, L.-L. Pu, C. Qu, M.
Ràmia, J. G. Reid, S. M. Rollmann, J. Rozas, N. Saada, L. Turlapati, K. C. Worley, Y.-
Q. Wu, A. Yamamoto, Y. Zhu, C. M. Bergman, K. R. Thornton, D. Mittelman, and R. A.
Gibbs. 2012. The *Drosophila melanogaster* Genetic Reference Panel. Nature 482:173–
178. Nature Publishing Group.

Manu, S. Surkova, A. V. Spirov, V. V. Gursky, H. Janssens, A.-R. Kim, O. Radulescu,
C. E. Vanario-Alonso, D. H. Sharp, M. Samsonova, and J. Reinitz. 2009a. Canalization
of gene expression and domain shifts in the *Drosophila* blastoderm by dynamical
attractors. PLoS Comp Biol 5:e1000303.

Manu, S. Surkova, A. V. Spirov, V. V. Gursky, H. Janssens, A.-R. Kim, O. Radulescu,
C. E. Vanario-Alonso, D. H. Sharp, M. Samsonova, and J. Reinitz. 2009b. Canalization
of gene expression in the *Drosophila* blastoderm by gap gene cross regulation. Plos
Biol 7:e1000049.

Markow, T. A., and P. O'Grady. 2005. *Drosophila*: a guide to species identification and
use. Academic Press.

Markow, T. A., S. BEALL, and L. M. MATZKIN. 2008. Egg size, embryonic development
time and ovoviviparity in *Drosophila* species. Journal of Evolutionary Biology 22:430–
434.

Masel, J., and M. L. Siegal. 2009. Robustness: mechanisms and consequences. Trends
in Genetics 25:395–403.

McKenzie, J. A., and S. W. McKechnie. 1979. A Comparative Study of Resource
Utilization in Natural Populations of *Drosophila melanogaster* and *D. simulans*.
Oecologia 40:299–309.

Miles, C. M., S. E. Lott, C. L. Luengo Hendriks, M. Z. Ludwig, Manu, C. L. Williams, and
M. Kreitman. 2010. ARTIFICIAL SELECTION ON EGG SIZE PERTURBS EARLY
PATTERN FORMATION IN *DROSOPHILA MELANOGASTER*. Evolution 65:33–42.

Nasiadka, A., B. H. Dietrich, and H. M. Krause. 2002. Anterior-posterior patterning in the
Drosophila embryo. Pp. 155–204 *in* Gene Expression at the Beginning of Animal
Development.

Nijhout, H. F., F. Sadre-Marandi, J. Best, and M. C. Reed. 2017. Systems Biology of
Phenotypic Robustness and Plasticity. Integrative and Comparative Biology 57:171–
184.

Nüsslein-Volhard, C., and E. Wieschaus. 1980. Mutations affecting segment number
and polarity in *Drosophila*. Nature 287:795–801.

Nüsslein-Volhard, C., H. G. Frohnhöfer, and R. Lehmann. 2008. Determination of
anteroposterior polarity in *Drosophila*. Science 238:1675–1681.

Obbard, D. J., J. Maclennan, K.-W. Kim, A. Rambaut, P. M. O'Grady, and F. M. Jiggins.
2012. Estimating Divergence Dates and Substitution Rates in the *Drosophila*
Phylogeny. Molecular Biology and Evolution 29:3459–3473.

Oliveira, D. C. S. G., F. C. Almeida, P. M. O'Grady, M. A. Armella, R. DeSalle, and W. J.
Etges. 2012. Monophyly, divergence times, and evolution of host plant use inferred from
a revised phylogeny of the *Drosophila repleta* species group. Molecular Phylogenetics
and Evolution 64:533–544.

Paaby, A. B., and M. V. Rockman. 2014. Cryptic genetic variation: evolution's hidden
substrate. Nat Rev Genet 15:247–258.

Payne, J. L., and A. Wagner. 2019. The causes of evolvability and their evolution. Nat
Rev Genet 20:24–38.

Petkova, M. D., G. Tkacik, W. Bialek, E. W. Cell, 2019. 2019. Optimal decoding of
cellular identities in a genetic network. Cell 176:844–855.e15.

Petkova, M. D., S. C. Little, F. Liu, and T. Gregor. 2014. Maternal Origins of
Developmental Reproducibility. Current Biology 24:1283–1288.

R Core Team (2019). R: A language and environment for statistical computing. R
Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Regier, J. C., J. W. Shultz, A. Zwick, A. Hussey, B. Ball, R. Wetzer, J. W. Martin, and C.
W. Cunningham. 2010. Arthropod relationships revealed by phylogenomic analysis of
nuclear protein-coding sequences. Nature 463:1079–1083.

Russo, C. A. M., B. Mello, A. Frazão, and C. M. Voloch. 2013. Phylogenetic analysis
and a time tree for a large drosophilid data set (Diptera: Drosophilidae). Zool J Linn Soc
169:765–775.

Russo, C. A., N. Takezaki, and M. Nei. 1995. Molecular phylogeny and divergence
times of drosophilid species. Molecular Biology and Evolution 12:391–404.

Rutherford, S. L., and S. Lindquist. 1998. Hsp90 as a capacitor for morphological
evolution. Nature 396:336–342.

Sánchez, L., and I. Guerrero. 2001. The development of the *Drosophila* genital disc.
Bioessays 23:698–707.

Siegal, M. L., and J.-Y. Leu. 2014. On the Nature and Evolutionary Impact of
Phenotypic Robustness Mechanisms. Annu. Rev. Ecol. Evol. Syst. 45:495–517.

Spieth, H. T. 1981. *Drosophila heteroneura* and *Drosophila silvestris*: Head shapes,
behavior and evolution. Evolution 35:921–930.

St Johnston, D., and C. Nüsslein-Volhard. 1992. The origin of pattern and polarity in the
Drosophila embryo. Cell 68:201–219.

Steel, R. G. D., J. H. Torrie, and D. D. A. 1997. Principles and Procedures of Statistics.
McGraw-Hill College.

Surkova, S., D. Kosman, K. Kozlov, Manu, E. Myasnikova, A. A. Samsonova, A. Spirov,
C. E. Vanario-Alonso, M. Samsonova, and J. Reinitz. 2008. Characterization of the
Drosophila segment determination morphome. Developmental Biology 313:844–862.

Surkova, S., E. Golubkova, L. Mamon, and M. Samsonova. 2018. Dynamic maternal
gradients and morphogenetic networks in *Drosophila* early embryo. Biosystems
173:207–213.

Tautz, D. 2004. Segmentation. Developmental Cell 7:301–312.

Turelli, M., B. S. Cooper, K. M. Richardson, P. S. Ginsberg, B. Peckenpaugh, C. X.
Antelope, K. J. Kim, M. R. May, A. Abrieux, D. A. Wilson, M. J. Bronski, B. R. Moore, J.-

J. Gao, M. B. Eisen, J. C. Chiu, W. R. Conner, and A. A. Hoffmann. 2018. Rapid Global
Spread of wRi-like Wolbachia across Multiple *Drosophila*. Current Biology 28:963–
971.e8.

Verd, B., E. Clark, K. R. Wotton, H. Janssens, E. Jiménez-Guri, A. Crombach, and J.
Jaeger. 2018. A damped oscillator imposes temporal order on posterior gap gene
expression in *Drosophila*. Plos Biol 16:e2003174–24.

Wagner, A. 2005. Robustness and Evolvability in Living Systems. Princeton University
Press.

Weber, K. E. 1992. How small are the smallest selectable domains of form? Genetics
130:345–353.

Wieschaus, E., and C. Nuesslein-Volhard. 2016. The Heidelberg Screen for Pattern
Mutants of *Drosophila*: A Personal Account. Annu. Rev. Cell Dev. Biol. 32:1–46.

Wunderlich, Z., and A. H. Depace. 2011. Modeling transcriptional networks in
Drosophila development at multiple scales. Current Opinion in Genetics & Development
21:711–718.

Wunderlich, Z., C. C. Fowlkes, K. B. Eckenrode, M. D. J. Bragdon, A. Abiri, and A. H.
Depace. 2019. Quantitative Comparison of the Anterior-Posterior Patterning System in
the Embryos of Five *Drosophila* Species. G3: Genesl Genomesl Genetics 9:2171–2182.

851
852
Figure 1. Example larval images and description of measurements. Example dark field
images of first instar larvae are shown for three species. *D. ananassae* is presented as
it is closest to the mean of all species for segment positions, while *D. persimilis* and *D.*
mojavensis are exceptional, especially in the length of their most posterior segments.
Red lines mark the anterior border and white lines mark the posterior border of denticle
belts, which are rows of bristles on the ventral surface of the animal and are recognized
by our image processing program. The anterior border of the denticle belt was used as
a proxy for segment border. To measure the position of an abdominal segment (e.g. A1)
relative to the body length (Y), the distance from the anterior border of the larvae to the
anterior border of that segment (X) is divided by the total body size (Y). Region
encompassing Head, T1, T2, T3 is referred to as "h+t" throughout the main text. T =
Thoracic Segment, A = Abdominal Segment.
865
Figure 2. Relative segment position and body size is highly variable among 12 species
of *Drosophila*. (A) 12 *Drosophila* species on an evolutionary tree. Adult male species
photos from Nicolas Gompel were downloaded from FlyBase. (B) Each black bar
represents the mean position in percent larval length for each segment in each species.
The color spread on the two sides of each black bar is 95% confidence interval. Each
segment is represented by a different color. x-axis is relative segment position in

percent larval length. A = Abdominal Segment. (C) Each black bar represents the
average first instar larval body length for each species. The gray shaded area around
the black bars is 95% confidence interval. x-axis is average body length in micrometers.
875
Figure 3. Coordinated direction of relative segment position changes in 12 *Drosophila*
species. Neighboring segments tend to shift together in a particular direction, toward the
anterior or posterior of the larva rather than shifting in opposite directions. Yellow
indicates anterior, blue indicates posterior shift in relative position as compared to the
mean position across all species for a given segment. The area of the circle is
proportional to the size of the shift in relative segment position.
882
Figure 4. Phylogenetic analysis of segment evolution in the 12 *Drosophila* species. (A)
Phylogeny inferred from nuclear loci with relative divergence times. Branches of the tree
are colored to indicate the overall rate of relative segment position evolution. Rates are
variable across the phylogeny, with some big differences in rate observed even
between closely related species. (B) Boxplots indicate the posterior distribution of the
evolutionary rate of relative segment position. The bar indicates the posterior mean rate;
the boxes and whiskers indicate the 50% and 95% posterior credible intervals,
respectively. The rate is higher for the most posterior segments, especially A7 and A8
(C) Marginal posterior probability distribution of the evolutionary rates of relative
segment position across branches. Most changes are small in magnitude, with
occasional larger changes.
894
Figure 5. Correlations between segment positions, within and between species. (A)
Within a species (here, *D. melanogaster* is shown as an example), correlations between
relative segment positions are highest for neighboring segments, and fall off as the
physical distance between two segments increases. Colors are used to distinguish
between segment pairs with different number of segments separating them. This
suggests local control of segment positioning. For plots for the rest of the species see
Fig. S10. (B) This plot is similar to that in part A, but instead shows averages of
correlation coefficients between segment positions over all species. Similarly,
correlations between relative segment positions are highest between neighboring
segments, but fall as distance between segments increase. In red are all comparisons
with segment A8. At any distance, correlations with this segment are lower (in black are
all other comparisons not involving A8). The gray and red shaded areas around the
black and red points, respectively, indicate their 95% confidence intervals. This
demonstrates that changes in the position of A8 are less correlated, and thus are more
independent, than changes in the position of the rest of the segments, given its distance
from any other segment. The unit of the x-axes in panels A and B, for mean relative
distance between segments, is percent larval length. (C) Between species, correlations
between evolutionary rates for a pair of segments are lower with increasing distance
between segment pairs. This plot represents a similar analysis to part B, but in the
phylogenetic framework and correlates rates of evolution for each segment. Color
indicates the first segment in the comparison, with the most-blue indicating comparisons

with segment 1, and the most yellow being comparisons with segment 7. Rates of
evolution are highly correlated between neighboring segments, and correlations
decrease with increasing distance between segments.
919
Figure S1. Testing the effect of larval mounting procedures on relative segment
position. For this experiment, relative segment position measurements were taken
before and after multiple stages of the mounting protocol, to test the effect of the
mounting protocol on segment position across multiple species (see Methods) (A) Bright
field image of a *D. melanogaster* first instar larva, hatched in water, iced for 2-3 minutes
before imaging. (B) The same *D. melanogaster* larva heated at 60°C for an hour prior to
bright field imaging. (C) Dark field image of the same *D. melanogaster* larva, mounted
using PVA, and incubated in the 60°C oven overnight. This represents the end stage of
our mounting protocol. White horizontal lines mark the full length of the larvae, vertical
lines mark the segment borders.
930
931 **Figure S2.** Comparing pairs of species, most segments were in different relative
932 positions. For the majority of species pair comparisons, five or more segments had
933 significantly shifted relative positions (t-tests with Bonferroni correction, p-value ≤
934 0.05). The graph shows the number of species pairs on the y-axis and number of
935 segments significantly shifted (see Methods) on the x-axis.
936
Figure S3. The posterior-most segments show the largest magnitude in differences
from mean across species. (A) Average absolute magnitude of shift in the relative
position of each segment, as compared to the across species mean, increases from
anterior to the posterior of the larvae. On the y-axis of this graph is absolute magnitude
of shift, in percent larval length, for each segment averaged over all species, on the x-
axis are abdominal segments A1 through A8. (B) From principal component analysis,
PC1 is plotted for all species. Positive values correspond to posterior shifts of
segments, negative values indicate anterior shifts of segments, relative to mean
centered positions for each segment. PC1 explains 88% of the variance and is highly
positively correlated with the positions of the most posterior segments (Pearson
correlation coefficients with each segment, A1-A8: A1= -0.15, A2=0.11, A3=0.28,
A4=0.49, A5=0.64, A6=0.77, A7=0.85, A8=0.88). Note the similarity in pattern to Figure
3, with *D. mojavensis* having the largest deviation from the rest of the species, with a
dramatic shift toward the anterior (negative PC1 values) and *D. persimilis* shifted toward
the posterior (positive PC1 values). (C) Absolute magnitude of shift changes from
anterior to posterior differently for each of the 12 *Drosophila* species. The graph shows
absolute magnitude of shift in percent larval length on the y-axis and abdominal
segments A1 through A8 on the x-axis. The legend in the graph shows the slopes
calculated for each trend form anterior to posterior. Lines in the graph and the legend
are color coordinated for each species. (D) Average absolute magnitude of shift in the
relative position of each segment, as compared to the across species mean, increases
from anterior to the posterior of the larvae even in the absence of segment position data
for *D. persimilis* and *D. mojavensis* (compare to Figure S3A, this representation is the

same, except with *D. persimilis* and *D. mojavensis* removed from the calculation of the
mean). On the y-axis of this graph is absolute magnitude of shift, in percent larval
length, for each segment averaged over all species, on the x-axis are abdominal
segments A1 through A8.
964
Figure S4. The A8+tail region is responsible for much of the number and magnitude of
significant differences between species. For all these plots, compare the effect of when
A8+tail is removed to when another terminal segment (h+t) is removed, to determine
whether the effect is A8+tail specific or simply an effect of removing a terminal segment.
(A) Comparison of numbers of significant position changes when segment positions
were calculated with all segments included, with h+t removed or with A8+tail removed.
Removal of A8+tail shows a decrease, whereas removal of h+t shows an increase, in
the number of segments that have significantly changed their relative position as
compared to when all eight abdominal segments were included in the position
calculations. Y-axis is total number of significant changes, normalized to the number of
segments included in position calculations. (B) This plot shows the same data as part A,
but illustrates how the differences in number change across the length of the larva.
When A8+tail was removed from position calculations, number of significant changes in
relative segment position went down sharply towards the posterior end of the larvae.
However, when h+t was removed in the same manner, number of significant changes in
relative segment position followed the same pattern from anterior to posterior as when
all abdominal segments were included in position calculations. y-axis is number of
significant changes in relative position normalized to the number of segments included
in position calculations. (C) Average absolute magnitude of shift (from the species
mean) for each segment went down sharply towards the posterior of the larvae when
A8+tail was removed from segment position calculations. On the other hand, removing
h+t followed the same pattern as when all segments were included in calculations, but
showed a bigger deviation from the species mean towards the posterior of the larvae. y-
axis is the average magnitude of shift (from the species mean) in segment position, in
percent larval length, ignoring the direction. X-axis is abdominal segments in both
panels B and C.
991
Figure S5. Features of body length. (A) Average body length difference between a
given pair of species is not correlated with the number of segments with significant
position shifts between the species. The y-axis in this graph shows average difference
in body length (in pixels) between each pair of species and the x-axis shows the number
of segments with significant position shifts in the same species pair. (B) This graph
shows differences in mean relative size for h+t, in percent larval length, among 12
Drosophila species. The black bars indicate the mean and the gray shaded area is 95%
confidence interval for each mean. (C) This graph shows differences in mean relative
size for A8+tail, in percent larval length, among 12 *Drosophila* species. The black bars
indicate the mean and the gray shaded area is 95% confidence interval for each mean.
Parts B and C represent transformations of the position and length measurements, to
highlight the length of the terminal portions of the larva.

1004
Figure S6. Coefficient of variation in segment positioning stays low throughout the
larva, indicating that the relative positions of all segments are equally as precisely
determined. However, standard deviation of segment positioning increases towards the
posterior end of the larva, as the measurement values increase (relative position is
measured in % of larval length from the anterior end). (A) Change in coefficient of
variation, averaged over 12 *Drosophila* species, for all segments from anterior to
posterior of the larva. The y-axis shows coefficient of variation, and the x-axis depicts
each abdominal segment. When segment positions were calculated with all segments
included (black dots), coefficient of variation (noise) did not change along the anterior-
posterior axis of the larva, and remained low. Removal of A8+tail (red dots) overall
decreased noise levels in segment positioning, whereas removal of h+t (blue dots)
increased noise levels towards the posterior of the larva. (B) Change in standard
deviation, averaged over 12 species, for all segments from anterior to posterior of the
larva. The y-axis shows standard deviation and the x-axis depicts each abdominal
segment. When segment positions were calculated with all segments included (black
dots), standard deviation of segment positions increased towards the posterior of the
larva. Removal of A8+tail (red dots) decreased standard deviations in the posterior,
whereas removal of h+t (blue dots) increased standard deviations towards the posterior
of the larva.
1024
Figure S7. Relative segment position and body size is highly variable among 12
species of *Drosophila,* though less so when A8+ tail is removed. These graphs have the
same format as Figure 2B and shows differences in relative segment position between
species. (A) Relative segment positions are calculated in the absence of A8+tail. Notice
a substantial number of nonsignificant differences as compared to Figure 2B. (B)
Relative segment positions are calculated in the absence of h+t and also with
measurements taken from the posterior of the larva instead of the anterior. As
compared to Figure 2B the segment order is flipped, but the pattern of change in
segment position is very similar. In both graphs, each black bar depicts the mean for
each segment in each species. The colored areas are 95% confidence intervals. The y-
axis depicts 12 *Drosophila* species and x-axis shows mean relative abdominal segment
position in percent larval length.
1037
Figure S8. Phylogenetic analysis of relative segment evolution in the 12 *Drosophila*
species, with posterior rate estimates for all branches. A) Like Figure 4A, the phylogeny
was inferred from nuclear loci with relative divergence times, and branches of the tree
are colored to indicate the overall rate of relative segment position evolution. Here, the
branches are numbered to correspond with the values pictured in part B. B) Box plot
depicting rate of evolution estimate for each branch of the phylogeny, x-axis
corresponds to the branch numbers indicated on the tree in part A.
1045
Figure S9. Phylogenetic analysis of relative segment position for each segment. Like
Figure 4A, this figure shows a phylogeny inferred from nuclear loci with relative

divergence times, and the branches of the tree are colored to indicate the overall rate of
segment position evolution. Like Figure S8, the tree diagrams also include numbers for
each linage, and are accompanied by a box plot showing the estimates for rates of
segment evolution by each numbered lineage. Here, there are 8 plots, to show rate
estimates for each abdominal segment separately.

1053

1054   **Figure S10**. Correlations between relative segment positions within each species. This
1055   series of graphs are constructed the same as Figure 5A, which was for *D.*
1056   *melanogaste*r, for each of the 12 *Drosophila* species. The y-axis shows correlation
1057   coefficients and the x-axis shows mean relative distance between pairs of segments in
1058   percent larval length.

1059

Figure S11. Correlation coefficient heat maps for each of the 12 *Drosophila* species.
This shows the correlation coefficient between shifts in the relative position (as
compared to the across species mean) of each pair of segments within a species. Red
indicates high positive correlation and white indicates low positive correlation. Pearson
correlation coefficients are printed on each square box. Notice that for most species, the
color scheme is lighter at the right edge of the triangle as compared to the bottom edge
of the triangle. This is consistent with other figures showing shifts in the position of A8
has a lower correlation with shifts in the position of other segments, regardless of the
distance in between any pair of segments.

1069

Figure S12. Correlations between segment positions, comparing correlations including
segment A8 with all other pairwise comparisons the same number of segments apart.
This series of graphs represent the information from Figure 5B, but broken out by how
many segments separate the two segments being compared. Each graph has mean
correlation coefficient over all species on the y-axis, and mean relative distance in
percent larval length over all species on the x-axis. Segments that are paired with A8
are in red. Error bars are 95% confidence intervals.

1077

Figure S13. When all segments are included in position calculations, mean correlation
coefficients between neighboring segments are highest, ever so slightly, in the middle of
the larva. In comparison, when A8+tail is removed from position calculations, correlation
between segment position shifts at the posterior of the larvae goes down significantly.
This is true for the anterior part of the larva, when h+t are removed from position
calculations.

1084

1085   **Figure S14.** This series of graphs are a continuation of Figure S13, as the distance
1086   between pairs of segments increase, from two to six segments apart. Notice that
1087   removal of h+t overall decreases correlation coefficients more than removal of A8+tail.

1088

1089   **Figure S15.** This series of graphs show the data represented in Figure S13 separately
1090   for each of the 12 *Drosophila* species. They show, for each species, how correlation
1091   coefficients change from the anterior to the posterior of the larvae, when relative

1092 segment position was calculated with all segments included (black) versus no A8+tail
1093 (red) or no h+t (blue).
1094
1095 **Table S1.** Experimental methods used for number of flies and days needed in a bottle
1096 to control population density of each species. Also includes, for each species,
1097 approximate embryonic developmental duration at 20°C and number of minutes needed
1098 in the 60°C oven for larvae to become straight.
1099
1100 **Table S2.** Lists the number of segments that are differentially positioned between pairs
1101 of species, and the divergence time between each pair in millions of years (Russo et al.
1102 1995; Obbard et al. 2012; Russo et al. 2013).
1103
1104 **Table S3.** Lists the deviation of the position of each segment in each species from the
1105 "across-species" mean as well as total number of significant segment position changes
1106 for each species over all species-pair comparisons.
1107
1108 **File S1.** Describes the observed changes in the position of posterior border of denticle
1109 belts and denticle width between species.
1110
1111 **File S2.** Methods and results of the phylogenetic analysis.
1112
File S3. Description of the method for removal of A8+tail or h+t from segment position
calculations, as well as some additional results from that analysis. In particular, this file
highlights the effects of the new calculations on differences in segment position along
the larva as well as the variation in segment position within a species. It also gives more
detail on the differences in the size of A8+tail and h+t between species when all
segments are included in segment position calculations.
1119
1120 **File S4.** Describes correlation between changes in the position of adjacent segments
1121 along the anterior-posterior axis for each species, highlighting species-specific patterns.
1122

Figure 1

D. ananassae

D. persimilis

D. mojavensis

Head T1   T2   T3   A1 A2   A3   A4   A5   A6   A7   A8 + Tail

X

Y

Relative Segment Position A1 = X/Y

Figure 2

A

B

C

Dvir
Dmoj
Dwil
Dper
Dpse
Dana
Dere
Dsan
Dyak
Dsec
Dsim
Dmel

40 30 20 10 0
Divergence time
(million years)

A1   A2   A3   A4   A5   A6   A7   A8

30   40   50   60   70   80   90
Mean Segment Position (% Larval Length)

900   1000   1100   1200
Average Body Length (μm)

Figure 3
0.0047% larval length
4.7% larval length
anterior shift
posterior shift

Figure 4

Figure 5

A Correlation between shifts in segment position (*D. melanogaster*)

$r^2 = 0.96$

Correlation Score

Mean Relative Distance Between Segments

B Average correlation between shifts in segment position

Mean Correlation Score

Mean Relative Distance Between Segments

C Correlations between evolutionary rates for pairs of segments

Correlation Score

Distance (number of segments)

Suplementary Figure 1

Supplementary Figure 2

Supplementary Figure 4

Supplementary Figure 6

Supplementary Figure 7

Supplementary Figure 8

Supplementary Figure 9
Segment 1

Supplementary Figure 9
Segment 2

Supplementary Figure 9
Segment 3

Supplementary Figure 9
Segment 4

Supplementary Figure 9

Segment 5

Supplementary Figure 9

Segment 6

Supplementary Figure 9

Segment 7

Supplementary Figure 9
Segment 8