Evolution of Larval Segment Position across 12 *Drosophila* Species

Running title: Evolution of Segmentation across Drosophila

Gizem Kalay[1], Joel Atallah[1*], Noemie C. Sierra[2], Austin M. Tang[1], Amanda E. Crofton[1], Mohan K. Murugesan[1], Sherri Wykoff-Clary[1], Susan E. Lott[1]

[1]The Department of Evolution and Ecology, University of California, Davis, One Shields Avenue, Davis, CA 95616
[2]Earth and Planetary Sciences Department, University of California, Davis, One Shields Avenue, Davis, CA 95616

Corresponding authors: Email Gizem Kalay (gizemkly@gmail.com) or Susan E. Lott (selott@ucdavis.edu).

Author contributions: G.K. and S.E.L. designed the study. G.K., A.M.T., A.E.C., M.K.M. and S.W. collected the data. J. A. wrote the Python script that took measurements from samples in an automated manner. G.K and N.C.S. analyzed the data. G.K. and S.E.L. wrote the manuscript with input from J.A. and N.C.S.

Acknowledgements: We thank Lott Lab members Emily L. Cartwright, Anna A. Feitzinger, Charles S. Omura for comments on the manuscript; and Graham Coop for comments and statistical help. We also thank The National Drosophila Species Stock Center for the Drosophila species used in this project. This work was financially supported in part by National Institutes of Health grants, R01GM111362, R00GM098448, and start-up funds provided to S.E.L. by the University of California, Davis.

Data Archiving: All data and R code used for phylogenetic analyses for segment position are deposited in FigShare Repository at: https://doi.org/10.6084/m9.figshare.8085206. Data tables for segment position measurements are deposited in FigShare Repository at: https://doi.org/10.6084/m9.figshare.8170787. Python script for taking larval segment measurements is available at J.A.'s github: https://github.com/joelatallah/larval_imaging.

Competing interests: The authors declare no competing interests.

---

[*] current address: The Department of Biological Sciences, University of New Orleans, 2000 Lakeshore Drive, New Orleans, LA 70148

Abstract

Many developmental traits that are critical to the survival of the organism are also robust. These robust traits are resistant to phenotypic change in the face of variation. This presents a challenge to evolution. In this paper, we asked whether and how a well-established robust trait, *Drosophila* segment patterning, changed over the evolutionary history of the genus. We compared segment position scaled to body length at the first instar larval stage among 12 *Drosophila* species. We found that relative segment position has changed many times between species. Changes were frequent, but primarily small in magnitude. Phylogenetic analysis demonstrated that rates of change in segment position are variable along the Drosophila phylogenetic tree, and that these changes can occur in short evolutionary timescales. Correlation between position shifts of segments decreased as the distance between two segments increased, suggesting local control of segment position. The posterior-most abdominal segment showed the highest magnitude of change on average, had the highest rate of evolution between species, and appeared to be evolving more independently as compared to the rest of the segments. This segment was exceptionally elongated in the cactophilic species in our dataset, raising questions as to whether this change may be adaptive.

Keywords: segment patterning, Drosophila, evolution, robustness, larval stage

Introduction

Many developmental phenotypes are critical to the survival and fitness of the organism. These phenotypes are often observed to be robust, in that they produce a stereotyped outcome despite variation encountered during development (Wagner 2005; Félix and Wagner 2006; Masel and Siegal 2009; Siegal and Leu 2014). The variation experienced in ontogeny can come in a variety of forms, including stochastic, genetic, and environmental variation (Wagner 2005; Félix and Wagner 2006; Masel and Siegal 2009; Siegal and Leu 2014). Some robust phenotypes have specific mechanisms in place to ensure they are produced faithfully (Félix and Wagner 2006; Masel and Siegal 2009; Siegal and Leu 2014; Nijhout et al. 2017). However, these same robust phenotypes do evolve over periods of evolutionary time (Félix et al. 2000; Arthur and Chipman 2005; Lott et al. 2007; 2010; Fowlkes et al. 2011; Félix 2012; Combs and Fraser 2018). This poses a fundamental question: how do robust traits evolve if the phenotypic variation necessary to selection is suppressed?

One such developmental phenotype is segmentation along the head to tail (anterior-posterior) axis. Segmentation is the periodic repetition of anatomical structures. It is a shared feature of three big animal phyla, annelids, arthropods, and chordates (Davis and Patel 1999; Tautz 2004). In Drosophila, the foundation of segmentation is laid during the beginning of embryogenesis (St Johnston and Nüsslein-Volhard 1992), when development is under control of the maternal gene products, before the zygotic genome is activated. Maternal gene products that are located in the anterior and posterior parts of the fertilized egg set up a concentration gradient along

the length of the embryo (St Johnston and Nüsslein-Volhard 1992; Surkova et al. 2018).

These maternal gene products regulate one another and regulate downstream targets

(St Johnston and Nüsslein-Volhard 1992; Surkova et al. 2018). This genetic network of

regulators, which has been well-established through decades of critical study (Nüsslein-

Volhard and Wieschaus 1980; Kornberg and Tabata 1993; Nasiadka et al. 2002;

Nüsslein-Volhard et al. 2008) precisely divide the embryo into progressively smaller

subsections over embryonic development, until the correct number of body segments is

reached. At the end of the embryonic stage, a first instar larva with a highly organized

segmented pattern of differentiated structures is produced.

Given the fundamental role of segmentation in Drosophila development, defects

in this process can be detrimental (or lethal) to the organism (Wieschaus and

Nuesslein-Volhard 2016). Indeed, the lethality of homozygous mutations in

segmentation genes was critical to their discovery in mutant screens (Wieschaus and

Nuesslein-Volhard 2016). However, studies have also shown that segmentation can

proceed correctly, while keeping its precision, in the face of substantial perturbations.

For instance, researchers showed that the stochastic variation between embryos in the

anterior-posterior gradient of the maternal gene *bicoid* (*bcd*), a master regulator for

embryonic patterning in Drosophila, is 2-5x higher than the variation observed in the

position of expression domain borders of downstream genes, particularly that of the gap

gene *hunchback* (*hb*) (Houchmandzadeh et al. 2002; Gregor et al. 2007; Manu et al.

2009b). The mechanism that cancels out the noise in Bcd gradient at the level of *hb*

expression is co-repression of *hb* by two other gap proteins, Kruppel and Knirps (Manu

et al. 2009b; Perry et al. 2012; Surkova et al. 2013). It has also been shown that in the

face of substantial genetic variation, such as mutations in important maternal genes or

absence of whole chromosomes, *hb* expression boundary was affected only

incrementally, if at all (Houchmandzadeh et al. 2002). Moreover, variation in the location

of *hb* expression boundary did not significantly increase (Houchmandzadeh et al. 2002).

Other studies have shown that changing Bcd dosage in the embryo resulted in

expansion or reduction of the head region as compared to the rest of the body (Driever

and Nüsslein-Volhard 1988; Namba et al. 1997). With some limitations, segment

patterning was corrected in these genetically perturbed embryos through increased or

decreased apoptosis in the enlarged versus reduced body regions, respectively (Namba

et al. 1997). We know that segmentation can also proceed precisely in the face of

substantial environmental perturbation. When whole (Houchmandzadeh et al. 2002) or

two halves (Lucchetta et al. 2005) of *D. melanogaster* embryos were raised at two

temperature extremes after fertilization, inducing different developmental rates

(Houchmandzadeh et al. 2002; Lucchetta et al. 2005), there were temperature induced

changes in Bcd gradient. However, by the gap or pair rule stage of embryonic

development, respectively, segments resumed their normal positions with the same

level of precision as control embryos (Houchmandzadeh et al. 2002; Lucchetta et al.

2005).

Drosophila segmentation is a complex trait that has been demonstrated to be

highly robust (Driever and Nüsslein-Volhard 1988; Namba et al. 1997;

Houchmandzadeh et al. 2002; Lucchetta et al. 2005; Manu et al. 2009b). It is perhaps

unsurprising then that with the exception of some Hawaiian Drosophila species (Spieth

1981), it is difficult to observe any gross differences in segment position, size and

shape, scaled to full body size, between adults of different *Drosophila* species.

However, over longer evolutionary time scales, segment patterning can vary

substantially between arthropod species (Regier et al. 2010). At the early embryonic

stage in Drosophila, previous studies have shown small quantitative differences in

relative segment position between both closely related (Lott et al. 2007; 2010; Fowlkes

et al. 2011; Combs and Fraser 2018) and more distantly related species (Fowlkes et al.

2011; Wunderlich et al. 2018). Additionally, small quantitative differences were found

within a species when *D. melanogaster* lines were subjected to strong artificial selection

for egg size (Miles et al. 2010) and also between naturally occurring strains of *D.*

melanogaster (Jiang et al. 2015). It is not known, however, whether these changes

persist beyond the embryonic stage, or if these trends appear beyond the limited

number of species examined. So while segment patterning is well documented to be

robust, it has also been demonstrated to have some small quantitative level of variation

within and between species. Evolution of this robust trait does occur, and the presence

of some variation within species suggests that evolution of small changes may be

possible without causing catastrophic failure of patterning. Thus, we hypothesized that

the evolution of segmentation may occur by relatively small shifts, rather than rare large

leaps, which we would need large sample sizes to detect.

In order to investigate the evolution of segment patterning systematically across

a range of divergence times, we characterized segment position across 12 species of

Drosophila. The twelve species of Drosophila in our experiment spanned the

evolutionary history of the genus (40-60 million years; Russo et al. 1995; Obbard et al.

2012; Russo et al. 2013) and included three pairs of sister species. We measured

position of each abdominal segment relative to body length in the 1st instar larvae.

Larval stage is the latest developmental stage where all segments are visible and easy

to measure, and the use of this stage facilitated the measuring of hundreds of larvae

from each species. We found that relative segment positions at the larval stage have

changed many times in the evolutionary history of Drosophila, and that these changes

are mostly small in magnitude, with some larger changes. Most species-pair

comparisons showed differences in the relative position of most abdominal segments.

Some sister species were significantly different at every segment, however some of the

most diverged species in our dataset showed no differences in their patterning. The

magnitude of differences in relative position increased towards the posterior segments

in most species, most strikingly in *D. persimilis* and *D. mojavensis*. Phylogenetic

modelling showed that rate of segment position change was highly variable among the

branches along the Drosophila phylogeny. Results of a correlation analysis for position

change between segments within species suggested local control of segment position,

as correlations decrease with physical distance. Correlations for changes in the rate of

evolution of segment position between branches of the phylogenetic tree also

decreased with physical distance between segments. Overall, these results

demonstrate that this complex and robust developmental trait does evolve, even over

short timescales, and it does so primarily by small frequent steps with occasional large

leaps. This may permit the evolution of novel patterns over long periods of evolutionary

time, without compromising the integrity of this critical developmental phenotype.

Materials and Methods

Species used

We took relative position measurements of abdominal segments from 12 *Drosophila*

species spanning the evolutionary history of the genus. These species were *D.*

melanogaster, *D. simulans*, *D. sechellia*, *D. yakuba*, *D. santomea*, *D. erecta*, *D.*

ananassae, *D. pseudoobscura*, *D. persimilis*, *D. willistoni*, *D. mojavensis wriglei* and *D.*

virilis (Figure 2A). We used the sequenced lines from 11 of these species (Clark et al.

2007). For *D. santomea* we used stock #14021-0271.01 (The National Drosophila

Species Stock Center).

Fly husbandry and population control

All flies were kept in plastic bottles in 20°C incubator with 60% humidity. Fly stocks were

raised on standard cornmeal food. The amount of time to sexual maturation varied

between species (Markow et al. 2008). Population control was conducted in a species-

specific manner through controlling the number of sexually mature adults added in a

given bottle, and how often the bottles were changed (Supplementary Table 1). As a

result, each bottle ended up with 40-50 pupae. Adult flies were discarded 14 days after

they became fertile, to control for parental age.

 Control experiment

In order to address whether the larval mounting procedure has species-specific effects

on measurements of relative segment position in first instar *Drosophila* larvae, we

examined relative segment position in *D. melanogaster*, *D. sechellia* and *D. virilis* at

three different stages of the larval prep procedure (Supplementary Figure 1). First instar

larvae that hatched in distilled water were temporarily immobilized using a 3-5 minute

incubation on ice, put on a microscope slide with water and then imaged using a Zeiss

AxioImager microscope in brightfield. These larvae were then put back in distilled water

incubated at 60°C for 50 minutes and imaged in water a second time using a brightfield

microscope. The larvae were then mounted in PVA Mounting Medium (BioQuip) (see

next section). The larvae were then imaged a third time using dark field microscopy (see

"Imaging" section). In the images of the iced (untreated control, the icing procedure

slows larvae enough for imaging) and then the heated larva, each of the 2nd and 3rd

thoracic, plus eight abdominal segments were marked by a node between bulges on the

larval cuticle (Supplementary Figure 1). In the dark field images, segment borders were

determined by the anterior border of each denticle belt. Denticle belts are rows of

bristles on the ventral side of the larvae that are used for traction while crawling

(Bejsovec 2013). The anterior border of each denticle belt is a proxy for the anterior

border of each abdominal segment (Lohs-Schardin et al. 1979). Using the Image J

(version 1.47t) "Line tool", measurements were taken from the anterior end of the larva

to the border of each segment. Segment position was then determined by dividing this

value by the full length of the larvae. Segment position data was analyzed using the

following linear models, implemented in R (R Core Team, 2017).

$Y = \mu + \text{prep} + \text{species} + \text{prep x species} + \varepsilon$.

$Y = \mu + \text{prep} + \text{species} + \text{segment} + \text{prep x species x segment} + \varepsilon$.

Preparing Drosophila 1st instar larvae for imaging

For each species, 20-50 newly emerged adults were obtained from a population-

controlled bottle and put in an egg collection bottle with a cap containing glucose-agar

food (100 grams of D-glucose, 15 gr of Agar, 1000 ml distilled H_2O with 1ml of Propionic

Acid added after autoclaving) that was sprinkled with Baker's yeast. For *D. sechellia,*

regular fly food was used in an egg lay cap with yeast sprinkled on it, as this prevents

this species from withholding egg laying, a particular problem with this species (Markow

et al. 2008). The bottle was then placed in 20°C incubator upside down. The next day

eggs were collected from egg lay caps, put onto a mesh and thoroughly washed using

distilled water to remove residual yeast and egg lay cap food. The eggs were then

placed in a petri dish filled with distilled water. This petri dish was placed in 20°C until

the eggs developed into first instar larvae. In addition to the variability in the number of

days necessary to reach sexual maturity, different species also varied in the number of

days necessary for a fertilized egg to develop into 1st instar larvae (Markow and

O'Grady 2005). The petri dish with water and larvae was then placed in 60°C oven for

~50 minutes (Supplementary Table 1). This incubation killed and straightened the

larvae. These larvae were then mounted on standard glass slides using PVA Mounting

Medium (BioQuip), standard coverslips and a dissection microscope. Larvae were

oriented such that their ventral side was facing up, their posterior spiracles were

protruding from their body and their left-right symmetry was protected. Once the slides

were ready, each coverslip was sealed with clear nail polish. The slides were then

incubated at 60°C oven overnight.

Imaging

The slides were imaged at 40X objective using a Zeiss AxioImager microscope and a

dark field filter. Using automated tiling, 64 high resolution images were taken for each

larva, which were later stitched using the ZEN 2012 (blue edition) software. Each image

was then exported to "tagged image format".

Image processing

Measurements for the position of each abdominal denticle belt were made using a

custom Python script (the program will be made available prior to publication). This

program rotated and positioned each larva horizontally, anterior to the left and posterior

to the right, and cropped the image at the anterior, posterior and lateral borders. The

program then marked the anterior and posterior borders of the abdominal denticle belts.

Afterwards, it measured the distance from the anterior-most point of the larva to the

anterior as well as posterior borders of each denticle belt, and from the anterior-most

point of the larva to the posterior-most point of the larva. Relative segment positions

were calculated as the distance of the anterior denticle boundary from the anterior of the

larva, divided by larval length. The number of larval samples, from which segment

position measurements were taken and used for data analysis, varied from 105 to 145

for each species (Supplementary Table 1).

Image editing

Some of the images were edited using ImageJ "Brush" tool to paint over bubbles around

the larvae and to adjust brightness and contrast when necessary. We found both of

these edits increased the number of successful runs by the custom image processing

program and did not change the measurements taken from these images.

Data analysis

The segment measurement data is deposited in FigShare Repository at

https://doi.org/10.6084/m9.figshare.8170787. A linear model ANOVA was fitted to the

data using R (R Core Team, 2017) with the effects of species and segment and an

interaction term. The formula follows:

Y = μ + species + segment + species x segment + ε

This was followed by Tukey's HSD (Honestly Significant Differences) (Steel et al. 1997)

test to conduct pairwise comparisons of the relative position of each segment between

species. Specifically, we used "HSD.test" function from the R package "agricolae". We

also replicated these results using t-tests between each species pair and then applying

Bonferroni multiple test correction for both multiple species and multiple segments. To

conduct these tests we used "t.test" and "p.adjust" functions in R.

For correlation analysis, we first calculated deviation from the between-species mean of

the relative position of each segment in each species. We then used "cor" and "cor.test"

functions in R to obtain Pearson correlation scores and the associated p-values.

Phylogenetic Ancestral State Estimation

We performed phylogenetic analyses to infer the evolution of relative segment position

in the 12 *Drosophila* species in our study. For this, we first inferred an ultrametric

phylogeny published in Turelli et al. (2018) under three candidate relaxed molecular-

clock models. For each of the resulting phylogenies, we assessed the fit of four

candidate models that variously describe how rates of segment evolution vary across

branches of the phylogeny. Finally, we jointly inferred the phylogeny, model of segment

evolution, and the ancestral states for each segment under the preferred model of

segment evolution. We summarized various aspects of the evolution of relative segment

position from the resulting joint posterior probability distribution of ancestral states. We

provide complete details of these analyses in the Supplementary File 1.

Results

The larval preparation procedure does not have species-specific effects on

segment position measurements

In order to test whether the larval preparation protocol had any species-specific

effects on relative segment position measurements, we conducted a preliminary

experiment using *D. melanogaster*, *D. sechellia* and *D. virilis*. These three species are

variable in their egg (Markow et al. 2008) and larval size (Figure 2C) and have varying

phylogenetic distances (Figure 2A). We therefore hypothesized that they might have a

higher chance of being differentially affected by the larval preparation protocol. We

imaged larvae from these species at three steps of the larval prep protocol: before

heating, after heating, and after mounting (See Methods). From these images we

manually took measurements of eight abdominal and two thoracic segments using

ImageJ (Supplementary Figure 1). Whether we included segment as a factor (p-value =

0.07 to 0.82) or not (p-value = 0.42 to 0.83) in our linear model formula, we did not find

any significant species-specific effects of the preparation protocol segment position

measurements.

The dataset: larval segment allometry is highly variable across Drosophila

species

To elucidate whether relative segment position has changed over *Drosophila*

evolutionary history, we compared relative abdominal segment position at the first instar

larval stage between 12 different species (Figure 2A). We found that there are many

significant differences in relative segment position among species (Figure 2B). Overall,

in 21 out of 66 species-pair comparisons, all eight segments had a significant difference

in their position. We note that species pairs do not represent independent observations

due to the underlying phylogeny (we perform a full phylogenetic analysis below). For the

majority of species-pair comparisons the relative position of five or more segments were

different (Supplementary Figure 2). In fact, there were only two pairs of species

compared (*D. simulans* - *D. ananassae* and *D. ananassae* - *D. willistoni*) where none of

the eight abdominal segments had a significant difference in their relative position

(Supplementary Table 2). Intriguingly, these species pairs are not closely related, with

15 and 32 million years of divergence, respectively. In all sister species comparisons in

our experiment, except for *D. yakuba* – *D. santomea*, the relative position of the majority

of abdominal segments were different. On the other hand, between divergent species

pairs, such as *D. melanogaster* and *D. virilis*, relative positions of only two segments

were different (Supplementary Table 2).

Next, we examined whether differences in segment position between species

were toward the head (anterior) or tail (posterior) of the larva. To do this, we calculated

mean positions of each segment across the 12 species in our study and characterized

whether a particular segment in a particular species were more anterior or more

posterior relative to this mean (Figure 3). We saw that, for the majority of the species,

the relative position of segments nearest each other differed in the same direction.

When the direction of the difference changed, it was between segments that had the

smallest (except for *D. willistoni*) magnitude of difference in a given species (Figure 3).

For seven out of 12 species, these differences were in one direction (anterior or

posterior) for all eight of their segments. In four species, *D. erecta*, *D. yakuba*, *D.*

santomea, *D. mojavensis,* the relative positions of all segments were more anterior,

whereas in *D. sechellia*, *D. ananassae* and *D. persimilis,* the relative positions of all

segments were more posterior as compared to the mean position for each segment

over 12 species. In *D. simulans* and *D. pseudoobscura*, there was only one segment

that was more anteriorly positioned, the rest of the segments were more posteriorly

positioned as compared to the mean position across the 12 *Drosophila* species in our

study.

Notably, averaged over all species, deviation from species mean in relative

segment position appeared to steadily increase toward the posterior of the larvae

(Supplementary Figure 3A). This trend was largely, but not exclusively, driven by the

segment position shifts in *D. mojavensis* and *D. persimilis* (Figure 3, Supplementary

Figure 3A, B)*,* which had the largest difference for segment A8 (which extends from the

anterior boundary of A8 to the tail of the larvae; we will refer to this segment as A8+tail)

among all the species. *D. mojavensis* has the largest magnitude of difference from the

mean over all species, with a much longer A8+tail segment, while *D. persimilis* has an

exceptionally short A8+tail segment. (Figures 1 and 2, Supplementary Figure 5C). We

also compared the rate of increase in magnitude of difference in relative segment

position along the length of the larvae between species. We saw that the steepest

increase was in *D. mojavensis*, followed by *D. persimilis*, *D. erecta* and *D.*

pseudoobscura (Figure 3, Supplementary Figure 3B). Interestingly, in all of the species

except for *D. sechellia*, *D. yakuba* and *D. willistoni*, magnitude of difference was highest

for the relative position of segment A8+tail. On the other hand, total number of

significant differences in relative position observed in species-pair comparisons did not

change much for segments A3 through A8, with segments A1 and A2 showing a slightly

lower number of significant differences in relative position (Supplementary Figure 4B,

black line).

Changes in the length of larval body, head, thorax and tail

We compared whole body length at the first instar larval stage among 12

Drosophila species (Figure 2C). *D. santomea* was the largest at this stage, followed by

D. sechellia and *D. virilis*. *D. perimilis* and *D. pseudoobscura* were the smallest,

followed by *D. mojavensis*. Body length for the rest of the species showed few and

smaller differences (Figure 2C). We had two comparisons between sister species with

large size differences, as *D. santomea* and *D. sechellia* were among the largest larvae,

while their sisters, *D. yakuba* and *D. simulans*, respectively, were of roughly average

size. Sister species *D. santomea* and *D. yakuba* had few differences in segment

position, whereas *D. sechellia* and *D. simulans* were different for every segment,

suggesting that larval size is not predictive of segment position differences. Overall,

there was no correlation between change in body length and the number of differences

in relative segment position between species (Supplementary Figure 5A). In other

words, having a bigger difference in body length was not correlated with having more

differences in relative segment position between species. Moreover, we detected no

difference between distributions of body lengths according to the number of more

anterior versus more posterior segments (Wilcoxon test, p-value=1). Thus, having a

shorter or a longer body was not correlated to having a bias towards higher number of

more anterior or more posterior segments.

We also compared the relative size of the head and thorax region (we will refer to

this region as h+t), i.e., head and three thoracic segments, normalized to the whole

body, among 12 Drosophila species and found only a few differences (Supplementary

Figure 5B). *D. virilis*, which is the third largest species at the first instar larval stage, had

the smallest h+t segments. The biggest h+t segments belonged to *D. persimilis*, the

smallest species and to *D. sechellia*, the second largest species at the first instar larval

stage. For the rest of the species, relative size of h+t seemed to fall in between, with

only few significant differences. Contrarily, we found many significant differences in the

relative size of A8+tail segment (Supplementary Figure 5C). Besides the largest and

smallest A8+tail segments observed in *D. mojavensis* and *D. persimilis*, respectively,

there are many differences of smaller magnitude between the rest of the species. We

also examined the width of our segment marker, the rows of bristles known as denticle

belts (see Methods) and re-performed our analysis for the posterior of the denticle belt

rather than the anterior. Briefly, the denticle belt marking segment A1 was the narrowest

(and most variable), with A8 being second narrowest, with the rest being of similar size,

with these patterns holding across species (Supplementary Figure 8A). Measuring

segment position from the posterior border of the denticle belt, we obtained very similar

results to those reported for the anterior border (Supplementary Figure 8B, C). For

details of these analyses see Supplementary File 2.

Changes in the size of A8+tail segment are responsible for a significant portion of

the shifts in relative segment position between species

Given that the relative position of segment A8+tail, as compared to the other

segments, had the largest deviation from the species experiment-average for most of

the species in our study (Supplementary Figures 3A, Supplementary Table 3) we asked

how much of the total variation in relative segment position is driven by A8+tail. In order

to address this, we recalculated relative segment position in the absence of A8+tail (see

Supplementary File 3). To determine that any changes in segment position we see was

due to the removal of A8+tail region specifically and not simply due to the removal of an

end segment, we made a separate recalculation of relative segment position after

removing the h+t region. In both cases we determined the total number of significant

differences in relative segment position between pairs of species and corrected it

according to the number of segments that were used in position calculations. We found

that the corrected total number of significant position differences was reduced from

51.25 to 38.42 (~25% decrease) when A8+tail was removed and increased from 51.25

to 53.71 when h+t was removed (Supplementary Figure 4A, compare also Figure 2B to

Supplementary Figure 6A and B). This suggests that changes in the size of A8+tail

drove a substantial portion of the differences in relative segment position between

species. On the other hand, changes in the size of h+t appear to have masked some of

the inter-species differences in relative position observed in the rest of the segments.

These results are consistent with the finding that the average difference in relative

position of A1 are the smallest in magnitude among all segments (Supplementary

Figure 3A), while A8 has the largest magnitude changes. For a more detailed analysis

of "end removal" and its effects see Supplementary File 3.

Is the A8+tail segment more variable within lines than the other segments?

As the posterior terminal segment (A8+tail) has the largest differences in

magnitude in our dataset and may be driving a significant proportion of the differences

we observe, we asked whether this segment was also the most variable within lines. To

determine if segment positioning in the anterior of the embryo is less noisy as compared

to the posterior, we measured coefficient of variation for the position of each segment

across species. When all segments are present in position calculations, then the

coefficient of variation averaged across all species does not vary considerably across

the length of the larva and stays low (~0.03) (Supplementary Figure 7A). This suggests

that while this segment has the largest magnitude differences in our dataset, it is not

positioned any less precisely.

Phylogenetic analysis shows variable rates of evolution

To investigate how segmentation has evolved over *Drosophila* evolutionary

history, we implemented phylogenetic methods to estimate the rates of morphological

evolution over the phylogeny. We employed a multivariate Brownian motion model of

evolution (Huelsenbeck and Rannala 2003; Lartillot and Poujol 2010), and tested

various morphological branch rate prior models that describe how rates of

morphological evolution vary across the branches of the tree. These analyses were

performed in RevBayes (v. 1.0.7; Höhna et al. 2016), and are outlined in detail in

Supplementary File 1. Of the candidate models of segment evolution that we explored,

the data significantly preferred the uncorrelated lognormal relaxed molecular clock

model, which allowed rates to vary episodically along ancestor-descendant branches

(*i.e.*, rates of segment evolution are not correlated between ancestor-descendant

branches). All the results in the main text are for this model, but other morphological

evolutionary models gave similar results (Supplementary File 1).

Our phylogenetic analyses indicate that evolutionary rates of relative segment

position are highly variable across branches of the *Drosophila* phylogeny (Figure 4A,

Supplementary Figure 9A, B). Many of the highest rates of evolution are on branches

for species with sister species included in the analysis (i.e. *D. sechellia* and *D. simulans*;

D. santomea and *D. yakuba*; *D. persimilis* and *D. pseudoobscura*). This points to

segment position evolving quickly between these sibling species. The patterns vary

among segments (Supplementary Figure 10). The rate of *D. sechellia* evolution

compared to *D. simulans* (and the rest of the species) is highest for the first four

abdominal segments (Supplementary Figure 10). The rate of evolution for *D. santomea*

is highest relative to *D. yakuba* in the center of the larvae, segments 4 and 5

(Supplementary Figure 10). Two lineages that are also evolving rapidly (Figure 4A),

those leading to *D. mojavensis* and *D. persimilis*, have the longest and shortest

posterior-most segment (A8+tail), respectively, in our dataset (Figure 2B,

Supplementary Figure 5C). The inclusion of the long branch leading to *D. mojavensis*

among branches with the fastest rates of evolution is of interest, because the other

lineages that we found to be evolving quickly are on shorter branches. This high rate

sustained over a long branch has resulted in the A8+tail segment of *D. mojavensis*

having the largest magnitude change in our dataset (Figure 2B, Supplementary Figure

3B). The rate of segment evolution in *D. mojavensis* and *D. persimilis* is highest in the

posterior half of the larvae (Supplementary Figure 10), giving rise to the large

magnitude of changes in the A8+tail segment (Figure 2B, Supplementary Figure 3B).

Given that several species had large changes in segment position toward the

posterior of the larvae (Figure 2B, Figure 3, Supplementary Figure 3B), we asked

whether the overall rate of evolution is also higher for some segments than it is for

others. We examined the rate of position evolution for each segment over all branches

of the tree. The evolutionary rate of relative segment position varies among segments,

with posterior segments exhibiting elevated rates (Figure 4B). Segment A8 had the

highest rate of change in segment position, with segment A7 also having a slightly

elevated rate (Figure 4B). The elevated rate of evolution for A8 is consistent with some

of the largest magnitude changes in our raw data (Figures 2B, 3) being found for this

segment.

To examine the magnitude of shifts in segment position, we examined the

distribution of the amount of change in relative segment position normalized across all

branches of the tree (i.e. amount of change per unit time; see Supplementary File 1,

S.3.4.5 and S.3.4.6 for details). The distribution of normalized magnitude of change in

segment position across all the branches of the phylogeny (Figure 4C) shows that most

segment positions on most branches have experienced small magnitude changes,

whereas some segment positions on some branches have experienced substantially

larger changes within a given amount of evolutionary time.

Is segment position along the anterior posterior axis controlled locally or globally?

Within each species, we asked how the shift in the relative position of one segment might be correlated to shifts in the relative position of other segments. This would show whether there is a focal point along the anterior-posterior axis controlling segment position or whether there is a more complex underlying regulatory mechanism. We found that within all of our species, the correlation between shifts in the relative positions of any two segments is inversely proportional to the distance between the two segments (Figure 5A, Supplementary Figures 11, 12). We did not observe a focal point governing the shift in segments, where it might be expected that correlations would drop off as the distance from that particular point increased. It appears that small adjustments in segment position are made locally. These correlations are looking at the relationship between segment positions across individuals within a species. As the variation within species reflects developmental variation in a genetically identical line, these correlations show that developmental variation in the positioning of one segment have stronger effects on nearby segments than more distant segments. This is consistent with evidence from the embryonic stage that showed correlation between deviation from mean in stripe boundary for various segmentation genes went down with increasing distance between their patterning domains of segmentation genes (Lott et al. 2007). Interestingly, averaged over all species, shift in the relative position of A8 was less well-correlated to shifts in the rest of the segments (Figure 5B). This relationship was strongest when adjacent segments were compared (Figure 5B), and less strong as

the distance between segments being compared increased (Supplementary Figure 13).

This suggests that shifts in the relative position of A8 are more independent than the

shifts in the relative position of other segments within species. For a report of correlation

scores between adjacent segments along the anterior-posterior axis for each species

see Supplementary File 4.

Next, we asked, given that the relative position of A8+tail changed more

independently within a species as compared to the other segments (Figure 5B), would

segment position be more tightly regulated in the absence of A8+tail or is A8+tail

essential for proper segment positioning? To address this question, we calculated

correlation scores using relative segment positions determined in the absence of

A8+tail. We saw that the absence of A8+tail resulted in lower correlation scores

between position shifts of adjacent segments as compared to when all segments were

present in position calculations. (Supplementary Figures 16 and 18, red line, *D.*

sechellia was exceptional, see Supplementary Figure 17). Similarly, when we removed

h+t from position calculations, averaged over all species, the correlation between

position shifts of adjacent segments was reduced as compared to when all segments

were present in position calculations. (Supplementary Figures 16 and 18, blue line).

These results suggest that both ends of the larvae are needed for proper segment

positioning, and removal of either one decreases the level of coordination between

segments.

Coordination of segment position evolution across the larva

Within a species, segment position seems to be controlled locally in response to

developmental variation, as correlations between segments decrease with physical

distance in the larva. How then are evolutionary segment position changes coordinated

across species? To address this question, we used the phylogenetic model to correlate

the rate of segment position change across species, across the larva. These

phylogenetic correlations may reflect underlying genetic correlations (in a quantitative

genetic sense) or be the result of constraints placed on the system by selection

pressures. Evolutionary rates of relative segment position are significantly correlated

across the branches of the phylogeny, and correlation decreases as the anatomical

distance between segments increases (Figure 5C). Additionally, all correlations are

positive, indicating that, for example, when rapid evolution occurs in one segment,

changes occur across all segments. However, across species, the rate of evolution of

segment A8 is not less correlated than expected, given its distance from other

segments. This is interesting to consider in light of the observation that within species

A8 is less correlated to the other segments than expected given its anatomical distance,

and perhaps surprising given that A8 also possesses a higher rate of evolution than the

other segments (Figure 4B). While this segment may evolve at a faster rate, on average

it does not seem to be evolving in a less coordinated fashion than the other segments,

when the whole phylogeny is considered.

Discussion

Many critical developmental processes and traits are known to be robust, in that

they are produced faithfully despite variation encountered in ontogeny. This tolerance of

variation does have limits, however, and exceeding these limits may disrupt the

development of these traits in such a way that development does not proceed. Hence,

robustness is desirable during development because it assures the precise production

of critical traits and processes within a range of developmental conditions.

Given the suppression of phenotypic variation in robust traits, their evolution has

been a subject of considerable interest to researchers over the years, and has produced

both theoretical and empirical work (Wagner 2005; Félix and Wagner 2006; Masel and

Siegal 2009; Siegal and Leu 2014; Payne and Wagner 2019). One proposal for the way

for a robust trait to evolve is for conditions to exceed the tolerance of variation for that

trait, and expose genetic variation that had previously been masked, i.e. cryptic genetic

variation, by the very robustness of the trait (Rutherford and Lindquist 1998; Gibson and

Dworkin 2004; Paaby and Rockman 2014). Unraveling any amount of genetic variation

of unknown consequence to a critical trait at a time of great stress seems like a

dangerous proposition in a multicellular animal.

Here, we focus on a well-studied robust process and trait, segmentation in

Drosophila. Previous studies have highlighted the ability of the segmentation network to

produce precisely localized segment markers in embryogenesis despite experimentally

produced perturbations in development (Félix and Wagner 2006; Masel and Siegal

2009). Our study extends this to measure the trait of segmentation in larvae, across 12

Drosophila species. Here, we demonstrate that segmentation varies considerably

across species, and that the rate of evolution of segment position varies considerably as

well. The variation in segment position consists mostly of small-magnitude differences,

suggesting that this trait evolves by many small steps, with some occasional larger-

scale changes. While it is difficult to compare sections in the embryo to segments in the

larva, this result is consistent with patterning differences observed between species in

the embryo (Lott et al. 2007; 2010). Our data shows that segment position differences

exist between species in later developmental stages as well. Additionally, we find that

each line of each species measured here precisely produces its characteristic

segmentation pattern in the larva. This points to the ability of segmentation to evolve

without losing its ability to be precisely localized (Lott et al. 2010). While we did not test

the robustness of each species to genetic and environmental variation specifically, this

suggests that segmentation can evolve and remain robust to stochastic variation. If

segmentation can evolve small quantitative changes over evolutionary time without

losing robustness, then this suggests that not all genetic variation in this trait is cryptic

(Miles et al. 2010; Jiang et al. 2015), and small-scale differences (and the occasional

larger-scale difference) in segmentation patterns may be available to selection (Weber

1992). Or, perhaps there is a neutral space where a range of segment positions are

tolerated, and stabilizing selection keeps them in that range.

The evolutionary patterns observed here may also be simply what might be

expected of complex traits generally, with changes in a number of genes producing

small quantitative changes in phenotype over evolutionary time. Here, we find that

segment position seems locally determined along the length of an individual, as position

shifts are highly correlated between local segments, and correlations drop off over

distance, consistent with a previous result within species in embryos (Lott et al. 2007).

This is also consistent with the known regulatory interactions in this network, where

patterns are refined over developmental time (Manu et al. 2009a) and regulation of

segment patterning becomes more local at each stage as compared to the previous one

(Nüsslein-Volhard and Wieschaus 1980; Kornberg and Tabata 1993; Nüsslein-Volhard

et al. 2008). This result suggests that a number of genes are involved in producing the

differences observed between species. This makes it difficult to determine the identity of

individual genetic changes that may cause the differences in segment position.

Across the larva, the most striking pattern we observed can be found in the

posterior-most segment, from A8 to the tail of the animal. We find that this segment has

the largest magnitude of differences between species. It has a higher rate of evolution

between species than the other segments. It is the most independently controlled

segment within a species, with correlations between its position and all other segments

being the lowest found. As the h+t region anterior of A1 does not share these properties

in our study, it is not simply a feature of terminal segments, but specific to the A8+tail

region. The A8+tail region contains the posterior spiracles, important breathing

structures in the larva (Hu and Castelli-Gair 1999), as well as the genital imaginal disc

from which the genital structures in the adult are produced (Sánchez and Guerrero

2001). While genital structures are known to evolve rapidly between species (Eberhard

2013), it is unclear whether this would produce differences in the entire segment in

which the genital imaginal disc is found. Alternatively, the differing conditions in which

larvae find themselves may produce differences in behavior that would require

differences in posterior spiracle length. We are currently exploring this possibility in

relatives of *D. mojavensis*, as this species has an exceptionally long A8+tail segment. In

our care, *D. mojavensis* larvae burrow deeply in food with only their long posterior

spiracles above the surface of the food. As *D. mojavensis* and its relatives are desert-

dwelling cactus specialists (Oliveira et al. 2012), perhaps the larvae burrow deeply into

cactus in their natural environment to find a more hospitable microclimate (McKenzie

and McKechnie 1979; Green et al. 1983).

Overall, our results show for the first time that segment position in Drosophila has

changed frequently throughout the evolutionary history of the genus. The changes were

mostly small in magnitude, presumably representing only small perturbations to the

development of the organism. Rate of evolution of this trait varies across the phylogeny,

with larger magnitude differences and a higher rate of evolution observed for the tail of

the larva. Changes in segment position within species, as well as rates of evolution

between species, are highly coordinated with changes in the closest segments,

indicating the highly coordinated nature of the genetic network underlying this trait.

Future studies are needed to unravel the nature of the genetic changes underlying

segment position differences between species and whether any of the observed

phenotypic differences between species might be adaptive.

References

648
649
Arthur, W., and A. D. Chipman. 2005. The centipede Strigamia maritima: what it can tell
us about the development and evolution of segmentation. Bioessays 27:653–660.

Bejsovec, A. 2013. Wingless/Wnt signaling in Drosophila: The pattern and the pathway.
Mol. Reprod. Dev. 80:882–894.

Clark, A. G., M. B. Eisen, D. R. Smith, C. M. Bergman, B. Oliver, T. A. Markow, T. C.
Kaufman, M. Kellis, W. Gelbart, V. N. Iyer, D. A. Pollard, T. B. Sackton, A. M.
Larracuente, N. D. Singh, J. P. Abad, D. N. Abt, B. Adryan, M. Aguade, H. Akashi, W.
W. Anderson, C. F. Aquadro, D. H. Ardell, R. Arguello, C. G. Artieri, D. A. Barbash, D.
Barker, P. Barsanti, P. Batterham, S. Batzoglou, D. Begun, A. Bhutkar, E. Blanco, S. A.
Bosak, R. K. Bradley, A. D. Brand, M. R. Brent, A. N. Brooks, R. H. Brown, R. K. Butlin,
C. Caggese, B. R. Calvi, A. Bernardo de Carvalho, A. Caspi, S. Castrezana, S. E.
Celniker, J. L. Chang, C. Chapple, S. Chatterji, A. Chinwalla, A. Civetta, S. W. Clifton, J.
M. Comeron, J. C. Costello, J. A. Coyne, J. Daub, R. G. David, A. L. Delcher, K.
Delehaunty, C. B. Do, H. Ebling, K. Edwards, T. Eickbush, J. D. Evans, A. Filipski, S.
Findeiß, E. Freyhult, L. Fulton, R. Fulton, A. C. L. Garcia, A. Gardiner, D. A. Garfield, B.
E. Garvin, G. Gibson, D. Gilbert, S. Gnerre, J. Godfrey, R. Good, V. Gotea, B. Gravely,
A. J. Greenberg, S. Griffiths-Jones, S. Gross, R. Guigo, E. A. Gustafson, W. Haerty, M.
W. Hahn, D. L. Halligan, A. L. Halpern, G. M. Halter, M. V. Han, A. Heger, L. Hillier, A.
S. Hinrichs, I. Holmes, R. A. Hoskins, M. J. Hubisz, D. Hultmark, M. A. Huntley, D. B.
Jaffe, S. Jagadeeshan, W. R. Jeck, J. Johnson, C. D. Jones, W. C. Jordan, G. H.
Karpen, E. Kataoka, P. D. Keightley, P. Kheradpour, E. F. Kirkness, L. B. Koerich, K.
Kristiansen, D. Kudrna, R. J. Kulathinal, S. Kumar, R. Kwok, E. Lander, C. H. Langley,
R. Lapoint, B. P. Lazzaro, S.-J. Lee, L. Levesque, R. Li, C.-F. Lin, M. F. Lin, K.
Lindblad-Toh, A. Llopart, M. Long, L. Low, E. Lozovsky, J. Lu, M. Luo, C. A. Machado,
W. Makalowski, M. Marzo, M. Matsuda, L. Matzkin, B. McAllister, C. S. McBride, B.
McKernan, K. McKernan, M. Mendez-Lago, P. Minx, M. U. Mollenhauer, K. Montooth,
S. M. Mount, X. Mu, E. Myers, B. Negre, S. Newfeld, R. Nielsen, M. A. F. Noor, P.
O'Grady, L. Pachter, M. Papaceit, M. J. Parisi, M. Parisi, L. Parts, J. S. Pedersen, G.
Pesole, A. M. Phillippy, C. P. Ponting, M. Pop, D. Porcelli, J. R. Powell, S. Prohaska, K.
Pruitt, M. Puig, H. Quesneville, K. Ravi Ram, D. Rand, M. D. Rasmussen, L. K. Reed,
R. Reenan, A. Reily, K. A. Remington, T. T. Rieger, M. G. Ritchie, C. Robin, Y.-H.
Rogers, C. Rohde, J. Rozas, M. J. Rubenfield, A. Ruiz, S. Russo, S. L. Salzberg, A.
Sanchez-Gracia, D. J. Saranga, H. Sato, S. W. Schaeffer, M. C. Schatz, T. Schlenke, R.
Schwartz, C. Segarra, R. S. Singh, L. Sirot, M. Sirota, N. B. Sisneros, C. D. Smith, T. F.
Smith, J. Spieth, D. E. Stage, A. Stark, W. Stephan, R. L. Strausberg, S. Strempel, D.
Sturgill, G. Sutton, G. G. Sutton, W. Tao, S. Teichmann, Y. N. Tobari, Y. Tomimura, J.
M. Tsolas, V. L. S. Valente, E. Venter, J. Craig Venter, S. Vicario, F. G. Vieira, A. J.
Vilella, A. Villasante, B. Walenz, J. Wang, M. Wasserman, T. Watts, D. Wilson, R. K.
Wilson, R. A. Wing, M. F. Wolfner, A. Wong, G. Ka-Shu Wong, C.-I. Wu, G. Wu, D.
Yamamoto, H.-P. Yang, S.-P. Yang, J. A. Yorke, K. Yoshida, E. Zdobnov, P. Zhang, Y.
Zhang, A. V. Zimin, J. Baldwin, A. Abdouelleil, J. Abdulkadir, A. Abebe, B. Abera, J.
Abreu, S. Christophe Acer, L. Aftuck, A. Alexander, P. An, E. Anderson, S. Anderson,
H. Arachi, M. Azer, P. Bachantsang, A. Barry, T. Bayul, A. Berlin, D. Bessette, T.
Bloom, J. Blye, L. Boguslavskiy, C. Bonnet, B. Boukhgalter, I. Bourzgui, A. Brown, P.
Cahill, S. Channer, Y. Cheshatsang, L. Chuda, M. Citroen, A. Collymore, P. Cooke, M.
Costello, K. D'Aco, R. Daza, G. De Haan, S. DeGray, C. DeMaso, N. Dhargay, K.

Dooley, E. Dooley, M. Doricent, P. Dorje, K. Dorjee, A. Dupes, R. Elong, J. Falk, A.
Farina, S. Faro, D. Ferguson, S. Fisher, C. D. Foley, A. Franke, D. Friedrich, L.
Gadbois, G. Gearin, C. R. Gearin, G. Giannoukos, T. Goode, J. Graham, E. Grandbois,
S. Grewal, K. Gyaltsen, N. Hafez, B. Hagos, J. Hall, C. Henson, A. Hollinger, T. Honan,
M. D. Huard, L. Hughes, B. Hurhula, M. Erii Husby, A. Kamat, B. Kanga, S. Kashin, D.
Khazanovich, P. Kisner, K. Lance, M. Lara, W. Lee, N. Lennon, F. Letendre, R. LeVine,
A. Lipovsky, X. Liu, J. Liu, S. Liu, T. Lokyitsang, Y. Lokyitsang, R. Lubonja, A. Lui, P.
MacDonald, V. Magnisalis, K. Maru, C. Matthews, W. McCusker, S. McDonough, T.
Mehta, J. Meldrim, L. Meneus, O. Mihai, A. Mihalev, T. Mihova, R. Mittelman, V.
Mlenga, A. Montmayeur, L. Mulrain, A. Navidi, J. Naylor, T. Negash, T. Nguyen, N.
Nguyen, R. Nicol, C. Norbu, N. Norbu, N. Novod, B. O'Neill, S. Osman, E. Markiewicz,
O. L. Oyono, C. Patti, P. Phunkhang, F. Pierre, M. Priest, S. Raghuraman, F. Rege, R.
Reyes, C. Rise, P. Rogov, K. Ross, E. Ryan, S. Settipalli, T. Shea, N. Sherpa, L. Shi, D.
Shih, T. Sparrow, J. Spaulding, J. Stalker, N. Stange-Thomann, S. Stavropoulos, C.
Stone, C. Strader, S. Tesfaye, T. Thomson, Y. Thoulutsang, D. Thoulutsang, K.
Topham, I. Topping, T. Tsamla, H. Vassiliev, A. Vo, T. Wangchuk, T. Wangdi, M.
Weiand, J. Wilkinson, A. Wilson, S. Yadav, G. Young, Q. Yu, L. Zembek, D. Zhong, A.
Zimmer, Z. Zwirko, P. Alvarez, W. Brockman, J. Butler, C. Chin, M. Grabherr, M. Kleber,
E. Mauceli, and I. MacCallum. 2007. Evolution of genes and genomes on the
Drosophila phylogeny. Nature 450:203–218.

Combs, P. A., and H. B. Fraser. 2018. Spatially varying cis-regulatory divergence in
Drosophila embryos elucidates cis-regulatory logic. PLoS Genet 14:e1007631–23.

Davis, G. K., and N. H. Patel. 1999. The origin and evolution of segmentation. Trends
Cell Biol. 9:M68–72.

Driever, W., and C. Nüsslein-Volhard. 1988. The bicoid protein determines position in
the Drosophila embryo in a concentration-dependent manner. Cell 54:95–104. Elsevier.

Eberhard, W. G. 2013. Sexual Selection and Animal Genitalia. Harvard University
Press.

Félix, M.-A. 2012. Caenorhabditis elegansvulval cell fate patterning. Phys. Biol.
9:045001–20.

Félix, M.-A., and A. Wagner. 2006. Robustness and evolution: concepts, insights and
challenges from a developmental model system. Heredity 100:132–140.

Félix, M.-A., P. De Ley, R. J. Sommer, L. Frisse, S. A. Nadler, W. K. Thomas, J.
Vanfleteren, and P. W. Sternberg. 2000. Evolution of Vulva Development in the
Cephalobina (Nematoda). Developmental Biology 221:68–86.

Fowlkes, C. C., K. B. Eckenrode, M. D. Bragdon, M. Meyer, Z. Wunderlich, L.
Simirenko, C. L. Luengo Hendriks, S. V. E. Keränen, C. Henriquez, D. W. Knowles, M.
D. Biggin, M. B. Eisen, and A. H. Depace. 2011. A Conserved Developmental

Patterning Network Produces Quantitatively Different Output in Multiple Species of
Drosophila. PLoS Genet 7:e1002346.

Gibson, G., and I. Dworkin. 2004. Uncovering cryptic genetic variation. Nat Rev Genet
5:681–690.

Green, C. H., B. Burnet, K. J. Connolly, 1983. 1983. Organization and patterns of inter-
and intraspecific variation in the behaviour of Drosophila larvae. Animal Behaviour
31:282–291.

Gregor, T., D. W. Tank, E. F. Wieschaus, and W. Bialek. 2007. Probing the Limits to
Positional Information. Cell 130:153–164.

Houchmandzadeh, B., E. Wieschaus, and S. Leibler. 2002. Establishment of
developmental precision and proportions in the early Drosophila embryo. Nature
415:798–802.

Höhna, S., M. J. Landis, T. A. Heath, B. Boussau, N. Lartillot, B. R. Moore, J. P.
Huelsenbeck, and F. Ronquist. 2016. RevBayes: Bayesian Phylogenetic Inference
Using Graphical Models and an Interactive Model-Specification Language. Syst Biol
65:726–736.

Hu, N., and J. Castelli-Gair. 1999. Study of the posterior spiracles of Drosophila as a
model to understand the genetic and cellular mechanisms controlling morphogenesis.
Developmental Biology 214:197–210.

Huelsenbeck, J. P., and B. Rannala. 2003. Detecting correlation between characters in
a comparative analysis with uncertain phylogeny. Evolution 57:1237–1247.

Jiang, P., M. Z. Ludwig, M. Kreitman, and J. Reinitz. 2015. Natural variation of the
expression pattern of the segmentation gene even-skipped in melanogaster.
Developmental Biology 405:173–181. Elsevier.

Kornberg, T. B., and T. Tabata. 1993. Segmentation of the Drosophila embryo. Current
Opinion in Genetics & Development 3:585–594.

Lartillot, N., and R. Poujol. 2010. A Phylogenetic Model for Investigating Correlated
Evolution of Substitution Rates and Continuous Phenotypic Characters. Molecular
Biology and Evolution 28:729–744.

Lohs-Schardin, M., C. Cremer, and C. Nüsslein-Volhard. 1979. A fate map for the larval
epidermis of Drosophila melanogaster: localized cuticle defects following irradiation of
the blastoderm with an ultraviolet laser microbeam. Developmental Biology 73:239–255.

Lott, S. E., M. Kreitman, A. Palsson, E. Alekseeva, and M. Z. Ludwig. 2007.
Canalization of segmentation and its evolution in Drosophila. Proceedings of the
National Academy of Sciences 104:10926–10931.

Lott, S. E., M. Z. Ludwig, and M. Kreitman. 2010. EVOLUTION AND INHERITANCE OF
EARLY EMBRYONIC PATTERNING IN DROSOPHILA SIMULANS AND D.
SECHELLIA. Evolution 65:1388–1399.

Lucchetta, E. M., J. H. Lee, L. A. Fu, N. H. Patel, and R. F. Ismagilov. 2005. Dynamics
of Drosophila embryonic patterning network perturbed in space and time using
microfluidics. Nature 434:1134–1138.

Manu, S. Surkova, A. V. Spirov, V. V. Gursky, H. Janssens, A.-R. Kim, O. Radulescu,
C. E. Vanario-Alonso, D. H. Sharp, M. Samsonova, and J. Reinitz. 2009a. Canalization
of gene expression and domain shifts in the Drosophila blastoderm by dynamical
attractors. PLoS Comp Biol 5:e1000303.

Manu, S. Surkova, A. V. Spirov, V. V. Gursky, H. Janssens, A.-R. Kim, O. Radulescu,
C. E. Vanario-Alonso, D. H. Sharp, M. Samsonova, and J. Reinitz. 2009b. Canalization
of gene expression in the Drosophila blastoderm by gap gene cross regulation. Plos
Biol 7:e1000049.

Markow, T. A., and P. O'Grady. 2005. Drosophila: a guide to species identification and
use. Academic Press.

Markow, T. A., S. BEALL, and L. M. MATZKIN. 2008. Egg size, embryonic development
time and ovoviviparity in Drosophila species. Journal of Evolutionary Biology 22:430–
434.

Masel, J., and M. L. Siegal. 2009. Robustness: mechanisms and consequences. Trends
in Genetics 25:395–403.

McKenzie, J. A., and S. W. McKechnie. 1979. A Comparative Study of Resource
Utilization in Natural Populations of Drosophila melanogaster and D. simulans.
Oecologia 40:299–309. Springer.

Miles, C. M., S. E. Lott, C. L. Luengo Hendriks, M. Z. Ludwig, Manu, C. L. Williams, and
M. Kreitman. 2010. ARTIFICIAL SELECTION ON EGG SIZE PERTURBS EARLY
PATTERN FORMATION IN DROSOPHILA MELANOGASTER. Evolution 65:33–42.

Namba, R., T. M. Pazdera, R. L. Cerrone, and J. S. Minden. 1997. Drosophila
embryonic pattern repair: how embryos respond to bicoid dosage alteration.
Development 124:1393–1403.

Nasiadka, A., B. H. Dietrich, and H. M. Krause. 2002. Anterior-posterior patterning in the
Drosophila embryo. Pp. 155–204 *in* Gene Expression at the Beginning of Animal
Development. Elsevier.

Nijhout, H. F., F. Sadre-Marandi, J. Best, and M. C. Reed. 2017. Systems Biology of
Phenotypic Robustness and Plasticity. Integrative and Comparative Biology 57:171–
184.

Nüsslein-Volhard, C., and E. Wieschaus. 1980. Mutations affecting segment number
and polarity in Drosophila. Nature 287:795–801.

Nüsslein-Volhard, C., H. G. Frohnhöfer, and R. Lehmann. 2008. Determination of
anteroposterior polarity in Drosophila. Science 238:1675–1681. American Association
for the Advancement of Science.

Obbard, D. J., J. Maclennan, K.-W. Kim, A. Rambaut, P. M. O'Grady, and F. M. Jiggins.
2012. Estimating Divergence Dates and Substitution Rates in the Drosophila
Phylogeny. Molecular Biology and Evolution 29:3459–3473.

Oliveira, D. C. S. G., F. C. Almeida, P. M. O'Grady, M. A. Armella, R. DeSalle, and W. J.
Etges. 2012. Monophyly, divergence times, and evolution of host plant use inferred from
a revised phylogeny of the Drosophila repleta species group. Molecular Phylogenetics
and Evolution 64:533–544.

Paaby, A. B., and M. V. Rockman. 2014. Cryptic genetic variation: evolution's hidden
substrate. Nat Rev Genet 15:247–258. Nature Publishing Group.

Payne, J. L., and A. Wagner. 2019. The causes of evolvability and their evolution. Nat
Rev Genet 20:24–38.

Perry, M. W., J. P. Bothma, R. D. Luu, and M. Levine. 2012. Precision of Hunchback
Expression in the Drosophila Embryo. Current Biology 22:2247–2252. Elsevier Ltd.

Regier, J. C., J. W. Shultz, A. Zwick, A. Hussey, B. Ball, R. Wetzer, J. W. Martin, and C.
W. Cunningham. 2010. Arthropod relationships revealed by phylogenomic analysis of
nuclear protein-coding sequences. Nature 463:1079–1083. Nature Publishing Group.

Russo, C. A. M., B. Mello, A. Frazão, and C. M. Voloch. 2013. Phylogenetic analysis
and a time tree for a large drosophilid data set (Diptera: Drosophilidae). Zool J Linn Soc
169:765–775.

Russo, C. A., N. Takezaki, and M. Nei. 1995. Molecular phylogeny and divergence
times of drosophilid species. Molecular Biology and Evolution 12:391–404.

Rutherford, S. L., and S. Lindquist. 1998. Hsp90 as a capacitor for morphological
evolution. Nature 396:336–342.

Sánchez, L., and I. Guerrero. 2001. The development of the Drosophila genital disc.
Bioessays 23:698–707.

Siegal, M. L., and J.-Y. Leu. 2014. On the Nature and Evolutionary Impact of
Phenotypic Robustness Mechanisms. Annu. Rev. Ecol. Evol. Syst. 45:495–517.

Spieth, H. T. 1981. Drosophila heteroneura and Drosophila silvestris: Head shapes,
behavior and evolution. Evolution 35:921–930. JSTOR.

St Johnston, D., and C. Nüsslein-Volhard. 1992. The origin of pattern and polarity in the
Drosophila embryo. Cell 68:201–219.

Steel, R. G. D., J. H. Torrie, and D. D. A. 1997. Principles and Procedures of Statistics.
McGraw-Hill College.

Surkova, S., E. Golubkova, L. Mamon, and M. Samsonova. 2018. Dynamic maternal
gradients and morphogenetic networks in Drosophila early embryo. Biosystems
173:207–213. Elsevier.

Surkova, S., E. Golubkova, Manu, L. Panok, L. Mamon, J. Reinitz, and M. Samsonova.
2013. Quantitative dynamics and increased variability of segmentation gene expression
in the Drosophila Krüppel and knirps mutants. Developmental Biology 376:99–112.
Elsevier.

Tautz, D. 2004. Segmentation. Developmental Cell 7:301–312.

R Core Team, 2017. R: A language and environment for statistical computing. R
Foundation for Statistical Computing, Vienna, Austria.

Turelli, M., B. S. Cooper, K. M. Richardson, P. S. Ginsberg, B. Peckenpaugh, C. X.
Antelope, K. J. Kim, M. R. May, A. Abrieux, D. A. Wilson, M. J. Bronski, B. R. Moore, J.-
J. Gao, M. B. Eisen, J. C. Chiu, W. R. Conner, and A. A. Hoffmann. 2018. Rapid Global
Spread of wRi-like Wolbachia across Multiple Drosophila. Current Biology 28:963–
971.e8. Elsevier Ltd.

Wagner, A. 2005. Robustness and Evolvability in Living Systems. Princeton University
Press.

Weber, K. E. 1992. How small are the smallest selectable domains of form? Genetics
130:345–353.

Wieschaus, E., and C. Nuesslein-Volhard. 2016. The Heidelberg Screen for Pattern
Mutants of Drosophila: A Personal Account. Annu. Rev. Cell Dev. Biol. 32:1–46.

Wunderlich, Z., C. C. Fowlkes, K. B. Eckenrode, M. D. J. Bragdon, A. Abiri, and A. H.
Depace. 2018. Quantitative comparison of the anterior-posterior patterning system in
the embryos of five Drosophila species. bioRxiv, doi: 10.1101/378430.

Figure 1

D. ananassae

D. persimilis

D. mojavensis

Head T1 | T2 | T3 | A1 | A2 | A3 | A4 | A5 | A6 | A7 | A8 + Telson

X

Y

Segment Position A1 = X/Y

868
869

Figure 1. Dark field images of first instar larvae from three species mounted as
described in Materials and Methods. Red lines mark the anterior border and white lines
mark the posterior border of abdominal denticle belts. Denticle belts were recognized by
our image processing program. The anterior border was used as a proxy for segment
border. To measure the position of an abdominal segment (e.g. A1) relative to the body
length (Y), the distance from the anterior border of the larvae to the anterior border of
that segment (X) is divided by the total body size (Y). T = Thoracic Segment, A =
Abdominal Segment.
878

Figure 2

Figure 2. Twelve Drosophila species show many differences in relative segment position and total body length. (A) Twelve Drosophila species on an evolutionary tree. Adult male species pictures were kindly provided by Nicolas Gompel and were downloaded from FlyBase. (B) Each black bar represents the mean position in percent larval length for each segment in each species. The color spread on the two sides side of each black bar is 95% confidence interval. Each segment is represented by a different color. x-axis is relative segment position in percent larval length. A = Abdominal Segment. (C) Each black bar represents the average first instar larval body length for each species. The gray shaded area around the black bars is 95% confidence interval. x-axis is average body length in micrometers.

Figure 3. Blocks of segments tend to shift together in a certain direction rather than alternating segments shifting in opposite direction. Yellow indicates anterior, blue indicates posterior shift in relative segment position as compared to the species mean for a given segment position. The area of the circle is proportional to the size of the shift in relative segment position as compared to the species mean for a given segment position.

Figure 4

Figure 4. Phylogenetic analysis of segment evolution in the 12 *Drosophila* species. (A) Phylogeny inferred from nuclear loci with relative divergence times. Branches of the tree are colored to indicate the overall rate of relative segment position evolution. (B) Boxplots indicate the posterior distribution of the evolutionary rate of relative segment position. The bar indicates the posterior mean rate; the boxes and whiskers indicate the 50% and 95% posterior credible intervals, respectively. (C) Marginal posterior probability distribution of the evolutionary rates of relative segment position across branches.

916
917
918

Figure 5. Correlations between segments, within and between species. (A) Within a species, correlation between the shift in the relative position of any two segments goes down as the distance between them gets bigger. The sample graph is for *D. melanogaster*. Colors are used to distinguish between segment pairs with different number of segments in between. (B) Over all species, average correlation score between the shift in the relative position of any two segments goes down as the distance between the two segments get bigger. Among all abdominal segments, A8 tends to have a worse correlation score with any given segment regardless of its distance to the other segment. Black dots represent the segment pairs that do not contain A8, red dots represent the segment pairs that contain A8. The gray and red shaded areas around the black and red dots, respectively, is their 95% confidence interval. The unit of the x-axes is in panels A and B is percent larval length. (C) Between species, correlations between evolutionary rates for a pair of segments are lower with increasing distance between segment pairs. Color indicates the first segment in the comparison, with the most blue indicating comparisons with segment 1, and the most yellow being comparisons with segment 7.

Suplementary Figure 1

A control

B heated

C permanently mounteed

937
938
Supplementary Figure 1. (A) Bright field image of a *D. melanogaster* first instar larva,
hatched in water, iced for 2-3 minutes before taking a picture. (B) The same *D.*
melanogaster larva heated at 60°C for an hour prior to taking a bright field picture. (C)
Dark field image of the same *D. melanogaster* larva, mounted using PVA, and
incubated in the 60°C oven overnight. White horizontal lines mark the full length of the
larvae, vertical lines mark the segment borders.
945
946

Supplementary Figure 2

947
948
949 **Supplementary Figure 2.** For the majority of species pair comparisons, five or more
950 segments had significantly shifted relative positions. The graph shows the number of
951 species pairs on the y-axis and number of segments significantly shifted on the x-axis.

957
958

Supplementary Figure 3. (A) Absolute magnitude of shift in relative segment position
as compared to the species mean for each segment position increases from anterior to
the posterior of the larvae. On the y axis of this graph is absolute magnitude of shift, in
percent larval length, for each segment averaged over all species, on the x axis are
abdominal segments A1 through A8. (B) Absolute magnitude of shift changes from
anterior to posterior differently for each of the 12 Drosophila species. The graph shows
absolute magnitude of shift in percent larval length on the y axis and abdominal
segments A1 through A8 on the x axis. The legend shows the slopes calculated for
each trend form anterior to posterior. Lines in the graph and the legend are color
coordinated for each species.

Supplementary Figure 4. (A) Comparison of significant position changes when segment positions were calculated with all segments included, with h+t removed or with A8+tail removed. Removal of A8+tail shows a decrease, whereas removal of h+t shows an increase, in the number of segments that have significantly changed their relative position as compared to when all eight abdominal segments were included in the position calculations. y-axis is total number of significant changes, normalized to the number of segments included in position calculations. (B) When A8+tail was removed from position calculations, number of significant changes in relative segment position went down sharply towards the posterior end of the larvae. However, when h+t was removed in the same manner, number of significant changes in relative segment position followed the same pattern from anterior to posterior as when all abdominal segments were included in position calculations. y-axis is total number of significant changes in segment position. (C) Average magnitude of shift (from the species mean) for each segment went down sharply towards the posterior of the larvae when A8+tail was removed from segment position calculations. On the other hand, removing h+t followed the same pattern as when all segments were included in calculations, but showed a bigger deviation from the species mean towards the posterior of the larvae. y-axis is the average magnitude of shift (from the species mean) in segment position, in percent larval length, ignoring the direction. x-axis is abdominal segments in both panels B and C.

Supplementary Figure 5

Supplementary Figure 5. (A) Average body length difference between a given pair of species is not proportional to the number of segments with significant position shifts. y axis in this graph shows average difference in body length (in pixels) between each pair of species and the x axis shows the number of segments with significant position shifts in the same species pair. (B) This graph shows differences in mean relative size for head+thorax, in percent larval length, among 12 Drosophila species. The black bars indicate the mean and the gray shaded area is 95% confidence interval for each mean. (C) This graph shows differences in mean relative size for A8+tail, in percent larval length, among 12 Drosophila species. The black bars indicate the mean and the gray shaded area is 95% confidence interval for each mean.

Supplementary Figure 6

Supplementary Figure 6. These graphs have the same format as Figure 2B and shows differences in relative segment position between species. (A) Relative segment positions are calculated in the absence of A8+tail. Notice a substantial number of nonsignificant differences as compared Figure 2B. (B) Relative segment positions are calculated in the absence of h+t and also with measurements taken from the posterior of the larva instead of the anterior. As compared to Figure 2B the segment order is flipped, but the pattern of change in segment position is very similar. In both graphs, each black bar depicts the mean for each segment in each species. The colored areas are 95% confidence intervals. The y axis depicts 12 species and x axis shows mean relative abdominal segment position.

Supplementary Figure 7

A

B

Supplementary Figure 7. (A) Change in coefficient of variation, averaged over 12 species, from anterior to posterior segments. The black dots represent the case when relative segment positions were calculated with all segments included. The y axis shows coefficient of variation and the y axis depicts each abdominal segment. (B) Change in standard deviation, averaged over 12 species, from anterior to posterior segments. The black dots represent the case when relative segment positions were calculated with all segments included. The y axis shows coefficient of variation and the y axis depicts each abdominal segment. In both graphs, the red and blue dots represent the cases when relative segment positions were calculated in the absence of A8+tail or h+t, respectively.

1054
1055

Supplementary Figure 8. (A) This graph shows differences in denticle width between
12 species. Blue represents segment A1, red represents segment A8 and gray
represents the rest of the segments. Black bars represent the mean and the colored
area around black bars are 95% confidence intervals. The y axis in this graph depicts
12 Drosophila species, the x axis shows mean denticle width. (B) This graph has the
same format as Figure 2B, however in addition to differences in relative segment
position (anterior border of denticle belts) it also shows changes in the posterior border
of abdominal denticle belts between 12 species. The y axis depicts 12 species and x-
axis shows mean relative abdominal segment position. Bars showing changes in
posterior border of denticle belts are shaded darker than bars representing changes in
the anterior borders. (C) This graph shows deviation from mean, averaged over all
species, for the anterior (black), posterior (blue) and width (red) of denticle belts for
each abdominal segment. The y axis show deviation from mean and the x axis depicts
each abdominal segment.

1070
1071
1072
1073

1074
1075

Supplementary Figure 9. Phylogenetic analysis of segment evolution in the 12 *Drosophila* species, with posterior rate estimates for all branches. A) Like Figure 4A, the phylogeny was inferred from nuclear loci with relative divergence times, and branches of the tree are colored to indicate the overall rate of relative segment position evolution. Here, the branches are numbered to correspond with the values pictured in part B. B) Box plot depicting rate of evolution estimate for each branch of the phylogeny, x-axis corresponds to the branch numbers indicated on the tree in part A.

Supplementary Figure 10
Segment 1

1088

1089

1090

1091

1092

Supplementary Figure 10
Segment 6

Supplementary Figure 10
Segment 7

1093

1094

1095
Supplementary Figure 10. Phylogenetic analysis of relative segment position for each
segment. Like Figure 4A, this figure shows a phylogeny inferred from nuclear loci with
relative divergence times, and the branches of the tree are colored to indicate the
overall rate of segment position evolution. Like Supplementary Figure 9, the tree
diagrams also include numbers for each linage, and are accompanied by a box plot
showing the estimates for rates of segment evolution by each numbered lineage. Here,
there are 8 plots, to show rate estimates for each abdominal segment separately.
1103
1104

1107   **Supplementary Figure 11.** This series of graphs are constructed the same as Figure
1108   6A, which was for *D. melanogaster*, for each of the 12 Drosophila species in our study.
1109   The y axis shows correlation scores and the x axis shows mean relative distance
1110   between pairs of segments.
1111
1112

Supplementary Figure 12

D.melanogaster Correlation Score Heatmap

Supplementary Figure 12

D.simulans Correlation Score Heatmap

Supplementary Figure 12

D.sechellia Correlation Score Heatmap

Supplementary Figure 12

D.yakuba Correlation Score Heatmap

Supplementary Figure 12

D.santomea Correlation Score Heatmap

Supplementary Figure 12

D.erecta Correlation Score Heatmap

1113

1114

Supplementary Figure 12. Series of correlation score heat maps for each of the 12 species in our study. There is a square showing the correlation score for the shift in the relative position of each segment pair within a species. Red means high positive correlation and white means low positive correlation. Actual correlation scores are printed on each square box. Notice that for most species, the color scheme is lighter at the right edge of the triangle as compared to the bottom edge of the triangle. This is consistent with other figures showing shifts in the position of A8 has worse correlation with shifts in the position of other segments, regardless of the distance in between.

1125

1126 **Supplementary Figure 13.** This series of graphs represent the information from Figure
1127 6B, in a zoomed in format. Each figure has mean correlation score over all species on
1128 the y axis and mean relative distance over all species on the x axis. Segments that are
1129 paired with A8 are in red. Error bars are 95% confidence intervals.
1130
1131

Supplementary Figure 14. This series of graphs show how correlation scores change
from anterior to posterior in all species. In each graph in the series, segment pairs
become further apart from each other (from one to seven segments apart). y axis
represents correlation scores and the x axis represents each of the eight abdominal
segments. Each species is represented by a different color. Notice, especially as the
pairs of segments become further apart, *D. erecta*, *D. mojavensis* and also *D. virilis*
stand out as the species with the highest correlation scores.
1140
1141

1144 **Supplementary Figure 15.** This series of graphs show how correlation scores change
1145 for each species from anterior to posterior. It is essentially a broken down version of the
1146 first graph in Supplementary Figure 12, without "A7.A8". Slope for each trend line is
1147 reported in the legend to give an idea of direction of change in correlation scores.

Supplementary Figure 16

1160
Supplementary Figure 16. When all segments are included in position calculations,
mean correlation scores between position shifts of segments that are next to each other
are highest ever so slightly in the middle of the larva. In comparison, when A8+tail is
removed from position calculations, correlation between segment position shifts at the
posterior of the larvae goes down significantly. This is true for the anterior part of the
larva when h+t are removed from position calculations.
1167
1168

1169

1171 **Supplementary Figure 17.** This series of graphs are a broken-down version of
1172 Supplementary Figure 16 into each species. They show, for each species, how
1173 correlation scores change from the anterior to the posterior of the larvae, when relative
1174 segment position was calculated with all segments included (black) versus no A8+tail
1175 (red) or no h+t (blue).
1176
1177

1178

1179 **Supplementary Figure 18.** This series of graphs are a continuation of Supplementary
1180 Figure 16, as the distance between pairs of segments increase, from two to six
1181 segments apart. Notice that removal of h+t overall decreases correlation scores more
1182 than removal of A8+tail.
1183
1184 **Supplementary Table 1.** Empirical data collected in the Lott Lab on number of flies and
1185 days needed in a bottle to control population density of each species. Also includes, for
1186 each species, approximate embryonic developmental duration at 20°C and number of
1187 minutes needed in the 60°C oven for larvae to become straight.

Species Name	Number of Male and Female flies per bottle	Number of days in bottle before removing adults	Number of days from egg to larva at 20°C (hours)	Number of minutes at 60° oven
D. melanogaster (Ore	15 each	4-5 days	~36	50
D. simulans	15 each	4-5 days	~24	50
D. sechellia	20 each	7 days	~36	90
D. yakuba	10 each	4-5 days	~24	50
D. santomea	25 each	7 days	~36	50
D. erecta	10 each	4-5 days	~24	50
D. ananassae	20 each	7 days	~36	50
D. pseudoobscura	10 each	7days	~36	50
D. persimilis	15 each	7 days	~36	50
D. willistoni	25 each	7 days	~48	50
D. mojavensis	25 each	7 days	~72	45
D. virilis	15 each	7 days	~60	90

1188
1189
1190
1191 **Supplementary Table 2.** Lists the number of segments that are differentially
1192 positioned between pairs of species
1193
1194 **Supplementary Table 3.** Lists the deviation of the position of each segment in each
1195 species from the "across-species" mean.
1196
1197 **Supplementary File 1.** Methods and results of the phylogenetic analysis.
1198
1199 **Supplementary File 2.** Additional information on the observed changes in the position
1200 of posterior border of denticle belts and denticle width between species.
1201
1202 **Supplementary File 3.** Description of the effects of removal of A8+tail or h+t from
1203 segment position calculations. In particular, effects of the new calculations on
1204 differences in segment position between species as well as the noise in segment
1205 position within a species.
1206