A protein-centric approach for exome variant aggregation enables sensitive association analysis with clinical outcomes

Ginny X.L. Li[1,2], Dan Munro[3], Damian Fermin[4], Christine Vogel[3], Hyungwon Choi[1,5,*]

1. Department of Medicine, Yong Loo Lin School of Medicine, National University of Singapore, Singapore

2. Saw Swee Hock School of Public Health, National University of Singapore, Singapore

3. Center for Genomics and Systems Biology, Department of Biology, New York University, New York, USA

4. Department of Pediatric Nephrology, University of Michigan Medical School, Ann Arbor, Michigan, USA

5. Institute of Molecular and Cell Biology, Agency for Science, Technology, Research, Singapore

*Corresponding author: Hyungwon Choi (hwchoi@nus.edu.sg)

Abstract

Somatic mutations are early drivers of tumorigenesis and tumor progression. However, the mutations typically occur at variable positions across different individuals, resulting in the data being too sparse to test meaningful associations between variants and phenotypes. To overcome this challenge, we devised a novel approach called Gene-to-Protein-to-Disease (GPD) which accumulates variants into new sequence units as the degree of genetic assault on structural or functional units of each protein. The variant frequencies in the sequence units were highly reproducible between two large cancer cohorts. Survival analysis identified 247 sequence units in which somatic mutations had deleterious effects on overall survival, including consensus driver mutations obtained from multiple calling algorithms. By contrast, around 75% of the survival predictive units had been undetected by conventional gene-level analysis. We demonstrate the ability of these signatures to separate patient groups according to overall survival, therefore providing novel prognostic tools for various cancers. GPD also identified sequence units with somatic mutations whose impact on survival was modified by the occupancy of germline variants in the surrounding regions. The findings indicate that a patient's genetic predisposition interacts with the effect of somatic mutations on survival outcome in some cancers.

Keywords: exome sequence variants, prognostic signatures, sequence units

Introduction

Accumulation of mutations in the genome can impair cellular function, lead to abnormal proliferation of cells, and eventually cause cancer (Martincorena and Campbell, 2015). Today's massively parallel sequencing technology can detect such variation in the whole genome or exome, and simultaneous analysis of paired tumor and normal samples enables reliable calling of somatic mutations and germline variants. Large-scale sequencing projects such as The Cancer Genome Atlas (TCGA) have successfully mapped the mutation landscape of >11,000 patients across 33 cancer types (Ding et al., 2018; Ellrott et al., 2018; Huang et al., 2018; Liu et al., 2018). The next logical step is to test for the association between cancer genome variants and clinical outcomes such as prognosis and response to first-line therapies (Castro et al., 2013; Gentles et al., 2015; Dellinger et al., 2016). For example, somatic driver mutations are responsible for tumor maintenance, progression, and metastasis (Reimand and Bader, 2013), which are robust indicators of patient survival.

Indeed, mutations at specific loci can make a marked difference in disease prognosis. For example, tumors from papillary thyroid carcinoma patients with the missense mutation V600E in the *BRAF* gene are often associated with more rapid tumor growth and higher death risk than the patients without the mutation (Xing, 2005), Further, missense mutations affect the transcriptional activity of tumor suppressor TP53 in different ways, as has been shown in a study of ~1,800 primary breast cancer patients (Olivier, 2006). Mutations found within the DNA-binding motif, especially those at codon 179 and 248, are associated with poorer patient survival outcome than mutations at other positions. Other than these well-known examples, however, systematic identification of variants useful for prognosis still remains a major undertaking of cancer genomics.

In the past, a plethora of algorithms have been developed to differentiate truly cancer causing, driver mutations from abundant passenger mutations. MuSic (Dees et al., 2012), OncodriveCLUST (Tamborero et al., 2013), and MutSigCV (Lawrence et al., 2013) identify the genes that harbor significantly more mutations than others based on cancer-specific

background mutations; SIFT (Kumar et al., 2009), FATHMM (Shihab et al., 2013),

PolyPhen2 (Adzhubei et al., 2010), and CHASM (Wong et al., 2011) identify mutations by

evaluating their functional impact; e-Driver (Porta-Pardo and Godzik, 2014) and ActiveDriver

(Reimand and Bader, 2013) focus on mutations residing at structurally important sites; and

Network-Based Stratification (Hofree et al., 2013), PARADIGM (Vaske et al., 2010), TieDIE

(Paull et al., 2013), and DriverNet (Bashashati et al., 2012) prioritize driver mutations using

network- and pathway-based approaches. In a recent massive meta-analysis of the Pan-

Cancer Atlas project, Bailey *et al. (Bailey et al., 2018)* applied 26 different computational

tools and identified 299 consensus driver genes and ~3,400 mutations across 33 cancer

types -- one of the most comprehensive such studies known to date.

Despite these developments, it remains challenging to test statistical association

between mutations and meaningful clinical outcomes such as patient prognosis. One reason

behind this challenge is the extreme heterogeneity of tumor genomes (McGranahan and

Swanton, 2015). In contrast to studies of single-nucleotide polymorphisms in healthy

individuals, somatic mutations detected in tumor specimens rarely reside at the same locus

across many patients (Pickrell et al., 2016). As a result, most tumor variants have very low

frequency, e.g. on average <0.011% across TCGA Pan-Cancer Atlas patients, with most of

them detected in a single patient. Further, the issue potentially reflects the varying genome

instability across patients, as is typical for cancer cells (Jeggo et al., 2016). As a

consequence, statistical analysis of somatic mutations at the locus level inherently lacks the

statistical power to detect prognostic markers (Kiiski et al., 2016).

A common remedy to this challenge is to count mutations *per* gene, or to classify the

mutation status of a gene as 'mutated' or 'not mutated' in each patient (Candido-dos-Reis et

al., 2015). A disadvantage of this approach is that it considers patients with different

mutation profiles as carriers of the same mutation burden, regardless of the positions of

mutations. An alternative systems-oriented approach, often referred to as network-based

stratification methods, propagates the binary mutation status of genes onto a protein-protein

interaction network and identifies interacting genes enriched with adjacent mutation events

as etiologically important players (Hofree et al., 2013; Zhong et al., 2015; Hristov and Singh,

2017; Le Morvan et al., 2017). While this method more sensitively identifies mutation

clusters than locus-based analysis, it ignores the relative positions of mutations in the gene

sequence and subsequently fails to provide directly testable hypotheses on the gain or loss

of protein function.

To counter these challenges and enable sensitive identification of prognostic exome

variants and preserve information on the location of the mutations, we developed a new

approach called Gene-to-Protein-to-Disease (GPD), which maps somatic mutations (and/or

germline variants) onto small sequence units located in protein coding or non-coding

regions. The approach therefore transforms the original variant data into substantially less

sparse count data and enables statistical analysis for disease prognosis. It successfully

recovers a handful of cancer driver mutations as prognostic markers, and also enables

identification of interactions between the effects of somatic mutations and germline variants

on survival outcomes.

Materials and Methods

Data Source. Exome sequencing data for somatic mutations in TCGA datasets were downloaded from the Genomic Data Commons (MC3 Public MAF) (Ellrott et al., 2018). Germline variant data were downloaded from the ISB-CGC web interface, where the data is deposited in a VCF file format (Huang et al., 2018). We further annotated the file with the snpEff tool (genome version GRch37.75) (Cingolani et al., 2012). In addition, we gathered 348,658 protein modification sites (244,852 phosphorylation, 46,352 ubiquitination, 25,807 acetylation, 20,691 methylation, 8,388 SUMOylation, and 2,568 glycosylation sites) from PhosphoSitePlus (Hornbeck et al., 2015), and 45,607 domains, families and repeats for 19,076 genes from Pfam (El-Gebali et al., 2019).

Survival outcome data were downloaded from TCGA Pan-Cancer Clinical Data Resource (Liu et al., 2018), which contains curated clinical information for 10,793 patients. Among these, 367 non-primary skin cutaneous melanoma patients with metastatic tumors were excluded. The data resource provides four types of clinical endpoints, including overall and disease-specific survival (OS, DSS), as well as disease-free and progression-free time interval (DSI, PFI). As advised by Liu et al., OS and PFI are the most reliable endpoints for the majority of the cancers. Hence, we report the results for the OS as the endpoint of interest in this study, while reporting the results for PFI in the **Supplementary Information**. Abbreviations for cancer types have been adapted from TCGA available at https://gdc.cancer.gov/resource-tcga-users/tcga-code-table/tcga-study-abbreviations.

Mapping scheme. GPD segments exons and adjacent genomic regions of each gene into three types of units (**Figure 1A**). First, Protein Information Units (PIU) refer to the genomic regions encoding protein domains, e.g., as defined by the Pfam database (El-Gebali et al., 2019)) or +/-5 amino acid-long windows around protein modification sites (e.g., as reported in the PhosphoSitePlus database). Sequence regions between PIUs are defined as linker units (LU). The LUs not only include linker regions between domains, but also cover

unannotated, repeat or disordered regions. The regions outside the protein-coding

sequences, including untranslated regions, introns, and regulatory regions are collectively

defined as non-coding units (NCU). NCUs are assigned to the closest gene in the genome.

Synonymous variants which do not result in amino acid changes in protein

sequences were excluded from the analysis. Missense mutations resulting in single amino

acid changes represent 68.2% of all genetic alterations in the data. These missense

mutations are unambiguously mapped to sequence units and counted. For truncating

mutations or those that can introduce large sequence changes (e.g. unexpected stop

codon), GPD counts them the same way as it counts missense mutations, since it is difficult

to quantify the effect of these mutations in terms of the mutational load on sequence units.

To understand the proximity between non-coding variants and DNA coding regions,

we downloaded a GTF file on GRCh37 human genome annotation from GENCODE website

(https://www.gencodegenes.org/human/release_32lift37.html). From this GTF file, we

extracted the coordinates of start and stop codons for each gene. For non-coding variants

outside of the translation interval, we calculated the distance (in base pairs) between the

variant loci and the first base of start codons if they lie in the upstream region of the

translation interval or the last base of stop codons if they are in the downstream region of the

translation interval.

Validation of mapping with MSK-IMPACT data. MSK-IMPACT cohort data was retrieved

from https://www.cbioportal.org/study/summary?id=msk_impact_2017. This data includes

exome-sequencing data from a total of 10 thousand patients across 14 cancer types. We

implemented GPD mapping scheme to map mutations occurred in primary tumor samples

collected in the study to PIUs, LUs and NCUs. Then we computed per sample mutation

counts (mutation frequency) in each common PIU and LU mapped by both TCGA data and

MSK-IMPACT data. Spearman correlations were calculated for each pair of mutation

frequencies to measure the correlation between mapping results generated from two

independent datasets.

Selection of prognostic sequence units (main effect model). The association between mutation count of each sequence unit and OS (or PFI) was evaluated using Cox proportional hazard regression. To focus on the association with long-term survival outcome, we excluded patients with less than 90 days survival time. For each cancer, we included sequence units with mutations in three or more patients for all subsequent analyses. We adjusted all regression models for the total mutation count, age, gender, race and tumor stage of each patient (available in ACC, BLCA, BRCA, CHOL, COAD, ESCA, HNSC, KICH, KIRC, KIRP, LIHC, LUAD, LUSC, MESO, PAAD, READ, SKCM, STAD, TGCT, THCA, UVM). The Cox model posits the hazard function as a product of the baseline risk function at time *t* and the relative risk based on covariates. For each sequence unit, the hazard function for subject *i* is given by:

$$\lambda(t|X_i) = \lambda_0(t)exp(X_i \cdot \beta)$$

Where $\lambda_0(t)$ denotes the baseline hazard function and $X_i = \{Count_i, TotalMutation_i, Age_i, Gender_i, Race_i, Stage_i\}$ are the covariates. For comparison, we repeated the analysis but counted mutations *per* gene (gene-level analysis), instead of *per* sequence unit. For each model, the proportional hazard assumption was tested by evaluating the association between transformed survival time and the scaled Schoenfeld residuals (Schoenfeld, 1982).

Validation of sequence unit signatures for the predictability of survival outcomes. We conducted 10-fold cross validation with supervised principal component method (Bair et al., 2006) for each cancer cohort to validate the robustness of predictability of the signatures. In each iteration of the 10-fold of validation, we used 9 partitions of randomly split samples as training data. We identified sequence units significantly associated with OS (*q*-value 0.05) and synthesized up to five principal components (PC) of the selected units and fit another cox proportional hazards model consisting of the covariates $X_i = \{PC1_i, PC2_i, PC3_i, PC4_i, PC5_i, TotalMutation_i, Age_i, Gender_i, Race_i, Stage_i\}$.

189   After fitting the models, we computed the predictive index (PI) for each training

190 sample as a measurement of survival risk.

$$PI = X \cdot \hat{\beta}$$

191 where $\hat{\beta}$ represents coefficients from the above models. Using the remaining one-fold of

192 the data as test data, we first projected them onto the PCs with the loadings of principal

193 components derived with training data, and computed the PIs for test samples as described

194 above. Finally, test samples were clustered into the high-risk group if their PI is higher than

195 the median of training samples' PIs or the low-risk group otherwise. After iterating the

196 process 10 times, we have assigned the samples into high- or low-risk groups. We then test

197 the difference in survival between these two groups with log-rank test.

198   To investigate the degree of overfitting, we shuffled the survival outcomes among the

199 samples 1,000 times. For each shuffled data, we conducted the 10-fold cross validation

200 described above. This process generates 1,000 log-rank statistics, allowing us to form the

201 empirical null distribution of the statistic, representing the assumption that there is no

202 association between sequence unit-level mutation counts and survival outcomes. Therefore,

203 the significance of survival difference in the original data can be derived by locating the log-

204 rank statistic in the empirical distribution.

206 ***Identification of interactions between effects****.* The hazard models accounting for the

207 interaction between somatic mutations and germline variants consider three main covariates:

208 the somatic mutation count of a unit, the total count of germline variants in the same gene,

209 and the interaction between the two. Similar to the main effect models above, models were

210 adjusted for total mutation count, gender, age, race and tumor stage. The hazard function is

211 identical to the above, but denotes a vector of covariates as:

212 $X_i = \{Count_{i_{somaticUnit}}, Count_{i_{germlineAllUnits}}, Count_{i_{somaticUnit}} * Count_{i_{germlineAllUnits}},$

213     $TotalMutation_i, Age_i, Gender_i, Race_i, Stage_i\}.$

GPD software implementation. The GPD framework was developed as an R package. The

package is publicly available at https://github.com/ginnyintifa/GPD, with user-defined GPD

units and control parameters for mapping and analysis. A user manual is available at the

aforementioned GitHub site. The GPD implementation maps mutations to sequence units.

The default genomic coordinates of protein domains and protein modification sites are

provided in the package, but the user can also provide the information from other sources in

the specified format. Users need to provide mutation data.

Results

GPD maps variants to "sequence units" encoding well-defined functional segments of

protein sequences

Figure 1A illustrates the concept of GPD's sequence units through a hypothetical exome

with somatic mutations: two mutations occurred within a protein domain (PIU1), one

mutation near a protein modification site (PIU2), two mutations in a LU outside the PIUs, and

three mutations in the non-coding unit. Using the protein domains from Pfam (El-Gebali et

al., 2019) database and the protein modification sites from PhosphoSitePlus (Hornbeck et

al., 2015) database as PIUs, GPD mapped all somatic mutations extracted from TCGA to a

total of 270,470 PIUs, 17,252 LUs, and 17,266 NCUs in 17,744 human genes, respectively.

Next, we counted the coding and non-coding variants in sequence units *per* patient,

where the somatic mutations were called by TCGA Pan-Cancer Atlas (Ellrott et al., 2018).

Mapping to GPD's sequence units revealed that different cancers showed unique somatic

mutation frequencies. For example, uterine cancer (UCEC) had on average >1,000

mutations *per* patient; skin and colon cancers (SKCM and COAD) had on average >500

mutations *per* patient. By contrast, the exomes in many other cancers (e.g. KICH, UVM,

TGCT, THCA and PCPG) had fewer than 30 somatic mutations *per* patient on average.

As well known in the literature, the overall distribution of somatic mutations in the

exomes of individuals was remarkably sparse and heterogeneous. Less than 8% of the

somatic mutations appeared in two or more patients at the same locus (**Figure 1B**). As

expected, the sparsity in the data due to the lack of common loci was significantly reduced

when accumulating mutations *per* gene or *per* sequence unit. While 99% of the genes

showed mutations, we found that 68% of PIUs, 96% of LUs, and 98% NCUs, respectively,

harbored somatic mutations in two or more patients. The variable mutation frequencies

across different sequence units within the same gene suggest that GPD's mapping approach

provides a higher-resolution map than gene-level mapping, as it can focus on smaller

sequence regions while still preserving the coverage and statistical power to compare

patient groups.

When examining the distribution of mutations across different genes, we noticed that

almost half the somatic mutations were within PIUs (49%), one third were within LUs (32%),

and 19% were in NCUs (**Figure 1C**). Among the mutations mapped to NCUs, 32% lie

between start codon and stop codon, indicating that they are located in introns. For the non-

intronic mutations in NCUs, 23% of the mutations are positioned upstream of a start codon

and 77% are positioned downstream of a stop codon. A closer investigation of the distance

between these non-intronic loci outside the coding regions to either the nearest start codon

or stop codon revealed that 78% of them are located within 1,000 base pairs from either start

codon or stop codon, and 92% are within 3,000 base pairs, implying that they are positioned

in promoter regions, 5' UTR, or 3' UTR (**Supplementary Figure 1**). In the current definition

of GPD's sequence units, 67% of the protein coding sequence is covered by PIUs and 33%

is covered by LUs. Accordingly, within the protein coding regions, both unit types have about

equal chance to carry a mutation: the expected frequency of a mutation is 0.17 and 0.16 for

PIUs and linker units, respectively.

Figure 1D shows the PIUs with the largest frequency of somatic mutations in TCGA

patients. Not surprisingly, these PIUs are located in well-known cancer genes. For example,

the PIU for the DNA-binding domain of *TP53* gene showed consistent somatic mutations

across the majority of the cancers. Other mutations surrounded the serine phosphorylation

site S269 of the protein, which is crucial for the conformational stability of the protein (Fraser

et al., 2010). In another gene, *KRAS*, we found the highest frequency of somatic mutations

in the PIU corresponding to the RAS domain of the protein in pancreas, lung, rectum, and

colon cancer (PAAD, LUAD, READ, and COAD). The RAS domain also accumulates most

mutations in the *NRAS* gene in 103 non-metastatic skin cancer (SKCM). Further, we find

frequent mutations in thyroid, skin, and colon cancers (THCA, SKCM, and COAD) in a

region in the *BRAF* gene densely populated with phosphorylation and ubiquitination sites

(T599, K601, S602, S605, and K601). Similarly, the cancer gene *PTEN* accumulated

mutations around lysine acetylation sites K125 and K128. **Supplementary Table 1**

enumerates the sequence units with high mutation rates in specific cancers.

Next, to assess the reproducibility of the mapping scheme by sequence units, we

validated the results above (see **Materials and Methods**) with somatic mutation data for

nine cancers from an independent cohort (MSK-IMPACT) (Zehir et al., 2017). As mutations

found outside protein coding regions (i.e. NCUs) were excluded in MSK-IMPACT, we

mapped variants to PIUs and LUs only. We found that the somatic mutation frequencies in

the sequence units are highly reproducible between the TCGA and MSK-IMPACT cohorts

(**Figure 1E**), suggesting that somatic mutations have the propensity to be enriched in

specific GPD units within each gene.

Prognostic signatures of somatic mutations for overall survival are validated in eight

cancer types

To identify sequence units harboring somatic mutations of prognostic potential, we tested

each sequence unit for positive association with OS in Cox proportional hazard models (see

Materials and Methods). We discovered 196, 170, and 107 unique PIUs, LUs and NCUs

that were associated with OS in at least one of 24 cancer types, with mutation(s) occurring in

at least three patients (*q*-value 0.05). Also, Schöenfeld residuals suggested no violation of

proportional hazards assumption in these sequence units. We performed 10-fold cross

validation on each cancer type (see **Materials and Methods**) to examine if the discovered

prognostic units are capable of predicting survival outcomes in a reproducible manner. In

eight cancer types (BLCA, BRCA, ESCA, GBM, KIRC, LGG, OV, UCEC), the predictive

indices can cluster patients into two risk groups with significantly different survival outcomes

(**Figure 2** and **Table 1**). Moreover, we conducted a permutation test to evaluate the degree

of overfitting in the cross-validation process (see **Materials and Methods**). These eight

cancers survived the test with no sign of overfitting (**Table 1** and **Supplementary Figure 2**).

Therefore, our final prognostic signatures consist of 103 PIUs, 86 linker and 59 non-coding

units discovered in these cancer cohorts (**Supplementary Table 2**).

Prognostic signatures associated with poorer survival outcomes are cancer-specific

The vast majority of Cox models with significant findings had positive regression coefficients

(245 out of 247), suggesting deleterious impact of mapped somatic mutations on patient

survival. **Figure 3A** illustrates that UCEC has the greatest number of prognostic signatures,

followed by BLCA, BRCA, and LGG. In sum, PIUs outnumber LUs and NCUs. Prognostic

PIUs uncovered by GPD consists of protein domains and PTM sites on genes implicated in

various biological processes (**Figure 3B**). A large proportion of PIUs are enzymes catalyzing

chemical reactions in cellular processes. For example, somatic mutation counts in the

hydrolase domain in *NIT2* gene for ovarian cancer patients (OV) and those in FAH and

ATP13A3 genes for uterine cancer (UCEC) patients were significantly associated with

overall survival. Mutations mapped to the peptidase domain in *FURIN*, *LGNM* and *CASP10*

genes in UCEC patients, and in *TINAG* gene in breast cancer (BRCA) patients are also

associated with poorer survival outcomes, highly likely by interfering with the catabolism of

peptides into amino acids. Domains involved in altering the genetic material constitute a

good proportion as well. For example, two histone domains in *H3F3A* and *H3F3B* have

mutations seen in three lower grade glioma (LGG) patients individually; N-terminal Myc

family in the transcription factor *MYC* gene harbours mutations in seven UCEC patients.

Mpp10 domain in *MPHOSPH10* gene is related to the processing pre-rRNA to mature rRNA,

and is a possible prognosis predictor in bladder cancer (BLCA) patients. Cadherin domains

are responsible for the formation of adherens junctions to bind cells with one another. GPD

revealed four cadherin-related domains in *PCDH11X* (LGG), *CHD9* (UCEC), *PCDHA7*

(BLCA) and *FAT1* (BLCA) that are significantly associated with overall survival. Five

sequence units encapsulating PTM sites were also discovered to be prognostic for overall

survival. These include a tyrosine phosphorylation site in *IDH1* (LGG), a threonine

phosphorylation site in *SLC9A4*, an arginine methylation site and a threonine

phosphorylation site in *TP53* (UCEC), and lastly, a threonine phosphorylation site in *TP53*

(BRCA).

It is well known that the isocitrate dehydrogenase *IDH1* is highly mutated in LGG

patients (78.3%). An examination of the mutational landscape brings attention to a hotspot

arginine 132 (R132), encapsulated the domain Iso_dh and a PTM site window for a tyrosine

phosphorylation site (position 130-140) mentioned above. In most cases, the frequent

missense mutation converts the arginine residue to histidine. A large body of evidence

suggests that the detection of mutated IDH in glioma affords a survival advantage compared

with wild-type IDH. (Lu and McDonald, 2018). GPD confirms this phenomenon with negative

log coefficients (-1.64) from the Cox proportional hazards model for the unit counts in both

the Iso_dh domain and the tyrosine phosphorylation site window. Epidermal growth factor

receptor *EGFR* is also highly mutated in different sequence regions in LGG patients. 19 out

of 457 patients have mutations in the Furin-like domain, 6 in GF_recep_IV domain and 5 in

Recep_L domain. Overall, we observed that prognostic markers are highly specific to one

cancer type, possibly suggesting that the most prognosis-related functional alterations

caused by somatic mutations are cancer specific.

GPD discovers associations overlooked by gene-level analysis

When somatic mutations are counted at the whole gene level, we obtained substantially

different sets of survival associated genes (**Supplementary Figure 3**). In sum, > 75% of the

genes (182 in 241) containing at least one significant sequence units were not discovered by

gene-level analysis (**Supplementary Table 3**). The extreme case is in oesophageal

carcinoma (ESCA), where two genes *ZNF459* and *RTN1* discovered by gene-level analysis

were seen in unit-level results, and three other prognostic units bearing genes *EML4*,

LPAL2, *REG3A* are novel findings. In glioma (LGG), *IDH1*, *EGFR*, *PTEN* and many other

known genes were discovered in both levels, but histone domains in *H3F3B* and *H3F3A*

were overlooked by the gene-level analysis. For the genes reported by both approaches,

unit-level analysis highlights specific regions associated with survival outcomes. Moreover,

unit-level analysis adds a total of 185 high-resolution discoveries in 182 novel genes to the

set of prognostic markers.

Survival-associated sequence units moderately overlap with driver mutations with

prognostic values

We next compared survival-associated sequence units against known driver mutations and

their corresponding genes (**Supplementary Tables 4** and **5**). To this end, we used the 299

cancer driver genes and ~3,400 driver mutations identified *in silico* by Bailey *et al.*, who

employed a multitude of software tools to obtain the consensus list with data from the TCGA

Pan-Cancer Atlas cohort (Bailey et al., 2018). We identified multiple PIUs and one LU within

eight known driver genes, namely *FAT1, TBX3, TP53, IDH1, EGFR, PTEN, PLCG1,* and

NF1 (**Supplementary Figure 3**). Further, using the position information for the core set of

579 missense driver mutations, we mapped the mutations to prognostic sequence units. 32

missense driver mutations were found in seven prognostic units, including Furin-like domain,

Receptor L-domain and GF_recep domain in *EGFR*, phosphorylation site T211 and

methylation site R213 in *TP53*, Iso_dh domain and phosphorylation site Y135 in *IDH1*. This

relatively scarce overlap between sequence units and driver genes/mutations is not

surprising, since driver mutations which confer a selective growth advantage to tumors are

not necessarily predictive of patient survival. Moreover, GPD only identified a small number

of sequence units with highly stringent criteria for association with survival (passing the

thresholds of q-value 0.05 and 10-fold cross-validation). However, the aforementioned genes

and sequence units overlapping with GPD's unit level analysis are structural components of

genes that can prognosticate clinical endpoints in respective cancers, adding reliability to the

reported discoveries by the GPD.

Interactions between somatic and germline variants in the association with overall

survival

We have shown that GPD allows for detection of sequence units enriched with variants that

are associated with OS. Next, we exploited GPD's capability to test for interactions between

germline variants and somatic mutations within each gene in the association with overall

survival. Such analysis is unique and of high clinical importance, as each subject's germline

genome may modulate the effects of somatic mutations.

We surveyed all sequence units in the eight cancer types, for which at least three

patients have both somatic mutations in each sequence unit and germline variants in the

gene containing the unit. This requirement rendered 8,240 PIUs, 9,641 LUs, and 5,209

NCUs in 8,821 genes eligible. For each sequence unit, we tested for the main effects of

somatic mutation counts and germline variant counts as well as the interaction between the

two, with adjustment of potential confounders (see **Materials and Methods)**. We did not

adjust for multiple testing because the test could be performed for a relatively small number

of units in each cancer cohort due to inclusion criteria explained above.

Figure 4A shows that bladder cancer (BLCA) had the largest number of interaction

effects, followed by uterine (UCEC), ovarian (OV), brain (GBM) cancers. **Supplementary**

Table 6 provides the list of 520 significant interactions across the 8 cancer types.

Interestingly, LUs appear to occupy the majority of significant interactions. In testing somatic

mutations alone, more PIUs are significant (**Figure 3A**). Around half the interactions (280 out

of 520) had positive coefficients, suggesting synergistic adverse effects of simultaneous

occurrence of somatic mutations and germline variants on overall survival in these cases.

Figure 4B shows examples demonstrating how the unit-level frequency of germline

variants modifies the hazard ratio of somatic mutation counts. For example, there were

multiple occurrences of both germline variants and somatic mutations in the PIU surrounding

the arginine methylation site R110 of the *TP53* gene. Fifteen out of 907 breast cancer

(BRCA) patients had at least one somatic mutation in this sequence unit and 234 patients

had more than ten germline variants on that gene. We found that the difference in the hazard

ratios between a patient with somatic mutation(s) and a patient without one was much

greater if the patients had more germline variants. Similar patterns were observed in the

DST and *CAMTA1* genes in their respective cancer types. In comparison, as seen in the

bottom panel of **Figure 4B**, having very few germline variants along with a somatic mutation

decreased the hazard ratio and elevated the risk of death substantially in patients with more

germline variants.

In sum, these findings suggest that substantial germline variation near the sequence

units predisposes patients to a higher risk of death. Somatic mutations coupled with more

germline variants in the same gene tend to have enhanced negative effects on survival,

while somatic mutations surrounded by few germline variations may have protective effects.

Discussion

We present a new approach, GPD, which summarizes sequence variation and mutations as frequency in small sequence units, providing two main advantages over alternative approaches. First, mutation count data for GPD's sequence units mitigate the sparsity problem of locus-level exome sequencing data, addressing the inherent drawback that most exome variants do not occur at the same position with reasonably high frequency across different subjects. The sequence units therefore allow for immediate downstream statistical analysis to test their association with clinical outcomes. However, we acknowledge the limitation that our mapping scheme ignores potentially different protein sequence alterations caused by different types of mutations (e.g. treating missense mutations and early stop codon changes the same).

Second, GPD exploits the fact that variant data are not randomly distributed across a gene's sequence but accumulates in specific regions. In contrast to conventional gene-level summaries, it does not sacrifice statistical power. At the same time, the sequence units associated with the OS inform on the direct functional consequences of the variants, facilitating the interpretation and enabling prioritization of targets for downstream experimental validation. Combined with survival analysis and association analysis for other clinical endpoints such as therapeutic response, the information can streamline patient-specific prognosis. To illustrate this, we have repeated the same analysis pipeline with PFI as the clinical outcome, where we report positive signatures validated in six cancer types. The results can be found in **Supplementary Information** document.

We first validated our findings through their reproducibility across a different large-scale dataset (MSK-IMPACT), confirmation by well-known cancer mutations, enrichment in protein domains implicated in cancer-related processes amongst survival associated genes, and overlap with an independently identified set of driver mutations. Furthermore, most sequence units with somatic mutations were highly patient-specific: they occurred in a small percentage of samples across cancers.

Another major advance in our work is the assessment of the combinatorial impact of

different types of sequence variation, identifying interactions between existing genetic

predisposition of a patient (germline variants) and tumorigenic events (somatic mutations)

with respect to disease prognosis. To date, only few other studies have explored the

interaction between germline and somatic variation, but in a limited fashion. For example,

Carter *et al.* (Carter et al., 2017) analysed ~6,000 tumors and identified 412 genetic

interactions between germline polymorphisms and major somatic events. However, the

analysis was limited to 138 known oncogenes and tumor suppressor genes and focused on

the effect of inherited polymorphisms in germline on the likelihood of somatic mutations,

rather than discovering the prognostic value of the interaction between the two classes of

sequence variation. Therefore, to the best of our knowledge, the interaction between genetic

pre-disposition and somatic mutations has not been systematically addressed through

statistical association analysis in the genome scale.

We found many genes in which patients with few germline variants and at least one

somatic mutation in the same gene had better OS. By contrast, in other genes with many

pre-existing germline variations, we found that additional somatic mutations almost

exclusively had deleterious effects on overall survival. This finding implies that genetic

predisposition (germline variants) affects the impact of subsequent mutation events on a

gene and their indirect impact on patient survival -- highlighting a new avenue for

individualized treatment based on the patient's exome data. Given that GPD provides a

highly flexible tool that can be applied to any definition of sequence units and large-scale

datasets, integrative analyses of such interactions between germline and somatic variants

can provide a new avenue for future individualized prognostic and therapeutic tools.

474   **Acknowledgments**

477 5R35GM127089-02 to C.V.). The results shown here are in whole or part based upon data

478 generated by the TCGA Research Network: https://www.cancer.gov/tcga.

480 **Conflict of Interest**

481 None of the authors have competing interests.
482

483 **Data Availability Statement**

484 Data sharing is not applicable to this article as no new variant data were created in this

485 study.

487 **Ethical Compliance**

488 This study describes re-analysis of publicly available variant data and no new prospective or

489 retrospective study was conducted.

492 **Tables**

Cancer Type	10-fold CV Significance	Permutation Significance	Number of Signatures	Average Consensus
BLCA	0.016	0.052	64	70.5%
BRCA	0.001	0.004	25	80.0%
ESCA	0.024	0.051	5	78.0%
GBM	<0.001	<0.001	12	75.0%
KIRC	0.009	0.023	3	70.0%
LGG	<0.001	<0.001	20	82.0%
OV	0.043	0.071	11	63.6%
UCEC	0.008	0.025	107	75.1%

495 **Table 1**. The results of cross-validation and permutation-based significance test for

496 overfitting of prognostic signatures in eight validated cancer types. Number of prognostic

497 signatures (including PIU, LU and NCU) with mutations occurring in at least three patients

498 along with their average consensus in cross validation folds. Consensus for each prognostic

499 unit is the number of times it being identified as prognostic unit in cross validation folds.

Figures

Figure 1. GPD mapping scheme and mapping results of somatic mutations across 33 cancers. (**A**) GPD's protein information units (PIUs) consist of domains and windows surrounding protein modification sites (+/- 5 amino acids around a PTM site) in proteins (yellow); Regions of the protein sequence that are not covered by a domain and a PTM site-containing window are called linker units (LUs) (blue). All sequence regions that do not encode the final protein product are named non-protein coding units (NCUs). Mutations (red triangles) are mapped to the GPD units defined above. GPD's unit-level data are per unit counts of mutations. Locus-level and gene-level counts are also illustrated for comparison. (**B**) Histograms of the mutation-carrying loci, genes, PIUs, LUs and NCUs in 10,793 subjects (x-axis) across all TCGA cancers. For example, a majority of mutations occurring at a specific locus have frequency of one or two individuals. By contrast, genes with any somatic mutations have a median count of 64. (**C**) Percentage of mutations mapped to one of the three types of sequences units across all genes. (**D**) Percentage of patients carrying mutations in the 50 most frequently mapped PIUs across 33 cancers. Stars indicate the PIUs mentioned in the main text. (**E**) Mutation frequency (logarithm base 2) of each PIU and LU among patients in TCGA Pan-Cancer cohort and MSK-IMPACT cohort.

Figure 2. Kaplan-Meier plot for eight cancer types with validated prognostic sigantures of sequence units. Predictive index (PI) was derived for each patient in the test group in each fold of the 10-fold cross validation. Patients who have PIs greater than the median in training group are clustered in the high-risk group (red), those with lower than the median PI are in the low-risk group (blue). P-values are derived from log-rank tests.

Figure 3. Association analysis between GPD somatic unit-level data and overall survival. (**A**) The total number and the proportion of significantly associated units (PIU, LU and NCU) for each cancer. (**B**) Significant PIUs across the seven cancers with prognostic signatures (no

significant PIU identified in ESCA). Height of bars indicates the fraction of patients with one

or more mutations on each PIU. Color of bars indicates the function each PIU is involved in.

Figure 4. Testing the interaction effect of somatic and germline variants in cancer prognosis.

(**A**) The number of statistically significant interaction effects in each cancer, among the

eligible units (see **Materials and Methods**). (**B**) Comparison of the hazard ratios between

patient groups with different somatic and germline variant levels. Patients are divided into

four groups based on whether they have zero or at least one somatic mutation in the unit

and whether they have fewer or more germline variants than the 75% quantile across all

patients.

References

Adzhubei IA, Schmidt S, Peshkin L, Ramensky VE, Gerasimova A, Bork P, Kondrashov AS, Sunyaev SR. 2010. A method and server for predicting damaging missense mutations. Nat Methods 7:248–249.

Bailey MH, Tokheim C, Porta-Pardo E, Sengupta S, Bertrand D, Weerasinghe A, Colaprico A, Wendl MC, Kim J, Reardon B, Kwok-Shing Ng P, Jeong KJ, et al. 2018. Comprehensive Characterization of Cancer Driver Genes and Mutations. Cell 174:1034–1035.

Bair E, Hastie T, Paul D, Tibshirani R. 2006. Prediction by Supervised Principal Components. Journal of the American Statistical Association 101, 119-137.

Bashashati A, Haffari G, Ding J, Ha G, Lui K, Rosner J, Huntsman DG, Caldas C, Aparicio SA, Shah SP. 2012. DriverNet: uncovering the impact of somatic driver mutations on transcriptional networks in cancer. Genome Biol 13:R124.

Candido-dos-Reis FJ, Song H, Goode EL, Cunningham JM, Fridley BL, Larson MC, Alsop K, Dicks E, Harrington P, Ramus SJ, Fazio A de, Mitchell G, et al. 2015. Germline mutation in BRCA1 or BRCA2 and ten-year survival for women diagnosed with epithelial ovarian cancer. Clin Cancer Res 21:652–657.

Carter H, Marty R, Hofree M, Gross AM, Jensen J, Fisch KM, Wu X, DeBoever C, Van Nostrand EL, Song Y, Wheeler E, Kreisberg JF, et al. 2017. Interaction Landscape of Inherited Polymorphisms with Somatic Events in Cancer. Cancer Discov 7:410–423.

Castro E, Goh C, Olmos D, Saunders E, Leongamornlert D, Tymrakiewicz M, Mahmud N, Dadaev T, Govindasami K, Guy M, Sawyer E, Wilkinson R, et al. 2013. Germline BRCA mutations are associated with higher risk of nodal involvement, distant metastasis, and poor survival outcomes in prostate cancer. J Clin Oncol 31:1748–1757.

Cingolani P, Platts A, Wang LL, Coon M, Nguyen T, Wang L, Land SJ, Lu X, Ruden DM. 2012. A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster strain w1118; iso-2; iso-3. Fly 6:80–92.

Dees ND, Zhang Q, Kandoth C, Wendl MC, Schierding W, Koboldt DC, Mooney TB, Callaway MB, Dooling D, Mardis ER, Wilson RK, Ding L. 2012. MuSiC: identifying mutational significance in cancer genomes. Genome Res 22:1589–1598.

Dellinger TH, Smith DD, Ouyang C, Warden CD, Williams JC, Han ES. 2016. L1CAM is an independent predictor of poor survival in endometrial cancer - An analysis of The Cancer Genome Atlas (TCGA). Gynecol Oncol 141:336–340.

Ding L, Bailey MH, Porta-Pardo E, Thorsson V, Colaprico A, Bertrand D, Gibbs DL, Weerasinghe A, Huang K-L, Tokheim C, Cortés-Ciriano I, Jayasinghe R, et al. 2018. Perspective on Oncogenic Processes at the End of the Beginning of Cancer Genomics. Cell 173:305–320.e10.

El-Gebali S, Mistry J, Bateman A, Eddy SR, Luciani A, Potter SC, Qureshi M, Richardson LJ, Salazar GA, Smart A, Sonnhammer ELL, Hirsh L, et al. 2019. The Pfam protein families database in 2019. Nucleic Acids Res 47:D427–D432.

Ellrott K, Bailey MH, Saksena G, Covington KR, Kandoth C, Stewart C, Hess J, Ma S, Chiotti KE, McLellan M, Sofia HJ, Hutter C, et al. 2018. Scalable Open Science Approach for

Mutation Calling of Tumor Exomes Using Multiple Genomic Pipelines. Cell Syst 6:271–
281.e7.

Fraser JA, Vojtesek B, Hupp TR. 2010. A novel p53 phosphorylation site within the MDM2
ubiquitination signal: I. phosphorylation at SER269 in vivo is linked to inactivation of p53
function. J Biol Chem 285:37762–37772.

Gentles AJ, Newman AM, Liu CL, Bratman SV, Feng W, Kim D, Nair VS, Xu Y, Khuong A,
Hoang CD, Diehn M, West RB, et al. 2015. The prognostic landscape of genes and
infiltrating immune cells across human cancers. Nat Med 21:938–945.

Hofree M, Shen JP, Carter H, Gross A, Ideker T. 2013. Network-based stratification of tumor
mutations. Nat Methods 10:1108–1115.

Hornbeck PV, Zhang B, Murray B, Kornhauser JM, Latham V, Skrzypek E. 2015.
PhosphoSitePlus, 2014: mutations, PTMs and recalibrations. Nucleic Acids Res 43:D512–
20.

Hristov BH, Singh M. 2017. Network-Based Coverage of Mutational Profiles Reveals Cancer
Genes. Cell Systems 5, 221–229.e4.
596
Huang K-L, Mashl RJ, Wu Y, Ritter DI, Wang J, Oh C, Paczkowska M, Reynolds S,
Wyczalkowski MA, Oak N, Scott AD, Krassowski M, et al. 2018. Pathogenic Germline
Variants in 10,389 Adult Cancers. Cell 173:355–370.e14.
Jeggo PA, Pearl LH, Carr AM. 2016.

Kiiski JI, Fagerholm R, Tervasmäki A, Pelttari LM, Khan S, Jamshidi M, Mantere T, Pylkäs K,
Bartek J, Bartkova J, Mannermaa A, Tengström M, et al. 2016. FANCM c.5101C>T mutation
associates with breast cancer survival and treatment outcome. Int J Cancer 139:2760–2770.

Kumar P, Henikoff S, Ng PC. 2009. Predicting the effects of coding non-synonymous
variants on protein function using the SIFT algorithm. Nat Protoc 4:1073–1081.

Lawrence MS, Stojanov P, Polak P, Kryukov GV, Cibulskis K, Sivachenko A, Carter SL,
Stewart C, Mermel CH, Roberts SA, Kiezun A, Hammerman PS, et al. 2013. Mutational
heterogeneity in cancer and the search for new cancer-associated genes. Nature 499:214–
218.

Le Morvan M, Zinovyev A, Vert J-P. 2017. NetNorM: Capturing cancer-relevant information
in somatic exome mutation data with gene networks for cancer stratification and prognosis.
PLoS Comput Biol 13:e1005573.

Liu J, Lichtenberg T, Hoadley KA, Poisson LM, Lazar AJ, Cherniack AD, Kovatich AJ, Benz
CC, Levine DA, Lee AV, Omberg L, Wolf DM, et al. 2018. An Integrated TCGA Pan-Cancer
Clinical Data Resource to Drive High-Quality Survival Outcome Analytics. Cell 173:400–
416.e11.

Lu VM, McDonald KL. 2018. Isocitrate dehydrogenase 1 mutation subtypes at site 132 and
their translational potential in glioma. CNS Oncol 7:41–50.

Martincorena I, Campbell PJ. 2015. Somatic mutation in cancer and normal cells. Science,
349(6255), 1483-1489.

McGranahan N, Swanton C. 2015. Biological and therapeutic impact of intratumor
heterogeneity in cancer evolution. Cancer Cell 27:15–26.

Olivier M. 2006. The Clinical Value of Somatic TP53 Gene Mutations in 1,794 Patients with
Breast Cancer. Clinical Cancer Research 12, 1157-1167.

Paull EO, Carlin DE, Niepel M, Sorger PK, Haussler D, Stuart JM. 2013. Discovering causal
pathways linking genomic events to transcriptional states using Tied Diffusion Through
Interacting Events (TieDIE). Bioinformatics 29:2757–2764.

Pickrell J, Berisa T, Segurel L, Tung JY, Hinds D. 2016. Detection and interpretation of
shared genetic influences on 42 human trails. Nat Genetics, 48(7), 709.

Porta-Pardo E, Godzik A. 2014. e-Driver: a novel method to identify protein regions driving
cancer. Bioinformatics 30:3109–3114.

Reimand J, Bader GD. 2013. Systematic analysis of somatic mutations in phosphorylation
signaling predicts novel cancer drivers. Mol Syst Biol 9:637.

Schoenfled, D. 1982. Partial residuals for the proportional hazards regression model.
Biometrika 69(1), 239-241.

Shihab HA, Gough J, Cooper DN, Day INM, Gaunt TR. 2013. Predicting the functional
consequences of cancer-associated amino acid substitutions. Bioinformatics 29:1504–1510.

Tamborero D, Gonzalez-Perez A, Lopez-Bigas N. 2013. OncodriveCLUST: exploiting the
positional clustering of somatic mutations to identify cancer genes. Bioinformatics 29:2238–
2244.

Vaske CJ, Benz SC, Sanborn JZ, Earl D, Szeto C, Zhu J, Haussler D, Stuart JM. 2010.
Inference of patient-specific pathway activities from multi-dimensional cancer genomics data
using PARADIGM. Bioinformatics 26:i237–45.

Wong WC, Kim D, Carter H, Diekhans M, Ryan MC, Karchin R. 2011. CHASM and SNVBox:
toolkit for detecting biologically important single nucleotide mutations in cancer.
Bioinformatics 27:2147–2148.

Xing M. 2005. BRAF mutation in thyroid cancer. Endocrine Related Cancer 12:245–262.

Zehir A, Benayed R, Shah RH, Syed A, Middha S, Kim HR, Srinivasan P, Gao J,
Chakravarty D, Devlin SM, Hellmann MD, Barron DA, et al. 2017. Mutational landscape of
metastatic cancer revealed from prospective clinical sequencing of 10,000 patients. Nat Med
23:703–713.

Zhong X, Yang H, Zhao S, Shyr Y, Li B. 2015. Network-based stratification analysis of 13
major cancer types using mutations in panels of cancer genes. BMC Genomics 16 Suppl
7:S7.