Ventral Pallidum is Essential for Cocaine Reinstatement After Voluntary Abstinence

[1]Mitchell R. Farrell*, B.A., [1]Christina M. Ruiz*, M.S., [1]Erik Castillo, B.S., [2]Lauren Faget, Ph.D., [1]Christine Khanbijian,

B.S., [1]Siyu Liu, B.S., [1]Hannah Schoch, Ph.D., [1]Gerardo Rojas, B.S., [2,3]Thomas S. Hnasko, Ph.D., [1]Stephen V. Mahler,

Ph.D.

Affiliations

[1]Department of Neurobiology & Behavior
University of California, Irvine
1203 McGaugh Hall
Irvine, CA 92697

[2]Department of Neurosciences
University of California, San Diego
9500 Gilman Drive
La Jolla, CA 92093
14
15   **[3]VASDHS Research Service**
16   **3350 La Jolla Village Dr**
17   **San Diego, CA 92161**
18

Corresponding Author: Stephen V. Mahler, Ph.D., Department of Neurobiology & Behavior University of

California, Irvine, 1203 McGaugh Hall, Irvine, CA 92697, (O) (949) 824-6128, (Fax) (949) 824-2447

Abstract

Addiction is a chronic relapsing disorder, and during recovery many people experience several relapse events as

they attempt to voluntarily abstain from drug. New preclinical relapse models have emerged which capture this

common human experience of relapse after voluntary abstinence, and mounting evidence indicates that

reinstatement of drug seeking after voluntary abstinence recruits neural circuits distinct from reinstatement

following experimenter-imposed abstinence, or abstinence due to extinction training. Ventral pallidum (VP), a key

limbic node involved in drug seeking, has well-established roles in conventional reinstatement models tested

following extinction training, but it is unclear whether this region also participates in more translationally-relevant

models of relapse. Here we show that chemogenetic inhibition of VP neurons strongly attenuates cocaine-,

context-, and cue-induced reinstatement tested after voluntary, punishment-induced abstinence. This effect was

strongest in the most compulsive, punishment-resistant rats, and reinstatement was associated with neural

activity in anatomically-defined VP subregions. VP inhibition also attenuated the propensity of rats to display

'hesitations,' a risk assessment behavior seen during punished drug taking that is likely due to concurrent

approach and avoidance motivations. These results indicate that VP, unlike other connected limbic brain regions,

is essential for reinstatement of drug seeking after voluntary abstinence. Since VP inhibition effects were

strongest in the most compulsively cocaine-seeking individuals, this could indicate that VP plays a particularly

important role in the most pathological, addiction-like behavior, making it an attractive target for future

therapeutic interventions.

43   **Introduction**

Addiction is characterized by persistent drug use despite negative consequences, and a lasting

vulnerability to relapse after protracted periods of abstinence [1-3]. Typically, human addicts eventually recognize

the negative consequences of their behavior, and choose to cease using drugs—a decision they usually renege

upon when tempted by drug cues, small doses of drug, or stressors [4]. In rodent relapse models, reinstatement

of seeking is triggered by analogous stimuli, usually following a period of imposed abstinence from drug

(incubation), or explicit extinction training. Recently, voluntary abstinence-based rodent models have emerged,

capturing the fact that in many cases addicted people choose to stop using drugs due to mounting negative life

consequences, rather than due to extinction training or external forces [5-9]. This is important because in rodents,

the neural substrates underlying reinstatement differ based upon how abstinence was achieved, be it

experimenter-imposed, through extinction training, or through voluntary cessation due to punishment or

availability of more attractive alternative reinforcers [10-15]. If the brain substrates of human relapse similarly

depend upon why a person stopped using drugs, then considering these factors in preclinical models will be

essential for developing effective interventions to treat addiction.

A hallmark of addiction is an inability to limit drug intake in the face of negative life consequences. This

can be modeled in rodents by training them to self-administer drugs, then introducing consequences to continued

use, such as co-delivered footshock [5, 6, 16-20]. As in humans, most rodents readily suppress their drug intake

when negative outcomes begin to result from their drug use. However, a subset of rodents show punishment-

resistant drug intake [17, 21-23], similar to the proportion of humans who use drugs that ultimately become

addicted [24]. Punishment-resistant animals also exhibit the most robust reinstatement of cocaine and

methamphetamine seeking [17, 25], suggesting that compulsive use and liability to relapse involve common

underlying neural mechanisms. Indeed, the circuitry underlying compulsive cocaine intake overlaps with the

limbic substrates of reinstatement behavior, at least when tested following extinction training [23, 26-29].

One brain region that has emerged as being crucial for motivated behavior is the ventral pallidum (VP),

the main efferent target of nucleus accumbens [30-35]. VP is thought to help translate motivation into action [36-

39], and accordingly, VP neural activity encodes reward motivation in rodents, monkeys, and humans [40-42],

including for cocaine [43]. VP is also required for seeking of several abused drugs [44-49], and for cocaine

reinstatement triggered by cues, stress, or cocaine following extinction training [47, 50, 51]. Notably, VP is a

heterogeneous structure, with functionally and anatomically distinct dorsolateral/ventromedial, and

rostral/caudal subregions that mediate distinct aspects of reward seeking, including cocaine reinstatement [43,

47, 52-59]. Given these results, and recent findings that VP contains phenotypically distinct populations of reward-

and aversion-related neurons [30-32, 60-62], its role in drug seeking under translationally-relevant mixed

motivation circumstances was of interest to us.

Here we explore effects of transiently and reversibly inhibiting VP neurons of punishment-resistant or

punishment-sensitive rats with designer receptors (DREADDs), determining effects on punished cocaine seeking,

context, discrete cue, and primed reinstatement after voluntary abstinence, as well as on cocaine-induced

locomotion. We also assessed reinstatement-related Fos in VP subregions. These studies shed light on the

functions of this essential, but understudied nucleus within cocaine addiction-related neural circuits.

Methods

Subjects. Male (*n*=50) and female (*n*=36) Long-Evans rats (220-250g at the start of experiments) were bred at the

University of California Irvine or obtained from Envigo, and were pair housed on a 12hr reverse light/dark cycle

with *ad libitum* food and water for all experiments. All training and testing was conducted in the dark period.

Procedures were approved by the UCI Institutional Animal Care and Use Committee, and are in accordance with

the NIH Guide for the Care and Use of Laboratory Animals [63].

Surgery. Animals were anesthetized with ketamine (56.5mg/kg), xylazine (8.7mg/kg), and the non-opioid

analgesic meloxicam (1.0mg/kg), and implanted with indwelling jugular catheters exiting the dorsal back. In the

same surgery, they also received bilateral viral vector injections (250-300nL) into VP with pressure injections using

a Picospritzer and glass micropipette. See **Figure 1** for schematic of procedures.

Viral Constructs. To transduce VP neurons with hM4Di inhibitory DREADDs, we used a human synapsin (hSyn)

promoter-driven AAV with mCitrine (*n*=44; U North Carolina vector core: *AAV2-hSyn-HA-hM4D(Gi)-IRES-mCitrine)*

or mCherry (*n*=16; Addgene: *AAV2-hSyn-hM4D(Gi)-mCherry*) reporter. To control for non-specific impact of viral

transduction and clozapine-N-oxide (CNO) in the absence of DREADDs, an eGFP-only reporter without DREADDs

(*n*=7; Addgene: *AAV2-hSyn-eGFP),* was employed in a group of control rats [64-66].

Figure 1. Schematic of experimental timeline. Following DREADD or control AAV injection, rats underwent cocaine self-administration, punishment training, followed by reinstatement and cocaine-induce locomotor testing. A final reinstatement test preceded sacrifice for neuronal activity (Fos) analysis.

Anatomical Analysis of DREADD Expression. hM4Di DREADD/reporter expression was visualized with

immunofluorescent amplification, with Substance P co-stain allowing demarcation of VP borders (see **Table S1** for

detailed list of antibodies). Rats with at least 40% of VP volume expressing DREADDs/reporter, and at least 40% of

virus expression localized within VP borders were considered hits (*n*=46). Rats with more than 60% of DREADD

expression localized outside VP (quantified blind to group and behavioral results) were considered misses (*n*=13).

Since rats with extra-VP DREADD expression did not behaviorally differ from fluorophore-only rats on

reinstatement tests (no main effect of group or CNO treatment on reinstatement types, *Fs*<1.29, *ps*>0.27; **Fig. S1**),

they were combined into a single control group (*n*=20) for subsequent analyses of CNO effects in the absence of

VP DREADDs.

RNAscope Analysis of DREADD Expression in VP Neurons. PFA-fixed brains were serially cut (16μm) on a cryostat

and mounted directly onto glass slides. Sections were stored at −80°C until processing for RNAscope Multiplex

Fluorescent assay (Advanced Cell Diagnostics). Briefly, sections were warmed on a hot plate for 30 minutes at

60°C then boiled at 100° for 6 min in target retrieval solution. Sections were then dehydrated in 100% ethanol and

treated with protease (pretreatment reagents, cat. No. 322380). RNA hybridization probes included antisense

probes against rat *gad1* (316401-C1) and *slc17a6* (317011-C3), respectively labeled with alexa488 and atto647

fluorophores. Slides were then incubated with rabbit anti-DsRed primary antibodies (632496, Clontech) and

donkey anti-rabbit AlexaFluor 594 secondary antibodies (711-585-152, Jackson ImmunoResearch), counterstained

with DAPI, and coverslipped using Fluoromount-G mounting medium. Images for cell counting were taken at 63x

(1.4 NA) magnification using a Zeiss AxioObserver Z1 widefield Epifluorescence microscope with a Zeiss ApoTome

2.0 for structured illumination and Zen Blue software. An average of 186 +/- 11 cells positive for AAV-hSyn-

hM4D(Gi)-mCherry were counted per brain (*n*=3 rats).

Drugs. Cocaine HCl (NIDA) was dissolved in 0.9% saline. Cocaine was available for self-administration at

0.2mg/50μL infusion for male rats, and 0.15mg/50μL infusion for female rats [67, 68]. Cocaine (10mg/kg) was

used for primed reinstatement and locomotion testing. CNO was dissolved in a vehicle of 5% DMSO in 0.9% saline,

and injected i.p. at 10mg/kg 30min prior to behavioral testing.

Behavioral Testing Apparatus. Self-administration training and testing took place in Med Associates operant

chambers within sound-attenuating boxes, equipped with two retractable levers with white lights above them,

and a tone generator. Cocaine-induced locomotion testing was conducted in 43×43×30.5cm Med Associates

locomotor testing chambers.

Self-Administration Training in Safe Context. We employed a punishment-induced abstinence/reinstatement

protocol modeled after previous reports [5, 20]. Initial self-administration occurred in a 'safe context,' signaled by

presence of a white or red house light, peppermint or orange scent, and plain or polka dot pattern walls

(randomly assigned). Intravenous cocaine was administered via a pump located outside the sound-attenuating

box. Rats received five daily 2hr sessions of fixed-ratio 1 (FR1) training where an active lever press delivered a

3.6sec cocaine infusion, and concurrent stimulus light + 2.9-kHz tone. A 20sec timeout period (signaled by

dimming of the house light) followed each infusion/cue presentation, during which additional lever presses did

not yield cocaine delivery. Pressing on an inactive lever was recorded but had no consequences. Following FR1

training to criterion (>10 infusions), rats then completed 3 days of variable-interval 5 schedule (VI5), on which an

active lever press initiated a timer with an average duration of 5sec, and another press after that interval

delivered a cocaine infusion with the light+tone cue. The VI schedule was increased to VI15 for the next 3 days,

then VI30 for an additional 3-6 days until rats had stable performance (**Fig. S2**).

Punishment Context Testing and Training. Following safe context self-administration training, rats (*n*=35) began

punishment training in a distinct chamber, with the opposite context used for safe context training. Rats

continued on a VI30 schedule, but 50% of cocaine infusions/cues were accompanied by a 0.3mA foot shock

(0.5sec). To test effects of inhibiting VP during punished cocaine intake, a subset of animals were injected with

CNO (*n*=22) or vehicle (*n*=13) prior to each of two daily shock punishment (0.3 mA) training sessions. In a

crossover design, these rats were administered the opposite treatment (vehicle/CNO) prior to a third punished

intake session 48hrs later, then a fourth punished cocaine intake training session with no vehicle or CNO injection

24hrs later. Another group of rats (*n*=31) received no injections during punishment context training. After 3-4 days

of shock training at 0.30mA, shock increased by 0.15mA every 2 training days, up to 0.75mA, until voluntary

abstinence criterion was met in all rats (<5 active lever presses for 2 consecutive days). Sensitivity to punishment

was determined in two ways. A suppression ratio (infusions on day 1 punishment/infusions on last day

unpunished; [17, 69]) was calculated as a measure of initial punishment sensitivity, with high ratios reflecting

relative insensitivity to shock-suppression of intake. Rats were also coded based on the maximum level of shock

they tolerated during punishment training, before meeting abstinence criterion.

Measuring Mixed Motivations During Punished Cocaine Intake: "Hesitation" Behavior. During punished cocaine

intake training sessions, rats exhibited a species-typical risk assessment behavior [70, 71] we term 'hesitation

behavior,' in which they stretch their trunk and extend their forepaw towards the active or inactive lever, but

rapidly retract it without completing the press to deliver cocaine and probabilistic shock. Hesitations directed at

the active and inactive levers were quantified using video analysis by a blinded observer on the final day of safe

context self-administration, and the first day of punishment context self-administration in CNO- and vehicle-

treated rats.

Reinstatement Tests. A series of 2hr reinstatement tests commenced 48hrs after rats met abstinence criterion,

with 48hrs elapsing between each test. Reinstatement tests occurred in: the safe context with response

contingent cues (*n*=66; vehicle and CNO administered on separate days in counterbalanced order), the safe

context without cues (vehicle/CNO, *n*=31), the safe context with no cues immediately after a cocaine priming

injection (10mg/kg; vehicle/CNO, *n*=38), the punishment context with cues (vehicle/CNO, *n*=35), and the

punishment context without cues (vehicle only, *n*=24). For reinstatement tests with cues, active lever presses

yielded 3.6sec cocaine-associated cue presentations (with no cocaine or shock), delivered on a VI-30 schedule,

and followed by a 15sec timeout period. For tests without discrete cues, lever presses were inconsequential, but

were recorded.

Cocaine Induced-Locomotion. Following reinstatement tests, a subset of rats (*n*=51) were habituated to a

locomotor testing chamber for 2 consecutive days, followed by two 2hr locomotor tests, 48hrs apart. Next, rats

were immediately placed in the chamber for 30min after vehicle/CNO, injected with cocaine, and returned to the

chamber for 90min. Horizontal locomotor activity and vertical rearing were recorded via infrared beam breaks.

Reinstatement-Related Fos. To examine VP neuronal activity during reinstatement, rats underwent a final drug-

free 2hr reinstatement test, 48hr after their last vehicle/CNO reinstatement test. They were tested in one of the

following reinstatement types: the safe context with response contingent cues (*n*=15), the safe context without

cues (*n*=16), the punishment context with cues (*n*=13), or no reinstatement (removed directly from their home

cage after equivalent self-administration/reinstatement training, *n*=7). After the 2hr test, rats were returned to

their home cages for 1hr, then perfused with saline (0.9%) and paraformaldehyde (4%), and brains sectioned

(40μm) following cryoprotection in 20% sucrose azide.

Fos Quantification. To allow Fos quantification within anatomically-defined VP subregions, we stained for Fos +

substance P to define VP borders. Ventromedial, ventrolateral, and dorsolateral subregions of substance P-

defined VP were delineated with reference to adjacent sections stained for substance P and neurotensin, defining

ventromedial/dorsolateral VP [54, 72] (**Table S1**). Images of VP were taken at 5x magnification, and one

section/animal was quantified bilaterally in rostral VP (+0.12 to +0.60 mm relative to Bregma), and another in

caudal VP (-0.48 to -0.24 mm; [73]). Fos+ neurons were identified using the Stereoinvestigator (Microbrightfield)

particle counter tool with thresholding parameters incorporating particle size (average size 100μm^2), minimum

distance between nuclei (150μm), and color relative to background. Fos density (Fos/mm^2) was computed for

each VP subregion on each slice (average of both hemispheres) of each rat. All structure delineation and

quantification was done blind to experimental conditions, and imaging/analysis settings were consistent across

animals.

Statistical Analyses. Effects of punishment on self-administration were examined with repeated measures

ANOVAs, including day (1 unpunished, 3 punished days) and behavioral output (active lever, inactive lever,

infusions) factors. Punishment sensitive versus resistant groups were compared on reinstatement with a two-way

ANOVA with punishment sensitivity group and reinstatement type as factors. Pearson correlation was used for

assessing relationships between suppression ratio and reinstatement behavior. Effects of punishment sensitivity

group on unpunished cocaine intake and cocaine-induced locomotion were examined with unpaired *t*-tests.

Effects of CNO in control and VP-hM4Di rats on hesitation behavior were computed with one-way ANOVA. Effects

of CNO on each reinstatement type in VP hM4Di-expressing and control rats were examined using separate

repeated measures ANOVAs with drug (vehicle/CNO) and lever (active/inactive) factors. Effects of VP inhibition on

reinstatement in punishment resistant and punishment sensitive rats were computed as change from vehicle day

behavior (CNO-vehicle), and compared with unpaired *t*-test. Separate one-way ANOVAs compared behavioral

groups on Fos in each VP subregion. Effects of rostrocaudal VP location on Fos was examined with a two-way

ANOVA with rostral/caudal site, and reinstatement type factors. Separate two-way ANOVAs were used to

compare CNO effects on cocaine-induced horizontal distance and rearing in control and VP-hM4Di rats. Tukey and

Bonferroni corrected *t*-tests were used for posthoc comparisons as appropriate.

206   **Results**

207   **Unpunished Self-Administration**. Rats readily discriminated between the inactive and active lever (Lever: $F_{(1,}$

208    $_{130)}$=55.3, *p*<0.0001), and daily cocaine intake was stable by the final 3 days of training ($F_{(2, 130)}$=0.87, *p*=0.42; **Fig.**

209   **S2**). Male and female rats did not differ in active lever presses or sex-adjusted cocaine doses self-administered

Figure 2. Punishment-resistant rats are more prone than punishment sensitive rats to cue, but not cocaine primed reinstatement. A) Probabilistic footshock reduces active lever pressing across all rats, while increasing the number of inactive presses. SA = self-administration. **B)** Individual variation in punishment sensitivity. Rats that reached at least 0.60mA footshock were considered punishment resistant (red). Punishment sensitive rats (gray) stopped taking cocaine at <0.60mA footshock intensities. **C)** Distribution of punishment resistance. Most rats cease cocaine intake at low shock levels (Punishment Sensitive: gray): 53/66 rats, 80.3%), but a subset reached the highest shock levels (Punishment Resistant: red): 13/66 rats, 19.7%). **D)** Punishment resistant rats reinstated more in the safe context with cues relative to punishment sensitive rats. This effect was specific to this reinstatement condition, and punishment sensitivity did not relate to context-only or cocaine primed reinstatment. **E)** Punishment effects on cocaine taking as quantified with suppression ratio [16,67] correlated with the degree of reinstatement in the safe context with cues. **F-G)** Punishment resistant rats were no different than sensitive rats on cocaine intake (F) or cocaine-induced locomotion (G). *** p<0.001. Panel E, squares = vehicle injection, triangles = CNO injection, circles = no injection.

during the last 3d of training (no main effect of sex (lever: $F_{(1, 64)}$=1.8, p=0.19, infusions: $F_{(1, 64)}$=0.29, p=0.59) or day

X sex interaction (Lever: $F_{(2, 128)}$=1.0, p=0.37, infusion: $F_{(2, 128)}$=0.48, p=0.62).

Individual Differences in Cocaine Seeking under Punishment. As expected, cocaine-coincident shock (50% of

infusions) in the punishment context suppressed cocaine self-administration overall (Day: $F_{(3, 585)}$=30.1, p<0.0001,

Fig. 2A). Most rats (80.3%; n=53) reached suppression criterion at the two lowest shock intensities (0.30-0.45mA:

'punishment sensitive' rats), but a subset of rats (19.7%, n=13) persisted in responding up to higher shock

intensities (0.60-0.75mA: 'punishment resistant' rats; **Fig. 2B-C**). In addition, punishment resistant rats had higher

suppression ratios (infusions on the first day in the punishment context/infusions on the last day in the safe

context [17, 69]; mean ± SEM = 0.48 ± 0.09) than punishment sensitive rats (mean ± SEM = 0.25 ± 0.02; t_{64}=4.4,

p<0.0001). Notably, of the 13 punishment resistant rats in this study, 8 were female (26.7% of tested females),

while 5 were male (13.9% of tested males).

Punishment Resistant Rats Reinstated More. Punishment resistant rats, once they received shock intensities high

enough to suppress even their seeking, showed greater cue-induced reinstatement than punishment sensitive

rats. However, this was only true in the "safe," unpunished context, and not when response contingent cues were

delivered in the "punishment" context (punishment sensitivity x reinstatement type interaction: $F_{(3, 166)}$=2.91,

p=0.036; punishment resistant vs. sensitive in safe context with cues: t_{166}=4.63, p<0.0001; **Fig. 2D**). Punishment

suppression ratio also correlated with the magnitude of cue reinstatement in the safe context (r=0.24, p=0.048;

Fig. 2E), further supporting a relationship between shock resistance and reinstatement propensity.

Punishment resistance was unrelated to total prior cocaine infusions (punishment resistant vs. sensitive total

unpunished infusions: t_{64}=0.67, p=0.50; **Fig. 2F**), or to cocaine's locomotor stimulating or reinstating effects

(horizontal distance traveled: t_{49}=1.45, p=0.15; **Fig. 2G**; rearing: t_{49}=1.77, p=0.084; cocaine primed reinstatement:

t_{36}=0.83, p=0.41), indicating that punishment resistance and cue-induced reinstatement likely involve underlying

individual differences in addiction-like compulsive cocaine seeking, rather than sensitivity to cocaine's effects per

se.

DREADD Expression in VP Neuronal Populations. Robust hM4Di-DREADD expression was observed throughout

the rostrocaudal extent of VP in this study (**Fig. 3A-B**). Fluorescent *in situ* hybridization (RNAscope) revealed

colocalization of hM4Di expression with $gad1^+$ neurons (85.3 +/- 3.2 %), with a smaller percentage colocalizing

with $gad1^-$ neurons (14.7 +/- 3.2 %; **Fig. 3C-E**), consistent with unbiased transfection of all VP neurons, as GABA

neurons represent the predominant neuronal phenotype in VP [31]. Non-GABA neurons likely represent

intermingled glutamatergic and cholinergic cell-types [31, 54].

Figure 3. Inhibitory DREADD localization in VP. **A)** Immunofluorescent co-stain for hM4Di-mCherry (red) within Substance P-expressing VP borders (green). **B)** Virus expression for individual animals on rostrocaudal VP axis. **C)** Wide-field and **D)** high magnification images of fluorescent *in situ* hybridization for *gad1* (green) combined with immunofluorescence for hM4Di-mCherry (red). Co-positive hM4Di⁺ *gad1*⁺ cells labeled with green stars. **E)** 85.3 +/- 3.2 % of hM4Di⁺ cells are co-positive for *gad1*. 14.7 +/- 3.2 % of hM4Di⁺ cells are *gad1⁻* (n=3 rats, total of 559 hM4Di⁺ cells counted). VM = ventromedial, VL = ventrolateral, DL = dorsolateral. AC = anterior commissure. Scale bars = 500μm (B), 200μm (C), and 20μm (D).

CNO Effects on Punishment-Induced Suppression of Cocaine Intake in VP-hM4Di rats. On day 1 of punished

cocaine self-administration, CNO in VP-hM4Di rats modestly, but non-significantly, decreased the number of

active and inactive lever presses relative to control rats (Treatment: $F_{(1, 33)}$=3.41, *p*=0.073; **Fig. 4A**), with no

interaction of Treatment x Lever ($F_{(1, 33)}$=0.27, p=0.61). CNO had no effect on lever pressing on day 2, though rats

decreased their responding relative to day 1 (Day: $F_{(1, 33)}$=20.56, p<0.0001). When vehicle and CNO treatments

were reversed on punishment day 3, no further changes were observed (p>0.05).

Footshock+cocaine also increased "hesitation" behaviors targeted toward both the active and inactive

levers, relative to the low levels seen during unpunished self-administration in the same context ($F_{(2, 40)}$=7.93,

p=0.0013; self-administration vs. vehicle: p=0.0022; **Fig. 4B**). Hesitations were highest in rats with the most active

lever presses (correlation of hesitations and presses after both CNO; r=0.61, p=0.027; and vehicle; r=0.61,

p=0.016), suggesting that hesitations may be a sensitive measure of deliberation about pursuing the now

dangerous cocaine. Accordingly, CNO strongly suppressed hesitations, returning them to unpunished levels in VP-

hM4Di rats (p=0.0013; CNO vs. vehicle: p=0.0082; **Fig. 4B**).

VP DREADD Inhibition Suppresses Cocaine Reinstatement After Voluntary Abstinence. CNO in VP-hM4Di rats

robustly suppressed context only-, cue-, and cocaine-induced reinstatement in the safe context, but failed to do

so in control rats without VP DREADDs. In VP-hM4Di rats, CNO (compared to vehicle) reduced cue-induced active,

but not inactive lever pressing in the safe context (Drug x Lever interaction: $F_{(1, 45)}$=18.53, p<0.0001; **Fig. 4C**), and

also suppressed safe context pressing without response-contingent cues (Drug x Lever interaction: $F_{(1, 20)}$=4.31,

p=0.05; **Fig. 4C**). Similarly, cocaine primed reinstatement (no cues) in the safe context was also suppressed by CNO

in VP hM4Di rats (Drug x Lever interaction: $F_{(1, 23)}$=7.94, p=0.01; **Fig. 4C**). Although we previously showed that

rostral and caudal VP differentially mediate cue- and primed reinstatement [47], in these experiments our viral

infection spanned most of the rostrocaudal axis of VP. In contrast to the safe context, CNO in VP-hM4Di rats failed

to reduce cue-induced reinstatement in the punishment context (Drug x Lever interaction: $F_{(1, 21)}$=0.19, p=0.66;

Fig. 4C). In control rats without VP DREADDs, CNO had no effects on lever pressing in any reinstatement test (p

values>0.05; **Fig. 4D**), suggesting that CNO effects here, as previously shown [45, 47, 74, 75], were specific to VP

DREADD-expressing rats.

Figure 4. VP inhibition reduces reinstatement especially in punishment resistant rats. A) Top panel: Example picture of a standard lever press. Bottom panel: CNO in VP-hM4Di rats modestly reduces active and inactive lever pressing for cocaine under threat of punishment. Control = vehicle-injected rats and CNO-injected misses. **B)** Top panel: Hesitation behavior in which the rat stretches its trunk towards the lever and extends its paw without depressing the lever. Bottom panel: Active and inactive lever hesitations quantified during safe (green shading) and punished (red shading). CNO in VP-hM4Di rats reduced hesitations to relative to control rats. VP inhibition reduced hesitations to unpunished levels. Control = vehicle-injected rats and CNO-injected misses. **C)** Within subjects comparisons of reinstatement for VP-hM4Di rats in safe (green shading) and punishment (red shading) contexts. CNO in VP-hM4Di rats reduced reinstatement in the safe context with cues, without cues, and with cocaine and no cues, but not in the punishment context with cues. **D)** CNO in control rats did not affect reinstatement under any condition. Control = eGFP-only rats and rats with hM4Di expression primarily outside VP. **E)** CNO in VP-hM4Di punishment resistant rats (red bars) elicited a greater decrease in reinstatement relative to punishment sensitive rats (gray bars). Data presented as change from vehicle test baseline. $p<0.05*$, $p<0.01**$, $p<0.001***$.

VP Inhibition Suppressed Reinstatement the Most in Punishment-Resistant Rats. VP inhibition reduced safe

context cue-induced reinstatement more in punishment-resistant rats than punishment-sensitive rats (t_{44}=2.23,

p=0.031; **Fig. 4E**). This effect was specific to the safe context with cues, as there was no such effect on other

reinstatement types (t scores<1.26, ps>0.22). This finding suggests that VP plays an especially important role in

relapse after punishment-imposed abstinence for the individual rats showing the most addiction-like behavior.

VP Inhibition did not Affect Cocaine-Induced Locomotion. CNO failed to affect the locomotor-activating effects of

cocaine in either VP-hM4Di or control groups (Treatment: $F_{(1, 49)}$=0.63, p=0.43; treatment x group interaction: $F_{(1,}$

$_{49)}$=0.58, p=0.45; **Fig. S3A**), though it did reduce rearing behavior after cocaine in VP-hM4Di rats, but not controls

(treatment x group interaction: $F_{(1, 49)}$=10.24, p=0.0024; **Fig. S3B**), further suggesting specificity of these findings to

cocaine seeking in particular. Moreover, CNO did not differentially reduce horizontal locomotion or rearing

behavior in punishment-sensitive versus punishment-resistant VP-hM4Di rats (group x treatment interaction;

locomotion: $F_{(3, 93)}$=0.70, p=0.55; Rearing: $F_{(3, 93)}$=0.61, p=0.61).

VP Subregion Fos Recruited During Reinstatement. Relative to cocaine/shock-experienced rats sacrificed from

their homecages, VP subregions showed strong Fos activation during all tested reinstatement conditions ($F_{(3,}$

$_{47)}$=3.93, p=0.014; punishment+cues, p=0.013 ; safe+cues, p=0.019; safe+no cues, p=0.043). Ventromedial VP was

selectively activated (relative to home cage) by the punishment context with cues, but not by either safe context

reinstatement test ($F_{(3, 47)}$=2.67, p=0.05; punishment+cues: p=0.048; safe+cues: p=0.28; safe+no cues: p=0.09; **Fig.**

5A-C). In contrast, ventrolateral and dorsolateral VP were activated in all reinstatement conditions relative to

homecage controls (ventrolateral: $F_{(3, 47)}$=5.98, p=0.0015; safe+cues: p=0.0051; safe+no cues: p=0.0011;

punishment+cues: p=0.0049; **Fig. 5D**; dorsolateral**:** $F_{(3, 47)}$=4.63, p=0.006; home cage vs. safe+cues: p=0.0043;

safe+no cues: p=0.017; punishment+cues: p=0.021; **Fig. 5E**). We then examined rostral and caudal sections of VP,

given known rostrocaudal functional and anatomical differences [47, 54-56]. Overall, rostral VP had greater Fos

density than caudal VP ($F_{(3, 47)}$=4.8, p=0.0051), though this did significantly differ between reinstatement types (no

reinstatement type X rostrocaudal position interaction; $F_{(3, 47)}$=1.42, p=0.25; **Fig. 5F**).

Figure 5. VP subregion Fos expression after reinstatement. **A)** Representative image of substance P (VP borders) and neurotensin (ventromedial VP marker) immunofluorescent co-stain. **B)** Representative image of substance P and Fos co-stained for Fos quantification. **C-E)** Comparison of ventromedial **(C)**, ventrolateral **(D)**, and dorsolateral **(E)** Fos density across reinstatement conditions. Green shading = safe context, red shading = punishment context, gray shading = home cage controls. **F)** Rostrocaudal Fos density across reinstatement conditions. Rostral VP reached significant only in the punishment context with cues. $p < 0.05$*; VM = ventromedial, VL = ventrolateral, DL = dorsolateral. AC = anterior commissure. Scale bar = 1000μm.

Sex Differences. Few sex differences with regard to VP manipulations were detected. Male and female rats

exhibited comparable levels of punishment-induced suppression of cocaine self-administration (suppression ratio:

t_{64}=1.65, p=0.10). As previously reported, female rats exhibited more cocaine-induced locomotion than males

(Sex; $F_{(1, 31)}$=4.91, p=0.034, [76]), though CNO (relative to vehicle) in VP-hM4Di did not differentially impact

locomotion between sexes (Sex x Treatment interaction; $F_{(1, 31)}$=0.33, p=0.57) or reinstatement under any

condition (F<2.5, p>0.13). Despite enhanced cocaine-induced locomotion, no sex differences were detected for

cocaine-primed reinstatement (Sex: $F_{(1, 22)}$=2.06, p=0.17), suggesting that cocaine promotes non-specific

locomotor activation in females, without enhancing the incentive motivational properties of cocaine under these

conditions.

<div align="center">

Discussion

</div>

These findings point to a crucial role for VP in cocaine reinstatement following voluntary abstinence, a

translationally-relevant model of humans who quit drugs due to mounting negative life outcomes. Indeed, we

found that VP plays an especially important role in the most compulsive cocaine-seeking individuals, i.e., the ~20%

of rats that tolerated significant footshock punishment to continue taking cocaine. We also found robust

reinstatement-related activity in anatomically-defined VP subregions. Our results suggest that unlike connected

limbic nuclei, VP plays a critical role in reinstatement regardless of how abstinence was achieved or how relapse

was initiated, thereby placing it amongst the most essential nodes within the neural circuits of cocaine addiction.

Persistent drug use despite negative consequences, and long-lasting relapse propensity are cardinal

features of addiction in humans [77]. Though compulsive drug intake despite punishment is common in rodents

following extended drug access [22, 23, 69], some rats seem to transition to compulsive use even after short

access to cocaine. Here, we observed such a subset of compulsive rats, and found that these same animals were

also more prone to cue+context reinstatement after voluntary abstinence, similar to prior findings [17, 25].

Importantly, VP inhibition in these compulsive rats had a greater reinstatement-suppressing effect than in

punishment sensitive rats, suggesting that VP plays a particularly important role in those rats which pathologically

seek drug. VP inhibition only modestly reduced punished cocaine self-administration, but instead selectively

reduced hesitations to press the cocaine/shock and inactive levers, which we interpret as reflecting motivation to

pursue cocaine, tempered by motivation to avoid being shocked. These results highlight the sensitivity of this

novel assay of conflicting motivations during cocaine seeking, and the importance of careful ethological analysis of

complex drug-seeking behaviors during such neural circuit manipulation experiments.

Relapse is not a unitary phenomenon, since brain circuits underlying drug reinstatement depend on the

drug of choice, mode of abstinence, and relapse trigger [10-12, 26, 78-81]. This said, we show that even under

maximally human-relevant conditions VP is broadly implicated in reinstatement regardless of trigger or mode of

abstinence. In contrast, other VP-connected limbic regions seem to be engaged differentially during

reinstatement after different modes of abstinence. For example, inhibition of basolateral amygdala *decreases*

reinstatement after extinction training, whereas the same manipulation during reinstatement following

punishment-induced abstinence *increases* drug seeking [10]. These results are consistent with the idea that VP

serves as a 'final common pathway' of drug seeking [38, 39]. Therefore, VP holds promise as a potential

therapeutic target for suppressing relapse in humans, especially since a prior human fMRI report found that

activity in the vicinity of VP predicts relapse propensity [82].

Interestingly, in the unpunished safe context, VP inhibition attenuated reinstatement with or without

cues, and also after a cocaine priming injection, yet VP inhibition did not reduce cue reinstatement in the

punishment context. As expected, conditioned suppression of seeking was observed in this context relative to the

safe context, but response-contingent cues nonetheless supported some pressing, reducing the likelihood of a

floor effect (**Fig. 4**). We therefore speculate that VP promotes conditioned drug seeking in a context-gated

manner, consistent with prior reports that VP is necessary for context-induced reinstatement of alcohol seeking

[44, 45, 49, 83].

VP is heterogeneous, with rostrocaudally- and mediolaterally located subregions, and functionally distinct,

genetically-defined neuronal subpopulations [30-32, 54, 60, 83]. We observed broad recruitment of Fos in VP

subregions during cue reinstatement tests in both the safe and punishment context, and even in the safe context

in the absence of response-contingent cues. This homecage-relative Fos recruitment was most pronounced in

rostral, relative to caudal VP. These results suggest a global recruitment of numerous VP subregions during both

context- and cue-induced reinstatement, including in the punished context where global VP inhibition failed to

suppress cocaine seeking. One possible explanation for this puzzling pattern of effects is that functionally opposed

VP cells are engaged in the safe- and punished-contexts, such as the intermingled VP GABA and glutamate

neurons which drive appetitive and aversive behavior, respectively [30-32, 61, 62]. Our pan-neuronal

chemogenetic approach primarily targeted reward-related VP GABA neurons (~85%), consistent with observed

reinstatement suppression in the safe context, where aversion-related glutamate neurons would be less relevant.

We speculate that in the punishment context, glutamate and GABA neurons were recruited (explaining Fos

results), but inhibiting both concurrently with DREADDs suppressed motivation as well as aversion, resulting in a

null effect. More work is needed to parse the specific behavioral roles for VP subregions and neuronal

subpopulations in addiction-related behaviors.

The present report firmly establishes VP as an essential node in the neural circuits of translationally-

relevant cocaine reinstatement behavior, especially in the most compulsive, addicted-like animals. By better

understanding how addiction-relevant behaviors map onto defined neural circuits in the addicted brain, we may

reveal neural signatures that could facilitate diagnosis and treatment of addiction in a personalized manner. These

and other results suggest that VP plays a key role across relapse triggers and modes of abstinence, making it a

promising target for future interventions to treat addiction.

Funding and Disclosures. This work was supported by PHS grants: R00DA035251, F31DA048578, P50DA044118,

R25GM055246, T32NS045540-15, and R21MH118748. The authors announce no conflicts of interest.

Acknowledgements. We thank Yavin Shaham for providing MED-PC code for punishment-induced abstinence

protocols. We thank Iohanna Pagnoncelli and Stephanie Lenogue for their assistance with behavioral testing.

References

1. Hunt, W.A., L.W. Barnett, and L.G. Branch, *Relapse rates in addiction programs.* Journal of clinical psychology, 1971. **27**(4): p. 455-456.
2. Wikler, A., *Dynamics of drug dependence: Implications of a conditioning theory for research and treatment.* Archives of general psychiatry, 1973. **28**(5): p. 611-616.
3. O'Brien, C.P., et al., *Classical Conditioning in Drug-Dependent Humans a.* Annals of the New York Academy of Sciences, 1992. **654**(1): p. 400-415.
4. Shaham, Y., et al., *The reinstatement model of drug relapse: history, methodology and major findings.* Psychopharmacology, 2003. **168**(1-2): p. 3-20.
5. Marchant, N.J., et al., *Context-induced relapse to alcohol seeking after punishment in a rat model.* Biological psychiatry, 2013. **73**(3): p. 256-262.
6. Panlilio, L.V., E.B. Thorndike, and C.W. Schindler, *Reinstatement of punishment-suppressed opioid self-administration in rats: an alternative model of relapse to drug abuse.* Psychopharmacology, 2003. **168**(1-2): p. 229-235.
7. Marchant, N.J., X. Li, and Y. Shaham, *Recent developments in animal models of drug relapse.* Current opinion in neurobiology, 2013. **23**(4): p. 675-683.
8. Economidou, D., et al., *High impulsivity predicts relapse to cocaine-seeking after punishment-induced abstinence.* Biological psychiatry, 2009. **65**(10): p. 851-856.
9. Venniro, M., D. Caprioli, and Y. Shaham, *Animal models of drug relapse and craving: from drug priming-induced reinstatement to incubation of craving after voluntary abstinence.* Progress in brain research, 2016. **224**: p. 25-52.
10. Pelloux, Y., et al., *Opposite effects of basolateral amygdala inactivation on context-induced relapse to cocaine seeking after extinction versus punishment.* Journal of Neuroscience, 2018. **38**(1): p. 51-59.
11. Fuchs, R.A., R.K. Branham, and R.E. See, *Different neural substrates mediate cocaine seeking after abstinence versus extinction training: a critical role for the dorsolateral caudate–putamen.* Journal of Neuroscience, 2006. **26**(13): p. 3584-3588.
12. Marchant, N.J., et al., *Context-induced relapse after extinction versus punishment: similarities and differences.* Psychopharmacology, 2018: p. 1-10.
13. Venniro, M., et al., *The anterior insular cortex→ central amygdala glutamatergic pathway is critical to relapse after contingency management.* Neuron, 2017. **96**(2): p. 414-427.
14. Venniro, M., et al., *Volitional social interaction prevents drug addiction in rat models.* Nature neuroscience, 2018. **21**(11): p. 1520.
15. Golden, S.A., M. Jin, and Y. Shaham, *Animal models of (or for) aggression reward, addiction, and relapse: behavior and circuits.* Journal of neuroscience, 2019: p. 0151-19.
16. Belin, D., et al., *Pattern of intake and drug craving predict the development of cocaine addiction-like behavior in rats.* Biological psychiatry, 2009. **65**(10): p. 863-868.
17. Deroche-Gamonet, V., D. Belin, and P.V. Piazza, *Evidence for addiction-like behavior in the rat.* Science, 2004. **305**(5686): p. 1014-1017.
18. Vanderschuren, L.J.M.J., et al., *Punishment models of addictive behavior.* Current opinion in behavioral sciences, 2017. **13**: p. 77-84.
19. Smith, R.J. and L.S. Laiks, *Behavioral and neural mechanisms underlying habitual and compulsive drug seeking.* Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2018. **87**: p. 11-21.
20. Pelloux, Y., et al., *Context-induced relapse to cocaine seeking after punishment-imposed abstinence is associated with activation of cortical and subcortical brain regions.* Addiction biology, 2018. **23**(2): p. 699-712.
21. Everitt, B.J., et al., *Neural mechanisms underlying the vulnerability to develop compulsive drug-seeking habits and addiction.* Philosophical Transactions of the Royal Society of London B: Biological Sciences, 2008. **363**(1507): p. 3125-3135.

22.   Vanderschuren, L.J.M.J. and B.J. Everitt, *Drug seeking becomes compulsive after prolonged cocaine self-*
administration. Science, 2004. **305**(5686): p. 1017-1019.

23.   Chen, B.T., et al., *Rescuing cocaine-induced prefrontal cortex hypoactivity prevents compulsive cocaine*
seeking. Nature, 2013. **496**(7445): p. 359.

24.   Anthony, J.C., L.A. Warner, and R.C. Kessler, *Comparative epidemiology of dependence on tobacco,*
alcohol, controlled substances, and inhalants: basic findings from the National Comorbidity Survey.
Experimental and clinical psychopharmacology, 1994. **2**(3): p. 244.

25.   Torres, O.V., et al., *Compulsive methamphetamine taking under punishment is associated with greater*
cue-induced drug seeking in rats. Behavioural brain research, 2017. **326**: p. 265-271.

26.   Farrell, M.R., H. Schoch, and S.V. Mahler, *Modeling cocaine relapse in rodents: Behavioral considerations*
and circuit mechanisms. Prog Neuropsychopharmacol Biol Psychiatry, 2018. **87**(Pt A): p. 33-47.

27.   Bock, R., et al., *Strengthening the accumbal indirect pathway promotes resilience to compulsive cocaine*
use. Nature neuroscience, 2013. **16**(5): p. 632.

28.   Ersche, K.D., et al., *Abnormal structure of frontostriatal brain systems is associated with aspects of*
impulsivity and compulsivity in cocaine dependence. Brain, 2011. **134**(7): p. 2013-2024.

29.   Yager, L.M., et al., *Chemogenetic inhibition of direct pathway striatal neurons normalizes pathological,*
cue-induced reinstatement of drug-seeking in rats. Addiction biology, 2019. **24**(2): p. 251-264.

30.   Stephenson-Jones, M., et al., *Opposing contributions of GABAergic and glutamatergic ventral pallidal*
neurons to motivational behaviours. Available at SSRN 3367001, 2019.

31.   Faget, L., et al., *Opponent control of behavioral reinforcement by inhibitory and excitatory projections*
from the ventral pallidum. Nature communications, 2018. **9**(1): p. 849.

32.   Tooley, J., et al., *Glutamatergic ventral pallidal neurons modulate activity of the habenula–tegmental*
circuitry and constrain reward seeking. Biological psychiatry, 2018. **83**(12): p. 1012-1023.

33.   Saga, Y., et al., *Ventral pallidum encodes contextual information and controls aversive behaviors.* Cerebral
Cortex, 2016. **27**(4): p. 2528-2543.

34.   Swanson, L.W. and W.M. Cowan, *A note on the connections and development of the nucleus accumbens.*
Brain research, 1975. **92**(2): p. 324-330.

35.   Williams, D.J., A.R. Crossman, and P. Slater, *The efferent projections of the nucleus accumbens in the rat.*
Brain Research, 1977. **130**(2): p. 217-227.

36.   Heimer, L., R.D. Switzer, and G.W. Van Hoesen, *Ventral striatum and ventral pallidum: Components of the*
motor system? Trends in Neurosciences, 1982. **5**: p. 83-87.

37.   Mogenson, G.J., D.L. Jones, and C.Y. Yim, *From motivation to action: functional interface between the*
limbic system and the motor system. Progress in neurobiology, 1980. **14**(2-3): p. 69-97.

38.   Kalivas, P.W. and N.D. Volkow, *The neural basis of addiction: a pathology of motivation and choice.*
American Journal of Psychiatry, 2005. **162**(8): p. 1403-1413.

39.   Smith, K.S., et al., *Ventral pallidum roles in reward and motivation.* Behavioural brain research, 2009.
196(2): p. 155-167.

40.   Tachibana, Y. and O. Hikosaka, *The primate ventral pallidum encodes expected reward value and regulates*
motor action. Neuron, 2012. **76**(4): p. 826-837.

41.   Pessiglione, M., et al., *How the brain translates money into force: a neuroimaging study of subliminal*
motivation. Science, 2007. **316**(5826): p. 904-906.

42.   Richard, J.M., et al., *Ventral pallidum neurons encode incentive value and promote cue-elicited*
instrumental actions. Neuron, 2016. **90**(6): p. 1165-1173.

43.   Root, D.H., et al., *Differential roles of ventral pallidum subregions during cocaine self-administration*
behaviors. Journal of Comparative Neurology, 2013. **521**(3): p. 558-588.

44.   Perry, C.J. and G.P. McNally, *A role for the ventral pallidum in context-induced and primed reinstatement*
of alcohol seeking. European Journal of Neuroscience, 2013. **38**(5): p. 2762-2773.

45.   Prasad, A.A. and G.P. McNally, *Ventral pallidum output pathways in context-induced reinstatement of*
alcohol seeking. Journal of Neuroscience, 2016. **36**(46): p. 11716-11726.

46. Rogers, J.L., S. Ghee, and R.E. See, *The neural circuitry underlying reinstatement of heroin-seeking*
behavior in an animal model of relapse. Neuroscience, 2008. **151**(2): p. 579-88.
47. Mahler, S.V., et al., *Designer receptors show role for ventral pallidum input to ventral tegmental area in*
cocaine seeking. Nature neuroscience, 2014. **17**(4): p. 577.
48. Hubner, C.B. and G.F. Koob, *The ventral pallidum plays a role in mediating cocaine and heroin self-*
administration in the rat. Brain research, 1990. **508**(1): p. 20-29.
49. Prasad, A.A. and G.P. McNally, *Ventral Pallidum and Alcohol Addiction*, in *Neuroscience of Alcohol*. 2019,
Elsevier. p. 163-170.
50. McFarland, K. and P.W. Kalivas, *The circuitry mediating cocaine-induced reinstatement of drug-seeking*
behavior. Journal of Neuroscience, 2001. **21**(21): p. 8655-8663.
51. McFarland, K., et al., *Limbic and motor circuitry underlying footshock-induced reinstatement of cocaine-*
seeking behavior. Journal of Neuroscience, 2004. **24**(7): p. 1551-1560.
52. Churchill, L. and P.W. Kalivas, *A topographically organized gamma-aminobutyric acid projection from the*
ventral pallidum to the nucleus accumbens in the rat. Journal of Comparative Neurology, 1994. **345**(4): p.
579-595.
53. Groenewegen, H.J., H.W. Berendse, and S.N. Haber, *Organization of the output of the ventral*
striatopallidal system in the rat: ventral pallidal efferents. Neuroscience, 1993. **57**(1): p. 113-42.
54. Root, D.H., et al., *The ventral pallidum: Subregion-specific functional anatomy and roles in motivated*
behaviors. Progress in neurobiology, 2015. **130**: p. 29-70.
55. Smith, K.S. and K.C. Berridge, *Opioid limbic circuit for reward: interaction between hedonic hotspots of*
nucleus accumbens and ventral pallidum. Journal of neuroscience, 2007. **27**(7): p. 1594-1605.
56. Smith, K.S. and K.C. Berridge, *The ventral pallidum and hedonic reward: neurochemical maps of sucrose*
"liking" and food intake. Journal of neuroscience, 2005. **25**(38): p. 8637-8649.
57. Calder, A.J., et al., *Disgust sensitivity predicts the insula and pallidal response to pictures of disgusting*
foods. European Journal of Neuroscience, 2007. **25**(11): p. 3422-3428.
58. Beaver, J.D., et al., *Individual differences in reward drive predict neural responses to images of food.*
Journal of Neuroscience, 2006. **26**(19): p. 5160-5166.
59. Stefanik, M.T., et al., *Optogenetic evidence that pallidal projections, not nigral projections, from the*
nucleus accumbens core are necessary for reinstating cocaine seeking. Journal of Neuroscience, 2013.
33(34): p. 13654-13662.
60. Knowland, D., et al., *Distinct ventral pallidal neural populations mediate separate symptoms of depression.*
Cell, 2017. **170**(2): p. 284-297.
61. Wulff, A.B., et al., *Ventral pallidal modulation of aversion processing.* Brain research, 2018.
62. Stephenson-Jones, M., *Pallidal circuits for aversive motivation and learning.* Current Opinion in Behavioral
Sciences, 2019. **26**: p. 82-89.
63. National Research, C., *Guide for the care and use of laboratory animals*. 2010: National Academies Press.
64. Mahler, S.V. and G. Aston-Jones, *CNO Evil? Considerations for the use of DREADDs in behavioral*
neuroscience. Neuropsychopharmacology, 2018. **43**(5): p. 934.
65. MacLaren, D.A.A., et al., *Clozapine N-oxide administration produces behavioral effects in Long–Evans rats:*
implications for designing DREADD experiments. eneuro, 2016. **3**(5).
66. Gomez, J.L., et al., *Chemogenetics revealed: DREADD occupancy and activation via converted clozapine.*
Science, 2017. **357**(6350): p. 503-507.
67. Zhou, L., et al., *Fos expression induced by cocaine-conditioned cues in male and female rats.* Brain
Structure and Function, 2014. **219**(5): p. 1831-1840.
68. Kohtz, A.S. and G. Aston-Jones, *Cocaine seeking during initial abstinence is driven by noradrenergic and*
serotonergic signaling in hippocampus in a sex-dependent manner. Neuropsychopharmacology, 2017.
42(2): p. 408.
69. Pelloux, Y., B.J. Everitt, and A. Dickinson, *Compulsive drug seeking by rats under punishment: effects of*
drug taking history. Psychopharmacology, 2007. **194**(1): p. 127-137.

70. Blanchard, R.J., et al., *The effects of ethanol and diazepam on reactions to predatory odors.* Pharmacology
Biochemistry and Behavior, 1990. **35**(4): p. 775-780.
71. Blanchard, D.C., et al., *Risk assessment as an evolved threat detection and analysis process.* Neuroscience
& Biobehavioral Reviews, 2011. **35**(4): p. 991-998.
72. Zahm, D.S. and L. Heimer, *Ventral striatopallidal parts of the basal ganglia in the rat: I. Neurochemical*
compartmentation as reflected by the distributions of neurotensin and substance P immunoreactivity.
Journal of Comparative Neurology, 1988. **272**(4): p. 516-535.
73. Paxinos, G. and C. Watson, *The rat brain in stereotaxic coordinates: hard cover edition*. 2006: Elsevier.
74. Chang, S.E., et al., *Chemogenetic manipulation of ventral pallidal neurons impairs acquisition of sign-*
tracking in rats. European Journal of Neuroscience, 2015. **42**(12): p. 3105-3116.
75. Mahler, S.V., et al., *Chemogenetic Manipulations of Ventral Tegmental Area Dopamine Neurons Reveal*
Multifaceted Roles in Cocaine Abuse. Journal of Neuroscience, 2019. **39**(3): p. 503-518.
76. Van Haaren, F. and M.E. Meyer, *Sex differences in locomotor activity after acute and chronic cocaine*
administration. Pharmacology Biochemistry and Behavior, 1991. **39**(4): p. 923-927.
77. American Psychiatric, A., *Diagnostic and statistical manual of mental disorders (DSM-5®)*. 2013: American
Psychiatric Pub.
78. Badiani, A., et al., *Opiate versus psychostimulant addiction: the differences do matter.* Nature Reviews
Neuroscience, 2011. **12**(11): p. 685.
79. Shalev, U., J.W. Grimm, and Y. Shaham, *Neurobiology of relapse to heroin and cocaine seeking: a review.*
Pharmacological reviews, 2002. **54**(1): p. 1-42.
80. Bossert, J.M., et al., *Neurobiology of relapse to heroin and cocaine seeking: an update and clinical*
implications. European journal of pharmacology, 2005. **526**(1-3): p. 36-50.
81. Mantsch, J.R., et al., *Stress-induced reinstatement of drug seeking: 20 years of progress.*
Neuropsychopharmacology, 2016. **41**(1): p. 335.
82. Li, Q., et al., *Predicting subsequent relapse by drug-related cue-induced brain activation in heroin*
addiction: an event-related functional magnetic resonance imaging study. Addiction biology, 2015. **20**(5):
p. 968-978.
83. Prasad, A.A., et al., *Complementary roles for ventral pallidum cell types and their projections in relapse.*
bioRxiv, 2019: p. 533554.

Figure Legends

Figure 1. Schematic of experimental timeline. Following DREADD or control AAV injection, rats underwent

cocaine self-administration, punishment training, followed by reinstatement and cocaine-induce locomotor

testing. A final reinstatement test preceded sacrifice for neuronal activity (Fos) analysis.

Figure 2. Punishment-resistant rats are more prone to reinstatement than punishment sensitive rats. A)

Probabilistic footshock reduces active lever pressing across all rats, while increasing the number of inactive

presses. SA = self-administration. **B)** Individual variation in punishment sensitivity. Rats that reached at least

0.60mA footshock were considered punishment resistant (red). Punishment sensitive rats (gray) stopped taking

cocaine at <0.60mA footshock intensities. **C)** Distribution of punishment resistance. Most rats cease cocaine

intake at low shock levels (Punishment Sensitive: gray): 53/66 rats, 80.3%), but a subset reached the highest shock

levels (Punishment Resistant: red): 13/66 rats, 19.7%). **D)** Punishment resistant rats reinstated more in the safe

context with cues relative to punishment sensitive rats. This effect was specific to this reinstatement condition. **E)**

Punishment effects on cocaine taking as quantified with suppression ratio [17, 69] correlated with the degree of

reinstatement in the safe context with cues. **F-G)** Punishment resistant rats were no different than sensitive rats

on cocaine intake (F) or cocaine-induced locomotion (G). *** p<0.001. Panel **E**, squares = vehicle injection,

triangles = CNO injection, circles = no injection.

Figure 3. Inhibitory DREADD localization in VP. **A)** Immunofluorescent co-stain for hM4Di-mCherry (red) within

Substance P-expressing VP borders (green). **B)** Virus expression for individual animals on rostrocaudal VP axis. **C)**

Wide-field and **D)** high magnification images of fluorescent *in situ* hybridization for *gad1* (green) combined with

immunofluorescence for hM4Di-mCherry (red). Co-positive hM4Di$^+$ *gad1$^+$* cells labeled with green stars. **E)** 85.3

+/- 3.2 % of hM4Di$^+$ cells are co-positive for *gad1.* 14.7 +/- 3.2 % of hM4Di$^+$ cells are *gad1$^-$* (n=3 rats, total of 559

hM4Di$^+$ cells counted). VM = ventromedial, VL = ventrolateral, DL = dorsolateral. AC = anterior commissure. Scale

bars=500μm (B), 200μm (C), and 20μm (D).

Figure 4. VP inhibition reduces reinstatement especially in punishment resistant rats. A) Top panel: Example

picture of a standard lever press. Bottom panel: CNO in VP-hM4Di rats modestly reduces active and inactive lever

pressing for cocaine under threat of punishment. Control = vehicle-injected rats and CNO-injected misses. **B)** Top

panel: Hesitation behavior in which the rat stretches its trunk towards the lever and extends its paw without

depressing the lever. Bottom panel: Active and inactive lever hesitations quantified during safe (green shading)

and punished (red shading). CNO in VP-hM4Di rats reduced hesitations to relative to control rats. VP inhibition

reduced hesitations to unpunished levels. Control = vehicle-injected rats and CNO-injected misses. **C)** Within

subjects comparisons of reinstatement for VP-hM4Di rats in safe (green shading) and punishment (red shading)

contexts. CNO in VP-hM4Di rats reduced reinstatement in the safe context with cues, without cues, and with

cocaine and no cues, but not in the punishment context with cues. **D)** CNO in control rats did not affect

reinstatement under any condition. Control = eGFP-only rats and rats with hM4Di expression primarily outside VP.

E) CNO in VP-hM4Di punishment resistant rats (red bars) elicited a greater decrease in reinstatement relative to

punishment sensitive rats (gray bars). Data presented as change from vehicle test baseline. $p<0.05^*$, $p<0.01^{**}$,

$p<0.001^{***}$.

Figure 5. VP subregion Fos expression after reinstatement. **A)** Representative image of substance P (VP borders)

and neurotensin (ventromedial VP marker) immunofluorescent co-stain. **B)** Representative image of substance P

and Fos co-stained for Fos quantification. **C-E)** Comparison of ventromedial **(C)**, ventrolateral **(D)**, and dorsolateral

(E) Fos density across reinstatement conditions. Green shading = safe context, red shading = punishment context,

gray shading = home cage controls. F) Rostrocaudal Fos density across reinstatement conditions. Rostral VP

reached significant only in the punishment context with cues. $p<0.05^*$; VM = ventromedial, VL = ventrolateral, DL

= dorsolateral. AC = anterior commissure. Scale Bar = 1000μm.

Supplemental Figures

Supplemental Figure 1. Self-administration in male and female rats. Males (black) did not differ from females

(gray) across 14d of self-administration on active lever presses (top), infusions obtained (middle), inactive lever

presses (bottom).

Supplemental Figure 2. Viral expression in hM4Di misses and eGFP controls. hM4Di misses (thin solid border)

and eGFP controls (thick dotted border) depicted on rostrocaudal VP sections.

Supplemental Figure 3. VP inhibition decreases cocaine-induced rearing, but not distance traveled. A) Within

subjects tests showed that CNO in VP-hM4Di rats fails to alter cocaine-induced horizontal locomotion. **B)** CNO

decreases rearing in VP-hM4Di rats. No effect of CNO in control rats for horizontal locomotion or rearing. Control

= hM4Di misses and eGFP-only rats. ***p<0.001

Supplemental Table 1. List of antibodies used for immunofluorescence (columns 1-3) and for

immunohistochemistry (column 4).

	mCherry+Substance P	GFP+Substance P	Neurotensin+Substance P	Fos+Substance P
Primary Antibodies	**Substance P** Immunostar (20064) Rabbit anti-substance P (1:5000) **mCherry** Clontech (632543) Mouse anti-mCherry (1:2000)	**Substance P** Immunostar (20064) rabbit anti-substance P (1:5000) **GFP** Abcam (AB13970) chicken anti-GFP (1:2000)	**Substance P** Abcam (ab14184) mouse anti-substance P (1:7500) **Neurotensin** Immunostar (24427) rabbit anti-neurotensin (1:5000)	**Substance P** Abcam (ab14184) mouse anti-substance P (1:5000 **Fos** Millipore (abe457) polyclonal rabbit anti-c-Fos (1:5,000)
Secondary Antibodies	**Substance P** Invitrogen (A21206) Donkey anti-rabbit AlexaFluor488 (1:500) **mCherry** Invitrogen (A21203) Donkey anti-mouse AlexaFluor594 (1:500)	**Substance P** Invitrogen (A-21207) donkey anti-rabbit AlexaFluor594 (1:500) **GFP** Jackson ImmunoReasearch (703-545-155) donkey anti-chicken AlexaFluor488 (1:500)	**Substance P** Invitrogen (A21202) donkey anti-mouse AlexFluor 488 (1:500) **Neurotensin** Invitrogen (A-21207) donkey anti-rabbit AlexaFluor594 (1:500)	**Substance P** Jackson ImmunoResearch (711-065-152) biotinylated goat anti-rabbit IgG (1:500) **Fos** Jackson ImmunoResearch (715-065-150) biotinylated donkey anti-mouse (1:500)

Supplemental Table 1. List of antibodies used for immunofluorescence (columns 1-3) and for immunohistochemistry (column 4).

Supplemental Figure 1. Viral expression in hM4Di misses and eGFP controls. hM4Di misses (thin solid border) and eGFP controls (thick dotted border) depicted on rostrocaudal VP sections.

Supplemental Figure 2. Self-administration in male and female rats.
Males (black) did not differ from females (gray) across 14d of self-administration on active lever presses (top), infusions obtained (middle), inactive lever presses (bottom).

Supplemental Figure 3. VP inhibition decreases cocaine-induced rearing, but not distance traveled. A) Within subjects tests showed that CNO in VP-hM4Di rats fails to alter cocaine-induced horizontal locomotion. **B)** CNO decreases rearing in VP-hM4Di rats. No effect of CNO in control rats for horizontal locomotion or rearing. Control = hM4Di misses and eGFP-only rats. ***p<0.001