Sub-lethal insecticide exposure affects host biting efficiency of *Kdr*-resistant

Anopheles gambiae

Diop Malal M [1], Chandre Fabrice [1], Rossignol Marie[1], Porciani Angélique[1], Chateau

Mathieu[1], Moiroux Nicolas[1,2*], Pennetier Cédric[1,3*]

[1] MIVEGEC, Univ Montpellier, CNRS, IRD, Montpellier, France

[2] Institut de Recherche en Sciences de la Santé (IRSS), Bobo-Dioulasso, Burkina Faso

[3] Institut Pierre Richet (IPR), Bouaké, Côte d'Ivoire

*contributed equally to this work

Contacts & correspondence: Nicolas Moiroux nicolas.moiroux@ird.fr ; Cédric Pennetier

cedric.pennetier@ird.fr

Abstract

The massive use of insecticide-treated nets (ITNs) has drastically changed the environment

for malaria vector mosquitoes, challenging their host-seeking behaviour and biting success.

Here, we investigated the effect of a brief exposure to an ITN on the biting behaviour of

Anopheles gambiae mosquitoes and the interaction between such behaviour and the *kdr*

mutation that confers resistance to pyrethroids. To this aim, we developed a video assay to

study the biting behaviour of mosquitoes with similar genetic background, but different *kdr*

locus genotypes (SS i.e. homozygous susceptible, RS i.e. heterozygous and RR i.e.

homozygous resistant), after a brief exposure to either control untreated nets or one of two

types of pyrethroid-treated nets (deltamethrin or permethrin). In presence of untreated nets,

the *kdr* mutation did not influence mosquito blood feeding success but caused differences in

feeding and prediuresis durations and blood meal size. Exposure to deltamethrin ITN

23 decreased the blood feeding success rate of RR and RS mosquitoes, whereas in presence of

24 permethrin ITN, the *kdr* mutation increased the blood-feeding success of mosquitoes.

25 Exposure to the two types of pyrethroid-treated nets reduced feeding duration, prediuresis

26 duration and blood meal size of all three genotypes. Our study demonstrates a complex

27 interaction between insecticide exposure and the *kdr* mutation on the biting behavior of

28 mosquitoes, which may substantially impact malaria vector fitness and disease transmission.

29 **Keywords:** pyrethroid insecticides, long-lasting treated nets, malaria, probing, blood meal,

30 prediuresis*, resistance, kdr*

Introduction

33 Malaria vector mosquitoes become infected by and then can transmit *Plasmodium spp.*

34 parasites during blood meals. *Plasmodium spp.* transmission is strongly dependent on the

35 mosquito biting rate on humans, as formalized by the Ross-MacDonald model of malaria (1).

36 The behaviour and host preferences of blood-feeding mosquitoes are influenced by several

37 factors, primarily their genetic background (2–5), but also environmental factors, such as host

38 diversity and availability (6,7), and the presence/absence of physical and chemical barriers,

39 such as Insecticide-Treated Nets (ITNs) (8–10).

40 ITNs should hamper the contacts between humans and nocturnal, anthropophilic (i.e. that bite

41 humans preferentially) and endophagic mosquitoes (i.e. that bite preferentially indoors), such

42 as *An. gambiae* (11). The widespread use of pyrethroid (PYR) insecticides in public health

43 (i.e. ITNs or indoor residual sprayings), crop protection and other selective pressures (such as

44 pollutants in mosquito environment) drove adaptations in malaria vectors to reduce the

45 insecticidal effect (12–17).

Physiological PYR resistance involves two main mechanisms: (i) metabolic resistance, due to

quantitative or qualitative changes in detoxification enzymes (cytochrome P450

monooxygenases, esterases and glutathione S-transferases), and (ii) target site resistance, due

to non-synonymous mutations in the voltage-gated sodium channels that are called knock-

down resistance (*kdr*) mutations (12,18).

Mosquitoes can reduce vector control tools efficacy also through behavioural adaptations. In

areas of sub-Saharan Africa where large-scale vector control programmes have been

implemented, several observations suggest that malaria vectors can avoid contacts with ITNs

or insecticides on walls by modulating their host-feeding activity. For instance, following the

large scale distribution of ITNs in Benin, some *An. funestus* populations started to feed

predominantly at dawn/early in the morning or in broad daylight (19,20), when most people

are outside the nets (20). Behavioural modulation might also be influenced by physiological

resistance mechanisms. Indeed, experimental hut trials showed that in *An. gambiae*, indirect

behavioural indicators in response to ITN presence (deterrence and induced exophily, i.e. the

proportion of mosquitoes that exit early and are found in exit traps relative to the untreated

hut) are related to the physiological tolerance to insecticide (21). Moreover, a study

demonstrated that *kdr* homozygous resistant mosquitoes have longer contacts with ITNs than

homozygous susceptible mosquitoes, which are more excited by PYR irritant effect (22).

Altogether, these findings indicate that insecticide treatments could affect the behaviour of

malaria vectors. However, the effects of insecticide exposure and *kdr* mutations on the biting

activity of *An. gambiae* remain poorly investigated.

The last step of the mosquito host-seeking behaviour after reaching a host protected by an

ITN is biting for taking a blood meal. During the host-seeking phase and the penetration

through a hole in the net, mosquitoes can be exposed to sub-lethal doses of insecticide (23–

25). Such doses do not cause death, but can have several physiological or behavioural effects

on host-seeking mosquitoes (26). For instance, sub-lethal doses of insecticide can affect the

feeding and reproductive behaviour of some blood-sucking insects (27). As the PYR

insecticide on ITNs acts on specific sites in the mosquito nervous system, it might alter some

physiological processes involved in the biting behaviour of malaria vectors. There is currently

scarce literature on this subject (28).

In the present study, we investigated whether pre-exposure to an ITN modulates the mosquito

ability to take a blood meal by using experimental conditions that mirror the exposure to

insecticide occurring when a mosquito passes through an ITN after having located a host. We

also assessed whether the *kdr* mutation (L1014F) modifies blood feeding success and biting

behaviour of *An. gambiae*.

Methodology

Ethical Considerations

This study was carried out in strict accordance with the recommendations of Animal Care and

Use Committee (29) named "Comité d'éthique pour l'expérimentation animale; Languedoc

Roussillon" and the protocol was approved by the Committee on the Ethics of Animal

Experiments (CEEA-LR-13002 for the rabbits).

Mosquito strains and rearing procedures

Two mosquito laboratory strains were used for this study. One is the insecticide-susceptible

Kisumu strain (KISUMU1, MRA-762, VectorBase stable ID VBS0000026 on

vectorbase.org), isolated in Kenya in 1975. This strain is PYR-susceptible and homozygous

(SS) for the L1014 codon. The second one is the Kdr-kis strain that is PYR-resistant and

homozygous (RR) for the L1014F *kdr* mutation. The Kdr-kis strain was obtained by

introgression of the *kdr* allele (L1014F) into the Kisumu genome through 19 successive back-

crosses between Kisumu and VKPer (30). The VKPer strain, originated from a rice growing

area named Vallée du Kou, less than 40 kms north of Bobo-Dioulasso (Burkina Faso) was

used to obtain Kdr-kis. VKPer displayed the same expression level of metabolic resistance

enzyme as Kisumu (31). Both Kisumu and Kdr-kis strains are maintained at the insectary of

IRD (Institut de Recherche pour le Développement) – WHO collaborating center FRA-72 in

Montpellier, France.

Polymorphisms between Kisumu and Kdr-kis strains are expected to be restricted in the

flanking region of the *Kdr* allele (15 cM for 19 backcrossing generations (32)) and the

observed phenotypes are therefore expected to be associated to this genetic area.

Heterozygous individuals (RS) for the L1014F *kdr* mutation were obtained by crossing once

Kisumu SS females (F1 progeny) with Kdr-kis RR males. Therefore, the three genotypes have

a common genetic background for most of their genome.

Mosquitoes were reared at $27 \pm 1°C$, 70-80% relative humidity under a 12h:12h (light : dark)

photoperiod in the insectary and fed on rabbits. Gravid females were allowed to lay eggs on

wet filter paper inside mesh-covered cages. Eggs were dispensed in plastic trays containing

osmotic water. Larvae were kept in trays and fed with TetraMin® fish food. Pupae were

removed and allowed to emerge inside 30x30x30cm cages. After emergence, adults could

feed *ad libitum* on 10% sucrose solution.

On each experimental day, an average of 8 (2 to 16 according to the insectary production)

female mosquitoes (7 to 9 days old) never fed with blood were randomly collected from the

rearing cages and placed in cups covered by gauze. By using 7-9 days old *An. gambiae*, we

expected to enhance the probability that these females were inseminated (33) and

physiologically active for host-seeking (34). However, mated status was not checked.

Mosquitoes were starved the day before the experiment because sucrose inhibits blood avidity

in mosquitoes (35).

Insecticide exposure

To simulate the contact with an ITN that may occur before the vector finds a way to reach a

host, starved females were individually exposed to PYR insecticide-treated netting using an

experimental setup in which insects can only walk or stand on the net surface and that was

initially designed to measure the median knock down time (36). We used three net types (i.e.

treatments): an untreated net (negative control, hereafter UTN), an Olyset Net[®] (impregnated

with 1000 mg/m^2 of permethrin, hereafter Olyset) and a PermaNet[®] 2.0 (coated with

55 mg/m^2 of deltamethrin, hereafter PermaNet). Based on previous experiments (25), each

mosquito was exposed for 30 seconds. This is the median time of contact with permethrin-

treated netting before PYR-susceptible anopheles (SS genotype) locate a hole to reach the

host (25). After exposure, a 1-min latency period was observed before releasing the insect in

the behavioural assay setup.

Behavioural assay

Experiments were conducted in a biting behavioural assay setup designed for video recording

the biting behaviour of the tested mosquitoes (Figure 1). The setup is made of a foam board

and composed of an observation tunnel (OT) and an observation zone (OZ) separated by a

transparent plastic (TP). The OZ is a triangular prism, closed on its base by a removable paper

sheet with a 1-cm diameter hole. The hole allows mosquitoes to bite the ear of a rabbit (R)

that is maintained immobile in a restraining cage (RC) to limit ear movement during blood

feeding. The same rabbit was used during all experiments. The experimental room was faintly

illuminated with a compact fluorescent lamp bulb placed at 15 cm from the OZ to allow the

acquisition of the biting behaviour sequence by the digital video system.

Figure 1 Experimental set-up to monitor the biting behavioural sequence.

OT: Observational tunnel, TP: Transparent plastic, OZ: Observational zone (mosquito biting the

rabbit ear), R: Rabbit, RC: Restraining cage

For each trial, a mosquito was individually released inside the behavioural assay through a

circular opening (CO) located on its lateral face. Cotton was used to plug the CO after

releasing. The number of released individual anopheles per genotype and treatment ranged

from 43 to 86. Each mosquito was filmed for 10 minutes using a Sony® Digital HD Video

Camera (HDR-XR550) placed on the top of the OT (Figure 1). The video camera was

connected to a computer screen placed outside the experimental room to allow real-time

monitoring of each mosquito, to avoid any disturbance of the rabbit and interference of the

experimenter on the mosquito behaviour. The MPEG-2 recordings (PAL video: 720x576

pixels at 25 frames/s) were analysed using the behavioural event recording program

EVENT01, version 1.2.4 ([©]R.D. Collins and M.K. Tourtellot, 1993-2002).

The experimenter used latex gloves to avoid any contamination by human skin odours. The

OZ was cleaned with ethanol and the removable paper was changed after each mosquito was

tested to avoid any contamination by insecticide residues between behavioural assays. The

feeding experiments were done under insectary conditions (temperature at $27 \pm 1°C$, and

relative humidity 70-80%), in a dark room during the night (according to the Light:Dark

photoperiod of the insectary).

Feeding success and behavioural parameters

As exposure to insecticide can induce a knockdown (KD) effect during the trial, mosquitoes

were recorded as KD, if they were lying on their side or their back. A mosquito (whatever its

KD phenotype) was scored as successful if it was fed at the end of the 10-min trial (whatever

the amount of blood it took) and unsuccessful if it did not. After the trial, successfully fed

mosquitoes were stored at -35°C for measuring the blood meal size (see below).

Analysis of the acquired images allowed quantifying the following variables in fed

mosquitoes: (1) number of probing events, (2) probing duration (the time from the

introduction of the stylet fascicule into the rabbit skin to blood appearance in the mosquito

abdomen), (3) feeding duration (from the blood appearance in the abdomen to the beginning

of the stylet fascicule withdrawal), and (4) prediuresis duration (the time during which

excretion of rectal fluid (plasma, water, metabolic wastes) is observed as red bright drops

during feeding and after proboscis withdrawal).

Blood meal volume measurement

The blood intake was evaluated by quantifying the haemoglobin amount, as described by

Briegel *et al.* (37). Each engorged mosquito was stored in one 1.5 ml Eppendorf tube at -

35°C. Then, the whole abdomen was ground in the presence of 0.5 ml of Drabkin's reagent

until it was completely disintegrated. Haemoglobin then reacted with the Drabkin's reagent

and was converted into haemoglobin cyanide (HiCN). Samples were incubated at room

temperature (25°C) for 20 min and then a chloroform solution was added in each tube.

Samples were centrifuged at 5600 rpm for 5 min and the aqueous supernatant (containing

HiCN) was placed in a new 1.5 ml Eppendorf tube. The absorbance was read at a wavelength

of 550 nm using a microplate spectrophotometer. Two replicates of the reading were done for

each mosquito and their absorbance values were averaged. A sample of the rabbit blood was

used as control for calibration curves.

As the blood meal volume is correlated with the mosquito size (38), the blood meal volume of

all mosquitoes from the same batch was divided by the average weight of five randomly

selected mosquitoes from the same rearing cage. The resulting ratio was expressed in µL of

blood per µg of weight and called weighted blood meal volume or blood meal size.

Statistical analysis

All statistical analyses were performed using the R software, version 3.5 (39).

We analysed the feeding success (coded as 1 for fed mosquitoes and 0 for unfed ones) with a

binomial logistic mixed-effect model using function 'glmmTMB' in the 'glmmTMB' package

(40). We analysed the number of probing events with a zero-truncated negative binomial

mixed-effect model using function 'glmmTMB'. We analysed durations (probing, feeding and

prediuresis) with a mixed effect Cox proportional hazard model using function 'coxme' of the

'coxme' package (41). We analysed weighted blood meal size with a linear mixed effect

model using function 'lmer' in the 'lme4' package (42). All models included the *kdr*

genotypes (SS, RS or RR), type of pre-exposure (UTN, Permethrin or Deltamethrin) and their

interactions as fixed terms explanatory variables and the date of the experiment as a random

intercept.

In order to discriminate between the effect of KD phenotype and other pleiotropic effects of

the *kdr* mutation on feeding success and behavioural parameters, we fitted all previously

described models on the dataset but for KD mosquitoes. To complement the later analysis, we

tested the effect of the KD phenotype on feeding success. For this task, we added KD

phenotype and its interactions as explanatory variables in the binomial model of feeding

success fitted only on mosquitoes exposed to the insecticide treatments (Permethrin and

Deltamethrin). This allowed us to compare feeding success between KD and non-KD

mosquitoes of each genotype and treatment.

In order to assess the relationship between KD phenotype and the genotypes for the *kdr*

mutation, KD (coded as 1 for KD mosquitoes and 0 for others) was analysed using a binomial

model with the *kdr* genotypes (SS, RS or RR), type of exposure (Permethrin or Deltamethrin)

and their interactions as fixed terms. Because the dataset showed data separation, we fitted the

model using the bias-reduction method developed by Firth (43). We used the 'brglmFit'

function of the 'brglm2' package (44) for this task.

We used Tukey's post-hoc test to perform multiple comparisons among genotypes and

treatments using 'emmeans' function (45). We computed Odds Ratios (OR) for the binomial

models, Hazard Ratios (HR) for Cox models, Rate Ratios (RR) for the negative binomial

models, Mean Differences (MD) for the linear models and their 95% confidence intervals.

We calculated the binomial confidence interval of feeding rates and knock-down rates with

the Wilson's score method using the 'binconf' command from the 'Hmisc' package (46).

Effect of feeding duration on blood meal size and prediuresis duration was analysed using a

linear mixed effect model and a mixed effect cox proportional hazard model, respectively.

Fixed terms included genotypes and treatment and interactions and date was set as a random

intercept. We used function 'emtrends' of the 'emmeans' package (45) to obtain estimates and

95% confidence intervals of the marginal slopes of the trends (between feeding duration and

blood meal size or prediuresis duration) for each genotypes and treatments.

Data and codes used for analyses and figures are available (47).

Results

In total, 511 *An. gambiae* females (182 SS, 156 RS and 173 RR) were released individually

for the study. The number of mosquitoes released, fed, unfed and knock-downed among

genotypes and treatments are shown in Table 1.

Table 1. Numbers of mosquito released per genotype and treatment

Genotype	Treatment	N tested	Fed		Unfed	
			KD	Non-KD	KD	Non-KD
SS	Untreated	86	0	41	0	45
SS	Permethrin	50	5	5	33	7
SS	Deltamethrin	46	1	15	18	12
RS	Untreated	70	0	40	0	30
RS	Permethrin	43	3	14	11	15
RS	Deltamethrin	43	1	9	15	18
RR	Untreated	70	0	40	0	30
RR	Permethrin	53	0	41	0	12
RR	Deltamethrin	50	4	12	12	22

SS: homozygous susceptible genotype, RS: heterozygous genotype, RR: homozygous resistant genotype, KD: Knockdown.

Impact of the kdr mutation on feeding success and biting behaviour

When female mosquitoes were pre-exposed to UTN (i.e., untreated netting), no difference in their feeding success was found among the three genotypes (OR_{RS-SS}=1.47 [0.67, 3.22]; OR_{RR-SS}=1.47 [0.68, 3.19]; OR_{RR-RS}=1.00 [0.44, 2.25]; Figure 2A).

Analysis of the biting behaviour of successful mosquitoes in absence of insecticide pre-exposure showed that feeding and prediuresis durations were shorter in RR than in both SS and RS mosquitoes (feeding duration: HR_{RR-SS} = 2.15 [1.26, 3.69]; HR_{RR-RS} = 2.46 [1.43, 4.23]; Figure 2B; prediuresis duration: HR_{RR-SS} = 1.94 [1.02, 3.71]; HR_{RR-RS} = 2.11 [1.11, 4.01]; Figure 2C). The weighted blood meal volume of RR mosquitoes was lower than that of RS mosquitoes (MD_{RS-RR} = 0.99 [0.24, 1.74]; Figure 2D)). Number of probing events and probing duration were not significantly different among genotypes (SS, RS and RR) (Supplementary Tables 1 and 2).

Figure 2 Feeding success and biting behavior in absence of insecticide of *Anopheles*

***gambiae* females of each *kdr* genotype**.

Feeding success of each genotype and 95% binomial confidence intervals of the proportions (error

bars) are shown in panel A (numbers n of mosquitoes exposed to the untreated net are indicated).

Panels B, C and D show boxes-and-whiskers plots of feeding duration, prediuresis duration and

blood-meal size, respectively (numbers n of blood-fed mosquitoes tested for behavioural parameters

are indicated in panel B). Boxes indicate 1st-3rd quartile and median values. Whiskers indicate 1.5

inter-quartile range. P-values according to Tukey's test after binomial mixed-effect model (panel A),

mixed-effect cox proportional hazard model (panels B and C) and after linear mixed effect model

(panel D) are indicated.

Impact of insecticide exposure on knockdown rates

Exposure to permethrin or deltamethrin induced 76 % [62.6, 85.7] and 41.3% [28.3, 55.7] KD

rates in SS mosquitoes, respectively. It induced 32.6% [20.5, 47.5] and 37.2% [24.4, 52.1]

KD rates in RS mosquitoes, respectively (Table 1). Among RR mosquitoes, permethrin

exposure did not induce any KD effect, whereas deltamethrin exposure led to 30.2% [20.8,

45.8] of KD mosquitoes (Table 1). *Kdr* genotype was highly correlated with KD rates of

mosquitoes exposed to permethrin (OR_{RR-RS} = 0.019 [6.1×10^{-4}, 0.597], OR_{RR-SS} = 0.003

[9.7×10^{-5}, 0.095], OR_{RS-SS} = 0.160 [5.41×10^{-2}, 0.471]) but not for mosquitoes exposed to

deltamethrin (OR_{RR-RS} = 0.80 [0.29, 2.22], OR_{RR-SS} = 0.67 [0.25, 1.83], OR_{RS-SS} = 0.85 [0.31,

2.34]).

Impact of insecticide exposure on feeding success

When compared to the UTN condition, exposure to permethrin reduced significantly the

feeding success of SS mosquitoes ($OR_{perm-UTN}$=0.28 [0.10, 0.73]; Figure 3A), but not that of

RS ($OR_{perm-UTN}$=0.49 [0.19, 1.26]; Figure 3B). Feeding success of RR mosquitoes was higher

(marginally not significant) ($OR_{perm-UTN}$=2.58 [0.97, 6.84]; Figure 3B) when exposed to

permethrin. Exposure to deltamethrin reduced significantly the feeding success of RS and RR

mosquitoes ($OR_{delta-UTN}$=0.23 [0.08, 0.64] and $OR_{delta-UTN}$=0.35 [0.14 – 0.88]; Figure 3B and

3C), but not of SS mosquitoes ($OR_{delta-UTN}$=0.58 [0.24, 1.43]; Figure 3A). When excluding

KD mosquitoes from this analysis, all trends were kept (supplementary Figure 1) except that

feeding success of SS mosquitoes was no longer reduced by permethrin exposure when

compared to UTN ($OR_{perm-UTN}$=0.78 [0.18, 3.40]; supplementary Figure 1A). Indeed among

SS mosquitoes, non-KD had higher feeding success than those recorded as KD (OR_{nonKD-}

$_{KD}$=4.71 [1.06, 20.9]; supplementary Table 3). The same was true for SS mosquitoes exposed

to deltamethrin ($OR_{nonKD-KD}$=22.5 [2.59, 195]; supplementary Table 3).

Figure 3 Feeding success after exposure to insecticides of *Anopheles gambiae* females of each *kdr* genotype.

Feeding success of SS, RS, and RR (panels A, B and C, respectively) genotypes when exposed to untreated, permethrin-treated (Olyset) and deltamethrin-treated (PermaNet) nettings are shown with 95% binomial confidence intervals of the proportions (error bars). Numbers n of mosquitoes exposed to each treatment and for each genotype are indicated. P-values according to Tukey's test after binomial mixed-effect model is indicated.

When comparing the feeding success among genotypes after insecticide exposure, the feeding rate of RR mosquitoes was higher than that of SS and RS mosquitoes after permethrin exposure (OR_{RR-SS}=13.82 [4.35, 43.92]; OR_{RR-RS}=5.29 [1.76, 15.86], Supplementary Figure 1B), whereas the feeding success of RS mosquitoes was although higher, not significantly

different than that of SS mosquitoes (OR_{RS-SS}=2.61 [0.85, 8.02], Supplementary Figure 2B).

In contrast, exposure to deltamethrin did not induce any difference in the feeding success of

the three genotypes (OR_{RR-SS}=0.891 [0.31, 2.53]; OR_{RR-RS}=1.55 [0.51, 4.76]; OR_{RS-SS}=0.57

[0.18, 1.80], Supplementary Figure 2C). We observed the same trends when excluding KD

mosquitoes from the analysis (supplementary Figure 3).

Impact of insecticide exposure on biting behaviour

After exposure to deltamethrin, feeding duration, prediuresis duration and weighted blood

meal size were significantly reduced in SS mosquitoes compared to the UTN condition

(feeding duration: $HR_{delta-UTN}$ = 4.38 [2.09, 9.15]; prediuresis: $HR_{delta-UTN}$ = 5.31 [2.11, 13.36];

blood meal size; $MD_{delta-UTN}$ = -1.02 [-1.82, -0.23]; Figure 4A, 4B and 4C, respectively). A

similar trend was observed after exposure to permethrin for feeding duration ($HR_{perm-UTN}$ =

2.87 [1.21, 6.81]; Figure 4A) and weighted blood meal volume ($MD_{perm-UTN}$ = -0.97 [-1.85, -

0.08]; Figure 4C) but not for prediuresis duration ($HR_{perm-UTN}$ = 2.27 [0.92, 5.63]; Figure 4B).

For this latter parameter, the non-significance was probably due to a lack of power.

Compared to UTN, permethrin and deltamethrin reduced the feeding and prediuresis

durations as well as blood meal size of both RS (feeding duration: $HR_{perm-UTN}$ = 7.42 [3.55,

15.53], $HR_{delta-UTN}$ = 4.57 [1.94, 10.76], Figure 4D; prediuresis duration: $HR_{perm-UTN}$ = 9.44

[3.36, 26.46] and $HR_{delta-UTN}$ = 7.22 [2.70, 19.29], Figure 4E, blood-meal size: $MD_{perm-UTN}$ = -

1.97 [-3.04, -0.90] and $MD_{delta-UTN}$ = -2.01 [-2.97, -1.05], Figure 4F) and RR genotypes

(feeding duration: $HR_{perm-UTN}$ = 2.06 [1.20, 3.51] and $HR_{delta-UTN}$ = 3.78 [1.83, 7.82], Figure

4G; prediuresis duration: $HR_{perm-UTN}$ = 1.71 [0.80, 3.64] and $HR_{delta-UTN}$ = 9.31 [3.57, 24.28],

Figure 4H; blood-meal size: $MD_{perm-UTN}$ = -1.52 [-2.23, -0.81] and $MD_{delta-UTN}$ = -1.41 [-2.28,

-0.54], Figure 4I). Number of probing events and probing duration were not significantly

different among treatments (UTN and ITNs) whatever the genotype (Supplementary Tables 4

and 5).

Figure 4 Biting behavior after exposure to insecticides of *Anopheles gambiae* females of each *kdr* genotype.

Boxes-and-whiskers plots of feeding duration (panels A, D and G), prediuresis duration (panels B, E and H) and blood-meal size (panels C, F and I) are shown for each genotypes SS (panels A, B and C), RS (panels, D, E and F) and RR (panels G, H and I). Boxes indicate 1^{st}-3^{rd} quartile and median values. Whiskers indicate 1.5 inter-quartile range. Numbers n of blood-fed mosquitoes for each genotypes and tested for behavioural parameters are indicated in panels A, D and G. P-values according to Tukey's test after mixed-effect cox proportional hazard model (panels A, B, D, E, G and H) and after linear mixed effect model (panels D, F and I) are indicated.

When comparing the biting behavior among genotypes after insecticide exposure, we found

that prediuresis duration of RS mosquitoes was shorter than that of SS mosquitoes after

permethrin exposure ($HR_{RS-SS} = 3.82$ [1.15, 12.7], supplementary Table 8). Moreover,

prediuresis duration of RR mosquitoes was shorter than that of SS after deltamethrin exposure

($HR_{RR-SS}=3.41$ [1.13, 10.29], supplementary Table 8). For all other parameters, we were not

able to evidence any differences among genotypes (Supplementary Tables 1, 2, 7 and 8).

In absence of exposure to insecticide (UTN), the blood meal size and the prediuresis duration

were positively correlated with the feeding duration for all genotypes (supplementary Tables

6 and 7). With the exception of blood meal size of RS exposed to permethrin, these

relationships were not observed when mosquitoes were exposed to insecticides

(supplementary Tables 9 and 10), suggesting a perturbation of processes underlying these

correlations.

Excluding 14 KD mosquitoes from the analyses of the biting behaviour parameters do not

significantly change the results (supplementary Tables 11 to 22).

Discussion

To investigate the influence of ITN exposure on the biting behaviour of *An. gambiae*

mosquitoes, we used mosquitoes that share the same genetic background, but for the *kdr* allele

locus, and then exposed them to ITN prior to blood feeding.

The blood-feeding success did not differ between the three genotypes in the absence of

insecticide exposure. Therefore, the *kdr* mutation was not associated with significant change

of blood meal success rate. However, feeding duration and blood meal size were different

between genotypes. RR mosquitoes spent less time taking their blood meal than RS and SS

mosquitoes. This might confer an advantage as fast feeding reduces the risk to be killed

because of the host defensive behaviour (48). On the other hand, RS mosquitoes took higher

blood volumes than RR females. This could improve the completion of oogenesis in RS

mosquitoes (49) and increase their fecundity compared to RR (50). However, large blood

meals reduce the flying ability, escape speed and agility required to avoid predators

(48,51,52). These different trades-offs between behavioural traits that might enhance

fecundity or survival in the three genotypes are of great interest and deserve further

investigations in relation with the ecological and vector control environment. Such trades-offs

possibly affect mosquito fitness and may therefore drive not only the evolution of insecticide

resistance in mosquitoes but also parasite transmission. For example, a decrease in blood meal

duration and size might increase the frequency of multiple feedings and consequently the risk

of *Plasmodium* transmission (53). Similarly, a bigger blood meal size might increase the

probability of mosquito infection by gametocytes (54).

Exposure to permethrin and deltamethrin induced opposite outcomes in term of blood feeding

success (increase and decrease, respectively) in RR mosquitoes. This opposite effect on the

feeding success rate of RR females might be linked to the different chemical properties of

permethrin and deltamethrin that induce two types of bursting activity of sodium channels

(55,56). Type II pyrethroids, such as deltamethrin, further delay the inactivation of the

voltage-gated sodium channel and in a less reversible way than type I pyrethroids, such as

permethrin (57). The lower PYR susceptibility of homozygous resistant mosquitoes could

lead to their over-stimulation compared to susceptible and heterozygous mosquitoes that are

more affected by the toxic effect of such insecticides (58).

In contrast, among females that have been successful in taking a blood meal, the behavioural

sequence was altered in the same way by both insecticides. They both induced a decrease of

feeding duration and prediuresis duration. This is in agreement with the results of Hauser *et*

al. (50) showing that mosquitoes biting trough permethrin+PBO nets (Olyset Plus) had less

feeding successes, shorter feeding duration and lower blood meal sizes compared to those

biting trough untreated nets. As discussed above, short feeding durations and small blood-

meal lead us to expect that exposure to insecticide may (i) reduce the risk of the vector to be

killed due to the defensive behaviour of the host or due to predators, (ii) reduce the possible

number of parasites ingested during one feeding attempt that may be compensated by (iii) the

increase of multiple feeding (and therefore the higher risk of human-to-mosquito and

mosquito-to-human parasite transmission).

Whatever the genotype, blood meal size and prediuresis duration were correlated to feeding

duration in absence of exposure to insecticides. This relationship was, in most cases, no

longer observed when mosquitoes were exposed to insecticides indicating that sub-lethal

contact with insecticides disrupt physiological processes involved in blood meal intake. Such

indirect evidence highlight the need to further investigate the consequences of sub-lethal

contact with insecticides.

Prediuresis duration was substantially reduced in RS and RR females after exposure to ITNs.

Prediuresis is an intestinal mechanism that plays a crucial role in protein concentration during

feeding (59) and contributes to thermoregulation (60,61). This perturbation of the prediuresis

phase by sub-lethal doses of PYR insecticides could lead to toxic accumulation of metabolic

wastes and products of oxidative stress in the haemolymph that might affect the lifespan of

mosquitoes. This results suggest that mosquitoes with the *kdr* mutation might be more

susceptible to new chemicals that target the mosquito renal system and that are currently

developed as an alternative to the currently used insecticides (62).

As expected, we found a strong relationship between *kdr* genotype and KD phenotype when

mosquitoes were exposed to Permethrin. However with deltamethrin, we were not able to find

such a relationship and mosquitoes carrying the *kdr* mutation experienced moderate levels of

KD. Indeed, deltamethrin is expected to induce higher knock-down rates than permethrin

against resistant populations of *Anopheles gambiae* (63). This observation is also true when

looking at mortality in experimental hut trial in areas with high frequencies of the *kdr*

mutation in the vector population (64–66). This difference between permethrin and

deltamethrin effect may be linked to the different chemical properties of permethrin and

deltamethrin (type I and Type II pyrethroids) as describe above.

We found that KD reduced the feeding success of mosquitoes exposed to PYR insecticides,

particularly in SS mosquitoes. However when analyzing feeding success and behavior of non-

KD mosquitoes carrying the *kdr* mutation, we observed the same trends than we get when

including KD mosquitoes. This indicates that observed differences in feeding success and

behavior are therefore directly linked to the presence of the mutation and not only a

consequence of the KD phenotype.

This work has some limitations. First, we were not able to randomize genotypes over time

because of rearing constraints. Consequently all RS mosquitoes were tested during the last

month of experiments. This did not induce any effect when analyzing the treatment effect

relative to the genotype but might have introduced a bias while comparing RS with RR and/or

SS mosquitoes (as experimental period is possibly a confounding factor for RS genotype).

The second limitation relies on body size measurement. Indeed, we chose to use an easy

method to get a proxy of mosquito size. We randomly selected 5 anopheles females from each

rearing cages used during the experiment, weighed them together and used the mean weight to

adjust blood-meal size. An individual wing length measurement would allow to avoid any

bias in developmental variability within each rearing cage.

To conclude, our study demonstrates a complex interaction between insecticide exposure and

the *kdr* mutation on the biting behaviour of mosquitoes. The behavioural modulation induced

by PYR-treated nets also raises concerns about the consequences of the *kdr* resistance-

insecticide interaction. In previous studies, we evidenced that RR mosquitoes prefer a host

protected by a permethrin-treated net rather than an untreated net (67) and that heterozygotes

RS mosquitoes have a remarkable ability to find a hole into a bet net (25). Herein, we have

completed the sequence by showing that *kdr* homozygous resistant *An. gambiae* displayed

enhanced feeding success when exposed to permethrin ITN. However here, we exposed all

our mosquitoes for a constant duration to the treated nets which does not reflect the variability

that happens in natural conditions. Indeed, we can expect from the literature (22,23,25) that

both the genotype for the *kdr* mutation and the type of pyrethroids on LLINs may affect the

contact duration. It would be of great interest to decipher with the relationship between the

time of contact with the insecticide and the feeding sequence.

Insecticide resistance genes in malaria vectors could modify vector competence and the

dynamics of infection by *P. falciparum*. For instance, recent studies have shown that parasite

infection increases insecticide susceptibility in mosquitoes carrying the *kdr* mutation (68) and

that insecticide exposure reduces parasite development in resistant mosquitoes (69). In

addition, malaria parasites have been shown to modify the feeding behaviour of their

mosquito vectors in ways that favour their transmission (70–72), but the role of insecticide

resistance that could modulate this phenomenon has not been investigated yet (73). It is

therefore urgent to decipher the links between insecticide exposure, resistance mechanisms

and infection by *P. falciparum* on the host-seeking and biting behaviour of malaria vectors to

better understand malaria transmission in areas where insecticidal tools for malaria prevention

are implemented. All these interactions should then be used as variables to include host-

seeking behavioural modulation by *kdr* resistance in models of resistance evolution and *P.*

falciparum transmission to better understand and/or predict the efficacy of vector control

strategies (74).

Acknowledgements

MD was supported by IRD's PhD student fellowship. The authors would like to acknowledge the technical/research platform dedicated on vectors at the IRD centre and all members of the platform for providing mosquitoes and technical support. This platform is a member of the Vectopole Sud network and of the LabEX CEMEB (Centre Méditerranéen de l'Environnement et de la Biodiversité) in Montpellier. We thank Dr Anna Cohuet and Dr Thierry Lefèvre for their valuable comments on the manuscript.

Conflict of interest disclosure

The authors of this preprint declare that they have no financial conflict of interest with the content of this article. NM and CP are *PCI Entomolgy* recommenders.

References

1. Garrett-Jones C, Shidrawi GR. Malaria vectorial capacity of a population of Anopheles gambiae: an exercise in epidemiological entomology. Bulletin of the World Health Organization. 1969;40(4):531.

2. Lefèvre T, Gouagna L-C, Dabire KR, Elguero E, Fontenille D, Costantini C, et al. Evolutionary lability of odour-mediated host preference by the malaria vector *Anopheles gambiae*. Tropical Medicine & International Health. 2009 Feb;14(2):228–36. doi:10.1111/j.1365-3156.2009.02206.x

3. Dana AN, Hong YS, Kern MK, Hillenmeyer ME, Harker BW, Lobo NF, et al. Gene expression patterns associated with blood-feeding in the malaria mosquito Anopheles gambiae. BMC Genomics. 2005 Jan 14;6(1):5. doi:10.1186/1471-2164-6-5

4. Rund SSC, Hou TY, Ward SM, Collins FH, Duffield GE. Genome-wide profiling of diel and circadian gene expression in the malaria vector Anopheles gambiae. Proceedings of the National Academy of Sciences. 2011 Jun 29;108(32):E421–30. doi:10.1073/pnas.1100584108

5. Zwiebel LJ, Takken W. Olfactory regulation of mosquito–host interactions. Insect Biochemistry and Molecular Biology. 2004 Jul;34(7):645–52. doi:10.1016/j.ibmb.2004.03.017

6.   Lyimo IN, Ferguson HM. Ecological and evolutionary determinants of host species
choice in mosquito vectors. Trends in Parasitology. 2009 Apr;25(4):189–96.
doi:10.1016/j.pt.2009.01.005

7.   Killeen GF, McKenzie FE, Foy BD, Bøgh C, Beier JC. The availability of potential
hosts as a determinant of feeding behaviours and malaria transmission by African
mosquito populations. Trans R Soc Trop Med Hyg. 2001 Oct;95(5):469–76.
doi:10.1016/s0035-9203(01)90005-7

8.   Mathenge EM, Gimnig JE, Kolczak M, Ombok M, Irungu LW, Hawley WA. Effect of
permethrin-impregnated nets on exiting behavior, blood feeding success, and time of
feeding of malaria mosquitoes (Diptera: Culicidae) in western Kenya. J Med Entomol.
2001;38(4):531–6. doi:10.1603/0022-2585-38.4.531

9.   Lefèvre T, Gouagna L-C, Dabiré KR, Elguero E, Fontenille D, Renaud F, et al. Beyond
nature and nurture: phenotypic plasticity in blood-feeding behavior of Anopheles
gambiae s.s. when humans are not readily accessible. Am J Trop Med Hyg. 2009
Dec;81(6):1023–9. doi:10.4269/ajtmh.2009.09-0124

10. Gatton ML, Chitnis N, Churcher T, Donnelly MJ, Ghani AC, Godfray HCJ, et al. The
importance of mosquito behavioural adaptations to malaria control in Africa. Evolution.
2013 Apr;67(4):1218–30. doi:10.1111/evo.12063

11. Pates H, Curtis C. Mosquito behavior and vector control. Annu Rev Entomol.
2005;50:53–70. doi:10.1146/annurev.ento.50.071803.130439

12. Ranson H, N'Guessan R, Lines J, Moiroux N, Nkuni Z, Corbel V. Pyrethroid resistance
in African anopheline mosquitoes: what are the implications for malaria control? Trends
Parasitol [Internet]. 2011;27. doi:10.1016/j.pt.2010.08.004

13. Corbel V, N'Guessan R. Distribution, mechanisms, impact and management of
insecticide resistance in malaria vectors: a pragmatic review. In: Manguin S, editor.
Anopheles mosquitoes: new insights into malaria vectors [Internet]. Rijeka: Intech;
2013 [cited 2020 Jan 28]. p. 579–633. Available from:
http://www.documentation.ird.fr/hor/fdi:010060052

14. Chouaïbou MS, Fodjo BK, Fokou G, Allassane OF, Koudou BG, David J-P, et al.
Influence of the agrochemicals used for rice and vegetable cultivation on insecticide
resistance in malaria vectors in southern Côte d'Ivoire. Malar J. 2016 24;15(1):426.
doi:10.1186/s12936-016-1481-5

15. Liu N. Insecticide Resistance in Mosquitoes: Impact, Mechanisms, and Research
Directions. Annual Review of Entomology. 2015;60(1):537–59. doi:10.1146/annurev-
ento-010814-020828

16. Poupardin R, Riaz MA, Jones CM, Chandor-Proust A, Reynaud S, David J-P. Do
pollutants affect insecticide-driven gene selection in mosquitoes? Experimental evidence
from transcriptomics. Aquatic Toxicology. 2012 Jun 15;114–115:49–57.
doi:10.1016/j.aquatox.2012.02.001

17. Carrasco D, Lefèvre T, Moiroux N, Pennetier C, Chandre F, Cohuet A. Behavioural
adaptations of mosquito vectors to insecticide control. Current Opinion in Insect Science
[Internet]. 2019 Mar 28; doi:10.1016/j.cois.2019.03.005

18. Hemingway J, Hawkes NJ, McCarroll L, Ranson H. The molecular basis of insecticide
resistance in mosquitoes. Insect Biochem Mol Biol. 2004 Jul;34(7):653–65.
doi:10.1016/j.ibmb.2004.03.018

19. Moiroux N, Gomez MB, Pennetier C, Elanga E, Djenontin A, Chandre F, et al. Changes
in Anopheles funestus Biting Behavior Following Universal Coverage of Long-Lasting
Insecticidal Nets in Benin. Journal of Infectious Diseases. 2012 Sep 10;206(10):1622–9.
doi:10.1093/infdis/jis565

20. Moiroux N, Damien GB, Egrot M, Djenontin A, Chandre F, Corbel V, et al. Human
Exposure to Early Morning Anopheles funestus Biting Behavior and Personal Protection
Provided by Long-Lasting Insecticidal Nets. PLoS ONE. 2014 août;9(8):e104967.
doi:10.1371/journal.pone.0104967

21. Corbel V, Chabi J, Dabiré RK, Etang J, Nwane P, Pigeon O, et al. Field efficacy of a
new mosaic long-lasting mosquito net (PermaNet 3.0) against pyrethroid-resistant
malaria vectors: a multi centre study in Western and Central Africa. Malar J. 2010 Apr
27;9:113. doi:10.1186/1475-2875-9-113

22. Chandre F, Darriet F, Duchon S, Finot L, Manguin S, Carnevale P, et al. Modifications
of pyrethroid effects associated with kdr mutation in Anopheles gambiae. Med Vet
Entomol. 2000 Mar;14(1):81–8. doi:10.1046/j.1365-2915.2000.00212.x

23. Parker JEA, Angarita-Jaimes N, Abe M, Towers CE, Towers D, McCall PJ. Infrared
video tracking of Anopheles gambiae at insecticide-treated bed nets reveals rapid
decisive impact after brief localised net contact. Scientific Reports. 2015 Sep 1;5:13392.
doi:10.1038/srep13392

24. Parker JEA, Angarita Jaimes NC, Gleave K, Mashauri F, Abe M, Martine J, et al. Host-
seeking activity of a Tanzanian population of Anopheles arabiensis at an insecticide
treated bed net. Malaria Journal. 2017 Jul 4;16:270. doi:10.1186/s12936-017-1909-6

25. Diop MM, Moiroux N, Chandre F, Martin-Herrou H, Milesi P, Boussari O, et al.
Behavioral Cost & Overdominance in Anopheles gambiae. PLoS ONE. 2015 Apr
1;10(4):e0121755. doi:10.1371/journal.pone.0121755

26. Desneux N, Decourtye A, Delpuech J-M. The sublethal effects of pesticides on
beneficial arthropods. Annu Rev Entomol. 2007;52:81–106.
doi:10.1146/annurev.ento.52.110405.091440

27. Haynes KF. Sublethal effects of neurotoxic insecticides on insect behavior. Annu Rev
Entomol. 1988;33:149–68. doi:10.1146/annurev.en.33.010188.001053

28. Glunt KD, Coetzee M, Huijben S, Koffi AA, Lynch PA, N'Guessan R, et al. Empirical
and theoretical investigation into the potential impacts of insecticide resistance on the
effectiveness of insecticide-treated bed nets. Evolutionary Applications.
2018;11(4):431–41. doi:10.1111/eva.12574

29. L'utilisation des animaux à des fins scientifiques [Internet]. Ministère de l'Enseignement
supérieur, de la Recherche et de l'Innovation. [cited 2019 Oct 11]. Available from:
//www.enseignementsup-recherche.gouv.fr/cid70597/l-utilisation-des-animaux-a-des-
fins-scientifiques.html

30. Alout H, Ndam NT, Sandeu MM, Djégbe I, Chandre F, Dabiré RK, et al. Insecticide
Resistance Alleles Affect Vector Competence of Anopheles gambiae s.s. for
Plasmodium falciparum Field Isolates. PLOS ONE. 2013 May 21;8(5):e63849.
doi:10.1371/journal.pone.0063849

31. N'Guessan R, Corbel V, Akogbéto M, Rowland M. Reduced efficacy of insecticide-
treated nets and indoor residual spraying for malaria control in pyrethroid resistance
area, Benin. Emerging Infect Dis. 2007 Feb;13(2):199–206.
doi:10.3201/eid1302.060631

32. Berticat C, Boquien G, Raymond M, Chevillon C. Insecticide resistance genes induce a
mating competition cost in Culex pipiens mosquitoes. Genet Res. 2002 Feb;79(1):41–7.
doi:10.1017/s001667230100547x

33. Sawadogo SP, Diabaté A, Toé HK, Sanon A, Lefevre T, Baldet T, et al. Effects of age
and size on Anopheles gambiae s.s. male mosquito mating success. J Med Entomol.
2013 Mar;50(2):285–93. doi:10.1603/me12041

34. Jones MDR, Gubbins SJ. Changes in the circadian flight activity of the mosquito
Anopheles gambiae in relation to insemination, feeding and oviposition. Physiological
Entomology. 1978 Sep 1;3(3):213–20. doi:10.1111/j.1365-3032.1978.tb00151.x

35. Jones JC, Madhukar BV. Effects of sucrose in blood avidity in mosquitoes. Journal of
Insect Physiology. 1976;22(3):357–60. doi:10.1016/0022-1910(76)90001-9

36. Skovmand O, Bonnet J, Pigeon O, Corbel V. Median knock-down time as a new method
for evaluating insecticide-treated textiles for mosquito control. Malaria Journal.
2008;7(1):114. doi:10.1186/1475-2875-7-114

37. Briegel H, Lea AO, Klowden MJ. Hemoglobinometry as a Method for Measuring Blood
Meal Sizes of Mosquitoes (Diptera: Culicidae)1. Journal of Medical Entomology. 1979
Mar 23;15(3):235–8. doi:10.1093/jmedent/15.3.235

38. Briegel H. Fecundity, metabolism, and body size in Anopheles (Diptera: Culicidae),
vectors of malaria. J Med Entomol. 1990 Sep;27(5):839–50.
doi:10.1093/jmedent/27.5.839

39. R Development Core Team. R: A Language and Environment for Statistical Computing.
3.5. Vienna, Austria: R Foundation for Statistical Computing; 2018.

40. Brooks ME, Kristensen K, Benthem KJ van, Magnusson A, Berg CW, Nielsen A, et al.
glmmTMB Balances Speed and Flexibility Among Packages for Zero-inflated
Generalized Linear Mixed Modeling. The R Journal. 2017;9(2):378–400.
doi:10.32614/RJ-2017-066

41. Therneau TM. coxme: Mixed Effects Cox Models [Internet]. 2018. Available from:
https://CRAN.R-project.org/package=coxme

42. Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models Using
lme4. Journal of Statistical Software. 2015 Oct 7;67(1):1–48. doi:10.18637/jss.v067.i01

43. Firth D. Bias Reduction of Maximum Likelihood Estimates. Biometrika. 1993;80(1):27–
38. doi:10.2307/2336755

44. Kosmidis I. brglm2: bias reduction in generalized linear models [Internet]. 2019.
Available from: https://CRAN.R-project.org/package=brglm2

45. Lenth R, Singmann H, Love J, Buerkner P, Herve M. emmeans: Estimated Marginal
Means, aka Least-Squares Means [Internet]. 2019. Available from: https://CRAN.R-
project.org/package=emmeans

46. Harrell FE, with contributions from Charles Dupont and many others. Hmisc: Harrell
Miscellaneous   [Internet].   2019.   Available   from: https://CRAN.R-
project.org/package=Hmisc

47. Moiroux N. Code and data to reproduce analyses in Diop et al. 2019. 2020 Jan 28 [cited
2020 Feb 4]; doi:10.5281/zenodo.2616599

48. Lyimo IN, Haydon DT, Mbina KF, Daraja AA, Mbehela EM, Reeve R, et al. The fitness
of African malaria vectors in the presence and limitation of host behaviour. Malar J.
2012 Dec 19;11:425. doi:10.1186/1475-2875-11-425

49. Fernandes L, Briegel H. Reproductive physiology of Anopheles gambiae and Anopheles
atroparvus. Journal of vector ecology. 2005;30(1):11.

50. Hauser G, Thiévent K, Koella JC. The ability of Anopheles gambiae mosquitoes to bite
through a permethrin-treated net and the consequences for their fitness. Sci Rep
[Internet]. 2019 May 31 [cited 2019 Oct 10];9. doi:10.1038/s41598-019-44679-1

51. Walker ED, Edman JD. The influence of host defensive behavior on mosquito (Diptera:
Culicidae)   biting   persistence.   J   Med   Entomol.   1985   Jul   26;22(4):370–2.
doi:10.1093/jmedent/22.4.370

52. Roitberg BD, Mondor EB, Tyerman JGA. Pouncing spider, flying mosquito: blood
acquisition increases predation risk in mosquitoes. Behavioral Ecology. 2003 Sep
1;14(5):736–40. doi:10.1093/beheco/arg055

53. Scott TW, Takken W. Feeding strategies of anthropophilic mosquitoes result in
increased risk of pathogen transmission. Trends in Parasitology. 2012 Mar;28(3):114–
21. doi:10.1016/j.pt.2012.01.001

54. Vézilier J, Nicot A, Gandon S, Rivero A. Insecticide resistance and malaria
transmission: infection rate and oocyst burden in Culex pipiens mosquitoes infected with
Plasmodium relictum. Malar J. 2010 Dec 31;9:379. doi:10.1186/1475-2875-9-379

55. Sattelle DB, Yamamoto D. Molecular Targets of Pyrethroid Insecticides. In: Evans PD,
Wigglesworth VB, editors. Advances in Insect Physiology [Internet]. Academic Press;
1988. p. 147–213. doi:10.1016/S0065-2806(08)60025-9

56. Wouters W, van den Bercken J. Action of pyrethroids. General Pharmacology: The
Vascular System. 1978 Jan 1;9(6):387–98. doi:10.1016/0306-3623(78)90023-X

57. O'Reilly AO, Khambay BPS, Williamson MS, Field LM, WAllace BA, Davies TGE.
Modelling insecticide-binding sites in the voltage-gated sodium channel. Biochemical
Journal. 2006 Jun 1;396(2):255. doi:10.1042/BJ20051925

58. Corbel V, Chandre F, Brengues C, Akogbeto M, Lardeux F, Hougard JM, et al. Dosage-
dependent effects of permethrin-treated nets on the behaviour of Anopheles gambiae and
the selection of pyrethroid resistance. Malaria journal. 2004 Jul 8;3:22.
doi:10.1186/1475-2875-3-22

59. Briegel H, Rezzonico L. Concentration of host blood protein during feeding by
anopheline mosquitoes (Diptera: Culicidae). J Med Entomol. 1985 Nov 27;22(6):612–8.
doi:10.1093/jmedent/22.6.612

60. Lahondère C, Lazzari CR. Mosquitoes cool down during blood feeding to avoid
overheating. Curr Biol. 2012 Jan 10;22(1):40–5. doi:10.1016/j.cub.2011.11.029

61. Lahondère C, Lazzari CR. Thermal Stress and Thermoregulation During Feeding in
Mosquitoes. In: New insights into malaria vectors [Internet]. Sylvie Manguin; 2013
[cited   2015   Feb   16].   p.   17.   Available   from:
http://cdn.intechopen.com/pdfs/44039/InTech-
Thermal_stress_and_thermoregulation_during_feeding_in_mosquitoes.pdf

62. Raphemot R, Rouhier MF, Hopkins CR, Gogliotti RD, Lovell KM, Hine RM, et al.
Eliciting Renal Failure in Mosquitoes with a Small-Molecule Inhibitor of Inward-
Rectifying Potassium Channels. PLoS ONE. 2013 May 29;8(5):e64905.
doi:10.1371/journal.pone.0064905

63. Hougard J-M, Duchon S, Darriet F, Zaim M, Rogier C, Guillet P. Comparative
performances, under laboratory conditions, of seven pyrethroid insecticides used for
impregnation of mosquito nets. Bull World Health Organ. 2003;81(5):324–33.

64. N'Guessan R, Asidi A, Boko P, Odjo A, Akogbeto M, Pigeon O, et al. An experimental
hut evaluation of PermaNet(®) 3.0, a deltamethrin-piperonyl butoxide combination net,
against pyrethroid-resistant Anopheles gambiae and Culex quinquefasciatus mosquitoes
in southern Benin. Trans R Soc Trop Med Hyg. 2010 Dec;104(12):758–65.
doi:10.1016/j.trstmh.2010.08.008

65. Toe KH, Müller P, Badolo A, Traore A, Sagnon N, Dabiré RK, et al. Do bednets
including piperonyl butoxide offer additional protection against populations of
Anopheles gambiae s.l. that are highly resistant to pyrethroids? An experimental hut
evaluation in Burkina Fasov. Medical and Veterinary Entomology. 2018;32(4):407–16.
doi:10.1111/mve.12316

66. Toé KH, Jones CM, N'Fale S, Ismail HM, Dabiré RK, Ranson H. Increased pyrethroid
resistance in malaria vectors and decreased bed net effectiveness, Burkina Faso.
Emerging Infect Dis. 2014 Oct;20(10):1691–6. doi:10.3201/eid2010.140619

67. Porciani A, Diop M, Moiroux N, Kadoke-Lambi T, Cohuet A, Chandre F, et al.
Influence of pyrethroïd-treated bed net on host seeking behavior of Anopheles gambiae

s.s. carrying the kdr allele. PLOS ONE. 2017 juil;12(7):e0164518.
doi:10.1371/journal.pone.0164518

68. Alout H, Yameogo B, Djogbenou LS, Chandre F, Dabire RK, Corbel V, et al. Interplay
Between Plasmodium Infection and Resistance to Insecticides in Vector Mosquitoes.
The Journal of infectious diseases [Internet]. 2014 May 14; doi:10.1093/infdis/jiu276

69. Alout H, Djegbe I, Chandre F, Djogbenou LS, Dabire RK, Corbel V, et al. Insecticide
exposure impacts vector-parasite interactions in insecticide-resistant malaria vectors.
Proceedings Biological sciences / The Royal Society [Internet]. 2014 Jul 7;281(1786).
doi:10.1098/rspb.2014.0389

70. Cator LJ, George J, Blanford S, Murdock CC, Baker TC, Read AF, et al. "Manipulation"
without the parasite: altered feeding behaviour of mosquitoes is not dependent on
infection with malaria parasites. Proceedings of the Royal Society B: Biological
Sciences. 2013 May 22;280(1763):20130711–20130711. doi:10.1098/rspb.2013.0711

71. Cator LJ, Lynch PA, Read AF, Thomas MB. Do malaria parasites manipulate
mosquitoes? Trends Parasitol. 2012 Nov;28(11):466–70. doi:10.1016/j.pt.2012.08.004

72. Vantaux A, Yao F, Hien DF, Guissou E, Yameogo BK, Gouagna L-C, et al. Field
evidence for manipulation of mosquito host selection by the human malaria parasite,
Plasmodium falciparum. bioRxiv. 2018 Nov 8;207183. doi:10.1101/207183

73. Lefevre T, Ohm J, Dabiré KR, Cohuet A, Choisy M, Thomas MB, et al. Transmission
traits of malaria parasites within the mosquito: Genetic variation, phenotypic plasticity,
and consequences for control. Evolutionary Applications. 2018 Apr 1;11(4):456–69.
doi:10.1111/eva.12571

74. Sherrard-Smith E, Griffin JT, Winskill P, Corbel V, Pennetier C, Djénontin A, et al.
Systematic review of indoor residual spray efficacy and effectiveness against
Plasmodium falciparum in Africa. Nature Communications. 2018 Nov 26;9(1):4982.
doi:10.1038/s41467-018-07357-w

Supplementary Figures:

Supplementary Figure 1: Feeding success after exposure to insecticides of non-KD

***Anopheles gambiae* females of each *kdr* genotype.**

Feeding success of non-KD SS, RS, and RR (panels A, B and C, respectively) genotypes when exposed

to untreated, permethrin-treated (Olyset) and deltamethrin-treated (PermaNet) nettings are shown

with 95% binomial confidence intervals of the proportions (error bars). Numbers n of mosquitoes

exposed to each treatment and for each genotype are indicated. P-values according to Tukey's test

after binomial mixed-effect model is indicated.

Supplementary Figure 2: Among *kdr* **genotypes comparison of feeding success of**
Anopheles gambiae **females after exposure to insecticides**.

Feeding success of each genotype and 95% binomial confidence intervals of the proportions (error

bars) are shown after exposure to untreated net (Panel A, same as Figure 2A), to permethrin treated

net (Olyset; panel B) and to deltamethrin treated net (Permanet; panel C). Numbers n of mosquitoes

exposed to each treatment and for each genotype are indicated. Significance according to Tukey's test

after binomial mixed-effect model is indicated.

Supplementary Figure 3: Among *kdr* genotypes comparison of feeding success of non-KD *Anopheles gambiae* females after exposure to insecticides.

Feeding success of non-KD mosquitoes of each genotype and 95% binomial confidence intervals of the proportions (error bars) are shown after exposure to permethrin treated net (Olyset; panel A) and to deltamethrin treated net (Permanet; panel B). Numbers n of mosquitoes exposed to each treatment and for each genotype are indicated. Significance according to Tukey's test after binomial mixed-effect model is indicated.