Sub-lethal insecticide exposure affects host biting efficiency of *Kdr*-resistant

Anopheles gambiae

Diop Malal M [1], Chandre Fabrice [1], Rossignol Marie[1], Porciani Angélique[1], Chateau

Mathieu[1], Moiroux Nicolas[1,2*], Pennetier Cédric[1,3*]

[1] MIVEGEC, Univ Montpellier, CNRS, IRD, Montpellier, France

[2] Institut de Recherche en Sciences de la Santé (IRSS), Bobo-Dioulasso, Burkina Faso

[3] Institut Pierre Richet (IPR), Bouaké, Côte d'Ivoire

*contributed equally to this work

Contacts & correspondence: Nicolas Moiroux nicolas.moiroux@ird.fr ; Cédric Pennetier

cedric.pennetier@ird.fr

Abstract

The massive use of insecticide-treated nets (ITNs) has drastically changed the environment

for malaria vector mosquitoes, challenging their host-seeking behaviour and biting success.

Here, we investigated the effect of a brief exposure to an ITN on the biting behaviour of

Anopheles gambiae mosquitoes and the interaction between such behaviour and the *kdr*

mutation that confers resistance to pyrethroids. To this aim, we developed a video assay to

study the biting behaviour of mosquitoes with similar genetic background, but different *kdr*

locus genotypes (SS i.e. homozygous susceptible, RS i.e. heterozygous and RR i.e.

homozygous resistant), after a brief exposure to either control untreated nets or one of two

types of pyrethroid-treated nets (deltamethrin or permethrin). In presence of untreated nets,

the *kdr* mutation did not influence mosquito blood feeding success but caused differences in

feeding and prediuresis durations and blood meal size. Exposure to deltamethrin ITN

decreased the blood feeding success rate of RR and RS mosquitoes, whereas permethrin ITN

increased the blood-feeding success of RR mosquitoes. Exposure to the two types of

pyrethroid-treated nets reduced feeding duration, prediuresis duration and blood meal size of

all three genotypes. We discussed the potential consequences of the observed behavioural

changes on malaria vector fitness and disease transmission.

Keywords: pyrethroid insecticides, long-lasting treated nets, malaria, probing, blood meal,

prediuresis*;*

30 Introduction

Malaria vector mosquitoes become infected by and then can transmit *Plasmodium spp.*

parasites during blood meals. *Plasmodium spp.* transmission is strongly dependent on the

mosquito biting rate on humans, as formalized by the Ross-MacDonald model of malaria [1].

The behaviour and host preferences of blood-feeding mosquitoes are influenced by several

factors, primarily their genetic background [2–5], but also environmental factors, such as host

diversity and availability [6,7], and the presence/absence of physical and chemical barriers,

such as Insecticide-Treated Nets (ITNs) [8–10].

ITNs should hamper the contacts between humans and nocturnal, anthropophilic and

endophagic mosquitoes, such as *An. gambiae* [11]. The widespread use of pyrethroid (PYR)

insecticides in public health (i.e. ITNs or indoor residual sprayings), crop protection and other

selective pressures (such as xenobiotics) drove adaptations in malaria vectors to reduce the

insecticidal effect [12–16].

Physiological PYR resistance involves two main mechanisms: (i) metabolic resistance, due to

quantitative or qualitative changes in detoxification enzymes (cytochrome P450

monooxygenases, esterases and glutathione S-transferases), and (ii) target site resistance, due

to non-synonymous mutations in the voltage-gated sodium channels that are called knock-

down resistance (*kdr*) mutations [12,17].

Mosquitoes can reduce vector control tools efficacy also through behavioural adaptations. In

areas of sub-Saharan Africa where large-scale vector control programmes have been

implemented, several observations suggest that malaria vectors can avoid contacts with ITNs

or insecticides on walls by modulating their host-feeding activity. For instance, following the

large scale distribution of ITNs in Benin, some *An. funestus* populations started to feed

predominantly at dawn/early in the morning or in broad daylight [18,19], when most people

are outside the nets [19]. Behavioural modulation might also be influenced by physiological

resistance mechanisms. Indeed, experimental hut trials showed that in *An. gambiae*, indirect

behavioural indicators in response to ITN presence (deterrence and induced exophily) are

related to the physiological tolerance to insecticide [20]. Moreover, a study demonstrated that

kdr homozygous resistant mosquitoes have longer contacts with ITNs than homozygous

susceptible mosquitoes, which are more excited by PYR irritant effect [21]. Altogether, these

findings indicate that insecticide treatments could affect the behaviour of malaria vectors.

However, the effects of insecticide exposure and *kdr* mutations on the biting activity of *An.*

gambiae remain poorly investigated.

The last step of the mosquito host-seeking behaviour after reaching a host protected by an

ITN is biting for taking a blood meal. During the host-seeking phase and the penetration

through a hole in the net, mosquitoes can be exposed to sub-lethal doses of insecticide [22–

24]. Such doses do not cause death, but can have several physiological or behavioural effects

on host-seeking mosquitoes [25]. For instance, sub-lethal doses of insecticide can affect the

feeding and reproductive behaviour of some blood-sucking insects [26]. As the PYR

insecticide on ITNs acts on specific sites in the mosquito nervous system, it might alter some

physiological processes involved in the biting behaviour of malaria vectors. There is currently

scarce literature on this subject [27].

In the present study, we investigated whether pre-exposure to an ITN modulates the mosquito

ability to take a blood meal by using experimental conditions that mirror the exposure to

insecticide occurring when a mosquito passes through an ITN after having located a host. We

also assessed whether the *kdr* mutation (L1014F) modifies blood feeding success and biting

behaviour of *An. gambiae*.

Methodology

Ethical Considerations

79   This study was carried out in strict accordance with the recommendations of Animal Care and

80   Use Committee named "Comité d'éthique pour l'expérimentation animale; Languedoc

81   Roussillon" and the protocol was approved by the Committee on the Ethics of Animal

82   Experiments (CEEA-LR-13002 for the rabbits).

Mosquito strains and rearing procedures

Two mosquito laboratory strains were used for this study. One is the insecticide-susceptible

Kisumu strain (KISUMU1, MRA-762, VectorBase stable ID VBS0000026 on

vectorbase.org), isolated in Kenya in 1975. This strain is PYR-susceptible and homozygous

(SS) for the L1014 codon. The second one is the Kdrkis strain that is PYR-resistant and

homozygous (RR) for the L1014F *kdr* mutation. The Kdrkis strain was obtained by

introgression of the *kdr*-west allele (L1014F) into the Kisumu genome through 19 successive

back-crosses between Kisumu and VKPer [28]. VKPer strain used to obtain Kdrkis displayed

the same expression level of metabolic resistance enzyme as Kisumu [29].

Polymorphisms between Kisumu and KdrKis strains are expected to be restricted in the

flanking region of the *Kdr* allele (15 cM for 19 backcrossing generations [30,31]) and the

observed phenotypes are therefore expected to be associated to this genetic area.

Heterozygous individuals (RS) for the L1014F *kdr* mutation were obtained by crossing once

Kisumu SS females (F1 progeny) with Kdrkis RR males. Therefore, the three genotypes have

a common genetic background for most of their genome.

Mosquitoes were reared at $27 \pm 1°C$, 70-80% relative humidity under a 12h:12h (light : dark)

photoperiod in the insectary. Gravid females were allowed to lay eggs on wet filter paper

inside mesh-covered cages. Eggs were dispensed in plastic trays containing osmotic water.

Larvae were kept in trays and fed with TetraMin® fish food. Pupae were removed and allowed

to emerge inside 30x30x30cm cages. After emergence, adults could feed *ad libitum* on 10%

sucrose solution.

On each experimental day, a batch of ten adult (7 to 9-day-old) female mosquitoes never fed

with blood were randomly collected from the rearing cages and placed in cups covered by

gauze. Mosquitoes were starved the day before the experiment because sucrose inhibits blood

avidity in mosquitoes [32].

Insecticide exposure

To simulate the contact with an ITN that may occur before the vector finds a way to reach a

host, starved females were individually exposed to PYR insecticide-treated netting using an

experimental setup in which insects can only walk or stand on the net surface and that was

initially designed to measure the median knock down time [33]. We used three net types (i.e.

treatments): an untreated net (negative control), an Olyset Net® (impregnated with

1000 mg/m^2 of permethrin, hereafter Olyset) and a PermaNet® 2.0 (coated with 55 mg/m^2 of

deltamethrin, hereafter PermaNet). Based on previous experiments [24], each mosquito was

exposed for 30 seconds. This is the median time of contact with permethrin-treated netting

before PYR-susceptible anopheles (SS genotype) locate a hole to reach the host [24]. After

exposure, a 1-min latency period was observed before releasing the insect in the behavioural

assay setup.

Behavioural assay

Experiments were conducted in a biting behavioural assay setup designed for video recording

the biting behaviour of the tested mosquitoes (Figure 1). The setup is made of a foam board

and composed of an observation tunnel (OT) and an observation zone (OZ) separated by a

transparent plastic (TP). The OZ is a triangular prism, closed on its base by a removable paper

sheet with a 1-cm diameter hole. The hole allows mosquitoes to bite the ear of a rabbit (R)

that is maintained immobile in a restraining cage (RC) to limit ear movement during blood

feeding. The same rabbit was used during all experiments.The experimental room was faintly

illuminated with a compact fluorescent lamp bulb placed at 15 cm from the OZ to allow the

acquisition of the biting behaviour sequence by the digital video system.

For each trial, a mosquito was individually released inside the behavioural assay through a

circular opening (CO) located on its lateral face. Cotton was used to plug the CO after

releasing. The number of released individual anopheles per genotype and treatment ranged

from 43 to 86. Each mosquito was filmed for 10 minutes using a Sony® Digital HD Video

Camera (HDR-XR550) placed on the top of the observational tunnel (Figure 1). The video

camera was connected to a computer screen placed outside the experimental room to allow

real-time monitoring of each mosquito, to avoid any disturbance of the rabbit and interference

of the experimenter on the mosquito behaviour. The MPEG-2 recordings (PAL video:

720x576 pixels at 25 frames/s) were analysed using the behavioural event recording program

EVENT01, version 1.2.4 (©R.D. Collins and M.K. Tourtellot, 1993-2002).

The experimenter used latex gloves to avoid any contamination by human skin odours .The

OZ was cleaned with ethanol and the removable paper was changed after each mosquito was

tested to avoid any contamination by insecticide residues between behavioural assays.

Feeding success and behavioural parameters

A mosquito was scored as successful if it was feed at the end of the 10-min trial (whatever the

amount of blood it took) and unsuccessful if it did not. After the trial, successfully fed

mosquitoes were stored at -35°C for measuring the blood meal size (see below). As pre-

exposure to insecticide can induce a knockdown (KD) effect during the trial, mosquitoes were

recorded as KD, if they were lying on their side or their back.

Analysis of the acquired images allowed quantifying the following variables in fed

mosquitoes: (1) number of probing events, (2) probing duration (the time from the

introduction of the stylet fascicule into the rabbit skin to blood appearance in the mosquito

abdomen), (3) feeding duration (from the blood appearance in the abdomen to the beginning

of the stylet fascicule withdrawal), and (4) prediuresis duration (the time during which

excretion of rectal fluid (plasma, water, metabolic wastes) is observed as red bright drops

during feeding and after proboscis withdrawal).

Blood meal volume measurement

The blood intake was evaluated by quantifying the haemoglobin amount, as described by

Briegel [34]. Each engorged mosquito was stored in one 1.5 ml Eppendorf tube at -35°C.

Then, the whole abdomen was ground in the presence of 0.5 ml of Drabkin's reagent until it

was completely disintegrated. Haemoglobin then reacted with the Drabkin's reagent and was

converted into haemoglobin cyanide (HiCN). Samples were incubated at room temperature

(25°C) for 20 min and then a chloroform solution was added in each tube. Samples were

centrifuged at 5600 rpm for 5 min and the aqueous supernatant (containing HiCN) was placed

in a new 1.5 ml Eppendorf tube. The absorbance was read at a wavelength of 550 nm using a

microplate spectrophotometer. Two replicates were done for each mosquito and their

absorbance values were averaged. A sample of the rabbit blood was used as control for

calibration curves.

As the blood meal volume is correlated with the mosquito size [35], the blood meal volume of

all mosquitoes from the same batch was divided by the average weight of five randomly

selected mosquitoes from the same rearing cage. The resulting ratio was expressed in µL of

blood per µg of weight and called weighted blood meal volume or blood meal size.

Statistical analysis

All statistical analyses were performed using the R software, version 3.5 [36]. We analysed

the feeding success (coded as 1 for fed mosquitoes and 0 for unfed ones) with a binomial

logistic mixed-effect model using 'glmer' function in the 'lme4' package [37]. We analysed

the number of probing events with a zero-truncated negative binomial mixed-effect model

using function 'glmmTMB' in the 'glmmTMB' package [38]. We analysed durations

(probing, feeding and prediuresis) with a mixed effect Cox proportional hazard model using

function 'coxme' of the 'coxme' package [39]. We analysed weighted blood meal size with a

linear mixed effect model using function 'lmer' in the 'lme4' package. All models included

the *kdr* genotypes (SS, RS or RR), type of pre-exposure (untreated, Olyset or PermaNet) and

their interactions as fixed terms explanatory variables and the date of the experiment as a

random intercept. We used Tukey's post-hoc test to perform multiple comparisons among

genotypes and treatments using 'emmeans' function [40]. We computed Odds Ratios (OR) for

the binomial model, Hazard Ratios (HR) for Cox models, Rate Ratios (RR) for the negative

binomial model, Mean Differences (MD) for the linear model and their 95% confidence

intervals.

We calculated the binomial confidence interval of feeding rates and knock-down rates with

the Wilson's score method using the 'binconf' command from the 'Hmisc' package [41].

Effect of feeding duration on blood meal size and prediuresis duration was analysed using a

linear mixed effect model and a mixed effect cox proportional hazard model, respectively.

Fixed terms included genotypes and treatment and interactions and date was set as a random

intercept. We used function 'emtrends' of the 'emmeans' package [40] to obtain estimates and

95% confidence intervals of the marginal slopes of the trends (between feeding duration and

blood meal size or prediuresis duration) for each genotypes and treatments.

Data and codes used for analyses and figures are available [42].

Results

In total, 511 *An. gambiae* females (182 SS, 156 RS and 173 RR) were released individually

for the study. The number of mosquitoes released, fed, unfed and knock-downed among

genotypes and treatments are shown in Table 1.

Table 1. Numbers of mosquito releases per genotype and treatment

Genotype	Treatment	N tested	Fed		Unfed	
			KD	Non-KD	KD	Non-KD
SS	Untreated	86	0	41	0	45
SS	Permethrin	50	5	5	33	7
SS	Deltamethrin	46	1	15	18	12
RS	Untreated	70	0	40	0	30
RS	Permethrin	43	3	14	11	15
RS	Deltamethrin	43	1	9	15	18
RR	Untreated	70	0	40	0	30
RR	Permethrin	53	0	41	0	12
RR	Deltamethrin	50	4	12	12	22

SS: homozygous susceptible genotype, RS: heterozygous genotype, RR: homozygous
resistant genotype, KD: Knockdown.

Impact of the kdr mutation on feeding success and biting behaviour

When female mosquitoes were pre-exposed to UTN (i.e., untreated netting), no difference in
their feeding success was found among the three genotypes (OR_{RS-SS}=1.47 [0.67, 3.22]; OR_{RR-SS}=1.47 [0.68, 3.19]; OR_{RR-RS}=1.00 [0.44, 2.25]; Figure 2A).

Analysis of the biting behaviour of successful mosquitoes in absence of insecticide pre-
exposure showed that feeding and prediuresis durations were shorter in RR than in both SS
and RS mosquitoes (feeding duration: $HR_{RR-SS} = 2.15$ [1.26, 3.69]; $HR_{RR-RS} = 2.46$ [1.43,
4.23]; Figure 2B; prediuresis duration: $HR_{RR-SS} = 1.94$ [1.02, 3.71]; $HR_{RR-RS} = 2.11$ [1.11,
4.01]; Figure 2C). The weighted blood meal volume of RR mosquitoes was lower than that of
RS mosquitoes ($MD_{RS-RR} = 0.99$ [0.24, 1.74]; Figure 2D)). Number of probing events and
probing duration were not significantly different among genotypes (SS, RS and RR) (Tukey's
p-value > 0.05).

Impact of insecticide exposure on knockdown rates

Pre-exposure to permethrin or deltamethrin induced 76 % [62.6, 85.7] and 41.3% [28.3, 55.7]
knock-down rates in SS mosquitoes, respectively. It induced 32.6% [20.5, 47.5] (14/43) and

37.2% [24.4, 52.1] (14/43) KD rates in RS mosquitoes, respectively (Table 1). Among RR

mosquitoes, permethrin pre-exposure did not induce any KD effect, whereas deltamethrin pre-

exposure led to 30.2% [20.8, 45.8] (16/50) of KD mosquitoes (Table 1).

Impact of insecticide pre-exposure on feeding success

When compared to the UTN condition, pre-exposure to permethrin (Olyset) reduced

significantly the feeding success of SS mosquitoes ($OR_{perm-UTN}$=0.28 [0.10, 0.73]; Figure 3A),

but not that of RS ($OR_{perm-UTN}$=0.49 [0.19, 1.26]; Figure 3B) or RR mosquitoes (OR_{perm-}

$_{UTN}$=2.58 [0.97, 6.84]; Figure 3B). Pre-exposure to deltamethrin (PermaNet) reduced

significantly the feeding success of RS and RR mosquitoes ($OR_{delta-UTN}$=0.23 [0.08, 0.64] and

$OR_{delta-UTN}$=0.35 [0.14 – 0.88]; Figure 3B and 3C), but not of SS mosquitoes ($OR_{delta-UTN}$=0.58

[0.24, 1.43]; Figure 3A).

When comparing the feeding success among genotypes after insecticide pre-exposure, the

feeding rate of RR mosquitoes was higher than that of SS and RS mosquitoes after permethrin

pre-exposure (OR_{RR-SS}=13.82 [4.35, 43.92]; OR_{RR-RS}=5.29 [1.76, 15.86], Supplementary

Figure 1B), whereas the feeding success of RS mosquitoes was not different than that of SS

mosquitoes (OR_{RS-SS}=2.61 [0.85, 8.02], Supplementary Figure 1B). In contrast, pre-exposure

to deltamethrin did not induce any difference in the feeding success of the three genotypes

(OR_{RR-SS}=0.891 [0.31, 2.53]; OR_{RR-RS}=1.55 [0.51, 4.76]; OR_{RS-SS}=0.57 [0.18, 1.80],

Supplementary Figure 1C).

Impact of insecticide exposure on biting behaviour

After exposure to deltamethrin (PermaNet), feeding duration, prediuresis duration and

weighted blood meal size were significantly reduced in SS mosquitoes compared to the UTN

condition (feeding duration: $HR_{delta-UTN}$ = 4.38 [2.09, 9.15]; prediuresis: $HR_{delta-UTN}$ = 5.31

[2.11, 13.36]; blood meal size; $MD_{delta-UTN}$ = -1.02 [-1.82, -0.23]; Figure 4A, 4B and 4C,

respectively). A similar trend was observed after exposure to permethrin (Olyset) for feeding

duration ($HR_{perm-UTN}$ = 2.87 [1.21, 6.81]; Figure 4A) and weighted blood meal volume

($MD_{perm-UTN}$ = -0.97 [-1.85, -0.08]; Figure 4C) but not for prediuresis duration ($HR_{perm-UTN}$ =

2.27 [0.92, 5.63]; Figure 4B).

Compared to UTN, permethrin and deltamethrin reduced the feeding and prediuresis

durations as well as blood meal size of both RS (feeding duration: $HR_{perm-UTN}$ = 7.42 [3.55,

15.53], $HR_{delta-UTN}$ = 4.57 [1.94, 10.76], Figure 4D; prediuresis duration: $HR_{perm-UTN}$ = 9.44

[3.36, 26.46] and $HR_{delta-UTN}$ = 7.22 [2.70, 19.29], Figure 4E, blood-meal size: $MD_{perm-UTN}$ = -

1.97 [-3.04, -0.90] and $MD_{delta-UTN}$ = -2.01 [-2.97, -1.05], Figure 4F) and RR genotypes

(feeding duration: $HR_{perm-UTN}$ = 2.06 [1.20, 3.51] and $HR_{delta-UTN}$ = 3.78 [1.83, 7.82], Figure

4G; prediuresis duration: $HR_{perm-UTN}$ = 1.71 [0.80, 3.64] and $HR_{delta-UTN}$ = 9.31 [3.57, 24.28],

Figure 4H; blood-meal size: $MD_{perm-UTN}$ = -1.52 [-2.23, -0.81] and $MD_{delta-UTN}$ = -1.41 [-2.28,

-0.54], Figure 4I). Number of probing events and probing duration were not significantly

different among treatments (UTN and ITN) (Tukey's p-value > 0.05).

In absence of pre-exposure to insecticide (UTN), the blood meal size and the prediuresis

duration were positively correlated with the feeding duration for all genotypes (supplementary

table 1 and 2). With the exception of blood meal size of RS pre-exposed to permethrin, these

relationships were not observed when mosquitoes were pre-exposed to insecticides

(supplementary table 1 and 2), suggesting a perturbation of processes underlying these

correlations.

Discussion

To investigate the influence of ITN exposure on the biting behaviour of *An. gambiae* mosquitoes, we used mosquitoes that share the same genetic background, but for the *kdr* allele locus, and then exposed them to ITN prior to blood feeding.

The blood-feeding success did not differ between the three genotypes in the absence of insecticide exposure. Therefore, the *kdr* mutation was not associated with significant change of blood meal success rate. However, feeding duration and blood meal size were different between genotypes. RR mosquitoes spent less time taking their blood meal than RS and SS mosquitoes. This might confer an advantage as fast feeding reduces the risk to be killed because of the host defensive behaviour [43]. On the other hand, RS mosquitoes took higher blood volumes than RR females. This could improve the completion of oogenesis in RS mosquitoes [44] and increase their fecundity compared to RR. However, large blood meals reduce the flying ability, escape speed and agility required to avoid predators [43,45,46]. These different trades-offs between behavioural traits that might enhance fecundity or survival in the three genotypes are of great interest and deserve further investigations in relation with the ecological and vector control environment. Such trades-offs possibly affect mosquito fitness and may therefore drive not only the evolution of insecticide resistance in mosquitoes but also parasite transmission. For example, a decrease in blood meal duration and size might increase the frequency of multiple feedings and consequently the risk of *Plasmodium* transmission [47]. Similarly, a bigger blood meal size might increase the probability of mosquito infection by gametocytes [48].

Exposure to permethrin and deltamethrin induced opposite outcomes in term of blood feeding success (increase and decrease, respectively) in RR mosquitoes. This opposite effect on the feeding success rate of RR females might be linked to the different chemical properties of

permethrin and deltamethrin that induce two types of bursting activity of sodium channels

[49,50]. Type II pyrethroids, such as deltamethrin, further delay the inactivation of the

voltage-gated sodium channel and in a less reversible way than type I pyrethroids, such as

permethrin [51]. The lower PYR susceptibility of homozygous resistant mosquitoes could

lead to their over-stimulation compared to susceptible and heterozygous mosquitoes that are

more affected by the toxic effect of such insecticides [52].

In contrast, among females that have been successful in taking a blood meal, the behavioural

sequence was altered in the same way by both insecticides. They both induced a decrease of

feeding duration and prediuresis duration. As discussed above, short feeding durations and

small blood-meal lead us to expect that exposure to insecticide may (i) reduce the risk of the

vector to be killed due to the defensive behaviour of the host or due to predators, (ii) reduce

the possible number of parasites ingested during one feeding attempt that may be

compensated by (iii) the increase of multiple feeding (and therefore the higher risk of human-

to-mosquito and mosquito-to-human parasite transmission).

Whatever the genotype, blood meal size and prediuresis duration were correlated to feeding

duration in absence of pre-exposure to insecticides. This relationship was, in most cases, no

longer observed when mosquitoes were pre-exposed to insecticides indicating that sub-lethal

contact with insecticides disrupt physiological processes involved in blood meal intake. Such

indirect evidence highlight the real need to decipher more deeply with consequences of sub-

lethal contact with insecticides.

Prediuresis duration was substantially reduced in RS and RR females after exposure to ITNs.

Prediuresis is an intestinal mechanism that plays a crucial role in protein concentration during

feeding [53] and contributes to thermoregulation [54,55]. This perturbation of the pre-

diuresisprediuresis phase by sub-lethal doses of PYR insecticides could lead to toxic

accumulation of metabolic wastes and products of oxidative stress in the haemolymph that

might affect the lifespan of mosquitoes. This results suggest that mosquitoes with the *kdr*

mutation might be more susceptible to new chemicals that target the mosquito renal system

and that are currently developed as an alternative to the currently used insecticides [56].

To conclude, our study demonstrates a complex interaction between insecticide exposure and

the *kdr* mutation on the biting behaviour of mosquitoes. The behavioural modulation induced

by PYR-treated nets also raises concerns about the consequences of the *kdr* resistance-

insecticide interaction. In previous studies, we evidenced that RR mosquitoes prefer a host

protected by a permethrin-treated net rather than an untreated net [31] and have a remarkable

ability to find a hole into a bet net [24]. Herein, we have completed the sequence by showing

that permethrin exposure enhances the feeding success of RR mosquitoes. This suggestsThe

whole picture suggests that permethrin ITN may increase vectorial capacity of *An. gambiae*

populations in areas where PYR resistance with *kdr* mutation is well established. Insecticide

resistance genes in malaria vectors could modify vector competence and the dynamics of

infection by *P. falciparum*. For instance, recent studies have shown that parasite infection

increases insecticide susceptibility in mosquitoes carrying the *kdr* mutation [57] and that

insecticide exposure reduces parasite development in resistant mosquitoes [58]. In addition,

malaria parasites have been shown to modify the feeding behaviour of their mosquito vectors

in ways that favour their transmission [59–61], but the role of insecticide resistance that could

modulate this phenomenon has not been investigated yet [62]. It is therefore urgent to

decipher the links between insecticide exposure, resistance mechanisms and infection by *P.*

falciparum on the host-seeking and biting behaviour of malaria vectors to better understand

malaria transmission in areas where insecticidal tools for malaria prevention are implemented.

All these interactions should then be used as variables to include host-seeking behavioural

modulation by *Kdr* resistance in models on *P. falciparum* transmission to better understand

and/or predict the efficacy of vector control strategies [63].

Acknowledgements

MD was supported by IRD's PhD student fellowship. The authors would like to acknowledge

the technical/research platform dedicated on vectors at the IRD centre and all members of the

platform for providing mosquitoes and technical support. This platform is a member of the

Vectopole Sud network and of the LabEX CEMEB (Centre Méditerranéen de

l'Environnement et de la Biodiversité) in Montpellier. We thank Dr Anna Cohuet and Dr

Thierry Lefèvre for their valuable comments on the manuscript.

References

1.   Garrett-Jones C, Shidrawi GR. Malaria vectorial capacity of a population of Anopheles
gambiae: an exercise in epidemiological entomology. Bulletin of the World Health
Organization. 1969;40(4):531.

2.   Lefèvre T, Gouagna L-C, Dabire KR, Elguero E, Fontenille D, Costantini C, et al.
Evolutionary lability of odour-mediated host preference by the malaria vector *Anopheles*
gambiae. Tropical Medicine & International Health. 2009 Feb;14(2):228–36.
doi:10.1111/j.1365-3156.2009.02206.x

3.   Dana AN, Hong YS, Kern MK, Hillenmeyer ME, Harker BW, Lobo NF, et al. Gene
expression patterns associated with blood-feeding in the malaria mosquito Anopheles
gambiae. BMC Genomics. 2005 Jan 14;6(1):5. doi:10.1186/1471-2164-6-5

4.   Rund SSC, Hou TY, Ward SM, Collins FH, Duffield GE. Genome-wide profiling of diel
and circadian gene expression in the malaria vector Anopheles gambiae. Proceedings of
the National Academy of Sciences. 2011 Jun 29;108(32):E421–30.
doi:10.1073/pnas.1100584108

5.   Zwiebel L., Takken W. Olfactory regulation of mosquito–host interactions. Insect
Biochemistry and Molecular Biology. 2004 Jul;34(7):645–52.
doi:10.1016/j.ibmb.2004.03.017

6.   Killeen GF, McKenzie FE, Foy BD, Bøgh C, Beier JC. The availability of potential
hosts as a determinant of feeding behaviours and malaria transmission by African
mosquito populations. Transactions of the Royal Society of Tropical Medicine and
Hygiene. 2001;95(5):469–476.

7. Lyimo IN, Ferguson HM. Ecological and evolutionary determinants of host species
choice in mosquito vectors. Trends in Parasitology. 2009 Apr;25(4):189–96.
doi:10.1016/j.pt.2009.01.005

8. Mathenge EM, Gimnig JE, Kolczak M, Ombok M, Irungu LW, Hawley WA. Effect of
permethrin-impregnated nets on exiting behavior, blood feeding success, and time of
feeding of malaria mosquitoes (Diptera: Culicidae) in western Kenya. Journal of medical
entomology. 2001;38(4):531–6.

9. Lefèvre T, Gouagna L-C, Dabiré KR, Elguero E, Fontenille D, Renaud F, et al. Beyond
nature and nurture: phenotypic plasticity in blood-feeding behavior of Anopheles
gambiae ss when humans are not readily accessible. The American journal of tropical
medicine and hygiene. 2009;81(6):1023–1029.

10. Gatton ML, Chitnis N, Churcher T, Donnelly MJ, Ghani AC, Godfray HCJ, et al. The
importance of mosquito behavioural adaptations to malaria control in Africa. Evolution.
2013;67(4):1218–1230.

11. Pates H, Curtis C. Mosquito behavior and vector control. Annu Rev Entomol.
2005;50:53–70.

12. Ranson H, N'Guessan R, Lines J, Moiroux N, Nkuni Z, Corbel V. Pyrethroid resistance
in African anopheline mosquitoes: what are the implications for malaria control? Trends
Parasitol [Internet]. 2011;27. doi:10.1016/j.pt.2010.08.004

13. Corbel V, N'Guessan R. Distribution, mechanisms, impact and management of
insecticide resistance in malaria vectors: a pragmatic review. 2013 [cited 2014 Dec 2];
Available from: http://www.intechopen.com/books/anopheles-mosquitoes-new-insights-
into-malaria-vectors/distribution-mechanisms-impact-and-management-of-insecticide-
resistance-in-malaria-vectors-a-pragmat

14. Chouaïbou MS, Fodjo BK, Fokou G, Allassane OF, Koudou BG, David J-P, et al.
Influence of the agrochemicals used for rice and vegetable cultivation on insecticide
resistance in malaria vectors in southern Côte d'Ivoire. Malaria Journal. 2016;15(1):426.

15. Liu N. Insecticide Resistance in Mosquitoes: Impact, Mechanisms, and Research
Directions. Annual Review of Entomology. 2015;60(1):537–59. doi:10.1146/annurev-
ento-010814-020828

16. Carrasco, D., Lefevre, T., Moiroux, N., Pennetier, C., Chandre, F., Cohuet, A.
Behavioural adaptations of mosquito vectors to insecticide control. Current Opinion in
Insect Science. 2019;Accepted.

17. Hemingway J, Hawkes NJ, McCarroll L, Ranson H. The molecular basis of insecticide
resistance in mosquitoes. Insect biochemistry and molecular biology. 2004;34(7):653–
65.

18. Moiroux N, Gomez MB, Pennetier C, Elanga E, Djenontin A, Chandre F, et al. Changes
in Anopheles funestus Biting Behavior Following Universal Coverage of Long-Lasting
Insecticidal Nets in Benin. Journal of Infectious Diseases. 2012 Sep 10;206(10):1622–9.
doi:10.1093/infdis/jis565

19.   Moiroux N, Damien GB, Egrot M, Djenontin A, Chandre F, Corbel V, et al. Human
409   Exposure to Early Morning Anopheles funestus Biting Behavior and Personal Protection
410   Provided by Long-Lasting Insecticidal Nets. PLoS ONE. 2014 août;9(8):e104967.
411   doi:10.1371/journal.pone.0104967

412   20.   Corbel V, Chabi J, Dabiré RK, Etang J, Nwane P, Pigeon O, et al. Field efficacy of a
413   new mosaic long-lasting mosquito net (PermaNet 3.0) against pyrethroid-resistant
414   malaria vectors: a multi centre study in Western and Central Africa. Malar J.
415   2010;9(113):113.

416   21.   Chandre F, Darriet F, Duchon S, Finot L, Manguin S, Carnevale P, et al. Modifications
417   of pyrethroid effects associated with kdr mutation in Anopheles gambiae. Medical and
418   Veterinary Entomology. 2000;14(1):81–8.

419   22.   Parker JEA, Angarita-Jaimes N, Abe M, Towers CE, Towers D, McCall PJ. Infrared
420   video tracking of Anopheles gambiae at insecticide-treated bed nets reveals rapid
421   decisive impact after brief localised net contact. Scientific Reports. 2015 Sep 1;5:13392.
422   doi:10.1038/srep13392

423   23.   Parker JEA, Angarita Jaimes NC, Gleave K, Mashauri F, Abe M, Martine J, et al. Host-
424   seeking activity of a Tanzanian population of Anopheles arabiensis at an insecticide
425   treated bed net. Malaria Journal. 2017 Jul 4;16:270. doi:10.1186/s12936-017-1909-6

426   24.   Diop MM, Moiroux N, Chandre F, Martin-Herrou H, Milesi P, Boussari O, et al.
427   Behavioral Cost & Overdominance in Anopheles gambiae. PLoS ONE. 2015 Apr
428   1;10(4):e0121755. doi:10.1371/journal.pone.0121755

429   25.   Desneux N, Decourtye A, Delpuech J-M. The sublethal effects of pesticides on
430   beneficial   arthropods.   Annu   Rev   Entomol.   2007;52:81–106.
431   doi:10.1146/annurev.ento.52.110405.091440

432   26.   Haynes KF. Sublethal effects of neurotoxic insecticides on insect behavior. Annual
433   review of entomology. 1988;33(1):149–168.

434   27.   Glunt KD, Coetzee M, Huijben S, Koffi AA, Lynch PA, N'Guessan R, et al. Empirical
435   and theoretical investigation into the potential impacts of insecticide resistance on the
436   effectiveness   of   insecticide-treated   bed   nets.   Evolutionary   Applications.
437   2018;11(4):431–41. doi:10.1111/eva.12574

438   28.   Alout H, Ndam NT, Sandeu MM, Djégbe I, Chandre F, Dabiré RK, et al. Insecticide
439   Resistance Alleles Affect Vector Competence of Anopheles gambiae s.s. for
440   Plasmodium   falciparum   Field   Isolates.   PLOS   ONE.   2013   mai;8(5):e63849.
441   doi:10.1371/journal.pone.0063849

442   29.   N'Guessan R, Corbel V, Akogbeto M, Rowland M. Reduced efficacy of insecticide-
443   treated nets and indoor residual spraying for malaria control in pyrethroid resistance
444   area, Benin. Emergent Infectious Diseases. 2007;13(2):199–206.

445   30.   Berticat C, Boquien G, Raymond M, Chevillon C. Insecticide resistance genes induce a
446   mating competition cost in Culex pipiens mosquitoes. Genet Res. 2002 Feb;79(1):41–7.

31. Porciani A, Diop M, Moiroux N, Kadoke-Lambi T, Cohuet A, Chandre F, et al.
Influence of pyrethroïd-treated bed net on host seeking behavior of Anopheles gambiae
s.s. carrying the kdr allele. PLOS ONE. 2017 juil;12(7):e0164518.
doi:10.1371/journal.pone.0164518

32. Jones JC, Madhukar BV. Effects of sucrose in blood avidity in mosquitoes. Journal of
Insect Physiology. 1976;22(3):357–60. doi:10.1016/0022-1910(76)90001-9

33. Skovmand O, Bonnet J, Pigeon O, Corbel V. Median knock-down time as a new method
for evaluating insecticide-treated textiles for mosquito control. Malaria Journal.
2008;7(1):114. doi:10.1186/1475-2875-7-114

34. Briegel H, Lea AO, Klowden MJ. Hemoglobinometry as a method for measuring blood
meal sizes of mosquitoes (Diptera: Culicidae). Journal of medical entomology.
1979;15(3):235–8.

35. Briegel H. Fecundity, metabolism, and body size in Anopheles (Diptera: Culicidae),
vectors of malaria. Journal of medical entomology. 1990;27(5):839–50.

36. R Development Core Team. R: A Language and Environment for Statistical Computing.
3.5. Vienna, Austria: R Foundation for Statistical Computing; 2018.

37. Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models Using
lme4. Journal of Statistical Software. 2015 Oct 7;67(1):1–48. doi:10.18637/jss.v067.i01

38. Brooks ME, Kristensen K, Benthem KJ van, Magnusson A, Berg CW, Nielsen A, et al.
glmmTMB Balances Speed and Flexibility Among Packages for Zero-inflated
Generalized Linear Mixed Modeling. The R Journal. 2017;9(2):378–400.

39. Therneau TM. coxme: Mixed Effects Cox Models [Internet]. 2018. Available from:
https://CRAN.R-project.org/package=coxme

40. Lenth R, Singmann H, Love J, Buerkner P, Herve M. emmeans: Estimated Marginal
Means, aka Least-Squares Means [Internet]. 2019. Available from: https://CRAN.R-
project.org/package=emmeans

41. Harrell FE, with contributions from Charles Dupont and many others. Hmisc: Harrell
Miscellaneous [Internet]. 2019. Available from: https://CRAN.R-
project.org/package=Hmisc

42. Moiroux N. Code and data to reproduce analyses in Diop et al. 2019 [Internet]. 2019.
Available from: https://github.com/Nmoiroux/diopKdrBiting

43. Lyimo IN, Haydon DT, Mbina KF, Daraja AA, Mbehela EM, Reeve R, et al. The fitness
of African malaria vectors in the presence and limitation of host behaviour. Malaria
journal. 2012;11(1):425.

44. Fernandes L, Briegel H. Reproductive physiology of Anopheles gambiae and Anopheles
atroparvus. Journal of vector ecology. 2005;30(1):11.

45. Walker ED, Edman JD. The influence of host defensive behavior on mosquito (Diptera:
Culicidae) biting persistence. Journal of Medical Entomology. 1985;22(4):370–2.

46. Roitberg BD, Mondor EB, Tyerman JG. Pouncing spider, flying mosquito: blood
acquisition increases predation risk in mosquitoes. Behavioral Ecology. 2003;14(5):736–
740.

47. Scott TW, Takken W. Feeding strategies of anthropophilic mosquitoes result in
increased risk of pathogen transmission. Trends in Parasitology. 2012 Mar;28(3):114–
21. doi:10.1016/j.pt.2012.01.001

48. Vézilier J, Nicot A, Gandon S, Rivero A. Insecticide resistance and malaria
transmission: infection rate and oocyst burden in Culex pipiens mosquitoes infected with
Plasmodium relictum. Malaria J. 2010;9:379.

49. Sattelle DB, Yamamoto D. Molecular Targets of Pyrethroid Insecticides. In: Evans PD,
Wigglesworth VB, editors. Advances in Insect Physiology [Internet]. Academic Press;
1988.   p.   147–213.   Available   from:
http://www.sciencedirect.com/science/article/pii/S0065280608600259

50. Wouters W, van den Bercken J. Action of pyrethroids. General Pharmacology: The
Vascular System. 1978 Jan 1;9(6):387–98. doi:10.1016/0306-3623(78)90023-X

51. O'Reilly AO, Khambay BPS, Williamson MS, Field LM, WAllace BA, Davies TGE.
Modelling insecticide-binding sites in the voltage-gated sodium channel. Biochemical
Journal. 2006 Jun 1;396(2):255. doi:10.1042/BJ20051925

52. Corbel V, Chandre F, Brengues C, Akogbeto M, Lardeux F, Hougard JM, et al. Dosage-
dependent effects of permethrin-treated nets on the behaviour of Anopheles gambiae and
the selection of pyrethroid resistance. Malaria journal. 2004 Jul 8;3:22.
doi:10.1186/1475-2875-3-22

53. Briegel H, Rezzonico L. Concentration of host blood protein during feeding by
anopheline mosquitoes (Diptera: Culicidae). Journal of medical entomology.
1985;22(6):612–8.

54. Lahondère C, Lazzari CR. Mosquitoes cool down during blood feeding to avoid
overheating. Current Biology. 2012;22(1):40–5.

55. Lahondère C, Lazzari CR. Thermal Stress and Thermoregulation During Feeding in
Mosquitoes. In: New insights into malaria vectors [Internet]. Sylvie Manguin; 2013
[cited   2015   Feb   16].   p.   17.   Available   from:
http://cdn.intechopen.com/pdfs/44039/InTech-
Thermal_stress_and_thermoregulation_during_feeding_in_mosquitoes.pdf

56. Raphemot R, Rouhier MF, Hopkins CR, Gogliotti RD, Lovell KM, Hine RM, et al.
Eliciting Renal Failure in Mosquitoes with a Small-Molecule Inhibitor of Inward-
Rectifying Potassium Channels. PLoS ONE. 2013 May 29;8(5):e64905.
doi:10.1371/journal.pone.0064905

57. Alout H, Yameogo B, Djogbenou LS, Chandre F, Dabire RK, Corbel V, et al. Interplay
Between Plasmodium Infection and Resistance to Insecticides in Vector Mosquitoes.
The Journal of infectious diseases [Internet]. 2014 May 14; doi:10.1093/infdis/jiu276

58. Alout H, Djegbe I, Chandre F, Djogbenou LS, Dabire RK, Corbel V, et al. Insecticide
exposure impacts vector-parasite interactions in insecticide-resistant malaria vectors.
Proceedings Biological sciences / The Royal Society [Internet]. 2014 Jul 7;281(1786).
doi:10.1098/rspb.2014.0389

59. Cator LJ, George J, Blanford S, Murdock CC, Baker TC, Read AF, et al. "Manipulation"
without the parasite: altered feeding behaviour of mosquitoes is not dependent on
infection with malaria parasites. Proceedings of the Royal Society B: Biological
Sciences. 2013 May 22;280(1763):20130711–20130711. doi:10.1098/rspb.2013.0711

60. Cator LJ, Lynch PA, Read AF, Thomas MB. Do malaria parasites manipulate
mosquitoes? Trends Parasitol. 2012 Nov;28(11):466–70. doi:10.1016/j.pt.2012.08.004

61. Vantaux A, Yao F, Hien DF, Guissou E, Yameogo BK, Gouagna L-C, et al. Field
evidence for manipulation of mosquito host selection by the human malaria parasite,
Plasmodium falciparum. bioRxiv. 2018 Nov 8;207183. doi:10.1101/207183

62. Lefevre T, Ohm J, Dabiré KR, Cohuet A, Choisy M, Thomas MB, et al. Transmission
traits of malaria parasites within the mosquito: Genetic variation, phenotypic plasticity,
and consequences for control. Evolutionary Applications. 2018 Apr 1;11(4):456–69.
doi:10.1111/eva.12571

63. Sherrard-Smith E, Griffin JT, Winskill P, Corbel V, Pennetier C, Djénontin A, et al.
Systematic review of indoor residual spray efficacy and effectiveness against
Plasmodium falciparum in Africa. Nature Communications. 2018 Nov 26;9(1):4982.
doi:10.1038/s41467-018-07357-w

Figure legends

Figure 1 Experimental set-up to monitor the biting behavioural sequence.

OT: Observational tunnel, TP: Transparent plastic, OZ: Observational zone (mosquito biting the rabbit ear), R: Rabbit, RC: Restraining cage

Figure 2 Feeding success and biting behavior in absence of insecticide of *Anopheles gambiae* females of each *kdr* genotype.

Feeding success of each genotype and 95% binomial confidence intervals of the proportions (error bars) are shown in panel A. Panels B, C and D show boxes-and-whiskers plots of feeding duration, prediuresis duration and blood-meal size, respectively. Boxes indicate 1^{st}-3^{rd} quartile and median values. Whiskers indicate 2.5-97.5 percentiles. Significance according to Tukey's test after binomial mixed-effect model (panel A), mixed-effect cox proportional hazard model (panel B and C) and after linear mixed effect model (panel D) are indicated (ns $p>0.05$, * $p<0.05$, ** $p<0.01$, *** $p<0.001$).

Figure 3 Feeding success after exposure to insecticides of *Anopheles gambiae* females of each *kdr* genotype.

Feeding success of SS, RS, and RR (panel A, B and C, respectively) genotypes when pre-exposed to untreated, permethrin-treated (Olyset) and deltamethrin-treated (PermaNet) nettings are shown with 95% binomial confidence intervals of the proportions (error bars). Significance according to Tukey's test after binomial mixed-effect model is indicated (ns $p>0.05$, * $p<0.05$, ** $p<0.01$, *** $p<0.001$).

Figure 4 Biting behavior after exposure to insecticides of *Anopheles gambiae* females of

each *kdr* genotype.

Boxes-and-whiskers plots of feeding duration (panels A, D and G), prediuresis duration

(panels B, E and H) and blood-meal size (panels C, F and I) are shown for each genotypes SS

(panels A,B and C), RS (panels, D, E and F) and RR (panels G, H and I). Boxes indicate 1^{st}-

3^{rd} quartile and median values. Whiskers indicate 2.5-97.5 percentiles. Significance according

to Tukey's test after mixed-effect cox proportional hazard model (panels A, B, D, E, G and H)

and after linear mixed effect model (panels D, F and I) are indicated (ns $p > 0.05$, * $p < 0.05$, **

$p < 0.01$, *** $p < 0.001$).

Supplementary Figure 1: Among *kdr* genotypes comparison of feeding success of

***Anopheles gambiae* females after exposure to insecticides**.

Feeding success of each genotype and 95% binomial confidence intervals of the proportions

(error bars) are shown after exposure to untreated net (Panel A, same as Figure 2A), to

permethrin treated net (Olyset; panel B) and to deltamethrin treated net (Permanet; panel C).

Significance according to Tukey's test after binomial mixed-effect model is indicated (ns

$p > 0.05$, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$).

Figure 1

Figure 2

Figure 3

Figure 4

Supp. Figure 1