1   Identification of new therapeutic targets in *CRLF2*-overexpressing B-ALL through discovery of
2   TF-gene regulatory interactions

Sana Badri[1,2,*], Beth Carella[1,3,*], Priscillia Lhoumaud[1], Dayanne M. Castro[2], Claudia Skok
Gibbs[4], Ramya Raviram[1, 5], Sonali Narang[6], Nikki Evensen[6], Aaron Watters[4], William Carroll[6],
Richard Bonneau[2,4,7,**], Jane A. Skok[1,**].

[1]Department of Pathology, New York University School of Medicine, New York, NY, USA.

[2]Department of Biology, Center for Genomics and Systems Biology, New York University, New
York, NY, USA.

[3]Division of Blood and Marrow Transplantation and Cellular Therapies, UPMC Children's
Hospital of Pittsburgh, Pittsburgh, PA.

[4]Flatiron Institute, Center for Computational Biology, Simons Foundation, New York, NY, USA.

[5]Department of Chemistry and Biochemistry, University of California, San Diego, CA, USA.

[6]Perlmutter Cancer Center, Departments of Pediatrics and Pathology, New York, NY, USA.

[7]Department of Computer Science, Courant Institute of Mathematical Sciences, New York, USA.

[*]These authors contributed equally to this manuscript

[**] Correspondence and requests for materials should be addressed to:

e-mail: rb133@nyu.edu (Richard Bonneau) jane.skok@med.nyu.edu (Jane A. Skok)

21
22
23

ABSTRACT

Although genetic alterations are initial drivers of disease, aberrantly activated transcriptional regulatory programs are often responsible for the maintenance and progression of cancer. *CRLF2*-overexpression in B-ALL patients leads to activation of JAK-STAT, PI3K and ERK/MAPK signaling pathways and is associated with poor outcome. Although inhibitors of these pathways are available, there remains the issue of treatment-associated toxicities, thus it is important to identify new therapeutic targets. Using a network inference approach, we reconstructed a B-ALL specific transcriptional regulatory network to evaluate the impact of *CRLF2*-overexpression on downstream regulatory interactions.

Comparing RNA-seq from *CRLF2*-High and other B-ALL patients (*CRLF2*-Low), we defined a *CRLF2*-High gene signature. Patient-specific chromatin accessibility was interrogated to identify altered putative regulatory elements that could be linked to transcriptional changes. To delineate these regulatory interactions, a B-ALL cancer-specific regulatory network was inferred using 868 B-ALL patient samples from the NCI TARGET database coupled with priors generated from ATAC-seq peak TF-motif analysis. CRISPRi, siRNA knockdown and ChIP-seq of nine TFs involved in the inferred network were analyzed to validate predicted TF-gene regulatory interactions.

In this study, a B-ALL specific regulatory network was constructed using ATAC-seq derived priors. Inferred interactions were used to identify differential patient-specific transcription factor activities predicted to control *CRLF2*-High deregulated genes, thereby enabling identification of new potential therapeutic targets.

Keywords: Transcriptional Regulatory Network, B-ALL, CRLF2, Leukemia, Transcription Factor Activity, TARGET, ATAC-seq, Network Inference

INTRODUCTION:

Acute lymphoblastic leukemia (ALL) is the most common cancer in children. Historically, clinical criteria have driven risk stratification for these patients, however over time many genetic alterations have been identified as prognostic predictors. Several groups have described varied genetic signatures associated with pediatric ALL with the aim of elucidating their contributions to leukemogenesis[1,2]. Improved risk stratification based on genetic signature has altered treatment and led to significant improvements in overall survival[3]. However, about 20% of patients fail current treatment strategies or die following relapse. Moreover, adults with ALL have a worse prognosis and an average overall survival of 35-50%[3]. Therefore, it is important to gain a better understanding of the mechanisms by which genetic alterations drive leukemogenesis to refine therapies that target the disease-essential pathways involved[4].

Genetic alterations that lead to overexpression of the cytokine receptor-like factor 2 (*CRLF2*) gene have been associated with a high-risk subset of pediatric patients with B-cell acute lymphoblastic leukemia (B-ALL). The *CRLF2* gene encodes the thymic stromal lymphopoietin receptor (TSLPR) which forms a heterodimer with IL-7 receptor alpha (IL7RA) to bind TSLP[5]. Binding of TSLP to the IL7RA-TSLPR complex signals the phosphorylation of Janus kinase 1 (JAK1) and Janus kinase 2 (JAK2), leading to the activation of the JAK-STAT signaling pathway[6]. Studies have shown that stimulation of TSLP in B-ALL not only induces activation of the JAK-STAT pathway, but also activates the PI3K/mTOR[7] and ERK/MAPK signaling pathways[3].

CRLF2 overexpression occurs in 5-15% of patients with B-ALL and in 50-60% of pediatric B-ALL patients with Down syndrome (DS)[6,8]. Overexpression can occur either from a chromosomal translocation between *CRLF2* and the immunoglobulin heavy chain locus (*IGH*) on chromosome 14 (*IGH-CRLF2*) or from an interstitial deletion of the pseudoautosomal region of the X/Y chromosomes resulting in the fusion of *CRLF2* to the *P2RY8* gene (*P2RY8-CRLF2*)[9] as shown in **Figure 1a**. *CRLF2* deregulation very rarely occurs via activating mutations of

CRLF2. *IGH-CRLF2* translocation occurs in precursor cells and is thought to result from

aberrant rejoining during V(D)J-recombination[10,11]. This translocation is most commonly found in

adolescents and adult patients and is typically associated with a poor prognoses, while the

P2RY8-CRLF2 fusion is found in younger patients[3].

CRLF2 chromosomal alterations are often accompanied by mutations in *JAK1*, *JAK2*

and *Ikaros* (*IKZF1*) genes. Several groups have suggested that aberrant *CRLF2* signaling

cooperates with mutant JAK and IKZF1 activity to promote the development of leukemia[9,12,13].

As a result, the focus has shifted towards the use of signal transduction inhibitors (STIs) to

target JAK-STAT, PI3K and MAPK signaling pathways[3]. Although, STIs have shown promise in

early clinical trials[14-16], broad application of signal transduction inhibitors is challenging due to

the interconnected roles of their targets in biological processes (i.e. JAK kinases) including

immunity and hematopoiesis[17]. Moreover, it has been shown that mutated *JAK2* is required for

the initiation of leukemia, but it is not necessary for its maintenance [18].

More recently, studies have constructed short and long chimeric antigen receptor (CAR)-

expressing T-cells to target *CRLF2* receptor and have demonstrated only the short CARs

targeting *CRLF2* can eliminate the leukemia in cell lines and xenograft models of *CRLF2*-

overexpressing ALL[19]. *CRLF2*-targeted CAR T cells represent a powerful immunotherapeutic

strategy for this cohort of B-ALL patients, nonetheless there are severe toxicity issues

associated with this type of therapy, related to damage of normal tissues that express this

antigen[20].

Exome and targeted sequencing of DS-ALL with *CRLF2* rearrangements has uncovered

driver mutations in *KRAS* and *NRAS* that are mutually exclusive to *JAK* mutations. *RAS* and

JAK mutations appear to occur at a similar frequency. Analysis of clonal architecture

demonstrates that the majority of clones initiated by *CRLF2* rearrangements are expanded into

distinct sub-clones by both *RAS* and *JAK2* mutations. Interestingly, the majority of relapsed

cases switch from a *JAK2*-mutated sub-clone to a *RAS*-mutated sub-clone, indicating there are

alternate drivers in relapsed *CRLF2*-overexpressing DS-ALL[21]. Although, this particular study

focused on DS-ALL, there is evidence that alternate genes can be responsible for the

maintenance of *CRLF2*-overexpressing cells[18], therefore, there is a need to more closely

investigate the impact of the altered *CRLF2* in B-ALL downstream of oncogenic signaling, to

identify alternate targets that can be evaluated for therapy.

To address the issue of alternate targets, we first sought to identify transcriptional

regulators that control differentially expressed genes associated with *CRLF2*-overexpression

through a comparative analysis of *CRLF2*-High cells versus other B-ALL samples that express

CRLF2 at low levels in primary patient samples. Differential analysis of RNA-seq samples

resulted in robust genome-wide gene expression changes, with an expected significant

enrichment in cytokine-mediated signaling pathways. Using ATAC-seq data, patient-specific

chromatin accessibility landscapes were defined and interrogated to identify altered regions that

could be linked to transcriptional changes. While strong changes in accessibility were detected

in patients, only a proportion of these were linked to transcriptional changes. As a result, we

hypothesized that differential binding of TFs at ATAC-seq peaks are influencing gene

expression changes.

Using 868 B-ALL patient samples from the NCI TARGET database, we constructed A B-

ALL transcriptional network to define the interactions between transcription factors (TFs) and

the genes they regulate[22-24]. With RNA-seq, ATAC-seq, and TF-motif analysis, we inferred the

targets of TFs linked to the gene set associated with the *CRLF2*-High cohort using the

Inferelator algorithm[25,26]. The network was then used to predict differential transcription factor

activity (TFA) in *CRLF2*-High versus other B-ALL (*CRLF2*-Low) patient samples. This approach

enabled the identification of a *CRLF2*-High associated sub-network consisting of thirty-four

potential regulators of differentially expressed genes (including FOXM1, ETS2, FOXO1).

Several of the differential gene targets of these TFs (i.e. *PIK3CD*, *TNFRSF13C, PTPN7, GAB1*

and *TCF4)* are conserved in CRLF2 High versus Low patient-derived cell lines and have been

previously implicated in leukemias. The network-based approach we applied has been similarly

implemented in several other systems to infer regulatory interactions that have been

experimentally validated[27-30]. To validate TF-gene inferred interactions, we analyzed CRISPRi,

siRNA knockdown, and ChIP-seq ENCODE datasets in K562 and GM12878 cells involving 9

individual TFs. This resulted in the validation of gene targets for each TF further supporting the

network-based approach applied in this study. Thus, we uncovered a number of genes along

with their regulators that could contribute to the maintenance of leukemia and which act as

potential candidates for therapeutic targeting in *CRLF2*-High B-ALL.

RESULTS

Genome-wide transcriptional changes in *CRLF2*-overexpressing primary patient samples

Traditional chemotherapy is non-specific and targets rapidly dividing cells. Thus, toxicity

results in injury to healthy cells, causing further morbidity[31]. To reduce overall toxicity and

improve prognosis, most research has been directed towards understanding the underlying

molecular pathology of the leukemia. In this study, we focused on identifying regulatory

interactions associated with overexpression of *CRLF2* with the goal of finding new potential

therapeutic targets in this subset of B-ALL patients.

Gene expression profiles associated with *CRLF2* overexpression were identified by

comparing RNA-sequencing from 15 CRLF2-overexpressing patients (CRLF2-High) and 15

patients with low levels of *CRLF2* (Other B-ALL). Principal component analysis was performed

on gene expression levels across all 30 primary patient samples. This revealed a clear

separation between the two groups of patients (**Fig 1b**). Principal component analysis with

samples labeled according to batch do not demonstrate a batch effect (**Supplementary Fig.**

1a).

To further analyze the changes in gene expression between *CRLF2*-High and other B-

ALL patient samples we performed differential expression analysis using DESeq2[32]. This

analysis uncovered a total of 3,586 de-regulated genes with an adjusted p-value of less than

0.01 and |log$_2$FoldChange| >2 (**Fig. 1c, Supplemental Table 1**). Of these 1470 genes (~41%)

were up-regulated and 2116 (~59%) down-regulated in the *CRLF2*-High patients. In addition,

hierarchical clustering of the 3,586 differentially expressed genes shown in the **Figure 1d**

heatmap highlights the prominent transcriptional changes associated with overexpression of

CRLF2.

CRLF2-High patient samples clearly exhibit up-regulation of *CRLF2* (log$_2$FoldChange=

10.12) compared to control B-ALL patient samples as shown by RNA-seq tracks on IGV

(**Supplementary Fig. 1b**). Based on previous findings we expected to find evidence for

activation of JAK-STAT signaling. Indeed, in *CRLF2*-High samples *SOCS2*, a downstream

target of JAK/STAT signaling which encodes a SOCS protein that inhibits cytokine signaling as

part of a classical feedback loop[33], was statistically significantly up-regulated (log$_2$FoldChange=

4.30) (**Supplemental Table 1**). Thus, consistent with previous findings[34], JAK-STAT[3,4] signaling

appears to be activated in these samples.

GSEA was applied on the normalized expression counts of the 3,586 differentially

expressed genes using the molecular signatures (MSigDB) database to identify enriched

hallmark gene sets that represent well-defined biological processes. Significant pathways with

p-value<0.01 and FDR=1% (**Supplemental Table 2**) were enriched including IL2-STAT5 and

cytokine mediated signaling (**Supplemental Fig. 1c-1h**).

***CRLF2*-overexpression leads to promoter-enriched ATAC-seq changes in patient**

samples.

To further investigate the impact of *CRLF2* overexpression, we asked whether CRLF2-

High associated transcriptional changes were accompanied by changes in chromatin

accessibility. For this we analyzed ATAC-sequencing and compared chromatin accessibility in

CRLF2-High versus other B-ALL patient samples (7 *CRLF2*-High and 6 other B-ALL). PCA

analysis on normalized ATAC-seq signal clearly separated the two conditions (**Figure 2a**).

Differential analysis of all the ATAC-seq peaks identified 700 regions that were more accessible

and 978 regions with reduced accessibility in the *CRLF2*-High versus low (**Fig. 2b-c**).

A total number of 115,457 ATAC-seq peaks across all patients were assigned to

promoters (within 3kb of TSS), UTRs, exons and introns of genes (**Fig. 2d**). Significant

differential ATAC-seq peaks (n=1,678) were similarly annotated (**Fig. 2e**) and more highly

enriched (~47%) at promoter regions compared to all peaks (~29%). There were 317 (18.9%)

differential ATAC-seq peaks associated with significant differentially expressed genes (n=268),

either on promoters or gene bodies. The majority of the differential ATAC-seq peaks (223) that

overlap differentially expressed genes were concordantly regulated, with the exception of 94

ATAC-gene pairs. Discordant ATAC-gene pairs could be representative of silencers and their

gene targets. For example, a decrease in accessibility can impair binding of a silencer to its

gene target and therefore lead to an increase in that gene's expression (**Fig. 2f**). Furthermore,

about 79% (177/223) of concordant pairs represented ATAC-seq peaks with decreasing

accessibility associated with genes exhibiting reduced expression. An example of this is shown

in **Figure 2g**, where an ATAC-seq peak with significantly decreased accessibility is associated

with down-regulation of receptor tyrosine kinase *KIT* in *CRLF2*-High patients. Importantly, the

majority of transcriptional changes were not associated with significant changes in chromatin

accessibility near the promoter or within the gene bodies (3328/3596 genes). In sum,

overexpression of *CRLF2* leads to genome-wide chromatin accessibility changes enriched at

promoter regions with a small proportion linked to significant transcriptional changes.

Construction of a B-ALL regulatory network using ATAC-motif derived priors

CRLF2 overexpression activates a signaling cascade that involves many TF regulators and gene targets. To examine these connections, we wanted to first identify which TFs could potentially be regulating the differentially expressed genes in *CRLF2*-High versus other B-ALL patient cohorts, and second to infer the relationship between these TFs and their target genes. To address this, we performed TF-motif analysis, using Analysis of Motif Enrichment (AME)[35] to identify enriched TF motifs at promoter-annotated differentially accessible ATAC-seq peaks compared to shuffled input control sequences. Using an adjusted fisher p-value$<1^{e-13}$, 138 enriched TF motifs were defined at differentially accessible promoter peaks that could be important for the *CRLF2*-High associated gene signature (**Supplemental Fig. 3a**).

To better understand the relationship between candidate TF regulators identified in **Supplemental Fig. 3a** and the genes they potentially regulate, we sought to infer a transcriptional regulatory network using the Inferelator algorithm[25,26]. The main limitation of network inference is the sample size (the limited number of samples with an *CRLF2*-High rearrangement). To model transcriptional interactions in a B-ALL specific context requires a large dataset that includes hundreds of B-ALL patient samples, not limited to samples with a specific genetic alteration. For this we made use of the TARGET initiative and analyzed 868 B-ALL RNA-seq samples that were used to obtain a normalized gene expression matrix that could be used for the construction of a B-ALL specific regulatory network.

There are three major steps involved in inferring a transcriptional regulatory network, including generating priors, estimating transcription factor activities, and modeling gene expression with regularized regression. To generate priors, recent studies have incorporated prior information from different data types, like ChIP-seq, ATAC-seq, and TF-motif analysis to associate TFs to their putative gene targets and this has considerably improved network inference [27-29,36]. Thus, we focused on combining chromatin accessibility data from the *CRLF2*-

High and Low patients with TF-motif analysis to generate priors and acquire the TF-gene

associations. First, all ATAC-seq peaks found within a 20kb window upstream of a gene's

transcription start site (TSS) were assigned to that gene. Second, FIMO was used to analyze all

known TF motifs enriched within gene-associated ATAC-seq peaks. Lastly, TF motifs were

aggregated across all ATAC-seq peaks for each gene. For each TF motif-gene association, a

score, a p-value and an adjusted p-value was computed. Only TF-gene associations that had an

adjusted p-value less than 0.01 were retained (**Fig. 3a**). With a prior matrix and a normalized

gene expression matrix from hundreds of B-ALL specific patient samples, we used the

Inferelator algorithm to infer regulatory interactions. Finally, to reduce the number of regulatory

hypotheses on TF-gene interactions and improve predictions, we tested only regulatory

interactions of TFs (n=138) that were significantly enriched at differentially accessible promoters

between *CRLF2*-High and other B-ALL patient samples **(Supplemental Fig 3a)**.

The second step of network inference involves estimating the activity of all TFs in our

network based on the cumulative effect of each TF on its target genes[29,37]. The prior matrix

represents a list of potential TF-gene interactions derived from ATAC-TF-motif analysis and the

activity of each TF can be estimated by taking the pseudo-inverse of the prior multiplied by the

gene expression matrix (**Fig. 3a**). Thus, transcription factor activities were estimated using the

ATAC-seq motif derived priors and the gene expression matrix obtained from the TARGET

database (**Fig. 3a**). Finally, gene expression was modeled as a weighted sum of its activities

using regression with regularization (see methods section for details). The output is a matrix

where each row represents a TF-gene interaction, with a β parameter that corresponds to the

magnitude and direction of a particular interaction and a computed confidence score (see

methods section for details).

Using the ATAC–motif prior, the performance of our model selection within the

Inferelator was evaluated by applying a 10-fold cross validation technique. Specifically, we split

the prior into 10 equal sized sets. One set was retained as the gold standard to test the model,

while the remaining 9 sets were used as the prior. The cross-validation procedure was repeated

10 times, until each of the 10 partitions was used exactly once as the gold standard. As a

negative control, we employed the same 10 fold cross-validation strategy after randomly

shuffling the prior. We computed the area under the precision-recall curve (AUPR) for each of

the 10 iterations and compared the results between the prior and the shuffled prior. Overall,

AUPR were significantly higher using the ATAC-seq derived motif prior compared to the

randomly shuffled prior (**Supplemental Fig 3b**). Thus, the final network was inferred using the

ATAC-seq derived motif prior and the number of significant regulatory interactions was limited to

TF-gene interactions with combined confidences > 0.9 (high-confidence interactions). This

generated a B-ALL specific regulatory network involving 135 TFs (that may be important

regulators of *CRLF2*-High gene signature) and 8,731 high confidence patient specific TF-gene

interactions (**Fig. 3a**).

Significant Transcription Factors altered by *CRLF2* overexpression

To identify TFs affected by *CRLF2* overexpression, we reapplied the TF activity

estimation strategy described above to estimate activity across the 30 *CRLF2*-High and other B-

ALL samples. By assigning the inferred 8,731 patient-specific TF-gene interactions as priors

with the normalized gene expression matrix including all 30 primary patient samples, we

estimated an activity matrix of all 135 TFs. Next, we evaluated TFA estimates using PCA

analysis across the 30 samples and demonstrated a clear separation of *CRLF2*-High and Low

patient samples (**Fig. 3b**). To define the top TFs that explain the variance associated with

overexpression of *CRLF2*, we extracted and ranked the absolute value of the PC1 values for

each TF (**Supplemental 4a**). As a secondary means to identify TFs altered at the activity level,

we computed the mean activity level of each TF between *CRLF2*-High and Low samples and

tested for statistical difference**.** To narrow the list of potential TFs altered as a result of *CRLF2*

overexpression, we ranked the most highly variable TFs and retained those with |PC1| > 0.075,

as well as TFs with significant differences in mean activity level with an adjusted p-value 0.001.

This resulted in 36 significant TFs as highlighted in the volcano plot in **Fig. 3c**. Boxplots

depicting highly variable TFs (n=36) with significant differential mean activities are represented

in **Supplemental Fig 4b**. As a result, we defined 36 significant TF regulators predicted to

regulate 2,410 gene targets from the inferred B-ALL regulatory network previously described,

with the number of targets for each TF shown in **Supplemental Fig 4c**. We note that many TFs

(FOXM1[38], ETS2[39,40], BCL11A[41], FOXO1[42], etc.) with significantly altered activity are implicated

in cancer and could therefore potentially contribute to tumorigenesis in *CRLF2*-High B-ALL

patients. Furthermore, 6 of the 36 significant TFs (*ETS2, SRY, FOXO1, EGR1, MEF2D, KLF5*)

exhibit a statistically significant difference in mRNA expression between the two patient groups

(**Supplemental Fig. 4d**). According to both transcriptional and predicted activity levels, these

TFs represent robust candidates that could be important downstream effectors of the *CRLF2*-

mediated signaling cascade driving leukemogenesis.

A *CRLF2*-associated sub-network of altered TFs and their differentially expressed targets

To filter regulatory interactions to a specific *CRLF2*-High sub-network, we computed the

number of gene targets that were differentially expressed (542/2410) in *CRLF2*-High versus

CRLF2-low patient samples (**Fig. 4a**). Ingenuity pathway analysis was performed on the 542

differentially expressed gene targets in the *CRLF2*-High associated sub-network. This

generated 22 significant pathways (**Fig. 4b**) including the activation of the canonical PI3K

pathway (**Fig. 4a**). The majority of the gene targets are involved in more than one signaling

pathway and many of these pathways are important in adaptive immunity and inflammatory

responses.

To define a set of genes in the *CRLF2*-High associated sub-network that could have the

highest potential as therapeutic targets, we turned to patient-derived cell lines to identify robust

changes across both model systems. The patient-derived cell lines used in our analysis were

MHH-CALL-4[43] (CALL) and MUTZ5[44] (MUTZ) that harbor *IGH-CRLF2* translocations leading to

CRLF2 overexpression, and a non-translocated B-ALL pre-B leukemic cell line, SMS-SB[45]

(SMS). As seen in the RNA-seq tracks of **Supplementary Fig. 5a**, *CRLF2* is clearly

overexpressed in the translocated CALL and MUTZ cell lines, compared to non-translocated

SMS. Principal component analysis of cell line RNA-seq samples (**Supplementary Fig. 5b**)

indicated that the majority of the variance (71%) lies between *CRLF2*-High cell lines and SMS

cell lines, while about 26% of variance separated the two *CRLF2*-High cell lines. Thus the PCA

analysis demonstrates overexpression of *CRLF2* is the major difference between the two

conditions, consistent with what was observed in patient samples.

Differential gene expression analysis of CALL versus SMS (1,631 DE genes) and MUTZ

versus SMS (1,484 DE genes) identified thousands of differentially expressed genes

(**Supplementary Fig. 5c**), with roughly equivalent numbers of up and down-regulated genes in

each case. The number of overlapping differentially expressed genes is shown in

Supplementary Fig. 5d. As shown in **Supplementary Fig. 5e,** the $\log_2$ fold change of

(CALL/SMS) and (MUTZ/SMS) demonstrated that about 97% of the genes were in convergent

orientation (561 up-regulated, 353 down-regulated). We compared the gene targets in the

CRLF2-associated regulatory sub-network with the convergent differentially expressed genes in

cell lines and identified 67 significant convergent transcriptional alterations that were conserved

across *CRLF2*-High patient and cell line samples. This is shown in **Figure 4c** by the $\log_2$ fold

change of all 67 convergent differentially expressed genes in the patients and each pairwise cell

line comparison (CALL versus SMS, and MUTZ versus SMS). Of the 67 gene targets, 84% are

up-regulated in *CRLF2*-High samples including *PIK3CD, DPEP1, PTPN7, TNFRSF13C*

(BAFFR) and Class II major histocompatibility complex (MHC) genes. Previous studies have

implicated the majority of these genes in hematologic malignancies including B-ALL[46-51] but their

affect specifically in *CRLF2*-overexpressing B-ALL is not known. Furthermore, thirteen of these

genes have available inhibitors (**Supplemental Table 3**).

Validating Inferred Gene Targets of TFs significantly altered by *CRLF2* overexpression

To support our network-based approach in identifying TF-gene regulatory interactions in

leukemic cells, we sought to validate our predictions of interactions involving TFs using publicly

available data in similarly related cells. We collected ENCODE data to support TF-gene

associations involving 9 significant TFs. The experiments analyzed include CRISPR

interference (CRISPRi) targeting of FOXM1 in myelogenous leukemic cells (K562), siRNA

targeting of MAZ and E2F6 in K562 cells, and ChIP-seq for 6 significant TFs (BCL11A, ETS2,

EGR1, PBX2, IRF1, TBP) in K562 cells and a B-lymphocyte derived cell line (GM12878).

First, we computed the number of gene targets we inferred for each of the 9 TFs (**Figure**

5a). To validate some of the predicted gene targets of FOXM1, we compared RNA-seq samples

in K562 cells treated with CRISPRi targeting of FOXM1 versus non-targeted cells. A PCA of

these samples distinctly separates the FOXM1 CRISPR targeted samples from the non-

targeting control samples (**Figure 5b**). DESeq2 was applied was used to examine differentially

expressed genes associated with FOXM1 targeting. These were overlapped with the network's

predicted gene targets of FOXM1. Of the 123 gene targets we predicted to be regulated by FOXM1, 50 (40%) had a significant change in their expression upon FOXM1 CRISPRi targeting including *CRLF2* itself (**Figure 5c**). FOXM1 is a forkhead box transcription factor functioning downstream of the PI3K/AKT/mTOR pathway, which is known to be induced in *CRLF2*-overexpressing B-ALL[7,52]. It is involved in cell proliferation, DNA repair, and self-renewal. Overexpression of FOXM1 has been detected not only in B-ALL[38] but also in a wide range of human cancers and its overexpression is linked to poor prognosis[38,53-62].

To evaluate predicted gene targets of another significant TF, we identified genes altered by siRNA knockdown of the E2F6 gene in K562 cells. E2F6 is one of the E2F transcription factors that play distinct and overlapping roles in regulating cell cycle progression[63]. E2F proteins can be transcriptional activators or repressors. In particular, E2F6, has been shown to repress the transcription of known E2F target genes and is considered a dominant negative inhibitor of the other E2F proteins. Yeast two-hybrid assays have determined that E2F6 associates with known components of the polycomb complex, suggesting a role for E2F6 in recruiting the polycomb transcriptional repressor to target genes[64]. To support some the inferred gene targets of E2F6 using the network based strategy in this study, we analyzed E2F6 targeted RNA-seq samples using siRNA versus non-targeted controls. A PCA analysis confirmed expected clustering of replicates for E2F6 targeted and control samples (**Figure 5d**). Differentially expressed genes were determined and overlapped with inferred gene targets of E2F6. This analysis supported 57 of the 272 (21%) predicted E2F6 gene targets (**Figure 5e**).

Next, we analyzed predicted gene targets of the MAZ TF, also known as Myc-associated zinc-finger protein. This TF is ubitiquously expressed in most tissues and has been shown to activate a variety of genes including c-Myc, insulin, VEGF, and MYB, and repress the p53 protein[65-67]. Specifically, it has been shown that the kinase Akt phosphorylates MAZ, which leads to the release of the MAZ TF from the p53 promoter followed by p53 transcriptional activation[66]. MAZ has also recently been identified as a factor that interacts with cohesin and

through this connection alter gene regulation as a result of its impact on TAD structure[68]. As

MAZ has been shown to have a dual role in both activation and repression of cancer-related

genes, it's role in different cancer types can vary greatly depending on its target genes. In this

particular study, we have predicted significant B-ALL specific regulatory interactions between

MAZ and 26 gene targets. To provide evidence supporting these predictions, we analyzed

siRNA knockdown of MAZ in K562 RNA-seq samples compared to control samples. A PCA on

MAZ knockdown and control K562 samples indicated reproducible replicates that cluster

according to their condition (**Figure 5f).** Genes that were differentially expressed were analyzed

and overlapped with predicted MAZ gene targets, resulting in 15% of predicted gene targets'

expression significantly altered by siRNA knockdown of MAZ (**Figure 5g).**

Finally, to support TF-gene regulatory interactions of other significant factors, we

downloaded available ENCODE optimal IDR (Irreproducible Discovery Rate) peaks for 6

important TFs (BCL11A, ETS2, EGR1, PBX2, IRF1, TBP) in K562 and GM12878 cell lines.

Optimal IDR peaks represent the most highly reproducible peaks across replicates. The aim

here was to delineate potential gene targets of each TF by determining binding sites according

to significant and reproducible ChIP-seq peaks at promoters. Thus, ChIP-seeker was applied to

annotate the optimal IDR peak sets of each TF to genes, using the same threshold applied to

generate TF-gene priors as previously described (20 KB window upstream of the TSS), the

number of genes associated with at least one ChIP-seq peak for each TF is shown in **Figure**

5h. For each individual TF of interest, we overlapped the network inferred gene targets of that

TF with the genes associated with ChIP-seq binding. Overall, we were able to validate 40.1% of

BCL11A gene targets, 69% of EGR1 gene targets, 6.1% of ETS2 gene targets, 26.3% of IRF1

gene targets, 57% PBX2 gene targets, and 47.8% of TBP gene targets that we infer using the

network-based approach (**Figure 5i**). Finally, individual hyper-geometric tests were performed

for each experiment; to test the significance of the proportion of validated gene targets

compared to random. The proportion of validated gene targets was significant for almost all 9

TFs, with the exception of ETS2, which could most likely be a matter of cell-type context (**Figure**

5i).

DISCUSSION

Patients with B-ALL who overexpress *CRLF2* are at increased risk for refractory disease

and relapse. This alteration leads to the activation of JAK-STAT and other associated

pathways[3]. Available treatments include non-specific cytotoxic chemotherapies which though

often effective are responsible for many of the toxicities seen in these patients. The addition of

Tyrosine-Kinase Inhibitors and JAK inhibitors in the treatment of leukemias has allowed for

targeted treatments but these treatments also affect the biology of healthy cells. Furthermore,

the use of *CRLF2*-targeted CAR T cells shows great promise in the eradication of the cancer,

yet limitations and toxicities associated with this treatment have yet to be studied in patients.

Thus, it is important to more deeply investigate the regulatory control underlying *CRLF2*-

overexpressing leukemia to identify and evaluate alternate gene therapeutic targets for tailored

treatment.

In this work, we sought to better understand the impact of *CRLF2* overexpression on

leukemogenesis, with the aim of delineating the regulatory interactions for patient-specific

CRLF2-High B-ALL. We analyzed RNA-seq from *CRLF2*-High and Other B-ALL patient samples

and identified hundreds of differentially expressed genes. We subsequently analyzed ATAC-seq

data to evaluate the effects of this genetic alteration on DNA accessibility. We found

accessibility to be altered across the genome with an enrichment of changes localized to

promoter regions. Only a small proportion of these altered promoter regions are linked with

gene expression changes.

To systematically annotate regulatory interactions, we inferred putative binding of

specific TF motifs at all ATAC-seq peaks found at the promoters of genes in the genome.

Leveraging hundreds of available B-ALL RNA-seq samples from the TARGET database along

with prior knowledge of TF-motifs at accessible promoters allowed us to infer a B-ALL specific

regulatory network involving TF motifs (135) enriched at differentially accessible promoters.

Deeper investigation of the TF regulators and robust differential gene targets specific to the

CRLF2-High sub-network identified a candidate list of potential therapeutic targets.

PIK3CD, one of the significantly overexpressed candidates identified in CRLF2-High

patients, encodes a class I phosphoinositide-3 kinase (PI3Ks), which is highly enriched in

leukocytes and known as p110δ. Up-regulation of this gene has been demonstrated to occur in

T-cell [46] acute lymphoblastic leukemia and it has emerged as a therapeutic target. Specifically,

the use of a p110δ inhibitor has led to reduced proliferation and apoptosis in T-ALL cell lines[46].

Furthermore, this gene target has been similarly implicated in Philadelphia chromosome–like

acute lymphoblastic leukemia (Ph-like ALL), another high-risk B-ALL subtype correlated with

poor prognosis[14]. CRLF2 overexpression occurs in about 50% of Ph-like ALL cases. Using

patient-derived xenograft (PDX) models of childhood Ph-like ALL, Tasian et al. demonstrate the

therapeutic efficacy of Idelalisib, a PI3Kδ inhibitor in vivo[14]. That PIK3CD has been previously

implicated in CRLF2-overexpressing B-ALL supports our approach in identifying relevant

CRLF2-associated gene targets.

Another gene target up-regulated in CRLF2-High patients is GAB1. This gene encodes

an adapter protein that recruits proteins, including PI3K to the plasma membrane to propagate

signals for many important biological processes such as cell proliferation, motility, and

erythrocyte development. GAB1 proteins lack enzymatic activity, but are phosphorylated at

tyrosine residues via interaction with PI3K, and this association is essential for activation of the

PI3K/AKT and MAPK/ERK pathways[69]. Up-regulation of GAB1 has been linked to breast cancer

progression[48] and implicated in BCR-ABL positive ALL cells[47]. Previous analysis of B-ALL adult

patients with the BCR-ABL alteration versus BCR-ABL–negative adult ALL patients, not only

identified overexpression of GAB1 but also up-regulation of several class II major

histocompatibility complex (MHC) genes and their trans activator *CIITA*[47]. These results are

similarly reproduced in our own data demonstrating conserved co-expression of *GAB1*, *CIITA,*

HLA-DQA1, HLA-DQB1, and *HLA-DRB1* across patients and cell lines. In addition, we

determined that 4 out of 5 of these genes (*GAB1*, *CIITA, HLA-DQA1, HLA-DQB1)* are predicted

to be gene targets of the BCL11A transcription factor. BCL11A is a zinc finger protein that is

expressed in hematopoietic cells and in the brain and whose presence is linked to

hematological malignancies[41,70]. This TF has been shown to play an important role in lymphoid

development. BCL11A expression in adult mouse is found in most hematopoietic cells but

enriched in B cells, early T cell progenitors, common lymphoid progenitors (CLPs) and

hematopoietic stem cells (HSCs). The deletion of BCL11A leads to early B-cell and CLP

apoptosis and impairs the development of HSCs into B, T, and natural killer (NK) cells[71].

BCL11A is a key regulator of fetal-to-adult hemoglobin switching. This regulator is also known to

interact with the chromatin remodeling SWI/SNF complex and the nucleosome remodeling and

deacetylase (NuRD) complex.[72]. The analysis described earlier, associating BCL11A ChIP-seq

binding sites to genes in GM12878 cells, validated 56 of BCL11A's predicted gene targets,

including overexpressed *GAB1*, *CIITA, HLA-DQA1, HLA-DQB1,* and *HLA-DRB1* in both patients

and cell lines. Another candidate gene target of BCL11A, validated by ChIP-seq binding, is

DPEP1. As a zinc-dependent peptidase, *DPEP1* is important for metabolism of the antioxidant

glutathione and was recently linked to relapse in B-ALL. Specifically, high levels of this gene

were linked with a higher incidence of relapse and worse overall survival[73].

GAB1 is also one of the few gene targets that we predict to be regulated by two

significantly altered TFs, BCL11A and PBX2. The pre-B-cell leukemia transcription factor 2,

PBX2 is a ubiquitously expressed protein and can regulate gene expression and effect

processes such as cell proliferation and differentiation[74]. It has also been shown to work as a

cofactor with the HOXA9 protein[75]. Knockdown of PBX2 leads to increased apoptosis in

adenocarcinoma and esophageal squamous cell carcinoma cancer cell lines[76]. PBX2 is also an

inferred regulator of the down-regulated *PTPN7* gene in *CRLF2*-overexpressing patients and

cell lines. This gene encodes a protein tyrosine phosphatase that is mainly expressed in

hematopoietic cells and known to negatively regulate the activation of ERK and MAPK

kinases[77]. Overexpression of *PTPN7* in T-cells down-regulates ERK signaling and reduces T-

cell activation and proliferation[78]. Thus, *PTPN7* represents another strong candidate gene target

as it can act as a negative regulator of one the canonical signaling pathways downstream of

CRLF2.

The up-regulation of *TNFRSF13C*, has also been linked to relapse in B-cell acute

lymphoblastic leukemia. *TNFRSF13C* encodes the receptor for B-cell activating factor (BAFFR)

and is important for B-cell maturation and survival. Expression of *TNFRSF13C* is primarily

detected in primary B-cell malignancies and B-cell related organs and absent in other healthy

related tissues[79,80]. There is evidence suggesting the expression of this receptor plays a role in

relapse as well as persistence of leukemic cells after early treatment[80]. Current drugs available

to target the BAFFR receptor include the Anti–BAFF-R antibody VAY-736[81]. In addition, a CAR-

based strategy has recently been engineered to target this receptor. Thus, *TNFRSF13C*

represents a potential target for *CRLF2*-overexpressing leukemias.

The network-driven approach also identified a number of significant TF regulator motifs

including ETS2[40], FOXO1[42], EGR1[82], and FOXM1[38,55,56], which are implicated in hematological

malignancies. FOXM1 could be an important regulator in *CRLF2*-overexpressing cells due to its

crucial role in cell proliferation and its implication in various cancer types, as described earlier.

Furthermore, our investigations predict that FOXM1 regulates *CRLF2* itself. This is interesting,

as FOXM1 is a known regulator functioning downstream of *CRLF2*-associated PI3K/AKT

pathway, yet it was not known that FOXM1 could regulate *CRLF2*.

Another candidate regulator, ETS2, a member of the *Ets* family of transcription factors, is

involved in cell proliferation and differentiation, and is a known target of the RAS/MAPK

signaling pathway[83]. In this study, ETS2 is differentially expressed and predicted to regulate the

aforementioned gene target *TNFRSF13C* or BAFFR. In AML, overexpression of ETS2 has been

shown to be associated with shorter overall survival and be a predictor of poor prognosis[39].

Analysis of overexpressed ETS2 in AML is associated with co-expression of *SOCS2* and *TCF4*,

both of which are significantly up-regulated in *CRLF2*-High B-ALL patient samples. *As*

mentioned previously, *SOCS2* is induced upon cytokine signaling. *TCF4*, also known as

TCF7L2, is a transcription factor that plays an important role in the Wnt signaling pathway and

its overexpression is required for proliferation and survival of AML cells[84]. Indeed, the results of

our present study identified *TCF4* as part of the *CRLF2*-associated sub-network and it is

overexpressed in both primary patients and patient derived cell lines. Given the role of *TCF4* in

the Wnt signaling pathway and its link to poor clinical outcome in AML, this gene represents

another putative gene target in *CRLF2*-overexpressing B-ALL.

FOXO1, another significantly altered TF, is one of the FOXO transcription factors

regulated by the PI3K/AKT signaling pathway. FOXO TFs are mostly considered as tumor

suppressors given their ability to inhibit cancer cell growth. Yet there is contrasting evidence for

their activation supporting tumor progression and inducing drug resistance[85]. In particular, the

TF FOXO1 plays a vital role in regulating proliferation and cell survival during B-cell

differentiation and it's activity, has been shown to be essential in B-cell precursor acute

lymphoblastic leukemia[42]. In this specific study in B-ALL, Wang et al. inhibited FOXO1 and

demonstrate reduced leukemic activity in cell lines, primary patients, and xenograft models[42].

The Early Growth Response 1 (EGR1) TF represents another candidate TF that is not

only significantly altered at the gene expression level but also predicted to have a change in

activity. This TF is induced in response to B cell receptor (BCR) signaling and mediated by the

Ras/Raf-1/MAPK pathway. Studies have also shown that EGR1 can promote differentiation in

pre-B cells[86]. High levels of EGR1 have been linked to human diffuse large B lymphoma cells

suggesting a role for EGR1 in B lymphoma cell growth[82]. In addition to previous evidence

supporting a role for FOXM1, ETS2, BCL11A, PBX2, EGR1, and FOXO1 in other malignancies,

the results of our analysis now implicate these TFs in *CRLF2*-overexpressing leukemia. We

speculate that downstream of *CRLF2*-mediated signaling, there are alterations to the

accessibility landscape that may be influencing TF binding and thereby deregulating

transcriptional programs. We have identified a set of TFs and their respect gene targets that we

propose may be important factors to consider in *CRLF2*-overexpressing leukemia (**Fig. 6a**).

Finally, using CRISPRi and siRNA RNA-seq data for FOXM1, E2F6, and MAZ, as well as ChIP-

seq for 6 other TFs from ENCODE, it was possible to validate a proportion of our predictions

linking these TFs and their predicted gene targets.

In summary, we have investigated the transcriptional impact of *CRLF2* overexpression in

a high-risk subset of B-ALL patients. Using an integrative approach, we derived regulatory

interactions that have enabled the identification of potential candidates for targeted therapies in

B-ALL patients with *CRLF2* overexpression. *PIK3CD, GAB1, DPEP1, TNFRSF13C, TFC4,*

PTPN7, BCL11A, FOXM1, ETS2, EGR1, PBX2, and *FOXO1* are amongst the most interesting

candidates as there is evidence supporting their role in hematological malignancies. The

strategy we have used here can further be adapted to elucidate important regulators and gene

targets responsible for additional subsets of B-ALL and other challenging malignancies.

543
544
545   **METHODS**

546
Cell Culture

The MUTZ5 (RRID:CVCL_1873) and MHH-CALL-4 (RRID:CVCL_1410) human B-cell leukemia

cell lines carrying CRLF2-High translocation were obtained, as well as the SMS-SB

(RRID:CVCL_AQ30) human B-cell leukemia control cell line. All cells were maintained in

RPMI1640 supplemented with 20% FBS and 1% penicillin-streptomycin under 5% CO_2 at 37

degrees. Ficoll-enriched, cryopreserved bone marrow samples were obtained from the

Children's Oncology Group (COG) from pediatric patients with B-precursor ALL from biology

protocols AALL03B1 and AALL05B1. Both *CRLF2*-HIGH and control patient samples were

used. All subjects (or their parents) provided written consent for banking and future research

use of these specimens in accordance with the regulations of the institutional review boards of

all participating institutions.

RNA-seq

RNA was extracted using the RNeasy plus kit from QIAGEN. For the cell lines, ribosomal

transcripts were depleted using the Illumina® Ribo Zero Module and for the patients, poly-

adenylated transcripts were positively selected using the NEBNext® Poly(A) mRNA Magnetic

Isolation Module, following the respective kit procedure. Libraries were prepared according to

the directional RNA-seq dUTP method adapted from

http://waspsystem.einstein.yu.edu/wiki/index.php/Protocol:directional_WholeTranscript_seq that

preserves information about transcriptional direction. Sequencing was performed with Illumina

Hi-Seq 2500 using 50 cycles paired-end mode.

ATAC-seq

Cell were counted to collect 50,000 cells per sample. The assay was performed as described

previously[87]. Cells were washed in cold PBS and resuspended in 50 µl of cold lysis buffer (10

mM Tris-HCl, pH 7.4, 10 mM NaCl, 3mM MgCl2, 0.1% IGEPAL CA-630). The tagmentation

reaction was performed in 25 µl of TD buffer (Illumina Cat #FC-121-1030), 2.5 µl Nextera Tn5

Transposase, and 22.5 µl of Nuclease Free H2O at 37oC for 30 min. DNA was purified on a

column with the Qiagen Mini Elute kit, eluted in 10 µl H2O. Purified DNA (10 µl) was combined

with 10 µl of H2O, 2.5 µl of each primer at 25 mM and 25 µl of NEB Next PCR master mix. DNA

was amplified for 5 cycles and a monitored quantitative PCR was performed to determine the

number of extra cycles needed as per the original ATAC-seq protocol[87]. DNA was purified on a

column with the Qiagen Mini Elute kit. Samples were quantified using Tapestation bioanalyzer

(Agilent) and the KAPA Library Quantification Kit and sequenced on the Illumina Hi-Seq 2000

using 50 cycles paired-end mode.

QUANTIFICATION AND STATISTICAL ANALYSIS

RNA-seq analysis in *CRLF2*-High and other B-ALL patient cohort

RNA-seq with replicates for individual patient samples were pooled. Paired-end reads were

mapped to the hg38 genome using TopHat2[88] (parameters:–no-coverage-search–no-

discordant–no-mixed–b2-very-sensitive–N 1). To Counts for Refseq genes were obtained using

htseq-counts[89]. PCA was performed using R to visualize the clustering of the patient samples.

Differential analysis was performed using DESeq2[32] and differential genes were considered

statistically significant if they had an adjusted p-value less than 0.01 and |log2 FoldChange| >2.

Hierarchical clustering was performed on significant differentially expressed genes (3,586) with

the manhattan distance and ward.d2 clustering method, using heatmap3 R package for

visualization. Bigwigs were obtained for visualization on individual as well as merged bam files

using Deeptools[90] (parameters bamCoverage–binSize 1–normalizeUsing RPKM). Gene Set

Enrichment Analysis (GSEA) for pathway enrichment was performed using the default

parameters of the GSEA desktop application on normalized reads counts for all patient samples

for the 3,586 differentially expressed genes between *CRLF2*-High and other B-ALL patients.

See below for information for each RNA-seq sample:

Original Label	Source	File Name	Sample Name
CRLF2-High_1	Skok Lab	CRLF2-High_1_RNA_counts.txt	CRLF2-High_1
CRLF2-High_2	Skok Lab	CRLF2-High_2_RNA_counts.txt	CRLF2-High_2
CRLF2-High_3	Skok Lab	CRLF2-High_3_RNA_counts.txt	CRLF2-High_3

CRLF2-High_4	Skok Lab	CRLF2-High_4_RNA_counts.txt	CRLF2-High_4
CRLF2-High_5	Skok Lab	CRLF2-High_5_RNA_counts.txt	CRLF2-High_5
CRLF2-High_6	Skok Lab	CRLF2-High_6_RNA_counts.txt	CRLF2-High_6
CRLF2-High_7	Skok Lab	CRLF2-High_7_RNA_counts.txt	CRLF2-High_7
CRLF2-High_8	Skok Lab	CRLF2-High_8_RNA_counts.txt	CRLF2-High_8
CRLF2-High_9	Skok Lab	CRLF2-High_9_RNA_counts.txt	CRLF2-High_9
CRLF2-High_10	Skok Lab	CRLF2-High_10_RNA_counts.txt	CRLF2-High_10
CRLF2-High_11	Skok Lab	CRLF2-High_11_RNA_counts.txt	CRLF2-High_11
CRLF2-High_12	Skok Lab	CRLF2-High_12_RNA_counts.txt	CRLF2-High_12
SRR457636	TARGET	CRLF2-High_13_RNA_counts.txt	CRLF2-High_13
SRR457642	TARGET	CRLF2-High_14_RNA_counts.txt	CRLF2-High_14
SRR457653	TARGET	CRLF2-High_15_RNA_counts.txt	CRLF2-High_15
SRR3162137	TARGET	BALL_1_RNA_counts.txt	BALL_1
SRR3162141	TARGET	BALL_2_RNA_counts.txt	BALL_2
SRR3162143	TARGET	BALL_3_RNA_counts.txt	BALL_3
SRR3162144	TARGET	BALL_4_RNA_counts.txt	BALL_4
SRR3162148	TARGET	BALL_5_RNA_counts.txt	BALL_5
SRR3162151	TARGET	BALL_6_RNA_counts.txt	BALL_6
SRR3162166	TARGET	BALL_7_RNA_counts.txt	BALL_7
SRR3162280	TARGET	BALL_8_RNA_counts.txt	BALL_8
SRR3162282	TARGET	BALL_9_RNA_counts.txt	BALL_9

SRR3162293	TARGET	BALL_10_RNA_counts.txt	BALL_10
SRR3162301	TARGET	BALL_11_RNA_counts.txt	BALL_11
SRR3162304	TARGET	BALL_12_RNA_counts.txt	BALL_12
SRR3162313	TARGET	BALL_13_RNA_counts.txt	BALL_13
SRR3162316	TARGET	BALL_14_RNA_counts.txt	BALL_14
SRR3162374	TARGET	BALL_15_RNA_counts.txt	BALL_15

ATAC-seq analysis in patient samples

ATAC-seq primary patient samples reads were mapped against the hg38 reference genome using Bowtie2[91]. ATAC-seq replicates for individual patient samples were pooled. ATAC-seq peaks were called using MACS2. A reference list of peaks or peakome representing all possible ATAC-seq peaks across patient samples was compiled by taking the union of all peaks and merged overlapping peaks. This resulted in a peakome of about 115,457 peaks. Differential analysis was performed using DESeq2[32] and differentially accessible regions were considered significant if they had an adjusted p-value < 0.01 and a |log2 fold change| >1. Hierarchical clustering was performed on significant differential ATAC-seq peaks (1678) with euclidean distance and complete clustering method with pheatmap R package. Genomic annotation of significant differential ATAC-seq peaks was done using ChIPseeker[92] with a 3kb window on either side of the TSS, to associate promoter peaks. Bigwigs were obtained for visualization using Deeptools/2.3.3[90] (parameters bamCoverage --binSize 1 --normalizeUsing RPKM). Heatmaps indicating average ATAC-signal at differentially expressed genes were generated using Deeptools.

RNA-seq analysis in patient derived cell lines

Paired-end reads were mapped to the hg38 genome using TopHat2[88] (parameters:–no-

coverage-search–no-discordant–no-mixed–b2-very-sensitive–N 1). Counts for genes were

obtained using htseq-counts[89]. Differential analysis was performed using DESeq2[32] and

differential genes were considered statistically significant if they had an adjusted p-value less

than 0.01 and |log2 FoldChange| >2.

TARGET RNA-seq analysis of 868 primary patient samples

We downloaded 868 B-ALL patient samples from the TARGET database and samples

were analyzed using the sns workflow (https://github.com/igordot/sns) to align and generate

counts for genes. Specifically, paired-end reads were mapped to the hg38 genome from

GENCODE using STAR aligner[93] and counts for the 57,316 GENCODE genes (includes coding

genes, lncRNAs and pseudogenes) were obtained. Data was normalized using DESeq2[32].

Identifying the most likely TF regulators used for network inference

To generate a list of candidate TF regulators, we identified a list of enriched TF motifs at

differentially expressed ATAC-seq peaks at promoters (793) previously annotated (**Figure 5e**) in

CRLF2-High versus Other B-ALL patients using Analysis of motif enrichment (AME[35]) to identify

enriched TF motif that are present in the Cis-BP[94] motif database. AME scores a set of regions

with a motif compared to a control sequence. Here we used randomly shuffled input sequences

as control sequences and used the one-tailed Fisher's Exact test to test for motif enrichment. TF

motifs with an adjusted p-value< $1e^{-13}$ were considered enriched TF motifs at promoters with

differentially accessible ATAC-seq peaks, which led to a list of 138 TFs. This allows us to limit

the number of TF-gene regulatory hypothesis tested.

Estimating transcription factor activities (TFA)

To generate the prior matrix from the *CRLF2*-high patient cohort, human TF binding

motifs (PWMs) were downloaded from the Cis-BP[94] motif database. Next, we used FIMO[95] to

scan for TF motifs at all ATAC-seq peaks (union of all ATAC-seq peaks in *CRLF2*-High cohort)

20kb upstream of any gene TSS. TF motifs found at each 20kb window per gene with a p-value

< 0.0001 were counted. . TF motifs were aggregated across all ATAC-seq peaks for each gene.

For each TF motif-gene association, a score, a p-value and an adjusted p-value was computed.

Only TF-gene associations that had an adjusted p-value less than 0.01 were retained.

To estimate activity, first DESeq2-normalized TARGET expression data was transposed

into matrix *X*, in which columns are individual patient samples and rows are genes. *P* is the TF-

gene matrix of known prior regulatory interactions between transcription factors (columns) and

genes (rows). $P_{i,k}$ is zero if there is no known regulatory interaction between transcription factor

k and gene *i*. *A* is the activity matrix, where the columns are the individual samples as in *X* and

rows are the TFs. We model the expression of gene *i* in individual samples *j* as a linear

combination of the activities of each TF, *k*, in condition *j*.

$$X_{i,j} = \sum_{k \in TFs} P_{i,k} A_{k,j}$$

658   This means that we use the known targets of a transcription factor to derive its activity.

659   The system of linear equations above lacks a unique solution, as there are more equations than

660   unknowns. Hence, to compute a "best fit" (least squares) to the system of linear equations

661   above that lacks a unique solution, we can compute the pseudo-inverse of the prior *P^{-1}*

662   multiplied by the expression matrix *X* to approximate *Â* :

$$\hat{A} \approx P^{-1}X$$

If a transcription factor has no prior targets present in *P*, we use the expression of that

transcription factor as a proxy for its activity.

B-ALL network inference using (Bayesian Best Subset Regression-Inferelator)

We use a bayesian best-subset regression (BBSR) method, previously described[36]

within the current Inferelator algorithm[25,26]. At steady state, we model the expression of a gene i

in individual cell j as a linear combination of the activities of its TF regulators in individual

samples j :

$$X_{i,j} = \sum_{k \in TFs} \beta_{i,k} A_{k,j}$$

The goal of the network inference procedure is to find a sparse solution for β, in which

non-zero entries define both the strength and direction (activation or repression) of a regulatory

relationship. This is because we expect a limited number of transcription factors to regulate a

particular gene. We select the model with the lowest Bayesian Information Criterion, which adds

a penalty term to the training error to account for model complexity and thereby reduce

generalization error. After this step, the output is a matrix of inferred regression parameters β,

where each entry corresponds to a regulatory relationship between transcription factor k and

gene i. Final TF-gene interactions were retained if they had a combined confidence > 0.9 and a

precision > 0.3, which resulted in 8,731 regulatory interactions between 135 TFs and 6,645

unique gene targets. Networks were visualized using jp_gene_viz, an interactive interface,

designed on iPython. Software is available at https://github.com/simonsfoundation/jp_gene_viz.

Evaluating Performance of the model

Performance of our model selection within the Inferelator is evaluated by applying a 10-

fold cross validation technique using the ATAC-motif prior. The prior matrix was split into 10

equal sized sets. One set is retained as the gold standard to test the model, while the remaining

9 sets are used as the prior. The cross-validation procedure is repeated 10 times, until each of

the 10 partitions is used exactly once as the gold standard. To generate a negative control, we

randomly shuffle the prior and employ the same 10 fold cross-validation strategy. Next, we

compute the area under the precision-recall curve (AUPR) for each of the 10 iterations and

compare the results between the prior and the shuffled prior. To test for a significant increase in

AUPR values using the ATAC-motif prior compared to a shuffled prior across the 10 iterations,

we applied a paired t-test (p-value=$4.5e^{-5}$).

in AUPR values   the Overall, AUPR were significantly higher using the ATAC-seq

derived motif prior compared to the randomly shuffled prior (**Supplemental Fig 4b**). Thus, the

final network was inferred using the ATAC-seq derived motif prior and the number of significant

regulatory interactions was limited to TF-gene interactions with combined confidences > 0.9

(high-confidence interactions). This generated a B-ALL specific regulatory network involving 135

TFs (that may be important regulators of *CRLF2*-High gene signature) and 8,731 high

confidence patient specific TF-gene interactions (**Fig. 5b**).

700
701
Estimating Transcription Factor Activity in *CRLF2*-High cohort

The B-ALL network was inferred as described above, but to approximate activity of 135

TFs in *CRLF2*-High cohort, we now use the inferred B-ALL network as prior information of TF-

gene interaction and gene expression the cohort to estimate activity. Activity values were

calculated for each *CRLF2*-High and other B-ALL patient samples. To identify the most likely

TFs altered as a result of *CRLF2* overexpression, we used to metrics to define this set of TFs.

First, we performed PCA analysis on the activity matrix of 135 TFs. The absolute PC1 values for

each TF were computed and ranked to identify the most highly variable TFs that best separate

CRLF2-High and other B-ALL patient samples, we randomly selected the absolute value of the

PC1 score > 0.75 as the most highly variable TFs. Next, we calculated the average TFA for

these TFs and a t-test was used to identify TFs with a significant difference in mean estimated

TFA between the two conditions (adjusted p-value < 0.001). As a result, we identify 36 TFs that

had a significant difference in mean estimated TFA and those with a |PC1| > 0.75.

Determining the *CRLF2*-associated sub-network affected by *CRLF2* overexpression

Using the inferred B-ALL regulatory interactions, we extract the subset of differentially expressed genes (542) regulated by the 36 significant TFs to derive the *CRLF2*-associated sub-network. Pathway analysis of these genes (Ingenuity Pathway Analysis) resulted in 22 significant pathways ($-\log_{10}$(B-H p-value) > 1.3 and a |z-score| >1). Differentially expressed gene targets in the CRLF2-associated sub-network were overlapped with convergent differentially expressed genes in patient derived cell-lines. This resulted in 67 gene targets within the *CRLF2*-associated subnetwork that are robust and convergently differentially expressed across patient and patient derived cell line samples. All sub-networks were visualized using jp_gene_viz (https://github.com/simonsfoundation/jp_gene_viz)

Validating inferred gene targets with ENCODE CRISPRi and siRNA RNA-seq samples

FOXM1 CRISPRi duplicate RNA-seq samples were downloaded from ENCSR701TVL experiment. Non-targeting control samples were downloaded from ENCSR095PIC experiment. Samples were analyzed using the sns workflow (https://github.com/igordot/sns) to align and generate counts for genes. Specifically, paired-end reads were mapped to the hg38 genome from GENCODE using STAR aligner[93] and counts for the 57,316 GENCODE genes (includes coding genes, lncRNAs and pseudogenes) were obtained. Differential expression analysis was performed using DESeq2[32] and genes that had an adjusted p-value < 0.05 and |log2FoldChange|>0.5 were considered significantly altered at the transcriptional levels. This set of genes was overlapped with the number of predicted FOXM1 gene targets to identify what percentage of predicted gene targets could be validated with the CRISPRi dataset. A hyper-geometric test was used to test whether the number of validated gene targets was statistically

significant, specifically using phyper in R as follows: phyper(50, 11918, 57316 -11918, 123,

lower.tail = FALSE)

E2F6 siRNA duplicate RNA-seq samples were downloaded from ENCSR820EGA

experiment. Non-targeting control samples were downloaded from ENCSR095PIC experiment.

Samples were analyzed using the sns workflow (https://github.com/igordot/sns) to align and

generate counts for genes. Specifically, paired-end reads were mapped to the hg38 genome

from GENCODE using STAR aligner[93] and counts for the 57,316 GENCODE genes (includes

coding genes, lncRNAs and pseudogenes) were obtained. Differential expression analysis was

performed using DESeq2[32] and genes that had an adjusted p-value < 0.05 and

|log2FoldChange|>0.5 were considered significantly altered at the transcriptional levels. This

set of genes was overlapped with the number of predicted E2F6 gene targets to identify what

percentage of predicted gene targets could be validated with the E2F6 siRNA dataset. A hyper-

geometric test was used to test whether the number of validated gene targets was statistically

significant, specifically using phyper in R as follows: phyper(57, 3836, 57316 -3836, 272,

lower.tail = FALSE)

MAZ siRNA duplicate RNA-seq samples were downloaded from ENCSR129WCZ

experiment. Non-targeting control samples were downloaded from ENCSR095PIC experiment.

Samples were analyzed using the sns workflow (https://github.com/igordot/sns) to align and

generate counts for genes. Specifically, paired-end reads were mapped to the hg38 genome

from GENCODE using STAR aligner[93] and counts for the 57,316 GENCODE genes (includes

coding genes, lncRNAs and pseudogenes) were obtained. Differential expression analysis was

performed using DESeq2[32] and genes that had an adjusted p-value < 0.05 and

|log2FoldChange|>0.5 were considered significantly altered at the transcriptional levels. This

set of genes was overlapped with the number of predicted MAZ gene targets to identify what

percentage of predicted gene targets could be validated with the MAZ siRNA dataset. A hyper-

geometric test was used to test whether the number of validated gene targets was statistically

significant, specifically using phyper in R as follows: phyper(4, 2599, 57316 -2599, 26, lower.tail

= FALSE)

Validating predicted gene targets with ENCODE ChIP-seq samples

Optimal IDR ChIP-seq peaks were downloaded for 6 TFs from ENCODE from the

following experiments:

TF	Experiment	Cells
ETS2	ENCSR596IKD	K562
TBP	ENCSR000EHA	K562
PBX2	ENCSR263DFP	K562
BCL11A	ENCSR000BHA	GM12878
IRF1	ENCSR000EGU	K562
EGR1	ENCSR211LTF	K562

Genomic annotation of optimal IDR peaks was done using ChIPseeker[92] with a 20kb window

upstream of the TSS, to associate peaks to genes. The same threshold was used to generate

TF-gene interactions for the prior matrix used for network inference. Genes with at least one

optimal IDR peak were overlapped with predicted gene targets for each TF from the B-ALL

network. This resulted in the number of predicted gene targets supported by ChIP-seq binding

of each TF of interest. Hyper-geometric tests were applied to test whether the number of

validated gene targets was statistically significant for each TF.

781
782
783

788 *Acknowledgements*
789 The authors thank all members of the Skok, Bonneau, and Carroll labs for discussions. The
790 Children's Oncology Group for providing us with the patient samples and metadata. New York
791 School of Medicine High Performance (HPC) for technical support. Adriana Heguy and the
792 Genome Technology Center (GTC) core for sequencing efforts. The TARGET database.
793
794 *Funding*
795 This work was supported by 1R21CA188968-01A1 and 1R35GM122515 (J.S), American
796 Society of Hematology Research Training Award for Fellows (B.C), AACR (P.L), National
797 Cancer Center (P.L).

798
799 *Author contributions*
800 B.C, S.B, R.B and J.S designed and managed the project. B.C, P.L, and N.E collected samples
801 and performed experiments. S.N and W.C provided experiments. S.B performed data analyses.
802 D.C, C.S, R.R, and A.W aided in analysis design. B.C, R.B, and J.S secured funding. S.B and
803 J.S wrote the manuscript. R.B and J.S revised the manuscript.
804
805 *Declaration of Interests*
806 The authors declare no competing interests.
807
808
809 **References**
810

1   Mullighan, C. G. *et al.* Genome-wide analysis of genetic alterations in acute

lymphoblastic leukaemia. *Nature* **446**, 758-764, doi:10.1038/nature05690 (2007).

2   Mullighan, C. G. *et al.* BCR-ABL1 lymphoblastic leukaemia is characterized by the

deletion of Ikaros. *Nature* **453**, 110-114, doi:10.1038/nature06866 (2008).

3   Tasian, S. K. & Loh, M. L. Understanding the biology of CRLF2-overexpressing acute

lymphoblastic leukemia. *Crit Rev Oncog* **16**, 13-24 (2011).

4   Mullighan, C. G. *et al.* JAK mutations in high-risk childhood acute lymphoblastic

leukemia. *Proc Natl Acad Sci U S A* **106**, 9414-9418, doi:10.1073/pnas.0811761106

(2009).

5   Pandey, A. *et al.* Cloning of a receptor subunit required for signaling by thymic stromal

lymphopoietin. *Nat Immunol* **1**, 59-64, doi:10.1038/76923 (2000).

6   Roll, J. D. & Reuther, G. W. CRLF2 and JAK2 in B-progenitor acute lymphoblastic

leukemia: a novel association in oncogenesis. *Cancer Res* **70**, 7347-7352,

doi:10.1158/0008-5472.CAN-10-1528 (2010).

7   Tasian, S. K. *et al.* Aberrant STAT5 and PI3K/mTOR pathway signaling occurs in human

CRLF2-rearranged B-precursor acute lymphoblastic leukemia. *Blood* **120**, 833-842

(2012).

8   Russell, L. J. *et al.* Characterisation of the genomic landscape of CRLF2-rearranged

acute lymphoblastic leukemia. *Genes Chromosomes Cancer* **56**, 363-372,

doi:10.1002/gcc.22439 (2017).

9   Russell, L. J. *et al.* Deregulated expression of cytokine receptor gene, CRLF2, is

involved in lymphoid transformation in B-cell precursor acute lymphoblastic leukemia.

Blood **114**, 2688-2698 (2009).

10   Tsai, A. G., Yoda, A., Weinstock, D. M. & Lieber, M. R.

t(X;14)(p22;q32)/t(Y;14)(p11;q32) CRLF2-IGH translocations from human B-lineage

ALLs involve CpG-type breaks at CRLF2, but CRLF2/P2RY8 intrachromosomal

deletions do not. *Blood* **116**, 1993-1994, doi:10.1182/blood-2010-05-286492 (2010).

11   Vesely, C. *et al.* Genomic and transcriptional landscape of P2RY8-CRLF2-positive

childhood acute lymphoblastic leukemia. *Leukemia* **31**, 1491-1501,

doi:10.1038/leu.2016.365 (2017).

12   Mullighan, C. G. *et al.* Rearrangement of CRLF2 in B-progenitor–and Down syndrome–

associated acute lymphoblastic leukemia. *Nature genetics* **41**, 1243-1246 (2009).

13   Harvey, R. C. *et al.* Rearrangement of CRLF2 is associated with mutation of JAK

kinases, alteration of IKZF1, Hispanic/Latino ethnicity, and a poor outcome in pediatric

B-progenitor acute lymphoblastic leukemia. *Blood* **115**, 5312-5321 (2010).

14   Tasian, S. K. *et al.* Potent efficacy of combined PI3K/mTOR and JAK or ABL inhibition in

murine xenograft models of Ph-like acute lymphoblastic leukemia. *Blood* **129**, 177-187,

doi:10.1182/blood-2016-05-707653 (2017).

15   Loh, M. L. *et al.* A phase 1 dosing study of ruxolitinib in children with relapsed or

refractory solid tumors, leukemias, or myeloproliferative neoplasms: A Children's

851   Oncology Group phase 1 consortium study (ADVL1011). *Pediatr Blood Cancer* **62**,

852   1717-1724, doi:10.1002/pbc.25575 (2015).

853   16   Maude, S. L. *et al.* Targeting JAK1/2 and mTOR in murine xenograft models of Ph-like

854   acute lymphoblastic leukemia. *Blood* **120**, 3510-3518, doi:10.1182/blood-2012-03-

855   415448 (2012).

856   17   Springuel, L., Renauld, J.-C. & Knoops, L. JAK kinase targeting in hematologic

857   malignancies: a sinuous pathway from identification of genetic alterations towards

858   clinical indications. *Haematologica* **100**, 1240-1253,

859   doi:papers3://publication/doi/10.3324/haematol.2015.132142 (2015).

860   18   Kim, S. K. *et al.* JAK2 is dispensable for maintenance of JAK2 mutant B-cell acute

861   lymphoblastic leukemias. *Genes Dev* **32**, 849-864, doi:10.1101/gad.307504.117 (2018).

862   19   Qin, H. *et al.* Eradication of B-ALL using chimeric antigen receptor-expressing T cells

863   targeting the TSLPR oncoprotein. *Blood* **126**, 629-639, doi:10.1182/blood-2014-11-

864   612903 (2015).

865   20   Brudno, J. N. & Kochenderfer, J. N. Toxicities of chimeric antigen receptor T cells:

866   recognition and management. *Blood* **127**, 3321-3330, doi:10.1182/blood-2016-04-

867   703751 (2016).

868   21   Nikolaev, S. I. *et al.* Frequent cases of RAS-mutated Down syndrome acute

869   lymphoblastic leukaemia lack JAK2 mutations. *Nat Commun* **5**, 4654,

870   doi:10.1038/ncomms5654 (2014).

871   22   Hecker, M., Lambeck, S., Toepfer, S., van Someren, E. & Guthke, R. Gene regulatory

872   network inference: data integration in dynamic models-a review. *Biosystems* **96**, 86-103,

873   doi:10.1016/j.biosystems.2008.12.004 (2009).

874   23   Alsadeq, A. *et al.* Zeta-chain-associated protein kinase-70 (ZAP-70) promotes acute

875   lymphoblastic leukemia (ALL) infiltration into the central nervous system by enhancing

876   chemokine receptor 7 (CCR7) expression. *Blood* **126**, 901 (2015).

24   Chai, L. E. *et al.* A review on the computational approaches for gene regulatory network

construction. *Comput Biol Med* **48**, 55-65, doi:10.1016/j.compbiomed.2014.02.011

(2014).

25   Bonneau, R. *et al.* The Inferelator: an algorithm for learning parsimonious regulatory

networks from systems-biology data sets de novo. *Genome Biol* **7**, R36, doi:10.1186/gb-

2006-7-5-r36 (2006).

26   Madar, A., Greenfield, A., Ostrer, H., Vanden-Eijnden, E. & Bonneau, R. The Inferelator

2.0: a scalable framework for reconstruction of dynamic regulatory network models. *Conf*

Proc IEEE Eng Med Biol Soc **2009**, 5448-5451, doi:10.1109/IEMBS.2009.5334018

(2009).

27   Miraldi, E. R. *et al.* Leveraging chromatin accessibility for transcriptional regulatory

network inference in T Helper 17 Cells. *Genome Res* **29**, 449-463,

doi:10.1101/gr.238253.118 (2019).

28   Castro, D. M., de Veaux, N. R., Miraldi, E. R. & Bonneau, R. Multi-study inference of

regulatory networks for more accurate models of gene regulation. *PLoS Comput Biol* **15**,

e1006591, doi:10.1371/journal.pcbi.1006591 (2019).

29   Arrieta-Ortiz, M. L. *et al.* An experimentally supported model of the Bacillus subtilis

global transcriptional regulatory network. *Mol Syst Biol* **11**, 839,

doi:10.15252/msb.20156236 (2015).

30   Ciofani, M. *et al.* A validated regulatory network for Th17 cell specification. *Cell* **151**,

289-303, doi:10.1016/j.cell.2012.09.016 (2012).

31   Vagace, J. M. & Gervasini, G. Chemotherapy toxicity in patients with acute leukemia.

INTECH Open Access Publisher (2011).

32   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion

for RNA-seq data with DESeq2. *Genome Biol* **15**, 550, doi:10.1186/s13059-014-0550-8

(2014).

33   Croker, B. A., Kiu, H. & Nicholson, S. E. SOCS regulation of the JAK/STAT signalling

pathway. *Semin Cell Dev Biol* **19**, 414-422, doi:10.1016/j.semcdb.2008.07.010 (2008).

34   Yoda, A. *et al.* Functional screening identifies CRLF2 in precursor B-cell acute

lymphoblastic leukemia. *Proc Natl Acad Sci U S A* **107**, 252-257,

doi:10.1073/pnas.0911726107 (2010).

35   McLeay, R. C. & Bailey, T. L. Motif Enrichment Analysis: a unified framework and an

evaluation on ChIP data. *BMC Bioinformatics* **11**, 165, doi:10.1186/1471-2105-11-165

(2010).

36   Greenfield, A., Hafemeister, C. & Bonneau, R. Robust data-driven incorporation of prior

knowledge into the inference of dynamic regulatory networks. *Bioinformatics* **29**, 1060-

1067, doi:10.1093/bioinformatics/btt099 (2013).

37   Schacht, T., Oswald, M., Eils, R., Eichmuller, S. B. & Konig, R. Estimating the activity of

transcription factors by the effect on their target genes. *Bioinformatics* **30**, i401-407,

doi:10.1093/bioinformatics/btu446 (2014).

38   Consolaro, F., Basso, G., Ghaem-Magami, S., Lam, E. W. & Viola, G. FOXM1 is

overexpressed in B-acute lymphoblastic leukemia (B-ALL) and its inhibition sensitizes B-

ALL cells to chemotherapeutic drugs. *Int J Oncol* **47**, 1230-1240,

doi:10.3892/ijo.2015.3139 (2015).

39   Zhang, G. *et al.* High ETS2 expression predicts poor prognosis in acute myeloid

leukemia patients undergoing allogeneic hematopoietic stem cell transplantation. *Ann*

Hematol **98**, 519-525, doi:10.1007/s00277-018-3440-4 (2019).

40   Fu, L. *et al.* High expression of ETS2 predicts poor prognosis in acute myeloid leukemia

and may guide treatment decisions. *J Transl Med* **15**, 159, doi:10.1186/s12967-017-

1260-2 (2017).

41   Xutao, G. *et al.* BCL11A and MDR1 expressions have prognostic impact in patients with

acute myeloid leukemia treated with chemotherapy. *Pharmacogenomics* **19**, 343-348,

doi:10.2217/pgs-2017-0157 (2018).

42   Wang, F. *et al.* Tight regulation of FOXO1 is essential for maintenance of B-cell

precursor acute lymphoblastic leukemia. *Blood* **131**, 2929-2942, doi:10.1182/blood-

2017-10-813576 (2018).

43   Tomeczkowski, J. *et al.* Absence of G-CSF receptors and absent response to G-CSF in

childhood Burkitt's lymphoma and B-ALL cells. *British journal of haematology* **89**, 771-

779 (1995).

44   Meyer, C. *et al.* Establishment of the B cell precursor acute lymphoblastic leukemia cell

line MUTZ-5 carrying a (12:13) translocation. *Leukemia* **15**, 1471-1474 (2001).

45   Smith, R. G., Dev, V. G. & Shannon, W. A., Jr. Characterization of a novel human pre-B

leukemia cell line. *J Immunol* **126**, 596-602 (1981).

46   Yuan, T. *et al.* Regulation of PI3K signaling in T-cell acute lymphoblastic leukemia: a

novel PTEN/Ikaros/miR-26b mechanism reveals a critical targetable role for PIK3CD.

Leukemia **31**, 2355-2364, doi:10.1038/leu.2017.80 (2017).

47   Juric, D. *et al.* Differential gene expression patterns and interaction networks in BCR-

ABL-positive and -negative adult acute lymphoblastic leukemias. *J Clin Oncol* **25**, 1341-

1349, doi:10.1200/JCO.2006.09.3534 (2007).

48   Ortiz-Padilla, C. *et al.* Functional characterization of cancer-associated Gab1 mutations.

Oncogene **32**, 2696-2702, doi:10.1038/onc.2012.271 (2013).

49   Poukka, M. *et al.* Acute Lymphoblastic Leukemia With INPP5D-ABL1 Fusion Responds

to Imatinib Treatment. *J Pediatr Hematol Oncol* **41**, e481-e483,

doi:10.1097/MPH.0000000000001267 (2019).

50   Laszlo, G. S. *et al.* High expression of myocyte enhancer factor 2C (MEF2C) is

associated with adverse-risk features and poor outcome in pediatric acute myeloid

leukemia: a report from the Children's Oncology Group. *J Hematol Oncol* **8**, 115,

doi:10.1186/s13045-015-0215-4 (2015).

51 Lech-Maranda, E. *et al.* Human leukocyte antigens HLA DRB1 influence clinical

outcome of chronic lymphocytic leukemia? *Haematologica* **92**, 710-711,

doi:10.3324/haematol.10910 (2007).

52 Yao, S., Fan, L. Y. & Lam, E. W. The FOXO3-FOXM1 axis: A key cancer drug target and

a modulator of cancer drug resistance. *Semin Cancer Biol* **50**, 77-89,

doi:10.1016/j.semcancer.2017.11.018 (2018).

53 Liao, G. B. *et al.* Regulation of the master regulator FOXM1 in cancer. *Cell Commun*

Signal **16**, 57, doi:10.1186/s12964-018-0266-6 (2018).

54 Li, X. *et al.* Forkhead box transcription factor 1 expression in gastric cancer: FOXM1 is a

poor prognostic factor and mediates resistance to docetaxel. *J Transl Med* **11**, 204,

doi:10.1186/1479-5876-11-204 (2013).

55 Huynh, K. M. *et al.* FOXM1 expression mediates growth suppression during terminal

differentiation of HO-1 human metastatic melanoma cells. *J Cell Physiol* **226**, 194-204,

doi:10.1002/jcp.22326 (2011).

56 Wang, I. C. *et al.* Foxm1 mediates cross talk between Kras/mitogen-activated protein

kinase and canonical Wnt pathways during development of respiratory epithelium. *Mol*

Cell Biol **32**, 3838-3850, doi:10.1128/MCB.00355-12 (2012).

57 Park, Y. Y. *et al.* FOXM1 mediates Dox resistance in breast cancer by enhancing DNA

repair. *Carcinogenesis* **33**, 1843-1853, doi:10.1093/carcin/bgs167 (2012).

58 Tan, G., Cheng, L., Chen, T., Yu, L. & Tan, Y. Foxm1 mediates LIF/Stat3-dependent

self-renewal in mouse embryonic stem cells and is essential for the generation of

induced pluripotent stem cells. *PLoS One* **9**, e92304, doi:10.1371/journal.pone.0092304

(2014).

59   Li, X. *et al.* FOXM1 mediates resistance to docetaxel in gastric cancer via up-regulating

Stathmin. *J Cell Mol Med* **18**, 811-823, doi:10.1111/jcmm.12216 (2014).

60   Carr, J. R., Park, H. J., Wang, Z., Kiefer, M. M. & Raychaudhuri, P. FoxM1 mediates

resistance to herceptin and paclitaxel. *Cancer Res* **70**, 5054-5063, doi:10.1158/0008-

5472.CAN-10-0545 (2010).

61   Liu, Y. *et al.* FOXM1 overexpression is associated with cisplatin resistance in non-small

cell lung cancer and mediates sensitivity to cisplatin in A549 cells via the

JNK/mitochondrial pathway. *Neoplasma* **62**, 61-71, doi:10.4149/neo_2015_008 (2015).

62   Kong, F. F. *et al.* FOXM1 regulated by ERK pathway mediates TGF-beta1-induced EMT

in NSCLC. *Oncol Res* **22**, 29-37, doi:10.3727/096504014X14078436004987 (2014).

63   Giangrande, P. H. *et al.* A role for E2F6 in distinguishing G1/S- and G2/M-specific

transcription. *Genes Dev* **18**, 2941-2951, doi:10.1101/gad.1239304 (2004).

64   Trimarchi, J. M., Fairchild, B., Wen, J. & Lees, J. A. The E2F6 transcription factor is a

component of the mammalian Bmi1-containing polycomb complex. *Proc Natl Acad Sci U*

S A **98**, 1519-1524, doi:10.1073/pnas.041597698 (2001).

65   Maity, G. *et al.* The MAZ transcription factor is a downstream target of the oncoprotein

Cyr61/CCN1 and promotes pancreatic cancer cell invasion via CRAF-ERK signaling. *J*

Biol Chem **293**, 4334-4349, doi:10.1074/jbc.RA117.000333 (2018).

66   Lee, W. P. *et al.* Akt phosphorylates myc-associated zinc finger protein (MAZ), releases

P-MAZ from the p53 promoter, and activates p53 transcription. *Cancer Lett* **375**, 9-19,

doi:10.1016/j.canlet.2016.02.023 (2016).

67   Tsutsui, H., Sakatsume, O., Itakura, K. & Yokoyama, K. K. Members of the MAZ family:

a novel cDNA clone for MAZ from human pancreatic islet cells. *Biochem Biophys Res*

Commun **226**, 801-809, doi:10.1006/bbrc.1996.1432 (1996).

68   Xiao, T., Li, X. & Felsenfeld, G. The Myc-Associated Zinc Finger Protein (MAZ) Works

Both Independently and Together with CTCF to Control Cohesin Positioning and

Genome Organization. *SSRN Electronic Journal*, doi:10.2139/ssrn.3549502 (2020).

69   Wang, W., Xu, S., Yin, M. & Jin, Z. G. Essential roles of Gab1 tyrosine phosphorylation

in growth factor-mediated signaling and angiogenesis. *Int J Cardiol* **181**, 180-184,

doi:10.1016/j.ijcard.2014.10.148 (2015).

70   Yin, J., Xie, X., Ye, Y., Wang, L. & Che, F. BCL11A: a potential diagnostic biomarker and

therapeutic target in human diseases. *Biosci Rep* **39**, doi:10.1042/BSR20190604 (2019).

71   Yu, Y. *et al.* Bcl11a is essential for lymphoid development and negatively regulates p53.

J Exp Med **209**, 2467-2483, doi:10.1084/jem.20121846 (2012).

72   Basak, A. & Sankaran, V. G. Regulation of the fetal hemoglobin silencing factor

BCL11A. *Ann N Y Acad Sci* **1368**, 25-30, doi:10.1111/nyas.13024 (2016).

73   Zhang, J. M. *et al.* DPEP1 expression promotes proliferation and survival of leukaemia

cells and correlates with relapse in adults with common B cell acute lymphoblastic

leukaemia. *Br J Haematol*, doi:10.1111/bjh.16505 (2020).

74   Selleri, L. *et al.* The TALE homeodomain protein Pbx2 is not essential for development

and long-term survival. *Mol Cell Biol* **24**, 5324-5331, doi:10.1128/MCB.24.12.5324-

5331.2004 (2004).

75   Schnabel, C. A., Jacobs, Y. & Cleary, M. L. HoxA9-mediated immortalization of myeloid

progenitors requires functional interactions with TALE cofactors Pbx and Meis.

Oncogene **19**, 608-616, doi:10.1038/sj.onc.1203371 (2000).

76   Qiu, Y. *et al.* Expression level of Pre B cell leukemia homeobox 2 correlates with poor

prognosis of gastric adenocarcinoma and esophageal squamous cell carcinoma. *Int J*

Oncol **36**, 651-663, doi:10.3892/ijo_00000541 (2010).

77   Inamdar, V. V., Reddy, H., Dangelmaier, C., Kostyak, J. C. & Kunapuli, S. P. The protein

tyrosine phosphatase PTPN7 is a negative regulator of ERK activation and thromboxane

generation in platelets. *J Biol Chem* **294**, 12547-12554, doi:10.1074/jbc.RA119.007735

(2019).

78   Seo, H. *et al.* Role of protein tyrosine phosphatase non-receptor type 7 in the regulation

of TNF-alpha production in RAW 264.7 macrophages. *PLoS One* **8**, e78776,

doi:10.1371/journal.pone.0078776 (2013).

79   Turazzi, N. *et al.* Engineered T cells towards TNFRSF13C (BAFFR): a novel strategy to

efficiently target B-cell acute lymphoblastic leukaemia. *Br J Haematol* **182**, 939-943,

doi:10.1111/bjh.14899 (2018).

80   Fazio, G. *et al.* TNFRSF13C (BAFFR) positive blasts persist after early treatment and at

relapse in childhood B-cell precursor acute lymphoblastic leukaemia. *Br J Haematol* **182**,

434-436, doi:10.1111/bjh.14794 (2018).

81   McWilliams, E. M. *et al.* Anti-BAFF-R antibody VAY-736 demonstrates promising

preclinical activity in CLL and enhances effectiveness of ibrutinib. *Blood Adv* **3**, 447-460,

doi:10.1182/bloodadvances.2018025684 (2019).

82   Ke, J. *et al.* The role of MAPKs in B cell receptor-induced down-regulation of Egr-1 in

immature B lymphoma cells. *J Biol Chem* **281**, 39806-39818,

doi:10.1074/jbc.M604671200 (2006).

83   Wasylyk, B., Hagman, J. & Gutierrez-Hartmann, A. Ets transcription factors: nuclear

effectors of the Ras-MAP-kinase signaling pathway. *Trends Biochem Sci* **23**, 213-216,

doi:10.1016/s0968-0004(98)01211-0 (1998).

84   Daud, S. S. *et al.* Identification of the Wnt Signalling Protein, TCF7L2 As a Significantly

Overexpressed Transcription Factor in AML. *Blood* **120**, 1281-1281,

doi:10.1182/blood.V120.21.1281.1281 (2012).

85   Hornsveld, M., Dansen, T. B., Derksen, P. W. & Burgering, B. M. T. Re-evaluating the

role of FOXOs in cancer. *Semin Cancer Biol* **50**, 90-100,

doi:10.1016/j.semcancer.2017.11.017 (2018).

86   Dinkel, A. *et al.* The transcription factor early growth response 1 (Egr-1) advances

differentiation of pre-B and immature B cells. *J Exp Med* **188**, 2215-2224,

doi:10.1084/jem.188.12.2215 (1998).

87   Buenrostro, J. D., Giresi, P. G., Zaba, L. C., Chang, H. Y. & Greenleaf, W. J.

Transposition of native chromatin for fast and sensitive epigenomic profiling of open

chromatin, DNA-binding proteins and nucleosome position. *Nat Methods* **10**, 1213-1218,

doi:10.1038/nmeth.2688 (2013).

88   Kim, D. *et al.* TopHat2: accurate alignment of transcriptomes in the presence of

insertions, deletions and gene fusions. *Genome Biol* **14**, R36, doi:10.1186/gb-2013-14-4-

r36 (2013).

89   Anders, S., Pyl, P. T. & Huber, W. HTSeq--a Python framework to work with high-

throughput sequencing data. *Bioinformatics* **31**, 166-169,

doi:10.1093/bioinformatics/btu638 (2015).

90   Ramirez, F. *et al.* deepTools2: a next generation web server for deep-sequencing data

analysis. *Nucleic Acids Res* **44**, W160-165, doi:10.1093/nar/gkw257 (2016).

91   Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat Methods*

9, 357-359, doi:10.1038/nmeth.1923 (2012).

92   Giannopoulou, E. G. & Elemento, O. An integrated ChIP-seq analysis platform with

customizable workflows. *BMC Bioinformatics* **12**, 277, doi:10.1186/1471-2105-12-277

(2011).

93   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21,

doi:10.1093/bioinformatics/bts635 (2013).

94   Weirauch, M. T. *et al.* Determination and inference of eukaryotic transcription factor

sequence specificity. *Cell* **158**, 1431-1443, doi:10.1016/j.cell.2014.08.009 (2014).

95   Grant, C. E., Bailey, T. L. & Noble, W. S. FIMO: scanning for occurrences of a given

motif. *Bioinformatics* **27**, 1017-1018, doi:10.1093/bioinformatics/btr064 (2011).

Figure 1: Genome-wide transcriptional changes in CRLF2-overexpressing primary patient samples. (a) Schematic of chromosomal alterations, IGH-CRLF2 (top) and P2RY2-CRLF2 (bottom) leading to CRLF2 overexpression. (b) PCA analysis of 15 CRLF2-High (purple, n=15) and 15 Other B-ALL (teal, n=15) patient RNA-seq samples. (c) Volcano plot of the differentially expressed genes (DESeq2: FDR=1%, |log2 fold change| >2) between CRLF2-High and other B-ALL patients. Red and blue points correspond to up and down-regulated genes (n=1470 and 2116, respectively) in CRLF2-High samples. (d) Expression heatmap of 3,586 differentially expressed genes across patient samples.

Figure 2: CRLF2-overexpression leads to promoter-enriched ATAC-seq changes in primary patient samples. (a) PCA analysis of 7 CRLF2-High (purple) and 6 Other B-ALL (teal) ATAC-seq samples. (b) Volcano plot of the differential accessible peaks (DESeq2: FDR=1%, |log2 fold change| >1) between CRLF2-High and other B-ALL patients. Red and blue points correspond to more accessible and less accessible ATAC-seq peaks, respectively (n=700 and 978,) in CRLF2-High samples. (c) Heatmap of 700 more accessible (purple bar) and 978 less accessible ATAC-seq peaks in CRLF2-High patients. (d) ChIPseeker genomic annotation of all ATAC-seq peaks or Peakome (n=115,457) across all patient samples. (e) ChIPseeker genomic annotation of significantly differential ATAC-seq peaks (n=1678). (f) Differentially ATAC-seq peaks (n=317), excluding peaks in distal intergenic regions, linked to differentially expressed genes (n=268) and the log2 fold change of gene expression (x-axis) plotted against the log2 fold change of ATAC-seq reads (y-axis). Linked genes are colored according to genomic annotation. (g) IGV screenshot of ATAC-seq tracks for individual samples at down-regulated KIT gene, RNA-seq tracks for CRLF2-High (orange) and other B-ALL (green) samples are overlayed. The highlighted region indicates a region of decreasing accessibility in CRLF2-High samples.

Figure 3: B-ALL network inference identifies significant TFs in the CRLF2-High patient cohort. (a) Workflow for construction of B-ALL regulatory network using Bayesian Best Subset Regression with the Inferelator algorithm. Inferelator resulted in a B-ALL regulatory network involving interactions (n=8,731) with combined confidences > 0.9 between 135 TF and 6,645 unique gene targets. (b) PCA of 30 primary patient samples computed according to estimated TF activity values of 135 TFs. (c) Volcano plot of the significant differential TFs (FDR=0.1%, |PC1 value| >0.075) between CRLF2-High and other B-ALL patients. Red points indicate highly variable TFs between two conditions with significant differential mean activity (t-test; q-value<0.001).

Figure 4: Significant TFs predicted to regulate conserved differentially expressed gene targets across patients and cell lines. (a) Filtered B-ALL network involving regulatory interactions between significant TFs (n=34) and their differentially expressed gene targets (n=542). TFs are the source nodes labeled in black and the gene targets are the blue target nodes. The size of the source nodes reflects the number of targets each TF has. TF activation and repression of a gene are represented by red and blue edges, respectively. (b) Significant pathways (–log10(B-H p-value > 1.3 & |z-score| > 1) identified from analysis of differential gene targets in the CRLF2 specific sub-network are shown at the top of the figure, ranked by the –log10(Benjamini-Hochberg p-value). Z-score in the bar plot indicates sign of the pathway with red and blue representing up- and down-regulation of the pathway. (c) Log2 fold changes of all the convergent differentially expressed gene targets in CRLF2-High vs other B-ALL patients and cell lines (CALL vs SMS, MUTZ vs SMS).

h

TF of Interest	Experiment	Predicted Gene Targets per TF	# Optimal IDR ChIP-seq Peaks	# of Genes associated with atleast one Peak	# of Validated Predicted Gene Targets with TF ChIP-seq peak
IRF1	ChIP-seq (K562)	19	1,483	1,289	5
BCL11A	ChIP-seq (GM12878)	136	20,663	8,452	56
TBP	ChIP-seq (K562)	253	22,063	11,411	121
PBX2	ChIP-seq (K562)	135	36,585	12,961	77
EGR1	ChIP-seq (K562)	323	30,044	13,161	223
ETS2	ChIP-seq (K562)	277	1,705	1,216	17

Figure 5: Validating inferred gene targets of TFs significantly altered by *CRLF2* overexpression. (a) Bar plot representing the number of predicted gene targets for 9 TFs, for which public data was available for validation. Exact numbers are represented at the top of the bar. (b) PCA analysis of 4 CRISPRi targeting FOXM1 (red) and non-targeting control (blue) RNA-seq samples. (c) Volcano plot of the FOXM1 predicted gene targets, those that are differentially expressed genes (FDR=5%, |log2 fold change| >0.5) between FOXM1 CRISPRi and non-targeting control samples are highlighted in red and labeled. (d) PCA analysis of 4 CRISPRi targeting E2F6 (red) and non-targeting control (blue) RNA-seq samples. (e) Volcano plot of the E2F6 predicted gene targets, those that are differentially expressed genes (FDR=5%, |log2 fold change| >0.5) between E2F6 CRISPRi and non-targeting control samples are highlighted in red and labeled. (f) PCA analysis of 4 CRISPRi targeting MAZ (red) and non-targeting control (blue) RNA-seq samples. (g) Volcano plot of the MAZ predicted gene targets, those that are differentially expressed genes (FDR=5%, |log2 fold change| >0.5) between MAZ CRISPRi and non-targeting control samples are highlighted in red and labeled. (h) Table representing the ENCODE ChIP-seq experiments for 6 TFs, including the TF of interest, the experiment, number of predicted gene targets from the B-ALL network, number of optimal IDR ChIP-seq peaks, number of annotated genes with at least one ChIP-seq peak, and finally the number of validated gene targets for each TF of interest. (i) Barplot representing the percentage of validated gene targets for each of the 9 TFs of interest. The percentage of validated inferred gene targets for each TF is labeled in white at the top of the bar. P-values obtained from hyper-geometric test are annotated as follows: NS, not significant; *, p < 0.05, p<0.01, p<0.001. Color of the bar represents the experiment.

Figure 6: Proposed Model. (a) Inferred transcriptional regulatory interactions downstream of CRLF2-mediated signaling in B-ALL.