1   Universal latent axes capturing Parkinson's patient deep phenotypic variation reveals patients

2   with a high genetic risk for Alzheimer's disease are more likely

3   to develop a more aggressive form of Parkinson's.

Cynthia Sandor DMV,PhD[1†], Stephanie Millin DPhil[2†], Andrew Dahl DPhil[3], Michael

Lawton PhD[4], Leon Hubbard PhD[5], Bobby Bojovic MS[3], Marine Peyret-Guzzon PhD[3],

Hannah Matten MS[3], Christine Blancher PhD[3], Nigel Williams PhD[5], Yoav Ben-Shlomo

PhD[4], Michele T. Hu MD, PhD[6,7], Donald G. Grosset MD, PhD[8], Jonathan Marchini PhD[3,9,*],

Caleb Webber PhD[1,2,*]

Affiliations:

[1]UK Dementia Research Institute, Cardiff University, Cardiff, UK

[2]Department of Physiology, Anatomy and Genetics, University of Oxford, Oxford, UK

[3]Wellcome Centre for Human Genetics, University of Oxford, Oxford, UK

[4]School of Social and Community Medicine, University of Bristol, Bristol, UK

[5]MRC Centre for Neuropsychiatric Genetics and Genomics, Institute of Psychological

Medicine and Clinical Neurosciences, School of Medicine, Cardiff University, Cardiff, UK

[6]Oxford Parkinson's Disease Centre, Department of Physiology, Anatomy and Genetics, Le

Gros Clark Building, University of Oxford, Oxford, UK

[7]Nuffield Department of Clinical Neurosciences, Division of Clinical Neurology, University

of Oxford, Oxford, UK.

[8]Department of Neurology, Institute of Neurological Sciences, Queen Elizabeth University

Hospital, Glasgow, United Kingdom

[9]Department of Statistics, University of Oxford, Oxford, UK

*To whom correspondence should be addressed: E-mail: webberc4@cardiff.ac.uk &

marchini@stats.ox.ac.uk

†Both authors contributed equally to this work.

Abstract

The generation of deeply phenotyped patient cohorts offers an enormous potential to identify

disease subtypes but are currently limited by the cohort size and the heterogeneity of the

clinical assessments collected across different cohorts. Identifying the universal axes of clinal

severity and progression is key to accelerating our understanding of how disease manifests

and progresses. These universal axes would accelerate our understanding of how Parkinson's

disease (PD) manifests and progresses through which patients may be appropriately

compared appropriately stratified, and personalised therapeutic strategies and treatments can

be developed and targeted. We developed a Bayesian multiple phenotype mixed model

incorporating the genetic relationships between individuals which is able to reduce a wide-

array of different clinical measurements into a smaller number of continuous underlying

factors named phenotypic axis. We identify three principal axes of PD patient phenotypic

variation which are reproducibly found across three independent, deeply and diversely

phenotyped cohorts. Together they explain over 75% of the observed clinical variation and

remain robustly captured with a fraction of the clinically-recorded features. The most

influential axis was associated with the genetic risk of Alzheimer's disease (AD) and involves

genetic pathways associated with neuroinflammation. Our results suggest PD patients with a

high genetic risk for AD are more likely to develop a more aggressive form of PD including,

but not limited to, dementia.

Introduction

A critical challenge in medicine is to understand why the clinical presentations of

each patient affected by the same disorder vary. This is especially true for Parkinson's disease

(PD), for which the age of onset, the rate of progression, type and severity of symptoms differ

across more than a million people worldwide living with this disease [1]. To accelerate the

identification of disease subtypes, large deeply phenotyped cohorts of PD patients have been

created, in which valuable clinical, imaging, biosample and genetic data has been collected,

and increasingly with longitudinal monitoring [2-4].

Recent studies exploiting these deeply phenotyped cohorts have classified patients

into discrete phenotypic subgroups, each displaying a characteristic set of symptoms [5-7]. To

define PD subtypes, most of these studies employ some form of variable selection to create a

distance matrix between individuals, followed by clustering methods such as k-means or

hierarchical clustering. These methods provide discrete phenotypic groups, which are

appealing in their categorical nature but have many shortfalls. Firstly, while selection

methods quantify how much variance each phenotype explains, no robust method was used to

define a threshold for this measure above which a phenotype contributes to the distance

matrix. Consequently, the definition of which phenotypes are essential to group patients and

which are irrelevant can be somewhat arbitrary. For example, two recent studies [5, 8], using the

same Parkinson's Progression Markers Initiative (PPMI) cohort show divergent results:

apathy and hallucinations were key subtype classifiers in the first study [8], but not in the

second one [5], because these variables were not included. Secondly, K-means clustering

requires the number of phenotypic groups to be prespecified, and this choice has the potential

to be biased towards preconceived expectations with smaller groups ignored or erroneously

joined with larger groups.   Finally, the creation of discrete groups may not reflect the

possibly continuous nature of phenotypic variability and ignores the greater statistical power

of continuous traits.

To overcome these limitations, we propose here an approach focused on the continuous variation of phenotypes. Rather than focusing on presence versus absence, or mild versus severe phenotypes, we incorporate the whole spectrum of severity displayed across the population. For this, we applied PHENIX (PHENotype Imputation eXpediated), a multiple phenotype mixed model (MPMM) approach initially developed to impute missing phenotypes [9], that can also be exploited for genetically-guided dimensionality reduction of multiple traits. This approach models the phenotypes as a combination of genetic and environmental factors and the genetic component is computed from the correlation matrix between the individual's genetic data.

Applying PHENIX to the deeply phenotyped UK-based *Discovery* cohort, we identify a small number of axes underlying individual PD patient phenotypic variation that explain the variation in the much larger number of clinically-observed phenotypes. We demonstrate the universality of these axes of phenotypic variation amongst PD patients by independently deriving similar axes in each of three cohorts: UK *Tracking* cohort including 1807 individuals, the UK *Discovery* cohort including 842 PD patients and US PPMI cohort including 439 PD patients that has a different clinical structure from the UK cohorts. We show that this reproducibility is not achieved by other commonly-used dimensionality-reduction methods. Finally, we demonstrate that the most influential axis was associated with the genetic risk of Alzheimer's disease (AD) suggesting PD patients with a high genetic risk for Alzheimer's disease are more likely to develop a more aggressive form of PD including dementia symptoms.

Materials and Methods

Discovery cohort

We considered 842 PD cases from the *Discovery* cohort constituted of 1700 subjects,

including over 1000 people with Parkinson's, plus 320 healthy controls and 340 individuals

thought to be 'at-risk' of developing future Parkinson's. Individuals were required to have at

least 90% chance of PD according to UK-Parkinson's disease brain bank criteria, no

alternative diagnosis and disease duration less than 3.5 years. All patients have a clinical

assessment repeated every eighteen months and have been already described[4,6]. Phenotype

data were collected for over a hundred clinical attributes, affecting autonomic, neurological

and motor phenotypes (**Supplementary Fig. 1**) and described in the **Supplementary Table**

1. Genotype data were generated using the Illumina HumanCoreExome-12 v1.1 and Illumina

InfiniumCoreExome-24 v1.1 SNP arrays.

UK Tracking Parkinson's study

We considered 1807 PD cases from the *Tracking* Parkinson's cohort, which was

already described in detail by Malek *et al.*[2] and was used to identify the impact of mutations

within glucocerebrosidase gene (*GBA*) on different PD clinicals manifestations[10]. Genotype

data were generated using the Illumina Human Core Exome array.

PPMI cohort

The PPMI cohort (http://www.ppmi-info.org) was already described in detail

(including PPMI protocol of recruitment and informed consent) by Marrek *et al.*[11]. We

downloaded data from the PPMI database on September 2017 in compliance with the PPMI

Data Use Agreement. We considered 472 newly-diagnosed typical PD subjects: subjects with

a diagnosis of PD for two years or less and who are not taking PD medications. We used the

baseline (t=0) of clinical assessments, described in detail in the **Supplementary Table 2**. We

excluded any individual with > 5% of missing data (437 individuals included). Participants

have been genotyped using two genotyping arrays, ImmunoChip [12] and NeuroX [13], [14]. As

more participants were genotyped on NeuroX array, we used the genotype data of the

NeuroX chip.

Methods

Genotype: quality control & Imputation

Quality control was carried out independently using PLINK v1.9 [15] (SI). Imputation of

unobserved and missing variants was carried out separately for each cohort (SI)

Phenotypic axis

Our continuous measures of severity are based on a multiple phenotypes mixed model

approach (MPMM) named PHENIX (PHENotype Imputation eXpediated) which includes

genetic relationships between individuals, and is designed to impute missing phenotypes [9].

To impute missing phenotypes, PHENIX reduces the variation within a cohort to a smaller

number of underlying factors that are then used to predict individual missing values. Here,

we exploit the identification of these underlying factors as providing the latent axes of patient

variation which underlie a larger number of clinically observed phenotypes. The outcome is

that the many clinical phenotypes (sometimes missing for some individuals) of each

individual are represented through a smaller number of underlying latent variables of

phenotypic variation that manifest the observed clinical phenotypes, which we name herein

as *phenotypic axes*.

PHENIX [9] use a Bayesian multiple-phenotype mixed model (MPMM), where the correlations

between clinical phenotypes (Y) are decomposed into a genetic and a residual component

with the following model: Y=U+e, where U represents the aggregate genetic contribution

(whole genotype) to phenotypic variance and e is idiosyncratic noise. As the estimation of

maximum likelihood covariance estimates can become computationally expensive with

increasing number of phenotypes, PHENIX uses a Bayesian low-rank matrix factorization

model for the genetic term U such as: $U = S\beta$, in which β is can be used to estimate the

genetic covariance matrix between phenotypes and S represents a matrix of latent

components that each follow $\sim N\,(0, G)$ where G is the Estimate of Relatedness Matrix from

genotypes. The resulting latent traits (S) are used as phenotypic axes, each representing the

severity of a number of non-independent clinical phenotypes. The details to run PHENIX and

extract the phenotypic axes are given in the **Supplemental Information.**

Disease phenotypic axis

We derived disease phenotypic axis consisting to replace the general genetic component in a

MPMM by a disease risk genetics component. To calculate a disease relatedness matrix, we

considered only genetic variants (after pruning) associated with human complex traits. For

different complex human traits with GWA results publically available (**Supplementary**

Table 01), we calculated a disease relatedness, that we used subsequently to derive

phenotypic axes (**Supplementary Information**).

Results

Three continuous measures capture 75% of the clinical variation.

Initially, we generated phenotypic axes from a cohort of 842 PD patients (*Discovery* cohort)

which had been genotyped and phenotypically characterised with 40 clinical assessments

(**Supplementary Table 1**). Each latent axis reflected a number of co-varying observed

clinical assessments. Among the phenotypic axes that explained more than 5%, Axes 1, 2 and

3 explained 39.6%, 28.7% and 6.8% of the clinical variation respectively. Together, these 3

top axes account for over 75% of the clinically-observed variation (**Supplementary Fig. 2**).

To examine whether similar phenotypic axes are obtained in different deeply phenotyped PD

cohorts, we derived phenotypic axes within an independent cohort of 1807 PD individuals

from the UK *Tracking* cohort [2] that had made similar clinical observations to the *Discovery*

cohort. We found significant Pearson's correlation coefficients between each cohort's first

three phenotypic axes: Axis 1 r=0.92 (p=3 x 10^{-13}), Axis 2 r=0.89 (p=4 x 10^{-11}), Axis 3

r=0.72 (p=5 x 10^{-6}) (**Fig. 1**). Nevertheless, a major concern was that the identification of the

same phenotypic axes might, at least in part, be due to the very similar structure of the

clinical phenotyping between the two UK cohorts. To address this, we examined the

independent US-based PPMI cohort consisting of 439 sporadic PD individuals that had been

clinically phenotyped following a substantially different protocol to the UK cohorts. After

deriving phenotypic axes in the PPMI cohort, we found significant similarities between the

first three phenotypic axes derived for both the *Discovery*-UK and PPMI-US cohorts: the

coefficients of determination (R^2) between three first axes across different categories of

clinical phenotypes from each cohort were: Axis1: 0.665 (p=0.048), Axis 2: 0.914 (p=0.003)

and Axis 3: 0.754 (p=0.025) (**Fig. 2 & Supplementary Figure 3**). These consistent

similarities in the axes of phenotypic variation independently derived for each of three

different PD cohorts demonstrates the reproducibility of these axes of phenotypic variation

amongst Parkinson's patients. Finally, by comparing PHENIX with other methods of

dimensionality   reduction,   specifically   Principle   Component   Analyses   (PCA),

Multidimensional Scaling (MDS) and Independent component analysis (ICA), only the

dimensions discovered by the MPMM model, PHENIX, were significantly correlated

between both cohorts and thus no other method was able to identify similar axes of

phenotypic variation across UK and US PD cohorts (**Fig. 2**).

Each phenotypic axis represents a distinct set of clinical features

To interpret the clinical relevance of each phenotypic axis, we examined the

correlation between individual clinical features and the phenotypic axes (**Table 1 & Fig. 1 &**

Supplementary Figure 4). We observed that each phenotypic axis corresponded to a subset

of clinical features, differing in both extents and directions of severity. Axis 1 represented

worsening non-tremor motor phenotypes, anxiety and depression accompanied by a decline

of the cognitive function (**Table 1 & Fig. 3**). Worsening anxiety and depression were also

features of Axis 2, in addition to increasing severity of autonomic symptoms and increasing

motor dysfunction. Axis 3 was associated with general motor symptom severity including

rigidity, bradykinesia and tremor of the whole body independently of non-motor features.

The contribution of different phenotypes to these axes was therefore highly variable. Specific

aspects of motor dysfunction were important factors in defining the majority of axes. Anxiety

and depression were also relatively important features, but only for axes explaining the

largest amounts of variation. Conversely, cognitive impairment was associated only with

Axis one. However, this observation must be weighted by the fact that cognitive

impairment/dementia are reported at a later disease stage and thus likely under-represented in

recently diagnosed cases.

Although each phenotypic axis is associated with a distinct set of clinical features,

they are not independent but instead strongly correlated (**Supplementary Figure 5**). We find

no significant relation between the phenotypic axes and principal components of genetic

ancestry (**Methods**) suggesting that the phenotypic axes are not biased by the population

structure (**Supplementary Figure 5, Supplementary Table 3**). However, as previously

reported, gender influences clinical symptoms [4] and we also observe a significant association

between gender and Axis 2 (**Supplementary Table 3,** p=4.5×10^{-5}).

To assess to what extent the phenotypic axes might be affected by the number of clinical

observations, within the *Discovery* cohort we compared the phenotypic axes built on all

clinical features with phenotypic axes generated with incomplete sets of randomly-selected

clinical features. We observed a strong correlation ($r > 0.8$) between each of the two first

phenotypic axes built with as few as 50% of the clinical variables and their respective

original phenotypic axes, suggesting that these two axes are extremely robust in terms of the

numbers of clinical variables considered (**Supplementary Figure 6**).

The integration of genetic relationships between patients improves capture of the

Parkinson's disease clinical variation and reproducibility.

The PHENIX MPMM approach employed here to derive phenotypic axes exploits the

genetic relatedness between individuals derived from genotypic similarity to further

decompose random effects into kinship effects between individuals. In its original application

to imputing missing phenotypes, PHENIX outperforms other imputation approaches when

the heritability (h^2) of a phenotype increased [9]. Similarly, when randomly removing and re-

imputing 10% of observed data, the quality of the imputation of PD clinical assessments was

in general better when considering the genetic relatedness between individuals as compared

to excluding this information (**Supplementary Figure 7**), suggesting that the resulting

phenotypic axes better capture PD heterogeneity when including genetic information.

Moreover, we found a higher agreement between the phenotypic axes derived by integrating

the genetic relationship between patients of different cohorts than when the phenotypic axes

were derived ignoring the genetic relationships (**Supplementary Figure 8**). Specifically, the

coefficient of determination reflecting the agreement between the axes derived from the

Discovery and those derived from the PPMI cohorts were from Axis 1 to 3: 0.665 (p=0.048),

0.914 (p=0.003) and 0.754 (p=0.025) when including the genetic similarity between patients

as compared to 0.604 (p=0.069), 0.908 (p=0.003) and 0.001 (p=0.991) without. Together,

these findings demonstrate that the integration of genetic relationship between patients

enhances the resulting phenotypic axes' ability to reproducibly capture PD clinical variation.

Metanalysis of Genome Wide Association Studies with phenotypic axes as unique and

universal quantitative traits

Each phenotypic axis provides a quantitative trait enabling the genetics underlying

patient variation to be studied by performing a Genome Wide Association Study (GWAS) via

a regression model with the covariates age, gender, and two genetic principal components (to

account for any underlying population substructure) in each individual cohort. As three

phenotypic axes were similar across each individual cohort (*Discovery*, *Tracking* and PPMI)

and to increase statistical power to detect an significant association, we conducted a meta-

analysis of each phenotypic axis genome-wide association studies using a common set of

4211937 variants across 3088 individuals. A significant departure from the expected

quantiles was observed for Axis 1 (meta-analysis combining the summary statistic of three

individual GWAS [*Discovery*-Tracking-PPMI]) (**Supplementary Figure 9**), but no variant

surpassed genome-wide significance (**Supplementary Figure 10**). Although we did not

observe a significant genome-wide association, the use of universal phenotypic axes

significantly unable us to conduct meta-analysis and thus to increase the statistical power to

identify genetic variants through their ability to align differently deeply phenotyped cohorts

and reduce the number of traits tested.

Next, we re-examined genetic associations for each of the three phenotypic axes for

three major PD risk genes, namely *SNCA*, *GBA* and *LRRK2*. We found a indicative local

association signal but however un-significant at GWA level with Phenotypic Axis 1 for a

variant in *SNCA*: 4: 90758437 (p-value=1.7×10^{-4}, **Supplementary Figure 11A**) which is in

high LD with rs1348224 ($r^2 > 0.8$), a SNP previously associated with PD with dementia and

dementia with Lewy bodies [20]. SNP rs1348224:G allele (minor allele) had a negative effect

on Phenotypic Axis 1, thus a protective effect for cognitive impairment, which is consistent

with a protective effect for PD with dementia and dementia with Lewy bodies previously

reported for this locus   [20] . We also found a indicative local association signal (p-

value=1.1x10^{-4}) with Phenotypic Axis 3 for an intronic variant in *LRRK2* (**Supplementary**

Figure 11B). Both *SNCA* and *LRRK2* variants were each nominally associated with only one

phenotypic axis (**Supplementary Table 4**), suggesting distinct pathogenic mechanisms.

Parkinson patients carrying a high genetic risk for Alzheimer's are more

likely to develop a more aggressive form of Parkinson's

To better understand the genetics risk factors influencing the phenotypic axis, we

calculated a disease-risk relatedness matrix in the MPMM, based on genetic variants

associated with different complex human traits. For example, replacing the overall genetic

similarity by how similar people are in their risk of diabetes or depression. By examining the

proportion of phenotypic variation explained by different phenotypic axis derived using these

different disease risks as compared to the original phenotypic axes derived using the entire

genotype, we showed that the phenotypic axis derived using Alzheimer's disease (AD)

genetic risk significantly outperforms (capture more patient phenotypic variation) the original

phenotypic axes (**Fig. 4**)

This result proposes that PD patients carrying a high genetic risk for AD are more likely to

develop a more aggressive form of PD including dementia symptoms: Axis 1 represents

worsening non-tremor motor phenotypes, anxiety and depression accompanied by a decline

of the cognitive function (preprint Table 1 & Fig. 2) Testing this hypothesis in the PPMI

cohort, we found a significant relationship between Phenotypic Axis 1 and CSF Aβ1-42, a

biomarker strongly associated with future conversion to dementia. Secondly, as for AD

genetics risk and unlikely to PD genetics risk, we found an association of phenotypic axis 1

risk variants with microglia-expressed genes, in both the SN and the cortex suggesting that

the neuro-inflammation play a key role in the development a more aggressive form of PD,

but not in the PD onset-risk. Finally we observed the phenotypic axis one is associated with

rapid progression of multiple clinical symptoms suggesting that AD genetic risk score in PD

patients could be used as a predictor of progression

Discussion

We propose here a novel approach to quantifying diverse patient phenotypes on a continuous scale via the use of phenotype axes. This approach overcomes many of the limitations associated with the clustering methods previously used to classify PD heterogeneity. By applying our approach to three independent and deeply phenotyped cohorts, we demonstrate the universality of these axes of phenotypic variation amongst PD patients. We also showed that our axes are robustly derived when reducing the number of clinical features considered and, unlike other dimensionality reduction methods, the PHENIX MPMM approach is the only method tested here that is able to identify the same phenotypic axes underlying PD patient variation between individuals from different cohorts. The phenotypic axes have multiple applications in PD precision medicine. We found that PD patients carry on a high genetic risk load for Alzheimer's disease can develop a more clinical aggressive PD form including dementia symptoms.

Our approach was able to identify representative quantitative variables that are clinically relevant to previously-defined categorical PD subtypes. A number of known comorbidities were represented among the phenotype axes. Anxiety and depression are highly correlated in PD patients, both of which are correlated with Axes 1 and 2 [26]. Rigidity and bradykinesia are also linked, possibly due to shared physiology [27], and varied in the same direction along Axis 3. Lawton *et al.* reported five PD subgroups, by using the same *Discovery* cohort but following a k-means clustering approach [6]. We examined the distribution of phenotypic axis score across these five PD subgroups (**Supplementary Figure 16**) and noted that the 5th subgroup of patients, characterised by severe motor, non-motor and cognitive disease, with poor psychological well-being clinical symptoms, were systematically associated with high severity score for all three of our phenotypic axes. Inversely, the first PD subgroup characterised by mild motor and non-motor disease (group affected by fewer

clinical symptoms) displayed a low severity score for our three phenotypic axes.

Furthermore, we observed that the individuals of subgroups 4 and 5, characterised by poor

psychological well-being, had high severity scores for phenotypic axis 2, the axis most

associated with depression and anxiety symptoms. These observations demonstrate some

consistency between subgroups defined with k-means and our phenotypic axis severity score.

The agreement of these phenotype axes with previously observed correlations provides

further support for underlying biological themes, but their reinterpretation as robust

continuous traits likely provides a better approximation of how the underlying biology

contributes, as opposed to a cut-off off for a phenotype. Specifically, the unimodal character

of the phenotypic axis distributions (**Supplementary Figure 17**) suggests here that the

development of continuous measures is more appropriate than clustering according to an

arbitrary threshold.

The phenotypic axes identified were robust in terms of the number of clinical features

considered and enable the alignment of patients from different cohorts with different clinical

phenotyping structures. The corollary is that Phenix did not require the variable selection

common in PD clustering approaches, and it can also guide clinicians in determining which

clinical assessments are essential to capture PD heterogeneity. Deep phenotyping is

burdensome to both patient and clinician and many of the measures exploited here are

compound scores summarising aspects of functioning. Further work identifying the

minimally burdensome observations that enable robust scoring of patients along these

phenotypic axes would facilitate their utility and adoption across the PD clinical community,

bringing increased power to the discovery of influencing factors. Finally, the MPMM

approach can be readily extended to include longitudinal data to determine the phenotypic

axes associated with disease progression while simultaneously dealing with missing data,

which is a common problem in longitudinal studies.

In conclusion, these universal axes have the potential to accelerate our understanding

of how PD presents in individual patients, providing more robust and objective quantitative

traits through which patients may be appropriately compared, through which the underlying

disease-modifying mechanism can be understood and appropriately stratified/personalised

therapeutic strategies and treatments can be developed.

Acknowledgments

The work was supported by the Monument Trust Discovery Award from Parkinson's UK.

Oxford Genomics Centre at the Wellcome Centre for Human Genetics, Oxford is Funded by

Wellcome Trust (grant reference 090532/Z/09/Z and MRC Hub grant G0900747 91070)

Samples and associated clinical data were supplied by the Oxford Parkinson's Disease Centre

(OPDC) study, funded by the Monument Trust Discovery Award from Parkinson's UK, a

charity registered in England and Wales (2581970) and in Scotland (SC037554), with the

support of the National Institute for Health Research (NIHR) Oxford Biomedical Research

Centre based at Oxford University Hospitals NHS Trust and University of Oxford, and the

NIHR Comprehensive Local Research Network. CW is supported by a UK DRI fellowship

funded by Medical Research Council (MRC), Alzheimer's Society and Alzheimer's Research

UK. CW and CS are supported by Computational Science Program funded by Michael J. Fox

Foundation. JM acknowledges funding for this work from the European Research Council

(ERC; grant 617306) . We thank the Oxford Genomics Centre at the Wellcome Centre for

Human Genetics, Oxford) for the generation genotyping data.

Conflict of interest

24   The authors declare that they have no competing interests.

References

1.   Foltynie T, Brayne C, Barker RA. The heterogeneity of idiopathic Parkinson's disease. *J Neurol* 2002; **249**(2)**:** 138-145.

2.   Malek N, Swallow DM, Grosset KA, Lawton MA, Marrinan SL, Lehn AC *et al.* Tracking Parkinson's: Study Design and Baseline Patient Data. *J Parkinsons Dis* 2015; **5**(4)**:** 947-959.

3.   PPMI. PPMI.

4.   Szewczyk-Krolikowski K, Tomlinson P, Nithi K, Wade-Martins R, Talbot K, Ben-Shlomo Y *et al.* The influence of age and gender on motor and non-motor features of early Parkinson's disease: initial findings from the Oxford Parkinson Disease Center (OPDC) discovery cohort. *Parkinsonism Relat Disord* 2014; **20**(1)**:** 99-105.

5.   Fereshtehnejad SM, Zeighami Y, Dagher A, Postuma RB. Clinical criteria for subtyping Parkinson's disease: biomarkers and longitudinal progression. *Brain* 2017; **140**(7)**:** 1959-1976.

6.   Lawton M, Baig F, Rolinski M, Ruffman C, Nithi K, May MT *et al.* Parkinson's Disease Subtypes in the Oxford Parkinson Disease Centre (OPDC) Discovery Cohort. *J Parkinsons Dis* 2015; **5**(2)**:** 269-279.

7.   Lawton M, Ben-Shlomo Y, May MT, Baig F, Barber TR, Klein JC *et al.* Developing and validating Parkinson's disease subtypes and their motor and cognitive progression. *Journal of Neurology, Neurosurgery & Psychiatry* 2018.

8.   Erro R, Picillo M, Vitale C, Palladino R, Amboni M, Moccia M *et al.* Clinical clusters and dopaminergic dysfunction in de-novo Parkinson disease. *Parkinsonism Relat Disord* 2016; **28:** 137-140.

9.   Dahl A, Iotchkova V, Baud A, Johansson A, Gyllensten U, Soranzo N *et al.* A multiple-phenotype imputation method for genetic studies. *Nat Genet* 2016; **48**(4)**:** 466-472.

10.   Malek N, Weil RS, Bresner C, Lawton MA, Grosset KA, Tan M *et al.* Features of GBA-associated Parkinson's disease at presentation in the UK Tracking Parkinson's study. *J Neurol Neurosurg Psychiatry* 2018.

11.   Parkinson Progression Marker I. The Parkinson Progression Marker Initiative (PPMI). *Prog Neurobiol* 2011; **95**(4)**:** 629-635.

12.   Parkes M, Cortes A, van Heel DA, Brown MA. Genetic insights into common pathways and complex relationships among immune-mediated diseases. *Nat Rev Genet* 2013; **14**(9)**:** 661-673.

13. Nalls MA, Bras J, Hernandez DG, Keller MF, Majounie E, Renton AE *et al.* NeuroX, a fast and efficient genotyping platform for investigation of neurodegenerative diseases. *Neurobiol Aging* 2015; **36**(3)**:** 1605 e1607-1612.

14. Nalls MA, Keller MF, Hernandez DG, Chen L, Stone DJ, Singleton AB *et al.* Baseline genetic associations in the Parkinson's Progression Markers Initiative (PPMI). *Mov Disord* 2016; **31**(1)**:** 79-85.

15. Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MA, Bender D *et al.* PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet* 2007; **81**(3)**:** 559-575.

16. Andreassen OA, Zuber V, Thompson WK, Schork AJ, Bettella F, Consortium P *et al.* Shared common variants in prostate cancer and blood lipids. *Int J Epidemiol* 2014; **43**(4)**:** 1205-1214.

17. Andreassen OA, Djurovic S, Thompson WK, Schork AJ, Kendler KS, O'Donovan MC *et al.* Improved detection of common variants associated with schizophrenia by leveraging pleiotropy with cardiovascular-disease risk factors. *Am J Hum Genet* 2013; **92**(2)**:** 197-209.

18. Zuber V, Jonsson EG, Frei O, Witoelar A, Thompson WK, Schork AJ *et al.* Identification of shared genetic variants between schizophrenia and lung cancer. *Sci Rep* 2018; **8**(1)**:** 674.

19. Winsvold BS, Bettella F, Witoelar A, Anttila V, Gormley P, Kurth T *et al.* Shared genetic risk between migraine and coronary artery disease: A genome-wide analysis of common variants. *PLoS One* 2017; **12**(9)**:** e0185663.

20. Guella I, Evans DM, Szu-Tu C, Nosova E, Bortnick SF, Group SCS *et al.* alpha-synuclein genetic variability: A biomarker for dementia in Parkinson disease. *Ann Neurol* 2016; **79**(6)**:** 991-999.

21. Andreassen OA, Thompson WK, Schork AJ, Ripke S, Mattingsdal M, Kelsoe JR *et al.* Improved detection of common variants associated with schizophrenia and bipolar disorder using pleiotropy-informed conditional false discovery rate. *PLoS Genet* 2013; **9**(4)**:** e1003455.

22. Noyce AJ, Kia DA, Hemani G, Nicolas A, Price TR, De Pablo-Fernandez E *et al.* Estimating the causal influence of body mass index on risk of Parkinson disease: A Mendelian randomisation study. *PLoS Med* 2017; **14**(6)**:** e1002314.

23. Brainstorm C, Anttila V, Bulik-Sullivan B, Finucane HK, Walters RK, Bras J *et al.* Analysis of shared heritability in common disorders of the brain. *Science* 2018; **360**(6395).

24. Nalls MA, Saad M, Noyce AJ, Keller MF, Schrag A, Bestwick JP *et al.* Genetic comorbidities in Parkinson's disease. *Hum Mol Genet* 2014; **23**(3)**:** 831-841.

1 25. Birtwistle J, Baldwin D. Role of dopamine in schizophrenia and Parkinson's disease.
2   *Br J Nurs* 1998; **7**(14)**:** 832-834, 836, 838-841.
3
4 26. Menza MA, Robertson-Hoffman DE, Bonapace AS. Parkinson's disease and anxiety:
5   comorbidity with depression. *Biol Psychiatry* 1993; **34**(7)**:** 465-470.
6
7 27. Berardelli A, Rothwell JC, Thompson PD, Hallett M. Pathophysiology of
8   bradykinesia in Parkinson's disease. *Brain* 2001; **124**(Pt 11)**:** 2131-2146.
9
10
11

Figure Legends

Fig. 1. **The clinical phenotypes of two independent deeply phenotyped Parkinson's**

disease cohorts identify the same phenotypic axes. Results were consistent in two

independents cohorts (842 *Discovery* and 1807 *Tracking* patients). Examination of

these two separate Parkinson's disease cohorts, using independent derivation of the

phenotypic axes in each, showed significant correlations between each cohort's first

three axes. Correlations between the axes from each cohort are Axis 1 r=0.92 (p=3 x

10-13), Axis 2 r=0.89 (p=4 x 10-11), Axis 3 r=0.72 (p=5 x 10-6). The correlation

coefficient (x-axis) between each axis derived in each cohort (blue: *Discovery* vs red:

Tracking) and each clinical observation (y-axis) is shown.

Fig. 2. **The reduced dimensions in other dimensionality reduction methods fail to align**

between differently but deeply phenotyped UK and US Parkinson's disease

cohorts. We compared the ability of different dimensionality reduction methods

(independent component analysis (ICA), Multidimensional scaling (MDS), Principal

component analysis (PCA) and phenotypic axis based on the PHENIX multiple

phenotype mixed model) to phenotypically align two deeply phenotyped Parkinson's

disease cohorts, specifically the *Discovery* (842 individuals) and PPMI (439 sporadic

Parkinson's disease) cohorts. The x-axis and y-axis represent the correlation

coefficient between each continuous variable with clinical observation associated with

a specific symptom category in *Discovery* and PPMI cohort respectively. Each

column panel and colour of points ("Axis") represents the dimension level of each

underlying dimension. All points on the diagonal would represent a perfect

phenotypic alignment of both cohorts. We examined the relationship between

correlation derived from both cohorts by performing a linear regression: R^2 and p

correspond to the coefficient of determination and the p-value respectively. Only the

dimensions discovered by the MPMM model, PHENIX, show a significant

relationship between both cohorts: MPMM phenotypic axes (R^2=0.86, p=2x10-8),

MDS (R^2=0.11, p=0.18), ICA (R^2=0.17, p=0.16) and PCA (R^2=0.31, p=0.06).

Fig. 3.   **The correlation of individual clinically-measured Parkinson's disease**

phenotypes with an underlying Phenotypic Axis 1. Modelling patient clinical

phenotypes as a combination of genetic and environmental factors revealed three

phenotypic severity axes (**Fig.1**), each representing a continuous pattern of variation

between multiple co-varying clinical phenotypes. In Axis 1 (shown), (A) clinical

measures relating to anxiety and depression and apathy are significantly and

positively correlated with an individual's score along this axis; patients with a higher

axis score have more severe mood and neuropsychiatric problems. (B) The severity of

motor phenotypes is positively correlated with this phenotypic axis; patients with a

higher axis score is associated with more severe motor phenotypes (C) Cognitive tests

were negatively correlated with this component (the patients that score high in these

cognitive tests have less cognitive impairments); individuals with a high score for this

component suffer from more severe anxiety, depression and displayed more cognitive

impairment and motor symptoms.

Fig. 4. Alzheimer's disease phenotypic axis significantly outperforms the original

phenotypic axis. These heatmap plots represents for two first phenotypic axes

(left=V1 and right=V2), in the 3 cohorts (row1=Discovery, row2=Tracking & row 3=

) the excess(red) or deficit of the phenotypic variance explained by the different

disease axes (columns) compared with the original phenotypic axes.

1   **Table1:** Correlation between each axis and each clinical phenotypic measure

	Clinical Observation			r		
				Axis1	Axis2	Axis3
Behavior	BDI total	Measure of the depression	+	0.60	0.60	0.01
	Leeds Anxiety Total	Measure of the anxiety	+	0.51	0.55	0.05
	Leeds Depression	Measure of the depression	+	0.51	0.62	0.00
	QUIP all	Impulsive-Compulsive Disorders	+	0.12	0.24	0.03
	UPDRS apathy	Apathy	+	0.40	0.39	0.18
	UPDRS fatigue	Fatigue	+	0.49	0.49	0.08
	UPDRS hallucinations	Hallucinations	+	0.17	0.17	-0.02
Autonomic	Constipation	Quantitative measure of constipation	+	-0.15	-0.07	0.01
	Orthostatic	Blood pression from sitting/lying to stand up	+	0.17	-0.09	-0.06
	UPDRS constipation	Constipation	+	0.38	0.33	-0.08
	UPDRS pain	Pain	+	0.47	0.47	-0.01
Cognitive	Education years	Number of years of education	-	-0.21	-0.23	0.16
	MMSE total	Measure of cognitive ability	-	-0.27	-0.07	0.17
	MOCA total	Measure of cognitive ability	-	-0.31	-0.06	0.23
	Phonemic fluency	Number of words beginning with a particular letter	-	-0.26	0.03	0.14
	Sementic fluency	Number of animals and the number of boy names	-	-0.28	0.09	0.15
	BMI	Body Mass index	+	0.16	0.09	-0.08
	CGIC	Clinical global impression of change	+	0.05	-0.07	0.08
	Disease Duration	Disease Duration	+	0.24	0.19	-0.07
Motors	Flamingo time	Time that a person can stand on one leg	-	-0.46	-0.03	0.16
	Getgo average	Time taken for an individual to get up from a chair, walk three meters, turn around, walk back to the chair and sit down.	+	0.52	0.04	-0.16
	Purdue assembly	Test to measure manual dexterity	-	-0.37	0.16	0.10
	Purdue total	Test to measure manual dexterity	-	-0.41	0.18	0.09
	UPDRS arms	Arms	+	0.63	-0.50	0.78
	UPDRS bradykinesia	Bradykinesia	+	0.63	-0.40	0.57
	UPDRS faceneck	Face/neck problems	+	0.26	-0.22	0.12
	UPDRS I	Non Motor Aspects of Experiences of Daily Living	+	0.68	0.67	0.02
	UPDRS II	Motor Aspects of Experiences of Daily Living	+	0.76	0.30	0.05
	UPDRS III	Motors Examination	+	0.71	-0.46	0.61
	UPDRS IV	Motors complications	+	0.16	0.16	0.05
	UPDRS laterality	Unilateral	+	-0.03	-0.01	0.15
	UPDRS legs	Legs	+	0.59	-0.31	0.44
	UPDRS postural	Postural	+	0.64	-0.02	-0.09
	UPDRS rigidity	Rigidity	+	0.51	-0.27	0.35

	UPDRS speech	Speech	+	0.22	-0.07	-0.04
	UPDRS tremor	Tremor	+	0.20	-0.40	0.58
Sleep	ESS total	Measure of daytime sleepiness	+	0.31	0.22	-0.07
	RBD total	Measure of REM Sleep behavior disorder	+	0.29	0.29	-0.03
e						
	Sniff total	Smell identifications	-	0.01	0.06	0.12
Drug	LEDD total	Quantitative measure of the amount of Parkinson's disease medication	+	0.31	0.27	-0.22

12
13
14 (1) A high score for a clinical measure indicates **more (+)** or **less (-)** issue for the patient.
15 (2) The correlation coefficient under and above |0.25| are indicated in gray or blue/red respectively
16 (3) Red and blue cells indicates when a high phenotypic axis score are associated with more and less
17   clinical issues for the patient respectively.
18
19

1   **Fig.1**
2

3
4

1   **Fig. 2**
2

1   **Fig.3**
2

1   **Fig. 4**

3
4
5