Amyloid-β induced membrane damage instigates tunneling nanotubes by exploiting

p21-activated kinase dependent actin remodulation

Aysha Dilna[1+], Deepak K.V[1+], Nandini Damodaran[1+], Claudia S. Kielkopf [2,3], Katarina

Kagedal [2], Karin Ollinger [2] and Sangeeta Nath[1]*

[1] Manipal Institute of Regenerative Medicine, Manipal Academy of Higher Education,

Bangalore, 560065, India.

[2] Experimental Pathology, Department of Biomedical and Clinical Sciences Linköping

University, 581 85 Linköping, Sweden.

[3] Current address: Novo Nordisk Foundation Center for Protein Research, University

of Copenhagen, Copenhagen, Denmark.

[+] Equally contributed

* **Corresponding to be addressed to** Sangeeta Nath, email: <u>sangeeta.nath@manipal.edu</u>

Running Title: Cell-to-cell transfer of oAβ in TNT

Abstract: Alzheimer's disease (AD) pathology progresses gradually via anatomically connected brain regions. Earlier studies have shown that amyloid-β_{1-42} oligomers (oAβ) can be directly transferred between connected neurons. However, the mechanism of transfer is not fully revealed. We observed formation of oAβ induced tunneling nanotubes (TNTs), nanoscaled f-actin containing membrane conduit, in differentially differentiated SH-SY5Y neuronal models. Time-lapse images showed that TNTs propagate oligomers from one cell to another. Preceding the TNT-formation, we detected oAβ induced plasma membrane (PM) damage and calcium-dependent repair through lysosomal-exocytosis and significant membrane surface expansion, followed by massive endocytosis to re-establish the PM. Massive endocytosis was monitored by an influx of the membrane-impermeable dye TMA-DPH and PM damage was quantified by propidium iodide influx in the absence of calcium. The massive endocytosis eventually caused accumulation of internalized oAβ in Lamp1 positive multi vesicular bodies/lysosomes via the actin cytoskeleton remodulating p21-activated kinase1 (PAK1) dependent endocytic pathway. Three dimensional quantitative and qualitative confocal imaging, structured illumination superresolution microscopy (SIM) and flowcytometry data revealed that oAβ induces activated phospho-PAK1, which modulates the formation of long stretched f-actin extensions between cells. Moreover, formation of TNTs can be inhibited by preventing PAK1 dependent internalization of oAβ using small-molecule inhibitor IPA-3, a highly selective cell permeable auto-regulatory inhibitor of PAK1. The present study gives insight that the TNTs are probably instigated as a consequence of oAβ induced PM damage and repair process, followed by PAK1 dependent endocytosis and actin remodeling, probably to maintain cell surface expansion and/or membrane tension in equilibrium.

Introduction.

Neurodegenerative diseases are propagating disorders characterized by accumulation of misfolded proteins that form aggregates, plaque and eventually cause neurodegeneration. A common hallmark of neurodegenerative diseases is prion-like self-propagation and gradual pathology progression in a predetermined pattern to different parts of the brain [1]. Several studies have shown that proteins involved in these diseases such as tau, Aβ, α-synuclein and huntingtin follow common patterns including misfolding, self-propagation and neuron-to-neuron transfer [2,3]. In a model of Alzheimer's disease (AD), we have previously shown that spreading of AD pathology is due to direct transfer of amyloidogenic oligomers between connected neurons [4]. Moreover, lysosomal stress due to gradual accumulation of toxic non-degradable oAβ enhances the cell-to-cell progression of pathology [5]. The studies [4,5] have provided a possible explanations of how intracellular soluble oAβ, reported as the potential initiator or driver of AD [6,7], could develop gradual pathology by propagating between connected cells. However, the mechanism of direct neuron-to-neuron propagation of neurodegenerative aggregates is not yet revealed.

Recently, several studies have demonstrated TNTs to transfer neurodegenerative proteins, such as PrP^{Sc}, α-synuclein, Aβ, tau, polyQ, from one cell-to-another [8–11]. Several of these studies implicated links between TNTs and the endo-lysosomal pathway in cell-to-cell spreading [12]. In addition, exosomes are investigated as means of cell-to-cell transfer of Aβ [13,14]. However, these studies could not explain the anatomically connected strict spatiotemporal pathology progression of AD. On the other hand, cell-to-cell transfers of both extracellular and intracellular monomers and protofibrils of Aβ$_{1-42}$ via tunneling nanotubes (TNTs) are demonstrated in primary cultures of neurons and astrocytes [8]. However, the molecular basis of TNTs formation remains underexplored.

TNTs are open-ended membrane nanostructures consisting of membrane actin protrusions

between neighbouring cells. Correlative cryo-electron microscopy has recently demonstrated

that TNTs are formed by 2-11 individual TNTs and their diameters vary between 145 to 700

nm [15]. TNTs are transient structures that can stay intact from minutes to hours. Membrane

protrusions like filopodium precede TNT formation and inhibition of actin polymerization

attenuates their formation [16]. TNT formation prevails in neuronal cells and primary neurons [17].

Successive studies also showed TNT formation in different cell types, such as immune cells,

fibroblast, epithelial cells, astrocytes, and neurons, as well as their implication in the spreading

of pathology in neurodegenerative diseases, HIV, herpes simplex virus (HSV) infections and

in cancer [18–20]. A growing number of studies have revealed that vesicle transfer, recycling

vesicles, lysosomes and molecules involved in membrane expansion play a role behind the

formation of TNTs, the actin membrane protrusions [12,21].

In this study, we show oAβ induced formation of TNTs and direct cell-to-cell propagation of

oAβ between neighbouring cells via TNTs. Preceding the formation of TNTs, we detect oAβ

induced PM damage and repair through lysosomal exocytosis, which is followed by massive

endocytosis via the membrane cytoskeleton actin remodulating kinase PAK1 engaged

endocytic pathway. Endocytosis of oAβ activates phospho-PAK1, which modulates long

stretched f-actin and formation of TNTs. Moreover, formation of TNTs can be prevented by

inhibiting PAK1 dependent endocytosis and actin remodulation. Altogether, these observations

give new insights that sprouting of TNTs might be instigated as a consequence of oAβ induced

PM damage and Ca^{2+} dependent PM repair through lysosomal exocytosis via PAK1 dependent

actin remodelling.

Material and Methods.

Preparation of soluble oAβ. Freshly made unlabelled oAβ and fluorescently labelled oAβ-TMR were prepared from lyophilized Aβ (Aβ$_{1-42}$) and Aβ-TMR (Aβ$_{1-42}$-5-tetramethyl rhodamin) suspended in 1,1,1,3,3,3-hexafluoro-2-propanol (AnaSpec). Lyophilized Aβ and Aβ-TMR were resuspended at a concentration of 5 mM in Me2SO and then diluted to a concentration of 100 μM in HEPES buffer, pH 7.4. The solution was immediately vortexed and sonicated for 2 min and then incubated at 4°C for 24 hours [4,5]. Oligomers were characterized before the experiments similarly as reported in our earlier papers [4,5], by electron microscopy imaging using a Jeol 1230 transmission electron microscope equipped with an ORIUS SC 1000 CCD camera, together with SDS-PAGE, Native-PAGE western blots and size exclusion chromatography.

Neuronal cells culture and differentiations. SH-SY5Y neuronal cells (ECACC; Sigma-Aldrich) were seeded on 10-mm glass-bottom Petri dishes (MatTek) at a concentration of 12,000 cells/cm^2. Cells were partially differentiated with 10 μM retinoic acid (RA; Sigma-Aldrich) for 7 days. Pre-differentiated or partially differentiated SH-SY5Y cells were further differentiated for additional 10 days in 10-mm glass-bottom Petri dishes (MatTek) with brain-derived neurotrophic factor, neuregulin-1, nerve growth factor, and vitamin D3. After 10 days of differentiation on glass, the cells form long, branched neurites and several neurospecific markers, as described previously [4,22].

Cell culture and transfections.

For actin, PM, Lamp1 and GFP-GPI, plasmid constructs were used. The plasmid mEGFP-lifeact-7 (Addgene # 54610) was a gift from Michael Davidson and Lamp1-mGFP (Addgene # 34831) was a gift from Esteban Dell'Angelica [23]. CAAX-mCherry (Ampicillin resistant)

original source is [24] and GFP-GPI construct previously used by others [25,26]. The competent

DH5α strain *E.coli* cells were used to transform the bacterial cells to be used for isolation of

the plasmids. Plasmid DNA isolation was carried out using the QIAGEN Plasmid Midi kit.

Transfections of SH-SY5Y cells were done using jetPRIME transfection reagent (Polyplus)

and also by using Lipofectamine 3000 (Invitrogen). The cells (30,000 per well) were seeded

on glass coverslips placed in a 24 well plate in 0.5 mL culture media. The plasmid DNA (0.5

µg) was diluted in 50 µL of jetPRIME buffer and vortexed for 10 seconds, followed by mixing

with 1.6 µL of jetPRIME reagent which was then added to the cells cultured at 70-90%

confluency and incubated between 24-48 hours before taking images. The transfection using

Lipofectamine 3000 (Invitrogen) were done using (0.5 µg) plasmid-DNA complex added to

the cells with fresh Opti-MEM (reduced serum) media which are at 70-90% confluency. The

mixture was left for two to three hours and then removed. Fresh DMEM media with serum was

added and microscopic images were taken after 24 - 48 hours.

oAβ internalization/uptake. To investigate the uptake mechanisms of oAβ in the used cell

system, differentiated SH-SY5Y cells were pre-treated with inhibitors in growth medium for

the indicated time and at the indicated concentration (Table 1). After washing with PBS, a final

concentration of 0.5 µM oligomerised Aβ-TMR was added to the cells for 1.5 h. After removal

of the reagents, the cells were kept in a growth medium for 30 min before flow cytometry

analysis. For flow cytometry, the cells were washed twice with PBS and trypsinised. Cells of

2 wells were resuspended in 300 µl PBS and filtered through a 50 µm nylon-mesh filter

(Partec). The cells were analysed on a FACS Aria III flow cytometer (BD Biosciences) using

FACSDiva acquisition software. Each treatment was carried out and analysed at least in

triplicates. The gating was set using control cells without fluorophore and cells treated with

Aβ-TMR only. The percentage of cells containing Aβ-TMR was calculated and normalised to

the mean of Aβ-uptake without any inhibitor, resulting in a fold change, to make results from

different experiments comparable. Kaluza Software (Beckman Coulter) was used for data

analysis.

Colocalization of internalized oAβ. To study if oAβ was internalized within LAMP1 positive

vesicles, different concentrations of oAβ-TMR of 250 nM, 500 nM and 1μM were added into

the cells transfected with Lamp1-mGFP for time periods of 15 min, 30 min, 1 hour and 2 hours.

oAβ (500nM and 1μM) internalization into the GPI positive vesicles were studied by

incubating the oligomers for 15 min, 30 min and 1 hour into the cells transfected with GFP-

GPI. Then images were taken after bleaching extracellular GFP-GPI by adding 0.4% Trypan

blue (Sigma Aldrich) for 15 min and fixing with 4% PFA (Paraformaldehyde, Sigma Aldrich)

for 15 min at room temperature and mounting with DABCO. oAβ (500nM and 1μM) and

dextran FITC (1mg/ml) were co-incubated with the SH-SY5Y cells for 15 min, 30 min and 1

h min at 37°C. Then images were captured for both live cells and cells fixed with 4% PFA.

Cells were visualised using fluorescence microscopy and % of colocalization from Mander's

overlapping coefficient were analyzed using ImageJ plugin Coloc 2 (open source by NIH,

USA).

Ca^{2+} dependent plasma membrane repair assay by propidium iodide staining.

Undifferentiated SH-SY5Y cells were incubated with 1 μM of oAβ for 1 h or 2 h at 37 °C in

the presence and absence of 5 mM EGTA (without Ca^{2+}) in DPBS buffer (PBS with Ca^{2+} and

Mg^{2+} chloride). Then the cells were washed and stained with propidium iodide (PI, 5 μg/ml)

for 5 min. Cells were washed again two times with PBS before applying 4% PFA as fixative

for 20 min at 4 °C. The fixed cells were observed by fluorescence microscopy or trypsinized

before flow cytometry analysis.

Immunocytochemistry. Lysosomal exocytosis was verified by immunocytochemical staining

of Lamp1 on the outer leaflet of the PM in unfixed cells, using an antibody directed to the

luminal part of Lamp1 in undifferentiated SH-SY5Y neuroblastoma cells as described earlier

[27], on unfixed cells. Cells were incubated for 15 and 30 min with 1 μM of oAβ in MEM media

without FCS (fetal calf serum) supplement. Then, endocytosis of the cells was blocked with

5% BSA + 10% FCS in PBS for 5 min at 4 °C. Then cells were incubated with Lamp1anti-goat

primary antibody (1:250, sc-8099, Santa Cruz Bio-technology; Santa Cruz, CA, USA; 2 h,

4 °C) in the blocking buffer for 45 min, followed by fixation of the cells in 4% PFA for 20 min

at 4 °C before labelling with the secondary anti-goat antibody conjugated to Alexa Fluor 488

(1:400 for 30 min; Molecular Probes, Eugene, OR, USA). Next, the cells were mounted in

ProLong Gold antifade reagent supplemented with 4′,6-diamidino-2-phenylindole (DAPI;

Molecular Probes). Conventional immunocytochemical staining was done to quantify the

activated PAK1 and actin in oAβ and IPA-3 treated cells using phospho-PAK1 (Thr423)/PAK2

(Thr402) antibody (Cell signalling #260; 1:150) and Phalloidin–Tetramethylrhodamine B

isothiocyanate (Sigma P1951, 1:500). Anti-rabbit Alexa 405 and FITC (1:500) were used as

secondary antibodies.

Live cell imaging of membrane dynamics. oAβ (1 μM) was added concurrently with 1 μM

of the membrane binding dye TMA-DPH (N,N,N-Trimethyl-4-(6-phenyl-1,3,5-hexatrien-1-yl)

phenylammonium p-toluenesulfonate; molecular formula: $C_{28}H_{31}NO_3S$) (10 mM of TMA-

DPH stock solution was made by dissolving in methanol) and membrane dynamics of the live

cells were followed by time-laps images using a confocal microscope. The excitation and

emission maximum of TMA-DPH is 384 and 430 nm, with a considerable tail of excitation

spectra at 405 nm. Therefore, time-lapse images were taken using a confocal-microscope by

exciting the TMA-DPH dye using the 405 nm laser and sufficient emission light was collected

using an opened pinhole. In this setup confocal microscope produces images similar to the

widefield or epifluorescence images. Cultures were carried to the microscope one by one in

500 μl of 20 mM HEPES buffer of pH 7.4 maintaining the temperature at 37°C. oAβ (1 μM)

was added concurrently with 1 μM of membrane binding dye TMA-DPH to HEK cells

similarly as above and membrane dynamics of the live cells were studied by images taken by

fluorescent microscope.

Cell viability assay. Viability of differentiated cells were measured in triplicates using MTT

reagent with the undifferentiated cells treated with oAβ (1μM) for 1 h with or without IPA-3

(20 μM) pretreated cells for 30 min at 37°C. Cells were incubated with MTT reagent for 2

hours at 37°C, then removed all the media and measured the DMSO solubilized formazan.

Formazan (bright orange in color), the reduced MTT product produced by viable cells after 2

h of incubation at 37°C was dissolved in DMSO and measured, at 570 nm using a Victor

Wallac (PerkinElmer) plate reader.

Confocal and fluorescence microscopy to image TNTs. Formation of oAβ induced TNTs

was observed in the cells treated with increasing concentrations of oAβ at different incubation

times, using either confocal or fluorescence microscopes. The treatments were done by

incubating the cells with oAβ in serum free medium and corresponding control cells were

treated at the same conditions, to nullify the serum starvation induced TNTs. Differentiated,

partially differentiated and undifferentiated cells on glass petri dishes were incubated with 100–

500 nM oAβ–TMR for 3 h at 37°C in a 5% CO_2 atmosphere. The cells were imaged after

incubating with LysoTracker (green; Invitrogen) 50-250 nM for 5-10 min and after extensive

PBS washing (two washes of 10 min each at 37°C with 5% CO_2). Images were acquired using

a Zeiss LSM laser scanning confocal microscope using 63X/1.4 NA or 40X/1.3 NA oil immersion plan-apochromatic objective (Carl Zeiss AG, Oberkochen, Germany). The time-laps image sequences of the live cells were taken at 37°C by capturing simultaneously differential interference contrast (DIC) and fluorescence modes. Fluorescence microscopy (IX73-Olympus) with 63X/1.3 NA and 100X1.4 NA was also used to do the experiments of SH-SY5Y cells transfected with Plenti-lifeact EGFP plasmid and CAAX-mCherry plasmid. To study the effect of IPA-3 on SH-SY5Y cells, the cells were first treated with IPA-3 (20 μM) for 30 minutes followed by oAβ (1 μM) for 1 - 3 hours respectively. The cells were then imaged under microscopes and numbers of TNTs were quantified and plotted in percentage by manual counting the TNTs with respect to the number of cells from each of the image frames.

Flow-cytometry. Internalization of oAβ-TMR in the presence and absence of inhibitors were quantified using BD FACS Aria ™ (BD Biosciences) and were analysed using BD FACS DIVA ™ (BD Bioscienc es) flow cytometer. Immunocytochemical stained cells with anti PAK1 and actin-phalloidin were fixed, trypsinised and suspended in PBS and quantified using BD LSR II (BD Biosciences) flow cytometer. Cells treated with propidium iodide (PI) were fixed, trypsinized and filtered using CellTrics 30 μm filters (Sysmex). Then re-suspended in PBS and quantified different sets either using BD FACS Aria ™ (BD Biosciences) or BD LSR II (BD Biosciences) flow cytometer and data were analysed using BD FACS DIVA ™ (BD Bioscienc es) flow cytometer.

Superresolution SIM images. All SIM (structured illumination microscopy) images were acquired using on DeltaVision OMX SR microscope from GE Healthcare using a 60X 1.42 NA objective and pco.edge sCMOS detector. The cells were fixed by 4% PFA by incubation

for 15 min and fixed cells were imaged. The widefield images were deconvolved using the

built-in algorithm.

Image analysis and statistics. Image analysis was done using ImageJ software (open source

by NIH, USA). Percentage of co-localization of oAβ (magenta) with lysosomes (green) was

performed by calculating the proportion of the magenta fluorescence pixels compared to the

co-localized pixels from the background subtracted images using the Coloc-2 plugin. The

number of TNTs were distinguished from neurites by comprising a 3D-volume view of cells

from confocal stacks using the volume view plugin in Image J software. Cells with blebs /

lamellipodia and TNTs were counted from images and normalized to the total number of cells

and represented in percentage. oAβ induced endocytosis was quantified by measuring the

integrated intensities of internalized TMA-DPH from the luminal part of each cell by drawing

ROI (region of interest) over sequences of time-laps confocal image stacks and comparing it

with the same quantification of control cells. The fusion of lysosomal membrane to reseal the

damaged membrane was detected as the appearance of LAMP1 (green) on the outer leaflet of

the PM. The outer leaflet LAMP1 was quantified measuring the integrated intensities from

drawn ROI and the proportionate percentage calculated comparing the total LAMP1 per cells.

One-way ANOVA tests were performed to validate statistical significance in all experiments.

Results

oAβ induced cellular stress instigates formation of TNTs and cell-to-cell pathology

propagation. Cell-to-cell propagation of oAβ between connected cells has been shown in

earlier studies [4,5,8,14]. In our previous work [4], we have shown asymmetric spreading of oAβ

along the dispersion of neurons path, when microinjected in a single neuron of a primary

hippocampal culture [4]. Additionally, we showed transfer of oAβ-TMR from connected donor

to acceptor cells using a 3D donor-acceptor co-culture model with differentiated SH-SY5Y cells. However, the clear mechanism behind the cell-to-cell propagation remains to be revealed. Differentiated cells both in 2D and 3D culture form neurites and express several neuronal markers and characteristics of mature neurons, where 3D differentiated cells express tau subtypes better comparable to mature human neurons than 2D differentiated [4,22]. Here, we have observed that differentiated cells in 2D culture upon treatment with oAβ-TMR, morphologically exhibit lamellipodia, cell membrane expansion or blebs (yellow arrows) as well as tunneling nanotube-like (TNT-like) long thin conduits (white arrows) between neighbouring cells (Fig. 1B, Supplementary Movie 1). We did not observe significant numbers of TNT-like structures and membrane expansions/lamellipodia in the differentiated control cells (Fig. 1A). We have quantified the number of cells with blebs/lamellipodia and counted the number of TNT-like structures from bright field images with respect to the total number of cells from each of the image frames and presented as a percentage. The results show a concentration-dependent increase in oAβ-TMR treated cells (200-500 nM) compared to the control cells (Fig. 1C, D). We have detected the direct transfer of oAβ-TMR between the cells prominently (indicated by blue arrows) via these TNT-like conduits (Fig. 1E-F, Supplementary Movie 3-4) by time-lapse imaging. Transfer of organelles and oAβ-TMR (blue arrows) via TNT-like conduits extended from lamellipodia between cells (Yellow arrows), were observed in the cells incubated for 3 h with 100, 250 and 500 nM of oAβ-TMR (Supplementary Movies 2-4).

In order to show that the studied TNT-like structures are indeed oAβ induced TNTs and not an artefact of differentiating reagents, we performed the experiments in partially differentiated (treated with retinoic acid for 7 days) and undifferentiated SH-SY5Y cells in parallel and confirmed that TNTs from 3D volumetric images obtained from confocal z-stacks. We have

observed thin TNT-like structures extended from expanded lamellipodia-like membrane protrusions in partially differentiated cells internalized with oAβ-TMR after incubation with 250 nM of oligomers for 3 h (Fig. 2A-B), in contrast to the control cells (Fig. 2C). Moreover, the partially differentiated SH-SY5Y cells formed networks of TNT-like conduits between neighbouring cells (yellow arrow, Fig. 2B). The cells make networks between 3 neighbouring cells via TNT-like conduits (yellow arrows), and those were also extended from expanded lamellipodia (black arrows, Fig. 2A). We have observed the transfer of oAβ-TMR colocalized with Lysotracker labelled lysosomes (yellow arrows) from one cell to another via the TNT-like conduits extended from expanded lamellipodia (black arrows; Fig. 2D). To confirm that the cell-to-cell conduits visible in bright field images are TNTs, we have composed 3D volume view from confocal z-stacks of phalloidin stained f-actin structures between neighbouring cells (yellow arrows, Fig. 2E-F). In the 3D volume view, TNTs were distinguished from neurites by the characteristics of their capacity to stay hanging without touching the substratum even after fixing the cells (Fig. 2F). The lengths of the TNTs were between 0.2-16 μm. Diameters of TNTs were measured from confocal z-stacks images of phalloidin stained cells and the values of majority (> 90%) of TNTs are within 240-960 nm measured at xy-plan (between 1-4 pixels, pixel size is 240 nm). However, around 10 % of TNTs are thicker and their diameters are in the range of 1.23 ± 0.27 μm. Finally, we have quantified the number of TNTs and neurites per cell in partially differentiated cells treated with 1 μM of oAβ-TMR over time (1-3 h), and compared to the control cells (Fig. 2G-H). The results show increasing numbers of TNTs over the time (1-3 h) of oAβ (1 μM) treatment (Fig. 2G). At the same time, we observed a significant decrease in the number of neurites per cells after 3 h of oAβ (1 μM) treatment (Fig. 2H). Similarly, oAβ showed toxicity to the neurites of differentiated cells, which caused significantly reduce number of neurites per cell detected in a concentration and time dependent manner (Supplementary Fig. 1A-D).

oAβ induced TNT formation precedes enhanced membrane activities and massive endocytosis. We have designed the experiments using undifferentiated SH-SY5Y cells to ensure that it is the toxicity of oAβ that causes induction of TNT-like structures, and not the differentiating reagents. Cells were co-transfected with lifeact-EGFP and CAAX-mCherry, to stain actin and peripheral membrane proteins. We found formation of numerous co-stained TNT-like structures (yellow arrows) after 1h of oAβ (1 μM) treatment, compared to the control cells (Fig. 3A-B). Further, the TNT-like structures were confirmed as a continuous extension of the PM and actin protrusions between two cells from 3D volume view from confocal z-stacks (Fig. 5B). Preceding the formation of oAβ induced TNTs, we have observed enhanced membrane activities along with the formation of membrane ruffles, filopodium, blebs and massive endocytosis compared to the control cells (Fig. 3C; Supplementary Movies 5-6). Enhanced membrane activities and endocytosis were detected immediately at the addition of oAβ (1 μM). The oAβ-induced endocytosis was quantified by measuring the internalization of TMA-DPH into the luminal part of the cells, compared to the control cells (Fig. 3D). TMA-DPH is membrane impermeable and able to enter the control cells and to the cells treated with oAβ (1 μM, for 15 min) only via endocytosis. On the other hand, the membrane impermeable dye TMA-DPH is able to enter the cells treated with oAβ (1 μM) for 1 h, and the dye stains the whole cells (oAβ treated for 1 h) immediately at the addition (Fig. 3E). The result suggests that oAβ induced changes in the membrane fluidity.

To understand if the oAβ induced TNT formation is specific to neuronal cells or if it is a basic cellular process, we incubated HEK-293 cells with oAβ (1 μM) for 1 h and observed formation of long TNTs hanging between distant neighbouring cells (Supplementary Fig. 2C). The number of TNT like conduits were significantly lower in control cells (Supplementary Fig. 2A-B). Similar to SH-SY5Y cells, the membrane of control HEK-293 cells found to be labelled

by TMA-DPH only at the periphery (Supplementary Fig. 2D). However, in the oAβ (1 μM for

1 h) treated HEK-293 cells, TMA-DPH labelled whole cells by entering the cells immediately

on addition. Additionally, these cells also showed TNT-like protrusions (Supplementary Fig.

2E). Results suggest oAβ induced change in membrane fluidity and increased number of TNT-

like structures.

oAβ$_{1-42}$ induces membrane damage and lysosomal exocytosis. oAβ induces TNT formation

in association with the substantial enhancement of membrane activities and massive

endocytosis, similarly as evident in Ca^{2+} dependent repair of injured PM by lysosomal

exocytosis [28]. We have observed that the oAβ-TMR (magenta) was efficiently internalized into

early endosomes (Rab5 positive organelles), followed by entry into multivesicular bodies

(MVB) or lysosomes (Lamp1 positive organelles) (Fig. 3F). The spontaneous internalizations

of extracellularly applied oAβ-TMR (250 nM to 1 μM, incubated for 15 min to 1 h) into

undifferentiated cells were quantified and results showed that 70 ± 9 % of oAβ-TMR (250 nM)

ended up in Lamp1 positive organelles after 15 min of incubation (Supplementary Fig. 3A-B).

Images in Fig. 3F, present the colocalization of oAβ-TMR (1 μM) after 30 min of incubation.

This inspired us to determine if oAβ caused PM damage. The Gold standard to detect PM repair

by lysosomal exocytosis and formation of a patch over the damaged membrane is to detect the

luminal part of the lysosomal membrane protein LAMP-1 exposed on the outer leaflet of the

plasma membrane. Accordingly, we have observed transiently transfected Lamp1-mGFP

distributed in large extent in the periphery of the PM within 15 to 30 min of oAβ (1 μM)

treatment, compared to the control cells (Fig. 3G upper panel). We have quantified oAβ (1 μM)

induced Lamp1 on the PM, in undifferentiated SH-SY5Y cells within 15-30 min of exposure

by surface staining of the cells (Fig. 3G lower panel & Fig. 3H). To verify that the process is

calcium dependent, we analysed the influx of the membrane-impermeant dye propidium iodide

(PI) in presence of 5 mM EGTA in PBS. The rationale behind this experiment is that if

lysosomal exocytosis-dependent PM repair is occurring, then chelation of Ca^{2+} will prevent the

repair and PI will be detected intracellularly. As seen in Fig. 3I-M, significant enhancement of

PI staining in absence of calcium was detected 30 min, 1h and 2 h after exposure to oAβ as

quantified by flow cytometry (Fig. 3K-M) and confocal imaging (Fig.3O). In the presence of

Ca^{2+} increased PI staining was not detected with oAβ treatments (Fig. 3I-J). The quantification

of internalized PI was compared by plotting median fluorescence intensity, as fluorescence

histogram profiles are bimodal in nature (Fig.3N). The presented dot plot of PI internalization

after 30 min of Ca^{2+} chelation in the presence and absence of oAβ (1 μM) (Fig. 3M). The results

show an increasing amount of internalization of PI with time even in the control cells, however

the differences between control and oAβ (1 μM) treated cells are significant at both 30 min and

1 h time point, and even at 2 h time point differences are detectable. The results suggest that

Ca^{2+} dependent membrane repair is a continuous process even in the control cells, and oAβ

induced membrane damage accelerates the process. Thus, we conclude that the addition of oAβ

causes damage to the PM and as a result of rapid membrane repair process occurs, by enhancing

the process of lysosomal exocytosis and fusion of lysosomal membrane with the PM.

Subsequent to the membrane repair process, re-establishment of the PM occurs by removing

membrane parts through endocytosis and as a consequence, oAβ is internalized into

MVB/lysosomes (Lamp1 positive organelles) (Fig. 3F and Supplementary Fig. 3A-B).

Involvement of the actin cytoskeleton remodeling kinase PAK1 engaged endocytosis in

the internalization of oAβ$_{1-42}$. We next wanted to determine the exact mechanism of the

massive internalization of oAβ. To identify the mechanisms of internalization of oAβ, partially

differentiated SH-SY5Y cells were pre-treated with different inhibitors against the uptake of

Aβ as suggested in the literature. We have used partially differentiated cells, because N-methyl-

D-aspartate (NMDA) receptors, a-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA) receptors and nicotinic acetylcholine receptors (nAChR) are known to be present on RA treated partially differentiated SH-SY5Y cells [29] and have previously been shown to mediate the internalization of Aβ [30–32]. To inhibit the function of these receptors, antagonists against NMDA receptors (AP-5), AMPA receptors (GYKI 52466) and nAChR (α-Bungaratoxin) were used and oAβ-TMR uptake was quantified by flow cytometry. However, no change in the uptake of oAβ was found (Fig. 4A).

To determine the mechanism of Aβ-uptake several inhibitors that affect different variants of endocytosis and macropinocytosis were tested. After pre-treatment with the inhibitors, the cells were treated with oAβ-TMR and quantified by flow cytometry (Fig. 4B-C). The percentage of oAβ-positive cells was normalized to the mean percentage of oAβ-uptake and the fold-change upon inhibitor treatment was calculated. MDA and PAO, which are inhibitors of clathrin-mediated endocytosis, [33] and DPA, which functionally inhibits acid sphingomyelinases [34] could not prevent the uptake. Similarly, the dynamin inhibitor dynasore [35], and an amiloride analogue, EIPA that acts as a specific inhibitor of macropinocytosis through inhibition of Na^+/H^+ exchangers [36] had no effect. However, we found significantly reduced oAβ-uptake when using bafilomycin A1 (Baf) and NH_4Cl, which both affect the lysosomal acidification and consequently fusing between vesicles [37] and the PAK1 inhibitor IPA-3 that inhibits actin regulated endocytosis similar as macropinocytosis by modulating PAK1 dependent actin polymerization [38,39]. PAK1 has an autoregulatory domain which is targeted by the inhibitor IPA-3, although the exact downstream inhibitory signalling pathway of IPA-3 is not known yet [38]. To establish if the oAβ internalization in undifferentiated SH-SY5Y cells occurs similarly via PAK1 dependent endocytosis, we also quantified the internalization in undifferentiated cells and the quantification showed similar results, Fig. 4D shows the representative confocal images.

Involvement of PAK1 has been reported mostly in membrane tension and cholesterol sensitive

clathrin-independent endocytosis (CIE) and in membrane modulating actin dependent

endocytosis such as, macropinocytosis, IL2Rβ endocytosis, CLIC/GEEC (clathrin-independent

carrier/ glycosylphosphatidylinositol-anchored protein enriched compartment) pathway [39,40]. It has

been shown that PAK1 specifically regulates macropinocytosis by following the uptake of 70k

Da dextran, a specific marker of fluid phase macropinocytosis [41]. Therefore, we have followed

the fate of internalization of Dextran-70 kDa with oAβ pulses. However, after 1 h of oAβ-TMR

(1 μM) treatment only small amounts of internalized dextran-FITC were detectable and there

was no significant colocalization with internalized oAβ-TMR (Fig. 4E). Similarly,

internalizations of oAβ were observed in the GFP-GPI (GFP- glycosylphosphatidylinositol)

transiently transfected SH-SY5Y cells, and the internalized oligomers were not colocalized

with GFP-GPI in the cells treated for 1 h with oAβ-TMR (1 μM) (Fig. 4E) and oligomers did

not enhance the GFP-GPI internalization observed after 15 - 30 min of oAβ (0.5 nM to 1 μM)

treatment. The results indicate that oAβ follows PAK1 dependent CIE machineries, which is

distinct from macropinocytosis or the CLIC/GEEC pathway. The same observations were also

reported recently [39] although the complete mechanism is not fully understood.

Active PAK1 in the oAβ induced formation of TNTs. ActivePAK1 regulates cortical actin

polymerization, directional movements and also polarized lamellipodia at the leading edge [42].

Actin depolymerizing drugs such as latrunculin and cytochalasin impair TNTs formation [17].

Therefore, the next step was to observe the role of PAK1 in the oAβ induced formation of

TNTs and if inhibition of oAβ uptake by IPA-3 also prevented the formation of TNTs. Thus,

oAβ induced formation of TNTs in undifferentiated SH-SY5Y cells was quantified by creating

3D volume view from z-stacks images (Fig. 5B) of phalloidin and anti PAK1/2-Thr423 stained

conduits between neighbouring cells at fixed condition (Fig. 5A). Number of TNTs was

quantified in cells treated with 1 μM of oAβ for 1-3 h, compared to the control cells (Fig. 5C)

and results show a significant time dependent increase in the number of TNTs after oAβ

treatment. Paraformaldehyde fixation might break or damage the thin TNTs. Thus, TNTs were

also quantified by staining f-actin in live cells by transiently transfecting with the lifeact-EGFP

plasmid (Fig. 5E). Transfection did not affect the ability of cell to form TNT-like structures

and the increased number of TNT-like structures in oAβ treated cells as compared to the

controls maintained. Cells that were treated with IPA-3 showed a decrease in number of TNTs

(Fig. 5D-E), images were quantified from live cells as well as from fixed condition (Fig. 5D).

The quantification of TNTs in the live cells was higher but the pattern is similar to the fixed

cells, which were quantified from phalloidin stained TNTs images by analysing in 3D volume

view from the confocal Z-stacks (Fig. 5E). Morphologically, IPA-3 treated cells were rounder

than the controls and oAβ treated cells, however MTT assay showed no significant change in

the cell viability (Supplementary Fig. 4A).

oAβ induced activation of phospho-PAK1 modulates f-actin and formation of TNTs.

Distinct differences in f-actin structures between oAβ treated and the oAβ + IPA-3 treated cells

were detected by observing the lifeact EGFP plasmid labelled f-actin structures at a resolution

of < 130 nm using Superresolution SIM (Structured Illumination Microscopy) images. TNTs

were observed to be a continuous extension of long cortical f-actin in oAβ treated cells (Fig.

5F and Fa). 3D volume view of the xyz-plane demonstrated that membrane actin protrusion

did not grow on the surface like neurites. Rather, they were 'hanging' between two

neighbouring cells, which is one of the characteristics distinguishing features of TNTs (Fig.

5Fb). In the Fig.5G, the disruptions of long actin fibres in IPA-3 pre-treated cells reveal PAK1

play an important role in f-actin modulation and thereby TNT formation. Experiments were

done with undifferentiated sparsely seeded SH-SY5Y cells to quantify actin stained TNTs like

structures between relatively distant cells (Supplementary Fig. 4B).

Images of immunocytochemical staining using phospho-PAK1 (Thr423)/PAK2 (Thr402)

antibody demonstrated increased levels of activated phospho-PAK1 with time (1-3 h) of oAβ

(1μM) treatment (Fig. 6A-B). The confocal images were represented as z-projected stacks and

the intensities were compared from z-projected images, taken at the same settings of laser

power and exposure. Co-staining of activated phospho-PAK1 with f-actin on TNT structures

were confirmed from the 3D-volume view analysis (Fig. 5A-B & 6A). Upregulation of

activated phospho-PAK1 by oAβ (1μM) treatment was quantified further from epi-

fluorescence images, where images were taken from more than 80 cells from each set using

20X magnification objective and at the same exposure and settings (Fig. 6C). The results show

a higher level of activated phospho-PAK1 in oAβ (1μM) after 1 h treatment of cells in

comparison to the control and IPA-3 pretreated cells (Fig. 6D). Further quantifications were

done by flow cytometry using immunocytochemical stained cells. The results obtained from

the histograms of flow cytometeric data showed higher expression of phospho-PAK1 and

phalloidin bound actin in oAβ treated cells as compared to control, IPA-3 and IPA-3 + oAβ

treated cells (Fig. 6F-G). Quantification of TNT-like cell-to-cell phalloidin stained actin

extensions with respect to the level of activated phospho-PAK1 showed a positive correlation

when quantified observing a larger number of cells (Fig. 6E).

Discussion.

Prion-like cell-to-cell propagation is a common characteristic of neurodegenerative diseases.

Several reports have consistently reported direct cell-to-cell propagation of neurodegenerative

protein aggregates and their implications in the gradual pathology progression [2,3]. Several

studies have suggested exosomes as a means of cell-to-cell transfer of Aβ [13,14]. On the other

hand, studies using the method of transwell assays have also shown the efficient cell-to-cell

transfer of prion proteins despite the blocking of exosome transfer [9,43]. Moreover, an increasing

number of reports show that cell-to-cell transfer of neurodegenerative proteins, such as PrP[Sc],

α-synuclein, tau, polyQ aggregates and Aβ, via TNTs instigate new avenues [8–11]. oAβ induced

formation of TNTs in primary neurons and astrocytes has already been reported [8]. Here we

have focused more on possible mechanism of oAβ induced formation of TNTs in differentially

differentiated SH-SY5Y neuronal cells. Due to its cancerous origin, the SH-SY5Y cell line

shows a number of genetic aberrations and different differentiation protocols generate

variations in neuronal properties, but most genes and pathways dysregulated in AD

pathogenesis stay intact even in undifferentiated SHSY5Y cells. Moreover, its differentiation

to cholinergic neurons like properties in retinoic acid treated SHSY5Y cells a widely accepted

model system in AD research [22].

Increasingly, clinical studies and animal models indicate that soluble oAβ is the disease

initiator and driver, rather than large extracellular depositions. Accumulation of amyloidogenic

proteins in lysosomes, abnormal lysosomal morphology and lysosomal membrane

permeabilization are major hallmark of neurodegenerative diseases. Notably, lysosomal stress

as well as damage due to accumulation of non-degradable amyloidogenic aggregates could

induce the formation of TNTs [12]. Stress signals from lysosomes dysregulate various cellular

processes and mediate increased oxidative stress [12,44–46]. Several studies have indicated that

ROS (reactive oxygen species) induced cellular stress enhances TNT formation [8,17].

In contrast, the study [47], demonstrated that the impaired processing of Aβ due to lysosomal

enzymatic inefficiency can enhance exocytosis. Additionally, a related protein of the exocyst

complex M-sec, involved in exosome fusion and membrane expansion, regulates formation of

TNTs [21]. PM recruitment of Ral-GTPase and filamin, both actin remodeling proteins, also

indicate positive regulating effects in TNT formation [21]. The study presented here also

demonstrates membrane expansion in the form of blebs, filopodium-like structures and

extension of TNTs from expanded lamellipodia. Previous reports have indicated that synthetic

oAβ makes ion-permeable pores in synthetic membranes [48,49]. Recently, it was also shown that

oAβ can induce a membrane repair response similar to that induced by exposure to the bacterial

pore-forming toxin produced by *B. thuringensis* [50]. Consequently, enhanced internalization of

Aβ occurs via endocytosis, which is independent of receptor interactions. Involvement of the

clathrin- and dynamin-independent endocytosis is also relevant in maintaining cellular

homeostasis by regulating membrane stress and cell surface expansion [39,51]. A recent study

reported that Aβ follows membrane tension sensitive and Rho GTPase family regulated actin-

dependent CIE [39]. Furthermore, Aβ uptake was earlier shown to be inhibited by nocodazole

and cytochalasin-D, the inhibitors of tubulin depolymerization and actin polymerization, by

preventing fluid phase-endocytosis in microglia and astrocytoma cells [52]. The results of this

study show that the internalization of oAβ through massive endocytosis is clathrin-

independent, but rather follows PAK1-dependent membrane actin modulating endocytosis

machineries.

PAK1 is a serine/threonine kinase found in the cytoplasm and nucleus of cells and PAK1 is

important in regulating cytoskeleton remodelling, phenotypic signalling, gene expressions and

it affects a variety of cellular processes [42]. In addition, PAK1 acts downstream of the small

GTPases Cdc42 and Rac1, which interact with many effector proteins, including Arp2/3, which

in turn can have an effect on cytoskeleton reorganization [53]. In addition, the role of CDC42 and

Rho-GTPases in TNT formation is not fully investigated. A report [54], demonstrated that the

activity of CDC42 and Rho-GTPases positively contributes to the formation of TNTs in

macrophages. In contrast, another study [55], found that CDC42/IRSp53/VASP negatively

regulates the formation of TNTs. PAK activation can occur independent of Rac and CDC42,

the specific lipids particularly sphingolipids can directly activate PAK [56]. Aβ internalization in

primary neurons in absence of apolipoprotein-E has been reported by a cholesterol and

sphingolipid sensitive lipids rafts mediated clathrin and caveolae independent but dynamin

dependent pathway [57,58]. The mechanism is similar to actin-dependent IL2Rβ endocytosis [40].

AD is highly associated with changes in lipid composition of neurons, and Aβ42 directly

downregulates sphingomyelin levels and the ratio of Aβ40/Aβ42 regulates membrane

cholesterol [59]. Here we have detected substantial changes in membrane fluidity upon oAβ

treatment using the membrane dye TMA-DPH together with rapid endocytosis. On the other

hand, oAβ is implicated in PAK1 dependent synaptic dysfunctions and PAK1 is aberrantly

activated and translocated from the cytosol to the membrane during the development of

pathology in the AD brain [60]. Interestingly, HIV and HSV viruses exploit PAK1 dependent

endocytosis *en route* and recent studies have also shown direct cell-to-cell spreading of HIV

and HSV via TNTs [19]. Another study has shown that the HIV-1 Nef protein mediated TNT

formation is associated with 5 proteins of the exocyst complex and that they are involved in a

PAK2 and Rab 11 dependent pathway [61]. Additionally, alpha herpes virus induced TNT-like

membrane actin projections depends on the conserved viral US3 serine/threonine protein

kinase-dependent modulations of the cytoskeleton. These modulations are caused by activation

of PAK1-dependent signalling and inhibition by IPA-3 attenuates these TNT-like projections

[62,63]. Exocytosis is involved in the expansion of cell surface area and results in decreased

membrane stress. Reduced membrane stress also arises during PM repair, due to extensive

exocytosis events [64]. Endosome recycling also plays a big role in maintaining membrane

surface area in equilibrium [65]. Interestingly, the role of vesicle recycling in TNT formation has

been evaluated in a recent study carried out on CAD cells, where an increased level of Rab

11a, Rab 8a and VAMP3 were reported in correlation to TNTs [66].

Aβ induced formation of TNTs in primary neurons and astrocytes has been reported earlier [8].

Here, we have shown the oAβ induced formation of TNTs can transfer aggregates directly

between differentiated neurons and the transferred oligomers develop gradual pathology. TNTs

as a mean of direct neuron-to-neuron transfer is a convincing model of how oligomers that are

suggested as an initiator or driver of AD pathology could gradually progress through the

anatomically connected brain regions. However, the possibility of pathology transfer via

exosomes could be a parallel mechanism, since we have observed that the formation of TNTs

is followed by oAβ induced enhanced lysosomal exocytosis. Therefore, further in-depth studies

are needed to understand how cells maintain homeostasis of intercellular communication by

balancing exosome release and TNTs in the stressed cells. Altogether, our results have

indicated that TNTs are characteristic membrane actin conduit that transfer oAβ aggregates

from one cell-to-another, and are formed as a consequence of oAβ induced membrane damage

and Ca^{2+} dependent lysosomal-exocytosis engaged rapid membrane repair process, followed

by PAK1-kinase dependent CIE and actin remodelling (Fig. 7), probably to maintain cell

surface area expansion and membrane stress in equilibrium.

Author Contributions. S.N conceived and conducted the research; S.N and K.O designed

research; S.N designed tunneling nanotubes, oAβ internalization and PAK1 experiments; K.O.

designed membrane dynamics and membrane repair experiments; A.D, D.K.V, N.D, C.K and

S.N performed experiments and analysed the data; S.N, K.O and K.K interpreted data; S.N

wrote the paper taking valuable inputs from all the authors.

Authors have no Competing Interests.

Acknowledgement. SN thanks to Manipal Academy of Higher Education intramural grants (India; 2019-2022), Magnus Bergvalls (Sweden, 2015-2016, #2014-00192), Gun och Bertil Stohnes research grants (Sweden, 2014-2015) and Alzheimer fonden (Sweden; 2012-2013). We thank our long-term collaborator Prof. Martin Hallbeck of Linkoping University, Sweden, for the valuable suggestions on designing of cell-to-cell transfer experiments. We thank Dr. Ravi Manjithaya of JNCASR, Bangalore, India, for giving us SH-SY5Y cell lines and giving access to the SIM-superresolution microscope. Thanks a lot to Likhesh Sharma, PhD, Product Manager GE Healthcare for assisting with imaging by SIM-superresolution microscope. We thank Prof. Satyajit Mayor of National centre for biological sciences, India, for his kind gift of GFP-GPI construct, CAAX-mCherry plasmid, sharing of some reagents and for his valuable inputs. The CAAX-mCherry plasmid was a kind gift to Prof. Satyajit Mayor by Prof. Jacco van Rheenen of Hubrecht Institute for Developmental Biology and Stem Cell Research. Thanks to Prof. Subba Rao Gangi Setty of Indian Institute of Science and Prof. Gopal Pande of Manipal institute of higher education, India, for their valuable inputs.

References.

1. Hardy, J. & Selkoe, D. J. The amyloid hypothesis of Alzheimer's disease: Progress and problems on the road to therapeutics. *Science* (2002). doi:10.1126/science.1072994

2. Hallbeck, M., Nath, S. & Marcusson, J. Neuron-to-neuron transmission of neurodegenerative pathology. *Neuroscientist* **19**, 560–566 (2013).

3. Brundin, P., Melki, R. & Kopito, R. Prion-like transmission of protein aggregates in neurodegenerative diseases. *Nature Reviews Molecular Cell Biology* (2010). doi:10.1038/nrm2873

4. Nath, S. *et al.* Spreading of neurodegenerative pathology via neuron-to-neuron transmission of β-amyloid. *J. Neurosci.* **32**, (2012).

5. Domert, J. *et al.* Spreading of amyloid-β peptides via neuritic cell-to-cell transfer is dependent on insufficient cellular clearance. *Neurobiol. Dis.* **65**, 82–92 (2014).

6. Gouras, G. K., Tampellini, D., Takahashi, R. H. & Capetillo-Zarate, E. Intraneuronal β-amyloid accumulation and synapse pathology in Alzheimer's disease. *Acta Neuropathologica* **119**, 523–541 (2010).

7. Walsh, D. M. & Selkoe, D. J. Deciphering the molecular basis of memory failure in

Alzheimer's disease. *Neuron* **44**, 181–93 (2004).

8. Wang, Y., Cui, J., Sun, X. & Zhang, Y. Tunneling-nanotube development in astrocytes depends on p53 activation. *Cell Death Differ.* **18**, 732–742 (2011).

9. Gousset, K. *et al.* Prions hijack tunnelling nanotubes for intercellular spread. *Nat. Cell Biol.* **11**, 328–336 (2009).

10. Tardivel, M. *et al.* Tunneling nanotube (TNT)-mediated neuron-to neuron transfer of pathological Tau protein assemblies. *Acta Neuropathol. Commun.* **4**, 117 (2016).

11. Dieriks, B. V. *et al.* α-synuclein transfer through tunneling nanotubes occurs in SH-SY5Y cells and primary brain pericytes from Parkinson's disease patients. *Sci. Rep.* **7**, (2017).

12. Victoria, G. S. & Zurzolo, C. The spread of prion-like proteins by lysosomes and tunneling nanotubes: Implications for neurodegenerative diseases. *Journal of Cell Biology* **216**, 2633–2644 (2017).

13. Rajendran, L. *et al.* Alzheimer's disease beta-amyloid peptides are released in association with exosomes. *Proc. Natl. Acad. Sci.* **103**, 11172–11177 (2006).

14. Sardar Sinha, M. *et al.* Alzheimer's disease pathology propagation by exosomes containing toxic amyloid-beta oligomers. *Acta Neuropathol.* **136**, 41–56 (2018).

15. Sartori-Rupp, A. *et al.* Correlative cryo-electron microscopy reveals the structure of TNTs in neuronal cells. *Nat. Commun.* (2019). doi:10.1038/s41467-018-08178-7

16. Gerdes, H. H., Rustom, A. & Wang, X. Tunneling nanotubes, an emerging intercellular communication route in development. *Mech. Dev.* **130**, 381–387 (2013).

17. Rustom, A., Saffrich, R., Markovic, I., Walther, P. & Gerdes, H. H. Nanotubular Highways for Intercellular Organelle Transport. *Science (80-.).* **303**, 1007–1010 (2004).

18. Onfelt, B., Nedvetzki, S., Yanagi, K. & Davis, D. M. Cutting edge: Membrane nanotubes connect immune cells. *J. Immunol.* **173**, 1511–3 (2004).

19. Jansens, R. J. J., Tishchenko, A. & Favoreel, H. W. Bridging the Gap: Virus Long-Distance Spread via Tunneling Nanotubes. *J. Virol.* (2020). doi:10.1128/jvi.02120-19

20. Davis, D. M. & Sowinski, S. Membrane nanotubes: Dynamic long-distance connections between animal cells. *Nature Reviews Molecular Cell Biology* **9**, 431–436 (2008).

21. Hase, K. *et al.* M-Sec promotes membrane nanotube formation by interacting with Ral and the exocyst complex. *Nat. Cell Biol.* **11**, 1427–1432 (2009).

22. Agholme, L., Lindström, T., Kgedal, K., Marcusson, J. & Hallbeck, M. An in vitro model for neuroscience: Differentiation of SH-SY5Y cells into cells with morphological and biochemical characteristics of mature neurons. *J. Alzheimer's Dis.* **20**, 1069–1082 (2010).

23. Falcón-Pérez, J. M., Nazarian, R., Sabatti, C. & Dell'Angelica, E. C. Distribution and dynamics of Lamp1-containing endocytic organelles in fibroblasts deficient in BLOC-3. *J. Cell Sci.* (2005). doi:10.1242/jcs.02633

24. Van Rheenen, J. *et al.* EGF-induced PIP2 hydrolysis releases and activates cofilin
locally in carcinoma cells. *J. Cell Biol.* (2007). doi:10.1083/jcb.200706206

25. Nichols, B. J. *et al.* Rapid cycling of lipid raft markers between the cell surface and
golgi complex. *J. Cell Biol.* (2001). doi:10.1083/jcb.153.3.529

26. Paladino, S., Pocard, T., Catino, M. A. & Zurzolo, C. GPI-anchored proteins are
directly targeted to the apical surface in fully polarized MDCK cells. *J. Cell Biol.*
(2006). doi:10.1083/jcb.200507116

27. Wäster, P., Eriksson, I., Vainikka, L., Rosdahl, I. & Öllinger, K. Extracellular vesicles
are transferred from melanocytes to keratinocytes after UVA irradiation. *Sci. Rep.* **6**,
(2016).

28. Idone, V. *et al.* Repair of injured plasma membrane by rapid Ca2+ dependent
endocytosis. *J. Cell Biol.* **180**, 905–914 (2008).

29. Xie, H. R., Hu, L. Sen & Li, G. Y. SH-SY5Y human neuroblastoma cell line: In vitro
cell model of dopaminergic neurons in Parkinson's disease. *Chinese Medical Journal*
(2010). doi:10.3760/cma.j.issn.0366-6999.2010.08.021

30. Bi, X., Gall, C. M., Zhou, J. & Lynch, G. Uptake and pathogenic effects of amyloid
beta peptide 1-42 are enhanced by integrin antagonists and blocked by NMDA
receptor antagonists. *Neuroscience* (2002). doi:10.1016/S0306-4522(02)00132-X

31. Zhao, W. Q. *et al.* Inhibition of calcineurin-mediated endocytosis and α-amino-3-
hydroxy- 5-methyl-4-isoxazolepropionic acid (AMPA) receptors prevents amyloid β
oligomer-induced synaptic disruption. *J. Biol. Chem.* (2010).
doi:10.1074/jbc.M109.057182

32. Yang, W. N. *et al.* Mitogen-activated protein kinase signaling pathways are involved
in regulating α7 nicotinic acetylcholine receptor-mediated amyloid-β uptake in SH-
SY5Y cells. *Neuroscience* (2014). doi:10.1016/j.neuroscience.2014.08.013

33. Guo, S. *et al.* Selectivity of commonly used inhibitors of clathrin-mediated and
caveolae-dependent endocytosis of G protein-coupled receptors. *Biochim. Biophys.*
Acta - Biomembr. (2015). doi:10.1016/j.bbamem.2015.05.024

34. Kornhuber, J. *et al.* Identification of novel functional inhibitors of acid
sphingomyelinase. *PLoS One* (2011). doi:10.1371/journal.pone.0023852

35. Macia, E. *et al.* Dynasore, a Cell-Permeable Inhibitor of Dynamin. *Dev. Cell* (2006).
doi:10.1016/j.devcel.2006.04.002

36. Mercer, J. & Helenius, A. Virus entry by macropinocytosis. *Nature Cell Biology*
(2009). doi:10.1038/ncb0509-510

37. Poëa-Guyon, S. *et al.* The V-ATPase membrane domain is a sensor of granular ph that
controls the exocytotic machinery. *J. Cell Biol.* (2013). doi:10.1083/jcb.201303104

38. Deacon, S. W. *et al.* An Isoform-Selective, Small-Molecule Inhibitor Targets the
Autoregulatory Mechanism of p21-Activated Kinase. *Chem. Biol.* (2008).
doi:10.1016/j.chembiol.2008.03.005

39. Wesén, E., Lundmark, R. & Esbjörner, E. K. Role of Membrane Tension Sensitive
Endocytosis and Rho GTPases in the Uptake of the Alzheimer's Disease Peptide

Aβ(1-42). *ACS Chem. Neurosci.* (2020). doi:10.1021/acschemneuro.0c00053

40.   Doherty, G. J. & McMahon, H. T. Mechanisms of Endocytosis. *Annu. Rev. Biochem.*
(2009). doi:10.1146/annurev.biochem.78.081307.110540

41.   Dharmawardhane, S. *et al.* Regulation of macropinocytosis by p21-activated kinase-1.
Mol. Biol. Cell (2000). doi:10.1091/mbc.11.10.3341

42.   Sells, M. A., Boyd, J. T. & Chernoff, J. p21-Activated kinase 1 (Pak1) regulates cell
motility in mammalian fibroblasts. *J. Cell Biol.* (1999). doi:10.1083/jcb.145.4.837

43.   Thayanithy, V. *et al.* A transwell assay that excludes exosomes for assessment of
tunneling nanotube-mediated intercellular communication. *Cell Commun. Signal.* **15**,
1–16 (2017).

44.   Freeman, D. *et al.* Alpha-Synuclein Induces Lysosomal Rupture and Cathepsin
Dependent Reactive Oxygen Species Following Endocytosis. *PLoS One* **8**, (2013).

45.   Eriksson, I., Nath, S., Bornefall, P., Giraldo, A. M. V. & Öllinger, K. Impact of high
cholesterol in a Parkinson's disease model: Prevention of lysosomal leakage versus
stimulation of α-synuclein aggregation. *Eur. J. Cell Biol.* **96**, 99–109 (2017).

46.   Gowrishankar, S. *et al.* Massive accumulation of luminal protease-deficient axonal
lysosomes at Alzheimer's disease amyloid plaques. *Proc. Natl. Acad. Sci.* **112**, E3699–
E3708 (2015).

47.   Annunziata, I. *et al.* Lysosomal NEU1 deficiency affects amyloid precursor protein
levels and amyloid-β secretion via deregulated lysosomal exocytosis. *Nat. Commun.* **4**,
(2013).

48.   Arispe, N., Pollard, H. B. & Rojas, E. Giant multilevel cation channels formed by
Alzheimer disease amyloid beta-protein [A beta P-(1-40)] in bilayer membranes. *Proc.*
Natl. Acad. Sci. U. S. A. **90**, 10573–7 (1993).

49.   Kagan, B. L. Membrane pores in the pathogenesis of neurodegenerative disease. *Prog.*
Mol. Biol. Transl. Sci. **107**, 295–325 (2012).

50.   Julien, C. *et al.* In vivo induction of membrane damage by β-amyloid peptide
oligomers. *Acta Neuropathol. Commun.* **6**, 131 (2018).

51.   Thottacherry, J. J. *et al.* Mechanochemical feedback control of dynamin independent
endocytosis modulates membrane tension in adherent cells. *Nat. Commun.* **9**, (2018).

52.   Mandrekar, S. *et al.* Microglia mediate the clearance of soluble aβ through fluid phase
macropinocytosis. *J. Neurosci.* (2009). doi:10.1523/JNEUROSCI.5572-08.2009

53.   Daniels, R. H., Hall, P. S. & Bokoch, G. M. Membrane targeting of p21-activated
kinase 1 (PAK1) induces neurite outgrowth from PC12 cells. *EMBO J.* (1998).
doi:10.1093/emboj/17.3.754

54.   Hanna, S. J. *et al.* The Role of Rho-GTPases and actin polymerization during
Macrophage Tunneling Nanotube Biogenesis. *Sci. Rep.* (2017). doi:10.1038/s41598-
017-08950-7

55.   Delage, E. *et al.* Differential identity of Filopodia and Tunneling Nanotubes revealed
by the opposite functions of actin regulatory complexes. *Sci. Rep.* (2016).
doi:10.1038/srep39632

56. Bokoch, G. M. *et al.* A GTPase-independent Mechanism of p21-activated Kinase
Activation. *J. Biol. Chem.* (1998). doi:10.1074/jbc.273.14.8137

57. Saavedra, L., Mohamed, A., Ma, V., Kar, S. & De Chaves, E. P. Internalization of β-
amyloid peptide by primary neurons in the absence of apolipoprotein E. *J. Biol. Chem.*
(2007). doi:10.1074/jbc.M701823200

58. Marshall, K. E., Vadukul, D. M., Staras, K. & Serpell, L. C. Misfolded amyloid-β-42
impairs the endosomal–lysosomal pathway. *Cell. Mol. Life Sci.* **77**, 5031–5043 (2020).

59. Grimm, M. O. W. *et al.* Regulation of cholesterol and sphingomyelin metabolism by
amyloid-β and presenilin. *Nat. Cell Biol.* (2005). doi:10.1038/ncb1313

60. Ma, Q. L. *et al.* p21-activated kinase-aberrant activation and translocation in
Alzheimer disease pathogenesis. *J. Biol. Chem.* (2008). doi:10.1074/jbc.M708034200

61. Mukerji, J., Olivieri, K. C., Misra, V., Agopian, K. A. & Gabuzda, D. Proteomic
analysis of HIV-1 Nef cellular binding partners reveals a role for exocyst complex
proteins in mediating enhancement of intercellular nanotube formation. *Retrovirology*
(2012). doi:10.1186/1742-4690-9-33

62. Van Den Broeke, C. *et al.* Alphaherpesvirus US3-mediated reorganization of the actin
cytoskeleton is mediated by group A p21-activated kinases. *Proc. Natl. Acad. Sci. U.
S. A.* (2009). doi:10.1073/pnas.0900436106

63. Jacob, T., Van den Broeke, C., Van Waesberghe, C., Van Troys, L. & Favoreel, H. W.
Pseudorabies virus US3 triggers RhoA phosphorylation to reorganize the actin
cytoskeleton. *J. Gen. Virol.* (2015). doi:10.1099/vir.0.000152

64. Andrews, N. W. & Corrotte, M. Plasma membrane repair. *Current Biology* (2018).
doi:10.1016/j.cub.2017.12.034

65. Grant, B. D. & Donaldson, J. G. Pathways and mechanisms of endocytic recycling.
Nature Reviews Molecular Cell Biology (2009). doi:10.1038/nrm2755

66. Bhat, S. *et al.* Rab11a–Rab8a cascade regulates the formation of tunneling nanotubes
through vesicle recycling. *J. Cell Sci.* (2018). doi:10.1242/jcs.215889

 Figure Legends

Figure legend 1: oAβ induced formation of TNT-like conduits in differentiated SH-SY5Y

neuronal cells. A-B) TNT-like conduits (indicated by white arrows) between neighbouring

cells were detected in differentiated SH-SY5Y cells incubated with 500, 250 and 200 nM of

oAβ-TMR (magenta) for 3 h, washed and stained with 50 nM of lysotracker (green) before the

capture of the image. The cells with long TNT-like conduits form noticeable blebs and cell

membrane expansion (indicated by yellow arrows), in contrast to control cells. A)

Differentiated control cells showed neurite like structures rather than TNT-like conduits. Note

that the control cells are also devoid of blebs and cell membrane expansion. C) Percentages of

cells with blebs / lamellipodia were quantified from the images taken with increasing

concentrations of oAβ-TMR (200-500 nM) and compared with the control cells. D) TNT-like

conduits were counted and plotted in percentage with respect to the number of cells from each

of the image frames. Quantifications were done from > 60 cells in each set. n > 3. Plots are

mean ± SD. One-way ANOVA tests were performed to validate statistical significance. E)

Differentiated SH-SY5Y cells were incubated with 500 nM of oAβ-TMR (magenta) for 3 h,

washed and labelled with 50 nM of lysotracker (green) and a sequence of images at different

time points were taken. The cells form TNT-like conduits and oAβ-TMR travels from one cell-

to-another as organelle like puncta structures through the conduits (blue arrows). The cells form

noticeable blebs (yellow arrows). F) The cells treated with 250 nM oAβ-TMR (magenta) for 3

h form lamellipodia (yellow arrows), show direct transfer of oAβ-TMR from one cell-to-

another as organelle like puncta structures (blue arrows) through the TNT-like conduits (white

arrows). Scale bars are 10 μm.

Figure legend 2: oAβ induced formation of TNTs in partially differentiated SH-SY5Y

neuronal cells. A-B) Partially differentiated SH-SY5Y cells were incubated with 250 nM of

oAβ-TMR (magenta) for 3 h, washed and labelled with 200 nM of lysotracker (green). A) oAβ-

TMR internalized cell makes network between 3 neighbouring cells via formation of TNT-like

connections (yellow arrows), extended from lamellipodia-like membrane protrusions (black

arrows). B) TNT-like conduits form networks between two neighbouring cells (yellow arrow).

C) Partially differentiated control cells devoid of blebs / lamellipodia. D) Partially

differentiated SH-SY5Y cells were incubated with 250 nM of oAβ-TMR (magenta) for 3 h,

washed and labelled with 200 nM of lysotracker (green). The cells form thin TNT-like conduits

(yellow arrows) extended from expanded lamellipodia-like membrane protrusions (black

arrows). oAβ-TMR co-localized with lysosomes that moves from one cell to another via

connected conduits (yellow arrows). E-F) 3D volume views were composed from confocal z-

stacks of phalloidin stained f-actin structures between neighbouring cells to validate whether

the cell-to-cell conduits which were visible in bright field images are TNTs or not. G-H) Then,

TNTs were distinguished from neurites from the characteristics of their capacity to stay

hanging without touching the substratum even after fixing the cells. Quantifications of G)

TNTs and H) neurites were done in the cells treated with 1 μM of oAβ for 1-3 h, compared to

the control cells, from > 60 cells in each set. Plots are mean ± SD. One-way ANOVA tests

were performed to validate statistical significance. Scale bars are 10 μm.

Figure Legend 3: oAβ induced PM damage and repair via coupled lysosomal-exocytosis

and endocytosis. A) TNT-like conduits (yellow arrows) were quantified in undifferentiated

SHSY5Y cells co-transfected with lifeact-EGFP (actin) and CAAX-mCherry (peripheral

membrane-protein). B) Quantification of TNT-like conduits from number of cells > 60 for each

set. C) Massive membrane activities and endocytosis were observed in the oAβ (1 μM) treated

cells together with the membrane dye TMA-DPH (0.5 μM), compared to control cells. D)

Endocytosis of PM labelled with TMA-DPH was quantified by measuring luminal part

intensities (plotted mean ± SD, quantified > 20 cells from each set, n=3). E) In oAβ (1h with 1

μM) treated cells, penetration of the membrane impermeable dye TMA-DPH on addition, and

the cells show TNT-like conduits (yellow arrows). F) Cells incubated with extracellularly

applied oAβ-TMR (1 μM) internalize (magenta) efficiently to early-endosomes (Rab 5; green)

and late-endosomes/lysosomes (Lamp1, green). G) Translocation of Lamp1-mGFP to cell

surface (Upper panels), n=3. H) Lamp1 surface staining (Lower panel) within 15-30 min of

exposure of oAβ (1 μM) was quantified from intensity (plotted mean ± SD) measurements from

defined ROI using ImageJ. (n=3, each dot represents the number of cells). I-J) oAβ (1 μM)

induced membrane damage detected as the uptake of the membrane-impermeable dye PI in

presence and absence of Ca^{2+}, were quantified by flowcytometry. Represented histograms I) In

presence and K-L) absence of Ca^{2+}. M) Representative dot-plot in absence of Ca^{2+}. N)

Penetration of PI presented as a change of median fluorescence in control and oAβ (1 μM, 30

min and 1 h) treated cells in presence and absence of Ca^{2+} (n = 6). O) Representative confocal

images of PI uptake after 2 h of oAβ treatment compared to control. Plotted mean ± SD. One-

way ANOVA were performed to validate statistical significance. Scale bars are 10 μm.

Figure Legend 4. Mechanisms of rapid internalization of oAβ-TMR were identified by

pre-treating the SH-SY5Y cells with different endocytosis and receptor inhibitors. The

percentages of oAβ-TMR containing cells in partially differentiated cells were analysed by

flow cytometry. A) Cells were pretreated with the receptor inhibitors α-Bungarotoxin, AP-5,

GYKI. n=1, triplicates. B) Cells were pretreated with endocytosis inhibitors Dynasore, DPA,

MDC, PAO, NH4Cl, Baf, IPA-3 and EIPA, followed by exposure to 0.5 μm oAβ-TMR for 1h.

n= 3, > minimum of duplicates in each set. The percentage values were normalized to the

respective mean of oAβ. C) Representative dot-plots with forward scatter vs fluorescence

intensities (TMR). D) Representative confocal images of undifferentiated SH-SY5Y cells

treated with endocytosis inhibitors and oAβ-TMR. E) Representative images are showing that

internalized oAβ-TMR (magenta; 1 μM incubated for 1 h) was not colocalized with 70 kDa

Dextran-FITC (green) and GFP-GPI (green). One-way ANOVA tests were performed to

validate statistical significance. Scale bar = 10 μm.

Figure Legend 5. Inhibition of rapid internalization of oAβ by the PAK 1 inhibitor IPA-

3 can also inhibit formation of TNTs. A) Represented oAβ induced TNTs from confocal z-

stack images in the undifferentiated SH-SY5Y cells co-stained by phalloidin (magenta) and

anti PAK1/2-Thr423 (green). B) TNTs were quantified creating 3D volume view from z-stacks

confocal images. C) Quantification of TNTs in the cells treated with 1 μM of oAβ for 1-3 h,

compared to the control cells. E) Fluorescence microscopy images of control cells, oAβ (1 μM)

treated, IPA-3 treated and cells treated with IPA-3 and then oAβ were captured using

undifferentiated SH-SY5Y cells transfected with lifeact EGFP (green) (white arrows). D)

Number of TNTs were counted with respect to the number of cells to quantify the inhibition

by IPA-3 and compared the TNT numbers both in live cells and fixed cells. Number of cells >

50 for each set. Plotted mean ± SD. One-way ANOVA tests were performed to validate

statistical significance. F) Superresolution images of lifeact stained undifferentiated SH-SY5Y

cells by DeltaVision™ OMX SR microscopy. SH-SY5Y cells were transfected with Plenti-

lifeact EGFP and treated with oAβ (1μM) and IPA-3 (20μM). F) In oAβ (1 h) treated cells,

TNTs appear to spread out as extensions from the long f-actin of the cytoskeleton and to

connect the two neighbouring cells. Fa) Zoomed image of a proper z-stack illustrated clearly

the f-actin labelled TNT. Fb) 3D volume view of z-stacks of SIM image shows that, in contrast

to neurons, TNTs do not touch the substratum, but rather stay "hanging" between the two

connected cells even after the fixation of the cells. G) In the IPA-3+oAβ   treated cells,

disruptions of long stretched f-actin of cytoskeleton and inhibition of TNTs are evident. 3D

reconstructions and analysis were done using ImageJ. Scale bars are 10 μm.

Figure Legend 6. oAβ induced activated phospho-PAK1 modulates f-actin extensions and

IPA-3 disrupts the long-stretched f-actin. A) Confocal images of immuno-stained cells

treated with oAβ (1 μM) over the time show localization of phospho-PAK1 (Thr423)/PAK2

(Thr402) antibody (green) on TNTs (indicated by white arrows). B) Intensity analysis of z-

projected confocal images show increase level of activated phospho-PAK1. C) Images of

immuno-stained cells using phospho-PAK1 (Thr423)/PAK2 (Thr402) antibody (green) and

actin-binding phalloidin-TMR (magenta) were taken by 20X 0.4 NA objective. D) Level of

active phospho-PAK1 were quantified measuring the intensities from each cell using image-j

ROI manager after background subtractions (number of cells >80 in each set) with oAβ, IPA-

3 and IPA-3+oAβ treated cells. E) Number of TNT like cell-to-cell connections (indicated by

white arrows) show a positive correlation with the intensities of activated phospho-PAK. F-G)

Level of active phospho-PAK1 and phalloidin labelled actin were quantified by flowcytometry

in the cells treated with oAβ, IPA-3 and IPA-3+oAβ in compared to the control. One-way

ANOVA tests were performed to validate statistical significance. Scale bars = 10 μm.

Figure Legend 7: Schematic summary to show that the sprouting of TNTs might be derived

as a result of oAβ induced PM damage and Ca^{2+} dependent lysosomal-exocytosis engaged rapid

membrane repair process, followed by PAK1-kinase dependent CIE and actin remodelling to

re-establish the cell surface area expansion or membrane stress in equilibrium.

Table 1

Pre-treatment conditions for inhibitors.

Inhibitor	Concentration	Treatment protocol
α-Bungaratoxin	100 nM	kept during experiment
AP-5 ((2R)-amino-5-phosphonopentanoate)	100 μM	kept during experiment
GYKI 52466 [4-(8-methyl-9H-1,3-dioxolo[4,5-h][2,3]benzodiazepin-5-yl)-benzenamine hydrochloride]	100 μM	kept during experiment
DPA (desipramine)	25 μM	2 h before experiment
Bafilomycin A1	10 nM	30 min before experiment
NH_4Cl	10 mM	30 min before experiment, no washing step, kept during experiment
MDC (monodansylcadaverine)	50 μM	30 min before experiment
PAO (phenylarsine oxide)	300 μM	30 min before experiment
Dynasore	80 μM	30 min before experiment

IPA-3 (1,1-Disulfanediyl-dinaphthalen-2-ol)	20 μM	30 min before experiment
EIPA (5-ethylisopropyl amiloride)	100 μM	30 min before experiment

916   Table 1

923 Figure 1

924 (*Dilna et al. 2021*)

926   Figure 2

927   (*Dilna et al. 2021*)

930   Figure 3

931   (*Dilna et al. 2021*)

933   Figure 4

934   (*Dilna et al. 2021*)

Figure 5

(Dilna et al. 2021)

Figure 6

(*Dilna et al. 2021*)

Figure 7

(*Dilna et al. 2021*)