High rate of sex chromosome turnover in muscid flies

Richard P Meisel[1]*, Pia U Olafson[2], Felix D Guerrero[2], Kranti Konganti[3], Joshua B Benoit[4]

1. Department of Biology and Biochemistry, University of Houston, Houston, TX

2. USDA-ARS Knipling-Bushland US Livestock Insects Research Laboratory, Veterinary Pest Genomics Center, Kerrville, TX

3. Texas A&M Institute for Genome Sciences and Society, College Station, Texas

4. Department of Biological Sciences, University of Cincinnati, Cincinnati, OH

*To whom correspondence should be addressed:
Richard P Meisel
Department of Biology and Biochemistry
University of Houston
3455 Cullen Blvd
Houston, TX 77204-5001
rpmeisel@uh.edu

Abstract

Sex chromosomes and the sex determining genes that reside on them can experience a high rate of evolutionary turnover in some taxa and evolutionary lineages. A substantial body of population genetics theory has been developed and tested to explain this rapid evolution. In contrast, we do not know why the rates of sex chromosome turnover differ across taxa. Addressing this question will require comparing closely related taxa with fast and slow rates of sex chromosome turnover to identify biological features that could explain these rate differences. Cytological karyotypes suggest that muscid flies (e.g., house fly) and blow flies are such a taxonomic pair. There appear to be multiple sex chromosome turnover events amongst muscids and very few in blow flies. However, we do not know the extent to which muscid sex chromosomes are independently derived along different evolutionary lineages. To address that question, we used genomic data to identify young sex chromosomes in two closely related muscid fly species, horn fly (*Haematobia irritans*) and stable fly (*Stomoxys calcitrans*). We provide evidence that the nascent sex chromosomes of horn fly and stable fly were derived independently from each other and from the young sex chromosomes of the closely related house fly (*Musca domestica*). We present three different scenarios that could have given rise to the sex chromosomes of horn fly and stable fly, and we describe how the scenarios could be distinguished. Distinguishing between these scenarios in future work could help to identify features of muscid genomes that promote sex chromosome turnover.

Introduction

In species where sex is determined by genetic differences between males and females, sex determining loci can reside on sex chromosomes, such as the male-limited Y chromosome in mammals that carries a male-determining gene (Swain and Lovell-Badge, 1999). Canonical Y chromosomes contain only a handful of genes with male-specific functions (Charlesworth *et al.*, 2005; Bachtrog, 2013). Sex determining genes and sex chromosomes often differ across species because of evolutionary turnover of sex chromosomes and sex determining pathways in common as species diverge (Bachtrog *et al.*, 2014; Beukeboom and Perrin, 2014). This sex chromosome turnover can convert autosomes into sex chromosomes and revert sex chromosomes back into autosomes (Carvalho and Clark, 2005; Larracuente *et al.*, 2010).

There are multiple ways in which sex chromosome turnover can occur, most of which can be divided into two broad categories. First, when a new sex determining locus arises on an autosome, it can convert the autosome into a "proto-sex-chromosome", and the ancestral sex chromosome can revert to an autosome (van Doorn, 2014). Second, autosomes can fuse with X, Y, Z, or W chromosomes to create "neo-sex-chromosomes" (Pennell *et al.*, 2015; Vicoso and Bachtrog, 2015). Population genetics theory suggests that sex-specific selection pressures and sexual antagonism are important contributors to sex chromosome turnover (Charlesworth and Charlesworth, 1980; Rice, 1986; van Doorn and Kirkpatrick, 2007, 2010). This theory has been tested in many plants and animals, and those tests have generally supported the hypothesis that sex-specific selection is important for sex chromosome turnover (e.g., Roberts *et al.*, 2009; Wang *et al.*, 2012; Qiu *et al.*, 2013; Zhou and Bachtrog, 2012).

Some taxa have high rates of sex chromosome turnover, whereas the same X & Y (or Z & W) chromosomes are shared across all species in other taxa (Beukeboom and Perrin, 2014; Blackmon and Demuth, 2014). Despite the well constructed theory explaining how sex chromosome turnovers can occur and the empirical work supporting that theory, we know very little about why rates of sex chromosome turnover differ across taxa (Bachtrog *et al.*, 2014; Abbott *et al.*, 2017). Contrasting closely related taxa with different rates of sex chromosome turnover could allow for the identification of biological factors that promote or inhibit sex chromosome turnovers.

Brachyceran flies (i.e., higher Diptera) are well-suited models for identifying factors that

promote or inhibit sex chromosome turnover because there are variable rates of sex chromosome turnover along different lineages (Vicoso and Bachtrog, 2015). The karyotype of the most recent common ancestor (MRCA) of Brachycera consists of five gene-rich autosomes (Muller elements A–E), a gene-poor heterochromatic X chromosome (Muller element F), and a Y chromosome that carries a male-determining locus (Boyes and Van Brink, 1965; Foster *et al.*, 1981; Weller and Foster, 1993; Vicoso and Bachtrog, 2013). Elements A–E correspond to the five gene-rich chromosome arms of *Drosophila melanogaster* (X, 2L, 2R, 3L, and 3R), and element F is homologous to the gene-poor *D. melanogaster* chromosome 4 (Muller, 1940). The ancestral brachyceran X chromosome has been conserved for ∼175 million years in some phylogenetic lineages, while new sex chromosomes have arisen along other lineages (Baker and Wilkinson, 2010; Wiegmann *et al.*, 2011; Vicoso and Bachtrog, 2013, 2015).

Within Brachycera, the sister families of Muscidae (house flies and their allies), Calliphoridae (blow flies), and Sarcophagidae (flesh flies) could be especially informative for comparative studies because they appear to have family-specific rates of sex chromosome turnover (**Fig 1**). Nearly all blow flies and flesh flies have the ancestral fly karyotype, with five autosomes, a heterochromatic X, and a male-determining locus on a Y chromosome (Boyes, 1961; Boyes and Van Brink, 1965; Scott *et al.*, 2014b; Vicoso and Bachtrog, 2015). The only exceptions are sex determination by maternal genotype in the blow fly *Chrysomya rufifacies* (Ullerich, 1963), and an expanded karyotype of 19–20 chromosomes in the flesh fly *Agria* (*Pseudosarcophaga*) *affinis* (Boyes, 1953). The X chromosomes of the Australian sheep blow fly (*Lucilia cuprina*) and gray flesh fly (*Sarcophaga bullata*) both correspond to element F (Vicoso and Bachtrog, 2013, 2015), suggesting that element F is the ancestral X of these families. In addition, the Y chromosomes of *L. cuprina* and *S. bullata* are extremely differentiated from their homologous X chromosomes (Vicoso and Bachtrog, 2013, 2015). The haploid X chromosome in *L. cuprina* males is up-regulated (i.e., dosage compensated) by an RNA-binding protein that is homologous to a *Drosophila* protein that binds nearly exclusively to element F (Linger *et al.*, 2015; Davis *et al.*, 2018). As expected because of the genetic differentiation between the *L. cuprina* X and Y, loss of function mutations in the *L. cuprina* gene encoding the dosage compensation protein are lethal specifically in males

(Davis *et al.*, 2018).

Figure 1: Cryptic sex chromosomes in Muscidae. Phylogenetic relationships and karyotypes of muscid flies and their relatives (Boyes *et al.*, 1964; Boyes and Van Brink, 1965; Schnell e Schuehli *et al.*, 2007; Vicoso and Bachtrog, 2015). The inferred mechanism of sex determination (Sex Detr) and the diploid chromosome number (2n) are listed for each species. "*M*" refers to a generic male-determining locus.

In contrast to flesh flies and blow flies, Muscidae seem to have a much higher rate of sex chromosome turnover. The iconic example of sex chromosome evolution in Muscidae is the house fly (*Musca domestica*), which has a well-characterized polygenic sex determination system (Hamm *et al.*, 2015). House fly appears to have the ancestral brachyceran karyotype (i.e., 5 pairs of euchromatic chromosomes and a heterochromatic sex chromosome pair), but the X and Y chromosomes are not differentiated (Meisel *et al.*, 2017). This is because the house fly male-determining locus (*Mdmd*) is a recently derived duplicated gene that can be found on at least 5 of the 6 chromosomes (Sharma *et al.*, 2017). The invasion and spread of this new male-determiner in the house fly genome since the divergence with closely related species is such that every house fly chromosome can be an undifferentiated proto-sex-chromosome pair.

In addition to house fly, other Muscidae have derived karyotypes that were evidenced by cytological examination (**Fig 1**). For example, the heterochromatic element F is missing from the karyotypes of the stable fly (*Stomoxys calcitrans*), horn fly (*Haematobia irritans*), and some other muscids (Boyes *et al.*, 1964; LaChance, 1964; Joslyn *et al.*, 1979; Avancini and Weinzierl, 1994; Parise-Maltempi and Avancini, 2007). These species have genetic sex determination with a dominant male-determining locus (Willis *et al.*, 1981; McDonald and Schmidt, 1987), but the specific identity of the X and Y chromosomes remains unresolved. We hypothesize that species with these derived karyotypes have cryptic sex chromosomes that arose when an ancestral sex chromosome (element F and/or the Y chromosome) recently fused to one of the other five chromosomes. Here, we describe the identification of the cryptic sex chromosomes in the stable fly and horn fly using genomic and transcriptomic sequence data. We provide evidence that the stable fly and horn fly sex chromosomes are young and of independent origins. These results demonstrate that muscid flies are a good model system for studying the factors that permit rapid sex chromosome turnover.

Methods

Assigning scaffolds to Muller elements

We used a homology-based approach that we had previously developed in house fly to map stable fly and horn fly scaffolds to Muller elements (Meisel *et al.*, 2015; Meisel and Scott, 2018). This approach works because Muller element gene content (synteny) is conserved across Brachycera (Foster *et al.*, 1981; Weller and Foster, 1993; Vicoso and Bachtrog, 2013; Sved *et al.*, 2016). For stable fly, we selected OrthoGroups from the OrthoDB annotation that contain a single *D. melanogaster* gene and a single *S. calcitrans* gene (Kriventseva *et al.*, 2018; Olafson *et al.*, 2019). For horn fly, we obtained annotated genes from the initial analysis of the genome, and we extracted the inferred *D. melanogaster* homologs for each gene (Konganti *et al.*, 2018). We assigned the stable fly and horn fly genes to the same Muller element as their *D. melanogaster* homologs. Each of these genes is part of a chromosomal scaffold. We used a majority-rules approach to assign those scaffolds to Muller elements if >50% of the genes on a scaffold are assigned to the same Muller element. This allowed us to assign 94.6% (1,482/1,566) of stable fly scaffolds and 97.5% (4,778/4,889) of horn fly

scaffolds containing annotated genes to Muller elements. All genes on a scaffold are then assigned to the Muller element of that scaffold regardless of the Muller element designation of their annotated ortholog.

Variant calling

Our approach to identifying sex chromosomes involves testing for Muller elements with increased heterozygosity in males, which is the expectation for young X-Y chromosome pairs (Vicoso and Bachtrog, 2015; Meisel *et al.*, 2017). To those ends, we used the Genome Analysis Toolkit (GATK) version 3.4-0 to identify heterozygous single nucleotide polymorphisms (SNPs) in stable fly and horn fly genomes and transcriptomes, following the GATK best practices (McKenna *et al.*, 2010; DePristo *et al.*, 2011; Van der Auwera *et al.*, 2013). The Illumina sequencing reads used to assemble the stable fly genome were generated from DNA extracted from males, which allows us to use those data to test for elevated male heterozygosity. The same cannot be done for horn fly because the sequences used to assemble the genome came from DNA isolated from mixed pools of males and females (Konganti *et al.*, 2018).

To quantify heterozygosity in stable fly males, we first mapped the sequencing reads to the assembled genomic scaffolds using the MEM algorithm implemented in BWA with the default parameters (Li and Durbin, 2009). Next, we used Picard Tools version 1.133 to identify and remove duplicate reads, and we realigned indels with GATK. Then, we performed naïve variant calling using the GATK HaplotypeCaller with a phred-scaled confidence threshold of 20, and we selected the highest confidence SNPs from that first-pass (QD < 2.0, MQ < 40, FS > 60, SOR > 4, MQRankSum < −12.5, ReadPosRankSum < −8). We used those high quality variants to perform base recalibration, we re-input those recalibrated bases into another round of variant calling, and we extracted the highest quality variants. We repeated the process so that we had performed three rounds of recalibration, which was sufficient for convergence of variant calls. We applied GenotypeGVCFs to the variant calls from all of the Illumina libraries for joint genotyping. We then used the GATK HaplotypeCaller to genotype all of the variable sites (phred-scaled confidence > 20), and we selected only the high quality variants (FS > 30 and QD < 2).

Comparing male and female heterozygosity, rather than only analyzing male heterozygosity, may be a better way to identify elevated heterozygosity in males. There are no genomic sequences available from female stable fly or from single-sex horn fly to use in such an analysis. However, Illumina RNA-seq data were collected from female and male tissues separately for both species. From stable fly, RNA-seq libraries were sequenced from female and male whole adults (SRX229930 and SRX229931), reproductive tissues (SRX995859 and SRX995857), and heads (SRX995858 and SRX995860). From horn fly, RNA-seq libraries were sequenced from ovary (SRX3340090) and testis (SRX3340086).

We used a modified GATK pipeline to identify SNPs in stable fly and horn fly RNA-seq data (Meisel *et al.*, 2017). First, RNA-seq reads were aligned to the reference genomes of the appropriate species using STAR version 2.4.0.1 (Dobin *et al.*, 2013). We used the aligned reads to create a new reference genome index from the inferred spliced junctions in the first alignment, and then we performed a second alignment with the new reference. We next marked duplicate reads and used SplitNCigarReads to reassign mapping qualities to 60 with the ReassignOneMappingQuality read filter for alignments with a mapping quality of 255. Indels were realigned, and three rounds of variant calling and base recalibration were performed as described above for the stable fly genomic sequencing reads. We applied GenotypeGVCFs to the variant calls from all tissues for joint genotyping of males and females from each species. Finally, we used the same filtering parameters that we applied to the stable fly genomic sequencing reads to extract high-quality SNPs from our variant calls.

Once we had identified sites in the sequence data that differ from the reference genome, we extracted heterozygous sites across the genome within annotated genes. We used those data to calculate the number of heterozygous SNPs per Mb within each annotated gene. Genes with no heterozygous sites were omitted from the results we present, but we obtain the same general patterns when these genes are included. For each gene, we calculated the fraction of all heterozygous sites in either sex that are heterozygous in males.

Gene expression analysis

We aligned the same RNA-seq data described above to the annotated transcripts in the stable fly and horn fly genomes and calculated transcripts per million reads (TPM) for each

transcript using kallisto version 0.44.0 (Bray *et al.*, 2016). We also used kallisto to align 454 GS FLX reads from whole adult female (SRR003191) and male (SRR003191) horn flies to the horn fly reference transcripts (Konganti *et al.*, 2018). In addition, we used the same approach to align Illumina RNA-seq reads from whole adult male (SRX208993 and SRX208994) and female (SRX208996 and SRX208997) house flies to the house fly reference genome (Scott *et al.*, 2014a). We then summed TPM for all transcripts from each gene for each sample type (e.g., stable fly female head) to get a gene-level estimate of expression in each sample. In house fly, where there are two RNA-seq libraries for each sex, we calculated the mean TPM for each gene across both libraries. Using these data, we calculated the $\log_2$-fold male:female expression level ($\log_2 \frac{M}{F}$) of each gene for each tissue type. In our analysis, we only considered genes with TPM > 0 in both males and females in a particular tissue and TPM > 1 in at least one sex.

Results

The stable fly sex chromosomes consist of elements D and F

We hypothesize that the stable fly has young, cryptic sex chromosomes (**Fig 1**). Nascent sex chromosomes can be identified based on elevated heterozygosity in the heterogametic sex (e.g., XY males) because the X and Y have begun to differentiate in the sequences of genes but the Y still retains most of the genes in common with the X chromosome (Vicoso and Bachtrog, 2015; Meisel *et al.*, 2017). The stable fly genome was sequenced from male DNA (Olafson *et al.*, 2019), allowing us to test for elevated heterozygosity using the genomic sequencing reads to identify the sex-linked element(s). To those ends, we first assigned most of the genes in the stable fly genome to Muller elements using homology relationships with *D. melanogaster* (see Methods). Next, we identified heterozygous SNPs in the sequencing reads generated from stable fly males. We found that genes assigned to Muller element D have more heterozygous SNPs than genes on scaffolds mapped to the other major elements ($P=10^{-138}$ in a Mann-Whitney test; **Fig 2A**). This suggests that element D is part of the stable fly X and Y chromosomes.

There are at least two concerns associated with our analysis of male heterozygosity based on the genome sequence reads. First, we may expect elevated male heterozygosity on ele-

Figure 2: Identifying stable fly and horn fly sex chromosomes. A. Boxplots show the number of heterozygous SNPs per megabase (Mb) identified using genomic sequencing reads within annotated stable fly genes mapped to each of the Muller elements. **B–C.** Boxplots show the fraction of heterozygous SNPs in males relative to females within annotated stable fly and horn fly genes mapped to each of the Muller elements. Dashed lines indicate the genome-wide average for all genes. Outliers were omitted from all plots. Inferred sex-linked elements are drawn in red.

ment F because it is the ancestral brachyceran X chromosome, but element F genes do not have an excess of heterozygous SNPs (**Fig 2A**). Our failure to detect elevated heterozygosity on element F is consistent with reduced variation along most of the length of *Drosophila* element F, likely because a lack of recombination enhances the diversity-reducing effects of selective sweeps and background selection (Berry *et al.*, 1991; Holly *et al.*, 1994; Jensen *et al.*, 2002; Wang *et al.*, 2002, 2004; Arguello *et al.*, 2010). Second, we expect elevated heterozygosity on a young sex chromosome in males relative to females, but we only assessed male heterozygosity without comparing to females. Comparing male and female heterozygosity may allow us to detect elevated male heterozygosity on element F.

To address those two concerns, we used available RNA-seq data to identify heterozygous SNPs in both males and females (Olafson *et al.*, 2019). We then calculated the fraction of all SNPs in each gene that are heterozygous in males. This value ranges from 0 if all heterozygous SNPs are in females to 1 if all heterozygous SNPs are in males. Consistent with our analysis of SNPs in male genomic DNA sequences, there is a higher fraction of male heterozy-

Figure 3: Fragmented assembly of the stable fly sex chromosome. A. The number of genomic scaffolds and **B.** the number of genes assigned to each stable fly Muller element. **C.** The composite length of all scaffolds assigned to each stable fly Muller element. **D.** The distributions of scaffold lengths and **E.** the distributions of genes per scaffold for scaffolds assigned to each stable fly Muller element are shown with boxplots. Inferred sex-linked elements are highlighted in red.

gous SNPs in genes on element D than in genes on the other major elements (P=0.047 in a Mann-Whitney test; **Fig 2B**). There is also a higher fraction of element D genes that only have heterozygous SNPs in males and not in females (518/1850 = 28.0%) when compared to genes on the other major elements (2168/8625 = 25.1%; z=2.56, P=0.005).

We also detect evidence for elevated male heterozygosity on stable fly element F (the ancestral X chromosome). Only 10 annotated stable fly genes are assigned to element F, of which five have heterozygous SNPs. Four of those five genes have more heterozygous SNPs in males than females. It is unlikely that $\geq$4/5 of element F genes would have >50% male heterozygous SNPs if heterozygosity were equal in males and female (z=1.34, P=0.09). In addition, three of the five element F genes only have heterozygous SNPs in males, which is more than the frequency of autosomal genes (25.1%) that only have male heterozygous SNPs (z=1.80, P=0.04). These results support the hypothesis that the stable fly X and Y chromosomes are young and minimally differentiated. They also suggest that the X chromosome consists of both elements D and F, as does the Y chromosome.

The genome assembly of stable fly provides a third line of support that element D has higher male heterozygosity and is therefore part of the X and Y chromosomes. We expect heterozygosity to interfere with genome assembly, leading to smaller contigs and scaffolds (Vinson *et al.*, 2005; Kajitani *et al.*, 2014; Pryszcz and Gabaldón, 2016). In the case of XY males with nascent sex chromosomes, this assembly fragmentation should be greater on sex-chromosome-derived scaffolds than autosomal scaffolds. More scaffolds in the stable fly assembly are assigned to element D than any of the other elements, even though element D does not have more genes or a larger inferred length than the other elements (**Fig 3A-C**). This suggests that the assembly of stable fly element D is more fragmented than the other four major elements. Consistent with this prediction, scaffolds assigned to stable fly element D are shorter ($P=10^{-15}$ in a Mann-Whitney test) and have fewer genes ($P=10^{-15}$ in a Mann-Whitney test) than scaffolds assigned to the other four major chromosomes (**Fig 3D-E**). Therefore, there are three lines of evidence that are all consistent with elevated heterozygosity on element D in stable fly males, providing a congruent picture that element D is part of a young X-Y chromosome pair. Stable fly element D is also enriched for a unique set of transposable elements (e.g., *Vingi* and *Dada*) that are not enriched on any other element (Olafson *et al.*, 2019), which is also expected for an evolving sex chromosome (Steinemann and Steinemann, 2005; Ellison and Bachtrog, 2013). The stable fly X and Y chromosomes also likely both contain element F, which probably fused to element D, because element F genes also have elevated male heterozygosity (**Fig 2B**).

The stable fly male-determining locus was previously mapped to chromosome 1 (Willis *et al.*, 1981). We therefore conclude that stable fly chromosome 1 corresponds to Muller elements D and F. The stable fly Y chromosome carries a male-determining locus, but we do not know the nature of this gene. The house fly male-determining gene (*Mdmd*) was not found in the stable fly genome or any other fly relatives (Sharma *et al.*, 2017). Therefore, either stable fly has an independently derived new male determiner or it has retained an ancestral male-determining locus that was replaced in house fly.

The horn fly sex chromosomes consist of elements A and F

We hypothesize that the horn fly also has a young X-Y chromosome pair. The horn fly genome was sequenced using DNA extracted from a mixed sample of males and females (Konganti *et al.*, 2018), which prevents us from using heterozygosity in the genome sequencing reads to identify the horn fly sex chromosomes. However, there is available RNA-seq data from horn fly testis and ovary, which we used to identify the horn fly sex chromosomes using the same approach as we did in stable fly. In horn fly, there is a higher fraction of male heterozygous SNPs in genes assigned to element A (P=0.024 in a Mann-Whitney test) and element F (P=0.0059 in a Mann-Whitney test) than in genes assigned to the other elements (**Fig 2C**). There is also a higher fraction of element A genes that only have heterozygous SNPs in males and not in females (385/889 = 43.3%) when compared to genes on the other major elements (1571/4163 = 37.7%; z=3.09, P=0.001).

Of the 44 horn fly genes assigned to element F, 11 have heterozygous SNPs. Of those 11 genes, 8 are only heterozygous in males, and the remaining 3 have more heterozygous SNPs in males than females. It is highly unlikely that >50% of heterogyzous SNPs would only be observed in males for all 11 of the horn fly element F genes if heterozygosity were equal in males and females (z=3.32, P=0.0005). There is also a higher fraction of element F genes that only have heterozygous SNPs in males when compared to genes on the autosomes (z=2.39, P=0.008). We therefore conclude that the horn fly X and Y chromosomes are likely both composed of elements A and F, and the sex chromosomes arose through a fusion of those two elements. We cannot test for a more fragmented assembly of the horn fly sex chromosomes because the entire horn fly genome assembly is fragmented. The scaffold N50 of the horn fly assembly is 23 Kb (Konganti *et al.*, 2018), and the vast majority ($4,112/4,778 = 86\%$) of horn fly scaffolds assigned to Muller elements have only 1 annotated gene.

Sex-biased gene expression on the sex chromosomes

Genes that are expressed at different levels between females and males are said to have "sex-biased" expression (Parsch and Ellegren, 2013). Genes with male-biased (female-biased) expression are often under- (over-) represented on fly X chromosomes as a result of the haploid dose of the X chromosome in males or sex-specific selection pressures that prevent

Figure 4: Sex-biased expression across Muller elements. The distributions of $\log_2$ male:female expression ($\log_2 \frac{M}{F}$) for genes on each Muller element in stable fly, horn fly, and house fly are shown with boxplots. Expression data are from either whole adult (all species), gonad (horn fly only), reproductive tissues (stable fly only), or head (stable fly only). Dashed lines indicate the genome-wide average for all genes. Outliers were omitted from all plots. Inferred sex-linked elements are drawn in red.

(favor) the evolution of male-biased (female-biased) expression on the X (Rice, 1984; Parisi *et al.*, 2003; Sturgill *et al.*, 2007; Meiklejohn and Presgraves, 2012; Meisel *et al.*, 2012). We cannot perform statistical tests for significant expression differences between males and females using available data from either stable fly or horn fly because only one replicate RNA-seq sample was collected for each tissue type from each sex for each species. However, we can calculate the relative expression of genes in males and females ($\log_2 \frac{M}{F}$) for each tissue type.

We compared the distributions of $\log_2 \frac{M}{F}$ for genes assigned to each Muller element in stable fly, horn fly, and the closely related house fly (**Fig 4**). Genes on stable fly element D (part of the X and Y chromosomes) have significantly lower $\log_2 \frac{M}{F}$ in reproductive tissues than genes on the other major elements ($P=0.0032$ in a Mann-Whitney test). The slight but significantly reduced $\log_2 \frac{M}{F}$ on element D is consistent with the "demasculinization" or "feminization" of the X chromosome observed in reproductive tissues of other flies (Parisi *et al.*, 2003; Sturgill *et al.*, 2007; Vicoso and Bachtrog, 2013, 2015). In contrast, genes on horn fly element F have higher $\log_2 \frac{M}{F}$ in gonad than genes on the autosomal elements ($P=10^{-4}$ in a Mann-Whitney test), suggesting a "masculinization" of the ancestral X chromosome. The small number of annotated element F genes in stable fly likely limit our ability to detect significant masculinization of stable fly element F (**Fig 4**). There is no significant difference in $\log_2 \frac{M}{F}$ between genes on horn fly element A (the new portion of the sex chromosomes) and the autosomes. House fly also has multiple young sex chromosome pairs, one of which includes the ancestral fly X chromosome, Muller element F (Meisel *et al.*, 2017). There is also no evidence for a difference in $\log_2 \frac{M}{F}$ between genes on the ancestral X chromosome (element F) and the autosomes in house fly. The minimal evidence for demasculinization or feminization of the X chromosome is consistent with young sex chromosomes in all three species.

Discussion

Stable fly and horn fly have derived karyotypes in which the ancestral X chromosome (Muller element F) and Y chromosome are not visible (**Fig 1**). We show, based on elevated male heterozygosity, that the X and Y chromosomes of stable fly both contain elements D and

Figure 5: Potential scenarios that could give rise to muscid cryptic sex chromosomes.
Ancestral autosomes (gray), X chromosome (red), and Y chromosome (black) are shown in both male and female karyotypes. Possible derived male and female genotypes are shown for three potential scenarios that could give rise to the cryptic sex chromosomes in stable fly and horn fly.

F (**Fig 2A-B**). The X and Y chromosomes of horn fly contain elements A and F (**Fig 2C**). Elevated male heterozygosity is a hallmark of a young sex chromosome pair (Vicoso and Bachtrog, 2015), suggesting that stable fly and horn fly have independently and recently derived nascent sex chromosomes. In addition, house fly also has multiple young proto-Y chromosomes (Meisel *et al.*, 2017). The minimal feminization/masculinization of the sex chromosomes across these three species is consistent with their recent origins (**Fig 4**). The slight masculinization of element F in horn fly suggests that a Y-linked copy of element F may have accumulated alleles that increase male expression (Zhou and Bachtrog, 2012), but additional work is necessary to test this hypothesis.

We propose three different scenarios that could have given rise to the stable fly and horn fly cryptic sex chromosomes (**Fig 5**). The order of most events in each scenario is arbitrary, and it is not necessary for the sex chromosomes of stable fly and horn fly to have arisen by the same scenario. All three scenarios assume that the MRCA of muscid flies had a karyotype with five euchromatic autosomes (elements A–E) and a heterochromatic sex chromosome pair (where element F is the X chromosome) because this is the ancestral karyotype of Brachycera (Vicoso and Bachtrog, 2013, 2015) and is still conserved in some Muscidae (**Fig 1**). We additionally assume that the Y chromosome of the MRCA of Muscidae carried a male-determining locus because that is the most common mechanism of sex determination in closely and distantly related families (Willhoeft and Franz, 1996; Bopp *et al.*, 2014; Scott *et al.*, 2014b).

In the first scenario (**Fig 5**), we hypothesize that the ancestral male-determining gene transposed from the ancestral Y chromosome to an autosome. This transposition event would convert the autosome into a proto-Y chromosome, and its homolog would be a proto-X chromosome. A transposing male-determining gene (*Mdmd*) was identified in house fly (Sharma *et al.*, 2017), demonstrating the feasibility of this scenario. We also hypothesize that the ancestral X chromosome (element F) fused to the same autosome containing the male-determining locus in this scenario. Importantly, element F must have fused to both a copy of the autosome with the male-determiner (the proto-Y), and a copy of the autosome without a male-determiner (the proto-X). This is because males must have two differentiated copies of element F (one X-linked and the other Y-linked) in order to explain the elevated

male heterozygosity of element F genes in both stable fly and horn fly (**Fig 2**). The order of the F-autosome fusion and the transposition of the male-determiner is arbitrary in this model. Finally, the ancestral Y chromosome either fused to one of the autosomes or it was lost from the genome. Fusion of an ancestral Y chromosome to an autosome (and reversion of the Y-linked genes to autosomal genes) happened following the creation of a neo-X chromosome in *Drosophila pseudoobscura* (Carvalho and Clark, 2005; Larracuente *et al.*, 2010). A Y-autosome fusion is possible in this scenario if the Y chromosome lost its male-determining activity, possibly via a pseudogenizing mutation. Alternatively, the ancestral Y chromosome could have fused to the element carrying the transposed male-determiner (the proto-Y), which would allow the ancestral Y to retain the male-determiner without creating an independently segregating second Y chromosome. If the ancestral Y chromosome was lost, we hypothesize that the ancestral Y did not contain any essential genes other than the male-determiner. Some *Drosophila* species have Y chromosomes that lack essential genes (Voelker and Kojima, 1971), demonstrating that it is feasible for a fly Y chromosome to not be essential for male viability or fertility. Moreover, the genetic differentiation of X and Y chromosomes in both blow fly and flesh fly could be explained by a lack of essential genes on their Y chromosomes other than the male-determiner (Vicoso and Bachtrog, 2013; Linger *et al.*, 2015; Vicoso and Bachtrog, 2015).

The second scenario differs from the first in that instead of the ancestral male-determiner transposing to an autosome, a new male-determiner arises on one of the autosomes (**Fig 5**). The new male-determiner would convert the autosome into a proto-Y chromosome, and its homolog would be a proto-X. The *Mdmd* gene in house fly arose from a highly conserved splicing factor that was duplicated after the divergence between house fly and stable fly (Sharma *et al.*, 2017), demonstrating that new male-determining genes can arise within Muscidae. As in the first scenario, element F would have fused to the same chromosome carrying the new male determiner (i.e., the proto-Y), and it must have also fused to the homologous proto-X to produce the elevated male heterozygosity we observe in element F genes (**Fig 2**). In addition, the ancestral Y was either lost or fused to an autosome, as in the first scenario.

In the third scenario we hypothesize that both the X and Y chromosomes of the MRCA of

Muscidae fused to the same ancestral autosome (**Fig 5**). These X- and Y-autosome fusions would convert one copy of the ancestral autosome into a neo-X chromosome and the other copy into a neo-Y chromosome. Element F genes have elevated heterozygosity in both stable fly and horn fly males (**Fig 2**), suggesting that males carry two copies of element F that have the same gene content and are only slightly differentiated at the sequence level. Therefore, this scenario requires the ancestral X and Y of Muscidae to be undifferentiated, with the Y chromosome essentially just a copy of element F that carries a male-determining locus. The X and Y chromosomes of house fly are very similar in gene content (Meisel *et al.*, 2017), but we do not know if this is the ancestral state of Muscidae or a derived condition in house fly.

Each of these scenarios makes predictions about the sex chromosomes of the MRCA of Muscidae, and testing those predictions is necessary in order to evaluate the scenarios. For example, if all muscid species with the ancestral karyotype have undifferentiated X and Y chromosomes (as in house fly), then we would infer that the MRCA had undifferentiated sex chromosomes, supporting scenario 3. Alternatively, if extant muscids with the ancestral karyotype have differentiated X and Y chromosomes, then we could infer that the MRCA had differentiated X and Y chromosomes. This would support scenarios 1 and 2.

In addition, identifying the male-determining genes across Muscidae would allow us to test if the same gene is used for male-determination across species or if new male-determiners have arisen in species other than the house fly (Sharma *et al.*, 2017). If the same male-determiner is used across species, then scenarios 1 and 3 would be supported. If new male-determiners arose in species with derived karyotypes, then scenario 2 would be supported. Therefore, by characterizing the sex chromosomes and male-determining genes across muscid flies, we can distinguish between all three scenarios. Distinguishing between the scenarios would provide valuable insights into the factors that promote high rates of sex chromosome turnover in muscid flies, which would serve as an informative model for understanding why rates of sex chromosome turnover differ across taxa. This goal should motivate future work in this system.

In summary, we have identified independently derived young sex chromosomes in stable fly and horn fly (**Fig 2**), which are different from the proto-sex chromosomes of house fly (Meisel *et al.*, 2017; Sharma *et al.*, 2017). Therefore, there are at least three independently

derived young sex chromosome systems in Muscidae, and probably more based on the derived karyotypes distributed across the family (**Fig 1**). In addition, we present three possible scenarios for the origins of the stable fly and horn fly sex chromosomes (**Fig 5**). Each scenario makes specific predictions about the male-determining genes and sex chromosomes in the MRCA of Muscidae and in other muscid species. Notably, the scenarios include two important factors that could allow for faster rates of sex chromosome turnovers in some taxa (e.g., Muscidae) than in closely related taxa (e.g., blow flies and flesh flies). First, the ancestral X and Y chromosomes of Muscidae could have been undifferentiated, in contrast to the differentiated blow fly and flesh fly sex chromosomes (Vicoso and Bachtrog, 2013; Linger *et al.*, 2015; Vicoso and Bachtrog, 2015; Davis *et al.*, 2018). Undifferentiated sex chromosomes could allow for a higher rate of sex chromosome turnover (Stöck *et al.*, 2011, 2013; Dufresnes *et al.*, 2015). Second, there could be a high rate of new or transposing male-determining genes across Muscidae, as is the case in house fly (Sharma *et al.*, 2017). These new or transposable male-determining genes could allow allow for a faster rate of sex chromosome turnover. Testing for undifferentiated sex chromosomes, new male-determining genes, or transposing male-determiners in Muscidae is therefore a promising approach to assess the relative importance of these factors in permitting a higher rate of sex chromosome turnover.

Acknowledgements

Some of the computational analyses reported were performed on the Maxwell Cluster that is part of the University of Houston Core facility for Advanced Computing and Data Science. We thank T. J. Raszick, S.-H. Sze, C. J. Coates, and A. M. Tarone for sharing unpublished results on repeat content in the stable fly genome.

References

Abbott, J. K., Nordén, A. K., and Hansson, B. 2017. Sex chromosome evolution: historical insights and future perspectives. *Proc. R. Soc. Lond., B, Biol. Sci.*, 284(1854): 20162806.

Arguello, J. R., Zhang, Y., Kado, T., Fan, C., Zhao, R., Innan, H., Wang, W., and Long, M. 2010. Recombination yet inefficient selection along the *Drosophila melanogaster* subgroup's fourth chromosome. *Mol. Biol. Evol.*, 27(4): 848–861.

Avancini, R. M. P. and Weinzierl, R. A. 1994. Karyotype of the horn fly, *Haematobia irritans* (L.) (Diptera, Muscidae). *Cytologia*, 59(3): 269–272.

Bachtrog, D. 2013. Y-chromosome evolution: emerging insights into processes of Y-chromosome degeneration. *Nat. Rev. Genet.*, 14(2): 113–124.

Bachtrog, D., Mank, J. E., Peichel, C. L., Kirkpatrick, M., Otto, S. P., Ashman, T.-L., Hahn, M. W., Kitano, J., Mayrose, I., Ming, R., Perrin, N., Ross, L., Valenzuela, N., and Vamosi, J. C. 2014. Sex determination: why so many ways of doing it? *PLoS Biol*, 12(7): e1001899.

Baker, R. H. and Wilkinson, G. S. 2010. Comparative genomic hybridization (CGH) reveals a neo-X chromosome and biased gene movement in stalk-eyed flies (genus *Teleopsis*). *PLoS Genet.*, 6(9): e1001121.

Berry, A. J., Ajioka, J. W., and Kreitman, M. 1991. Lack of polymorphism on the Drosophila fourth chromosome resulting from selection. *Genetics*, 129: 1111–1117.

Beukeboom, L. and Perrin, N. 2014. *The Evolution of Sex Determination*. Oxford University Press, New York, NY.

Blackmon, H. and Demuth, J. P. 2014. Estimating tempo and mode of Y chromosome turnover: Explaining Y chromosome loss with the fragile Y hypothesis. *Genetics*, 197(2): 561–572.

Bopp, D., Saccone, G., and Beye, M. 2014. Sex determination in insects: variations on a common theme. *Sex. Dev.*, 8(1-3): 20–28.

Boyes, J. W. 1953. Somatic chromosomes of higher diptera: II. Differentiation of sarcophagid species. *Can. J. Zool.*, 31(6): 561–576.

Boyes, J. W. 1961. Somatic chromosomes of higher diptera: V. Interspecific and intraspecific variation in the Calliphoridae. *Can. J. Zool.*, 39(4): 549–570.

Boyes, J. W. and Van Brink, J. M. 1965. Chromosomes of calyptrate diptera. *Can. J. Genet. Cytol.*, 7(4): 537–550.

Boyes, J. W., Corey, M. J., and Paterson, H. E. 1964. Somatic chromosomes of higher diptera IX. Karyotypes of some muscid species. *Can J Cytol*, 42: 1025–1036.

Bray, N. L., Pimentel, H., Melsted, P., and Pachter, L. 2016. Near-optimal probabilistic RNA-seq quantification. *Nat. Biotech.*, 34: 525–527.

Carvalho, A. B. and Clark, A. G. 2005. Y chromosome of *D. pseudoobscura* is not homologous to the ancestral *Drosophila* Y. *Science*, 307: 108–110.

Charlesworth, D. and Charlesworth, B. 1980. Sex differences in fitness and selection for centric fusions between sex-chromosomes and autosomes. *Genet. Res.*, 35: 205–214.

Charlesworth, D., Charlesworth, B., and Marais, G. 2005. Steps in the evolution of heteromorphic sex chromosomes. *Heredity*, 95: 118–128.

Davis, R. J., Belikoff, E. J., Scholl, E. H., Li, F., and Scott, M. J. 2018. *no blokes* is essential for male viability and X chromosome gene expression in the Australian sheep blowfly. *Curr. Biol.*, 28(12): 1987–1992.

DePristo, M. A., Banks, E., Poplin, R., Garimella, K. V., Maguire, J. R., Hartl, C., Philippakis, A. A., del Angel, G., Rivas, M. A., Hanna, M., McKenna, A., Fennell, T. J., Kernytsky, A. M., Sivachenko, A. Y., Cibulskis, K., Gabriel, S. B., Altshuler, D., and Daly, M. J. 2011. A framework for variation discovery and genotyping using next-generation DNA sequencing data. *Nat. Genet.*, 43(5): 491–498.

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., and Gingeras, T. R. 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*, 29(1): 15–21.

Dufresnes, C., Borzée, A., Horn, A., Stöck, M., Ostini, M., Sermier, R., Wassef, J., Litvinchuck, S. N., Kosch, T. A., Waldman, B., Jang, Y., Brelsford, A., and Perrin, N. 2015. Sex-chromosome homomorphy in Palearctic tree frogs results from both turnovers and X-Y recombination. *Mol. Biol. Evol.*, 32(9): 2328–2337.

Ellison, C. E. and Bachtrog, D. 2013. Dosage compensation via transposable element mediated rewiring of a regulatory network. *Science*, 342(6160): 846–850.

Foster, G. G., Whitten, M. J., Konovalov, C., Arnold, J. T. A., and Maffi, G. 1981. Autosomal genetic maps of the Australian sheep blowfly, *Lucilia cuprina dorsalis* R.-D. (Diptera: Calliphoridae), and possible correlations with the linkage maps of *Musca domestica* L. and *Drosophila melanogaster* (Mg.). *Genet. Res.*, 37: 55–69.

Hamm, R. L., Meisel, R. P., and Scott, J. G. 2015. The evolving puzzle of autosomal versus Y-linked male determination in *Musca domestica*. *G3*, 5: 371–384.

Holly, H., M., K. R., and Jody, H. 1994. Using hitchhiking genes to study adaptation and divergence during speciation within the *Drosophila melanogaster* species compelx. *Evolution*, 48(6): 1900–1913.

Jensen, M. A., Charlesworth, B., and Kreitman, M. 2002. Patterns of genetic variation at a chromosome 4 locus of Drosophila melanogaster and D. simulans. *Genetics*, 160: 493–507.

Joslyn, D. J., Seawright, J. A., and Willis, N. L. 1979. The karyotype of the stable fly, *Stomoxys Calcitrans* (L.) (Diptera: Muscidae). *Caryologia*, 32(3): 349–354.

Kajitani, R., Toshimoto, K., Noguchi, H., Toyoda, A., Ogura, Y., Okuno, M., Yabana, M., Harada, M., Nagayasu, E., Maruyama, H., Kohara, Y., Fujiyama, A., Hayashi, T., and Itoh, T. 2014. Efficient de novo assembly of highly heterozygous genomes from whole-genome shotgun short reads. *Genome Res.*, 24(8): 1384–1395.

Konganti, K., Guerrero, F. D., Schilkey, F., Ngam, P., Jacobi, J. L., Umale, P. E., Perez de Leon, A. A., and Threadgill, D. W. 2018. A whole genome assembly of the horn fly, *Haematobia irritans*, and prediction of genes with roles in metabolism and sex determination. *G3*, 8(5): 1675–1686.

Kriventseva, E. V., Kuznetsov, D., Tegenfeldt, F., Manni, M., Dias, R., Simão, F. A., and Zdobnov, E. M. 2018. OrthoDB v10: sampling the diversity of animal, plant, fungal, protist, bacterial and viral genomes for evolutionary and functional annotations of orthologs. *Nucl. Acids Res.*, 47(D1): D807–D811.

LaChance, L. E. 1964. Chromosome studies in three species of Diptera (Muscidae and Hypodermatidae). *Ann. Entomol. Soc. Am.*, 57(1): 69–73.

Larracuente, A. M., Noor, M. A. F., and Clark, A. G. 2010. Translocation of Y-linked genes to the dot chromosome in *Drosophila pseudoobscura*. *Mol. Biol. Evol.*, 27(7): 1612–1620.

Li, H. and Durbin, R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics*, 25: 1754–1760.

Linger, R. J., Belikoff, E. J., and Scott, M. J. 2015. Dosage compensation of X-linked Muller element F genes but not X-linked transgenes in the Australian sheep blowfly. *PLoS ONE*, 10(10): e0141544.

McDonald, P. T. and Schmidt, C. D. 1987. Genetics of permethrin resistance in the horn fly (Diptera: Muscidae). *J. Econ. Entomol.*, 80(2): 433–437.

McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., Garimella, K., Altshuler, D., Gabriel, S., Daly, M., and DePristo, M. A. 2010. The Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res.*, 20(9): 1297–1303.

Meiklejohn, C. D. and Presgraves, D. C. 2012. Little evidence for demasculinization of the *Drosophila* X chromosome among genes expressed in the male germline. *Genome Biol. Evol.*, 4(10): 1007–1016.

Meisel, R. P. and Scott, J. G. 2018. Using genomic data to study insecticide resistance in the house fly, *Musca domestica*. *Pestic. Biochem. Physiol.*, page 10.1016/j.pestbp.2018.01.001.

Meisel, R. P., Malone, J. H., and Clark, A. G. 2012. Disentangling the relationship between sex-biased gene expression and X-linkage. *Genome Res.*, 22: 1255–1265.

Meisel, R. P., Scott, J. G., and Clark, A. G. 2015. Transcriptome differences between alternative sex determining genotypes in the house fly, *Musca domestica*. *Genome Biol. Evol.*, 7(7): 2051–2061.

Meisel, R. P., Gonzales, C. A., and Luu, H. 2017. The house fly Y Chromosome is young and minimally differentiated from its ancient X Chromosome partner. *Genome Res.*, 27(8): 1417–1426.

Muller, H. J. 1940. Bearings of the 'Drosophila' work on systematics. In J. Huxley, editor, *The New Systematics*, pages 185–268. Clarendon Press, Oxford, Oxford.

Olafson, P. U., Aksoy, S., Attardo, G. M., Buckmeier, G., Chen, X., Coates, C. J., Davis, M., Dykema, J., Emrich, S. J., Friedrich, M., Holmes, C. J., Ioannidis, P., Jansen, E. N., Jennings, E. C., Lawson, D., Martinson, E. O., Maslen, G. L., Meisel, R. P., Murphy, T. D., Nayduch, D., Nelson, D. R., Oyen, K. J., Raszick, T. J., Ribeiro, J. M. C., Robertson, H. M., Rosendale, A. J., Sackton, T. B., Swiger, S. L., Sze, S.-H., Tarone, A. M., Taylor, D. B., Warren, W. C., Waterhouse, R. M., Weirauch, M. T., Werren, J. H., Wilson, R. K., Zdobnov, E. M., and Benoit, J. B. 2019. Functional genomics of the stable fly, *Stomoxys calcitrans*, reveals mechanisms underlying reproduction, host interactions, and novel targets for pest control. *bioRxiv*, 623009.

Parise-Maltempi, P. and Avancini, R. 2007. Comparative cytogenetic study in Muscidae flies. *Braz. J. Biol.*, 67: 945–950.

Parisi, M., Nuttall, R., Naiman, D., Bouffard, G., Malley, J., Andrews, J., Eastman, S., and Oliver, B. 2003. Paucity of genes on the *Drosophila* X chromosome showing male-biased expression. *Science*, 299: 697–700.

Parsch, J. and Ellegren, H. 2013. The evolutionary causes and consequences of sex-biased gene expression. *Nat. Rev. Genet.*, 14(2): 83–87.

Pennell, M. W., Kirkpatrick, M., Otto, S. P., Vamosi, J. C., Peichel, C. L., Valenzuela, N., and Kitano, J. 2015. Y fuse? sex chromosome fusions in fishes and reptiles. *PLoS Genet.*, 11(5): e1005237.

Pryszcz, L. P. and Gabaldón, T. 2016. Redundans: an assembly pipeline for highly heterozygous genomes. *Nucl. Acids Res.*, 44(12): e113.

Qiu, S., Bergero, R., and Charlesworth, D. 2013. Testing for the footprint of sexually antagonistic polymorphisms in the pseudoautosomal region of a plant sex chromosome pair. *Genetics*, 194(3): 663–672.

Rice, W. R. 1984. Sex chromosomes and the evolution of sexual dimorphism. *Evolution*, 38: 735–742.

Rice, W. R. 1986. On the instability of polygenic sex determination: the effect of sex-specific selection. *Evolution*, 40(3): 633–639.

Roberts, R. B., Ser, J. R., and Kocher, T. D. 2009. Sexual conflict resolved by invasion of a novel sex determiner in Lake Malawi cichlid fishes. *Science*, 326: 998–1001.

Schnell e Schuehli, G., de Carvalho, C. J. B., and Wiegmann, B. M. 2007. Molecular phylogenetics of the Muscidae (Diptera: Calyptratae): new ideas in a congruence context. *Invertebrate Systematics*, 21(3): 263–278.

Scott, J. G., Warren, W. C., Beukeboom, L. W., Bopp, D., Clark, A. G., Giers, S. D., Hediger, M., Jones, A. K., Kasai, S., Leichter, C. A., Li, M., Meisel, R. P., Minx, P., Murphy, T. D., Nelson, D. R., Reid, W. R., Rinkevich, F. D., Robertson, H. M., Sackton, T. B., Sattelle, D. B., Thibaud-Nissen, F., Tomlinson, C., van de Zande, L., Walden, K. K. O., Wilson, R. K., and Liu, N. 2014a. Genome of the house fly, *Musca domestica* L., a global vector of diseases with adaptations to a septic environment. *Genome Biol.*, 15(10): 466.

Scott, M. J., Pimsler, M. L., and Tarone, A. M. 2014b. Sex determination mechanisms in the Calliphoridae (blow flies). *Sex. Dev.*, 8(1-3): 29–37.

Sharma, A., Heinze, S. D., Wu, Y., Kohlbrenner, T., Morilla, I., Brunner, C., Wimmer, E. A., van de Zande, L., Robinson, M. D., Beukeboom, L. W., and Bopp, D. 2017. Male sex in houseflies is determined by *Mdmd*, a paralog of the generic splice factor gene *CWC22*. *Science*, 356(6338): 642–645.

Steinemann, S. and Steinemann, M. 2005. Y chromosomes: born to be destroyed. *BioEssays*, 27(10): 1076–1083.

Stöck, M., Horn, A., Grossen, C., Lindtke, D., Sermier, R., Betto-Colliard, C., Dufresnes, C., Bonjour, E., Dumas, Z., Luquet, E., Maddalena, T., Sousa, H. C., Martinez-Solano, I., and Perrin, N. 2011. Ever-young sex chromosomes in european tree frogs. *PLoS Biol.*, 9(5): e1001062.

Stöck, M., Savary, R., Betto-Colliard, C., Biollay, S., Jourdan-Pineau, H., and Perrin, N. 2013. Low rates of X-Y recombination, not turnovers, account for homomorphic sex chromosomes in several diploid species of Palearctic green toads (*Bufo viridis* subgroup). *J. Evol. Biol.*, 26(3): 674–682.

Sturgill, D., Zhang, Y., Parisi, M., and Oliver, B. 2007. Demasculinization of X chromosomes in the *Drosophila* genus. *Nature*, 450: 238–241.

Sved, J. A., Chen, Y., Shearman, D., Frommer, M., Gilchrist, A. S., and Sherwin, W. B. 2016. Extraordinary conservation of entire chromosomes in insects over long evolutionary periods. *Evolution*, 70(1): 229–234.

Swain, A. and Lovell-Badge, R. 1999. Mammalian sex determination: a molecular drama. *Genes Dev.*, 13(7): 755–767.

Ullerich, F.-H. 1963. Geschlechtschromosomen und Geschlechtsbestimmung bei einigen Calliphorinen (Calliphoridae, Diptera). *Chromosoma*, 14(1): 45–110.

Van der Auwera, G. A., Carneiro, M. O., Hartl, C., Poplin, R., del Angel, G., Levy-Moonshine, A., Jordan, T., Shakir, K., Roazen, D., Thibault, J., Banks, E., Garimella, K. V., Altshuler, D., Gabriel, S., and DePristo, M. A. 2013. From fastq data to high-confidence variant calls: The genome analysis toolkit best practices pipeline. *Curr. Protoc. Bioinformatics.*, 43: 1–33.

van Doorn, G. S. 2014. Patterns and mechanisms of evolutionary transitions between genetic sex-determining systems. *Cold Spring Harbor Perspectives in Biology*, 6(8): a017681.

van Doorn, G. S. and Kirkpatrick, M. 2007. Turnover of sex chromosomes induced by sexual conflict. *Nature*, 449(7164): 909–912.

van Doorn, G. S. and Kirkpatrick, M. 2010. Transitions between male and female heterogamety caused by sex-antagonistic selection. *Genetics*, 186(2): 629–645.

Vicoso, B. and Bachtrog, D. 2013. Reversal of an ancient sex chromosome to an autosome in *Drosophila*. *Nature*, 499(7458): 332–335.

Vicoso, B. and Bachtrog, D. 2015. Numerous transitions of sex chromosomes in Diptera. *PLoS Biol.*, 13(4): e1002078.

Vinson, J. P., Jaffe, D. B., O'Neill, K., Karlsson, E. K., Stange-Thomann, N., Anderson, S., Mesirov, J. P., Satoh, N., Satou, Y., Nusbaum, C., Birren, B., Galagan, J. E., and Lander, E. S. 2005. Assembly of polymorphic genomes: Algorithms and application to *Ciona savignyi*. *Genome Res.*, 15(8): 1127–1135.

Voelker, R. A. and Kojima, K.-i. 1971. Fertility and fitness of XO males in *Drosophila* i. qualitative study. *Evolution*, 25(1): 119–128.

Wang, J., Na, J.-K., Yu, Q., Gschwend, A. R., Han, J., Zeng, F., Aryal, R., VanBuren, R., Murray, J. E., Zhang, W., Navajas-Pérez, R., Feltus, F. A., Lemke, C., Tong, E. J., Chen, C., Man Wai, C., Singh, R., Wang, M.-L., Min, X. J., Alam, M., Charlesworth, D., Moore, P. H., Jiang, J., Paterson, A. H., and Ming, R. 2012. Sequencing papaya X and Yh chromosomes reveals molecular basis of incipient sex chromosome evolution. *Proc. Natl. Acad. Sci. U.S.A.*, 109(34): 13710–13715.

Wang, W., Thornton, K., Berry, A., and Long, M. 2002. Nucleotide variation along the *Drosophila melanogaster* fourth chromosome. *Science*, 295: 134–137.

Wang, W., Thornton, K., Emerson, J. J., and Long, M. 2004. Nucleotide variation and recombination along the fourth chromosome in *Drosophila simulans*. *Genetics*, 166(4): 1783–1794.

Weller, G. L. and Foster, G. G. 1993. Genetic maps of the sheep blowfly *Lucilia cuprina*: linkage-group correlations with other dipteran genera. *Genome*, 36(3): 495–506.

Wiegmann, B. M., Trautwein, M. D., Winkler, I. S., Barr, N. B., Kim, J.-W., Lambkin, C., Bertone, M. A., Cassel, B. K., Bayless, K. M., Heimberg, A. M., Wheeler, B. M., Peterson, K. J., Pape, T., Sinclair, B. J., Skevington, J. H., Blagoderov, V., Caravas, J., Kutty, S. N., Schmidt-Ott, U., Kampmeier, G. E., Thompson, F. C., Grimaldi, D. A., Beckenbach, A. T., Courtney, G. W., Friedrich, M., Meier, R., and Yeates, D. K. 2011. Episodic radiations in the fly tree of life. *Proc. Natl. Acad. Sci. U.S.A.*, 108(14): 5690–5695.

Willhoeft, U. and Franz, G. 1996. Identification of the sex-determining region of the *Ceratitis capitata* Y chromosome by deletion mapping. *Genetics*, 144(2): 737–745.

Willis, N. L., Seawright, J. A., Nickel, C., and Joslyn, D. J. 1981. Reciprocal translocations and partial correlation of chromosomes in the stable fly. *J. Hered.*, 72(2): 104–106.

Zhou, Q. and Bachtrog, D. 2012. Sex-specific adaptation drives early sex chromosome evolution in *Drosophila*. *Science*, 337(6092): 341–345.