Limits in the detection of m6A changes using MeRIP/m6A-seq

Alexa B.R. McIntyre[1,2]*, Nandan S. Gokhale[3], Leandro Cerchietti[4], Samie R. Jaffrey[5], Stacy M. Horner[3,6]*,

Christopher E. Mason[1,7,8,9]*

[1]Department of Physiology and Biophysics, Weill Cornell Medicine, New York City, NY 10065

[2]Tri-Institutional Program in Computational Biology and Medicine, New York City, NY 10065

[3]Department of Molecular Genetics and Microbiology, Duke University Medical Center, Durham, NC

27710

[4]Division of Hematology and Medical Oncology. Weill Cornell Medicine, New York City, NY 10065

[5]Department of Pharmacology, Weill Cornell Medicine, New York City, NY 10065

[6]Department of Medicine, Duke University Medical Center, Durham, NC 27710

[7]The HRH Prince Alwaleed Bin Talal Bin Abdulaziz Alsaud Institute for Computational Biomedicine, Weill

Cornell Medicine, New York, NY 10021

[8]The Feil Family Brain and Mind Research Institute, Weill Cornell Medicine, New York, NY 10065

[9]The WorldQuant Initiative for Quantitative Prediction, Weill Cornell Medicine, New York, NY 10021

*corresponding authors:

Alexa B.R. McIntyre, abm237@cornell.edu

Stacy M. Horner, stacy.horner@duke.edu

Christopher E. Mason chm2042@med.cornell.edu

Abstract

Many cellular mRNAs contain the modified base m^{6}A, and recent studies have suggested that various

stimuli can lead to changes in m^{6}A. The most common method to map m^{6}A and to predict changes in m^{6}A

between conditions is methylated RNA immunoprecipitation sequencing (MeRIP-seq), through which

methylated regions are detected as peaks in transcript coverage from immunoprecipitated RNA relative to

input RNA. Here, we generated replicate controls and reanalyzed published MeRIP-seq data to estimate

reproducibility across experiments. We found that m^{6}A peak overlap in mRNAs varies from ~30 to 60%

between studies, even in the same cell type. We then assessed statistical methods to detect changes in

m^{6}A peaks as distinct from changes in gene expression. However, from these published data sets, we

detected few changes under most conditions and were unable to detect consistent changes across

studies of similar stimuli. Overall, our work identifies limits to MeRIP-seq reproducibility in the detection

both of peaks and of peak changes and proposes improved approaches for analysis of peak changes.

Keywords

RNA base modifications, *N*6-methyladenosine, MeRIP-seq, m^{6}A-seq, RNA immunoprecipitation

sequencing, statistical methods, bioinformatics

Introduction

Methylation at the *N*6 position in adenosine (m^{6}A) is the most common internal modification in

eukaryotic mRNA. A methyltransferase complex composed of METTL3, METTL14, WTAP, VIRMA, and

other cofactors catalyzes methylation at DRACH/DRAC motifs, primarily in the last exon (1,2). Most m^{6}A

methylation occurs during transcription (3). The modification then affects mRNA metabolism through

recognition by RNA-binding proteins that regulate processes including translation and mRNA degradation

(4–9). However, whether m^{6}A is lost and gained in response to various cellular changes remains

contentious (3,10–15). To assess the evidence for proposed dynamic changes in m^{6}A, a reliable and

reproducible method to detect changes in methylation as distinct from changes in gene expression is

necessary.

The first and most widely-used method to enable transcriptome-wide studies of m⁶A, MeRIP-seq

or m⁶A-seq, involves the immunoprecipitation of m⁶A-modified RNA fragments followed by peak detection

through comparison to background gene coverage (16,17). A second method was developed in 2015,

miCLIP or m⁶A-CLIP, which involves crosslinking at the site of antibody binding to induce mutations

during reverse transcription for single-nucleotide detection of methylated bases (2,18). MeRIP-seq is still

more often used than miCLIP, despite less precise localization of m⁶A to peak regions of approximately

50-200 base pairs that can contain multiple DRAC motifs, since it follows a simpler protocol, requires less

starting material, and generally produces higher coverage of more transcripts. Antibodies for m⁶A can

also detect a second base modification, N^6,2′-O-dimethyladenosine (m⁶A$_m$), found at a lower abundance

than m⁶A and located at the 5′ ends of select transcripts (15,18). We thus refer to the base modifications

detected through MeRIP-seq collectively as m⁶A$_{(m)}$, although most are likely m⁶A. As of late 2018, over

fifty studies used MeRIP-seq to detect m⁶A$_{(m)}$ in mammalian mRNA (**Additional File 1: Supplementary**

Table 1).

Although MeRIP-seq can reveal approximate sites of m⁶A$_{(m)}$, it cannot be used to quantitatively

measure the fraction of transcript copies that are methylated (19). Studies of m⁶A variation in response to

stimuli instead estimate differences at individual loci through changes in peak presence or peak height.

Using these approaches, studies have reported changes to m⁶A with heat shock, microRNA expression,

transcription factor expression, cancer, oxidative stress, human immunodeficiency virus (HIV) infection,

Kaposi's sarcoma herpesvirus (KSHV) infection, and Zika virus infection, including hundreds to thousands

of changes in enrichment at specific sites (20–29). Statistical approaches to analysis have only recently

been published and there have been no comprehensive evaluations of methods to detect changes in m⁶A

based on MeRIP-seq data (30,31). Thus, while studies have suggested that m⁶A shows widespread

changes in response to diverse stimuli, they have applied inconsistent analysis methods to detect

changes in m⁶A and often don't control for differences in RNA expression between conditions or typical

variability in peak heights between replicates. In some cases, these studies have reported m⁶A changes

based on simple differences in peak count (24,26,27,32). However, others have applied statistical tests or

thresholds for differences in immunoprecipitated (IP) over input fraction enrichment and visual analysis of

coverage plots, and have reported fewer m^6A changes or suggested that m^6A is a relatively stable mark

(33,34). As in RNA-seq, there is noise in MeRIP-seq, and multiple replicates are therefore necessary to

estimate variance and statistically identify the effects of experimental intervention (35–37). To date, only

one MeRIP-seq study has used more than three replicates per condition (34), while ten have used only

one (17,20,32,33,38–43), suggesting that most studies may not have enough power to detect changes in

$m^6A_{(m)}$.

To re-evaluate the evidence for $m^6A_{(m)}$ changes under various conditions, we first examined the

variability in $m^6A_{(m)}$ detection across replicates, cell lines, and experiments using our own negative

controls (12 replicates) as well as 24 published MeRIP-seq data sets. We then compared statistical

methods to detect differences in IP enrichment using biological negative and positive controls for m^6A

changes. We found that these methods are limited by noise, including biological variability from changes

in RNA expression and technical variability from immunoprecipitation and sequencing that limits

reproducibility across studies. Our results suggest that the scale of statistically detectable $m^6A_{(m)}$ changes

in response to various stimuli is orders of magnitude lower than the scale of changes reported in many

studies. However, we also found that statistical detection could miss the majority of changed sites when

using only 2-3 replicates. We use our results to propose approaches to MeRIP-seq experimental design

and analysis to improve reproducibility and more accurately measure differential regulation of $m^6A_{(m)}$ in

response to stimuli. These data and analyses emphasize the need for further research and alternative

assays, for example recently developed endoribonuclease-based sequencing methods (44,45) or direct

RNA nanopore sequencing (46), to resolve the extent to which m^6A changes in response to specific

conditions.

Results

Detection of peaks across replicates, experiments, and cell types

The first steps in MeRIP-seq data analysis are to align sequencing reads to the genome or

transcriptome of origin and to identify peaks in transcript coverage in the IP fraction relative to the input

control. Several methods have been developed for MeRIP-seq peak detection, including exomePeak,

MeTPeak, MeTDiff, and bespoke scripts. Another method often used for MeRIP-seq peak detection is

MACS2, which was originally designed to detect protein binding sites in DNA from chromatin

immunoprecipitation sequencing (ChIP-seq). We compared $m^6A_{(m)}$ peak detection by exomePeak,

MeTPeak, MeTDiff, and MACS2 (31,47–49) in seven replicates of MeRIP-seq data obtained from mouse

cortices under basal conditions (34), and in 12 replicates of MeRIP-seq data we generated from human

liver Huh7 cells (50). The intersect between all tools tested was high, and we saw minimal differences in

DRAC motif enrichment, which we use to provide an estimate of tool precision in the absence of true

positive m^6A sites (**Additional File 2: Supplementary Figure 1a**). In addition, we assessed the

METTL3/METTL14-dependence of specific peaks identified by single tools using MeRIP-RT-qPCR. We

found that of these peaks, 4/4 from MACS2, 5/5 from MeTPeak, and 4/5 from MeTDiff showed decreased

$m^6A_{(m)}$ enrichment following METTL3/METTL14 depletion, suggesting that these are true m^6A sites. By

comparison, only 1/5 of the peaks uniquely called by exomePeak showed statistically significant

decreases ($p < 0.05$), although replicate variance was high and 4/5 showed a downward trend

(**Additional File 2: Supplementary Figure 1b**). Since MACS2 was the most commonly used tool for

peak calling and was previously found to perform well in comparison with a graphical user interface tool

and several other peak callers (51), we used MACS2 for the remainder of our analyses. Repeating the

analyses shown in Figures 2-4 using the MeTDiff peak caller instead of MACS2 did not affect any of our

conclusions (**Additional File 3**).

For $m^6A_{(m)}$ peak detection, a transcript must be sufficiently expressed for enrichment by the

$m^6A_{(m)}$ antibody and for adequate sequencing coverage in both the IP and input fractions. Previous

reports have suggested that $m^6A_{(m)}$ presence does not decrease with lower mRNA expression level, and,

if anything, is higher in mRNAs with lower expression as methylated transcripts tend to be less stable

(9,38). Peak callers, however, identify fewer peaks in genes at low expression, which we therefore

assume reflects inadequate coverage for peak calling. To estimate the level of coverage necessary for

peak detection, we analyzed the percent of genes with at least one, two, or three peaks relative to mean

input transcript coverage in both the mouse cortex and Huh7 cell data (**Figure 1a**). Based on the upper

shoulders of the sigmoidal curves as the percent of genes with peaks begins to plateau, we estimate that

mean gene coverage of approximately 10-50X is necessary to avoid missing peaks based on insufficient

coverage. Including a wider array of samples in this analysis likewise showed an increase in the percent

of transcripts with ≥1 peak as coverage rose to 10X (**Additional File 2: Supplementary Figure 1c**). Our

analysis of the input RNA-seq coverage of peak regions alone again supported a similar threshold; few

peaks are detected with median input read counts below 10 across replicates (**Additional File 2:**

Supplementary Figure 1d). These thresholds do not mean that peaks in genes with mean coverage <

10X or peaks with fewer than 10 input reads are false positives, but that the likelihood of false negatives

rises with lower coverage (**Additional File 2: Supplementary Figure 1e**).

Figure 1: Thresholds for peak detection. a) $m^6A_{(m)}$ site detection in MeRIP-seq data from mouse cortex (left) and human liver cells (Huh7, right) shows saturation of peak detection as transcript coverage approaches 10-50X for replicates at basal conditions, with peaks merged from all replicates. **b)** The total number of peaks captured increases with more replicates, with single replicates capturing a median of 66-78% of total peaks depending on study. Boxes span the 1st to 3rd quartiles of distributions for random subsamples of replicates, with lines indicating the median number of peaks, and whiskers showing the minimum and maximum points within ±1.5x the interquartile distance from the boxes. Jittered points show results for each random subsample (a total of 6 subsamples per replicate number for the mouse cortex data and 12 for the Huh7 data). **c)** The percent of peaks detected in at least r replicates for the same data sets.

To evaluate the reproducibility of MeRIP-seq data, we next examined the consistency of $m^6A_{(m)}$

peak calling between replicates. Previous studies have reported that peak overlap between replicates is

approximately 80% (9,16,52,53). Similarly, we found that between two replicates, $\log_2$ fold enrichment of

IP over input reads at detected peaks showed a Pearson correlation of approximately 0.81 to 0.86

(**Additional File 2: Supplementary Figure 1f**). A single sample captured a median of 78% of the peaks

found in seven replicates of mouse cortex data and 66% of peaks found in twelve replicates of Huh7 cell

data. The number of detected peaks increased log-linearly with the addition of more replicates, such that

with three replicates, 84-92% of the peaks found with 7-12 replicates were detected (**Figure 1b**).

Conversely, the number of peaks in common across replicates decreased as the number of replicates

increased, such that while ~80% of peaks were detected in at least two replicates, only ~60% were

detected in six replicates for both data sets and ~25% in all twelve replicates of Huh7 cell data (**Figure**

1c). Detection of peaks in more replicates did not increase DRAC motif enrichment (**Additional File 2:**

Supplementary Figure 1g). These results suggest that many $m^6A_{(m)}$ sites may be missed in studies that

use one to three replicates, and that increasing replicates could enable detection of more peaks.

However, not all peaks correspond to true $m^6A_{(m)}$ sites. A recent comparison to data from an

endoribonuclease-based method for m^6A detection suggested MeRIP-seq has a false positive rate of

~11%, although this would differ by study and detection threshold (3,54).

The number of peaks detected across studies varies. Given that coverage affects peak detection,

we hypothesized that variation in sequencing depth could contribute to differences in peak count. Zeng et

al. (2018) reported that peak count begins to saturate by around 20 million reads by subsampling data

within individual studies (42). However, we found that there is no positive correlation between peak count

and input or IP sequencing depth across data sets from different published studies, each of which had 3-

81M reads per replicate (input Pearson's R = -0.37, p = 0.015; IP Pearson's R = -0.17, p = 0.28)

(**Additional File 1: Supplementary Table 2, Additional File 2: Supplementary Figure 2a-b**). This

implies that experimental factors beyond sequencing depth contribute to the variability of peak counts

across studies.

Figure 2: Reproducibility of peak detection. a) Peak detection between studies that used the same cell type shows variable overlap. Overlap was calculated as the percent of peaks detected in Experiment 1 with an overlap of ≥ 1 base pair with peaks from Experiment 2. **b)** Peak detection across tissue and cell types shows samples from the same study cluster better together than samples from the same tissue. Median overlap was 46%. **c)** Peak detection across tissue types for data from the same study (Xiao et al., 2019). Median overlap was 72%. Studies used in (a) and (b) are described in **Additional File 1: Supplementary Table 2**.

We next analyzed the overlap of peaks among studies and found inconsistency in peak

localization on transcripts as well. Within four commonly used cell types, the percent of peaks detected in

one experiment that were also detected in a second varied among pairs of studies from as low as 2% of

peaks to as high as 90% (median = 45%), after filtering for transcripts expressed above a mean of 10X

input coverage in both to ensure sufficient expression for peak detection (**Figure 2a**). In fact, peaks

showed higher overlap within different cell types from the same study than within the same cell type from

different studies, suggesting that MeRIP-seq data is prone to strong batch effects (**Figure 2b**). While this

could be due to differences among experimental protocols used (summarized in **Additional File 1:**

Supplementary Table 2), we were unable to identify such a link. Overall, most percent overlaps of $m^6A_{(m)}$

peaks fell between ~30% (1st quartile) and ~60% (3rd quartile) (**Figure 2b**). With rare exceptions (e.g. that

described by Ke et al., 2017 in their Supplementary Figure 8 (3)), most MeRIP-seq data sets do show

enrichment of the m^6A motif DRAC. These results indicate, however, that multiple labs running MeRIP-

seq on the same cell type will detect different subsets of $m^6A_{(m)}$ sites. Possible contributing factors in the

differences among studies include cell state (e.g. different stages of the cell cycle), experimental

conditions, and sequencing depth. Despite predictions that tissue or cell type would be a large factor in

differences among samples, though, peaks detected in different tissues analyzed in a single experiment

showed high overlap and little clustering by tissue type (**Figure 2c**) (55). This suggests that although

there is evidence that m^6A levels vary by tissue (19), modified sites are consistent.

Detection of changes in peaks between conditions

Following $m^6A_{(m)}$ peak detection, many studies compare the expression of peaks between two

conditions to predict peak changes. While looking at plots of IP and input gene coverage under different

conditions can help evaluate the evidence for these changes (33), statistical or heuristic methods are first

necessary to narrow down a list of candidate sites to plot. Several tools used for statistical analysis by

the studies in **Additional File 1: Supplementary Table 1** or for other types of RNA IP sequencing assays

model peak counts using either (a) the Poisson distribution, in which the variance of a measure (here,

read counts) is assumed to be equal to the mean (MeTDiff), or (b) the negative binomial distribution, in

which a second parameter allows for independent adjustment of mean and variance (QNB and two implementations of a generalized linear model approach using DESeq2 or edgeR, **Table 1**) (30,31,56– 58). In the mouse cortex and Huh7 cell data, we found that, similar to RNA-seq data (24,57,59), the variance in read counts under peaks exceeded their mean, indicative of overdispersion (**Additional File 2: Supplementary Figure 3a**). The log likelihood (the probability of an observation given a distribution with known parameters) for our sample also fell within the distribution of expected log likelihoods for the negative binomial distribution (bottom) but not the Poisson distribution (top) (**Figure 3a**). Thus, the negative binomial distribution captures the mean-variance relationship in MeRIP-seq data, suggesting that tools that account for overdispersion better model the distribution of read counts at $m^6A_{(m)}$ peaks than tools that do not.

Table 1: Statistical methods for the detection of peak changes

Method	Read count distribution	Publication
MeTDiff	Poisson	Cui et al. (2018)
Quad-negative binomial (QNB)	Negative binomial	Liu et al. (2017)
GLM (DESeq2)	Negative binomial	based on Park et al. (2014)
GLM (edgeR)	Negative binomial	method for HITS-CLIP

We next defined positive and negative controls to evaluate tool performance for detection of changes in $m^6A_{(m)}$ peaks. Past publications describing new methods to detect $m^6A_{(m)}$ peak changes have used data sets in which methylation machinery genes or the methyl donor were disrupted compared to baseline conditions as positive controls, and have simulated negative controls by randomly swapping labels in the positive controls (30,31). However, swapping labels for conditions that may feature differences in gene expression in addition to differences in m^6A levels could unrealistically increase variance in read counts within groups. Therefore, we instead used the two data sets from mouse cortex and Huh7 cells, which each comprised many replicates at baseline conditions (n=7 and n=12,

 respectively), as negative controls. We randomly divided the mouse cortex data into two groups of three

 to four replicates for comparison and divided the Huh7 replicates by lab of incubation, which did not affect

 sample clustering (**Additional File 2: Supplementary Figure 3b**). We would expect to see minimal

 changes in IP enrichment at m6A peaks between groups for our negative controls, whereas our positive

 controls, which featured genetic or chemical interference with the m6A machinery, should show

 discernible differences in peaks when compared to baseline or wildtype conditions in the same cell lines

 (summarized in **Additional File 1: Supplementary Table 3**). Indeed, the absolute difference in $\log_2$ fold

 change between peaks and genes was centered around 0 for the negative controls and showed small

 shifts that varied in magnitude and direction for the positive controls (**Additional File 2: Supplementary**

 Figure 3c).

Figure 3: Analysis of methods to detect peak changes disproportional to gene expression changes. a) A comparison of Poisson (above) and negative binomial (below) models for read counts under peaks. The negative binomial mean log likelihood of the sample data fell within the 74th and 89st percentiles of 500 simulations for mouse cortex and Huh7 cell data, respectively, while the Poisson model failed to capture the sample distributions. **b)** The percent of sites below an unadjusted p-value threshold of 0.05 for different methods (described in **Table 1**) to detect differential methylation in negative controls between two groups at baseline conditions and positive controls in which methylation processes were disrupted with respect to baseline conditions (**Additional File 1: Supplementary Table 3**). The line at 5% indicates the expected proportion of sites given a uniform p-value distribution (see **Additional File 2: Supplementary Figure 2c**), while colours indicate negative (orange) and positive (purple) control experiments. **c)** The correlation between change in gene expression and change in peak expression between conditions for sites identified as differentially methylated in the eight positive control experiments. Pearson's R = 0.22, 0.10, 0.55, and 0.14 for edgeR, DESeq2, MeTDiff, and QNB, respectively, with p = 0.05, 0.09, 5.8E-87, and 2.4E-11. **d)** Coverage plots showing changes in peak expression are proportional to changes in gene expression for genes identified as differentially methylated by Bertero et al. (2018) using MeTDiff with activin signaling and an activin-NODAL inhibitor, SB431542 (SB). Lines show the mean coverage across three replicates, while shading shows the standard deviation. Peaks detected as significantly changed are highlighted in yellow. Coding sequences are shown in grey. **e)** The intersect and union of peaks with p < 0.05 from DESeq2, edgeR, and QNB from (b), coloured as in (b).

Using statistical methods to detect changes in peak enrichment, we found that the percent of

changes called below a p-value threshold of 0.05 were similar in the positive and negative controls

(**Figure 3b**). With all tools except MeTDiff, a knockout of Mettl3 showed the largest effects on m^6A (60),

while fewer significant peaks in other positive controls suggested variable effects of the positive control

conditions on $m^6A_{(m)}$, possibly related to efficiency of the methylation machinery knockdown or

overexpression (7,33,61–65). In the absence of true differences between groups, p-value distributions

should be uniform for well-calibrated statistical tests, meaning that ~5% of peaks should have p-values <

0.05 for the negative controls. MeTDiff reported an excess number of sites with p-values below 0.05

(**Additional File 2: Supplementary Figure 3d**) and identified a higher percentage of sites as differentially

methylated in the mouse cortex negative control data set than in all but two positive controls (**Figure 3b**).

On the other hand, the generalized linear models (GLMs) and QNB showed uniform to conservatively

shifted p-value distributions, with differences between the mouse cortex and Huh7 data sets (**Additional**

File 2: Supplementary Figure 3d), suggesting that these tools detect fewer false positives.

To ensure significant peak changes detected by each of the tools reflected changes in IP

enrichment independent of differential gene expression, we measured the correlation between changes in

IP read counts at peak sites and changes in input read counts across their encompassing genes. For

significant peaks (FDR-adjusted p-value < 0.05) from the positive controls, correlation between $\log_2$ fold

change in peak IP and gene input read counts was low for the GLMs and QNB (Pearson's R = 0.10 to

0.22) but reached 0.55 (p = 5.8E-87) for MeTDiff (**Figure 3c**). The higher correlation for MeTDiff was

driven by peaks with proportional changes in IP and input levels, which suggests that MeTDiff often

detects differential expression of methylated genes rather than differential methylation. Therefore,

published studies that have used MeTDiff may actually be detecting differential expression and not

differential methylation (22,66). Indeed, plotting coverage for genes reported as differentially methylated

in one of these studies, with the y-axis scaled separately per condition, confirmed that changes in m^6A

identified by MeTDiff were proportional to changes in gene expression (**Figure 3d**) (22). Given these

results, QNB or the GLM implementations are better methods than MeTDiff to detect differential

methylation. Taking the intersect of significant peaks for the GLMs and QNB may help determine the

most probable sites of m^6A changes, while taking the union of predictions provides a less conservative

approach to selecting sites for further validation (**Figure 3e**). However, additional filters are needed for

robust peak change detection as there were still significant peaks for which the difference between peak

$\log_2$ fold change and gene $\log_2$ fold change was close to zero, particularly with QNB (**Additional File 2:**

Supplementary Figure 3e). For microarray and RNA-seq data, a filter of absolute $\log_2$ fold change > 1

has been recommended to reduce false positive rates (67); in the remainder of our analyses, we

implemented a similar filter for absolute difference in peak and gene $\log_2$ fold change ≥ 1 to the combined

predictions from QNB and the two GLMs, with an additional filter where noted for peak read counts ≥ 10

across all replicates and conditions to ensure sufficient coverage for consistent peak detection (as

discussed in Figure 1a).

Reanalyzing peak changes between conditions

We next estimated the scale of statistically detectable peak changes under various conditions

using our approaches and compared these results to previously reported estimates of these changes

(**Figure 4a**, **Additional File 1: Supplementary Table 4**). We identified fewer peaks as differentially

methylated than originally reported under most conditions, with zero to hundreds of peaks significantly

changed (depending on experiment and method), versus hundreds to over ten thousand described in

publications (22–26,34,63,66,68–71). Notably, knockdown of Zc3h13 did appear to disrupt m^{6}A$_{(m)}$,

suggesting the gene does participate in methylation as recently described (69). Another study reported

that activin treatment of human pluripotent stem cells led to differential methylation of genes that encode

pluripotency factors (22). However, our reanalysis only found a few peak changes that passed our filters

for significance, fold change, and expression (minimum input read count across peaks ≥ 10), and no

enrichment for pluripotency factors among affected genes. Even when we removed the thresholds for fold

change and expression, the adjusted p-value for "signaling pathways regulating pluripotency of stem

cells" was still 0.15 and driven by only three genes, LEFTY2, FZD28, and FGFR3 (**Additional File 2:**

Supplementary Figure 4a). Interestingly, the minimum read threshold made a particularly dramatic

difference in the case of a recent study that looked at the effects of knocking down the histone

methyltransferase SETD2 on m^{6}A in mRNA. For this data, of the 2065 sites predicted by QNB, 2064 fell

below the minimum read threshold due to low input coverage in the first and second replicates (**Figure**

4a, **Additional File 2: Supplementary Figure 4b-e**) (70). We could not compare our approach to results

reported by Su et al. (2018), who found 6,024 peaks changed with R2HG treatment, Zeng et al. (2018),

who found 465-599 peaks changed between tumour samples, or Ma et al. (2018), who found 12,452

peaks were gained and 11,192 lost between P7 and P20 mouse cerebella, as each relied on a single

sample per condition, with no replicates (40–42).

Figure 4: Changes in peaks between conditions. a) Detected m6A(m) changes in thirteen published data sets that measured m6A(m) peak changes between two conditions **(Additional File 1: Supplementary Table 4)**. The number of peaks detected as changed in the original published analyses are compared to the number of peaks with FDR-adjusted p-values < 0.05 in our reanalysis using DESeq2, edgeR, or QNB, and taking the union of results from these three tools with additional filters for log2 fold difference in peak and gene changes of ≥1 and peak read counts ≥10 across all replicates and conditions ("filtered"). **b)** Gene coverage plots for *Hspa1a* in mouse embryonic fibroblasts (MEFs) and *HSPA1A* in human cells (HepG2 and BCL) before and after heat shock. Input coverage is shown in black and IP coverage in raspberry, with putative m6A peaks changed highlighted in yellow and marked by arrows. miCLIP coverage for an experiment in HepG2 cells is shown in orange. **c)** Coverage plots for *PSIP1*, which was reported to have a change in 5′ UTR m6A with HIV infection by Lichinchi et al (2016). **d)** Coverage plots for *SUN2*, in which we detected changes in m6A with HCMV infection and dsDNA treatment suggesting a possible increase in methylation under higher interferon conditions (after 12 h of dsDNA treatment compared to untreated controls and after 6 h post-HCMV infection compared to 72 h, when interferon levels have declined). Lines in coverage plots (**b-d**) show the mean across all replicates for each experiment, while shading shows the standard deviation. Coding sequences are shown in grey.

Multiple studies have investigated m6A(m) in the context of heat shock, HIV infection, KSHV

infection, and dsDNA treatment or human cytomegalovirus (HCMV) infection (**Additional File 1:**

Supplementary Table 5). Since each step in MeRIP-seq analysis risks introducing false negatives, we

cannot rule out consistent changes between studies that used similar experimental interventions based

on statistical detection alone. Therefore, we started by plotting coverage for specific genes reported as

differentially methylated to evaluate reproducibility across these studies. Zhou, et al. (2015) reported 5′

UTR methylation of *Hspa1a* with heat shock (20). Coverage was too low for untreated controls to

determine if *Hspa1a* was simply newly expressed or was actually newly methylated with heat shock

based on our alignment of their data using STAR (72). We were also unable to detect a change in

methylation of *HSPA1A* using data from other heat shock studies, including a new data set from a B-cell

lymphoma cell line and a published miCLIP data set, although coverage was again low (**Figure 4b**)

(4,73). Lichinchi, et al. (2016) reported that 56 genes showed increased methylation with HIV infection in

MT4 T-cells, with enrichment for genes involved in viral gene expression (25). Specific genes, for

example *PSIP1*, in which we also detected a peak using MACS2 and see a change in the peak when

plotting coverage using the data from Lichinchi et al. (2016), did not show the same changes in data from

two other CD4+ cell types, primary CD4+ cells and Jurkat cells (**Figure 4c**) (74). Two other studies both

used MeRIP-seq to establish the presence of m^{6}A in *IFNB1* induced through dsDNA treatment or by

infection with the dsDNA virus HCMV (75,76). While these studies did not discuss changes in m^{6}A, we

used these data sets to examine the replicability of m^{6}A$_{(m)}$ changes in response to dsDNA sensing and

interferon induction. Although different dsDNA stimuli, time points, and use of a fibroblast cell line versus

primary foreskin fibroblasts make it difficult to compare between the two experiments, using QNB and the

GLM approaches, we found four peaks in three genes (*AKAP8*, *SUN2*, and *TMEM140*) that showed

significant changes with higher interferon (**Figure 4d**). Overall, we were unable to detect the same

changes in m^{6}A$_{(m)}$ across studies of heat shock or HIV, and we detected only a few common changes in

the response to dsDNA. However, we do note that cell line-specific differences in m^{6}A$_{(m)}$ regulation and

differences in experimental protocols could account for some of the variability among these studies.

321   We did not have MeRIP-seq data for two studies from exactly the same conditions and cell lines

to compare, but two studies both used cell lines derived from iSLK to study the effects of KSHV on host

m^{6}A (27,28). Both suggested that KSHV infection could decrease the number of m^{6}A sites in host

transcripts. Hesser et al. (2018) found that lytic KSHV infection decreased the number of peaks on host

transcripts by >25%; Tan et al. (2018) suggested a loss of 17-59% of peaks in two different cell types, but

that m^{6}A$_{(m)}$ peak fold enrichment showed better clustering by cell type than by infection status. Neither of

these studies discussed specific genes that showed differential methylation with lytic infection. For our

comparison of $m^6A_{(m)}$ peak changes in these data sets, we identified probable changes in peaks based on

statistical significance using QNB or the GLMs with $\log_2$ fold change difference between peaks and genes

of ≥ 1. We detected 80 peak changes in the data from Hesser et al. (2018) and 18 in the data from Tan et

al. (2018) but found no peaks that changed in both iSLK data sets with lytic KSHV infection. Applying the

same statistical approaches, we were likewise unable to detect any shared peak changes between the

studies of HIV infection, and there were insufficient replicates to compare heat shock studies

(16,20,25,73,74). Thus, in our reanalysis of m^6A changes in response to stimuli, we detected only four

statistically reproducible peak changes, all in response to dsDNA.

Disparities between experiments were not simply due to significance thresholding or differences

in peak detection. Taking the union of peaks called in two experiments for KSHV, HIV, and dsDNA

treatment, we found minimal to negative correlations in changes in m^6A enrichment induced by treatment

at the same sites, further showing that changes with similar treatments are not reproducible (**Additional**

File 2: Supplementary Figure 4e).

MeRIP-RT-qPCR validation

Although statistical approaches revealed fewer changes in $m^6A_{(m)}$ with various stimuli than

published estimates, and we were unable to confirm changes in $m^6A_{(m)}$ methylation of specific genes

across studies of similar conditions, many of the studies we looked at do include additional validation of

$m^6A_{(m)}$ changes from MeRIP-seq using MeRIP-RT-qPCR. Recently it was shown that MeRIP-RT-qPCR

can capture differences in m^6A:A ratios at specific sites (34), but it is unknown how MeRIP-RT-qPCR is

affected by changes in gene expression. To test this, we ran MeRIP-RT-qPCR on in vitro transcribed

RNA oligonucleotides that lacked or contained m^6A spiked into total RNA extracted from Huh7 cells

(**Additional File 1: Supplementary Table 6**). We found that MeRIP-RT-qPCR detected the direction of

change in m^6A levels at different concentrations of spike-in RNAs (**Figure 5a-b**). However, technical

variation could also lead to spuriously significant differences. For example, a comparison of m^6A

enrichment between two dilutions (0.1 fmol and 10 fmol) of a 30% methylated spike-in mixture returned a

p-value of 0.004 (unpaired Student's *t*-test).

Figure 5: MeRIP-RT-qPCR validation and replicates necessary for the detection of peak changes.
a) Relative enrichment of the indicated amounts of an in vitro transcribed standard containing unmodified
A or m^6A, as measured by MeRIP-RT-qPCR. Data are shown for two independent replicates of three
technical replicates each as IP enrichment over input relative to pulldown of a positive control spike-in,
with the 0.1 fmol (0.01 m^6A: 0.09 A) sample normalized to 1. Bars represent mean ± SEM of two
independent replicates. *** p ≤ 0.005 by unpaired Student's *t*-test. **b)** Linear regression of relative m^6A
enrichment from (a). Points and error bars mark mean ± SEM of two independent replicates. **c)** Change
in MeRIP-RT-qPCR vs. MeRIP-seq enrichment for peaks detected as significantly differentially expressed
with infection of Huh7 cells by dengue virus, Zika virus, and hepatitis C virus. **d)** Number of replicates of
infected vs. uninfected cells needed to detect the peaks in (c). Replicates were randomly subsampled 10
times to calculate the fraction of subsamples in which peaks were called as significant by the GLMs or
QNB. Boxes span the 1st to 3rd quartiles, with medians indicated. Whiskers show the minimum and
maximum points within ±1.5x the interquartile distance from the boxes. Results for each subsample of
replicates are shown as jittered points.

We next assessed the correlation between m6A enrichment observed using MeRIP-seq and

MeRIP-RT-qPCR using data from our recent work that identified 58 peak changes in m6A in Huh7 cells

following infection by four different viruses (50). For those experiments, we again selected peaks that

change based on results from the union of QNB and the GLM approaches. We found that the magnitude

of changes in common among viruses correlated between MeRIP-seq and MeRIP-RT-qPCR, both across

peaks (Pearson's R = 0.57, p = 3.7E-6) and within single peaks across viruses (13 out of 19 peaks

showed positive correlations, four of which had p-values < 0.05 with three data points) (**Figure 5c,**

Additional File 2: Supplementary Figure 5). Given the correlation we found between MeRIP-seq and

MeRIP-RT-qPCR, it is unclear why changes in IP over input sequencing reads were undetectable at the

peaks reported by Bertero et al. (2018) and Huang et al. (2019) but differences in peaks were

successfully validated using MeRIP-RT-qPCR (22,70). Based on these discrepancies, while MeRIP-RT-

qPCR can be used as an initial method of validation for predicted peak changes, additional methods are

necessary to confirm quantitative differences in m6A levels and to resolve points where the assays do not

agree.

We next used our peaks validated with MeRIP-RT-qPCR to estimate the number of replicates

necessary for detection of changes with either the GLM or QNB methods. Using a permutation test, we

downsampled infected and uninfected replicates and reran statistical detection of changes. We found that

approximately 6-9 replicates were necessary for consistent detection (in at least 50% of subsamples) of

most peak changes (**Figure 5d**). Schurch et al. (2016) and Conesa et al. (2016) produced similar

recommendations for basic RNA-seq studies, finding that 6-12 replicates were necessary to detect most

changes in gene expression and that changes of 1.25 were detectable 25% of the time with five

replicates, rising to 44% with ten replicates, respectively (36,77). While our findings broadly agree with

these recommendations for RNA-seq, they also suggest that almost all published MeRIP-seq studies to

date are underpowered.

Discussion

In the eight years since MeRIP-/m6A-seq was first published (16,17), many studies have used

these methods to examine the function of m6A, its distribution along mRNA transcripts, and how it might

be regulated under various conditions. While 35 out of 64 of the MeRIP- and miCLIP-seq papers we

surveyed **(Additional File 1: Supplementary Table 1)** refer to m6A as "dynamic", and, by contrast, only

two describe the modification as "static", the literature is unclear on what is meant by the word "dynamic".

There is mixed evidence as to whether m6A is reversible through demethylation by the proposed

demethylases FTO and ALKBH5 (71,78–80). Recent research using an endoribonuclease-based method

for m6A detection suggests that ALKBH5 has only a mild suppressive effect on m6A levels and FTO no

effect (54). Although m6A does not appear to change over the course of an mRNA's lifetime at steady-

state (3), whether it changes in response to a particular stimulus and at what point is less clear. Some

studies have suggested that m6A may be modulated through changes in methyltransferase and

demethylase expression, producing consistent directions of change across transcripts (8,23,34), through

alternative mechanisms involving microRNA, transcription factors, promoters, or histone marks

(21,22,66,70,81), or through indeterminate mechanisms (17,20,25–28,52). However, based on our

reanalysis of available MeRIP-seq data, there is still only meagre support for widespread changes in m6A

across the transcriptome independent of changes in the expression of methylation machinery (e.g.

increases or decreases in METTL3 expression).

In particular, replication of peaks and changes in peaks across studies is limited. As with other

RNA IP-based methods, MeRIP-seq data contains noise, owing to technical and biological variation (82).

In fact, while peak overlaps reach ~80% between replicates of the same study, they decrease to a

median of 45% between studies, most of which use 2-3 replicates each **(Figure 1)**. Given that the

detection of peaks is so variable and that peak heights differ among replicates, it is perhaps not surprising

that peak changes have yet to be reproduced between multiple studies of similar conditions. Indeed,

variability in MeRIP-seq could also mask differences in m6A regulation among cell types, which have

been described in mouse brains (34) and in cell lines exposed to KSHV (28). To distinguish biological and

technical variation, it will therefore be particularly important to test if multiple groups using the same cell

line and conditions can better reproduce changes in m6A.

Figure 6: Proposed approach to identify candidates for m6A(m) changes for further validation using MeRIP-seq data. We suggest predicting changes in m6A(m) using DESeq2, edgeR, and QNB, and have implemented the DEQ package in R to facilitate this.

Disparities in the methods used to detect changes in m6A(m) peaks also play a role in differing

conclusions among studies. Here, we analyzed four statistical methods to detect changes in peaks and

found that three of these methods showed uniform or conservatively shifted p-value distributions and

were able to identify changes in m6A(m) independent of changes in gene expression. We therefore

suggest that these statistical methods, in combination with filters for input levels in both conditions and

the difference in $\log_2$ fold change between peaks and genes, can be used to identify candidate m6A(m)

sites from MeRIP-seq data for further analysis and validation (**Figure 6**). Based on our results, while

MeTDiff works for peak detection, we do not recommend MeTDiff for peak change detection as it does

not control well for differences in gene expression **(Figure 3)**. Similar to others (33), we found that plotting

predicted m6A changes was invaluable and that appropriate scaling for gene coverage could reveal

changes proportional to gene expression. In addition, plotting the standard deviation in transcript

coverage can help assess typical variation in peak height among replicates. We note that both differential

methylation of a gene and methylation of a gene that is differentially expressed could be important, but

they should not be conflated when considering the role of m6A in transcript regulation.

The extent to which m6A changes on particular transcripts and whether it changes in binary

presence/absence or in degree is unclear. MeRIP-RT-qPCR could detect methylation differences in in

vitro transcribed RNA. Further, we found that these changes correlated with differences in MeRIP-seq

enrichment. However, neither MeRIP-seq nor MeRIP-RT-qPCR can reveal the precise fraction of

transcript copies modified by m6A. In general, antibody-based methods are subject to biases, including

from differences in binding efficiencies based on RNA structure and motif preferences (83). There is an

oft-cited but little-used method for quantification of m6A, site-specific cleavage and radioactive-labeling

followed by ligation-assisted extraction and thin-layer chromatography (SCARLET) (19). However, this

method can be challenging, works only for highly abundant transcripts, and is impractical for

transcriptome-wide analysis. A recently developed endoribonuclease-based, antibody-independent

approach for m6A detection is promising in terms of quantification of m6A, but its use is limited to a subset

of m6A sites within DRAC motifs ending in ACA (~16% of all sites) (44,45). So far, comparison to this data

suggests that antibody-based approaches may underestimate the number of m6A sites (54). Alternative

methods to detect m6A based using single-molecule sequencing (including direct RNA sequencing and

real-time cDNA synthesis) are under development and may offer ways to detect, quantify, and phase m6A

sites, but these have not yet been shown to accurately detect m6A across a cellular transcriptome (84–

86). For now, site-specific SCARLET is the only option to biochemically validate proposed changes in

m6A at most motifs.

Conclusions

Our work reveals the limits of MeRIP-seq reproducibility for the detection of m6A$_{(m)}$ and in

particular suggests caution when using MeRIP-seq for the detection of changes in m6A$_{(m)}$. To increase

confidence in predicted changes in m6A$_{(m)}$, we propose statistical approaches that account for differences

in gene expression between conditions and variability among replicates. These methods can be used to

gain insight into the regulation and function of m6A$_{(m)}$ and to predict specific sites for validation before the

development of high-throughput alternatives to MeRIP-seq, and similar strategies may be applicable to

other types of RNA sequencing assay.

Methods

New MeRIP-seq data

- Huh7 data

Total RNA was extracted from Huh7 cells using Trizol (Thermo-Fisher). mRNA was purified from 200 μg

total RNA using the Dynabeads mRNA purification kit (Thermo-Fisher) and concentrated by ethanol

precipitation. Purified mRNA was fragmented using the RNA Fragmentation Reagent (Thermo-Fisher) for

15 minutes followed by ethanol precipitation. Then, MeRIP was performed using EpiMark N6-

methyladenosine Enrichment kit (NEB). 25 μL Protein G Dynabeads (Thermo-Fisher) per sample were

washed three times in MeRIP buffer (150 mM NaCl, 10 mM Tris-HCl, pH 7.5, 0.1% NP-40) and incubated

with 1 μL anti-m6A antibody (NEB) for 2 hours at 4°C with rotation. After washing three times, anti-m6A

conjugated beads were incubated with purified mRNA with rotation at 4°C overnight in 300 μL MeRIP

buffer with 1 μL RNAse inhibitor (recombinant RNasein; Promega). Beads were then washed twice with

500 μL MeRIP buffer, twice with low salt wash buffer (50 mM NaCl, 10 mM Tris-HCl, pH 7.5, 0.1% NP-

40), twice with high salt wash buffer (500 mM NaCl, 10 mM Tris-HCl, pH 7.5, 0.1% NP-40), and once

again with MeRIP buffer. m6A-modified RNA was eluted twice in 100 μL MeRIP buffer containing 5mM

m6A salt (Santa Cruz Biotechnology) for 30 minutes at 4°C with rotation and concentrated by ethanol

precipitation. RNA-seq libraries were prepared from eluate and the 10% of RNA set aside as input using

the TruSeq mRNA library prep kit (Illumina) and checked for fragment length using the Agilent 2100

Bioanalyzer. Single-end 50 base pair reads were sequenced on an Illumina HiSeq 2500.

- Heat shock

Early passage OCI-Ly1 diffuse large B-cell lymphoma cells were grown in Iscove's modified Eagle

Medium (IMDM) with 10% fetal bovine serum (FBS). OCI-Ly1 cells were obtained from the Ontario

Cancer Institute and regularly tested for *Mycoplasma* contamination by PCR and identified by single

nucleotide polymorphism. Cells were maintained with 1% penicillin/streptomycin in a 37°C, 5% CO_2,

humidified incubator. In these growing conditions, heat shocked cells were exposed to 43 °C for 1 hour,

followed by 1 hour of recovery at 37°C while control cells were maintained at 37°C. Following treatment,

cells were processed at 4°C to obtain total cell lysates. Lysates were immunoprecipitated for $m^6A_{(m)}$ using

Synaptic Systems antibody (SYSY 202 003) following the protocol described in Meyer, et al (2012) and

sequenced on an Illumina HiSeq 2500 (16).

Read processing

Reads were trimmed using Trimmomatic (87) and aligned to the human genome (hg38) or the mouse

genome (mm10), as appropriate, using STAR, a splice-aware aligner for RNA-seq data (72). We used the

flag "--outFilterMultimapNmax 1" to keep only uniquely aligned reads. Scripts used for alignment are

provided with the rest of the analysis scripts at https://github.com/al-mcintyre/merip_reanalysis_scripts.

Peak detection and comparison

IP over input peaks were called using MACS2 callpeak using the parameters "--nomodel --extsize 100 (or,

if available, the approximate fragment size for a specific experiment to extend reads at their 3' ends to a

fixed length) --gsize 100e6 (the approximate size of mouse and human transcriptomes based on gencode

annotations)" (49). No filter for coverage was applied at the stage of peak detection. Transcript coverage

was estimated using Kallisto (88) with an index construct 31mers, except for the Schwartz et al (2014)

data set, where the reads were too short and an alternative index based on 29mers was constructed (33).

For **Figure 1b**, the full union of unique peaks was taken and the percent of that set detected in single

replicates calculated. Intersects between peaks that overlapped for transcripts with ≥10X mean coverage

in both samples were taken using bedtools (89) for **Figure 2**, allowing a generous minimum of 1

overlapping base. Heatmaps for peak overlaps were generated using the ComplexHeatmap package in R

(90). MeRIP-seq data sets in **Figure 2b** included those for human cell lines in **Figure 2a**, other data sets

from the same studies and any data sets that shared the same cell lines, and other data sets that looked

at multiple human cell types. We considered only data sets from baseline conditions in **Figure 2**

(untreated cells and knockdown controls).

Poisson and negative binomial fits

Reads aligned to peaks were counted using featureCounts from the Rsubread package (91). Poisson and

negative binomial models were fit to input and IP read counts at peaks using maximum likelihood

estimation. Simulated read counts were generated with Poisson or negative binomial distributions based

on estimated parameters from the sample, with 500 random generations per model. The log likelihood of

seeing read counts from the sample and the simulations given the model parameters was then calculated

and the mean taken across all peaks.

Peak change detection and generalized linear models

516   Generalized linear models to detect changes in IP coverage while controlling for differences in input

517   coverage were implemented based on a method previously applied to HITS-CLIP data (58). Full and

518   reduced models were constructed as follows:

$$\log \mu_{ij} = \beta_i^0 + \beta_i^{IP} X_j^{IP} + \beta_i^{STIM} X_j^{STIM} + \beta_i^{STIM:IP} X_j^{STIM:IP}$$

$$\log \mu_{ij} = \beta_i^0 + \beta_i^{IP} X_j^{IP} + \beta_i^{STIM} X_j^{STIM}$$

Where μ_{ij} is the expected read count for peak i in sample j, modelled as a negative binomial distribution,

$X_j^{IP} = 1$ for IP samples and 0 for input samples, and $X_j^{STIM} = 1$ for samples under the experimental

intervention and 0 for control samples.

Statistical significance was then assessed using a chi-squared test (df=1) for the difference in deviances

between the full and reduced models, with the null hypothesis that the interaction term ($\beta_i^{STIM:IP}$) for

differential antibody enrichment driven by the experimental intervention is zero. The likelihood ratio test

was implemented through DESeq2 (56) and edgeR (57), two programs developed for RNA-seq analysis

that differ in how they filter data and in how they estimate dispersions for negative binomial distributions.

Generalized linear models implemented through edgeR included a term for the normalized library size of

sample j.

QNB was run as suggested for experiments with biological replicates, where each IP and input variable

("ip1", etc.) consisted of a matrix of peak counts for either condition 1 or condition 2:

> qnbtest(ip1, ip2, input1, input2, mode="per-condition")

We extracted functions from MeTDiff so that we could supply our own peaks and thus control for

differences in peak detection among tools. The main post-peak calling function, diff.call.module, was run

as follows using the same count matrices as for QNB:

> diff.call.module(ip1, input1, ip2, input2)

Gene and peak expression changes were estimated as $\log_2$ fold changes from DESeq2 based on

differences in input read counts aligned to genes and IP read counts aligned to peaks, respectively, and

the change in peak relative to gene enrichment was calculated as the absolute difference in $\log_2$ fold

change between those values.

Comparison to published studies

The sources for published estimates of m^6A peak changes included in our comparison are listed in

Additional File 1: Supplementary Table 4. Significant (FDR-adjusted $p < 0.05$) peaks were considered

for DESeq2, edgeR, and QNB, run as described above. We also considered a filtered set of peaks

derived from the union of significant peaks from the three tools with additional filters for location within

exons, |$\log_2$ fold change between peak IP and gene input| ≥ 1, and a minimum peak read count of 10

across replicates and conditions. We used gProfiler to calculate enrichment of functional categories (92).

In **Figure 4b-c**, we selected *Hspa1a/HSPA1A* as our representative gene for heat shock because it was

the primary example cited by Zhou et al. (2015) and Meyer et al. (2015) (4,20). For HIV, we selected

PSIP1 because it was among the 56 genes reported by Lichinchi et al. (2016a) (25), it plays a known role

in HIV infection, and we detected a peak in the gene using MACS2.

For KSHV, we compared significant results (adjusted p < 0.05) from QNB and GLMs (DESeq2 and

edgeR), with additional filtering for |peak IP – gene input $\log_2$ fold change| ≥ 1 (lowering this threshold to

0.5 did not change results), for data from Hesser et al. (2018) (27) in lytic vs. latent iSLK.219 cells and

data from Tan et al. (2018) (28) in lytic vs. latent iSLK BAC16 cells. We used the same approach to

compare data from Rubio et al. (2018) and Winkler et al. (2019) (75,76) for response to dsDNA. Data sets

used for site-specific comparisons are summarized in **Additional File 1: Supplementary Table 5**.

Gene coverage was plotted using CovFuzze (https://github.com/al-mcintyre/CovFuzze), which

summarizes mean and standard deviation in coverage across available replicates (93). Pearson's

correlations were taken for **Supplementary Figure 4** for peaks expressed above a minimum input peak

read count of 10 across replicates and conditions.

Spike-in controls and MeRIP-RT-qPCR

In vitro transcribed (IVT) controls were provided by the Jaffrey Lab and consisted of 1001 base long RNA

sequences with three adenines in GAC motifs (**Additional File 1: Supplementary Table 6**) either fully

methylated or unmethylated. m^6A and A controls were mixed in various ratios (1:9, 3:7, and 9:1) that

approximate the variation in m^6A levels detected by SCARLET (m^6A levels at specific sites have been

reported to vary from 6-80% of transcripts (19)). Modified and unmodified standards were mixed at the

indicated ratios to yield a final quantity of 0.1 fmol, 1 fmol, and 10 fmol. Mixed RNA standards were added

to 30 µg total RNA from Huh7 cells, along with 0.1 fmol of positive (m^6A-modified *Gaussia* luciferase

RNA, "GLuc") and negative control (unmodified *Cypridina* luciferase, "CLuc") spike-in RNA provided with

the N6-methyladenosine Enrichment kit (EpiMark). Following MeRIP as described above, cDNA was

synthesized from eluate and input samples using the iScript cDNA synthesis kit (Bio-Rad), and RT-qPCR

was performed on a QuantStudio Flex 6 instrument. Data was analyzed as a percent of input of the spike-

in RNA in each condition relative to that of the provided positive control spike-in.

For MeRIP-RT-qPCR to test peak callers, Huh7 cells plated in 6-well plates were transfected with

siRNAs against METTL3 and METTL14 (Qiagen; SI04317096 and SI00459942) or non-targeting control

siRNa (SI03650318) using Lipofectamine RNAiMax (Thermo Fisher) twice, 24 hours apart. 48 hours

following the second round of siRNA transfection, cells were harvested in TRIzol reagent and total RNA

was extracted. 30 μg total RNA was fragmented for 3 mins at 75°C, concentrated by ethanol precipitation,

and MeRIP-RT-qPCR was performed as described above. Primers used for RT-qPCR are provided in

Additional File 1: Supplementary Table 7 and siRNA sequences in **Additional File 1: Supplementary**

Table 8.

Cell culture and infection (data used for MeRIP-RT-qPCR comparisons)

Huh7 cells were grown in DMEM (Mediatech) supplemented with 10% fetal bovine serum (HyClone), 2.5

mM HEPES, and 1X non-essential amino acids (Thermo-Fisher). The identity of the Huh7 cell lines was

verified using the Promega GenePrint STR kit (DNA Analysis Facility, Duke University), and cells were

verified as mycoplasma free by the LookOut Mycoplasma PCR detection kit (Sigma). Infectious stocks of

a cell culture-adapted strain of genotype 2A JFH1 HCV were generated and titered on Huh7.5 cells by

focus-forming assay (FFA), as described (94). Dengue virus (DENV2-NGC), West Nile virus (WNV-

NY2000), and Zika virus (ZIKV-PRVABC59) viral stocks were generated in C6/36 cells and titered on

Vero cells as described (94). All viral infections were performed at a multiplicity of infection of 1 for 48

hours.

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of Data and Materials

MeRIP-seq data for the Huh7 negative controls is available in the GEO repository, under accession

number GSE130891. MeRIP-seq data for heat shock in B-cell lymphoma is available under accession

number GSE130892. Accession numbers for all other data sets reanalyzed in the study are included in

Additional File 1: Supplementary Tables 1-5. Scripts used for analysis are available at

https://github.com/al-mcintyre/merip_reanalysis_scripts and a pipeline implementing generalized linear

models through DESeq2 and edgeR, as well as QNB, is provided at https://github.com/al-mcintyre/deq.

Competing interests

619   C.E.M. is a cofounder and board member for Biotia and Onegevity Health, as well as an advisor or

620   compensated speaker for Abbvie, Acuamark Diagnostics, ArcBio, BioRad, DNA Genotek, Genialis,

621   Genpro, Karius, Illumina, New England Biolabs, QIAGEN, Whole Biome, and Zymo Research.

Funding

We are grateful for funding from the Starr Cancer Consortium (I9-A9-071), the Bert L and N. Kuggie Vallee

Foundation, the WorldQuant Foundation, The Pershing Square Sohn Cancer Research Alliance, NASA

(NNX14AH50G), the National Institutes of Health (R01AI125416, R21AI129851, R01MH117406), the

Leukemia and Lymphoma Society (LLS) grants (LLS 9238-16, Mak, LLS-MCL-982, Chen-Kiang), the

Burroughs Wellcome Fund (S.M.H.), the National Science and Engineering Research Council of Canada

(A.B.R.M. PGS-D funding), and the American Heart Association (N.S.G. Pre-doctoral Fellowship,

17PRE33670017).

Authors' contributions

A.B.R.M. and C.E.M. conceived the study. A.B.R.M. developed and ran the analyses and wrote the

manuscript with N.S.G. and S.M.H. S.R.J. provided in vitro controls for MeRIP-RT-qPCR. N.S.G.

prepared MeRIP-seq libraries for the Huh7 controls and ran MeRIP-RT-qPCR tests. L.C. contributed

additional heat shock data. All authors read and edited the manuscript.

Acknowledgements

We would like to thank Christina Leslie and Jonathan Victor for statistical advice, Aashiq Mirza for the

MeRIP-RT-qPCR controls, and Helen Lazear for WNV infection. We would also like to thank the

Epigenomics Core at Weill Cornell for preparing MeRIP-seq libraries, the Scientific Computing Unit

(SCU), and New England Biolabs for donating anti-m^6A antibodies.

References

1.   Balacco DL, Soller M. The m6A Writer: Rise of a Machine for Growing Tasks. Biochemistry. 2019;

2.   Ke S, Alemu EA, Mertens C, Gantman EC, Fak JJ, Mele A, et al. A majority of m6A residues are in
the last exons, allowing the potential for 3′ UTR regulation. Genes & development. 2015;

3.   Ke S, Pandya-Jones A, Saito Y, Fak JJ, Vågbø CB, Geula S, et al. m6A mRNA modifications are
deposited in nascent pre-mRNA and are not required for splicing but do specify cytoplasmic
turnover. Genes & Development. 2017;31(10):990–1006.

4.   Meyer KD, Patil DP, Zhou J, Zinoviev A, Skabkin MA, Elemento O, et al. 5′ UTR m6A promotes cap-
independent translation. Cell. 2015;163(4):999–1010.

5.   Wang X, Zhao BS, Roundtree IA, Lu Z, Han D, Ma H, et al. N6-methyladenosine modulates
messenger RNA translation efficiency. Cell. 2015;161(6):1388–99.

6.   Lin S, Choe J, Du P, Triboulet R, Gregory RI. The m6A methyltransferase METTL3 promotes
translation in human cancer cells. Molecular cell. 2016;62(3):335–45.

7.   Wang Y, Li Y, Toth JI, Petroski MD, Zhang Z, Zhao JC. N 6-methyladenosine modification
destabilizes developmental regulators in embryonic stem cells. Nature cell biology. 2014;16(2):191.

8.   Molinie B, Wang J, Lim KS, Hillebrand R, Lu Z, Van Wittenberghe N, et al. m 6 A-LAIC-seq reveals
the census and complexity of the m 6 A epitranscriptome. Nature methods. 2016;13(8):692.

9. Yoon K-J, Ringeling FR, Vissers C, Jacob F, Pokrass M, Jimenez-Cyrus D, et al. Temporal control of
mammalian cortical neurogenesis by m6A methylation. Cell. 2017;171(4):877–89.

10. Meyer KD, Jaffrey SR. Rethinking m6A readers, writers, and erasers. Annual review of cell and
developmental biology. 2017;33:319–42.

11. Roundtree IA, Evans ME, Pan T, He C. Dynamic RNA modifications in gene expression regulation.
Cell. 2017;169(7):1187–200.

12. Rosa-Mercado NA, Withers JB, Steitz JA. Settling the m6A debate: methylation of mature mRNA is
not dynamic but accelerates turnover. Genes & development. 2017;31(10):957–8.

13. Darnell RB, Ke S, Darnell JE. Pre-mRNA processing includes N6 methylation of adenosine residues
that are retained in mRNA exons and the fallacy of "RNA epigenetics." RNA. 2018;24(3):262–7.

14. Zhao BS, Nachtergaele S, Roundtree IA, He C. Our views of dynamic N6-methyladenosine RNA
methylation. RNA. 2018;24(3):268–72.

15. Mauer J, Jaffrey SR. FTO, m6Am, and the hypothesis of reversible epitranscriptomic mRNA
modifications. FEBS letters. 2018;592(12):2012–22.

16. Meyer KD, Saletore Y, Zumbo P, Elemento O, Mason CE, Jaffrey SR. Comprehensive analysis of
mRNA methylation reveals enrichment in 3′ UTRs and near stop codons. Cell. 2012;149(7):1635–
46.

17. Dominissini D, Moshitch-Moshkovitz S, Schwartz S, Salmon-Divon M, Ungar L, Osenberg S, et al.
Topology of the human and mouse m 6 A RNA methylomes revealed by m 6 A-seq. Nature.
2012;485(7397):201.

18. Linder B, Grozhik AV, Olarerin-George AO, Meydan C, Mason CE, Jaffrey SR. Single-nucleotide-
resolution mapping of m6A and m6Am throughout the transcriptome. Nature methods.
2015;12(8):767.

19. Liu N, Parisien M, Dai Q, Zheng G, He C, Pan T. Probing N6-methyladenosine RNA modification
status at single nucleotide resolution in mRNA and long noncoding RNA. Rna. 2013;

20. Zhou J, Wan J, Gao X, Zhang X, Jaffrey SR, Qian S-B. Dynamic m 6 A mRNA methylation directs
translational control of heat shock response. Nature. 2015;526(7574):591.

21. Chen T, Hao Y-J, Zhang Y, Li M-M, Wang M, Han W, et al. m6A RNA Methylation Is Regulated by
MicroRNAs and Promotes Reprogramming to Pluripotency. Cell Stem Cell. 2015 Mar 5;16(3):289–
301.

22. Bertero A, Brown S, Madrigal P, Osnato A, Ortmann D, Yiangou L, et al. The SMAD2/3 interactome
reveals that TGFβ controls m 6 A mRNA methylation in pluripotency. Nature. 2018;555(7695):256.

23. Liu J, Eckert MA, Harada BT, Liu S-M, Lu Z, Yu K, et al. m 6 A mRNA methylation regulates AKT
activity to promote the proliferation and tumorigenicity of endometrial cancer. Nature cell biology.
2018;20(9):1074.

24. Anders M, Chelysheva I, Goebel I, Trenkner T, Zhou J, Mao Y, et al. Dynamic m6A methylation
facilitates mRNA triaging to stress granules. Life science alliance. 2018;1(4):e201800113.

25. Lichinchi G, Gao S, Saletore Y, Gonzalez GM, Bansal V, Wang Y, et al. Dynamics of the human and
viral m6A RNA methylomes during HIV-1 infection of T cells. Nature Microbiology. 2016;1.

26. Lichinchi G, Zhao BS, Wu Y, Lu Z, Qin Y, He C, et al. Dynamics of human and viral RNA methylation
during Zika virus infection. Cell host & microbe. 2016;20(5):666–73.

27. Hesser CR, Karijolich J, Dominissini D, He C, Glaunsinger BA. N6-methyladenosine modification and
the YTHDF2 reader protein play cell type specific roles in lytic viral gene expression during Kaposi's
sarcoma-associated herpesvirus infection. PLoS pathogens. 2018;14(4):e1006995.

28. Tan B, Liu H, Zhang S, da Silva SR, Zhang L, Meng J, et al. Viral and cellular N 6-methyladenosine
and N 6, 2′-O-dimethyladenosine epitranscriptomes in the KSHV life cycle. Nature microbiology.
2018;3(1):108.

29. Vu LP, Pickering BF, Cheng Y, Zaccara S, Nguyen D, Minuesa G, et al. The N 6-methyladenosine (m
6 A)-forming enzyme METTL3 controls myeloid differentiation of normal hematopoietic and
leukemia cells. Nature medicine. 2017;23(11):1369.

30. Liu L, Zhang S-W, Huang Y, Meng J. QNB: differential RNA methylation analysis for count-based
small-sample sequencing data with a quad-negative binomial model. BMC bioinformatics.
2017;18(1):387.

31. Cui X, Zhang L, Meng J, Rao MK, Chen Y, Huang Y. MeTDiff: a novel differential RNA methylation
analysis for MeRIP-seq data. IEEE/ACM Transactions on Computational Biology and Bioinformatics
(TCBB). 2018;15(2):526–34.

32. Zhao X, Yang Y, Sun B-F, Shi Y, Yang X, Xiao W, et al. FTO-dependent demethylation of N6-
methyladenosine regulates mRNA splicing and is required for adipogenesis. Cell research.
2014;24(12):1403.

33. Schwartz S, Mumbach MR, Jovanovic M, Wang T, Maciag K, Bushkin GG, et al. Perturbation of m6A
writers reveals two distinct classes of mRNA methylation at internal and 5′ sites. Cell reports.
2014;8(1):284–96.

34. Engel M, Eggert C, Kaplick PM, Eder M, Röh S, Tietze L, et al. The role of m6A/m-RNA methylation
in stress response regulation. Neuron. 2018;99(2):389–403.

35. Zhou J, White KP, Liu Y. RNA-seq differential expression studies: more sequence or more
replication? Bioinformatics. 2013 Dec 4;30(3):301–4.

36. Schurch NJ, Schofield P, Gierliński M, Cole C, Sherstnev A, Singh V, et al. How many biological
replicates are needed in an RNA-seq experiment and which differential expression tool should you
use? Rna. 2016;22(6):839–51.

37. Su Z, Łabaj PP, Li S, Thierry-Mieg J, Thierry-Mieg D, Shi W, et al. A comprehensive assessment of
RNA-seq accuracy, reproducibility and information content by the Sequencing Quality Control
Consortium. Nature biotechnology. 2014;32(9):903.

38. Geula S, Moshitch-Moshkovitz S, Dominissini D, Mansour AA, Kol N, Salmon-Divon M, et al. m6A
mRNA methylation facilitates resolution of naïve pluripotency toward differentiation. Science.
2015;347(6225):1002–6.

39. Cui Q, Shi H, Ye P, Li L, Qu Q, Sun G, et al. m6A RNA methylation regulates the self-renewal and
tumorigenesis of glioblastoma stem cells. Cell reports. 2017;18(11):2622–34.

40. Ma C, Chang M, Lv H, Zhang Z-W, Zhang W, He X, et al. RNA m 6 A methylation participates in
regulation of postnatal development of the mouse cerebellum. Genome biology. 2018;19(1):68.

41. Su R, Dong L, Li C, Nachtergaele S, Wunderlich M, Qing Y, et al. R-2HG exhibits anti-tumor activity
by targeting FTO/m6A/MYC/CEBPA signaling. Cell. 2018;172(1–2):90–105.

42. Zeng Y, Wang S, Gao S, Soares F, Ahmed M, Guo H, et al. Refined RIP-seq protocol for
epitranscriptome analysis with low input materials. PLoS biology. 2018;16(9):e2006092.

43. Zhou J, Wan J, Shu XE, Mao Y, Liu X-M, Yuan X, et al. N6-methyladenosine guides mRNA
alternative translation during integrated stress response. Molecular cell. 2018;69(4):636–47.

44. Garcia-Campos MA, Edelheit S, Toth U, Shachar R, Nir R, Lasman L, et al. Deciphering the'm6A
code'via quantitative profiling of m6A at single-nucleotide resolution. BioRxiv. 2019;571679.

45. Zhang Z, Chen L-Q, Zhao Y-L, Yang C-G, Roundtree IA, Zhang Z, et al. Single-base mapping of m6A
by an antibody-independent method. bioRxiv. 2019;575555.

46. Liu H, Begik O, Lucas MC, Mason CE, Schwartz S, Mattick JS, et al. Accurate detection of m6A RNA
modifications in native RNA sequences. bioRxiv. 2019;525741.

47. Meng J, Lu Z, Liu H, Zhang L, Zhang S, Chen Y, et al. A protocol for RNA methylation differential
analysis with MeRIP-Seq data and exomePeak R/Bioconductor package. Methods.
2014;69(3):274–81.

48. Cui X, Meng J, Zhang S, Chen Y, Huang Y. A novel algorithm for calling mRNA m 6 A peaks by
modeling biological variances in MeRIP-seq data. Bioinformatics. 2016;32(12):i378–85.

49. Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, et al. Model-based analysis of
ChIP-Seq (MACS). Genome biology. 2008;9(9):R137.

50. Gokhale NS, McIntyre ABR, Mattocks MD, Holley CL, Lazear HM, Mason CE, et al. Altered m6A
modification of specific cellular transcripts affects Flaviviridae infection. bioRxiv. 2019 Jan 1;670984.

51. Antanaviciute A, Baquero-Perez B, Watson CM, Harrison SM, Lascelles C, Crinnion L, et al.
m6aViewer: software for the detection, analysis, and visualization of N6-methyladenosine peaks
from m6A-seq/ME-RIP sequencing data. RNA. 2017;23(10):1493–501.

52. He S, Wang H, Liu R, He M, Che T, Jin L, et al. mRNA N6-methyladenosine methylation of postnatal
liver development in pig. PloS one. 2017;12(3):e0173421.

53. Tao X, Chen J, Jiang Y, Wei Y, Chen Y, Xu H, et al. Transcriptome-wide N 6-methyladenosine
methylome profiling of porcine muscle and adipose tissues reveals a potential mechanism for
transcriptional regulation and differential methylation pattern. BMC genomics. 2017;18(1):336.

54. Garcia-Campos MA, Edelheit S, Toth U, Safra M, Shachar R, Viukov S, et al. Deciphering the "m6A
Code" via Antibody-Independent Quantitative Profiling. Cell. 2019 Jul 25;178(3):731-747.e16.

55. Xiao S, Cao S, Huang Q, Xia L, Deng M, Yang M, et al. The RNA N6-methyladenosine modification
landscape of human fetal tissues. Nature Cell Biology. 2019 May 1;21(5):651–61.

56. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data
with DESeq2. Genome biology. 2014;15(12):550.

57. Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential expression
analysis of digital gene expression data. Bioinformatics. 2010;26(1):139–40.

58. Park S-M, Deering RP, Lu Y, Tivnan P, Lianoglou S, Al-Shahrour F, et al. Musashi-2 controls cell
fate, lineage bias, and TGF-β signaling in HSCs. Journal of Experimental Medicine.
2014;211(1):71–87.

59. Marioni JC, Mason CE, Mane SM, Stephens M, Gilad Y. RNA-seq: an assessment of technical
reproducibility and comparison with gene expression arrays. Genome research. 2008;18(9):1509–
17.

60. Batista PJ, Molinie B, Wang J, Qu K, Zhang J, Li L, et al. m6A RNA modification controls cell fate
transition in mammalian embryonic stem cells. Cell stem cell. 2014;15(6):707–19.

61. Fustin J-M, Doi M, Yamaguchi Y, Hida H, Nishimura S, Yoshida M, et al. RNA-methylation-dependent
RNA processing controls the speed of the circadian clock. Cell. 2013;155(4):793–806.

62. Hess ME, Hess S, Meyer KD, Verhagen LA, Koch L, Brönneke HS, et al. The fat mass and obesity
associated gene (Fto) regulates activity of the dopaminergic midbrain circuitry. Nature
neuroscience. 2013;16(8):1042.

63. Li Z, Weng H, Su R, Weng X, Zuo Z, Li C, et al. FTO plays an oncogenic role in acute myeloid
leukemia as a N6-methyladenosine RNA demethylase. Cancer cell. 2017;31(1):127–41.

64. Huang H, Weng H, Sun W, Qin X, Shi H, Wu H, et al. Recognition of RNA N 6-methyladenosine by
IGF2BP proteins enhances mRNA stability and translation. Nature cell biology. 2018;20(3):285.

65. Zhong X, Yu J, Frazier K, Weng X, Li Y, Cham CM, et al. Circadian Clock Regulation of Hepatic Lipid
Metabolism by Modulation of m6A mRNA Methylation. Cell reports. 2018;25(7):1816–28.

66. Barbieri I, Tzelepis K, Pandolfini L, Shi J, Millán-Zambrano G, Robson SC, et al. Promoter-bound
METTL3 maintains myeloid leukaemia by m 6 A-dependent translation control. Nature.
2017;552(7683):126.

67. SEQC/MAQC-III Consortium, Su Z, Łabaj PP, Li S, Thierry-Mieg J, Thierry-Mieg D, et al. A
comprehensive assessment of RNA-seq accuracy, reproducibility and information content by the
Sequencing Quality Control Consortium. Nature Biotechnology. 2014 Aug 24;32:903.

68. Li M, Zhao X, Wang W, Shi H, Pan Q, Lu Z, et al. Ythdf2-mediated m 6 A mRNA clearance
modulates neural development in mice. Genome biology. 2018;19(1):69.

69. Wen J, Lv R, Ma H, Shen H, He C, Wang J, et al. Zc3h13 regulates nuclear RNA m 6 A methylation
and mouse embryonic stem cell self-renewal. Molecular cell. 2018;69(6):1028–38.

70. Huang H, Weng H, Zhou K, Wu T, Zhao BS, Sun M, et al. Histone H3 trimethylation at lysine 36
guides m 6 A RNA modification co-transcriptionally. Nature. 2019;1.

71. Zheng G, Dahl JA, Niu Y, Fedorcsak P, Huang C-M, Li CJ, et al. ALKBH5 is a mammalian RNA
demethylase that impacts RNA metabolism and mouse fertility. Molecular cell. 2013;49(1):18–29.

72. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast universal RNA-
seq aligner. Bioinformatics. 2013;29(1):15–21.

73. Dominissini D, Moshitch-Moshkovitz S, Schwartz S, Salmon-Divon M, Ungar L, Osenberg S, et al.
Topology of the human and mouse m 6 A RNA methylomes revealed by m 6 A-seq. Nature.
2012;485(7397):201.

74. Tirumuru N, Zhao BS, Lu W, Lu Z, He C, Wu L. N6-methyladenosine of HIV-1 RNA regulates viral
infection and HIV-1 Gag protein expression. Elife. 2016;5:e15528.

75. Rubio RM, Depledge DP, Bianco C, Thompson L, Mohr I. RNA m6 A modification enzymes shape
innate responses to DNA by regulating interferon β. Genes & development. 2018;32(23–24):1472–
84.

76. Winkler R, Gillis E, Lasman L, Safra M, Geula S, Soyris C, et al. m 6 A modification controls the
innate immune response to infection by targeting type I interferons. Nature immunology.
2019;20(2):173.

77. Conesa A, Madrigal P, Tarazona S, Gomez-Cabrero D, Cervera A, McPherson A, et al. A survey of
best practices for RNA-seq data analysis. Genome Biology. 2016 Jan 26;17(1):13.

78. Jia G, Fu Y, Zhao X, Dai Q, Zheng G, Yang Y, et al. N6-methyladenosine in nuclear RNA is a major
substrate of the obesity-associated FTO. Nature chemical biology. 2011;7(12):885.

79. Wei J, Liu F, Lu Z, Fei Q, Ai Y, He PC, et al. Differential m6A, m6Am, and m1A demethylation
mediated by FTO in the cell nucleus and cytoplasm. Molecular cell. 2018;71(6):973–85.

80. Mauer J, Luo X, Blanjoie A, Jiao X, Grozhik AV, Patil DP, et al. Reversible methylation of m 6 A m in
the 5′ cap controls mRNA stability. Nature. 2017;541(7637):371.

81. Aguilo F, Zhang F, Sancho A, Fidalgo M, Di Cecilia S, Vashisht A, et al. Coordination of m6A mRNA
methylation and gene transcription by ZFP217 regulates pluripotency and reprogramming. Cell stem
cell. 2015;17(6):689–704.

82. Chakrabarti AM, Haberman N, Praznik A, Luscombe NM, Ule J. Data science issues in studying
Protein–RNA interactions with CLIP technologies. Annual Review of Biomedical Data Science.
2018;1:235–61.

83. Liu B, Merriman DK, Choi SH, Schumacher MA, Plangger R, Kreutz C, et al. A potentially abundant
junctional RNA motif stabilized by m6A and Mg2+. Nature Communications. 2018 Jul 17;9(1):2761.

84. Saletore Y, Meyer K, Korlach J, Vilfan ID, Jaffrey S, Mason CE. The birth of the Epitranscriptome:
deciphering the function of RNA modifications. Genome Biology. 2012;13(10):175.

85. Garalde DR, Snell EA, Jachimowicz D, Sipos B, Lloyd JH, Bruce M, et al. Highly parallel direct RNA
sequencing on an array of nanopores. Nature Methods. 2018 Jan 15;15:201.

86. Liu H, Begik O, Lucas MC, Ramirez JM, Mason CE, Wiener D, et al. Accurate detection of m6A RNA
modifications in native RNA sequences. Nature Communications. 2019 Sep 9;10(1):4079.

87. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data.
Bioinformatics. 2014;30(15):2114–20.

88. Bray NL, Pimentel H, Melsted P, Pachter L. Near-optimal probabilistic RNA-seq quantification. Nature
biotechnology. 2016;34(5):525.

89. Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic features.
Bioinformatics. 2010;26(6):841–2.

90. Gu Z, Eils R, Schlesner M. Complex heatmaps reveal patterns and correlations in multidimensional
genomic data. Bioinformatics. 2016;32(18):2847–9.

91. Liao Y, Smyth GK, Shi W. The R package Rsubread is easier, faster, cheaper and better for
alignment and quantification of RNA sequencing reads. Nucleic Acids Research. 2019 Feb
20;47(8):e47–e47.

92. Reimand J, Arak T, Adler P, Kolberg L, Reisberg S, Peterson H, et al. g: Profiler—a web server for
functional interpretation of gene lists (2016 update). Nucleic acids research. 2016;44(W1):W83–9.

93. Imam H, Khan M, Gokhale NS, McIntyre AB, Kim G-W, Jang JY, et al. N6-methyladenosine
modification of hepatitis B virus RNA differentially regulates the viral life cycle. Proceedings of the
National Academy of Sciences. 2018;115(35):8829–34.

94. Gokhale NS, McIntyre ABR, McFadden MJ, Roder AE, Kennedy EM, Gandara JA, et al. N6-
methyladenosine in Flaviviridae viral RNA genomes regulates infection. Cell host & microbe. 2016
Nov;

872 **Additional file 1:** Supplementary Tables. (xls)

873 **Additional file 2:** Supplementary Figures. (pdf)

874 **Additional file 3**: Results with MeTDiff peak caller. (pdf)

875 **Additional file 4**: Peaks and statistical test results for experiments in Figures 3-4. (zip)

876 **Additional file 5**: Response to reviewers. (txt)