**A live-cell imaging system for visualizing the transport of Marburg virus nucleocapsid-like structures**

Yuki Takamatsu[1, 3], Takeshi Noda[3,4*] and Stephan Becker[1,2*]

[1]Institut für Virologie, Philipps-Universität Marburg, Germany;

[2]German Center of Infection Research (DZIF), partner site Giessen-Marburg-Langen, Marburg, Germany;

[3]Laboratory of Ultrastructural Virology, Institute for Frontier Life and Medical Sciences, Kyoto University, Japan

[4]Laboratory of Ultrastructural Virology, Graduate School of Biostudies, Kyoto University, Japan

*Corresponding authors:

Takeshi Noda

Laboratory of Ultrastructural Virology

Institute for Frontier Life and Medical Sciences

Kyoto University

Japan

Tel.: +49 6421/28-66254

Fax.: +49 6421/28-68962

Email: t-noda@kyoto-uni.ac.jp

Stephan Becker

Institute of Virology

Philipps-University Marburg

Hans-Meerwein-Straße

35043 Marburg

Germany

Tel.: +49 6421/28-66254

Fax.: +49 6421/28-68962

Email: becker@staff.uni-marburg.de

**Abstract**

Live-cell imaging is a powerful tool for visualization of the spatio-temporal dynamics of living organisms. Although this technique is utilized to visualize nucleocapsid transport in Marburg virus (MARV)- or Ebola virus-infected cells, the experiments require biosafety level-4 (BSL-4) laboratories, which are restricted to trained and authorized individuals. To overcome this limitation, we developed a live-cell imaging system to visualize MARV nucleocapsid-like structures using fluorescence-conjugated viral proteins, which can be conducted outside BSL-4 laboratories. Our experiments revealed that nucleocapsid-like structures have similar transport characteristics to nucleocapsids observed in MARV-infected cells. This system provides a safe platform to evaluate antiviral drugs that inhibit MARV nucleocapsid transport.

**Introduction**

Marburg virus (MARV), together with the Ebola virus (EBOV), belongs to

the family *Filoviridae*, and has a roughly 19 kb non-segmented, single-stranded,

negative-sense RNA genome. It causes severe hemorrhagic fever with high

fatality rates. MARV epidemics have occasionally been reported in Central Africa,

with the largest one, having a 90% fatality rate, being reported in Angola between

2004 and 2005 (CDC, 2014). Currently, there are no approved vaccines or

antiviral therapeutics available to prevent or treat MARV infection. Therefore,

understanding the interplay between viral and host proteins during MARV

replication is necessary to establish countermeasures for the diseases. For

example, revealing the mechanisms for the assembly and transport of

nucleocapsids, which comprise viral genomic RNA, nucleoprotein (NP), viral

proteins (VP24, VP30, VP35), and polymerase L, and are responsible for the

transcription and replication of the viral genome, might contribute to the

development of new therapeutic options.

The main nucleocapsid protein of MARV is NP, which is responsible for

the encapsidation of single-stranded viral genomic RNA (Bharat *et al.*, 2012,

Muhlberger *et al.*, 1999). In addition to NP, MARV nucleocapsids also contain the

minor matrix protein VP24 and the polymerase cofactor VP35, both of which are

essential structural elements that directly interact with NP to build a helical

nucleocapsid approximately 1000 nm in length and 50 nm in diameter (Huang *et al.*, 2002, Watanabe *et al.*, 2006, Bharat et al., 2012). Furthermore, the viral

polymerase L and the transcription factor VP30 are also associated with the

nucleocapsid (Hartlieb *et al.*, 2003, Biedenkopf *et al.*, 2013). The core complex

of the nucleocapsid, formed by the NP, VP35, and VP24 proteins together with

the viral RNA, is defined as a nucleocapsid-like structure (NCLS, Fig. 1a). In

MARV- and EBOV-infected cells, immunofluorescence microscopy, as well as

live-cell imaging, have been used to visualize and analyze nucleocapsids by

fluorescently labeling the nucleocapsid proteins (Schudt *et al.*, 2013, Schudt *et al.*, 2015, Becker *et al.*, 1996, Kolesnikova *et al.*, 2004a, Kolesnikova *et al.*, 2004b). According to these reports, nucleocapsid formation occurs in the

perinuclear inclusion bodies, following which they are transported into the

cytoplasm and redistributed prior to budding through the cell surface. The velocity

of nucleocapsid movement inside cells ranges from 100 nm/s to 500 nm/s in

MARV- or EBOV-infection (Schudt et al., 2013, Schudt et al., 2015). Application

of specific cytoskeleton inhibitors revealed that the transport of MARV

nucleocapsids was dependent on actin polymerization. Viral matrix protein VP40,

which is a peripheral membrane protein and plays a pivotal role in filamentous

virion formation, is essential for the recruitment of nucleocapsids to the cell

periphery and for their incorporation into progeny virions (Noda *et al.*, 2007,

Bharat et al., 2012, Noda *et al.*, 2002, Dolnik *et al.*, 2010). The surface

glycoprotein GP, which is an integral membrane protein and is responsible for cell

entry, forms the filamentous virions together with VP40 (Fig. 1b) (Becker et al.,

1996, Mittler *et al.*, 2007, Beniac & Booth, 2017, Booth *et al.*, 2013).

Live-cell imaging is a powerful tool for visualization of the spatio-temporal

dynamics of living organisms. In addition to immunofluorescence microscopy,

live-cell imaging microscopy has been utilized to visualize the localization of viral

proteins, and interactions between viral and host proteins in various virus-infected

cells (Nanbo *et al.*, 2013, Hoenen *et al.*, 2012, Lakdawala *et al.*, 2014, Becker et

al., 1996). However, because of its high pathogenicity, MARV must be handled

under the highest biosafety conditions [biosafety level 4 (BSL-4)], which

complicates and delays research using live-cell imaging (Falzarano *et al.*, 2011).

In this study, we developed a safe, live-cell imaging system, following a previously

established method for EBOV (Takamatsu *et al.*, 2018), to visualize MARV

100     nucleocapsid-like structures (NCLSs) in cells expressing viral proteins, outside of

101     BSL-4 laboratories. By using this live-cell imaging system, we were able to

102     analyze interactions between NCLSs and the cellular cytoskeleton, as well as

103     intracellular transport of NCLSs.

## Materials and methods

### Cell culture.

Huh-7 (human hepatoma) cells were maintained at 37 °C and 5% $CO_2$ in Dulbecco's Modified Eagle Medium (DMEM, Life Technologies) supplemented with 10% (vol/vol) Fetal bovine serum (FBS , PAN Biotech), 5 mM L-glutamine (Q; Life Technologies), 50 U/mL penicillin, and 50 µg/mL streptomycin (PS; Life Technologies).

### Plasmids and transfection.

Plasmids encoding the MARV structural proteins (NP, VP35, VP24, L, VP40 and GP): pCAGGS-NP, pCAGGS-VP35, pCAGGS-VP24, pCAGGS-L, pCAGGS-VP40, and pCAGGS-GP, and a MARV minigenome-expressing plasmid which encodes a *Renilla* luciferase were used (Wenigenrath *et al.*, 2010, Hoenen *et al.*, 2011). The plasmid pCAGGS-VP30-GFP, coding for the green fluorescent protein-VP30 fusion protein, was produced as previously described (Schudt et al., 2013). The transfection was performed in 50 µL Opti-MEM without phenol red (Life Technologies) using TranSIT (Mirus) according to the manufacturer's instructions.

**Lice cell imaging microscopy.**

A total of $2 \times 10^4$ Huh-7 cells were seeded onto a µ-Slide 4 well (Ibidi) and cultivated in DMEM/PS/Q with 10% FBS. Each well was transfected with the following plasmids, encoding all MARV structural proteins: (250 ng of pCAGGS-NP, 50 ng of pCAGGS-VP35, 125 ng of pCAGGS-VP30-GFP, 50 ng of pCAGGS-VP24, 500 ng of pCAGGS-L, 125 ng of pCAGGS-VP40 and 125 ng of pCAGGS-GP), together with a T7-driven, MARV minigenome-expressing plasmid, which encodes a *Renilla* luciferase, and a T7 polymerase-coding plasmid (pCAGGS-T7) (Wenigenrath et al., 2010, Hoenen et al., 2011). The inoculum was removed at 1 h post-transfection (p.t.), and 500 µL $CO_2$-independent Leibovitz's medium (Life Technologies) with PS/Q, non-essential amino acid solution, and 20% (vol/vol) FBS were added. Live-cell time-lapse experiments were recorded with a Nikon ECLIPSE TE2000-E using a 63× oil objective or a GE healthcare Delta Vision Elite using a 60× oil objective in biosafety level-2 laboratories.

**Treatment of cells with cytoskeleton-modulating drugs.**

Cells were treated with 15 µM nocodazole (Sigma), 0.3 µM cytochalasin D

(Sigma), or 0.15% dimethyl sulfoxide (DMSO, Sigma), following previous

publication (Schudt et al., 2013). The chemicals were added to the cell culture

medium 3 hours prior to observation.

**Image processing and analysis.**

Acquired pictures and movie sequences were processed using the Fiji plugin

"TrackMate" (Tinevez *et al.*, 2017, Schindelin *et al.*, 2012).

**Results and Discussion**

**Establishment of a live-cell imaging system for MARV NCLSs transport.**

The Marburg virus VLP system, which models a complete, single infectious cycle, has been developed and used to analyze transcription and replication, as well as the budding processes (Wenigenrath et al., 2010, Hoenen et al., 2011). In this study, we attempted to visualize MARV NCLSs transport in Huh-7 cells, which is a suitable method to determine the sequential intracytoplasmic movement of MARV nucleocapsids (Schudt et al., 2013), by using this VLP system following the procedure established in EBOV (Takamatsu et al., 2018). Transfection of the VLP components with the following modifications: replacing VP30 with VP30-GFP, was performed as described in Figure 2a. We employed this VLP-based system in all subsequent experiments in this study.

At 18 h p.t., we detected a large number of signals with various shapes (Supplementary Movie 1). Over 1,000 signals in the acquired movie sequences were analyzed. The representing sequence was expressed as the maximum intensity projection, in which the image projected maximum intensity of each time-lapse were plotted. (Fig. 2b). Among the signals, we focused on the signals

showing long-distance and directional transport, which represent NCLSs

trajectories (Takamatsu et al., 2018). We detected trajectories with lengths

ranging from 100 nm to 20 µm, with a mean length of 1.89 ± 0.91 µm (Fig. 2c).

The direction of movement of each NCLS also varied, though the cause of this

variation remains unclear. The speed of MARV NCLSs transport ranged from 10

nm/s to 500 nm/s, with a mean velocity of 167 ± 96 nm/s (Fig. 2d), which is

comparable to the velocity of nucleocapsids transport in MARV-infected cells

analyzed in BSL-4 laboratories (Schudt et al., 2013). In summary, the movement

characteristics of NCLSs in our system is similar to those of nucleocapsids in

MARV-infected cells.

**Actin polymerization is required for MARV NCLSs transport.**

In MARV-infected cells, the microtubule depolymerizing drug nocodazole

does not alter the movement of nucleocapsids, whereas the actin depolymerizing

drug cytochalasin D arrests it (Schudt et al., 2013). To confirm the relevance of

the live-cell imaging system we developed in this study, we analyzed NCLSs

movement after treatment with cytoskeletal modulation drugs. Huh-7 cells were

transfected with plasmids as described in Figure 2a. The culture medium was

replaced at 15 h p.t. with Leibovitz's medium containing either 0.15 % DMSO

(control), 0.15 M nocodazole, or 0.3 µM cytochalasin D (Fig. 3a-d, Supplementary

Movie 2-4) (Schudt et al., 2013). After incubation the cells with cytoskeletal

modulating drugs for 3h, time-lapse images were acquired. Nocodazole

treatment did not alter the trajectory length of NCLSs transport in comparison to

the control, whereas cytochalasin D treatment induced immediate cessation of

long-distance transport (Fig. 3e-g). The mean velocity of NCLSs transport in the

control or nocodazole-treated cells was 188 ± 98 nm/s and 219 ± 99 nm/s,

respectively (Fig. 3h). On the other hand, only a few non-specific NCLSs

movements were detectable in cytochalasin D-treated cells, with a mean velocity

of 2.73 ± 23 nm/s (Fig. 3i-j). These results confirmed that NCLSs transport is

dependent on actin polymerization, as well as the availability of our assay to test

candidate drugs which target nucleocapsid intracellular transport.

**Concluding remarks**

In the present study, we developed a live-cell imaging system for cells

expressing viral proteins, which can be safely used in BSL-2 laboratories.

Furthermore, we demonstrated the relevance of our system as a substitute for

the analysis of nucleocapsids transport in MARV-infected cells. Currently, cellular

factors involved in the nucleocapsid transport have not been fully understood.

The combined approach of gene silencing and inhibitor screening using the

system we developed can be utilized to identify the key host factors for the

intracellular transport of MARV nucleocapsids. Moreover, it is noteworthy that the

technical approach developed here might be applicable to study the nucleocapsid

transport of other mononegaviruses, as well as to characterize antivirals inhibiting

nucleocapsid transport.

**Acknowledgments**

The authors are grateful to Olga Dolnik and Gordian Schudt (Philipps University

Marburg, Germany) for fruitful discussion.

**Funding**

The work was supported by Japan Society for the Promotion of Science JSPS

Grant number 18J01631, 19K16666 (to Y.T.), by AMED, Research Program on

Emerging and Re-emerging Infectious Diseases, by AMED Japanese Initiative

for Progress of Research on Infectious Disease for global Epidemic, by JSPS

Core-to-Core Program A, the Advanced Research Networks, by Grant for Joint

Research Project of the Institute of Medical Science, University of Tokyo, by Joint

Usage/Research Center program of Institute for Frontier Life and Medical

Sciences Kyoto University, by the Daiichi Sankyo Foundation of Life Science, and

by the Takeda Science Foundation (to T.N.), and by the Deutsche

Forschungsgemeinschaft (DFG, German research foundation) Project number

197785619-SFB 1021 (to S.B.).

## References

Becker, S., Klenk, H. D. & Muhlberger, E. (1996) Intracellular transport and processing of the Marburg virus surface protein in vertebrate and insect cells. *Virology,* 225, 145-155.

Beniac, D. R. & Booth, T. F. (2017) Structure of the Ebola virus glycoprotein spike within the virion envelope at 11 A resolution. *Scientific reports,* 7, 46374.

Bharat, T. A., Noda, T., Riches, J. D., Kraehling, V., Kolesnikova, L., Becker, S., Kawaoka, Y. & Briggs, J. A. (2012) Structural dissection of Ebola virus and its assembly determinants using cryo-electron tomography. *Proceedings of the National Academy of Sciences of the United States of America,* 109, 4275-4280.

Biedenkopf, N., Hartlieb, B., Hoenen, T. & Becker, S. (2013) Phosphorylation of Ebola virus VP30 influences the composition of the viral nucleocapsid complex: impact on viral transcription and replication. *The Journal of biological chemistry,* 288, 11165-11174.

Booth, T. F., Rabb, M. J. & Beniac, D. R. (2013) How do filovirus filaments bend without breaking? *Trends in microbiology,* 21, 583-593.

CDC (2014) Chronology of Marburg Hemorrhagic Fever Outbreaks.

Dolnik, O., Kolesnikova, L., Stevermann, L. & Becker, S. (2010) Tsg101 is recruited by a late domain of the nucleocapsid protein to support budding of Marburg virus-like particles. *Journal of virology,* 84, 7847-7856.

Falzarano, D., Geisbert, T. W. & Feldmann, H. (2011) Progress in filovirus vaccine development: evaluating the potential for clinical use. *Expert review of vaccines,* 10, 63-77.

Hartlieb, B., Modrof, J., Muhlberger, E., Klenk, H. D. & Becker, S. (2003) Oligomerization of Ebola virus VP30 is essential for viral transcription and can be inhibited by a synthetic peptide. *The Journal of biological chemistry,* 278, 41830-41836.

Hoenen, T., Groseth, A., de Kok-Mercado, F., Kuhn, J. H. & Wahl-Jensen, V. (2011) Minigenomes, transcription and replication competent virus-like particles and beyond: reverse genetics systems for filoviruses and other negative stranded hemorrhagic fever viruses. *Antiviral research,* 91, 195-208.

Hoenen, T., Shabman, R. S., Groseth, A., Herwig, A., Weber, M., Schudt, G., Dolnik, O., Basler, C. F., Becker, S. & Feldmann, H. (2012) Inclusion bodies are a site of ebolavirus replication. *Journal of virology,* 86, 11779-11788.

Huang, Y., Xu, L., Sun, Y. & Nabel, G. J. (2002) The assembly of Ebola virus nucleocapsid requires virion-associated proteins 35 and 24 and posttranslational modification of nucleoprotein. *Molecular cell,* 10, 307-316.

Kolesnikova, L., Bamberg, S., Berghofer, B. & Becker, S. (2004a) The matrix protein of
Marburg virus is transported to the plasma membrane along cellular membranes:
exploiting the retrograde late endosomal pathway. *Journal of virology,* 78, 2382-2393.
Kolesnikova, L., Berghofer, B., Bamberg, S. & Becker, S. (2004b) Multivesicular bodies as a
platform for formation of the Marburg virus envelope. *Journal of virology,* 78, 12277-
12287.
Lakdawala, S. S., Wu, Y., Wawrzusin, P., Kabat, J., Broadbent, A. J., Lamirande, E. W., Fodor,
E., Altan-Bonnet, N., Shroff, H. & Subbarao, K. (2014) Influenza a virus assembly
intermediates fuse in the cytoplasm. *PLoS pathogens,* 10, e1003971.
Mittler, E., Kolesnikova, L., Strecker, T., Garten, W. & Becker, S. (2007) Role of the
transmembrane domain of marburg virus surface protein GP in assembly of the viral
envelope. *Journal of virology,* 81, 3942-3948.
Muhlberger, E., Weik, M., Volchkov, V. E., Klenk, H. D. & Becker, S. (1999) Comparison of
the transcription and replication strategies of marburg virus and Ebola virus by
using artificial replication systems. *Journal of virology,* 73, 2333-2342.
Nanbo, A., Watanabe, S., Halfmann, P. & Kawaoka, Y. (2013) The spatio-temporal
distribution dynamics of Ebola virus proteins and RNA in infected cells. *Scientific*
*reports,* 3, 1206.
Noda, T., Sagara, H., Suzuki, E., Takada, A., Kida, H. & Kawaoka, Y. (2002) Ebola virus VP40
drives the formation of virus-like filamentous particles along with GP. *Journal of*
*virology,* 76, 4855-4865.
Noda, T., Watanabe, S., Sagara, H. & Kawaoka, Y. (2007) Mapping of the VP40-binding
regions of the nucleoprotein of Ebola virus. *Journal of virology,* 81, 3554-3562.
Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch,
S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J. Y., White, D. J., Hartenstein, V.,
Eliceiri, K., Tomancak, P. & Cardona, A. (2012) Fiji: an open-source platform for
biological-image analysis. *Nature methods,* 9, 676-682.
Schudt, G., Dolnik, O., Kolesnikova, L., Biedenkopf, N., Herwig, A. & Becker, S. (2015)
Transport of Ebolavirus Nucleocapsids Is Dependent on Actin Polymerization: Live-
Cell Imaging Analysis of Ebolavirus-Infected Cells. *The Journal of infectious diseases,*
212 Suppl 2, S160-166.
Schudt, G., Kolesnikova, L., Dolnik, O., Sodeik, B. & Becker, S. (2013) Live-cell imaging of
Marburg virus-infected cells uncovers actin-dependent transport of nucleocapsids
over long distances. *Proceedings of the National Academy of Sciences of the United*
*States of America,* 110, 14402-14407.
Takamatsu, Y., Kolesnikova, L. & Becker, S. (2018) Ebola virus proteins NP, VP35, and VP24

are essential and sufficient to mediate nucleocapsid transport. *Proceedings of the*
*National Academy of Sciences of the United States of America,* 115, 1075-1080.

Tinevez, J. Y., Perry, N., Schindelin, J., Hoopes, G. M., Reynolds, G. D., Laplantine, E.,
Bednarek, S. Y., Shorte, S. L. & Eliceiri, K. W. (2017) TrackMate: An open and
extensible platform for single-particle tracking. *Methods,* 115, 80-90.

Watanabe, S., Noda, T. & Kawaoka, Y. (2006) Functional mapping of the nucleoprotein of
Ebola virus. *Journal of virology,* 80, 3743-3751.

Wenigenrath, J., Kolesnikova, L., Hoenen, T., Mittler, E. & Becker, S. (2010) Establishment
and application of an infectious virus-like particle system for Marburg virus. *The*
*Journal of general virology,* 91, 1325-1334.

**Figure legends**

**Figure 1. Orientation of MARV nucleocapsid and its transport pathway.** (a)

The viral genome is encapsidated by NP. Nucleocapsids additionally contain

VP24, VP30, VP35, and L. Among them, NP, VP24, and V35 form the core

structure of the nucleocapsid called a "nucleocapsid-like structure (NCLS)". (b)

Nucleocapsids are formed in the perinuclear inclusion bodies and are

subsequently transported along polymerized actin filaments to the plasma

membrane, where budding and release of virions takes place. The VP40 protein

forms filamentous virions together with GP.

**Figure 2. Live-cell imaging system of MARV nucleocapsid-like structures.**

(a) The experimental setting for detection of NCLS transport. Huh-7 cells were

transfected with plasmids encoding NP, L, VP35, VP24, VP40, GP, Marburg virus

specific minigenome, T7 polymerase, and VP30-GFP. (b) Plasmid-transfected

Huh-7 cells were observed at 18 h p.t. The image shows the maximum-intensity

projection of time-lapse images of cells, recorded for 90 seconds; images were

captured every 2 s. (c, d) Over 100,000 moving signals were captured and over

1,000 selected signals were analyzed using the Fiji plugin "TrackMate". (c) The

length of the NCLS trajectories was evaluated. The y-axis represents the number

of signals in each range (x-axis). The numbers indicate mean ± SD (µm). (d) The

velocity of NCLSs transport was evaluated. The y-axis represents the number of

signals in each range (x-axis). The numbers indicate mean ± SD (nm/s).

**Figure 3. Effect of cytoskeleton-modulating drugs on MARV NCLSs**

**transport.** (a) Experimental setting to observe the effects of cytoskeleton

modulating drugs on MARV NCLSs transport. Huh-7 cells were transfected with

plasmids as described in Fig. 2(a), and treated at 15 h p.t. with either 0.15 %

DMSO (control), 0.15 µM nocodazole, or 0.3 µM cytochalasin D. After 3 hours of

treatment, observation of the cells began (18 h p.t.). (b-d) Time-lapse images

were acquired for each of the drug treated cells  (b: DMSO, c: Nocodazole, d:

Cytochalasin D). The pictures show the maximum-intensity projection of time-

lapse images of cells, recorded for 90 seconds; images were captured every 2 s.

(e-i) Over 100,000 moving signals were captured and over 1,000 selected signals

were analyzed using Fiji plugin "TrackMate". (e-g) The length of the NCLS

trajectories was evaluated in each of the drug treated cells (e: DMSO, f:

Nocodazole, g: Cytochalasin D). The y-axis represents the number of signals in

each range (x-axis). The numbers indicate mean ± SD (µm). (h-i) The velocity of

NCLSs transport was evaluated in each of the drug treated cells (h: DMSO, i:

Nocodazole, j: Cytochalasin D). The y-axis represents the number of signals in

each range (x-axis). The numbers indicate mean ± SD (nm/s).

(a)

**nucleocapsid**

L

VP30

vRNA

**nucleocapsid-like structure**

- 🔴 NP
- 🟢 VP24
- ⚪ VP35

(b)

polymerized actin

virion

Inclusion bodies

nucleocapsid

nucleus

🟡 VP40

GP

**Figure 1**

**(a)**

NP  VP35  VP30-GFP  L  T7  minigenome

VP24  VP40  GP

Huh-7 cells

**(b)**

10 μm

**(c)**

1.89 ± 0.91 (μm)

count

Length of trajectories (μm)

**(d)**

167 ± 96 (nm / s)

count

Velocity (nm / s)

**Figure 2**

**Figure 3**