Associations between neonatal cry acoustics and visual attention during the first year

Aicha Kivinummi[1], Gaurav Naithani[2], Outi Tammela[3], Tuomas Virtanen[2], Enni Kurkela[1], Miia

Alhainen[5], Dana JH Niehaus[4], Anusha Lachman[4], Jukka M. Leppänen[5], and Mikko J. Peltola[1]

[1] Faculty of Social Sciences, Tampere University, Tampere, Finland

[2] Faculty of Information Technology and Communication Sciences, Tampere University, Tampere,
Finland

[3] Department of Pediatrics, Tampere University Hospital, Tampere, Finland

[4] Department of Psychiatry, Faculty of Health Sciences, University of Stellenbosch, Cape Town,
Republic of South Africa

[5] Faculty of Medicine and Health Technology, Tampere University, Tampere, Finland

Corresponding author: Mikko Peltola, Faculty of Social Sciences, Tampere University, 33014

Tampere, Finland. E-mail: mikko.peltola@tuni.fi, tel: +358503186120

Abstract

24
25
26

It has been suggested that early cry parameters are connected to later cognitive abilities. The present

study is the first to investigate whether the acoustic features of infant cry are associated with

cognitive development already during the first year, as measured by oculomotor orienting and

attention disengagement. Cry sounds for acoustic analyses (fundamental frequency; F0) were

recorded in two neonatal cohorts at the age of 0-5 days (Tampere, Finland) or at 6 weeks (Cape

Town, South-Africa). Eye-tracking was used to measure oculomotor orienting to peripheral visual

stimuli and attention disengagement from central stimuli at 8 months (Tampere) or at 6 months

(Cape Town) age. In the Tampere cohort, a marginal positive correlation between fundamental

frequency of cry (F0) and visual attention disengagement was observed; infants with a higher

neonatal F0 were slower to shift gaze away from the central stimulus to the peripheral stimulus.

However, a similar correlation was not observed in the Cape Town cohort. No associations between

F0 and oculomotor orienting were observed in either cohort. We discuss possible factors

influencing the discrepancy in results between the cohorts and suggest directions for future research

investigating the potential of early cry analysis in predicting later cognitive development.

Introduction

Monitoring children's cognitive development is an important part of pediatric practice. Traditional psychometric methods enable detailed and reliable measurements especially after two years of age. However, the assessment of cognitive development during the neonatal period and early infancy is faced with multiple challenges, such as age-related changes in how different cognitive functions can be measured. For example, motor abilities are emphasized in most early measures, but their relation to cognitive functioning may vary at different ages, which may weaken the predictive validity of early assessments of cognitive development [1,2]. These challenges make research on the early markers of later cognitive functioning important.

The current study is focused on the connection between the acoustic characteristics of infant cry and visual attention within the first year of life. Cry is a vital, reflex-like means of communication present from the very beginning of life [3] and it indicates the neurological status of the neonate [4,5] . Variations in visual attention during the first year have been shown to be associated with later cognitive development [6-9], thus providing an outcome measure that emerges early in development and is meaningfully associated with more complex and later-emerging cognitive functions. As both the acoustic characteristics of cry and visual attention can be measured reliably at an early age, examining their interrelations may help to determine the potential of neonatal cry analysis as an accessible marker of later development.

A limited number of small-scale studies are available to suggest that early cry parameters are associated with later cognitive abilities. Lester [10] observed that preterm and term infants having high average fundamental frequency of cry (F0; heard as the pitch of the cry) and larger variability in F0 within 2 weeks of term postconceptional age were more likely located in the group with lower scores on the Bayley Scales of Infant Development at 18 months of age compared to infants with lower average F0 and less variable F0. In addition, high variation in neonatal F0 predicted lower scores in the McCarthy Scales of Children's Abilities at 5 years of age. Similarly,

studying infants prenatally exposed to methadone, Huntington et al. [11] found that higher and

more variable F0 in the first days of life were related to lower mental development scores measured

by the Bayley scales at 8, 12, and 18 months. However, the association was only marginal at 24

months. Both Lester [10] and Huntington et al. [11] observed that the relation between the F0

variables and the cognitive development outcomes remained after controlling for risk factors

including prematurity and prenatal exposure to drugs. Other studies have observed that the acoustic

characteristics of cry in children with neurodevelopmental disorders such as CNS abnormalities and

developmental delays (for reviews, see [4,12] and children with autism spectrum disorders [13,14]

are associated with an atypical and more unstable F0. The similarities in the findings of studies

investigating at-risk infants, as well as the scarce findings among normally developing infants

emphasize the importance of further research on the connections between the acoustic features of

newborn cry and cognitive functioning during the first year of life among typically developing

infants.

85   Regarding the neural basis of variation in neonatal cry, it is suggested that imperfect

functioning of the vagus nerve and inhibitory parasympathetic nervous system activity are reflected

in infant cry as higher and more variable F0 [15,16]. Neonatal cry requires versatile integration of

respiratory and larynx area organs, which are suggested to be controlled in early, reflex-like crying

exclusively by the brainstem. No higher areas than the midbrain are required for producing cry

during the first months of life [3]. Consequently, the integrity of the brainstem is suggested to

influence the acoustic features of cry [10,17] and the vagus nerve of the parasympathetic nervous

system is considered as a crucial pathway through which the brainstem regulates neonatal and

infant cry [3,4,18]. The vagus nerve originates from the medulla oblongata in the brainstem and is

densely connected with the vocal cords and respiratory muscles, on which it has an inhibitory

function. In addition, neonatal F0 has been shown to vary as a function of cardiac vagal tone

(Shinya et al., 2016). Thus, imperfect inhibitory parasympathetic control of the cry organs is likely

reflected in higher and more variable infant F0. These findings highlight the role of the brainstem

and vagal system in affecting neonatal cry acoustics.

In addition to influencing infant cry production, integrity of the brainstem may have a

pivotal impact on visual attention and the processing of visual information [19-23]. The

physiologically based model of attention suggests that vagus nerve mediated organizational

processes of the brainstem are required to integrate sensory and motor information on attention. As

a part of the parasympathetic nervous system, the vagus nerve has an inhibitory function on arousal.

Further on, attention is typically modulated by arousal state and it has been observed that infants

with brainstem dysfunction show atypical modulation of attention by arousal. Healthy neonates

preferred familiar visual stimuli during increased arousal and novel stimuli during decreased

arousal, whereas this pattern was less pronounced in neonates with brainstem dysfunction [24].

Karmel et al. [25] observed that a similar atypical arousal-modulated attention pattern persists at

least until 4 months of age in infants with brainstem dysfunction. In addition, sensitivity to social

and non-social content of visual stimuli is connected to brainstem integrity. Geva et al. [26]

detected that children with discrete neonatal brainstem dysfunction preferred an active nonsocial

content (a flock of birds flying) or a neutral social content (a person listening) over an active social

content (a person expressing affective narratives). These findings were opposite to the pattern of

attention in normal controls and the difference remained even in a follow up at 8 years of age.

Thus, it appears that neonatal cry and infant visual attention share partially overlapping

neural mechanisms, with particularly the brainstem implicated in both. In the current study, we

therefore investigated the associations between infant cry acoustics (F0 and its variation) during the

neonatal period (0-6 weeks) and visual attention at 6-8 months of age. As the indicator of infant

visual attention, we measured oculomotor orienting and attention disengagement, both

operationalized as saccadic reaction times in separate eye-tracking paradigms.

The ability to produce rapid shifts of gaze to stimuli appearing in a new spatial location,

typically measured as saccadic reaction time, matures in early infancy and its variations correlate

with later cognitive development [7]. For example, infants with longer latency of saccades to

peripheral target stimuli were shown to score lower on standardized IQ measures at 4 years of age

(Dougherty & Haith, 1997). In addition, saccadic reaction times have been shown to be affected by

neurodevelopmental risk factors such as prenatal alcohol exposure [27], preterm birth [28] and

familial risk for autism spectrum disorder [29]. Hence, studying infant visual attention with

paradigms measuring saccadic reaction times may provide objective and applicable markers for

examining neurocognitive development in a variety of populations.

We used two infant attention paradigms for measuring saccadic reaction times. First,

oculomotor orienting was measured as saccadic reaction time to a new peripheral stimulus

following the offset of a stimulus on the center of the screen. Second, we measured disengagement

of attention as the duration of gaze shift from centrally presented face and non-face stimuli to

laterally presented competing stimuli. Particularly oculomotor orienting has been associated with

later cognitive development, with faster saccadic reaction times in infancy predicting higher

childhood IQ (Dougherty & Haith, 1997; Rose, Feldman, Jankowski, & Van Rossem, 2012).

Attention disengagement paradigms have been especially used for investigating infants' attention to

social stimuli such as faces [30] with recent findings indicating that variations in infants' attention

to faces in this paradigm are associated with later social development [31]. Furthermore, given that

oculomotor orienting and attention disengagement appear to mature at different speed during the

first year (with attention disengagement maturing later) [19,32,33] and rely on different neural

mechanisms [19,34] we considered it important to include both measures of infant attention in the

current study.

Taken together, as infant cry and visual attention may have a partially overlapping neural

basis and as the basic functions of infant visual attention may be important in understanding later-

emerging more complex cognitive and social abilities, infant visual attention is an ideal outcome

measure for attempts to determine whether the acoustic features of newborn cry are associated with

normative developmental variations in basic cognitive functions.

Due to the rather explorative nature of our study, we implemented the analysis across two

independent samples of infants from Tampere, Finland and Cape Town, South Africa, to assess the

reliability of the findings. Based on previous findings showing that both F0 of infant cry [11,35]

and saccadic reaction times (Dougherty & Haith, 1997; Rose et al., 2012) are associated with

childhood cognitive development, we hypothesized that higher and more variable F0 would be

associated with longer saccadic reaction times in the oculomotor orienting task. Regarding the

attention disengagement paradigm, the paucity of previous research prevented strong directional

hypotheses. However, given the links between neonatal cry production and brainstem integrity on

the one hand [3,17,35] and between brainstem integrity and attention to social stimuli on the other

hand (Geva et al., 2017), it could be tentatively hypothesized that higher and more variable F0

would be associated with less attention to faces in the attention disengagement paradigm (i.e., faster

saccadic reaction times in shifting attention away from faces).

Materials and Methods

Ethics Statement

164
165   The study adhered to ethical guidelines and the study protocol was approved by the Ethical

166 Committee of Tampere University Hospital and on behalf of Cape Town participants by the

167 Institutional Review Board of the University of Stellenbosch. Infants were included in the study

168 after a written consent from their parent was obtained.

Participants

170
Data for the current analyses were obtained from two cohorts, one in Tampere, Finland, and

the other in Cape Town, South Africa. Our interest was to study the associations between acoustic

variables of cry during the neonatal period (Tampere: 0-5 days after birth; Cape Town: 6 weeks of

age) and infant visual attention measured by eye-tracking methods (Tampere: 8 months; Cape

Town: 6 months of age).

Tampere cohort

Participants in the Tampere cohort were enrolled in the study at birth (0-5 days) and

subsequently invited for a follow-up visit at 8 months of age (mean = 253.58 days, SD = 12.34

days). A total of 104 neonates were recruited from the Tampere University Hospital Maternity

Ward Units and the Neonatal Ward Unit between April and August 2014. Infants were considered

eligible to be enrolled in the study if the infant did not have any kind of abnormalities in the neck

area, nor a history of recent airway suctioning, which may influence the voice, and if one of the

parents understood Finnish or English sufficiently to be able to be adequately informed about the

study.

From the original pool of 104 infants, 6 infants did not cry when the researcher was present,

hence cry recording was captured from 98 infants. Out of these, 73 infants attended the eye-tracking

follow-up visit at 8 months. Eye-tracking data were not obtained from 5 infants because of technical

problems and from 4 infants because the infant showed inconvenience about the registration

situation. Two infants were further excluded because of not being full-term ($\geq$ 37 weeks), 3 because

of low birth weight ($\geq$ 2500 grams), 2 because of perinatal asphyxia, and 1 because of an abnormal

brain MRI finding.

From this sample of 56 infants, 2 infants needed to be excluded because of the inclusion

criteria of the cry analysis. At least 3 expiratory cry utterances from the beginning of the recording

had to be present for analyzing the F0. For the analyses of oculomotor orienting, 2 infants were

excluded because of for the number of less than 4 trials with good gaze data for calculating

oculomotor orienting. For the analyses of attention disengagement, 8 infants were excluded for

having less than 3 scorable trials in one of the stimulus conditions (i.e., a non-face control stimulus

and happy and fearful face stimuli).

Consequently, 52 infants (22 female, 30 male) and 46 infants (female 19, male 27) were

included in the oculomotor orienting and attention disengagement analyses, respectively. The

included infants did not have known health or developmental disadvantages nor known prenatal

mother-related risk factors, such as drug abuse, untreated diabetes or high blood pressure. Data

about health, risk factors, and medical therapies were collected from the medical records of the

hospital.

In addition to the primary analyses, we ran additional analyses including only those infants

whose cry was captured from the beginning of a cry bout. This was done to enable more direct

comparisons with the Cape Town cohort in which the cry recordings were always captured from the

beginning of the cry bout. Thus, this analysis controlled the possible influence of the location of the

cry sample in the recording. For these analyses, 29 and 28 infants were included in the oculomotor

orienting and attention disengagement analyses, respectively.

Cape Town cohort

The data for the Cape Town cohort was obtained from a larger research project in

collaboration with the Department of Psychiatry at the University of Stellenbosch. Participants were

enrolled in this study at the age of 6 weeks (mean = 44,73 days, SD = 3,34 days) between January

2014 and May 2015 and they participated in the follow-up registration at 6 months (mean = 187.85

days, SD = 13.838 days) of age. Similarly to the Tampere cohort, the infants did not have any kind

of abnormalities in the neck area, nor a history of recent airway suctioning, which may influence

the voice. Likewise, the other exclusion criteria were concordant with the Tampere cohort.

Cry recordings were obtained from a total of 100 infants. The follow-up assessment was

completed by 73 infants. Of these, 5 infants needed to be excluded because of the age criteria at cry

recording and 4 because there were less than 3 expiratory cry utterances in the recording. Three

infants were excluded because of not being full-term ($\geq$ 37 weeks). Birthweight data was not

available for this cohort, and none of the infants had perinatal asphyxia or a brain MRI finding. The

infants did not have other health or developmental disadvantages. Considering the inclusion criteria

of mother-related risk factors, we excluded 6 infants due to prenatal alcohol use. For the eye-

tracking data, identical inclusion criteria were used as in the Tampere cohort. Consequently, 47

infants (18 female, 29 male) and 53 infants (19 female, 34 male) were included in the oculomotor

orienting and disengagement analyses, respectively.

Within the cohort in the primary analysis, there were 14 infants exposed to prenatal SSRI

medication. Therefore, to control the possible influence of SSRI medication, additional analysis

were conducted after excluding these participants, resulting in 37 and 42 infants in the oculomotor

orienting and attention disengagement analyses, respectively. In both the analyses a proportion of

mothers had psychiatric problems such as depression and anxiety disorders. This was due to the

interests in the larger research project, from where the data was captured.

Measurements
237

Cry recordings in Tampere

We recorded one cry sample from each infant in normal clinical settings at the age of 0 to 5

days. During the recording, infants were laying on supine position. The location of the captured

sample in the cry bout varied since we recorded spontaneous cries which were not launched by a

distinct painful trigger. Soothing was performed by the parent whenever needed.

The duration of the samples in the whole data ranged from 14.4 s to 430.5 s (mean =114.2 s;

SD = 81.7) The recording microphone was a Tascam DR-100MK Linear PCM Recorder, Røde, M3

cardioid microphone. The samples were stored with a 48-kHz sampling rate, in a 24-bit Waveform

Audio file (WAV) format. The microphone was at a stand at 30-cm distance directly in front of the

infant's mouth.

Cry recordings in Cape Town

Recording of cry was done as a part of a standardized protocol at a routine 6-weeks baby

clinic visit to the nurse. The room was always the same and the cry was elicited by vaccination or

measuring the infant's weight on a scale. During the vaccination, the infant was laying on mother's

lap. Soothing was performed in the way the mother preferred. We captured one cry sample at the

beginning of the cry bout from every infant.

The duration of the samples in the whole data ranged from 16.9 s to 150 s (mean = 80.7 s;

SD = 18.8 s). The audio recorder used was a Zoom H4n recorder with built-in condenser

microphones. Recordings were stored with a 48-kHz sampling rate, two-channel audio in a 24-bit

Waveform Audio file (WAV) format. The distance between the infant's mouth and the microphone

was 1.3 m for infants being vaccinated and 70 cm for infants being weighed. The microphone was

fixed on a wall stand.

Acoustic variables

As the cry samples were captured in clinical settings containing varying amounts of

extraneous sounds, the first task was to separate the cry vocalizations from these extraneous sounds

which are not useful for the purpose of acoustic analysis. We used an in-house hidden Markov

model (HMM) [36] based audio segmentation system (Naithani et al., 2018) for cry segment

extraction, and the acoustic analysis was performed in MATLAB (Mathworks, Natick, MA). The

cry data was divided to three acoustic regions; expiratory phases, inspiratory phases, and residual,

where residual included all the vocalizations that are not expiratory or inspiratory cries. We

included only expiratory phases in the analysis, based on the frequent praxis in earlier studies as

well as the results of our preliminary study showing that expiratory cries were detected more

reliably than inspiratory cries by the analysis tool [37].

In the analyses, we used the first 5 expiratory phases from the beginning of the captured cry

sample. The missing values of the used variables were imputed based on other scorable phases if

there was a minimum of 3 scorable phases. Only phases with duration longer than 500 ms were

included, in line with previous studies [38,39]. The identified expiratory phases were then divided

into overlapping frames of 25-ms duration and 50 % overlap, and fundamental frequency (F0) was

extracted for each such frame.

As cry variabes we used fundamental frequency (F0) and the variation of fundamental

frequency (F0var), which we estimated using the YIN algorithm (De Cheveigné & Kawahara, 2002).

In line with earlier studies (Michelsson & Michelsson, 1999; Rothgänger, 2003), F0 was defined as

the mean of first five within-sample means of frame-wise fundamental frequencies, and F0var was

defined as mean of the first five within-sample standard deviations of frame-wise fundamental

frequencies. Cry phases with more than 70% inharmonic frames selected via absolute threshold

parameter of the YIN algorithm (here 0.3), were excluded (for more information, see Fuamenya et

al., 2015). Descriptive characteristics of F0 in different analyses are presented in Table 1.

Table 1. Descriptive data of F0 in Tampere and Cape Town cohorts.

Cohort	F0 (Hz)	F0 SD	F0var	F0var SD
Tampere ($N = 54$)	463.45	62.52	65.18	26.10
Cape Town ($N = 58$)	475.34	48.20	54.63	21.49

Eye tracking assessments in Tampere

In the 8-month follow-up visit, the infants participated in an eye-tracking assessment of

saccadic reaction times measured during separate oculomotor orienting and attention

disengagement paradigms. During the assessment, the infant sat on the parent's lap in a dimly lit

room watching the test stimuli on a 23-in monitor located at 60±5 cm distance from the infant's

eyes. The parent was asked to close her/his eyes during the recording to avoid false registrations

[40]. The session was paused or terminated in case the infant expressed inconvenience or was not

willing to co-operate. Eye tracking data were collected by using a Tobii TX300 remote eye tracking

camera (Tobii Technology, Stockholm, Sweden) with a 300-Hz sampling rate.

Before recording the data, the eye-tracker was calibrated by using a 5-point calibration

script and Tobii SDK calibration algorithms. We repeated the calibration two times if one or more

calibration points were missing or inadequate. We used the final calibration in case none of the

calibrations produced an adequate result.

The tasks (i.e., oculomotor orienting and attention disengagement) were both divided in two

blocks that were presented in alternate order starting with the first block of the oculomotor orienting

task, followed by the first disengagement block, and ending with the second oculomotor orienting

and disengagement blocks. Both oculomotor orienting blocks included 16 trials and the

disengagement blocks included 24 trials (i.e., a total of 32 and 48 trials in the oculomotor orienting

and disengagement tasks, respectively). Cheerful music with smiling children was played between

the blocks. Stimulus presentation was controlled by an open-source software written in Python

programming language (https://github.com/infant-cognition-tampere/drop), together with Psychopy

functions and a Tobii SDK plug-in. For both tasks, all data processing from the raw *x-y* gaze

position coordinates to the parameters reflecting oculomotor orienting and disengagement were

completed automatically using gazeAnalysisLib, a library of MATLAB routines for offline analysis

of raw gaze data [41].

Oculomotor orienting. Each trial in the oculomotor orienting task started with a 1000-ms

blank screen followed by an attention-grabbing animation presented on the center of the screen. The

animation was a checkerboard pattern and it rotated into its mirror image after each 250 ms. The

pattern remained visible at least 250 ms until the infant looked at it. When the infant looked at the

central cue, the cue disappeared and the target (a checkerboard pattern) appeared randomly in one

of the four corners of the monitor. The target remained visible 750 ms after the infant's point of

gaze was first recorded in the target area. After that, a picture of a cartoon image or a toy combined

by a joyful audio stimulus was presented on the center of the screen as a rewarding stimulus. The

eye tracking data was analyzed with the gazeAnalysisLib [42] to determine the latency of infants'

orienting response (saccadic reaction time) from the central to the target stimulus. Gaze shifts away

from the screen and reaction times longer than 1000 ms were excluded.

Infants with 4 or more scorable oculomotor orienting trials were included. The criteria for

included trials were: adequate fixation on the central stimulus (i.e., >70 % of the time) during the

time preceding a gaze shift, sufficient amount of non-missing data samples (i.e., no gaps longer than

200 ms), and valid information about the time of the gaze transition from the central to the

peripheral stimulus (i.e., the transition did not occur during a period of missing data). Trials with

gaze shifts within a period that started 150 ms after the onset of the peripheral stimulus and ended

1000 ms after the lateral stimulus onset were used to calculate mean oculomotor orienting latency

for each participant [42].

Attention disengagement. Attention disengagement was measured with a gaze-contingent

Overlap paradigm, which has been commonly used for measuring infant attention disengagement

from various stimuli (e.g., Peltola et al., 2008, 2018). First, a fixation point was presented on the

center of the screen until the infant looked at it for 500 ms. After that, a happy or a fearful face, or a

non-face stimulus (a phase-scrambled rectangle containing the amplitude and color spectra of the

original face stimuli) appeared on the center of the screen for 1000 ms. A distractor stimulus was

then presented on the left or right side of the screen parallel to the central stimulus for 500 ms. At

250 ms the distractor stimuli changed to its mirror image. After 500 ms, the distractor stimulus

disappeared and the central stimulus remained on the screen for 1000 ms, followed by a 1000-ms

blank screen before the next trial. The central stimulus conditions (happy or fearful face, or the non-

face stimulus) were presented in random order with the restriction that the same stimulus condition

was not repeated more than 3 times in a row.

The corneal-reflection eye-tracker measured the reaction times for gaze shifts between the

central stimulus and the distractor stimulus. A *dwell time index* reflecting attention disengagement

was calculated if there were at least 3 scorable trials in each stimulus condition. We calculated the

dwell time index using the trials with a gaze shift and trials without a gaze shift, excluding non-

scorable trials, and the time interval from the shortest to the longest acceptable saccadic reaction

time. Trials with insufficient fixation on the central stimulus (i.e., >70 % of the time) during the

time preceding a gaze shift or the end of the analysis period, > 200 ms gaps of valid gaze data, or

invalid information about the gaze shift (i.e., with the eye movement occurring during a period of

missing gaze data) were excluded from the analyses. For the included trials, attention dwell time

on the central stimulus was determined for the period starting 150 ms from the onset of the

distractor stimulus and ending 1000 ms after distractor onset. The duration was then converted to

a normalized dwell time index score by using the following formula:

$$\text{Dwell time index} = \frac{\sum_{i=1}^{n} \left(1 - \frac{1000 - x_i}{850}\right)}{n},$$

where x is the time point of the saccadic eye movement (i.e., last gaze point in the area of the

central stimulus preceding a saccade toward the distractor stimulus) and n is the number of

scorable trials in a given stimulus condition. In this index, the shortest acceptable saccadic

reaction time (150 ms) results in 0, and the longest possible saccadic reaction time (or lack of

saccade, which is equal to the last measured time point of the central stimulus at 1000 ms) results

in a score of 1 [41]. The dwell time index was calculated as the mean of the stimulus conditions

(happy or fearful face, or the non-face stimulus) from the accepted trials, and used as the dependent

variable in the analyses of attention disengagement. The inclusion criterion was a minimum of 3

acceptable trials in each stimulus condition.

Eye tracking assessments in Cape Town

Similar to the assessments in Tampere, infants were assessed in a dimly lit room at a private

health clinic or a state hospital. A custom MATLAB script (Mathworks, Natick, MA), running on

MacBook OS (Apple Inc., CA), and interfacing via Psychtoolbox and Tobii SDK plug-in with the

eye tracking system was used for stimulus presentation and data collection (see Pyykkö et al., 2019

for full description of the tests). Eye tracking data were collected by using a Tobii X2-60 or TX60

screen-based eye tracking system (Tobii Technology, Stockholm, Sweden).

The procedure for eye tracker calibration was similar for that used in the study in Tampere.

Oculomotor orienting was measured with a visual search task (adapted from [43] and explained in

detail in [44]). In this task, an image of a red apple (5° visual angle) was presented on the center of

the screen, accompanied by an *oh* sound. After the infant had looked at the stimulus and 2000 ms

had elapsed (or a maximum wait period of 4000 ms had elapsed), the infant saw a blank screen for

500 ms, followed by the re-appearance of the apple in a randomly chosen location. The apple was

presented alone (*one-object condition*), among four or eight identical distractors (*multiple-objects*

condition), or among four to eight different types of distractors (*conjunction condition*). There was

a total of eight trials per condition. To obtain a similar measure for oculomotor orienting as in the

Tampere cohort, we extracted orienting latencies from the data collected in the one-object

condition, and averaged these latencies for each child to obtain a measure of oculomotor orienting.

385   In the **disengagement task**, infants saw two stimuli with a 1000 ms onset asynchrony. The

first was a picture of a non-face pattern (as above) or a face of a South African female expressing

happiness or fear. The ethnicity of the model matched that of the child. The second stimulus was a

geometric shape (black and white circles or a checkerboard pattern) laterally on the left or right side

of the screen, which was superimposed by a still picture of the first frame of a child-friendly

cartoon animation. When the infant shifted gaze to the lateral image or 2000 ms had elapsed, still

picture turned into a dynamic cartoon for 4000 ms. Infants saw a total of 16 trials with non-faces

and 16 with faces. As above, the dwell time indices were calculated separately for each of the

stimulus conditions by using an automated script.

Statistical Analyses

395   We performed the statistical analyses with SPSS version 25.0 (IBM Corp., Armonk, NY).

The alpha level for statistical significance was set at .05. All the variables were normally distributed

according to Kolmogorov-Smirnov tests. We studied the associations between infant cry and later

visual attention with Pearson's two-tailed correlation coefficients. In the Tampere cohort, we

conducted additional analyses to control for the location of the cry sample in the cry bout by

including only those infants whose cry sequence was from the beginning of a bout. In the Cape

Town cohort, we conducted an additional analysis to control for the possible influence of prenatal

SSRI medication by including only those infants without prenatal SSRI exposure. In the Tampere

cohort there were no infants with SSRI exposure and in the Cape Town cohort all the cry samples

were from the beginning of a cry bout.

405 **Results**

406
As can be observed from Table 2, the analysis within the Tampere cohort showed a marginal

positive correlation between F0 and attention disengagement. When F0 was higher, the dwell time

index was larger, i.e., the infants were slower to shift gaze away from the central stimulus to the

distractor stimulus. A similar but a larger correlation was observed in the additional analysis

including only those infants whose cry sequence was captured from the beginning of a bout. A

similar correlation between F0 and attention disengagement was, however, not found in the Cape

Town cohort, neither in the primary analyses nor in the analysis where infants with exposure to

prenatal SSRI medication were excluded.

There were no statistically significant correlations between F0 and oculomotor orienting.

Further on, there were no statistically significant correlations between F0 variability and either of

the attentional outcome variables. The correlations are presented in table 2.

Table 2. Correlations (Pearson's *r*) between the F0 parameters and attention outcomes in the two cohorts.

		Oculomotor orienting	Attention disengagement
Tampere	F0	*r* = .047, *p* = .741	*r* =.288, *p* = .052
	F0var	*r* = -.102, *p* = .471	*r* = -.048, *p* = .750
Tampere, from the beginning of the cry	F0	*r* = .249, *p* .194	*r* = .497, *p* = .008
	F0var	*r* = .115, *p* = .553	*r* = -.041, *p* = .840
Cape Town	F0	*r* = -.053, *p* = .726	*r* = .160, *p* = .252
	F0var	*r* = -.141, *p* =.343	*r* = .038, *p* = .785
Cape Town, prenatal SSRI excluded	F0	*r* = -.077, *p* = .652	*r* = .141, *p* = .373
	F0var	*r* = -.073, *p* = .668	*r* = -.075, *p* = .636

Discussion

The present study is the first to explore the relation of infant cry acoustics to visual attention during the first year. It integrates two previously separate research lines of infant cry analysis and eye-tracking based measurement of infant visual attention. Saccadic reaction times during infancy have been suggested to be a basic cognitive skill which mediates the development of more advanced

cognitive functions [7]. Using these methods, we studied whether higher and more unstable

fundamental frequency (F0) of neonatal cry would be connected to visual attention measured with

eye-tracking in later infancy.

For the most part, the results did not indicate that the F0 of neonatal cry is associated with

visual attention at 6 to 8 months of age. The only exception was an association between higher

neonatal F0 and slower attention disengagement from central stimuli to peripheral distractor stimuli

at 8 months of age in the Tampere cohort. This result was replicated when we controlled for the

location of the analyzed cry sample within the cry bout, with a similar correlation observed when

the cry was captured from the beginning of the cry bout. The same association was not, however,

observed in the Cape Town cohort where the cry was captured at 6 weeks and attention

disengagement at 6 months. Furthermore, oculomotor orienting turned out to have no correlation

with F0 in either sample. Finally, the variation of F0 did not correlate with the eye-tracking

outcomes in any of the analyses.

The lack of correlation between neonate cry and oculomotor orienting in the current cohorts

could be due to the early maturation of oculomotor orienting, which could result in rather low

variability in saccadic reaction times in infants from typically developed cohorts at 6 and 8 months.

Indeed, the variability in oculomotor orienting times were moderately low in the current cohorts,

and the variability of saccadic reaction times in the attention disengagement task was significantly

higher than variability in the oculomotor orienting task. Both F0 and oculomotor orienting have

been indicated to reflect later cognitive functions, and the brainstem and vagal nerve to have

important roles in their regulation. A distinctive difference between the two attention

measurements, i.e., oculomotor orienting and disengagement, is that in the disengagement task,

attention needs to be disengaged from a stimulus at fixation, whereas in the oculomotor orienting

task, shifts of attention indicate a reaction to a new stimulus appearing on a blank screen. This

difference may render the disengagement task cognitively more demanding. Matsuzawa et. al. [33]

observed significant differences in the maturation of these two components of attention.

Disengagement times decreased markedly from 2 ½ months to 6 months of age, and changed only

little from 6 months up to 1 year. In contrast, oculomotor orienting seemed to have matured before

the first measurement at 2 ½ months and showed only minor changes within the studied age period.

Additionally, saccadic reaction times in the disengagement task at 2 ½ months were approximately

double compared to saccadic reaction times in the oculomotor orienting task, and approached the

speed of oculomotor orienting at 6 months. Thus, differences in the maturational timing of the two

components of attention may have influenced our results, considering that the eye-tracking

registrations were captured at 6 or 8 months. Future studies may benefit from measuring

oculomotor orienting at an earlier age.

Nevertheless, the difference between the maturation of oculomotor orienting and

disengagement does not explain why there was a correlation between F0 and attention

disengagement only in the Tampere cohort but not in the Cape Town cohort. Considering the lack

of correlation between F0 and attention disengagement in the Cape Town cohort we will first pay

attention to the age differences at the time of cry recording between the cohorts. Secondly, we

discuss the potential influence of early caregiver-infant interaction on the observed pattern of

results.

There is a consensus that early cry reflects the development of the central nervous system

[4,10]. The regulation of cry goes through prominent changes within the first months of life.

Neonatal cry is reflex-like, regulated by the brainstem and not dependent on higher cortical areas,

the influence of which increase with age [3]. Due to the intense early development of cry

regulation, it is thus possible that the variation in cry acoustics at 6 weeks of age does not reflect

activity of areas important for visual attention (e.g., brainstem and vagal nerve) to the same extent

as cry captured at the neonatal age. Further on, it has been detected that F0 changes as a function of

age in normal development. F0 decreases in the first days and weeks, which is usually attributed to

the growth of vocal cords. Decrease of F0 is followed by an increase of F0, which is suggested to

relate to progressive control of vocalizations due to neurological maturation. However, the exact

time course of this process is not clear [45,46]. Consequently, the prominent changes in the

regulation of cry as well as normal age-related changes in F0 during the first weeks and months of

life highlight the importance for standardizing the age of recording data for cry analysis. For future

studies, we suggest recording cry samples at the neonatal period, as this will also make the data

comparable to the earlier studies showing associations between neonatal F0 and later cognitive

development [11,35] .

Likewise, interaction with the social and physical environment has the potential to influence

infants' cognitive development and cry acoustics. Studies with neonates have shown that the

amount and the acoustic qualities of cry may vary as a function of caregiver-infant interaction

[47,48], with tactile stimulation and reinforcement learning playing a significant role in the

variation of early cry [49]. Consequently, infants learn to adjust their cry to the most effective level

according to situational factors. In the current study, the age difference between the cry recordings

in Tampere (0-5 days) and Cape Town (6 weeks) may be considered a significant timespan in

postnatal development. Additionally, a proportion of Cape Town mothers had psychiatric problems

such as depression and anxiety disorders, which are known to influence to mother-child interaction

and the development of the child [50-52], and may have further influenced the results. Thus, in

future studies, in addition to standardizing the age of cry recordings, measures of early interaction

quality are likely important.

The other cry feature that has been associated with later development in previous studies,

i.e., variation of fundamental frequency (F0var), was not related to the eye-tracking outcomes. A

vast amount of studies has observed more varying F0 among preterm and/or low birthweight

infants, infants with prenatal risk factors, as well as infants with severe medical or developmental

abnormalities. Instead, autism spectrum disorder has been associated with reduced variation in F0.

While it is possible that the absence of associations between F0 variation and the outcomes was due

to the infants in our study being full term, with normal birth weight, and without any known health

or developmental problems, it should be noted that two earlier studies [11,35] detected that more

varying F0 was related to lower scores in cognitive outcome measures at several ages even after

controlling for the risk factors, including prematurity and prenatal exposure to drugs in the cohorts.

As there is currently a little methodological consensus about the determination of F0var, an

important line for future studies is to examine the predictive significance of different possibilities in

determining F0var, such as index variables or micro-variation measures, e.g., fluctuation or

interquartile range.

The influence of the trigger of the cry and the location of the analyzed cry sample in

cry bout need to be considered further. Our additional analysis in the Tampere cohort suggests that

the correlation between F0 and attention disengagement is stronger when the cry is captured from

the beginning of the cry bout. In a cry triggered by a sudden pain, the F0 is high in the beginning

and decreases by time. In contrast, in an inconvenience cry (e.g., hunger, cold, or loneliness) the F0

is lower in the beginning and increases by time if the response to the cry is delayed [53,54] In our

study the trigger in the Cape Town cohort was either a vaccination or a weight measurement on a

scale. In Tampere, the triggers ranged from venipuncture to unknown inconvenience. However, the

venipuncture did not trigger a cry in all cases, and there were no other triggers in Tampere which

could be considered as painful. In general, attempts to make a taxonomy of different cry triggers

face prominent challenges, because evaluating the trigger is more or less subjective except in case

of strong pain caused by an external operator. Future studies could benefit from measuring infants'

arousal levels during cry paradigms (e.g., with heart rate or skin conductance measurements) as

autonomic arousal is known to influence F0 [55,56] and correlate with the experience of pain and

inconvenience [57] .

The reliability of the correlation estimates in the current study was limited by the number of

infants included in the analyses, and it will be important to replicate the findings with a larger

sample. Additionally, the tasks in measuring oculomotor orienting and attention disengagement

were not completely identical in Tampere and Cape Town, which may have influenced the results.

Notwithstanding these limitations, a clear strength of the study is the integration of two previously

separate research lines of infant cry analysis and eye-tracking based measurement of infant visual

attention. Both methods enable registrations at a very early age which is a prominent advantage in

evaluating early cognitive development. Methodological development (e.g., Naithani et al., 2018)

has also enabled reliable segmentation and acoustic analyses of cry sounds captured in various

settings, broadening the possibilities of neonatal cry analysis.

Monitoring early cognitive development is an important, yet challenging part of pediatric

practice, which increases the need for applicable methods that can be administered at an earlier age

than more elaborate psychological assessments. Research on the potential of using infant cry to

predict cognitive development is still scarce and often conducted with small samples. In the present

study with typically developing infants, the results suggested a small association between neonatal

cry F0 and later visual attention disengagement. Whether variations in infant cry characteristics are

sufficiently distinct in typically developing populations to be used for monitoring cognitive skills

requires further research. Nevertheless, detecting normative variation in infant cognition is vitally

important especially for identifying children with milder forms of atypical cognitive development.

To reach this population for appropriate support, rehabilitation and health care as early as possible

would be especially important.

548
549

References

1. Colombo J. Infant cognition: Predicting later intellectual functioning: Sage publications; 1993.

2. McCall RB. The development of intellectual functioning in infancy and the prediction of later IQ.
Handbook of infant development. 1979: 707-741.

3. Newman JD. Neural circuits underlying crying and cry responding in mammals. Behav Brain
Res. 2007;182: 155-165.

4. LaGasse LL, Neal AR, Lester BM. Assessment of infant cry: acoustic cry analysis and parental
perception. Ment Retard Dev Disabil Res Rev. 2005;11: 83-93.

5. Lester BM. Developmental outcome prediction from acoustic cry analysis in term and preterm
infants. Pediatrics. 1987;80: 529-534.

6. Colombo J. Infant attention grows up: The emergence of a developmental cognitive neuroscience
perspective. Current Directions in Psychological Science. 2002;11: 196-200.

7. Dougherty TM, Haith MM. Infant expectations and reaction time as predictors of childhood
speed of processing and IQ. Dev Psychol. 1997;33: 146.

8. Rose SA, Feldman JF, Jankowski JJ. The building blocks of cognition. J Pediatr. 2003;143: 54-
61.

9. Rose SA, Feldman JF, Jankowski JJ. Implications of infant cognition for executive functions at
age 11. Psychological science. 2012;23: 1345-1355.

10. Lester BM. Developmental Outcome Prediction From Acoustic Cry Analysis in Term and
Preterm Infants. Pediatrics. 1987;80: 529-534.

11. Huntington L, Hans SL, Zeskind PS. The relations among cry characteristics, demographic
variables, and developmental test scores in infants prenatally exposed to methadone. Infant
Behavior and Development. 1990;13: 533-538. doi: //doi.org/10.1016/0163-6383(90)90021-Y.

12. Michelsson K, Michelsson O. Phonation in the newborn, infant cry. Int J Pediatr
Otorhinolaryngol. 1999;49, Supplement 1: S301.

13. Esposito G, Del Carmen Rostagno M, Venuti P, Haltigan JD, Messinger DS. Brief Report:
Atypical expression of distress during the separation phase of the strange situation procedure in
infant siblings at high risk for ASD. J Autism Dev Disord. 2014;44: 975-980.

14. Sheinkopf SJ, Iverson JM, Rinaldi ML, Lester BM. Atypical cry acoustics in 6-month-old
infants at risk for autism spectrum disorder. Autism Res. 2012;5: 331-339.

15. Porter FL, Porges SW, Marshall RE. Newborn pain cries and vagal tone: parallel changes in
response to circumcision. Child Dev. 1988: 495-505.

16. Lester BM. Spectrum analysis of the cry sounds of well-nourished and malnourished infants.
Child Dev. 1976: 237-241.

17. Vohr BR, Barry L, Rapisardi G, O'Dea C, Brown L, Peucker M, et al. Abnormal brain-stem
function (brain-stem auditory evoked response) correlates with acoustic cry features in term infants
with hyperbilirubinemia. The Journal of Pediatrics. 1989;115: 303-308. doi:
//doi.org/10.1016/S0022-3476(89)80090-3.

18. Lester BM, Dreher M. Effects of marijuana use during pregnancy on newborn cry. Child Dev.
1989: 765-771.

19. Colombo J. The development of visual attention in infancy. Annu Rev Psychol. 2001;52: 337-
367.

20. Geva R, Feldman R. A neurobiological model for the effects of early brainstem functioning on
the development of behavior and emotion regulation in infants: Implications for prenatal and
perinatal risk. Journal of Child Psychology and Psychiatry. 2008;49: 1031-1041.

21. Gazzaniga MS. The cognitive neurosciences: MIT press; 2004.

22. Geva R, Schreiber J, Segal-Caspi L, Markus-Shiffman M. Neonatal brainstem dysfunction after
preterm birth predicts behavioral inhibition. Journal of Child Psychology and Psychiatry. 2014;55:
802-810.

23. Hunnius S, Geuze RH. Gaze shifting in infancy: A longitudinal study using dynamic faces and
abstract stimuli. Infant Behavior and Development. 2004;27: 397-416.

24. Gardner JM, Karmel BZ, Flory MJ. Arousal modulation of neonatal visual attention:
Implications for development. Perspectives on fundamental processes in intellectual functioning.
2003;2: 125-153.

25. Karmel BZ, Gardner JM, Freeland RL. Arousal-modulated attention at four months as a
function of intrauterine cocaine exposure and central nervous system injury. J Pediatr Psychol.
1996;21: 821-832.

26. Geva R, Dital A, Ramon D, Yarmolovsky J, Gidron M, Kuint J. Brainstem as a developmental
gateway to social attention. Journal of Child Psychology and Psychiatry. 2017;58: 1351-1359.

27. Green CR, Lebel C, Rasmussen C, Beaulieu C, Reynolds JN. Diffusion tensor imaging
correlates of saccadic reaction time in children with fetal alcohol spectrum disorder. Alcoholism:
Clinical and Experimental Research. 2013;37: 1499-1507.

28. Hunnius S, Geuze RH, Zweens MJ, Bos AF. Effects of preterm experience on the developing
visual system: A longitudinal study of shifts of attention and gaze in early infancy. Dev
Neuropsychol. 2008;33: 521-535.

29. Elsabbagh M, Fernandes J, Webb SJ, Dawson G, Charman T, Johnson MH. Disengagement of
visual attention in infancy is associated with emerging autism in toddlerhood. Biol Psychiatry.
2013;74: 189-194.

30. Peltola MJ, Leppänen JM, Palokangas T, Hietanen JK. Fearful faces modulate looking duration
and attention disengagement in 7-month-old infants. Developmental Science. 2008;11: 60-68.

31. Peltola MJ, Yrttiaho S, Leppänen JM. Infants' attention bias to faces as an early marker of social
development. Developmental science. 2018;21: e12687.

32. Hood BM, Atkinson J. Disengaging visual attention in the infant and adult. Infant Behavior and
Development. 1993;16: 405-422.

33. Matsuzawa M, Shimojo S. Infants' fast saccades in the gap paradigm and development of visual
attention. Infant behavior and Development. 1997;20: 449-455.

34. Rafal R, Robertson L. The neurology of visual attention. . 1995.

35. Lester BM. Developmental outcome prediction from acoustic cry analysis in term and preterm
infants. Pediatrics. 1987;80: 529.

36. Schuster-Böckler B, Bateman A. An introduction to hidden Markov models. Current protocols
in bioinformatics. 2007;18: A. 3A. 9.

37. Naithani G, Kivinummi J, Virtanen T, Tammela O, Peltola MJ, Leppänen JM. Automatic
segmentation of infant cry signals using hidden Markov models. EURASIP Journal on Audio,
Speech, and Music Processing. 2018;2018: 1.

38. Fuamenya NA, Robb MP, Wermke K. Noisy but effective: crying across the first 3 months of
life. Journal of voice : official journal of the Voice Foundation. 2015;29: 281-286.

39. Reggiannini B, Sheinkopf SJ, Silverman HF, Li X, Lester BM. A flexible analysis tool for the
quantitative acoustic assessment of infant cry. J Speech Lang Hear Res. 2013;56: 1416-1428.

40. Gredebäck G, Johnson S, von Hofsten C, Humanistisk-samhällsvetenskapliga
vetenskapsområdet, Samhällsvetenskapliga fakulteten, Uppsala universitet, et al. Eye Tracking in
Infancy Research. Dev Neuropsychol. 2010;35: 1-19.

41. Leppänen JM, Forssman L, Kaatiala J, Yrttiaho S, Wass S. Widely applicable MATLAB
routines for automated analysis of saccadic reaction times. Behavior Research Methods. 2015;47:
538-548.

42. Leppänen JM, Forssman L, Kaatiala J, Yrttiaho S, Wass S. Widely applicable MATLAB
routines for automated analysis of saccadic reaction times. Behavior Research Methods. 2015;47:
538-548.

43. Kaldy Z, Kraper C, Carter AS, Blaser E. Toddlers with autism spectrum disorder are more
successful at visual search than typically developing toddlers. Developmental science. 2011;14:
980-988.

44. Pyykkö J, Forssman L, Maleta K, Ashorn P, Ashorn U, Leppänen JM. Early development of
visual attention in infants in rural Malawi. Developmental science. 2018: e12761.

45. Baeck HE, de Souza MN. Longitudinal Study of the Fundamental Frequency of Hunger Cries
Along the First 6 Months of Healthy Babies. Journal of Voice. 2007;21: 551-559.

46. Gilbert HR, Robb MP. Vocal fundamental frequency characteristics of infant hunger cries: birth
to 12 months. Int J Pediatr Otorhinolaryngol. 1996;34: 237-243.

47. Belsky J, Rovine M, Taylor DG. The Pennsylvania Infant and Family Development Project, III:
The Origins of Individual Differences in Infant-Mother Attachment: Maternal and Infant
Contributions. Child Dev. 1984;55: 718-728.

48. Lotem A, Winkler DW. Can reinforcement learning explain variation in early infant crying?
Behav Brain Sci. 2004;27: 468. doi: 10.1017/S0140525X04310101.

49. Cecchini M, Lai C, Langher V. Communication and crying in newborns. Infant behavior and
development. 2007;30: 655-665.

50. Field T. Postpartum depression effects on early interactions, parenting, and safety practices: a
review. Infant Behavior and Development. 2010;33: 1-6.

51. Field TM. Early interactions between infants and their postpartum depressed mothers. Infant
Behavior & Development. 1984.

52. Bakeman R, Brown JV. Early interaction: Consequences for social and mental development at
three years. Child Dev. 1980: 437-447.

53. Green JA, Gustafson GE, McGhie AC. Changes in infants' cries as a function of time in a cry
bout. Child Dev. 1998;69: 271-279.

54. Zeskind PS, Sale J, Maio ML, Huntington L, Weiseman JR. Adult Perceptions of Pain and
Hunger Cries: A Synchrony of Arousal. Child Dev. 1985;56: 549-554.

55. Porter FL, Porges SW, Marshall RE. Newborn pain cries and vagal tone: parallel changes in
response to circumcision. Child Dev. 1988: 495-505.

56. Shinya Y, Kawai M, Niwa F, Myowa-Yamakoshi M. Preterm birth is associated with an
increased fundamental frequency of spontaneous crying in human infants at term-equivalent age.
Biology letters. 2014;10: 20140350.

57. Stevens B, McGrath P, Gibbins S, Beyene J, Breau L, Camfield C, et al. Determining
behavioural and physiological responses to pain in infants at risk for neurological impairment. Pain.
2007;127: 94-102.

683