Title: Community turnover by composition and climate affinity across scales in an alpine system

Authors: Brian V. Smithers[1,2,3],*, Meagan F. Oldfather[1,4],*, Michael J. Koontz[1,3,5], Jim Bishop[1], Catie

Bishop[1], Jan Nachlinger[1], Seema N. Sheth[1,6]

Affiliations:

[1]Gloria Great Basin, Oroville, CA, USA

[2]Department of Ecology, Montana State University, Bozeman, MT, USA

[3]Department of Plant Sciences, University of California, Davis, CA, USA

[4]Department of Ecology and Evolutionary Biology, University of Colorado, Boulder, CO, USA

[5]Earth Lab, University of Colorado Boulder, CO, USA

[6]Department of Plant and Microbial Biology, North Carolina State University, Raleigh, NC, USA

*Corresponding authors

Short title: Alpine plant community-climate relationships with elevation

ABSTRACT

Premise of the study: Examining community turnover across climate gradients at multiple scales is vital

to understanding biogeographic response to climate change. This approach is especially important for

alpine plants in which the relative roles of topographic complexity and non-climatic or stochastic factors

vary across spatial scales.

Methods: We examined the structure of alpine plant communities across elevation gradients in the

White Mountains, California. Using community climatic niche means (CCNMs) and measures of

community dissimilarity, we explored the relationship between community composition and elevation

gradients at three scales: the mountain range, individual peaks, and within elevation contours.

Key Results: At the mountain range scale, community turnover and CCNMs showed strongly significant

relationships with elevation, with an increase in the abundance of cooler and wetter-adapted species at

higher elevations. At the scale of a single peaks, we found weak and inconsistent relationships between

CCNMs and elevation, but variation in community composition explained by elevation increased. Within

the elevation contours, the range of CCNMs was weakly positively correlated with turnover in species

identity, likely driven by microclimate and other site-specific factors.

Conclusions: Our results suggest that there is strong environmental sorting of alpine plant communities

at broad scales, but microclimatic and site-specific, non-climatic factors together shape community

turnover at finer scales. In the context of climate change, our results imply that community-climate

relationships are scale-dependent, and predictions of local alpine plant range shifts are limited by a lack

of topoclimatic and habitat information.

INTRODUCTION

Mountain environments contain important ecosystems and are heralds of species distribution

shifts in response to climate change (Körner, 2003). Pronounced warming at higher elevations

exacerbates this potential for mountains to exhibit early signs of biotic responses to climate change

(Pepin et al., 2015). Species have responded by generally moving upslope in response to warming

(Parmesan and Yohe, 2003; Lenoir et al., 2008; Pauli et al., 2012; Millar et al., 2015; Smithers et al.,

2018; Malanson et al., 2019), with this upslope advance having accelerated in recent years (Steinbauer

et al., 2018). Mountain geometry generally, but not always, also means that species will face decreasing

land area as they track their climatic niche to higher elevations (Elsen and Tingley, 2015). However, it is

increasingly recognized that climate change expectations of wholesale upward or poleward shifts of

species distributions are overly simplistic (Lenoir and Svenning, 2015), especially in topographically

complex landscapes (Rapacciuolo et al., 2014). Further, alpine species, often cool-adapted and long-

lived, may be both directly sensitive to temperature change due to their physiology and indirectly

sensitive due to competitive exclusion by species recruiting from lower elevations (Alexander et al.,

2015; Rumpf et al., 2018). Predicting how these direct and indirect impacts of climate change will

together shape future communities requires a better understanding of how complex mountain

landscapes shape community assembly.

A long-standing debate in community ecology is whether community assembly is repeatable

and predictable. If environmental filtering shapes community assembly, then there should be repeated,

deterministic patterns of community composition across similar climate and geologic conditions

(Clements, 1916, 1936). On the other hand, less predictable, or stochastic, processes in the form of

chance colonization, random extinction, ecological drift, and dispersal limitation could be more

important in explaining patterns of community assembly (Gleason, 1927; Slatkin, 1974). While this

debate has often been waged as either deterministic or stochastic forces being the driver of community

assembly, it is more likely that communities are structured by both processes (Chase and Myers, 2011).

In the context of climate change, this debate can be further distilled to a debate about whether climatic

or non-climatic factors structure community assembly. Importantly, the relative importance of

deterministic climatic factors versus deterministic non-climatic or stochastic factors (hereafter referred

to as non-climatic factors) can change depending on the system or scale at which we examine it. In

alpine systems, the relative importance of these factors in shaping plant communities is particularly

unclear, and this knowledge gap limits our ability to predict the rate and magnitude of plant community

change in alpine systems in response to climate warming.

A first step to understanding the relative importance of these factors in shaping plant

communities is to examine how communities are currently distributed across multiple scales of

climatologic complexity. In mountain systems, scales for describing plant community response to

climate change can range from a single elevation at one location to an entire mountain range. Elevation

is a useful geographic feature when studying deterministic and stochastic effects on community

assembly because at different scales of elevation gradients, the relative importance of each process may

change. Across an entire mountain range, the physical isolation of mountaintops may amplify the

stochastic processes of dispersal and colonization as low-lying areas between suitable high-elevation

habitat can create geographic barriers to species movement (Dirnböck et al., 2011). However at the

scale of a single peak where dispersal is likely not limiting, the deterministic process of environmental

filtering may play a larger role with high community turnover across steep climate gradients driven by

elevation (Körner, 2003, 2007). Lastly, small-scale topographic complexity in the alpine zone can also

drive variation in community composition (Van de Ven et al., 2007; Scherrer and Körner, 2011; Lenoir et

al., 2013; Winkler et al., 2016). At this microclimatic scale, community variation within an elevation

contour may be high where topographic complexity is high; the variety of potential habitats may

deterministically facilitate close proximity of species and communities with a variety of climatic niches.

However, within an elevation contour, other factors such as the geology, habitat stability (e.g., scree

moves more than bedrock) or species interactions may also play a larger role (Graae et al., 2018).

Comparing different types of community characteristics allows us to disentangle how climatic

and non-climatic processes are shaping the community across scales. Community climatic niche affinities

are metrics of the community (similar to any community-weighted mean traits) derived from climatic

conditions found across the component species' regional distributions (Landolt et al., 2010; Scherrer and

Körner, 2011; Lenoir et al., 2013). When considering community climatic niche affinities, environmental

filtering processes may drive species turnover. Conversely, community similarity metrics based on

species composition alone may be more dependent on the stochastic order and timing of species

arrivals into the community (Fukami et al., 2005). Similar patterns of climatic niche affinities and

community composition across elevation for a single scale would indicate that deterministic processes

are driving much of the community turnover. For a counter example, if the range of climatic niche

affinities encompassed within elevation contours is not correlated with turnover in the community

composition (ie., where species turnover is high, the range of climatic niche affinity values among

species is low), this points to non-climatic factors shaping the community variation at this scale. These

types of comparisons could be scaled up to look at different elevation bands on a single peak, among

peaks in a mountain range, or even among mountain ranges.

Management of large areas of public lands for resilience with a changing climate challenges us

to understand how communities will shift both locally and regionally (Pecl et al., 2017). Examining

relationships between plant communities and climate gives key insights into the potential for species

and communities to respond to climate change by tracking their climatic niche across different scales

and may inform whether moving upslope in elevation is a predictable method for species to remain in

their climatic niche in response to climate change. Evaluating changes in biodiversity and species

distributional shifts requires long-term monitoring across multiple scales (Gottfried et al., 2012). The

Global Observation Research Initiative in Alpine Environments (GLORIA) is an international collaboration

assessing global distributional shifts of alpine plant species in response to changing climate

(http://www.gloria.ac.at/). This international program was founded to provide cost-effective, universal

monitoring protocols and a unifying network to investigate the rate and magnitude of the changes in

alpine communities through time and spatial patterning at local, regional, and global scales (Grabherr et

al., 2010). This network has led to key findings about the vulnerability of montane systems to changing

climate, including upward distributional shifts of many alpine species and loss of cool-adapted species in

mountains (Pauli et al., 2007, 2012; Gottfried et al., 2012; Lamprecht et al., 2018; Rumpf et al., 2018;

Steinbauer et al., 2018). The North American chapter was founded in 2004 (Millar and Fagre, 2007) with

installations in California and Montana, and now has installations in many mountain regions of North

America. As part of this effort, GLORIA Great Basin (www.gloriagreatbasin.org) established a series of

downslope transects on five peaks in the White Mountains, California (Fig. 1) extending from alpine

mountaintops into the subalpine zone to monitor community change in the local species pool at the

individual mountain and at the mountain range scale.

In this study, we evaluate the role of climatic versus non-climatic processes in explaining plant

community turnover across elevation gradients at three scales: across an entire mountain range, within

each of five peaks spread across the mountain range, and within elevation contours of each peak, in the

White Mountains, California, USA (Fig. 1). We use metrics of community climatic niche means and

community composition to examine relationships among species turnover, community niche affinities,

and elevation across scales. Consistent with environmental filtering, community niche affinities may be

highly dependent on the elevation gradient, with the presence and abundance of cooler and wetter-

adapted species (dependent on their climatic niche) increasing with elevation, especially with climate

metrics that vary the most predictably across the elevation gradient, such as temperature and

precipitation. Given that the relative importance of deterministic and stochastic processes may vary

with spatial scale and the level of community organization under consideration (Fukami et al., 2005), the

relationships between elevation and climatic niche affinities may be strongest at broader spatial scales,

while the relationship between raw community composition and elevation may be stronger at smaller

scales (i.e., within a single peak). Specifically, we propose the following hypotheses.

1) At broad spatial scales (mountain range), non-climatic or stochastic differences will be more

important in explaining variation in species composition and turnover. We predict that elevation

explains less variation in species composition and turnover across the entire range versus within each

peak.

2) At broad spatial scales, climatic factors as environmental filters will be more important in

explaining plant community variation in climatic niche space. We predict that elevation will be better at

predicting community climatic niche affinities at large spatial scales (mountain range) than at smaller

scales (individual peak).

3) At the smallest scales (within elevation contour on a single peak), microclimate, habitat

differences, and other unmeasured factors facilitate niche partitioning, thereby driving patterns of

community assembly. We predict a positive relationship between the range of community climatic niche

affinities and species turnover within elevation contours.

METHODS

Study Area—The White Mountains are a cold and dry mountain range along the California-

Nevada border at the western margin of the Great Basin in the eastern rain-shadow of the Sierra

Nevada (Powell and Klieforth, 1991). At the White Mountains Research Center's Barcroft Station (3800

m), mean annual precipitation is approximately 455 mm and mean annual temperature is -1.7 $^{\circ}$C

(Western Regional Climate Center). Most of the precipitation in the range comes as winter snow,

although monsoonal rain events also occur during the summer. The alpine zone ranges in elevation from

treeline at roughly 3500 m to the top of White Mountain Peak at 4300 m. The alpine flora in the White

Mountains is made up of obligate alpine species, lower montane species, and cold-desert species

associated with the Sierra Nevada, Great Basin, and Rocky Mountains (Rundel et al., 2008).

Field Surveys—We sampled alpine community composition along elevation gradients using 100-

meter belt transects on five peaks in the White Mountains: White Mountain Peak (WMT), Mount

Barcroft (BAR), Sheep Mountain East (SME), Patriarch Grove South (PGS), and Campito Mountain (CPT)

(Bishop, 2011). On each peak, the first transect was established on the southeast aspect (east aspect for

Campito Peak), 35 vertical meters below the summit. Two, 50m x 1m belt transects were extended from

this central point following the elevation contour to create a 100-meter x 1-meter belt transect at one

elevation. Additional 100m x 1m transects were established on each peak in intervals of 25 vertical

meters below the previous transect on the same aspect (Fig. 1). Transects were established across 3

consecutive years from 2011-2013 (Table 1). Each transect was divided into ten continuous, 10m x 1m

segments. In each segment, we recorded the presence of every species. We assessed cover of plant

species and substrate (rock, scree, litter, bare ground) along the entire transect using a point-intercept

method with 40 points per 10m x 1m segment. A pair of points was located at every half-meter along

the transect, with the points in each pair 0.5 m apart (0.25 m from the center of the belt transect). The

substrate or plant species was recorded as a "hit" if a point intersected with it.

Community Metrics—To investigate how communities varied across spatial gradients, we used

community-weighted climatic niche means (CCNM) and community composition. A CCNM is the

abundance-weighted mean of each component species' average species niche space across its range (De

Frenne et al., 2013). For community composition we used the Bray-Curtis dissimilarity metric of β

diversity, which incorporates both species presence and relative abundance (Anderson et al., 2011). Our

measure of abundance used for both metrics was the number of segments in which it was found for

each transect. We consider how these two community metrics change across three nested scales: the

elevation gradient of the entire mountain range, the elevation gradient of a single peak, and within a

single elevation contour of a peak.

A CCNM is a property of a single community, or in our specific case, a transect for mountain

range- and peak-scale analyses and a 10m segment within transects for the elevation contour-scale

analysis. Any climate metric can be used to quantify a community's CCNM. We focused on the following

climatic measurements as they are likely major structuring factors for alpine plant communities: climatic

water deficit (CWD), mean annual precipitation (MAP), July maximum temperature (July T_{max}), and

January minimum temperature (Jan T_{min}). CWD is an integrative measure of how much energy

availability (temperature) exceeds water supply (precipitation), with higher values indicating hotter,

drier conditions, and lower values indicating cooler, wetter conditions. Using a recently-compiled

database of California herbarium specimens derived from the Consortium of California Herbaria as well

as additional natural history museums and herbaria (Baldwin et al., 2017), we extracted locations from

all vouchered specimens in the database for each surveyed species. We used those database locations

to extract 30-year (1961-1990) mean climate values for CWD, MAP, July T_{max}, and Jan T_{min} from the

TerraClimate gridded dataset (Abatzoglou et al., 2018), which is itself an amalgamation of the WorldClim

dataset (Fick and Hijmans, 2017) and other data. After culling the herbaria records to one record per

species per grid cell to reduce sampling bias, we extracted the four mean climate values for each

species, representing the mean climatic niche of the species for each climate metric (Appendix S1; see

Supplemental Data with this article). This 30-year period was chosen as the temporal range of most

likely establishment for the majority of the generally long-lived individual plants sampled. For each

transect on each of the five peaks, we calculated a weighted mean of the mean climatic niche value of

each species present in that transect. The mean was weighted by the abundance of each species in that

community, in our case the number of segments in which it was found when the community was

defined as the whole transect and the number of point-intercept hits when the community was defined

as the 10m x 1m segments within the transect. The abundance weighting allows more dominant species

to have a larger effect on the community's CCNM. This final calculation produces one CCNM value for

each climate metric in each community, allowing for examination of how the overall climatic niche of

the community changes across the elevation gradients or ranges within elevation contours. For this

system, regional CCNM values are representative of local climatic niche means derived from

microclimate field data (Oldfather and Ackerly, in press).

218   To examine community patterns within the elevation contours we used two metrics to

represent how species identity and climatic niches vary across the contour: β diversity and CCNM range.

We calculated per-transect β diversity as the total species richness in each transect divided by the mean

species richness across the 10 segments within each transect. Therefore, the minimum possible value is

a β diversity of 1, which would indicate that every species in the whole transect is represented within

each segment of that transect. This metric gives a measure of the amount of species variation that can

be found within a single elevational band. The range of CCNM within a single transect was determined

by calculating the CCNM per segment, and then subtracting the minimum from the maximum of each

transect for each climate variable. The CCNM range gives a measure of how variable the climatic niches

of the communities are along the elevation contour; a low range indicates that elevation is

predominantly driving any variation in the climatic niche of the communities, and a high range indicates

that microclimate may be shaping the patterns of community climatic niches across the landscape as

well.

232   ***Statistical Analyses***—To examine how CCNMs changed across the entire mountain range, we

constructed linear mixed effects models for each of the four CCNMs (CWD, MAP, July T_{max}, and Jan T_{min})

as the response variable, elevation as the predictor variable, and with a random intercept effect of peak.

We compared models with and without a quadratic term for elevation and used Akaike's Information

Criterion (AIC) to choose the model with the best fit. For each of the four CCNMs, we also built linear

models for each peak with CCNM as the response variable and elevation as the predictor variable.

In order to quantify the community turnover explained by the elevational gradient at the scale

of the mountain range and all peaks separately, we performed permutation-based MANOVAs using the

'adonis' function (999 permutations) in the vegan package (Oksanen et al., 2008). We built separate

MANOVAs for the entire mountain range and for each of the peaks. For each of these models, the

dependent variable is a Bray-Curtis dissimilarity matrix (with the matrix rows representing single

transects) and the independent variable is elevation across either the entire mountain range or across

each peak, respectively. To visualize these relationships, we performed an ordination based on non-

metric dimensional scaling (NMDS) that included community data from each peak across the entire

White Mountain range.

To examine community composition and species turnover at the scale of each elevation

contour, we used linear models to determine whether the range in CCNM within an elevation contour

(transect) significantly explained variation in β diversity. All analyses were performed using R version

3.6.0 (R Core Team, 2018). Data manipulation and visualization were performed using the R packages

dplyr (Wickham et al., 2019), sf (Pebesma, 2018), and ggplot2 (Wickham, 2016).

RESULTS

Community Composition—We observed a total of 123 species in 70 genera and 25 families across the

entire mountain range, with individual peak species richness ranging from 50-69 (Table 1). Species

richness increased with elevation at PGS, and steeply declined with elevation at WMT, but did not vary

linearly with elevation on the remaining peaks (Appendix S2).

Across the mountain range, elevation explained 30% of the variation in community composition

($P < 0.001$, Fig. 2). When looking at each peak independently, there was an approximate 10-20%

increase in the community variation explained by elevation (BAR 42% $P < 0.001$, CPT 37% $P = 0.013$, PGS

40% $P < 0.001$, SME 42% $P < 0.001$, WMT 52% $P < 0.001$). The NMDS ordination visualizes these

relationships by reducing the community dimensionality into 2 dimensions. First, the ordinations show a

clear range-wide relationship between the first axis (NMDS1) and elevation (95% $P < 0.001$), as indicated

by the arrow. Therefore, species such as *Erigeron vagus* are associated with higher elevation sites

and *Erigeron tenor* is associated with lower elevation sites. Within the summits, the community

composition had stronger, but unique relationships with elevation; the ellipses that encompass the

standard deviation of points associated with each summit line up well with the first axis and vary across

the second axis (NMDS2).

Community-weighted Climatic Niche Mean—All CCNM metrics (CWD, MAP, Jan T_{min}, and July

T_{max}) showed highly significant linear relationships with elevation at the mountain range scale ($P < 0.001$

for all metrics, Fig. 3, Appendix S3). For each CCNM at the mountain range scale, the non-quadratic

model performed best. CWD, Jan T_{min}, and July T_{max} CCNMs showed a negative relationship with

elevation while annual precipitation (MAP) showed a positive relationship with elevation. Collectively,

these results indicate that the presence and abundance of species with cooler, wetter niches increased

with elevation. At the individual mountain scale, the relationships between the CCNM and elevation

were weak, inconsistent, or absent (Fig. 3, Table S2). On some of the peaks, the CCNM trend ran counter

to the range-wide trend, although those peak-scale countertrends were insignificant.

Variation in community composition and CCNMs across elevation contours—The mean β

diversity per transect across the elevation contours was 2.1 [range, 1.3-3.2]. A β diversity of 2

indicates that, on average across the segments of the transect, there are twice as many species

represented in the whole transect than in each segment. The mean ranges of CCNM per

transect across the elevation contours were CWD 173 mm [0 - 465], MAP 15.3 mm [0 - 47.9],

July T_{max} 1.61°C [0 - 3.88], and Jan T_{min} 2.01°C [0 - 5.75]. Interestingly, this variation is slightly

lower, but similar in magnitude as the range of CCNM values observed across the entire

mountain range (Fig. 3).

289   For CCNM MAP and July T_{max}, the CCNM range was significantly and positively correlated with

290   the per-transect β diversity (MAP: adjusted R^2 = 0.138, *P* = 0.003; July T_{max}: adjusted R^2 = 0.073, *P* =

291   0.027; Fig. 4). For CWD and Jan T_{min}, there was no significant relationship between CCNM range and per-

292   transect β diversity (adjusted R^2 = -0.016, *P* = 0.683; Jan T_{min}: adjusted R^2 = -0.014, *P* = 0.590).

294   DISCUSSION

295   Investigating the dynamics of community turnover across climate gradients is a fruitful avenue

for understanding where and how species distributions may shift with a changing climate (Scherrer and

Körner, 2011; Lenoir et al., 2013). Using a detailed alpine plant community dataset across steep

elevation and climate gradients, we investigated how climate associations shape species distributions

and community assembly across multiple scales over which range shifts have been predicted and

quantified. Across the entire mountain range, both community metrics (climate-derived CCNM and

composition-derived Bray-Curtis dissimilarity) exhibited strong relationships with elevation, but the

relationships between CCNM and elevation at the individual peak scale was weak and inconsistent (Figs.

2, 3). In contrast, there was a stronger relationship between elevation and community composition at

the scale of the individual peak relative to the entire mountain range (Fig. 2). Lastly, the range of

community climatic niche means was positively, but weakly correlated with the per-transect β diversity

(Fig. 4). Below, we discuss in more depth how the processes that shape alpine plant distributions and

community assembly vary across scales and explore potential impacts for using these community

metrics for predicting the effects of climate change on mountain communities.

Community Composition and Turnover Across Scales—Elevation was a strong predictor of

community dissimilarity and CCNM across the entire White Mountain range (Figs. 2, 3), suggesting that

elevation structures both community composition and niche affinities across broad spatial scales.

However, we found varying responses to elevation between the two community metrics at the scale of

individual peaks. Consistent with our first hypothesis, more variation in community composition was

explained by elevation at the individual peak scale relative to the entire range despite decreased

elevation range within each individual summit relative to the mountain range. At the mountain range

scale, non-climatic differences between peaks (e.g., biotic interactions or geology) or stochastic

differences likely reduce the community variation explainable by elevation. Community dissimilarity has

a strong, but unique relationship with elevation within each summit (Fig. 2), which muddles the broader-

scale relationship between elevation and dissimilarity across summits.

At the scale of individual peaks, the elevation gradient may encompass variation in additional

factors besides climate, such as available space, underlying changes in geology, biotic interactions, and

microtopography. Thus, these additional non-climatic factors may influence patterns of species

dissimilarity across the elevation gradients within each peak. Further, dispersal limitation is another

potentially important non-climatic factor that may weaken the relationship between elevation and

community composition at the scale of the entire mountain range (Dirnböck et al., 2011). On the one

hand, if stochastic processes such as dispersal effects explain community assembly, then species

dissimilarity should be invariant across environmental gradients such as elevation. On the other hand, if

deterministic processes such as environmental filtering are more important in shaping community

assembly, then species turnover should be invariant across spatial gradients (Chase and Myers, 2011).

Because elevation across the entire mountain range encompasses both spatial and climatic gradients

(Fig. 1; Appendix S4), then our ability to assess the relative importance of stochastic vs. deterministic

processes in shaping species turnover may be limited. Nonetheless, if dispersal limitation is reduced

within individual peaks relative to among peaks across the range, then variation in community

composition within peaks may be better explained by elevation as species are filtered with respect to

the climatic conditions across the gradient (Svenning and Sandel, 2013; Graae et al., 2018).

Climatic Niche Means Across Scales—If climatic filtering is primarily responsible for shaping

community assembly, then the presence and abundance of cooler, wetter-adapted species should

increase with elevation, particularly at the scale of the entire mountain range. Indeed, we found that

CCNMs based on all four climate metrics were significantly related to elevation at the mountain range

scale (Fig. 3). Encouragingly, these relationships at this scale all also matched the predicted patterns of

climatic niche means and elevation based on how the actual climate is predicted to vary across elevation

gradients (Appendix S4). Species associated with cooler and wetter conditions were found more often

and/or in higher abundance at higher elevations (Fig. 3). We also found support for our second

hypothesis that elevation should more strongly predict climatic niche affinities at large spatial scales

(mountain range) than at smaller scales (individual peak). Despite the strong, consistent relationship

between CCNMs and elevation across the entire mountain range, dependent on the peak and the

climate variable used to quantify CCNM, there were variable results for whether elevation was

predictive of community niche affinities for individual peaks. Therefore, at finer scales, elevation is less

consistently predictive of the climatic niche means of communities. This may be due to the reduced

range of elevation change, and therefore climatic variation, within each summit, allowing non-climatic

factors to play a larger role in community assembly. Within elevation contours, the range of community

climatic niche affinities influenced variation in species turnover with the CCNMs based on July maximum

temperatures and precipitation (Fig. 4), suggesting that for at least these climate metrics, microclimate

may facilitate niche partitioning and species coexistence at the within-contour scale. Future studies

should more formally examine the mechanisms of how microclimate drives patterns of community

assembly across these three spatial scales.

One important caveat of our use of CCNMs is that due to data availability, we quantified CCNMs

with species' occurrences from California distributions, not the species' global distributions. For many

species, California encompasses the range of climatic conditions that the species experience across their

ranges. For other species, however, the occurrences in California are a subset of the species' larger

distribution, and hence their climatic niches. Niche estimates are likely particularly inaccurate for species

whose western range limits are found in eastern California. Further, all CCNM calculations assumed no

variation among populations across species ranges and thus do not account for the possibility that the

populations surveyed in the White Mountains may have different climatic niches relative to those found

in other parts of the species range. CCNMs also disregard influences of fine-scale climate on species'

distributions due to the coarse resolution of the climatic and herbarium specimen data (Bramer et al.,

2018), potentially helping to explain the weaker and less consistent relationships found between

elevation and CCNMs within each peak relative to across the entire mountain range.

Drivers of Community Variation Within Elevation Contours—The maximum range in CCNMs observed

within a single elevation contour was on par with the overall maximum range of CCNM values observed

across the entire mountain range. This points to at least some role of microclimate in shaping species

distributions and community assembly. We found CCNM differences within elevation contours that

were equivalent to those found across 100 m in elevation! While in general, species are expected to

move upslope in response to warming, at the local scale there will likely be suitable microclimate within

the same elevation. Species movement may be lateral or even downslope in order to track newly

suitable microclimate. These findings are supported by research in alpine plant biology which has found

that microclimate has a large influence on species' distributions (Körner, 2003). However, the range of

community climatic niche means was positively, but weakly, correlated with the β diversity within

elevation contour (Fig. 4), indicating minor support for our third hypothesis. Therefore, other non-

climatic processes are also shaping the species turnover within the elevation contour. As dispersal

limitation at this scale is unlikely, two major potential factors that may be instead be shaping community

distributions are habitat type (e.g., presence of moveable scree) and species interactions. In addition to

microclimate, substrate type (rock, scree, organic soil) likely heavily influence species distributions, as

well as species' responses to climate change (Kulonen et al., 2018). Species interactions directly, or in

conjunction with habitat type and microclimate, can also influence species distributions at fine spatial

scales (Alexander et al., 2015; Blonder et al., 2018; Graae et al., 2018). For example, lower elevation

species may be more competitive on organic soils than on rock or scree whereas higher elevation

obligate alpine plants potentially have more adaptive traits (e.g., contractile roots) for persisting on the

dynamic scree habitat (Kulonen et al., 2018). As we build our temporal dataset, future studies will

examine whether interactions between habitat segregation and elevation influence species and

community responses to climate change over time.

Conclusions/Implication for Climate Change Research—Our findings suggest that the effects of

elevation on community niche affinities and community dissimilarity varies with spatial scale. At the

scale of the entire mountain range, there is climatic filtering across the elevation gradient, with an

increase in the presence and abundance of cool, wet-adapted species at higher elevations. Within

individual peaks and elevation contours, however, the signal of climatic filtering breaks down. This

points to differences in how climatic and non-climatic factors may shape species composition at

different scales. At broad spatial scales, deterministic climate appears to be a main driver of plant

community assembly since the large, consistent elevation gradient can mask stochastic factors such as

chance colonization and extinction and non-climatic factors like geology. At smaller scales where

dispersal is less limiting and microsite climatic variation is common even within the same elevation

contour, synoptic climatic factors appear less important than non-climatic factors in structuring

community. We did not directly measure those non-climatic factors, but they are likely to be related to

habitat and substrate type. Since climatic niche metrics are often used for quantifying community

responses to climate change, it is important to understand their relative strengths in explaining

community turnover across scales. The breakdown of correlations between CCNM and elevation at

finer-scales, even though species turnover based on identity is still explained by elevation at this scale,

suggests an important caveat for using climate-based community measures at these scales. Perhaps we

would not find this pattern if our climate data used to calculate the species climatic niches were at finer

resolutions, which is supported by the positive correlation per-transect β diversity and range of CCNM

within a transect. However, this correlation is weak indicating that higher resolutions may not be

sufficient to understand species distributions at these micro-topographic scales. Within the context of

climate change, predictions of local alpine plant range shifts will be limited by availability of

topoclimatic, as well as habitat, information.

Acknowledgements

The authors thank the UC Natural Reserve System's White Mountain Research Center, USFS

Pacific Southwest Research Station, Consortium for Integrated Climate Research in Western Mountains

(CIRMOUNT), the Global Observation Research Initiative in Alpine Environments (GLORIA), and USFS

Inyo National Forest for their support and collaboration. Our data collection teams consisted of many

citizen scientists that include graduate, undergraduate, and high school students as well as members of

NGOs such as GLORIA Great Basin, the California Native Plant Society, the Nevada Native Plant Society,

and the Davis Botanical Society. We especially thank Connie Millar and Adelia Barber for their leadership

and support of the early stages of this research. This work was supported by USDA National Institute of

Food and Agriculture Hatch project 1016272.

Author Contributions

Brian Smithers, Meagan Oldfather, and Michael Koontz participated in data collection, data processing,

analysis, and in the writing of the manuscript.

Jim Bishop and Catie Bishop created the downslope transect protocol, participated in data collection,

data processing, and in editing of the manuscript.

Jan Nachlinger participated in data collection, data processing, and in the editing of the manuscript.

Seema Sheth participated in data processing, analysis and in the writing of the manuscript.

Data Accessibility

Data and R code from the project are available on the GLORIA Great Basin website

(https://www.gloriagreatbasin.org/documents-archives) as well as in a permanent Figshare repository

(DOI to come).

Literature Cited

Abatzoglou, J. T., S. Z. Dobrowski, S. A. Parks, and K. C. Hegewisch. 2018. TerraClimate, a high-resolution

global dataset of monthly climate and climatic water balance from 1958–2015. *Scientific Data* 5:

170191.

Alexander, J. M., J. M. Diez, and J. M. Levine. 2015. Novel competitors shape species' responses to

climate change. *Nature* 525: 515–518.

Anderson, M. J., T. O. Crist, J. M. Chase, M. Vellend, B. D. Inouye, A. L. Freestone, N. J. Sanders, et al.

2011. Navigating the multiple meanings of β diversity: a roadmap for the practicing ecologist.

Ecology Letters 14: 19–28.

Baldwin, B. G., A. H. Thornhill, W. A. Freyman, D. D. Ackerly, M. M. Kling, N. Morueta-Holme, and B. D.

Mishler. 2017. Species richness and endemism in the native flora of California. *American Journal*

of Botany 104: 487–501.

Bishop, J. 2011. Using alpine flora to measure ecological effects of climate change. *Fremontia* 38: 2–9.

Blonder, B., R. E. Kapas, R. M. Dalton, B. J. Graae, J. M. Heiling, and Ø. H. Opedal. 2018.

Microenvironment and functional-trait context dependence predict alpine plant community

dynamics R. Salguero-Gómez [ed.],. *Journal of Ecology* 106: 1323–1337.

Bramer, I., B. J. Anderson, J. Bennie, A. J. Bladon, P. De Frenne, D. Hemming, R. A. Hill, et al. 2018.

Advances in Monitoring and Modelling Climate at Ecologically Relevant Scales. Advances in

Ecological Research, 101–161. Elsevier.

Chase, J. M., and J. A. Myers. 2011. Disentangling the importance of ecological niches from stochastic

processes across scales. *Philosophical Transactions of the Royal Society B: Biological Sciences*

366: 2351–2363.

Clements, F. E. 1936. Nature and Structure of the Climax. *Journal of Ecology* 24: 252–284.

Clements, F. E. 1916. Plant succession: an analysis of the development of vegetation. Carnegie

Institution of Washington, Washington, DC.

De Frenne, P., F. Rodríguez-Sánchez, D. A. Coomes, L. Baeten, G. Verstraeten, M. Vellend, M. Bernhardt-

Römermann, et al. 2013. Microclimate moderates plant responses to macroclimate warming.

Proceedings of the National Academy of Sciences 110: 18561–18565.

Dirnböck, T., F. Essl, and W. Rabitsch. 2011. Disproportional risk for habitat loss of high-altitude endemic

species under climate change. *Global Change Biology* 17: 990–996.

Elsen, P. R., and M. W. Tingley. 2015. Global mountain topography and the fate of montane species

under climate change. *Nature Climate Change* 5: 772–776.

Fick, S. E., and R. J. Hijmans. 2017. Worldclim 2: New 1-km spatial resolution climate surfaces for global

land areas. *International Journal of Climatology*.

Fukami, T., T. M. Bezemer, S. R. Mortimer, and W. H. van der Putten. 2005. Species divergence and trait

convergence in experimental plant community assembly. *Ecology Letters* 8: 1283–1290.

Gleason, H. A. 1927. Further Views on the Succession-Concept. *Ecology* 8: 299–326.

Gottfried, M., H. Pauli, A. Futschik, M. Akhalkatsi, P. Barančok, J. L. Benito Alonso, G. Coldea, et al. 2012.

Continent-wide response of mountain vegetation to climate change. *Nature Climate Change* 2:

111–115.

Graae, B. J., V. Vandvik, W. S. Armbruster, W. L. Eiserhardt, J.-C. Svenning, K. Hylander, J. Ehrlén, et al.

2018. Stay or go – how topographic complexity influences alpine plant population and

community responses to climate change. *Perspectives in Plant Ecology, Evolution and*

Systematics 30: 41–50.

Grabherr, G., M. Gottfried, and H. Pauli. 2010. Climate Change Impacts in Alpine Environments: Climate

change impacts in alpine environments. *Geography Compass* 4: 1133–1153.

Körner, C. 2003. Alpine Plant Life: Functional Plant Ecology of High Mountain Ecosystems. 2nd ed.

Springer-Verlag, Berlin Heidelberg.

Körner, C. 2007. Climatic treelines: conventions, global patterns, causes. *Erdkunde*: 316–324.

Kulonen, A., R. A. Imboden, C. Rixen, S. B. Maier, and S. Wipf. 2018. Enough space in a warmer world?

Microhabitat diversity and small-scale distribution of alpine plants on mountain summits J. Diez

[ed.],. *Diversity and Distributions* 24: 252–261.

Lamprecht, A., P. R. Semenchuk, K. Steinbauer, M. Winkler, and H. Pauli. 2018. Climate change leads to

accelerated transformation of high-elevation vegetation in the central Alps. *New Phytologist*

220: 447–459.

Landolt, E., B. Bäumler, A. Erhardt, O. Hegg, F. A. Klötzli, W. Lämmler, M. Nobis, et al. 2010. Flora

Indicativa: Ecological Indicator Values and Biological Attributes of the Flora of Switzerland and

the Alps / Ökologische Zeigerwerte und Biologische Kennzeichen zur Flora der Schweiz und der

Alpen. Haupt Verlag, Stuttgart, Vienna.

Lenoir, J., J. C. Gegout, P. A. Marquet, P. de Ruffray, and H. Brisse. 2008. A Significant Upward Shift in

Plant Species Optimum Elevation During the 20th Century. *Science* 320: 1768–1771.

Lenoir, J., B. J. Graae, P. A. Aarrestad, I. G. Alsos, W. S. Armbruster, G. Austrheim, C. Bergendorff, et al.

2013. Local temperatures inferred from plant communities suggest strong spatial buffering of

climate warming across Northern Europe. *Global Change Biology* 19: 1470–1481.

Lenoir, J., and J.-C. Svenning. 2015. Climate-related range shifts – a global multidimensional synthesis

and new research directions. *Ecography* 38: 15–28.

Malanson, G. P., L. M. Resler, D. R. Butler, and D. B. Fagre. 2019. Mountain plant communities:

Uncertain sentinels? *Progress in Physical Geography: Earth and Environment*:

0309133319843873.

Millar, C. I., and D. B. Fagre. 2007. Monitoring alpine plants for climate change: The North American

GLORIA project. *Mountain Views Chronicles* 1: 12–15.

Millar, C. I., R. D. Westfall, D. L. Delany, A. L. Flint, and L. E. Flint. 2015. Recruitment patterns and growth

of high-elevation pines to in response to climatic variability (1883-2013), western Great Basin,

USA. *Canadian Journal of Forest Research* 45: 1299–1312.

Oksanen, J., R. Kindt, P. Legndre, B. O'Hara, G. L. Simpson, P. Solymos, H. H. Stevens, and H. Wagner.

2008. The vegan Package.

Oldfather, M., and D. Ackerly. in press. Increases in Thermophilus Plants in an Arid Alpine Community in

Response to Experimental Warming. *Arctic, Antarctic, and Alpine Research* NA: NA.

Parmesan, C., and G. Yohe. 2003. A globally coherent fingerprint of climate change impacts across

natural systems. *Nature* 421: 37–42.

Pauli, H., M. Gottfried, S. Dullinger, O. Abdaladze, M. Akhalkatsi, J. L. B. Alonso, G. Coldea, et al. 2012.

Recent Plant Diversity Changes on Europe's Mountain Summits. *Science* 336: 353–355.

Pauli, H., M. Gottfried, K. Reiter, C. Klettner, and G. Grabherr. 2007. Signals of range expansions and

contractions of vascular plants in the high Alps: observations (1994–2004) at the GLORIA*

master site Schrankogel, Tyrol, Austria. *Global Change Biology* 13: 147–156.

Pebesma, E. 2018. Simple Features for R: Standardized Support for Spatial Vector Data. *The R Journal* 10:

439--446.

Pecl, G. T., M. B. Araújo, J. D. Bell, J. Blanchard, T. C. Bonebrake, I.-C. Chen, T. D. Clark, et al. 2017.

Biodiversity redistribution under climate change: Impacts on ecosystems and human well-being.

Science 355.

Pepin, N., R. Bradley, H. Diaz, M. Baraer, E. Caceres, N. Forsythe, H. Fowler, et al. 2015. Elevation-

dependent warming in mountain regions of the world. *Nature Climate Change* 5: 424–430.

Powell, D. R., and H. E. Klieforth. 1991. Weather and climate. *In* Hall, Clarence A. [ed.], Natural History of

the White-Inyo Range, Eastern California. California Natural History Guide,.

Rapacciuolo, G., S. P. Maher, A. C. Schneider, T. T. Hammond, M. D. Jabis, R. E. Walsh, K. J. Iknayan, et al.

2014. Beyond a warming fingerprint: individualistic biogeographic responses to heterogeneous

climate change in California. *Global Change Biology* 20: 2841–2855.

Rumpf, S. B., K. Hülber, G. Klonner, D. Moser, M. Schütz, J. Wessely, W. Willner, et al. 2018. Range

dynamics of mountain plants decrease with elevation. *Proceedings of the National Academy of*

Sciences: 201713936.

Rundel, P. W., A. C. Gibson, and M. R. Sharifi. 2008. The Alpine Flora of the White Mountains, California.

Madroño 55: 202–215.

Scherrer, D., and C. Körner. 2011. Topographically controlled thermal-habitat differentiation buffers

alpine plant diversity against climate warming: Topographical control of thermal-habitat

differentiation buffers alpine plant diversity. *Journal of Biogeography* 38: 406–416.

Slatkin, M. 1974. Competition and Regional Coexistence. *Ecology* 55: 128–134.

Smithers, B. V., M. P. North, C. I. Millar, and A. M. Latimer. 2018. Leap frog in slow motion: Divergent

responses of tree species and life stages to climatic warming in Great Basin subalpine forests.

Global Change Biology 24: e442–e457.

Steinbauer, M. J., J.-A. Grytnes, G. Jurasinski, A. Kulonen, J. Lenoir, H. Pauli, C. Rixen, et al. 2018.

Accelerated increase in plant species richness on mountain summits is linked to warming.

Nature 556: 231–234.

Svenning, J.-C., and B. Sandel. 2013. Disequilibrium vegetation dynamics under future climate change.

American Journal of Botany 100: 1266–1286.

Van de Ven, C. M., S. B. Weiss, and W. G. Ernst. 2007. Plant Species Distributions under Present

Conditions and Forecasted for Warmer Climates in an Arid Mountain Range. *Earth Interactions*

11: 1–33.

Wickham, H. 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag, New York.

Wickham, H., R. François, L. Henry, and K. Müller. 2019. dplyr: A Grammar of Data Manipulation.

Winkler, M., A. Lamprecht, K. Steinbauer, K. Hülber, J.-P. Theurillat, F. Breiner, P. Choler, et al. 2016. The

rich sides of mountain summits - a pan-European view on aspect preferences of alpine plants.

Journal of Biogeography 43: 2261–2273.

	Sample				#	Total
Peak	**Year**	**Elevation (m)**	**Soil Type**	**Aspect**	**Transects**	**Richness**
White Mountain Peak (WMT)	2012	3971-4296	Igneous	SE	14	50
Mount Barcroft (BAR)	2011	3625-3950	Igneous	SE	14	69
Sheep Mountain (SME)	2011	3475-3725	Dolomitic	SE	11	57
Campito Mountain (CPT)	2013	3325-3500	Quartzitic	E	8	51
Patriarch Grove South (PGS)	2012	3299-3478	Dolomitic	SE	8	51

Table 1: Sampled peaks and associated data in the White Mountains, CA. Total richness is the number of

unique species found on each peak.

FIGURE LEGENDS

Figure 1. Geographic extent and terrain of the study area in the White Mountains, east of the Sierra

Nevada mountain range in California (red diamond in inset). Points mark the centers for each transect,

with 50m long and 1m wide belt transects extending on either side of these center points following the

elevation contour. Terrain is rendered as a hillshade using the ⅓ arc second (~10m) digital elevation

model from the USGS National Elevation Dataset with a sun angle of 40 degrees and a sun direction of

270 degrees. WMT = White Mountain Peak, BAR = Mount Barcroft, SME = Sheep Mountain East, PGS =

Patriarch Grove South, and CPT = Campito Mountain. All transects are centered along the southeast

aspect of each peak, except Campito Mountain (CPT), which is centered on the east aspect.

Figure 2. Non-metric dimensional scaling (NMDS) ordination of Bray-Curtis dissimilarity. The arrow

indicates the directional influence of elevation. In the event of species name overlap, only the more

common species is shown here, but all species were included in the analysis.

Figure 3: Community-weighted Climatic Niche Means for (A) Climatic Water Deficit (CWD), (B) Mean

Annual Precipitation (MAP), (C) Average July maximum temperature (July T_{max}), and (D) Average January

minimum temperature (Jan T_{min}) along the elevation gradient. The black solid lines represent the linear

model fit for the entire range, all of which showed highly significant trends ($P < 0.001$). Shaded areas

represent the standard error around the linear model. Each point is 1 transect and is color-coded by

peak. Peaks with a significant ($P < 0.05$) relationship between CCNM and elevation are shown with a

solid line, and those with a non-significant ($P > 0.05$) relationship are shown with a dashed line.

Figure 4. Relationship between community climatic niche mean (CCNM) range and per-transect β

diversity. Climate metrics with a significant ($P < 0.05$) relationship between CCNM range and β diversity

per transect are shown with a solid line, and those with a non-significant ($P > 0.05$) relationship are

shown with a dashed line.

N

Summit
- WMT (14 transects)
- BAR (14 transects)
- SME (11 transects)
- PGS (8 transects)
- CPT (9 transects)

0   1   2   3 km