The activity-dependent transcription factor Npas4 regulates IQSEC3 expression in somatostatin interneurons to mediate anxiety-like behavior

Seungjoon Kim[1,#], Dongseok Park[1,#], Jinhu Kim[1], Dongsoo Lee[2], Dongwook Kim[1], Hyeonho Kim[1], Sookyung Hong[3], Jongcheol Jeon[3], Jaehoon Kim[3], Eunji Cheong[2], Ji Won Um[1,4,*] & Jaewon Ko[1,*]

[1]Department of Brain and Cognitive Sciences, Daegu Gyeongbuk Institute of Science and Technology (DGIST), 333 Techno Jungangdae-Ro, Hyeonpoong-eup, Dalseong-gun, Daegu 42988, Korea. [2]Department of Biotechnology, College of Life Science and Biotechnology, Yonsei University, Seoul 03722, Korea. [3]Department of Biological Sciences, Korea Advanced Institute of Science and Technology (KAIST), Daejeon 34141, Korea. [4]Core Protein Resources Center, Daegu Gyeongbuk Institute of Science and Technology (DGIST), 333 Techno Jungangdae-Ro, Hyeonpoong-eup, Dalseong-gun, Daegu 42988, Korea.

[#]These authors contributed equally to this study

*Email: jiwonum@dgist.ac.kr (J.W.U.) or jaewonko@dgist.ac.kr (J.K.)

Keywords: ARF; GABA; inhibitory synapse; IQSEC3; Npas4; somatostatin interneuron

Abstract

Organization of mammalian inhibitory synapses is thought to be crucial for normal brain

functions, but the underlying molecular mechanisms have been still incompletely understood.

IQSEC3 (IQ motif and Sec7 domain 3) is a guanine nucleotide exchange factor for ADP-

ribosylation factor (ARF-GEF) that directly interacts with gephyrin. Here, we show that

GABAergic synapse-specific transcription factor, Npas4 (neuronal PAS domain protein 4)

directly binds to the promoter of *Iqsec3* and regulates its transcription. Strikingly, an enriched

environment (EE) induced Npas4 upregulation and concurrently increased IQSEC3 protein

levels specifically in mouse CA1 *stratum oriens* layer somatostatin (SST)-expressing

GABAergic interneurons, which are compromised in *Npas4*-knockout (KO) mice. Moreover,

expression of wild-type (WT) IQSEC3, but not a dominant-negative (DN) ARF-GEF–inactive

mutant, rescued the decreased GABAergic synaptic transmission in Npas4-deficient SST

interneurons. Concurrently, expression of IQSEC3 WT normalized the altered GABAergic

synaptic transmission in dendrites, but not soma, of Npas4-deficient CA1 pyramidal neurons.

Furthermore, expression of IQSEC3 WT, but not IQSEC3 DN, in SST-expressing interneurons

in CA1 SST *Npas4*-KO mice rescued the altered anxiety-like behavior. Collectively, our results

suggest that IQSEC3 is a key GABAergic synapse component that is directed by Npas4 activity-

and ARF activity-dependent gene programs in SST-expressing interneurons to orchestrate the

functional excitation-to-inhibition balance.

Introduction

An essential function of the nervous system is to generate optimal behavioral responses to diverse environmental information; moreover, combinatorial interactions between genetic components and sensory cues are essential for driving discrete steps in brain development[1, 2, 3, 4]. Neuronal activity and subsequent calcium influx activate specific signaling cascades that cause nuclear translocation of various transcription factors, resulting in the rapid induction of an early-response gene-expression program that, in turn, induces transcription of late-response genes that locally regulate synaptic development and plasticity. This paradigm has been well established at both glutamatergic-producing and γ-aminobutyric acid-producing (GABAergic) synapses[3, 5]. Transcription factors that are commonly induced by synaptic activity regulate distinct sets of target genes to control the strength, location, and number of both glutamatergic and GABAergic inputs[6]. However, which specific transcription factors couple with key synapse-organizing proteins to orchestrate the regulation of synapse development at both transcriptional and post-transcriptional levels remains incompletely understood[3].

One prominent transcription factor that specifically controls GABAergic synapse development is Npas4 (neuronal PAS domain protein 4)[7, 8, 9, 10, 11, 12], which is required for neuronal excitability in the adult dentate gyrus (DG) and for new and reactivated fear memories[13, 14]. Npas4 recruits RNA polymerase II to promoters and enhancers of target genes

that are regulated by neuronal activity in CA3 hippocampal subfields, thereby contributing to short- and long-term contextual memory[15]. Deletion of *Npas4* abolishes depolarization-induced mRNA expression of immediate-early genes, such as Arc/Arg3.1, c-Fos, Zif268/Egr1, and brain-derived neurotrophic factor (BDNF)[15]. Importantly, Npas4 expression levels dictate the degree of inhibition in specific neuronal compartments by organizing distinct neuron-type–specific genetic programs[8, 9]. Notably, BDNF regulates somatic, but not dendritic, inhibition in neural circuits of CA1 pyramidal neurons, and is responsive only to increased inhibitory inputs in glutamatergic neurons[8, 9]. In addition, Npas4 mediates experience-dependent spine development in olfactory bulb interneurons (INs) through control of Mdm2 expression, confers neuroprotection against kainic acid-induced excitotoxicity in hippocampal neurons through control of synaptotagmin-10 (Sy10) expression, and organizes the structure and strength of hippocampal mossy fiber-CA3 synapses during contextual memory formation through control of polo-like kinase 2 (Plk2) expression[16, 17, 18]. Moreover, in response to heightened neuronal activity, Npas4 is recruited to activity-dependent regulatory elements by ARNT2, which is normally complexed with NCoR2, a co-repressor of neuronal activity-regulated gene expression under basal states[19]. These prior studies suggest that synaptic roles of Npas4 vary in brain region-, synaptic-, and cellular context-dependent manner. Despite these insights, the sheer number of Npas4 target genes has hampered efforts to fully understand the mechanisms

underlying Npas4-mediated GABAergic synapse development. IQSEC3 (IQ motif and Sec7 domain 3; also known as BRAG3 or SynArfGEF) was previously isolated as a putative Npas4 target[8]. However, direct functional coupling of Npas4 with IQSEC3 in the context of GABAergic synapse development *in vivo* has not been investigated.

Postsynaptic scaffolding proteins organize functional synapses, provide platforms for postsynaptic receptors, and regulate downstream signaling cascades[20, 21]. Arguably the most extensively studied scaffolding protein at GABAergic synapses is gephyrin[22, 23]. We recently reported that gephyrin directly interacts with IQSEC3 to promote GABAergic synapse formation[24, 25]. IQSEC3, a member of the brefeldin A-resistant ADP ribosylation factor guanine exchange factor (ARF-GEF) family, is exclusively localized at GABAergic synapses[24, 25, 26, 27]. IQSEC3 overexpression increases the number of postsynaptic gephyrin and presynaptic GAD67 puncta, whereas IQSEC3 knockdown (KD) specifically decreases gephyrin cluster size in cultured hippocampal neurons, without altering gephyrin puncta density[24]. In addition, IQSEC3 KD in the dentate gyrus induced severe seizure activity, an effect that was reversed by restricted expression of somatostatin (SST) peptides in SST[+] neurons[28], suggesting that IQSEC3 may be critical for maintenance of network activities *in vivo*. However, the brain-region specific, neuronal activity-dependent role of IQSEC3 *in vivo* has yet to be identified.

Here, we provide the first evidence demonstrating that IQSEC3 mediates activity-

dependent GABAergic synapse development, coupled with Npas4-dependent transcriptional programs, specifically in SST-expressing interneurons. Npas4 binds to the promoter of *Iqsec3* to regulate *Iqsec3* expression levels in a manner that depends on synaptic activity, which concurrently increase Npas4 levels. Remarkably, EE-induced upregulation of IQSEC3 proteins is prominent in SST-expressing interneurons in the hippocampal CA1 stratum oriens layer, suggesting a unique role for IQSEC3 in shaping hippocampal circuitries. In keeping with this observation, IQSEC3 regulates Npas4-dependent spontaneous GABAergic synaptic transmission in SST-expressing interneurons, and further modulates the mode of Npas4-mediated GABAergic synaptic transmission in apical dendrites (but not soma) of CA1 pyramidal neurons, a function that also requires its ARF-GEF activity. Furthermore, expression of IQSEC3 WT, but not an ARF-GEF activity-deficient IQSEC3 mutant, in SST-expressing interneurons restored the altered anxiety-like behaviors observed in conditional knockout (KO) mice lacking *Npas4* in SST interneurons. We further suggest that IQSEC3 is a downstream factor for Npas4 in sculpting specific neuronal inhibition in distal dendritic domains by providing inhibitory inputs onto excitatory neurons, hinting at its crucial roles in shaping activity-dependent specific inhibitory neural circuits and controlling a selective set of mouse behaviors.

Results

Npas4 binds to the *IQSEC3* promoter.

The structural and functional plasticity of GABAergic synapses are dynamically modulated in a neuronal activity-dependent manner[29, 30]. We thus hypothesized that IQSEC3 might be involved in activity-dependent GABAergic synapse development. Among GABAergic synapse-specific transcription factors, Npas4 appears to be a prominent candidate factor with the potential to play this role[7, 31]. Npas4 orchestrates the expression of a wide variety of target genes, notably including that encoding BDNF, which regulates GABAergic synapse development to mediate domain-specific inhibition in particular cell-types[7, 8, 9]. Intriguingly, deep sequencing of Npas4-bound DNA revealed IQSEC3 as a putative Npas4 target[8], and a recent genome-wide ChIP-seq analysis showed strong enrichment of Npas4 at the upstream region near the transcription start site of the *IQSEC3* gene[32] (**Figure 1a**). Intriguingly, we noted that this Npas4 peak contains a single Npas4 consensus CACG motif (**Figure 1a**). Thus, we tested whether IQSEC3, like BDNF, could also be an Npas4 target that acts to promote GABAergic synapse development. To this end, we cotransfected HEK293T cells with a *Iqsec3* promoter-luciferase reporter construct and a vector expressing Npas4. Ectopic expression of Npas4 increased luciferase expression ~3-fold; notably, this increase was abolished by mutation

of the Npas4 consensus site in the *Iqsec3* promoter region (**Figure 1b**). We further performed

luciferase reporter assays in cultured cortical neurons transfected with Npas4 wild-type (WT) or

control, and treated with 55 mM KCl to trigger membrane depolarization (**Figure 1c**). We found

that Npas4 WT similarly increased luciferase expression of IQSEC3 by ~4-fold (**Figure 1c**).

Electrophoretic mobility shift assays (EMSAs) confirmed these results, showing that a

radiolabeled *Iqsec3* WT oligoduplex probe containing purified Npas4 protein yielded a single

bound complex; consistent with promoter-reporter assays, this DNA-protein interaction was

abolished by mutation of the Npas4 consensus site (**Figure 1d**). Chromatin immunoprecipitation

(ChIP) assays also demonstrated that Npas4 binds to the *Iqsec3* promoter in primary

hippocampal cultured neurons treated with 55 mM KCl to trigger membrane depolarization

(**Figure 1e**). No Npas4-binding was detected in untreated hippocampal cultured neurons

(**Figure 1e**). These data suggest that Npas4 can function as a DNA-binding transcriptional

activator of *IQSEC3*.

IQSEC3 functions as an Npas4 target to promote GABAergic synapse development.

To further determine whether IQSEC3, analogous to BDNF[7], is involved in Npas4

actions on GABAergic synapse development, we infected cultured neurons with lentiviruses

expressing shRNAs against *Iqsec3*, *Bdnf*, or control shRNA at 3 days *in vitro* (DIV3);

transfected them at DIV10 with EGFP alone, or EGFP and myc-Npas4; and then immunostained

them at DIV14 for gephyrin. Overexpression of Npas4 significantly increased gephyrin puncta

density in both dendrites and soma, an effect that was completely abolished by IQSEC3 KD

(**Supplementary Figure 1**). These effects were distinct from the previously reported BDNF-KD

effects on an Npas4 minigene, which led to altered GABAergic synaptic transmission

specifically in the soma, but not in the dendritic compartment[8] (**Supplementary Figure 1**).

These data reinforce the conclusion that IQSEC3 acts downstream of Npas4 to regulate activity-

dependent GABAergic synapse development in cultured hippocampal neurons.

IQSEC3 expression is specifically upregulated in somatostatin-positive interneurons of

hippocampal CA1 region.

It was previously reported that expression of *Npas4* and *Bdnf* mRNAs changes in

response to neuronal depolarization[7, 8]. Thus, we asked whether *Iqsec3* mRNA expression is also

responsive to changes in neuronal activity (**Figure 2** and **Supplementary Figure 2**). To test this,

we manipulated the sensory experiences of juvenile wild-type (WT) mice by housing littermates

in an enriched environment (EE), consisting of a running wheel and several novel objects that

were regularly refreshed (**Figure 2a**, see '**Methods**' for details). After 5 days in an EE setting,

hippocampi were removed, sectioned, and immunostained for c-Fos as an indirect marker of

neuronal activity. As expected, EE exposure drove robust c-Fos expression in the hippocampal

CA1 region (**Figure 2b**). We then performed immunohistochemical analyses on sections from

age-matched mice housed in a standard environment (SE) as a control using previously

validated antibodies recognizing Npas4 and IQSEC3[7, 27]. Basal levels of IQSEC3 in the

hippocampal CA1 region from mice housed under SE conditions were low (**Figure 2c, 2d**).

There was a significant increase in the number of Npas4$^+$ neurons in CA1 *stratum oriens* (SO)

layers from mice allowed to explore an EE relative to those maintained in an SE (**Figure 2c**).

Notably, IQSEC3 expression was also concomitantly increased in Npas4-positive neurons, an

effect that was prominent in SO layers of the CA1 hippocampus (**Figure 2c, 2d**). EE-induced

upregulation of IQSEC3 in the SO layer of the CA1 hippocampus was prominently observed in

SST, but not parvalbumin (PV) or cholecystokinin (CCK), GABAergic interneurons (**Figure 2e,

2f**). RNAscope-based *in situ* hybridization analyses also showed that EE induced marked

increases in the number of SST$^+$ neurons in the SO layer, accompanied by significant increases

in IQSEC3 mRNA expression, without affecting the number of pyramidal neurons in the

hippocampal CA1 region (**Figure 2e–g**), suggesting the operation of a specific transcriptional

mechanism for IQSEC3 mRNAs selectively in SST$^+$ neurons.

       To determine whether an activity-dependent increase in Npas4 level is responsible for

selective upregulation of IQSEC3 in the SO layer of the hippocampal CA1 region, we expressed

either active Cre-combinase fused to EGFP (Cre) or inactive Cre-recombinase fused to EGFP

(ΔCre, as a control) in floxed Npas4 mice (Npas4$^{f/f}$) via adeno-associated virus (AAV)-mediated

stereotactic injection, and housed the resulting mice in either SE or EE conditions (**Figure 3a**).

Quantitative immunohistochemical analyses showed that EE induced a significant increase in

the number of Npas4$^+$ or IQSEC3$^+$ neurons and Npas4 deletion blunted the this effect, reducing

the number of IQSEC3$^+$ neurons to a level comparable to that observed in SE-housed *Npas4-*

cKO mice (**Figure 3b–e**). This blunting of the number of IQSEC3$^+$ neurons was also

recapitulated by specific deletion of Npas4 in SST$^+$ neurons (**Figure 3f, 3g**; see **Supplementary**

Figure 2 for validation of the EE-induced upregulation of Npas4 level in Npas4-deficient SST$^+$

brain sections), further supporting the idea that Npas4 induces upregulation of IQSEC3

specifically in SST$^+$ neurons.

To further investigate the expression profile of IQSEC3 under pathological conditions

in which network activity is severely heightened, we performed immunohistochemistry using

~11 week-old adult mice that were either injected with saline (SA) or 15 mg/kg of KA

(**Supplementary Figure 3a**). KA injections significantly increased Npas4$^+$ cell numbers in both

SP and SO layers of the hippocampal CA1 region, GCL and hilus of the DG (**Supplementary**

Figure 3b). Strikingly, KA injections resulted in an increase in the number of IQSEC3$^+$ cells,

specifically in the SO layer, but not in the SP layer, molecular layer, or GCL of the DG

(**Supplementary Figure 3b**). In addition, quantitative analyses using KA-injected Npas4 cKO

mice showed that Npas4 deletions decreased the number of IQSEC3$^+$ cells to a level comparable

to that observed in SA-injected WT mice (**Supplementary Figure 3c, 3d**). Furthermore, the KA

injection-dependent upregulation of IQSEC3 in the SO layer of the CA1 hippocampus was

prominently observed in SST, but not PV or CCK, GABAergic interneurons (**Supplementary**

Figure 3e, 3f). Collectively, these data show that IQSEC3 levels are preferentially altered

specifically in SST interneurons following induction of neuronal activity *in vivo*.

IQSEC3 is required for postsynaptic GABAergic synaptic transmission in Npas4-deficient

SST-expressing interneurons.

To determine whether Npas4-induced increases in IQSEC3 levels in SST-expressing

interneurons has an impact on postsynaptic GABAergic transmission, we crossed mice carrying

the Cre-driver SST-Cre[33] and the Ai9 reporter mouse line[34] with Npas4 floxed mice (Npas4$^{f/f}$

mice), and exposed the resulting mice to EE conditions to induce Npas4 protein expression, as

previously described[8, 35] (**Figure 4a**).

We then performed voltage-clamp recordings from SST$^+$ (tdTomato$^+$) neurons to assess

predicted changes in GABAergic synaptic currents. Recordings from SST$^+$ neurons of the

hippocampal CA1 region revealed drastic reductions in the frequencies (but not amplitudes) of

miniature inhibitory postsynaptic currents (mIPSCs) in SST-*Npas4*–KO mice (**Figure 4b–d**), in

keeping with the loss of postsynaptic GABA$_A$Rs. Again, restricted introduction of IQSEC3 WT,

but not IQSEC3 E749A, into SST-*Npas4*–KO mice rescued the deficits in GABAergic synaptic

transmission (**Figure 4b–d**). The input resistance, resting membrane potentials, and resting

membrane potentials of Npas4-deficient SST$^+$ neurons were comparable to those of control

neurons (**Supplementary Table 2**).

In addition, to determine whether IQSEC3 levels in SST$^+$ neurons have an impact on

the number of GABAergic synapses, we stereotactically injected AAV-sh-IQSEC3 into the

hippocampal CA1 of SST-Cre; Ai9 mice (**Figure 4e**). Semi-quantitative immunohistochemical

analyses showed that the overlap in the area of punctate immunoreactivity of the GABA$_A$R γ2

subunit (GABA$_A$Rγ2) with the tdTomato$^+$ somata of SST$^+$ neurons in the SO layer of the

hippocampus CA1 region was markedly reduced in SST-*Npas4*–KO mice compared with that in

control mice (**Figure 4f, 4g**). Remarkably, transduction of Npas4-deficient SST$^+$ neurons with

double-floxed, inverted open reading frame (DIO), Cre-dependent AAVs expressing IQSEC3

WT, but not those expressing the ARF-GEF–inactive IQSEC3 E749A mutant completely

reversed the decreased density of GABA$_A$Rγ2 puncta in the SO (**Figure 4f, 4g**). Parallel

analyses using VGAT or GAD67 antibodies yielded similar results (**Figure 4f, 4g**). These data

suggest that IQSEC3 in SST$^+$ CA1 hippocampal neurons postsynaptically regulates Npas4-

mediated activity-dependent GABAergic synaptic transmission *in vivo*.

IQSEC3 is required for Npas4-mediated GABAergic synaptic transmission in dendrites of

hippocampal CA1 pyramidal neurons.

Next, to determine if IQSEC3 is an Npas4 target in regulating somatic and/or dendritic

inhibition of pyramidal neurons *in vivo*, we caused excision of Npas4 *in vivo* by injecting AAV-

EGFP/Cre (Npas4 KO) into Npas4$^{f/f}$ mice; mice injected with AAV-EGFP/ΔCre served as

controls. We then exposed these mice to EE for 4–5 days (**Figure 5a**), and recorded layer-

specific, monosynaptically isolated, evoked inhibitory postsynaptic currents (eIPSCs) from

AAV-infected CA1 pyramidal neurons (**Figure 5b–i**). Consistent with a previous report[8],

stimulation of axons in the SP layer produced marginally, but significantly, smaller eIPSCs in

Npas4 KO neurons than in neighboring Npas4 WT neurons, whereas stimulation of axons in the

stratum radiatum (SR) layer generated significantly larger eIPSCs in Npas4 KO neurons

(**Figure 5b–e**). Strikingly, expression of IQSEC3 WT in Npas4 KO neurons completely

reversed the increase in eIPSC amplitude induced by stimulation of axons in the SR layer, but

did not affect the decrease in eIPSC amplitude induced by stimulation of axons in the SP layer,

compared with uninfected Npas4 KO neurons (**Figure 5f, 5g**). Expression of IQSEC3 E749A in

Npas4 KO neurons did not affect the amplitudes of eIPSCs induced by stimulation of axons in

the SR or SP layer (**Figure 5h, 5i**). Paired-pulsed ratios (PPRs) measured following stimulation

of axons in SR and SP layers of Npas4 WT and Npas4 KO mice housed under enriched-

environment conditions and infected with AAVs expressing IQSEC3 WT or IQSEC3 E749A

were not different among groups (**Supplementary Figure 4**), indicating that the Npas4-

dependent changes in eIPSCs are not caused by altered presynaptic neurotransmitter release.

These data indicate that Npas4 acts through IQSEC3 in a manner that depends on IQSEC3

ARF-GEF activity to mediate dendritic GABAergic synaptic inhibition *in vivo*.

To corroborate the electrophysiological data obtained from CA1 Npas4 KO neurons

(**Figure 5b–i**), we performed semi-quantitative analyses of endogenous $GABA_AR\gamma2$ puncta in

the soma (SP layer), apical dendrites (SR layer), and basal dendrites (SO layer) of CA1

hippocampal neurons of Npas4$^{f/f}$ mice coinjected with AAVs encoding IQSEC3 WT and either

EGFP-Cre (Npas4 KO) or EGFP-ΔCre (Control) (**Figure 5j**). Infected neurons were labeled by

expressing AAV-EGFP-Cre or AAV-EGFP-ΔCre, whereas the differential effects of IQSEC3

expression in both somatic (SP layer) and dendritic (SR or SO layer) compartments on

GABAergic synapse development were quantified by expressing AAV-IQSEC3 WT at moderate

levels. The number of $GABA_AR\gamma2$ puncta in the SP layer of Npas4 KO neurons from mice

exposed to enriched environment (EE) was significantly decreased compared with that in

control neurons (**Figure 5k–n**), with a concomitant increase in the number of apical dendritic

GABA$_A$Rγ2 puncta in the SR (but not SO) layer of Npas4 KO neurons, consistent with a

previous report[8]. Remarkably, IQSEC3 WT expression in Npas4 KO neurons completely

abrogated the increase in the density of GABA$_A$Rγ2 puncta in the SR only in mice exposed to

EE conditions, without altering SP GABA$_A$Rγ2 puncta density (**Figure 5k–n**), consistent with

the electrophysiology data. These data suggest that IQSEC3 exerts differential Npas4

expression-dependent inhibition *in vivo*.

IQSEC3 is required for controlling hippocampal CA1- and Npas4-dependent anxiety-like

behavior.

Lastly, we investigated the behavioral effects of IQSEC3 expression in Npas4 KO mice.

Npas4 KO mice have been reported to exhibit diverse cognitive, emotional, and social

behavioral deficits[10, 18, 36, 37, 38]. To link the hippocampal CA1 electrophysiological phenotypes in

Npas4 KO mice with behavioral abnormalities, we performed a number of behavioral tasks,

including elevated plus maze (EPM), Y-maze, open-field (OF), novel object-recognition (NOR),

and forced swim (FS) tests using SST-*Npas4* KO and littermate control mice. SST-*Npas4* KO

mice were additionally injected in the hippocampal CA1 region with AAV-DIO-mCherry

(Npas4 cKO), AAV-DIO-IQSEC3 WT or AAV-DIO-IQSEC3 E749A (**Figure 6a**). SST-specific

Npas4-KO mice exhibited decreased anxiety/exploration-related behavior (increased entries

with similar time spent in open arms in the EPM test; **Figure 6b–d**), but normal depressive-related behavior (as assessed by FS test; **Supplementary Figure 5**), normal locomotor activity (as assessed by the OF test; **Supplementary Figure 6**), normal working memory (as assessed by the Y-maze test; **Supplementary Figure 7**), and normal recognition memory (as assessed by the NOR test; **Supplementary Figure 8**) phenotypes that are slightly different from those of constitutive Npas4 KO mice[37, 38]. Strikingly, expression of IQSEC3 WT in CA1 SST neurons was sufficient to rescue the altered anxiety-related behavior (EPM test) observed in SST-specific *Npas4*-KO mice (**Figures 6b–d**). Intriguingly, expression of IQSEC3 E749A failed to normalize the increased anxiety-related behavior, suggesting that the ARF-GEF activity of IQSEC3 is required for mediating the ability to form a subset of Npas4- and CA1 SST-dependent mouse behaviors (**Figures 6b–d**). Expression of BDNF in CA1 SST neurons in the SST-specific *Npas4*-KO mice induced enormous hyperactivity and anxiety, consistent with previous reports[39, 40], thus, we were unable to perform Y-maze, EPM, or FS tests on BDNF-expressing SST-specific *Npas4*-KO mice (**data not shown**). Taken together, together with previous studies[15, 18], these data suggest that Npas4 employs distinct sets of downstream components in its roles in various types of cognitive tasks involving different brain regions.

Discussion

In this study, we provide evidence from various experimental approaches to support the conclusion that IQSEC3 organizes activity-dependent GABAergic synapse development and controls hippocampal CA1-dependent anxiety-like behavior through intricate molecular mechanisms involving Npas4 and ARFs.

IQSEC3 is linked to Npas4-mediated post-transcriptional programs that orchestrate activity-dependent GABAergic synapse development. The activity-dependent regulation of IQSEC3 expression level prompted us to dissect how IQSEC3 is related to the previously elucidated mechanisms underlying activity-dependent GABAergic synapse development. A prime candidate was Npas4, a well-known transcription factor and putative upstream regulator of IQSEC3[8, 10]. Npas4 binds to the promoter region of *Iqsec3*, which contains Npas4 consensus binding sites; IQSEC3 protein, in turn, functions downstream of Npas4 in regulating GABAergic inhibition in both somatic and dendritic compartments of cultured hippocampal neurons (**Figure 1** and **Supplementary Figure 1**). IQSEC3 expression is specifically induced by manipulation of synaptic activity *in vivo* (**Figure 2** and **Supplementary Figure 3**), consistent with the idea that IQSEC3, whose corresponding gene is an Npas4 target, promotes activity-dependent GABAergic synapse development. This interpretation is further supported by the observations that upregulated IQSEC3 protein levels in SST[+] neurons of the hippocampal

CA1 SO layer were completely compromised in SST-specific *Npas4*-KO mice (**Figure 2, 3** and

Supplementary Figure 3). Although prior screens using GABAergic neuron-enriched cultures

to profile activity-dependent transcriptomics and single-cell analyses in the mouse visual cortex

failed to identify IQSEC3 as a stimulus-responsive gene[9, 41], experimental conditions employed

in these previous studies might have been unsuitable for identifying responsive genes, that are

specifically regulated in hippocampal CA1 SST interneurons. Our results also suggest that

context-dependent regulatory mechanisms may underlie Npas4-mediated transcriptional and

post-transcriptional control of IQSEC3 levels *in vivo*[42]. Notably, Npas4 differentially dictates

gene regulatory mechanisms tailored to the type of activity patterns along subcellular

compartments of a neuron[43].

IQSEC3 performs its inhibitory actions, distinct from other known Npas4 downstream

components that are involved in regulating neural circuit-wide homeostasis (BDNF and

Homer1a), neuroprotective signaling (Syt10), and synaptic plasticity during memory formation

(Plk2) across various brain regions, neuron types and synapse types[9, 17, 18, 44]. Loss of Npas4

function results in reduced GABAergic synapse density in both perisomatic and dendritic

regions of cultured hippocampal neurons, and blunts the inhibitory mode, concurrent with an

abnormal distribution of GABAergic synapses at both soma and apical dendrites of CA1

pyramidal neurons[7]. BDNF is specifically expressed in excitatory 'cortical' neurons and serves

to specifically promote perisomatic inhibition of them, whereas Plk2 is strongly expressed in excitatory 'hippocampal' CA3 pyramidal neurons that regulate thorny excrescence structures at mossy fiber-CA3 synapses[9, 18]. Anatomical analyses in the current study revealed that IQSEC3 KD abolished the Npas4-induced increase in inhibitory synaptic puncta density at both perisomatic and apical dendritic regions of hippocampal cultured neurons, whereas IQSEC3 overexpression in SST-specific *Npas4*-KO mice resulted in selective restoration of altered inhibitory synaptic transmission and structure in apical dendritic regions in CA1 hippocampal pyramidal neurons (**Figures 4, 5**, and **Supplementary Figure 1**). This mechanism of IQSEC3 action is in stark contrast to the specific mode of action of BDNF and neuroligin-2 at perisomatic regions of hippocampal neurons[7, 8, 45]. These observations might be integrated under a scenario in which altered inhibitory synapse development in both somatic and dendritic compartments of IQSEC3-deficient neurons involves at least two distinct pathways *in vivo*: an Npas4-dependent IQSEC3 pathway that mediates dendritic inhibition (**Figure 7b**), and an Npas4-independent IQSEC3 pathway that mediates somatic and/or dendritic inhibition (**Figure 7a**). Consistent with this speculation, activity-dependent upregulation of IQSEC3 was notable in SST interneurons (but not excitatory neurons) of the hippocampal CA1 region (**Figure 2**), suggesting that one plausible role of IQSEC3 in SST neurons is to increase inhibitory inputs onto them enabling Npas4 to manage optimal responses and enhance feedback inhibition of

principal neurons within local neural circuits in a context-dependent fashion (**Figure 7**). Given that IQSEC3 is expressed in various types of interneurons (PV^+, SST^+, or CCK^+) in mouse hippocampal regions (**data not shown**), it is likely that IQSEC3 differentially shapes specific circuit features that modulate homeostatic plasticity at distinct principal neuron-interneuron synapses. Indeed, SST^+ interneurons also provide inhibitory input to PV^+ interneurons, which preferentially innervate the perisomatic region and axon initial segment of pyramidal cells, thereby controlling their spike output[45]. Thus, it is possible that increased inhibition of PV^+ cells by disinhibited SST+ neurons might also contribute to disinhibition of pyramidal cells in SST-*Npas4* KO mice. In support of this conjecture, increased eIPSC amplitudes were observed in Npas4-deficient pyramidal cells of the SP layer (**Figure 5**). Future studies in which Npas4 is conditionally deleted in PV^+ interneurons are warranted to clarify this issue.

Expression of IQSEC3 in CA1 SST interneurons completely normalized the altered anxiety-like behavior of SST-specific *Npas4*-KO mice (**Figure 6**), suggesting that a distinct set of factors downstream of Npas4 is required for the modulation of a spectrum of hippocampus-dependent cognitive and behavioral tasks (e.g., IQSEC3 for CA1-dependent anxiety-like behavior, and BDNF and Plk2 for CA3-dependent contextual memory formation). More importantly, our results suggest that the altered anxiety-like behavioral phenotype of SST-*Npas4* KO mice reflects differential GABAergic inhibition of pyramidal cells by the context-dependent,

balanced activity of SST[+] interneurons and PV[+] interneurons. Conditional transgenic mice in which IQSEC3 is selectively deleted in specific interneurons should be further analyzed to address how various neural circuits involving different cell types are altered in the context of the GABAergic synaptic functions of IQSEC3 proposed in the current study. This further study is especially meaningful given the implications of SST[+] GABAergic neurons in modulating anxiolytic brain states and major depressive disorders through the control of inhibition at specific synapse types formed in discrete neural circuits[47, 48, 49, 50]. Other intriguing questions include whether hyper-excitability of SST[+] neurons leads to increased expression and release of SST peptides that might contribute to the behavioral phenotype of SST-*Npas4* KO mice, and how ARF signaling pathways are differentially involved in cell type-specific manner.

In conclusion, our observations underscore the significance of biochemical complexes involving IQSEC3, Npas4, ARFs, and other components in organizing activity-dependent GABAergic synapse development, and in shaping specific neural circuit architectures involved in a specific behavior.

References

1. Katz LC, Shatz CJ. Synaptic activity and the construction of cortical circuits. *Science* 274, 1133-1138 (1996).

2. Wong RO, Ghosh A. Activity-dependent regulation of dendritic growth and patterning. *Nat Rev Neurosci* 3, 803-812 (2002).

3. West AE, Greenberg ME. Neuronal activity-regulated gene transcription in synapse development and cognitive function. *Cold Spring Harb Perspect Biol* 3, pii:a005744 (2011).

4. Chen LF, Zhou AS, West AE. Transcribing the connectome: roles for transcription factors and chromatin regulators in activity-dependent synapse development. *J Neurophysiol* 118, 755-770 (2017).

5. West AE, Griffith EC, Greenberg ME. Regulation of transcription factors by neuronal activity. *Nat Rev Neurosci* 3, 921-931 (2002).

6. Mann EO, Paulsen O. Role of GABAergic inhibition in hippocampal network oscillations. *Trends Neurosci* 30, 343-349 (2007).

7. Lin Y*, et al.* Activity-dependent regulation of inhibitory synapse development by Npas4. *Nature* 455, 1198-1204 (2008).

8. Bloodgood BL, Sharma N, Browne HA, Trepman AZ, Greenberg ME. The activi

ty-dependent transcription factor NPAS4 regulates domain-specific inhibition. *Natu re* 503, 121-125 (2013).

9. Spiegel I*, et al.* Npas4 regulates excitatory-inhibitory balance within neural circui ts through cell-type-specific gene programs. *Cell* 157, 1216-1229 (2014).

10. Sun X, Lin Y. Npas4: Linking Neuronal Activity to Memory. *Trends Neurosci* 3 9, 264-275 (2016).

11. Ko J, Choii G, Um JW. The balancing act of GABAergic synapse organizers. *Tr ends Mol Med* 21, 256-268 (2015).

12. Kim S, Kim H, Um JW. Synapse development organized by neuronal activity-reg ulated immediate-early genes. *Exp Mol Med* 50, 11 (2018).

13. Ploski JE, Monsey MS, Nguyen T, DiLeone RJ, Schafe GE. The neuronal PAS domain protein 4 (Npas4) is required for new and reactivated fear memories. *PL oS One* 6, e23760 (2011).

14. Sim S, Antolin S, Lin CW, Lin Y, Lois C. Increased cell-intrinsic excitability in duces synaptic changes in new neurons in the adult dentate gyrus that require N pas4. *J Neurosci* 33, 7928-7940 (2013).

15. Ramamoorthi K*, et al.* Npas4 regulates a transcriptional program in CA3 require d for contextual memory formation. *Science* 334, 1669-1675 (2011).

16. Yoshihara S*, et al.* Npas4 regulates Mdm2 and thus Dcx in experience-dependent dendritic spine development of newborn olfactory bulb interneurons. *Cell Rep* 8, 843-857 (2014).

17. Woitecki AM, Muller JA, van Loo KM, Sowade RF, Becker AJ, Schoch S. Identification of Synaptotagmin 10 as Effector of NPAS4-Mediated Protection from Excitotoxic Neurodegeneration. *J Neurosci* 36, 2561-2570 (2016).

18. Weng FJ*, et al.* Npas4 Is a Critical Regulator of Learning-Induced Plasticity at Mossy Fiber-CA3 Synapses during Contextual Memory Formation. *Neuron* 97, 1137-1152 e1135 (2018).

19. Sharma N*, et al.* ARNT2 Tunes Activity-Dependent Gene Expression through NCoR2-Mediated Repression and NPAS4-Mediated Activation. *Neuron* 102, 390-406 e399 (2019).

20. Fritschy JM, Panzanelli P, Tyagarajan SK. Molecular and functional heterogeneity of GABAergic synapses. *Cell Mol Life Sci* 69, 2485-2499 (2012).

21. Arancibia-Carcamo IL, Moss SJ. Molecular organization and assembly of the central inhibitory postsynapse. *Results Probl Cell Differ* 43, 25-47 (2006).

22. Papadopoulos T, Soykan T. The role of collybistin in gephyrin clustering at inhibitory synapses: facts and open questions. *Front Cell Neurosci* 5, 11 (2011).

23. Choii G, Ko J. Gephyrin: a central GABAergic synapse organizer. *Exp Mol Med* 47, e158 (2015).

24. Um JW*, et al.* IQ Motif and SEC7 Domain-containing Protein 3 (IQSEC3) Inter acts with Gephyrin to Promote Inhibitory Synapse Formation. *J Biol Chem* 291, 10119-10130 (2016)

25. Um JW. Synaptic functions of the IQSEC family of ADP-ribosylation factor gua nine nucleotide exchange factors. *Neurosci Res* 116, 54-59 (2017).

26. Sakagami H*, et al.* Distinct synaptic localization patterns of brefeldin A-resistant guanine nucleotide exchange factors BRAG2 and BRAG3 in the mouse retina. *J Comp Neurol* 521, 860-876 (2013).

27. Fukaya M*, et al.* SynArfGEF is a guanine nucleotide exchange factor for Arf6 a nd localizes preferentially at post-synaptic specializations of inhibitory synapses. *J Neurochem* 116, 1122-1137 (2011).

28. Kim S*, et al.* Loss of IQSEC3 Disrupts GABAergic Synapse Maintenance and Decreases Somatostatin Expression in the Hippocampus. *Cell Rep* In Press.

29. Flores CE, Mendez P. Shaping inhibition: activity dependent structural plasticity of GABAergic synapses. *Front Cell Neurosci* 8, 327 (2014).

30. Oh WC, Smith KR. Activity-dependent development of GABAergic synapses. *Br*

ain Res 1707, 18-26 (2019).

31. Shamloo M*, et al.* Npas4, a novel helix-loop-helix PAS domain protein, is regula

    ted in response to cerebral ischemia. *Eur J Neurosci* 24, 2705-2720 (2006).

32. Kim TK*, et al.* Widespread transcription at neuronal activity-regulated enhancers.

    Nature 465, 182-187 (2010).

33. Taniguchi H*, et al.* A resource of Cre driver lines for genetic targeting of GAB

    Aergic neurons in cerebral cortex. *Neuron* 71, 995-1013 (2011).

34. Madisen L*, et al.* A robust and high-throughput Cre reporting and characterizatio

    n system for the whole mouse brain. *Nat Neurosci* 13, 133-140 (2010).

35. Hartzell AL*, et al.* Npas4 recruits CCK basket cell synapses and enhances canna

    binoid-sensitive inhibition in the mouse hippocampus. *Elife* 7, pii: e35927 (2018).

36. Shepard R, Heslin K, Coutellier L. The transcription factor Npas4 contributes to

    adolescent development of prefrontal inhibitory circuits, and to cognitive and em

    otional functions: Implications for neuropsychiatric disorders. *Neurobiol Dis* 99, 3

    6-46 (2017).

37. Jaehne EJ, Klaric TS, Koblar SA, Baune BT, Lewis MD. Effects of Npas4 defic

    iency on anxiety, depression-like, cognition and sociability behaviour. *Behav Brai*

    n Res 281, 276-282 (2015).

38. Coutellier L, Beraki S, Ardestani PM, Saw NL, Shamloo M. Npas4: a neuronal transcription factor with a key role in social and cognitive functions relevant to developmental disorders. *PLoS One* 7, e46604 (2012).

39. Kernie SG, Liebl DJ, Parada LF. BDNF regulates eating behavior and locomotor activity in mice. *EMBO J* 19, 1290-1300 (2000).

40. Govindarajan A, *et al.* Transgenic brain-derived neurotrophic factor expression ca uses both anxiogenic and antidepressant effects. *Proc Natl Acad Sci U S A* 103, 13208-13213 (2006).

41. Hrvatin S, *et al.* Single-cell analysis of experience-dependent transcriptomic states in the mouse visual cortex. *Nat Neurosci* 21, 120-129 (2018).

42. Tyssowski KM, *et al.* Different Neuronal Activity Patterns Induce Different Gene Expression Programs. *Neuron* 98, 530-546 e511 (2018).

43. Brigidi GS, *et al.* Genomic Decoding of Neuronal Depolarization by Stimulus-Sp ecific NPAS4 Heterodimers. *Cell* 179, 373-391 e327 (2019).

44. Shan W, *et al.* Neuronal PAS domain protein 4 (Npas4) controls neuronal home ostasis in pentylenetetrazole-induced epilepsy through the induction of Homer1a. *J Neurochem* 145, 19-33 (2018).

45. Gibson JR, Huber KM, Südhof TC. Neuroligin-2 deletion selectively decreases in

hibitory synaptic transmission originating from fast-spiking but not from somatost atin-positive interneurons. *J Neurosci* 29, 13883-13897 (2009).

46. Pelkey KA, Chittajallu R, Craig MT, Tricoire L, Wester JC, McBain CJ. Hippoc ampal GABAergic Inhibitory Interneurons. *Physiol Rev* 97, 1619-1747 (2017).

47. Klausberger T, *et al.* Brain-state- and cell-type-specific firing of hippocampal inte rneurons in vivo. *Nature* 421, 844-848 (2003).

48. Lin LC, Sibille E. Somatostatin, neuronal vulnerability and behavioral emotionalit y. *Mol Psychiatry* 20, 377-387 (2015).

49. Fuchs T, Jefferson SJ, Hooper A, Yee PH, Maguire J, Luscher B. Disinhibition of somatostatin-positive GABAergic interneurons results in an anxiolytic and anti depressant-like brain state. *Mol Psychiatry* 22, 920-930 (2017).

50. Fee C, Banasr M, Sibille E. Somatostatin-Positive Gamma-Aminobutyric Acid Int erneuron Deficits in Depression: Cortical Microcircuit and Therapeutic Perspective s. *Biol Psychiatry* 82, 549-559 (2017).

51. Li K, *et al.* betaCaMKII in lateral habenula mediates core symptoms of depressi on. *Science* 341, 1016-1020 (2013).

52. Xu W, Südhof TC. A neural circuit for memory specificity and generalization. *S cience* 339, 1290-1295 (2013).

53. Soler-Llavina GJ, Fuccillo MV, Ko J, Südhof TC, Malenka RC. The neurexin lig ands, neuroligins and leucine-rich repeat transmembrane proteins, perform converg ent and divergent synaptic functions in vivo. *Proc Natl Acad Sci U S A* 108, 16 502-16509 (2011).

54. Kim J*, et al.* RAD6-Mediated transcription-coupled H2B ubiquitylation directly st imulates H3K4 methylation in human cells. *Cell* 137, 459-471 (2009).

55. Ko J, Na M, Kim S, Lee JR, Kim E. Interaction of the ERC family of RIM-bi nding proteins with the liprin-alpha family of multidomain proteins. *J Biol Chem* 278, 42377-42385 (2003).

56. Ko J*, et al.* Organization of the presynaptic active zone by ERC2/CAST1-depend ent clustering of the tandem PDZ protein syntenin-1. *J Neurosci* 26, 963-970 (20 06).

57. Sztainberg Y, Chen A. An environmental enrichment model for mice. *Nat Protoc* 5, 1535-1539 (2010).

Acknowledgements. We are grateful to Drs. Yingxi Lin (MIT, USA) and Hiroyuki Sakagami (Kitasato University, Japan) for kind gifts of reagents. This study was supported by grants from the National Research Foundation of Korea (NRF) funded by the Ministry of Science and

Future Planning (2019R1A2B5B02069324 to J.K.; 2019R1A2C1086048to J.W.U), the Brain

Research Program through the NRF funded by the Ministry of Science, ICT & Future Planning

(2017M3C7A1023470 to J.K. and 2017M3C7A1023471 to E.C.). The authors declare no

competing financial interests.

Author contributions. J.W.U. and J.K. conceived and directed the project; S.K., D.P., Jinhu

K., D.L., D.K., H.K., S.H. and J.J. performed the experiments; S.K., D.P., Jinhu K., D.L.,

Jaehoon K., E.C., J.W.U. and J.K. analyzed the data; and E.C., J.W.U. and J.K. wrote the

manuscript. All authors read and commented on the manuscript.

Competing interests. The authors declare no competing interests.

Figure legends

Fig. 1 Npas4 binds to the IQSEC3 promoter in an activity-dependent manner.

a Schematic representation of the Iqsec3 locus indicating the three amplicons used for RT-qPCR. An Npas4-binding site identified by ChIP-seq analysis[32] is indicated as an Npas4 responsive element (Npas4 RE).

b HEK293T cells were cotransfected with a luciferase reporter construct (100 ng) containing a *Iqsec3* proximal promoter region (–280 to +36) together with a vector expressing Npas4 (40 ng), as indicated. MT indicates a promoter-reporter construct harboring a mutation in the consensus Npas4-binding sequence. Luciferase activity in cell extracts was assayed and expressed as relative to luciferase activity in cells transfected with a luciferase-reporter construct containing the WT *Iqsec3* promoter region (defined as 1). Values are expressed as means ± SEM from three independent experiments (***$p < 0.001$ vs. Mock-transfected controls; Mann-Whitney U Test).

c Cortical cultured neurons derived from Npas4$^{f/f}$ mice were infected with lentiviruses expressing ΔCre or Cre at DIV3, and transfected with a luciferase reporter construct containing an *Iqsec3* proximal promoter region at DIV10. Luciferase activity in extracts of neurons left untreated or treated with 55 mM KCl for 2 hours was assayed at DIV14. Values are expressed as

means ± SEM from three independent experiments (*$p < 0.05$; non-parametric ANOVA with Kruskal-Wallis test, followed by *post hoc* Dunn's multiple comparison test.).

d EMSAs employing purified FLAG-tagged Npas4 protein (10–90 ng) and a probe corresponding to nucleotide -110 to -69 within the *Iqsec3* proximal promoter are shown. An inset shows an image of a Coomassie Blue-stained SDS-PAGE gel of the purified FLAG-Npas4 protein used in experiments.

e ChIP-seq binding profiles for Npas4 at the *Iqsec3* locus in KCl-treated mouse cortical neurons. Data were downloaded from the GEO website (http://www.ncbi.nlm.nih.gov/geo); GEO accession number, GSE21161[31]. The total number of tags was normalized to the corresponding input.

Fig. 2. IQSEC3 protein level is regulated by neuronal activity *in vivo*.

a The enriched environment (EE) setting, including tunnels, shelters, toys and running wheels for voluntary exercise, used in the current study.

b Representative immunofluorescence images of hippocampal CA1 regions from adult (11-week-old) WT mice in a standard environment (SE) or EE setting were immunostained for c-Fos (red) and counterstained with DAPI (blue).

c Representative immunofluorescence images of the hippocampal CA1 SO layer from WT mice

in an SE or EE setting were immunostained for Npas4 (red) and IQSEC3 (green), and

counterstained with DAPI (blue).

d Bar graphs summarizing the density of IQSEC3/Npas4 double-positive neurons shown in (**c**).

Data are presented as means ± SEMs (*$p < 0.05$; Mann-Whitney U test; n = 24 sections/4 mice

for all conditions).

e Representative immunofluorescence images of the hippocampal CA1 SO layer from WT mice

in an SE or EE setting, immunostained for IQSEC3 (green), DAPI (blue), and SST (red; **left**),

PV (red; **middle**), or CCK (red; **right**). Yellow arrows indicate SST-positive interneurons with

upregulated IQSEC3 levels. Scale bar, 50 µm (applies to all images).

f Bar graphs summarizing quantitative results shown in (**e**). Data are presented as means ±

SEMs (**$p < 0.01$; Mann-Whitney U test; n = 18−24 sections/3−4 mice).

g, h Expression of *Iqsec3* (red) and *Sst* (green) mRNAs in adult WT mice in an SE or EE setting

was visualized by *in situ* hybridization using RNAscope technology. Nuclei were stained with

DAPI (blue); representative images of hippocampal CA1 neurons from the pyramidal (**g**) and

SO layer (**h**) from mice reared under the indicated conditions are shown. Scale bar, 20 µm

(applies to all images).

i Quantitative analyses of RNAscope data measuring *Iqsec3* mRNA expression in pyramidal

neurons or SST[+] interneurons. Data are presented as means ± SEMs (**$p < 0.01$, nonparametric

one-way ANOVA with Kruskal-Wallis test followed by *post hoc* Dunn's multiple comparison

test; n = 3−5 mice).

Fig. 3. Upregulation of IQSEC3 protein level *in vivo* depends on the presence of Npas4.

a Experimental scheme for immunohistochemistry. The CA1 region of the hippocampus of ~9-

week-old *Npas4*[f/f] mice was bilaterally injected with AAVs expressing ΔCre or Cre. Mice were

housed for 5 days in an SE or EE setting 9 days after AAV injections, and then analyzed by

immunohistochemistry.

b, d Representative immunofluorescence images of the hippocampal CA1 SO layer from

Npas4[f/f] mice was bilaterally injected with AAVs expressing ΔCre or Cre in an SE or EE setting,

immunostained for Npas4 (red) (**b**) or IQSEC3 (red) (**d**) and DAPI (blue). Green fluorescence

indicates neurons infected with AAV-Cre or AAV-ΔCre. Scale bar, 50 μm (applies to all images).

c, e Bar graphs summarizing the percentage of Npas4$^+$ (**c**) or IQSEC3$^+$ (**e**) cells shown in (**b** and

d). Data are presented as means ± SEMs (*$p < 0.05$; **$p < 0.01$, ****$p < 0.0001$; non-

parametric ANOVA with Kruskal-Wallis test, followed by *post hoc* Dunn's multiple comparison

test).

f, Representative immunofluorescence images of the hippocampal CA1 SO layer from Control

or SST-*Npas4* mice under SE or EE conditions, immunostained for SST (red), IQSEC3 (green),

and DAPI (blue). Scale bar, 50 μm (applies to all images).

g, Bar graphs summarizing the number of IQSEC3$^+$ cells expressed in SST$^+$ interneurons shown

in (**f**). Data are presented as means ± SEMs (**$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$; non-

parametric ANOVA with Kruskal-Wallis test, followed by *post hoc* Dunn's multiple comparison

test).

Fig. 4 IQSEC3 is postsynaptically required for GABAergic synaptic transmission in

somatostatin-expressing interneurons.

a Experimental configuration for *ex vivo* electrophysiology experiments (**b–f**).

b–d Representative mIPSC traces (**b**) and quantification of results (**c**, **d**) in neurons from SST-

Npas4–KO mice exposed to EE conditions. Control, n = 29; SST-*Npas4,* n = 25; + DIO-

IQSEC3 WT, n = 26; and +DIO-IQSEC3 DN, n = 34. Data shown are means ± SEMs (*$p <$

0.05; paired two-tailed t-test). See **Supplementary Table 2** for intrinsic electrophysiological

properties of WT and SST-*Npas4*–KO neurons.

e Experimental configuration for immunohistochemistry experiments (**h**, **i**).

f, g Representative images (**f**) and quantification (**g**) of somatic GABA$_A$Rγ2 (left), VGAT

(middle), or GAD67 (right) puncta in the SO of neurons from SST- *Npas4*–KO mice exposed to

EE conditions. Data shown are means ± SEMs (*$p < 0.05$; **$p < 0.01$, ***$p < 0.001$; non-

parametric ANOVA with Kruskal-Wallis test, followed by *post hoc* Dunn's multiple comparison

test; n = 4 mice for all conditions). Scale bar, 10 μm (applies to all images).

Fig. 5 IQSEC3 regulates behaviorally induced, Npas4-mediated dendritic, but not somatic,

inhibition.

a Experimental configuration for *ex vivo* electrophysiology experiments. SL, stratum

lacunosum; SR, stratum radiatum; SP, stratum pyramidale; and SO, stratum oriens.

b–i eIPSC measurements in mice exposed to EE. Representative traces (**b**, **d**, **f**, **h**) of responses

in neurons infected with the indicated AAVs (blue or red) to stimulation in the SR or SP layer,

normalized pairwise to responses in uninfected neighboring neurons (black). Quantification

results are presented in (**c**, **e**, **g**, **i**). Open circles represent Npas4 KO/Npas4 WT pairs; black

circles indicate means ± SEM (SR [blue]: Control (ΔCre), n = 7 pairs; Npas4 KO (Cre), n = 7

pairs; WT rescue (Cre + IQSEC3 WT), n = 9 pairs; E749A rescue (Cre + IQSEC3 E749A), n =

10 pairs. SP [red]: Control (ΔCre), n = 7 pairs; Npas4 KO (Cre), n = 7 pairs; WT rescue (Cre +

IQSEC3 WT), n = 8 pairs; and E749A rescue (Cre + IQSEC3 E749A), n = 7 pairs). Data shown

are means ± SEMs (*$p < 0.05$; paired two-tailed t-test). See **Supplementary Table 3** for

intrinsic electrophysiological properties of Npas4 WT and Npas4 KO neurons.

j Experimental configuration for *in vivo* imaging experiments. SL, stratum lacunosum; SR,

stratum radiatum; SP, stratum pyramidale; and SO, stratum oriens.

k–n Representative images (**k**) and quantification of dendritic GABA$_A$Rγ2 puncta in the SO (**l**)

or SR (**m**), and somatic GABA$_A$Rγ2 puncta in the SP (**n**) of neurons obtained from mice housed

under EE conditions and infected with either AAV-ΔCre-EGFP (Control), AAV-Cre-EGFP

(Npas4 KO), or AAV-Cre-EGFP plus AAV-IQSEC3 WT (IQSEC3 WT res.). Data are presented

as means ± SEMs (*$p < 0.05$; non-parametric ANOVA with Kruskal-Wallis test, followed by

post hoc Dunn's multiple comparison test. n = 18–24 sections/3 mice for all conditions). Scale

bar, 10 μm (applies to all images).

Fig. 6 Expression of IQSEC3 WT, but not IQSEC3 E749A, in the hippocampal CA1

normalizes altered anxiety-like behavior observed in KO mice lacking Npas4 in

somatostatin-positive neurons.

a Schematic diagram of mouse behavioral analyses. The CA1 region of the hippocampus of ~6-

week-old SST-*Npas4* mice was bilaterally injected with AAVs expressing mCherry (SST-*Npas4*

KO), or either of two IQSEC3 DIO-AAVs [AAV-DIO-IQSEC3 WT or AAV-DIO-IQSEC3 DN

(E749A)]. WT littermate mice were used as controls. Injected mice were subjected to five

behavioral tasks 4 weeks after the injections in the indicated order. Abbreviations: EPM,

elevated plus-maze test; FS, forced swim test; NOR, novel object-recognition test; and OF,

open-field test.

b–d Analysis of anxiety/exploration-related behavior by EPM test in SST-*Npas4* mice injected

with the indicated AAVs. Representative heat maps of the time spent in the open arms of the

EPM (**b**) are presented. Red represents increased time spent, and blue represents minimal time

spent during the test. Number of open arm entries (**c**) and time spent (**d**) in open arms were

measured. Data are presented as means ± SEMs (Control, n = 14; SST-*Npas4*, n = 15; Rescue

[+IQSEC3 WT], n = 14; and Rescue [+IQSEC3 E749A], n = 15; *****$p < 0.05$; ******$p < 0.01$, one-

way ANOVA with Bonferroni's *post hoc* test).

Fig. 7 Molecular model of IQSEC3 action in regulating Npas4 activity-dependent neural

circuit-wide homeostasis.

a Npas4-independent IQSEC3 action in controlling activity-dependent GABAergic synapse

development across universal synapse types. **b** Npas4-dependent IQSEC3 action in controlling

activity-dependent GABAergic synapse development in specific synapse types. Neuronal

activity-induced transcriptional programs are orchestrated by coordinated actions of various

Npas4 downstream genes in distinct neuron types to maintain proper network activity. In WT

mice (**b1**), a specific pattern of neuronal activity (①) specifically upregulates IQSEC3

expression in SST interneurons, possibly inducing increased inhibitory input onto SST

interneurons (②), and act in conjunction with decreased inhibitory drive onto excitatory

neurons to maintain circuit activity (③). This occurs despite widespread upregulation of Npas4

expression in both excitatory and inhibitory neurons, which may activate other Npas4 target

genes in both excitatory and inhibitory neurons to maintain circuit activity. In contrast, neuronal

activity fails to induce IQSEC3 upregulation in SST interneurons of Npas4 KO mice (**b2)** (④),

possibly resulting in reduced recruitment of inhibitory inputs onto SST interneurons (⑤), and

collectively contributing to increased inhibitory drive onto a specific dendritic domain in

excitatory neurons (⑥), namely the dendritic domain in the SR layer, but not the SL or SO layer,

of the hippocampal CA1 region).

Methods

Cell culture. HEK293T cells were cultured in Dulbecco's Modified Eagle's Medium (DMEM; WELGENE) supplemented with 10% fetal bovine serum (FBS; Tissue Culture Biologicals) and 1% penicillin-streptomycin (Thermo Fisher) at 37°C in a humidified 5% CO_2 atmosphere. Cultured primary hippocampal neurons were prepared from embryonic day 18 (E18) Sprague-Dawley rat brains (KOATECK). Neurons were seeded on 25-mm poly-L-lysine (1 mg/ml)–coated coverslips and cultured in Neurobasal media (Gibco) containing penicillin-streptomycin and 0.5 mM GlutaMax (Thermo Fisher) supplemented with 2% B-27 (Thermo Fisher) and 0.5% FBS (Hyclone). All procedures were conducted according to the guidelines and protocols for rodent experimentation approved by the Institutional Animal Care and Use Committee of DGIST.

Animals. All C57BL/6N mice were maintained and handled in accordance with protocols approved by the Institutional Animal Care and Use Committee of DGIST under standard, temperature-controlled laboratory conditions, or in an enriched environment with free access to colored tunnels, mazes, climbing materials, and running wheels. Mice were kept on a 12:12 light/dark cycle (lights on at 6:00 am), and received water and food *ad libitum*. Npas4 KO

(Npas4[f/f]) mice were previously described[7]. Ai9 reporter mice (007909, Jackson Research Laboratories) and Sst-ires-Cre mice (013044, Jackson Research Laboratories) were gifts of Dr. Eunjoon Kim (KAIST, Korea). All experimental procedures were performed on male mice. Pregnant rats purchased from Daehan Biolink were used *in vitro* for dissociated cultures of cortical or hippocampal neurons.

Construction of expression vectors. *1*. IQSEC3. AAVs encoding full-length rat *Iqsec3* WT and the E749A point mutant were generated by amplification of the full-length region by PCR and subsequent subcloning into the pAAV-T2A-tdTomato vector (a gift from Hailan Hu[51]) at *Xba*I and *BamH*I sites. DIO-rat *Iqsec*3 WT and E749A fragments were PCR amplified and subcloned into the pAAV-hSynI-DIO-mCherry vector at *Nhe*I sites. The shRNA AAV against mouse *Iqsec3* (GenBank accession number: NM_207617.1) was constructed by annealing, phosphorylating, and cloning oligonucleotides targeting mouse *Iqsec3* (5'-GAA CTG GTG GTA GGC ATC TAT GAG A-3') into the *BamH*I and *EcoR*I sites of the pAAV-U6-GFP vector (Cell BioLabs, Inc.) or *Avr*II and *EcoR*I sites of the pAAV-Ef1α-DSE-mCherry-PSE vector (Addgene). *2*. Others. The shRNA lentiviral expression vector against *Bdnf* was constructed by annealing, phosphorylating, and cloning oligonucleotides targeting rat *Bdnf* (Genbank accession number: AY559250; 5'-ACA TCA GAC CAT CGG CCA CGA CATT A-3') into the *Xho*I and *Xba*I sites of the L-315

vector. AAV encoding mouse *Bdnf* (Genbank accession number: AY231132) was generated by amplification of the full-length region by PCR and subsequent subcloning into the pAAV-hSynI-DIO-mCherry vector at *Nhe*I sites. *3*. Previously published reagents. The following constructs were as previously described: pcDNA3.1-Myc-Npas4 [a gift from Yingxi Lin][7]; L-315 rat sh-IQSEC3, pcDNA3.1-Myc-IQSEC3 (shRNA-resistant) WT and E749A[24]; pAAV2/9-Cre-IRES-EGFP and pAAV2/9-ΔCre-IRES-EGFP[52]; and pFSW-Cre and pFSW-ΔCre[53].

Antibodies. The following commercially available antibodies were used: goat polyclonal anti-EGFP (Rockland), mouse monoclonal anti-Npas4 (clone N408/79; Neuromab), rabbit polyclonal anti-cholecystokinin-8 (Immunostar), mouse monoclonal anti-GAD67 (clone 1G10.2; Millipore), guinea pig polyclonal anti-VGLUT1 (Millipore), rabbit polyclonal anti-VGAT (Millipore), mouse monoclonal anti-gephyrin (clone 3B11; Synaptic Systems), rabbit polyclonal anti-GABA$_A$ receptor γ2 (Synaptic Systems), mouse monoclonal anti-PSD-95 (clone 7E3-1B8; Thermo Fischer Scientific), mouse monoclonal anti-β-actin (clone C4; Santa Cruz Biotechnology), mouse monoclonal anti-parvalbumin (clone PARV-19; Swant), and mouse monoclonal anti-somatostatin (clone YC7; Millipore). The following antibodies were previously described: rabbit polyclonal anti-IQSEC3 [JK079][24]; guinea pig polyclonal anti-IQSEC3/SynArfGEF [a gift from Hiroyuki Sakagami][27]; and rabbit polyclonal anti-Npas4 [a

gift from Michael Greenberg][8].

Iqsec3 promoter-luciferase reporter analysis. A DNA fragment bracketing the proximal promoter and transcription start site of the *Iqsec3* gene (–280 to +36) was PCR-amplified from mouse genomic DNA and subcloned into the pGL3-Basic vector (Promega). Full-length mouse Npas4 cDNA was subcloned in pcDNA3 (Gibco-Invitrogen). For luciferase assays, 5×10^4 HEK293T cells or cultured hippocampal neurons in 24-well plates were transfected with an Npas4 expression vector and reporter plasmid using iN-fect (iNtRON Biotechnology; for HEK293T cells) or CalPhos (Clontech; for cultured hippocampal neurons). Cells were harvested after 48 hours and analyzed for luciferase activity as described by the manufacturer (Promega).

Electrophoretic mobility shift assay. Full-length mouse Npas4 cDNA was subcloned into the pFASTBAC1 vector containing an N-terminal FLAG-tag, and baculovirus was generated according to the manufacturer's instructions (Thermo Fisher). After baculovirus infection into Sf9 insect cells, Npas4 protein was affinity purified on M2 agarose (Sigma) as described previously[54]. An oligonucleotide containing the consensus sequence for Npas4 binding (5'-CGG GAA GCA GGG TTC CTC ACG AAG CCC AGA GGT GGG AGG TGG-3') and its point mutant (5'-CGG GAA GCA GGG TTC CTA AAA AAG CCC AGA GGT GGG AGG TGG-3';

mutated nucleotides are underlined) within the *Iqsec3* proximal promoter region were end-labeled using polynucleotide kinase and [γ-^{32}P]ATP and then annealed to complementary oligonucleotides. Reactions containing purified Npas4 protein in 19 µl of reaction buffer (10 mM Tris-Cl pH 7.5, 5 mM MgCl$_2$, 1 mM EDTA, 50 mM K-glutamate, 1 mM DTT and 5% glycerol) supplemented with 25 ng/µl poly (dI-dC) were pre-incubated on ice for 15 minutes. After addition of 1 µl of ^{32}P-labeled probe containing approximately 3 ng of oligonucleotide duplex, reactions were incubated on ice for 30 minutes. Protein in samples were resolved by electrophoresis at 4°C on 6% polyacrylamide gels in 0.25x TBE buffer and subjected to autoradiography.

Chromatin immunoprecipitation assays. Cultured cortical neurons at DIV14 were treated with 55 mM KCl for 6 hours, after which chromatin immunoprecipitation assays were performed according the manufacturer's instructions (Millipore). Detailed information about primers used for quantitative PCR is provided in **Supplementary Table 1**.

Chemicals. Amino-5-phosphonopentanoic acid (D-APV; Cat. No. 0106), 2,3-dihydroxy-6-nitro-7-sulfamoyl-benzo[f] quinoxaline-2,3-dione (NBQX; Cat. No. 0373), 6,7-dinitroquinoxaline-2,3(1H,4H)-dione (DNQX; Cat. No. 0189), tetrodotoxin (TTX; Cat. No. 1078), and QX-314

(Cat. No. 1014) were purchased from Tocris Bioscience. 6-Cyano-7-nitroquinoxaline-2,3-dione

(CNQX; Cat. No. C239), picrotoxin (Cat. No. P1675), 2,2,2-tribomoethyl alcohol (Cat. No.

T4840-2), and Tert-amylalcohol (Cat. No. 24048-6), and kainic acid (Cat. No. K0250) were

purchased from Sigma. Diazepam (Cat. No. 117) was purchased from Daewon Pharm.

Neuron culture, transfections, imaging, and quantitation. Cultured hippocampal neurons

were prepared from E18 rat brains, as previously described[55, 56], cultured on coverslips coated

with poly-D-lysine (Sigma), and grown in Neurobasal medium supplemented with B-27

(Thermo Fisher), 0.5% FBS (WELGENE), 0.5 mM GlutaMAX (Thermo Fisher), and sodium

pyruvate (Thermo Fisher). For overexpression of IQSEC3 in cultured neurons, hippocampal

neurons were transfected with pCAGGS-FLAG-IQSEC3 or its various derivatives, as indicated

in the individual figures, or with EGFP (Control) using a CalPhos Transfection Kit (Takara) at

DIV10 and immunostained at DIV14. For KD of IQSEC3 in cultured neurons, hippocampal

neurons were transfected with L-315 alone (Control), L-315 rat sh-IQSEC3 (IQSEC3 KD), or

cotransfected with IQSEC3 KD and shRNA-resistant myc-IQSEC3 using a CalPhos

Transfection Kit (Takara) at DIV8 and immunostained at DIV14. For immunocytochemistry,

cultured neurons were fixed with 4% paraformaldehyde/4% sucrose, permeabilized with 0.2%

Triton X-100 in phosphate buffered saline (PBS), immunostained with the indicated primary

antibodies, and detected with the indicated Cy3- and fluorescein isothiocyanate (FITC)-conjugated secondary antibodies (Jackson ImmunoResearch). Images were acquired using a confocal microscope (LSM780, Carl Zeiss) with a 63 x objective lens; all image setting were kept constant. Z-stack images were converted to maximal intensity projection and analyzed to obtain the size, intensity, and density of puncta immunoreactivities derived from marker proteins. Quantification was performed in a blinded manner using MetaMorph software (Molecular Devices).

AAV production, stereotactic surgery and virus injection. Recombinant AAVs were packaged with pHelper and AAV1.0 (serotype 2/9) capsids for high efficiency. HEK293T cells were cotransfected with pHelper and pAAV1.0, together with pAAV-U6-EGFP alone (Control), pAAV-U6-shNpas4 (Npas4 KD), pAAV-U6-sh-IQSEC3 (IQSEC3 KD), pAAV-DIO-IQSEC3 WT, pAAV-DIO-IQSEC3 E749A, pAAV-IQSEC3 WT or pAAV-IQSEC3 E749A. Cells were harvested 72–108 hours post transfection; after adding 0.5 M EDTA to the media, cells were washed three times with PBS, and collected by centrifugation. Cells were then resuspended in PBS and lysed by subjecting them to four freeze-thaw cycles in an ethanol/dry ice bath (7 minutes each) and 37 °C water bath (5 minutes). Lysates were centrifuged and supernatants were collected and incubated with a solution containing 40% poly(ethylene glycol) (Sigma) and

2.5 M NaCl on ice for 1 hour and centrifuged at 2000 rcf for 30 minutes. The pellets were

resuspended in HEPES buffer (20 mM HEPES pH 8.0, 115 mM NaCl, 1.2 mM $CaCl_2$, 1.2 mM

$MgCl_2$, 2.4 mM KH_2PO_4), mixed with chloroform, and centrifuged at 400 rcf for 10 minutes.

The supernatant was collected and concentrated using Amicon Ultra Centrifugal Filters (0.5 ml,

3K MWCO; Millipore). Viruses were assessed for infectious titer by RT-PCR, and used for

infections at 1×10^{10}–10^{11} infectious units/μl. For stereotactic delivery of recombinant AAVs, 6–

8-week-old C57BL/6N mice were anesthetized by inhalation of isoflurane (3–4%) or

intraperitoneal injection of 2% Avertin solution (2,2,2-tribromoethyl alcohol dissolved in Tert-

amylalcohol (Sigma)) dissolved in saline, and secured in the stereotactic apparatus. Viral

solutions were injected with a Hamilton syringe using a Nanoliter 2010 Injector (World

Precision Instruments) at a flow rate of 100 nl/min (injected volume, 0.6 μl). The coordinates

used for stereotactic injections into mice are as follows: hippocampal DG injections (anterior-

posterior [AP] -2.2 mm, medial-lateral [ML] ± 1.3 mm, and dorso-ventral [DV] 2.35 mm) and

hippocampal CA1 injections (AP -2.2 mm, ML ± 1.3 mm, and DV 1.9 mm). Each injected

mouse was returned to its home cage and used for scoring seizure-like behaviors,

immunohistochemical analyses, electrophysiological recordings or behavioral analyses after 2–4

weeks.

qRT-PCR. Cultured rat cortical neurons were infected with recombinant lentiviruses at DIV3 and harvested at DIV10 for qRT-PCR using SYBR green qPCR master mix (Takara). Total RNA was extracted from rat cortical neurons using the TRIzol reagent (Invitrogen) according to the manufacturer's protocol. Briefly, one well of a 12-well plate of cultured neurons was harvested and incubated with 500 μl of TRIzol reagent at room temperature for 5 minutes. After phenol-chloroform extraction, RNA in the upper aqueous phase was precipitated. cDNA was synthesized from 500 ng of RNA by reverse transcription using a ReverTra Ace-α kit (Toyobo). Quantitative PCR was performed on a CFX96 Touch Real-Time PCR system (BioRad) using 1 μl of cDNA. The ubiquitously expressed β-actin was used as an endogenous control. The sequences of the primer pairs used are as follows: rat I*qsec3*, 5'-GGA GCA GAT TCG GAT AGA ATG G-3' (forward) and 5'–GGG TGA TCC TTG CTT TGA CT-3' (reverse); rat *Npas4*, 5'-GAG GCT GGA CAT GGA TTT ACT–3' (forward) and 5'-GGG TGA TCC TTG CTT TGA CT-3' (reverse); rat *Bdnf*, 5'-CTG AGC GTG TGT GAC AGT ATT A-3' (forward) and 5'-CTT TGG ATA CCG GGA CTT TCT C-3'; mouse *Npas4*, 5'-CCT CCA AAG AGC TGG ACT TC-3' (forward) and 5'-ATC CTT GCT CAG GTC TGC TT-3' (reverse); mouse *Iqsec3*, 5'-ATC ACC ACC AAC ATC ACC AC–3' (forward) and 5'-CTT GAC AAT CTG CTG GCA CT-3' (reverse); and mouse *Bdnf*, 5'-CTG AGC GTG TGT GAC AGT ATT A–3' (forward) and 5'-CTT TGG ATA CCG GGA CTT TCT C-3' (reverse)

Activity alteration protocols. *1. Environmental enrichment*. Male Npas4$^{f/f}$ mice (P28–P30)

injected with the indicated AAVs into the CA1 region were kept for 4 weeks and then exposed

to an enriched environment for 4–5 days before functional analyses. The enriched environment

consisted of a large cage containing a running wheel, hut, tunnel and several other novel objects,

as previously described[57]. *2. Induced seizure protocol*. Seizures were induced by intraperitoneal

administration of 11–12-week-old male mice with KA (15 mg/kg); saline injection was used as

a control. Mice were decapitated at the indicated times (3 or 6 hours) after injection, and the

injected mouse brains were prepared immediately for further analyses.

Fluorescent *in situ* hybridization (RNAscope assay). Frozen sections (14-µm thick) were cut

coronally through the hippocampal formation and thaw-mounted onto Superfrost Plus

microscope slides (Advanced Cell Diagnostics). Sections were fixed in 4% formaldehyde for 10

minutes, dehydrated in increasing concentrations of ethanol for 5 minutes, air-dried, and then

pretreated with protease for 10 minutes at room temperature. For RNA detection, sections were

incubated in different amplifier solutions in a HybEZ hybridization oven (Advanced Cell

Diagnostics) at 40°C. Synthetic oligonucleotides complementary to the sequence corresponding

to nucleotide residues 697–1045 of NM-001033354.3 (Mm–lqsec3–tv1), 1908–3117 of NM-

001033354.3 (Mm-lqsec3–C2), 793-1795 of NM_153553.4 (Mm-Npas4-C2), and 18-407 of

NM_009215.1 (Mm-SST-C3) (Advanced Cell Diagnostics) were used as probes. The labeled

probes were conjugated to Alexa Fluor 488, Altto 550 or Altto 647, after which labeled probe

mixtures were hybridized by incubating with slide-mounted sections for 2 hours at 40 °C.

Nonspecifically hybridized probes were removed by washing the sections three times for 2

minutes each with 1X wash buffer at room temperature, followed by incubation with Amplifier

1-FL for 30 minutes, Amplifier 2–FL for 15 minutes, Amplifier 3–FL for 30 minutes, and

Amplifier 4 Alt B–FL for 15 minutes at 40°C. Each amplifier was removed by washing with 1X

wash buffer for 2 minutes at room temperature. The slides were imaged with an LSM780

microscope (Zeiss) and analyzed using Image J (NIH).

Immunohistochemistry. Three-month-old mice were anaesthetized and immediately perfused,

first with PBS for 3 minutes, and then with 4% paraformaldehyde for 5 minutes. Brains were

dissected out, fixed in 4% paraformaldehyde overnight, and then incubated with 30% sucrose

(in PBS) overnight, and sliced into 30-μm-thick coronal sections using a vibratome (Model

VT1200S; Leica Biosystems) or a cryotome (Model CM-3050-S; Leica Biosystems). Sections

were permeabilized by incubating with 1% Triton X-100 in PBS containing 5% bovine serum

albumen and 5% horse serum for 30 minutes. For immunostaining, sections were incubated for

8–12 hours at 4 °C with primary antibodies diluted in the same blocking solution. The following primary antibodies were used: anti-Npas4 (1:100), anti-IQSEC3 (JK079; 2 μg/ml), anti-GAD67 (1:100), anti-PV (1:500), anti-CCK (1:100), and anti-SST (1:50). Sections were washed three times in PBS and incubated with appropriate Cy3- or FITC-conjugated secondary antibodies (Jackson ImmunoResearch) for 2 hours at room temperature. After three washes with PBS, sections were mounted onto glass slides (Superfrost Plus; Fisher Scientific) with Vectashield mounting medium (H-1200; Vector Laboratories).

***Ex vivo* electrophysiology.** *1. SST interneuron recordings.* Whole-cell voltage-clamp recordings were obtained from acute brain slices. Brain slices were transferred to a recording chamber and perfused with a bath solution of aerated (O_2 95%/CO_2 5% mixed gas) artificial cerebrospinal fluid (aCSF) consisting of 124 mM NaCl, 3.3 mM KCl, 1.3 mM NaH_2PO_4, 26 mM $NaHCO_3$, 11 mM D-glucose, 2.5 mM $CaCl_2$, and 1.5 mM $MgCl_2$ at 30°C. Patch pipettes (open pipette resistance, 3–5 MΩ) were filled with an internal solution consisting of 145 mM CsCl, 5 mM NaCl, 10 mM HEPES, 10 mM EGTA, 4 mM Mg-ATP, and 0.3 mM Na-GTP. Whole-cell recordings of mIPSCs were performed on CA1 *stratum oriens* SST⁺ interneurons, voltage clamped at -70 mV, and currents were pharmacologically isolated by bath application of 50 μM D-APV (Tocris), 10 μM CNQX (Sigma), and 1 μM TTX (Tocris). Electrophysiological

data were acquired using pCLAMP software and a MultiClamp 700B (Axon Instruments), and were digitized using an Axon DigiData 1550B data acquisition board (Axon Instruments). Data were sampled at 10 kHz and filtered at 4 kHz. Experiments were discarded if the holding current was greater than -250 pA, the mean series resistance of pairs was greater than 30 MΩ, or the series resistance differed by more than 20% between the two recordings. *2. Pyramidal neuron recordings.* Whole-cell voltage-clamp recordings were obtained from acute brain slices. Brain slices were transferred to a recording chamber and perfused with a bath solution of aerated (O_2 95%/CO_2 5% mixed gas) artificial cerebrospinal fluid (aCSF) consisting of 124 mM NaCl, 3 mM KCl, 1.3 mM $MgSO_4$, 1.25 mM NaH_2PO_4, 26 mM $NaHCO_3$, 2.4 mM $CaCl_2$-$2H_2O$, and 10 mM glucose at 32°C. Patch pipettes (open pipette resistance, 2–5 MΩ) were filled with an internal solution consisting of 130 mM CsCl, 2 mM $NgCl_2$, 10 mM HEPES, 5 mM Mg-ATP, 0.5 mM Na-GTP and 0.1 mM EGTA. For experiments in which eIPSCs were recorded, currents were pharmacologically isolated by bath application of 50 µM D-APV (Tocris) and 50 µM CNQX (Sigma), and inclusion of 5 mM QX-314 (Sigma) in the internal solution. Responses in hippocampal CA1 pyramidal neurons in the pyramidal layer evoked by stimulation of neurons in the SR layer were measured from a holding potential of -70 mV. Extracellular stimulation within specific layers of the hippocampus was achieved by current injection through a stimulating electrode placed in the relevant layer within 100–200 µm laterally of the patched

pair. The stimulus strength used was the minimum required to generate an eIPSC in both Npas4

WT and Npas4 KO neurons. Electrophysiological data were acquired using pCLAMP software

and a MultiClamp 700B (Axon Instruments), and were digitized using an Axon DigiData 1550B

data acquisition board (Axon Instruments). Data were sampled at 10 kHz and filtered at 4 kHz.

Experiments were discarded if the holding current was greater than -500 pA, the mean series

resistance of pairs was greater than 30 MΩ, or if the series resistance differed by more than 30%

between the two recordings. In experiments in which direct comparisons were made between

two neurons, all data were normalized to the peak amplitude of Npas4 WT neurons. The

amplitude of eIPSCs was calculated by averaging the normalized peak amplitude. Paired-pulse

ratios were calculated by recording the amplitudes of paired eIPSC at stimulation intervals of 25,

50, 100 and 200 ms.

Mouse behavioral tests. Male *Npas4*$^{f/f}$ mice (9–11 weeks old) crossed with the SST-*ires*-Cre

driver line were injected with the indicated AAVs were used for all behavioral tests. Tests were

performed in the following order: Y-maze, open-field, novel object-recognition, elevated-plus

maze, and forced swim test. Mice were excluded for quantitative analyses if one or both of the

injections were off-target, as demonstrated by *post hoc* immunostaining after behavioral

analyses. <u>*1. Y-maze test.*</u> A Y-shaped white acrylic maze with three 40-cm–long arms at a 120°

angle from each other was used. Mice were introduced into the center of the maze and allowed

to explore freely for 8 minutes. An entry was counted when all four limbs of a mouse were

within the arm. The movement of mice was recorded by a top-view infrared camera, and

analyzed using EthoVision XT 10 software (Noldus). *2. Open-field test.* Mice were placed into a

white acrylic open-field box ($40 \times 40 \times 40$ cm), and allowed to freely explore the environment

for 60 minutes in the dark (0 lux). The traveled distance moved and time spent in the center

zone of freely moving mice were recorded by a top-view infrared camera, and analyzed using

EthoVision XT 10 software (Noldus). *3. Novel object-recognition test.* An open field chamber

was used in this test. Mice were habituated to the chamber for 10 minutes. For training sessions,

two identical objects were placed in the center of the chamber at regular intervals, and mice

were allowed to explore the objects for 10 minutes. After the training session, mice were

returned to their home cage for 6 hours. For novel object-recognition tests, one of the two

objects was exchanged for a new object, placed in the same position of the chamber. Mice were

returned to the chamber and allowed to explore freely for 10 minutes. The movement of mice

was recorded by infrared camera, and the number and duration of contacts were analyzed using

EthoVision XT 10 (Noldus). *4. Elevated plus-maze test.* The elevated plus-maze is a plus-

shaped (+) white acrylic maze with two open arms ($30 \times 5 \times 0.5$ cm) and two closed arms

($30 \times 5 \times 30$ cm) positioned at a height of 75 cm from the floor. Light conditions around open

and closed arms were ~300 and ~30 lux, respectively. For the test, mice were introduced into the center zone of the elevated plus-maze and allowed to move freely for 10 minutes. All behaviors were recorded by a top-view infrared camera, and the time spent in each arm and the number of arm entries were measured and analyzed using EthoVision XT 10 software (Noldus).

5. Forced swim test. Mice were individually placed in a glass cylinder (15×30 cm) containing water ($24°C \pm 1°C$; depth, 15 cm). All mice examined were forced to swim for 6 minutes, and the duration of immobility was recorded and measured during the final 4 minutes of the test. The latency to immobility from the start of the test (delay between the start of the test and appearance of the first bout of immobility, defined as a period of at least 1 second without any active escape behavior), and the duration of immobility (defined as the time not spent actively exploring the cylinder or trying to escape from it) were measured. Immobility time was defined as the time the mouse spent floating in the water without struggling, making only minor movements that were strictly necessary to maintain its head above water.

Data analysis and statistics. All data are expressed as means ± SEM. All experiments were repeated using at least three independent cultures, and data were statistically evaluated using a Mann-Whitney *U* test, analysis of variance (ANOVA) followed by Tukey's *post hoc* test, Kruskal-Wallis test (one-way ANOVA on ranks), paired two-tailed t-test (for electrophysiology

experiments), or one-way ANOVA with Bonferroni's *post hoc* test (for behavior experiments), as appropriate. Prism7 (GraphPad Software) was used for analysis of data and preparation of bar graphs. *P* values < 0.05 were considered statistically significant (individual *p* values are presented in figure legends).

a

Mouse mm9_chr6

Npas4 ChIP-Seq

IQSEC3

Npas4 RE · exon 1 · intron 1 · exon 2

A · B · C · PCR primers

start

b *Luciferase reporter analyses*

HEK293T cells

Relative Luciferase Activity

***

Ctrl · WT · MT · IQSEC3 · Npas4

c

Relative Luciferase Activity

Luc-Iqsec3 promoter -expressing cultured neurons

Npas4^{f/f} + ΔCre
Npas4^{f/f} + Cre

KCl

d *EMSA analyses*

FLAG-Npas4

Luc-*Iqsec3* WT · MT

kDa

Npas4

Free DNA

e *ChIP analyses*

-10K · Promoter · +14K

Enrichment (% of input)

Control
Npas4

KCl

a *Enriched environment (EE)*

b
SE
c-Fos/DAPI

EE
c-Fos/DAPI

c

SE	*EE*
IQSEC3/Npas4/DAPI	IQSEC3/Npas4/DAPI
Npas4	Npas4
IQSEC3	IQSEC3
DAPI	DAPI

d
IQSEC3+/Npas4+ cells/mm^2

SE EE *

e

SE	EE	SE	EE	SE	EE
SST	SST	PV	PV	CCK	CCK
IQSEC3	IQSEC3	IQSEC3	IQSEC3	IQSEC3	IQSEC3
DAPI	DAPI	DAPI	DAPI	DAPI	DAPI

f
% of SST+IQSEC3 cells/IQSEC3 cells

**

SST **
PV
CCK
etc

SE EE

g *CA1 pyramidal cell*
Iqsec3/DAPI

SE

EE

h *CA1 SO layer SST interneuron*
Iqsec3/*Sst*/DAPI

SE

EE

i
iqsec3 mRNA puncta/mm^2 (x10^4)

Pyr   *SST*

SE EE SE EE

**

a *Npas4f/f mice*

Stereotaxic injection of AAV-ΔCre or AAV-Cre into hippocampal CA1 area of mice

5 weeks   SE vs. EE   **Immunohistochemistry**

7 weeks

b

| *Npas4f/f + ΔCre* | | *Npas4f/f + Cre* | |
| SE | EE | SE | EE |

ΔCre/Npas4/DAPI | ΔCre/Npas4/DAPI | Cre/Npas4/DAPI | Cre/Npas4/DAPI

Npas4 | Npas4 | Npas4 | Npas4

ΔCre | ΔCre | Cre | Cre

DAPI | DAPI | DAPI | DAPI

c *Summary for panel b*

% of Npas4 cells/Infected cells

**   **

****

SE EE | SE EE
ΔCre | Cre

d

| *Npas4f/f + ΔCre* | | *Npas4f/f + Cre* | |
| SE | EE | SE | EE |

ΔCre/IQSEC3/DAPI | ΔCre/IQSEC3/DAPI | Cre/IQSEC3/DAPI | Cre/IQSEC3/DAPI

IQSEC3 | IQSEC3 | IQSEC3 | IQSEC3

ΔCre | ΔCre | Cre | Cre

DAPI | DAPI | DAPI | DAPI

e *Summary for panel d*

% of IQSEC3 cells/Infected cells

*
**   ****

SE EE | SE EE
ΔCre | Cre

f

| Npas4 floxed mice (Control) | | SST-*Npas4* KO mice | |
| SE | EE | SE | EE |

SST/IQSEC3/DAPI | SST/IQSEC3/DAPI | SST/IQSEC3/DAPI | SST/IQSEC3/DAPI

IQSEC3 | IQSEC3 | IQSEC3 | IQSEC3

SST | SST | SST | SST

DAPI | DAPI | DAPI | DAPI

g *Summary for panel f*

IQSEC3+/SST+ cells/mm²

**
***   ****

SE EE | SE EE
Ctrl | SST-*Npas4*

a SST-Cre or Npas4^f/f;SST-Cre

AAV-DIO-mCherry (SST-*Npas4*)
AAV-DIO-mCherry + AAV-DIO-IQSEC3 WT
AAV-DIO-mCherry + AAV-DIO-IQSEC3 DN (E749A)

Stereotaxic injection of AAV-DIO
into hippocampal CA1 area

Enriched environment (EE)
Electrophysiology

5 weeks   9 weeks

b *mIPSC measurement*

Control

SST-*Npas4*

SST-*Npas4* + DIO-IQSEC3 WT

SST-*Npas4* + DIO-IQSEC3 DN (E749A)

40 pA
200 ms

c *mIPSC frequency*

Frequency (Hz)

d *mIPSC amplitude*

Amplitude (pA)

- Control
- SST-*Npas4*
- + DIO-IQSEC3 WT
- + IQSEC3 DN

e SST-Cre; Ai9-tdTomato

AAV-DIO-sh-Control-mCherry
AAV-DIO-sh-IQSEC3-mCherry
AAV-DIO-sh-IQSEC3-mCherry + AAV-DIO-IQSEC3 WT
AAV-DIO-sh-IQSEC3-mCherry + AAV-DIO-IQSEC3 DN (E749A)

Stereotaxic injection of AAV-DIO
into hippocampal CA1 area

Enriched environment (EE)
Immunohistochemistry

5 weeks   9 weeks

f

DIO-sh-Control / GABA_ARγ2 ; DIO-sh-Control / VGAT ; DIO-sh-Control / GAD67
DIO-sh-IQSEC3 / GABA_ARγ2 ; DIO-sh-IQSEC3 / VGAT ; DIO-sh-IQSEC3 / GAD67
+ IQSEC3 WT / GABA_ARγ2 ; + IQSEC3 WT / VGAT ; + IQSEC3 WT / GAD67
+ IQSEC3 DN / GABA_ARγ2 ; + IQSEC3 DN / VGAT ; + IQSEC3 DN / GAD67

g

Integrated intensity (AU)

GABA_ARγ2

VGAT

GAD67

- DIO-sh-Control
- DIO-sh-IQSEC3
- + IQSEC3 WT (res.)
- + IQSEC3 DN (res.)

a *Enriched environment (CA1 hippocampal neuron)*

SL
SR
SP
SO

Stimulating electrode

Patch electrode

AAV-uninfected; Npas4 f/f   *AAV-infected; Npas4 f/f*

Screen captures of a CA1 slice in DIC and epifluorescence

Uninfected   Infected

DIC

EGFP

b **Control (ΔCre)**

SR   *SP*

Uninf.   Inf.   Uninf.   Inf.

300 pA / 50 ms   150 pA / 50 ms

c

○ Pairs   ● Average

d **Npas4 KO (Cre)**

SR   *SP*

Uninf.   Inf.   Uninf.   Inf.

300 pA / 50 ms   100 pA / 50 ms

e

○ Pairs   ● Average

f **IQSEC3 WT res. (Cre+WT)**

SR   *SP*

Uninf.   Inf.   Uninf.   Inf.

150 pA / 50 ms   100 pA / 50 ms

g

○ Pairs   ● Average

h **IQSEC3 E749A res. (Cre+E749A)**

SR   *SP*

Uninf.   Inf.   Uninf.   Inf.

100 pA / 50 ms   150 pA / 50 ms

i

○ Pairs   ● Average

j *Enriched environment (CA1 hippocampal neurons)*

SO
SP
SR
SL

Npas4 f/f + CA1 AAV-infections of
1. *ΔCre-EGFP:* Control
2. *Cre-EGFP:* Npas4 KO
3. *Cre-EGFP + IQSEC3 WT:* IQSEC3 WT res.

k *Enriched environment (EE)*

Control (ΔCre)   Npas4 KO (Cre)   IQSEC3 WT res. (Cre+WT)

GABA$_A$Rγ2   GABA$_A$Rγ2   GABA$_A$Rγ2   SO

GABA$_A$Rγ2   GABA$_A$Rγ2   GABA$_A$Rγ2   SP

GABA$_A$Rγ2   GABA$_A$Rγ2   GABA$_A$Rγ2   SR

l *CA1 SO* — Puncta density/μm²

m *CA1 SP* — Puncta density/μm²
p=0.0896
p=0.0691

n *CA1 SR* — Puncta density/μm²

■ Control
■ Npas4 KO (SO)
■ Npas4 KO (SP)
■ Npas4 KO (SR)
■ IQSEC3 WT res. (SO)
■ IQSEC3 WT res. (SP)
■ IQSEC3 WT res. (SR)

a

SST-Cre or Npas4^{f/f};SST-Cre

Stereotaxic injection of AAV-DIO into hippocampal CA1 area

	Y-maze	OFT	NOR	EPM	FST
Day 1	Day 28	Day 29	Day 30	Day 31	Day 33

AAV-DIO-mCherry (SST-Npas4)
AAV-DIO-IQSEC3 WT
AAV-DIO-IQSEC3 DN (E749A)

b *Elevated plus maze test*

Control   SST-Npas4   + DIO-IQSEC3 WT   + DIO-IQSEC3 DN

Open arms
Closed arms

c

Open arm entries in elevated plus maze

d

Time in open arms in elevated plus maze (sec)

* ** *

● Control ■ SST-Npas4
▲ SST-Npas4 + DIO-IQSEC3 WT
▼ SST-Npas4 + DIO-IQSEC3 DN

a *Npas4-independent IQSEC3 action*

b1 WT **b2** KO — *Npas4-dependent IQSEC3 action*

SO
SP
SR
SL

SST+ interneuron (i.e., *O-LM neuron*)
PV+ interneuron (i.e., *Basket neuron*)
Excitatory neuron
IQSEC3 protein
Neuronal activity

Inhibitory input
Inhibitory input
Inhibitory input
Excitatory input

① NPAS4 IQSEC3↑
④ NPAS4 IQSEC3↓