Recombinant rotaviruses rescued by reverse genetics reveal the role of NSP5 hyperphosphorylation in the assembly of viral factories

Guido Papa[1,*], Luca Venditti[1], Francesca Arnoldi[1,2], Elisabeth M. Schraner[3], Christiaan Potgieter[4,5], Alexander Borodavka[6], Catherine Eichwald[3] and Oscar R. Burrone[1,*]

[1] International Centre for Genetic Engineering and Biotechnology, ICGEB, Padriciano 99, 34149 Trieste, Italy
[2] Department of Medicine, Surgery and Health Sciences, University of Trieste, Trieste, Italy
[3] Institute of Virology, University of Zürich, Zürich, Switzerland
[4] Department of Biochemistry, Centre for Human Metabolomics, North-West University, Potchefstroom, South Africa
[5] Deltamune (Pty) Ltd, 248 Jean Avenue, Lyttelton, Centurion, South Africa
[6] Astbury Centre for Structural Molecular Biology, School of Molecular and Cellular Biology, University of Leeds, Leeds, UK

* Corresponding authors, guido.papa@icgeb.org, burrone@icgeb.org

ABSTRACT

Rotavirus (RV) replicates in round-shaped cytoplasmic viral factories although how they assemble remains unknown.

During RV infection, NSP5 undergoes hyperphosphorylation, which is primed by the phosphorylation of a single serine residue. The role of this post-translational modification in the formation of viroplasms and its impact on the virus replication remains obscure. Here we investigated the role of NSP5 during RV infection by taking advantage of a modified fully tractable reverse genetics system. An NSP5 trans-complementing cell line was used to generate and characterise several recombinant rotaviruses (rRVs) with mutations in NSP5. We demonstrate that a rRV lacking NSP5, was completely unable to assemble viroplasms and to replicate, confirming its pivotal role in rotavirus replication.

A number of mutants with impaired NSP5 phosphorylation were generated to further interrogate the function of this post-translational modification in the assembly of replication-competent viroplasms. We showed that the rRV mutant strains exhibit impaired viral replication and the ability to assemble round-shaped viroplasms in MA104 cells. Furthermore, we have investigated the mechanism of NSP5 hyper-phosphorylation during RV infection using NSP5 phosphorylation-negative rRV strains, as well as MA104-derived stable transfectant cell lines expressing either wt NSP5 or selected NSP5 deletion mutants. Our results indicate that NSP5 hyper-phosphorylation is a crucial step for the assembly of round-shaped viroplasms, highlighting the key role of the C-terminal tail of NSP5 in the formation of replication-competent viral factories. Such a complex NSP5 phosphorylation cascade may serve as a paradigm for the assembly of functional viral factories in other RNA viruses.

IMPORTANCE

Rotavirus (RV) double-stranded RNA genome is replicated and packaged into

virus progeny in cytoplasmic structures termed viroplasms. The non-structural

protein NSP5, which undergoes a complex hyperphosphorylation process during

RV infection, is required for the formation of these virus-induced organelles.

However, its roles in viroplasm formation and RV replication have never been

directly assessed due to the lack of a fully tractable reverse genetics (RG)

system for rotaviruses. Here we show a novel application of a recently developed

RG system by establishing a stable trans-complementing NSP5-producing cell

line required to rescue rotaviruses with mutations in NSP5. This approach

allowed us to provide the first direct evidence of the pivotal role of this protein

during RV replication. Furthermore, using recombinant RV mutants we shed light

on the molecular mechanism of NSP5 hyperphosphorylation during infection and

its involvement in the assembly and maturation of replication-competent

viroplasms.

INTRODUCTION

Rotavirus (RV) is the most common cause of viral gastroenteritis in young children and infants worldwide (1, 2). It is a non-enveloped RNA virus with a genome composed of 11 segments of double-stranded RNA (dsRNA), which replicates in cytoplasmic structures primarily composed of viral proteins (3–5). During infection, the first steps of viral morphogenesis and genome replication occur within cytoplasmic viral replication factories known as viroplasms. (3, 5–7). The assembly of viroplasms requires co-expression of at least two non-structural proteins, NSP5 and NSP2 (8, 9), however, how these virus-induced organelles are formed remains unknown.

Other viral proteins also found in viroplasms include RNA-dependent RNA polymerase (RdRp) VP1, the main inner-core protein VP2, guanyltransferase/ methylase VP3, and the middle layer (inner capsid) protein VP6 (10, 11). Biochemical evidence suggests that viroplasms are essential for RV replication since the virus production is highly impaired upon silencing of either NSP2 or NSP5 (12–15).

Rotavirus NSP5, encoded by genome segment 11, is a small serine (Ser)- and threonine (Thr)-rich non-structural protein that undergoes multiple post-translational modifications in virus-infected cells, including O-linked glycosylation (16), N-acetylation (17), SUMOylation (18) and crucially hyperphosphorylation that involves several distinct Ser residues (19, 20). The NSP5 hyperphosphorylation is a complex process, which gives rise to multiple phosphorylation states ranging from the most abundant 28 kDa phospho-isoform, up to the hyperphosphorylated 32-34 kDa states (19, 20). All these forms have been found to be more stable in viroplasms, while chemical disruption of viroplasms results in NSP5 de-phosphorylation (21). The mechanism of NSP5 phosphorylation is not yet wholly understood, but it involves interactions with other viral proteins. When expressed alone in non-infected cells, NSP5 is not phosphorylated, while co-expression with NSP2 or VP2 results in NSP5 hyperphosphorylation and formation of viroplasm-like structures (VLS) (8, 22, 23). NSP5 hyperphosphorylation involves the phosphorylation of Serine 67

(Ser67) by Casein Kinase 1α (CK1α) to initiate the phosphorylation cascade (24,

25) and it is considered to be essential for the assembly of viroplasms (26).

Although the structure of NSP5 remains unknown, it readily forms higher

molecular weight oligomeric species in solution, potentially providing a larger

interface for interacting with multiple components of viroplasms (27).

In addition, the C-terminal region ('a tail' including amino acids 180-198) is

required for NSP5 decamerisation *in vitro* (27) and VLS formation *in vivo* (7).

However, due to the lack of a fully tractable reverse genetics (RG) system for

RVs until recently, previous studies on NSP5 have been carried out using the

mutants expressed in the absence of a complete set of viral proteins. Here, we

took advantage of the novel plasmid only-based, helper-virus free RG systems

for rotaviruses (28, 29) to gain new insights into the mechanisms of NSP5

hyperphosphorylation and its role in viroplasm assembly and virus replication

during viral infection.

To achieve this, we generated and characterised several viable recombinant

rotaviruses (rRVs) with mutations in NSP5. Using these mutants, we show the

role of NSP5 hyperphosphorylation for viroplasm assembly and in genome

replication. These studies shed light on a complex hierarchical mechanism of

NSP5 hyperphosphorylation during rotaviral infection.

MATERIALS AND METHODS

Cells and viruses. MA104 (embryonic African green monkey kidney cells, ATCC

CRL-2378.1 from *Chlorocebus aethiops*), U2OS (Human bone osteosarcoma

epithelial cells), Caco-2 (colorectal adenocarcinoma human intestinal epithelial

cell line, ATCC®HTB-37) and HEK293T (embryonic human kidney epithelial,

ATCC®CRL-3216) cells were cultured in Dulbecco's Modified Eagle's Medium

(DMEM) (Life Technologies) supplemented with 10% Fetal Bovine Serum (FBS)

(Life Technologies) and 50 µg/ml gentamycin (Biochrom AG).

MA104-NSP5-EGFP cells (MA-NSP5-EGFP) (7) were cultured in DMEM
supplemented with 10% FBS (Life Technologies), 50 µg/ml gentamycin
(Biochrom AG) and 1 mg/ml geneticin (Gibco-BRL, Life Technologies).
MA104-NSP2-mCherry (MA-NSP2-mCherry), MA104-Δ3 (MA-Δ3), MA104-Δtail
(MA-ΔT) and MA104-NSP5wt (MA-NSP5) stable transfectant cell lines
(embryonic African green monkey kidney cells, ATCC® CRL-2378) were grown
in DMEM (Life Technologies) containing 10% FBS, 50 µg/ml gentamycin
(Biochrom AG) and 5 µg/ml puromycin (Sigma-Aldrich).
BHK-T7 cells (Baby hamster kidney stably expressing T7 RNA polymerase) were
cultured in Glasgow medium supplemented with 5% FBS, 10% Tryptose
Phosphate Broth (TPB) (Sigma-Aldrich), 50 µg/ml gentamycin (Biochrom AG),
2% Non-Essential Amino Acid (NEAA), 1% Glutamine.
Recombinant simian RV strain SA11 (rRV-wt), rescued using reverse genetics
system using cDNA clones encoding the wild-type SA11 (G3P[2]) virus (28), was
propagated in MA104 cells cultured in DMEM supplemented with 0.5 µg/ml
trypsin (Sigma Aldrich).
174

Recombinant RVs Titration
Recombinant NSP5 mutant RVs were grown in MA-NSP5 cells and the lysate
was 2-fold serially diluted and used to infect MA-NSP5 cells, seeded in 24-wells
plates with coverslips. After 1 hour of adsorption, virus was removed, and cells
were incubated at 37°C. At 5 hours post-infection (hpi), cells were fixed with 4%
paraformaldehyde (PFA) in phosphate buffer saline (PBS) [137 mM NaCl; 2.7
mM KCl; 8.1 mM Na_2HPO_4 and 1.74 mM KH_2PO_4 pH7.5] for 15 min at room
temperature and permeabilized for 5 min with PBS containing 0.01% Triton X-
100. Next, cells were incubated for 30 min with PBS supplemented with 1%
bovine serum albumin (PBS-BSA) at room temperature and then with anti-NSP5
(1:1000) or anti-VP2 (1:200) or anti-NSP2 (1:200) guinea pig serum diluted in
PBS-BSA. After washing three times with PBS, cells were incubated for 1 h at
room temperature with TRITC-conjugated Anti-guinea Pig IgG (Jackson
ImmunoResearch) (1:500) diluted in PBS-BSA.

Nuclei were stained with ProLong™ Diamond Antifade Mountant with DAPI
(Thermo Scientific). Samples were imaged using a confocal setup (Zeiss
Airyscan equipped with a 63x, NA=1.3 objective). Each viroplasms-containing
cell was counted as one focus-forming unit (FFU). The average of cells with
viroplasms of six fields of view per each virus dilution was determined and the
total number of cells containing viroplasms in the whole preparation was
estimated. The virus titre was determined as:

$$Virus\ Titer\ \left(\frac{FFU}{ml}\right) = \frac{N * Dilution\ Factor}{V(ml)}$$

where *N* is a total number of cells containing 1 or more viroplasms, and *V* is the
volume of virus inoculum added.

Replication Kinetics of recombinant viruses
MA104 cell line (ATCC CRL-2378.1) or stably transfected MA104 cells (NSP5;
Δ3, ΔT) were seeded into 24-well plates and subsequently infected with
recombinant RVs at MOI (FFU/cell) of 0.5 for multi-step growth curve
experiments and MOI 5 for a single-step growth curve experiment. After
adsorption for 1 h at 37°C, cells were washed twice with PBS and the medium
replaced with DMEM without trypsin. After incubation at 37°C, cells were
harvested after 8, 16, 24, 36 hours post virus adsorption. The cell lysates were
freeze-thawed three times and activated with trypsin (1 μg/ml) for 30 min at 37°C.
The lysates were used to infect monolayers of MA-NSP5 cells seeded in μ-Slide
8 Well Chamber Slide-well (iBidi GmbH, Munich, Germany). The cells were then
fixed 5 hours post infection for 15 min with 4% paraformaldehyde and
permeabilized for 5 min with PBS containing 0.01% Triton X-100. Next, cells
were incubated for 30 min with PBS-BSA at room temperature and then with anti-
NSP5 serum (1:1000) diluted in PBS-BSA. After three washes with PBS, cells
were incubated for 1 h at room temperature with TRITC-conjugated Anti-guinea
Pig IgG (Jackson ImmunoResearch) (1:500) diluted in PBS containing 1% BSA
(PBS-BSA).

The number of infected cells was counted, and the virus titres were expressed in
Focus-Forming Units per mL (FFU/mL).
220
Plasmid construction.
RV plasmids pT$_7$-VP1-SA11, pT$_7$-VP2-SA11, pT$_7$-VP3-SA11, pT$_7$-VP4-SA11,
pT$_7$-VP6-SA11, pT$_7$-VP7-SA11, pT$_7$-NSP1-SA11, pT$_7$-NSP2-SA11, pT$_7$-NSP3-
SA11, pT$_7$-NSP4-SA11, and pT$_7$-NSP5-SA11 (28) were used to rescue
recombinant RVs by reverse genetics. pT$_7$-NSP5/S67A carrying a mutation in the
nucleotide T220G in the gs11 and pT$_7$-NSP5/Tyr18Stop harbouring a nucleotide
substitution T75G were generated by QuikChange II Site-Directed Mutagenesis
(Agilent Technologies). pT$_7$-NSP5/ΔT was generated from pT$_7$-NSP5-SA11 by
deleting the last 18 C-terminal amino acids (FALRMRMKQVAMQLIEDL) using
substitution of the F181 encoding triplet with a stop codon. pT$_7$-NSP5/Δ176-180
were obtained deleting the amino acids 176 to 180 (YKKKY). The described
deletions were performed using the QuikChange II Site-Directed Mutagenesis kit
(Agilent Technologies).
For the generation of lentiviral plasmids, NSP5 and NSP2-mCherry were
amplified by PCR and inserted into the plasmid pAIP (Addgene #74171; (30) at
the *NotI-EcoRI* restriction enzymes sites to yield pAIP-NSP5 and pAIP-NSP2-
mCherry, which were then used to generate lentiviruses for the MA104-stable
transfectant cell lines (MA-NSP5 and MA-NSP2-mCherry) (31). NSP5/ΔT was
amplified from the pT$_7$-NSP5/ΔT by PCR and inserted into the pPB-MCS (Vector
Builder) at restriction enzyme sites *NheI-BamHI* to generate pPB-NSP5/ΔT for
the production of the stable transfectant MA-ΔT.
pPB-NSP5/Δ3, for MA-Δ3 stable cell line establishment, was generated with a
GenParts™ DNA Fragment (GenScript) containing NSP5 ORF lacking the amino
acids 80-130 (VKTNADAGVSMDSSAQSRPSSNVGCDQVDFSLNKG
LKVKANLDSSISIST) and inserted into the *NheI-BamHI* restriction sites of pPB-
MCS vector.
247
Generation of stable cell lines

MA-NSP5-EGFP were generated as previously described (7).

MA-NSP2-mCherry and MA-NSP5 cell line were generated using lentiviral vector

system (31). Briefly, HEK293T cells were maintained in DMEM (Life

Technologies) supplemented with 10% FBS (Life Technologies), and 50 µg/ml

gentamycin (Biochrom AG). Approximately 7×10^6 HEK293T cells were seeded

in 10 cm^2 tissue culture dishes 24 hours before transfection. For each well, 2.4

µg of pMD2-VSV-G, 4 µg of pMDLg pRRE, 1.8 µg of pRSV-Rev and 1.5 µg of

plasmid containing pAIP-NSP2-mCherry or pAIP-NSP5 and the human

immunodeficiency virus long terminal repeats were co-transfected with

Lipofectamine 3000 (Sigma-Aldrich) according to the manufacturer's instructions.

After 48 h, the virus was collected by filtration with a 0.45-µm polyvinylidene

fluoride filter and was immediately used or stored at −80 °C. For lentiviral

transduction, MA104 cells were transduced in six-well plates with 1 ml of lentiviral

supernatant for 2 days.

MA-Δ3 and MA-ΔT were generated using the PiggyBac Technology (32). Briefly,

10^5 MA104 cells were transfected with the pCMV-HyPBase (32) and the

transposon plasmid pPB-NSP5/Δ3 and pPB-NSP5/ΔT using a ratio of 1:2.5 with

Lipofectamine 3000 (Sigma-Aldrich) according to the manufacturer's instructions.

The cells were maintained in DMEM supplemented with 10% FBS for 3 days and

then the cells were incubated with DMEM supplemented with 10% FBS and 5

µg/ml puromycin (Sigma-Aldrich) for 4 days to allow the selection of cells

expressing the gene of interest.

Rescue of recombinant RVs (rRVs) from cloned cDNAs.

To rescue recombinant RV strain SA11 (rRV-WT), monolayers of BHK-T7 cells

(4×10^5) cultured in 12-well plates were co-transfected using 2.5 µL of TransIT-

LT1 transfection reagent (Mirus) per microgram of DNA plasmid. Each mixture

comprised 0.8 µg of SA11 rescue plasmids: pT$_7$-VP1, pT$_7$-VP2, pT$_7$-VP3, pT$_7$-

VP4, pT$_7$-VP6, pT$_7$-VP7, pT$_7$-NSP1, pT$_7$-NSP3, pT$_7$-NSP4, and 2.4 µg of pT$_7$-

NSP2 and pT$_7$-NSP5 (29). Furthermore 0.8 µg of pcDNA3-NSP2 and 0.8 µg of

pcDNA3-NSP5, encoding NSP2 and NSP5 proteins, were also co-transfected to
increase rescue efficiency.
To rescue recombinant rRVs encoding NSP5 mutants, pT$_7$ plasmids encoding
NSP5/S67A, NSP5/Y18Stop, NSP5/Δ180-198 segments were used instead of
pT$_7$-NSP5. At 24 h post-transfection, MA-NSP5 cells (5 × 10^4 cells) were added
to transfected cells to provide a functional NSP5 for the virus rescue. The cells
were co-cultured for 3 days in FBS-free medium supplemented with trypsin (0.5
µg/mL) (Sigma Aldrich). After incubation, transfected cells were lysed by freeze-
thawing and 200 µl of the lysate was transferred to fresh MA-NSP5 cells. After
adsorption at 37°C for 1 hour, cells were washed three times with PBS and
further cultured at 37°C for 4 days in FBS-free DMEM supplemented with 0.5
µg/mL trypsin (Sigma Aldrich, 9002-07-7) until a clear cytopathic effect was
visible. The recombinant viruses were then checked by RT-PCR.
292
Immunofluorescence microscopy
Immunofluorescence experiments were performed using µ-Slide 8 Well Chamber
Slide-well (iBidi GmbH, Munich, Germany) and the following antibody dilutions:
anti-NSP5 guinea pig serum 1:1,000; anti-NSP2 guinea pig serum 1:200; anti-
VP2 guinea pig serum 1:500; anti-VP6 mouse monoclonal antibody 1:1,000;
Alexa Fluor 488-conjugated anti-mouse, 1:500 (Life Technologies), and TRITC-
conjugated anti-guinea pig, 1:500 (Life Technologies).
300
5-Ethynyl-Uridine (EU) labeling
Newly synthesized RNAs were labeled by including 2 mM 5-ethynyl uridine (EU)
into the cell culture medium, and modified incorporated nucleotides were reacted
with an azide-conjugated fluorophore Alexa-488 following the manufacturer's
protocol for Click-iT RNA Alexa Fluor 488 imaging kit (Thermo Fisher Scientific).
Cell nuclei were stained with ProLong™ Diamond Antifade Mountant with 4',6-
diamidino-2-phenylindole (DAPI, Thermo Scientific). Samples were imaged using
a confocal setup (Zeiss Airyscan equipped with a 63x, NA=1.3 objective), and the
images were processed using ZEN lite software.

RNA Fluorescence in situ Hybridization (FISH)

Rotavirus-infected MA104 cells were fixed with 4% (v/v) paraformaldehyde in nuclease-free Dulbecco's phosphate saline buffer (DPBS) for 10 min at room temperature. Samples were washed twice with DPBS, and then permeabilized with 70% (v/v) ethanol in RNAse-free water at +4°C for at least 1 hour prior to hybridization. Permeabilized samples were re-hydrated for 5 min in a pre-hybridization buffer (300 mM NaCl, 30 mM trisodium citrate, pH 7.0 in nuclease-free water, 10 % v/v formamide, supplemented with 2 mM vanadyl ribonucleoside complex). Re-hydrated samples were hybridized with 62.5 nM of an equimolar mixture of Cy3-labelled DNA probes designed to target the coding region of the gene segment 6 of simian rotavirus A/SA11 (Genbank Acc. AY187029.1) using Stellaris Probe Designer v2 software (LCG Biosearch Technologies), in a total volume of 200 μl of the hybridization buffer (Stellaris RNA FISH hybridization buffer, SMF-HB1-10, Biosearch Technologies, supplemented with 10% v/v of deionized formamide). After 4-8 hours of incubation at 37°C in a humidified chamber, samples were briefly rinsed with the wash buffer (300 mM NaCl, 30 mM trisodium citrate, pH 7.0, 10 % v/v formamide in nuclease-free water), after which a fresh aliquot of 300 μl of the wash buffer was applied to each sample and incubated twice at 37°C for 30 min. After 2 washes, nuclei were briefly stained with 300 nM DAPI solution in 300 mM NaCl, 30 mM trisodium citrate, pH 7.0, and the samples were finally rinsed with and stored in the same buffer without DAPI prior to imaging.

Transmission electron microscopy. MA104 cells were seeded at 1×10^5 cells in 2 cm^2 wells onto sapphire discs and infected at MOI of 75 FFU/cell. At 10 hpi, cells were fixed with 2.5% glutaraldehyde in 100 mM Na/K phosphate buffer, pH 7.4 for 1 h at 4°C and kept in that buffer overnight at 4°C. Afterward, samples were postfixed with 1% osmium tetroxide in 100 mM Na/K phosphate buffer for 1h at 4°C and dehydrated in a graded ethanol series starting at 70%, followed by two changes in acetone, and embedded in Epon. Ultrathin sections (60 to 80 nm)

were cut and stained with uranyl acetate and lead citrate (33). Samples were

analysed in a transmission electron microscope (CM12; Philips, Eindhoven, The

Netherlands) equipped with a charge-coupled-device (CCD) camera (Ultrascan

1000; Gatan, Pleasanton, CA, USA) at an acceleration of 100 kV.

λ-Protein phosphatase and Calf Intestinal Alkaline Phosphatase Assays

Cellular extracts were incubated with 2,000 units of λ-protein phosphatase (λ-

Ppase) (New England Biolabs) in 50 mM Tris-HCl (pH 7.5), 0.1 mM EDTA, 5 mM

DTT, 0.01% Brij 35, and 2 mM $MnCl_2$. The mixture was incubated at 30°C for 2 h.

Samples were loaded in SDS-PAGE and analysed by Western blotting.

For the Calf Intestinal Alkaline Ppase (CIP) assay, cellular extracts were

incubated with 1,000 units of CIP (New England Biolabs) in CutSmart™ reaction

buffer (New England Biolabs) and incubated at 30°C for 2h. Samples were

subjected to SDS-PAGE and analysed by Western blotting.

Electrophoresis of Viral dsRNA Genomes.

Cells were infected with the recombinant viruses at MOI of 5 and were lysed 16

hours post infection. Total RNA was extracted from lysed cells with RnaZol[®]

(Sigma-Aldrich) according the manufacturer's protocol and the dsRNA segments

were resolved on 10% (wt/vol) poly-acrylamide gels (PAGE) for 2 hours at 180

Volts and visualized by ethidium bromide staining (1μg/ml).

Protein Analysis

Proteins derived from rRVs-infected cellular extract were separated on an SDS-

PAGE for 2 hours at 35 mA and transferred to polyvinylidene difluoride

membranes (Millipore; IPVH00010) for 1 hour and 30 minutes at 300 mA (24).

For protein analysis membranes were incubated with the following primary

antibodies: anti-NSP5 (1:5,000) (16), anti-VP2 (1:5,000) (10), anti-NSP2 guinea

pig sera (1:2,000). The membranes were then incubated with the corresponding

horseradish peroxidase (HRP)-conjugated goat anti-guinea pig (1:10,000;

Jackson ImmunoResearch). Mouse HRP-conjugated anti-actin mAb (1:35,000)

(clone AC-15, Sigma-Aldrich) was used as loading control.

Signals were detected using the enhanced chemiluminescence system (Pierce

ECL Western blotting substrate; Thermo Scientific).

Statistics used

Statistical analysis and plotting were performed using GraphPad Prism 6

software (GraphPad Prism 6.0, GraphPad Software Inc., La Jolla, CA, USA).

Error bars represent standard deviation. Data were considered to be statistically

significant when $p < 0.05$ by Student's t test.

RESULTS

Generation of gs11 mutated recombinant RVs

Recombinant rotaviruses (simian rotavirus A strain SA11) carrying different

mutations in the NSP5 coding region of gs11 were obtained using a recently

developed reverse genetics protocol (29). Two essential additional modifications

were introduced to *trans*-complement the potential loss of NSP5 function in the

rRV mutants: i) an additional T7-driven plasmid encoding the ORF of wt NSP5

(i.e., without gs11 5' and 3' UTRs) was included in the transfection step of BHK-

T7 cells; and ii) each rescued rRV was amplified in a stable transfectant cell line

MA-NSP5, supplying the wt NSP5 in trans. Crucially, we have successfully

established a novel MA-NSP5 cell line to support the replication of NSP5-

deficient recombinant viruses to enable their further in-depth characterisation.

Both steps (i) and (ii) were absolutely required for rescuing the NSP5 KO virus,

confirming the essential role of NSP5 for RV replication. These mutants, as well

as additional stable cell lines generated for this study as described in Materials

and Methods, are summarised in Table 1.

<u>**rRV-NSP5/KO**</u>

NSP5 expression and localisation to viroplasms in virus-infected cells have been

considered essential for virus replication (12, 13, 25, 34). Previous studies, using

siRNA targeting gs11 mRNA have shown strong impairment of RV replication

(12, 13). In order to investigate the effects of point mutations and deletions within

the NSP5 gene, we took advantage of the established trans-complementing

MA104 cell line stably expressing NSP5. In addition, we also made two cell lines

expressing NSP2-mCherry and NSP5-EGFP fusions, which are rapidly and

efficiently recruited into viroplasms upon virus infection (Table 1) (7, 22).

Here we provide direct demonstration of the role of NSP5 in RV replication, using

an NSP5 knock out rRV (termed rRV-NSP5/KO) generated by reverse genetics.

To rescue the NSP5/KO strain, a stop codon at position 18Y was introduced (Fig.

1A-B). Analysis of MA104-virus-infected cell extracts confirmed the presence of

NSP2 and VP2, but not of NSP5 (Fig. 2A). Moreover, we did not detect

viroplasms containing NSP2, VP2 or VP6 in both MA104-infected cells and in

stable transfectant cell lines expressing the fluorescent fusion proteins NSP2-

mCherry or NSP5-EGFP (Fig. 2B). Interestingly, this indicates that the NSP5-

EGPF fusion protein is not able to trans-complement the lack of NSP5, and

indeed the rRV-NSP5/KO strain does not replicate in MA-NSP5-EGFP cells.

Furthermore, both genomic dsRNA synthesis and infectious progeny virus

production were completely abrogated in MA104 cells, but not in the trans-

complementing MA-NSP5 cell line (Fig. 2C). Together, these data confirm that

NSP5 is essential for RV replication.

rRVs with impaired NSP5 phosphorylation

To address the role of NSP5 hyperphosphorylation, we then generated a number

of rRVs harbouring NSP5 mutations previously known to impact NSP5

phosphorylation. We first generated an rRV carrying an S67A mutation (rRV-

NSP5/S67A) (Table 1 and Fig. 3A-B). Its replication in wild type MA104 cells was

strongly impaired (Fig. 3B right panel), resulting in approximately a 100-fold

reduction of the infectious progeny virus titre at different time points post-infection

(Fig. 3C). Despite the overall reduction of replication fitness, the rRV-NSP5/S67A

mutant virus was stable after 10 passages in the wild-type MA104 cells,

confirmed by the sequencing of the progeny virus. Consistent with our previous

results, the NSP5-S67A mutant was not hyper-phosphorylated in all cell lines

tested, including MA104, U2OS and Caco-2 cells (Fig. 3D), further confirming the

role of Ser67 in the initiation of NSP5 phosphorylation cascade (24). While the wt

rRV yielded multiple hyper-phosphorylated NSP5 isoforms, the NSP5/S67A

mutant mostly produced a single, homogeneous form of NSP5 with an apparent

mass of 26 kDa that could be detected in the virus-infected cell extracts at 5 or

10 hpi (Fig. 3E). Enzymatic de-phosphorylation with λ-Ppase and alkaline Calf

Intestinal Ppase (CIP), previously used to discriminate phosphorylated from non-

phosphorylated NSP5 (19), further corroborated the observed lack of

NSP5/S67A phosphorylation (Fig. 3F). Because of the differences in the

molecular weight markers used, the NSP5 band with the fastest PAGE mobility

has been traditionally described as 26 kDa, and the most abundant one as 28

kDa. Since this nomenclature has been used in many publications, we have

preferred to maintain it, despite current PAGE migrations do not correspond to

the markers presently used.

We then generated two additional rRVs harbouring truncated versions of NSP5,

one lacking the 18 AA long C-terminal tail (rRV-NSP5/ΔT), and the second one

with a 5 AA deletion (176-YKKKY-180) just upstream of the tail region (rRV-

NSP5/Δ176-180) (Table 1 and Fig. 1A-B). Despite the presence of Ser67, and

the lack of Thr and Ser residues within the deletions, both mutants did not show

a classical hyperphosphorylation pattern (Fig. 4A) and failed to replicate,

confirmed by the absence of *de novo* synthesis of genomic dsRNA (Fig. 4B).

We also investigated viroplasm formation in cells infected with these rRV

mutants. At early infection (5 hpi), the rRV-NSP5/S67A mutant produced

structures resembling viroplasms that appeared smaller and more

heterogeneous in shape compared to the regular, spherical ones produced

during the wt rRV infection (Fig. 5A upper panel and Fig. 5B). Remarkably,

during late infection (10-12 hpi), the rRV-NSP5/S67A mutant produced multiple

NSP5-containing aberrant structures, as well as fibre-like structures that became

more apparent during late infection (12 hpi) (Fig. 5A lower panel and Fig. 5B).

Interestingly, these structures produced by rRV-NSP5/S67A were
morphologically similar to those observed during wt RV infection of MA104 cells
silenced for cellular kinase CK1α, previously shown to be required for NSP5
phosphorylation (25). NSP2-mCherry and NSP5-EGFP fusion proteins were also
recruited to both types of these structures (Fig. 5C), suggesting that the observed
lack of phosphorylation does not affect NSP2/NSP5 interactions. Furthermore,
the NSP5/S67A mutant and additional viroplasmic proteins NSP2, VP6 and VP2
(Fig. 5D) were all present in the aberrant structures, suggesting that they could
also represent sites of virus replication.

In contrast, no viroplasms containing NSP5, NSP2 or VP2 were observed when
MA104 cells were infected with the rRV-NSP5/ΔT and rRV-NSP5/Δ176-180
deletion mutants. Interestingly, these mutants yielded fibre-like structures
containing only VP6 protein (Fig. 5E-F). Similar VP6-fibres are normally formed
when VP6 is over-expressed in cells in the absence of other viral proteins (35,
36).

Taken together, these results confirm the role of NSP5 hyper-phosphorylation for
controlling the assembly of regular-shaped viroplasms, highlighting the key role
of the C-terminal tail in the formation of RV viral factories.

RNA accumulation in aberrant structures

Having examined the viral protein composition of the aberrant viroplasms formed
during infection with rRVs exhibiting impaired NSP5 phosphorylation, we then
assessed their RNA content. Viral RNA transcripts were labelled by incorporation
of 5-ethynyl uridine (5-EU) in actinomycin D-treated RV-infected cells, and total
viral ssRNA was visualized by reacting with Alexa-488-azide, as described in
Materials and Methods. As expected, most viral transcripts localised in
viroplasms of rRV-wt infected cells (Fig. 6A), consistent with the roles of
viroplasms in supporting the viral replication and assembly. In contrast, no viral
RNA transcripts could be detected in the aberrant structures in both wt MA104 or
MA-NSP2-mCherry cells infected with the rRV-NSP5/S67A mutant (Fig. 6A).

RNA accumulation in viroplasms was instead restored when infecting MA-NSP5
cells that supply wild type NSP5 in *trans* (Fig. 6A-lower panel).

Although we did not detect RNA in aberrant structures (or viroplasms) in rRV
NSP5/S67A-infected MA104 cells using 5-EU staining, this mutant could still
replicate (Fig. 3C). We then explored whether rRV-NSP5/S67A transcripts could
accumulate in viroplasms, albeit with much lower efficiency, i.e., beyond the
sensitivity limit of 5-EU staining. For this purpose, we used single molecule RNA
fluorescence in situ hybridisation (smFISH) to identify the sites of RV
transcription. At 10 hpi, abundant gs6 transcripts could be detected in all
viroplasms identified in MA-NSP5-EGFP cells infected with the rRV-wt (Fig. 6B).
Conversely, the rRV-NSP5/S67A-infected cells had sparse EGFP-tagged
structures, but less than 20% of them contain gs6 RNA (Fig. 6B-C). Interestingly,
the less frequently occurring rod-like aberrant structures also showed gs6 RNA
accumulation, further suggesting that these structures could represent
replication-functional organelles.

In contrast, smFISH performed on cells infected with the rRV-NSP5/ΔT showed
diffuse distribution of gs6 RNA that did not localise to any structures resembling
viroplasms (Fig. 6B), also failing to support the virus genome replication (Fig.
4B).

We then examined the ultrastructures of viroplasms with altered morphologies in
the RV mutant rRV-NSP5/S67A using electron microscopy. Upon infection with
rRV-wt (Fig. 7A, left panel), multiple membrane-less electron dense inclusions
encircled by the well-defined endoplasmatic reticulum (ER) filled with triple-
layered particles were present in cells. At late infection points (10 hpi), filled with
triple-layered particles (TLPs) ER appeared to adopt a more tubular morphology,
suggesting a successive step in the virus egress. In contrast, the rRV-
NSP5/S67A-infected cells contained only few immature viroplasms that lacked
the ER network filled with TLPs (Fig. 7A, right panel). Only few immature
particles containing transient lipid membranes could be identified in cells infected
with rRV-NSP5/S67A mutant (Fig. 7A, left panel). Furthermore, the observed
immature viroplasms also appeared to be less-electron dense, likely due to their

lower RNA composition and, subsequently, decreased number of available

phosphate groups that bind UO^{2+} ions during the EM staining procedure.

Together, these data strongly support the role of NSP5 phosphorylation in

maintaining the viral RNA production and genome replication in viroplasms.

The mechanism of NSP5 hyper-phosphorylation

We have previously proposed a model of the hierarchical NSP5

hyperphosphorylation associated with the assembly of viroplasms that involves a

three-step mechanism: i) initial interaction of non-phosphorylated NSP5 with

NSP2 (or VP2); ii) phosphorylation of Ser67 by CK1α. This step does not take

place when NSP5 is expressed alone; iii) hyper-phosphorylation of NSP5

triggered by Ser67 phosphorylation that requires the 18 AA long C-terminal tail

(24).

Here, we have investigated the phosphorylation mechanism of NSP5 during RV

infection using a number of NSP5 phosphorylation-negative rRV strains and

MA104-derived stable transfectant cell lines (Table 1). We demonstrated that

despite the presence of Ser67, deletion mutant NSP5/ΔT was not phosphorylated

and failed to form viroplasms. We have previously shown that co-expression of

NSP5/ΔT was also unable to trigger the phosphorylation cascade of NSP5/S67A,

while other NSP5 mutants referred hereafter as activators of phosphorylation,

e.g., NSP5/Δ3 did (24). Interestingly, upon co-infection with two rRVs

NSP5/S67A and NSP5/ΔT, both mutated NSP5 variants were not phosphorylated

(Fig. 8A) (8, 19, 24, 26). This result was supported further by infecting the MA104

transfectant cell line stably expressing NSP5-ΔT (MA-ΔT) with the rRV-

NSP5/S67A strain (Table 1 and Fig. 8B, lanes 3-4). The NSP5-Δ176-180 mutant

was also unable to induce phosphorylation of NSP5/S67A following the co-

infection with the two rRVs, despite both containing Ser67 and the C-terminal tail.

This result suggests that the 'activator' NSP5 needs to be also hyper-

phosphorylated (Fig. 8C).

In MA-ΔT cells infected with rRV-NSP5/S67A mutant strain aberrant viroplasms

were produced (Fig. 8E), similar to those seen in wt MA104 cells, which did not

support virus replication compared to the wt virus (Fig. 8F).

In contrast to NSP5/ΔT, the NSP5 deletion mutant lacking amino acids 80-130,

(NSP5/Δ3) becomes hyper-phosphorylated when expressed alone and can

function as an activator of the NSP5/S67A phosphorylation (19, 24). We

therefore asked whether an MA104 stable cell line expressing the deletion

mutant NSP5/Δ3 (MA-Δ3) and infected with the rRV-NSP5/S67A strain was able

to trigger hyperphosphorylation of NSP5/S67A and as a consequence, sustain

replication of the mutant rRV strain. As shown in Fig. 8B (lanes 5-6) NSP5/S67A

was hyper-phosphorylated in presence of NSP5/Δ3 mutant and confirmed by the

λPpase treatment (Fig. 8D), although it did not completely rescue the

phosphorylation pattern of NSP5 normally observed in rRV-wt infection. When

loading large amounts of NSP5/S67A, we have occasionally observed a second

faint band with reduced mobility, as in Fig. 8B, lane 2 (indicated with *).

Interestingly, regular round-shaped structures resembling viroplasms, containing

NSP5, NSP2 and VP2 with peripheral localisation of VP6, were recovered in

these cells infected with rRV-NSP5/S67A, and yet viral replication was

nevertheless impaired (Fig. 8E-F). Consistently, the electron microscopy images

showed structures of aberrant viroplasms similar to those obtained in MA104 wt

cells (Fig. 7B, right panel).

The NSP5 phosphorylation negative mutants of the two other rRVs (NSP5/ΔT

and NSP5/Δ176-180), did not undergo hyper-phosphorylation in MA-ΔT or MA-Δ3

cell lines (Figs. 9A-B). In both cases, viroplasms formation and virus replication

were not rescued (Figs. 9C-D and 9E-F). Interestingly, apparently normal round-

shaped structures were observed in MA-Δ3 cells, which, however, recruited

significantly less VP6 (Fig. 9C-D). In addition, the previously observed VP6 spiky

structures were not detected in these cells (Figs. 9C-D). Similar results were

obtained with the rRV-NSP5/KO virus strain, which in MA-Δ3 cells did not

replicate and also showed spherical structures that contained NSP5 and NSP2,

but not VP6 (Fig. 10A-C).

Taken together, our data support a model of NSP5 hyper-phosphorylation, which
absolutely requires the presence of the C-terminal tail, and the AA residues 176-
180. Furthermore, NSP5 hyperphosphorylation requires Ser67 phosphorylation
to initiate the phosphorylation cascade, thus playing a key role in the assembly of
replication-competent viral factories.
589

DISCUSSION
Rotaviruses replicate within cytoplasmic viral factories, or viroplasms. Most RV
assembly intermediates, i.e., single layered particles (cores) and double-layered
particles (DLPs) are primarily concentrated in viroplasms. Following budding of
DLPs into the lumen of the endoplasmatic reticulum, the immature particles
acquire a transient envelope, as well as the outer capsid proteins VP4 and VP7,
resulting in a mature triple-layered virion. Moreover, downregulation of
expression of the most abundant viroplasm-forming proteins NSP5 and NSP2
severely impacts on the formation of viroplasms and production of virus progeny
(13, 15, 25, 34). In light of these observations, viroplasms have long been
recognised as essential compartments supporting the RV replication.
Of the two non-structural proteins involved in viroplasm assembly, NSP5 appears
to play a crucial role by potentially providing a scaffold that allows for recruitment
of additional viral proteins. Only when NSP5 is co-expressed with NSP2 and/or
VP2, these proteins assemble into the viroplasm-like structures (VLS), which are
also capable to recruit additional structural proteins including VP1, VP2 and VP6
(8, 10). Given these observations, we hypothesised that complete removal of
NSP5 would be lethal for RV replication. Using a modified reverse genetics
system for rotaviruses (29), here we provide the first direct evidence of the
essential role of NSP5 in viroplasm formation and viral replication. In order to
characterise replication-deficient NSP5-negative mutants, we have established a
trans-complementing system that provides NSP5 to the virus both transiently in
BHK-T7 cells, and stably in NSP5-producing MA104 cell line (MA-NSP5), thus
enabling facile isolation of rRVs lacking functional NSP5. Using this approach,
we have demonstrated that NSP5-deficient rRV was unable to form viroplasms

and replicate in the wt MA104 cells, while the viroplasm formation and viral
replication were efficiently rescued in the trans-complementing MA-NSP5 cell
line. Interestingly, the rRVs generated using this method also failed to
incorporate dsRNA originating from the NSP5-encoding mRNA lacking the 5' and
3' untranslated regions (UTRs), further suggesting the essential roles of UTRs for
genome packaging in RVs (37, 38).

NSP5 hyper-phosphorylation has been previously implicated in the regulation of
NSP5 assembly into viroplasms. This phosphorylation, however, requires the
interaction of NSP5 with, either NSP2 or VP2, as NSP5 is not phosphorylated
when expressed alone (8, 22, 23). Previous studies suggest that activation of
NSP5 hyper-phosphorylation may require a conformational change that leads to
its efficient hyper-phosphorylation via a positive feedback loop mechanism (8, 23,
24). Two regions comprising the N-terminal amino acids 1-33 (region 1) and the
central region amino acids 81-130 (region 3) have been reported to prevent
NSP5 phosphorylation in the absence of other viral proteins, while the 18 amino
acids long C-terminal tail was found to be essential for its phosphorylation (8, 23,
24).

Here, we have shown that all three rRV mutants S67A, ΔT and Δ176-180
expressing the phosphorylation-negative NSP5 variants were unable to form
round-shaped viroplasms upon infection of MA104 cells. Interestingly, further
analysis of these mutants reveals some key differences between each NSP5
variant. While rRV-NSP5/S67A strain formed aberrant structures resembling
viroplasms that poorly support RV replication, this variant was still capable of
producing the infectious progeny, in contrast to the two other rRV mutant strains.
Interestingly, the phenotype observed with the NSP5 mutant S67A was
essentially the same as the one previously reported with the wt virus infecting
MA104 cells silenced for expression of CK1α, which is involved in
phosphorylating Ser67 and initiating the hyper-phosphorylation cascade (24, 25).
It has recently been shown that CK1α is also involved in phosphorylating NSP2,
controlling the formation of rotavirus viral factories (39). Our data obtained with
the rRV-NSP5/S67A mutant strongly suggest that the lack of NSP5 hyper-

phosphorylation determines both the morphogenesis of viroplasms and their
capacity to support RV genome replication. We cannot rule out the role of NSP2
phosphorylation in assembly of viroplasms since NSP2 is also likely to be
phosphorylated by CK1α upon infection of the rRV-NSP5/S67A strain. Despite
the formation of aberrant structures resembling viroplasms, the amount of RNA
produced within those structures in rRV-NSP5/S67A-infected MA104 cells was
practically below the detection limit of the 5-EU labelling, while the RNA
replication was fully rescued in the trans-complementing MA-NSP5 cell line (Fig.
5). It is unlikely that the viral mRNAs produced in MA104 wt cells infected with
the rRV-NSP5/S67A strain was degraded faster, as most of the ssRNA
synthesised is a consequence of the secondary round of transcription from the
newly made dsRNA-containing particles. Indeed, very low amounts of dsRNA
were detected during the infection of MA104 cells. smFISH results confirmed the
presence of small amounts of RV (+)ssRNA in some of these aberrant structures.
This result is consistent with the finding that the rRV-NSP5/S67A strain did
replicate, albeit at much lower levels than the rRV-wt. Thus, these structures are
likely to sustain virus replication with decreased efficiency, which was further
confirmed by the electron microscopy analysis of MA104 cells infected with the
rRV-NSP5/S67A strain.

The important role of NSP5 hyper-phosphorylation was further supported by the
results obtained with the two phosphorylation-negative mutant strains NSP5/ΔT
and NSP5/Δ176-180 that possess Ser67 and yet failed to form viroplasms in
MA104 cells. Surprisingly, the Δ176-180 mutant, containing the C-terminal tail
has also failed to form viroplasms in MA104 cells, despite the absence of Ser or
Thr within the chosen 176-180 region. Both phosphorylation-negative mutants
tested did not replicate in MA104 cells and we could not detect any structures
containing viral RNA.

Using the NSP5 mutant strains described above and the established stable
transfectant MA104 cells we were able to investigate the molecular mechanism
that leads to NSP5 hyperphosphorylation. We showed that the NSP5/S67A
mutant from the rRV was indeed hyperphosphorylated, albeit not completely,

when infecting MA-Δ3 cells, restoring the round-shape morphology of the
structures resembling viroplasms with a complete absence of the aberrant
structures observed in MA104 cells. This finding strongly suggests that
impairment of NSP5 phosphorylation is the direct cause of the formation of the
aberrant structures in the cytosol of the infected cell. Despite the fact that these
structures appeared morphologically similar to the classical round-shaped
viroplasms in MA-Δ3 cells, the presence of VP6 around these structures was only
observed with the hyperphosphorylated NSP5/S67A mutant, in contrast to the
other NSP5 phosphorylation-negative mutants. One possibility is that, during RV
infection, accumulation of VP6 in round-shaped viroplasms requires a full-length
hyperphosphorylated NSP5, as well as phosphorylation of multiple serine
residues likely by CK2 (17, 26). Moreover, the round-shaped structures found in
MA-Δ3 cells infected with rRV-NSP5/KO failed to contain VP6.

The observed failure to rescue the replication of the rRV-NSP5/S67A strain to the
wt levels in MA-Δ3 cells could be the consequence of the incomplete recovery of
the complex pattern of phosphorylated isoforms of wt NSP5. This suggests that
some intermediate isoforms might be important for the formation of fully
functional replication-competent viroplasms.

We propose a model of the complex hierarchical mechanism of NSP5 hyper-
phosphorylation during RV infection (Fig. 11). It involves (a) interaction of NSP5
with either NSP2 or VP2, required to (b) make Ser67 available for CK1α
phosphorylation. This initial step is then sequentially completed (c) by CK2-
mediated phosphorylation of other serines to generate the NSP5 'activator',
during a step dependent (d) on the interaction of the Ser67-phosphorylated
molecules with the non-phosphorylated partners in the NSP5 oligomeric
complexes. Alternatively (c'), oligomers could be formed before the activation
step. NSP5 interactions mediated by the carboxy-terminal tails T result in (e)
substrate activation and a fully hyperphosphorylated NSP5. This process leads to
(f) the assembly of viroplasms scaffolds containing NSP2, and (g) recruitment of
VP6, as well as the other viroplasmic proteins, VP1, VP2, VP3 to assemble
replication-competent viroplasms.

The hierarchical phosphorylation of proteins appears to be a common
mechanism regulating many cellular processes. In mammalian cells, hierarchical
phosphorylation has been described for β-catenin, in which Ser 45
phosphorylated by CK1α primes it for hyperphosphorylation by glycogen
synthase kinase-3 (GSK-3) (40, 41), which triggers its ubiquitination and
proteasomal degradation (42). A similar phosphorylation mechanism has recently
been described for a non-structural protein NS5A of the Hepatitis C Virus (HCV),
with the hyper-phosphorylation cascade primed by the initial phosphorylation of
Serine 225 by CK1α, and the subsequent phosphorylation of neighbouring
residues involving other kinases (43, 44). Moreover, the NS5A phosphorylation
was shown to play a key role in controlling the establishment of replication
complexes during HCV infection (45). Similarly, a number of other non-structural
viral proteins have been shown to undergo multiple phosphorylation events
during virus infection, suggesting that this complex post-translational
modifications play a pivotal role in orchestrating the assembly of replication-
competent viral factories (46–48).
724
725

ACKNOWLEDGEMENTS
We thank John T. Patton, Asha Ann Philipp and Jin Dai (Indiana University,
Bloomington, USA), Yuta Kanai and Takeshi Kobayashi (Osaka University,
Osaka) and Albie Van Dijik (North-West University, South Africa) for helpful
discussions on the establishment of the reverse genetics system. We are grateful
to Naoto Ito for providing BHK-T7 cells.

FIGURE LEGENDS

Figure 1. A) Schematic representation of mutations in gs11 of the corresponding

rRV strains. Sequences mutated or deleted in NSP5 are indicated. B), Profiles of

viral dsRNAs of the different rRV strains generated grown in MA-NSP5 cells.

gs11, indicated in red with *.

Figure 2. Characterisation of rRV-NSP5/KO. A) Western blot of MA104 cells

extracts infected with rRV-wt and rRV-NSP5/KO strains (MOI of 1 FFU/cell). B)

Confocal immunofluorescence microscopy of MA104, MA-NSP2-mCherry and

MA-NSP5-EGFP cells infected with rRV-wt or rRV-NSP5/KO (MOI of 1 FFU/cell),

using antibodies for NSP5, NSP2, VP2 and VP6, as indicated. Scale bar, 15 μm.

C) Electrophoretic migration pattern of dsRNAs extracted from rRV-NSP5/KO

strain grown in MA-NSP5 or MA104 cells. Genome segments 1-11 are indicated

to the left.

Figure 3. Characterisation of rRV-NSP5/S67A. A) Schematic representation of

rRV-NSP5/S67A gs11 and sequence of NSP5 wt and S67A mutant (highlighted).

B) Electrophoretic pattern of dsRNA genome segments of rRV-wt and rRV-

NSP5/S67A strains grown in MA-NSP5 cells (left panel) and in MA104 wt cells

(right panel). C) Replication kinetics of rRV-wt and rRV-NSP5/S67A in MA104

cells. Data are expressed as the means +/- standard deviations (n=3); ***, $p <$

0.001 (Student's t test). D) Western blot of extracts of U2OS, Caco-2 and MA104

cells infected with rRV-wt or rRV-NSP5/S67A strains. E) Western blot of NSP5

phosphorylation pattern in MA104 cells infected with rRV-wt or rRV-NSP5/S67A

(MOI of 1 FFU/cell) at 5 and 10 hpi. F) Western blot of λ-Ppase and alkaline

Ppase treatment of lysates of MA104 cells infected with rRV-wt or rRV-
NSP5/S67A. Protein bands corresponding to NSP5 are shown.

Figure 4. Characterisation of rRV-NSP5/ΔT and rRV-NSP5/Δ176-180. A)
Western blot of NSP5 and NSP5 mutants from MA104 cells infected with rRV-wt,
rRV-NSP5/ΔT or rRV- NSP5/Δ176-180 at 5 hpi. B) Electrophoretic dsRNA
migration pattern of rRV-NSP5/ΔT and rRV-NSP5/Δ176-180 grown in MA-NSP5
and MA104 cells.

Figure 5. Viroplasms morphology in MA104 cells infected with rRV-NSP5 wt
and mutants. Representative confocal immunofluorescence micrographs of
MA104 cells infected with rRV-wt and rRV-NSP5/S67A (MOI of 15 FFU/cell) at 5
hpi (upper panels) or 10 hpi (lower panels). Cells were stained with anti-NSP5
and DAPI. B) Quantitative analysis of viroplasms size (μm^2) of MA104 cells
infected with rRV-wt and rRV-NSP5/S67A at 5 hpi and 10 hpi. ***, $p < 0.001$; **, p
< 0.01. C) Confocal immunofluorescence micrographs of MA-NSP5-EGFP and
MA-NSP2-mCherry cells infected with rRV-NSP5/S67A (MOI of 15 FFU/cell). D)
Confocal immunofluorescence micrographs of rRV-wt and rRV-NSP5/S67A
infected MA104 cells (MOI 15 FFU/cell) and stained at 10 hpi with the indicated
antibodies. E), F), Confocal immunofluorescence of MA104 cells infected with
rRV-NSP5/Δ176-180 (E) or rRV-NSP5/ΔT (F) and stained with the indicated
antibodies and DAPI. Scale bar, 15 μm.

Figure 6. Viral RNA detection in rRV NSP5/S67A infected cells. A)
Representative confocal immunofluorescence micrographs of MA104 (upper
panel), MA-NSP5 (middle panel) and MA-NSP2-mCherry cells (lower panel)
infected with rRV-wt or rRV-NSP5/S67A strains and stained with anti-NSP5 (red,
in MA104 and MA-NSP5 cells) and EU (green). B) Single molecule RNA
Fluorescence In Situ Hybridisation (smFISH) on MA-NSP5-EGFP cells infected
with rRV-wt, rRV-NSP5/S67A or rRV-NSP5/ΔT strains. Viroplasms detected with
NSP5-EGFP (green) and viral RNA (probe specific for gs6 was Cy3-conjugated,

red). Colocalising viroplasms and RNAs are indicated by white arrows. Scale bar,

15 μm. C) Quantitative analysis of NSP5-EGFP positive structures (viroplasms)

for RNA (gs6) in MA-NSP5-EGFP cells infected with rRV-wt or rRV-NSP5/S67A.

***, p < 0.001

Figure 7. Electron microscopy of cells infected with rRV-NSP5/S67A. High–

definition electron micrographs of MA104 (A) and MA-Δ3 (B) cells infected with

rRV-wt (left panel) and rRV-NSP5/S67A (right panel) (MOI, 75 FFU/cell). At 10

hpi, cells were fixed with glutaraldehyde and processed for transmission electron

microscopy. V, viroplasm; m, mitochondria; mt, microtubule-bundles; Nu, nucleus.

The white open box indicates the immediate right magnified image. White arrows

indicate the endoplasmatic reticulum surrounding viroplasms; black arrowheads

indicate viral particles with an envelope. Scale bar, 500 nm.

Figure 8. Phosphorylation of NSP5/S67A. A) Western blot of extracts of

MA104 cells infected with rRV-NSP5/S67A or co-infected with rRV-NSP5/S67A

and rRV-NSP5/ΔT. Blue arrowhead indicates NSP5/S67A. The faster migrating

band corresponds to NSP5-ΔT. B) Western blot of MA104, MA-ΔT and MA-Δ3

cells infected with rRV-NSP5/S67A. Blue arrowhead indicates NSP5/S67A. C)

Western blot of extracts of MA104 cells co-infected with rRV-NSP5/S67A and

rRV-NSP5/Δ176-180. Blue arrowhead indicates NSP5/S67A.   D) λ-Ppase

treatment of extracts of MA-Δ3 cells infected with the rRV-NSP5/S67A strain.

Filled and open arrowheads indicate de-phosphorylated NSP5-S67A and NSP5-

Δ3, respectively. E) Representative confocal immunofluorescence micrographs of

MA104, MA-ΔT and MA-Δ3 cells infected with rRV-NSP5/S67A. Cells were

stained with the indicated antibodies. Scale bar, 15 μm. F) Yield of infectious

virus of rRV-NSP5/S67A grown in MA-NSP5, MA104, MA-ΔT and MA-Δ3 cells at

24 hpi.*, $p < 0.05$; **, $p < 0.01$.

Figure 9. Phosphorylation of NSP5/-ΔT and NSP5/Δ176-180 and viroplasm

formation. Western blots (A, B) and confocal immunofluorescence micrographs

(C, D) of MA104, MA-ΔT and MA-Δ3 cells infected with rRV-NSP5/ΔT (A, C) or

with rRV-NSP5/Δ176-180 (B, D). Blue arrowhead indicates NSP5/ΔT and

NSP5/Δ176-180, respectively. Open arrowheads indicate NSP5-Δ3 and NSP5-

ΔT. Cells were stained with the indicated antibodies and DAPI. Scale bar, 15 μm.

E) Single step growth of rRV-NSP5/ΔT (left) and rRV-NSP5/Δ176-180 (right) in

MA-NSP5, MA104, MA-ΔT and MA-Δ3 cells, as indicated. The experiment was

terminated at 24 hpi. **, $p < 0.01$; #, Viral titer < 300 FFU/ml.

Figure 10. rRV NSP5/KO in infected cells. Western blot (A) and confocal

immunofluorescence (B) of MA104, MA-ΔT and MA-Δ3 cells infected with the

rRV-NSP5/KO strain. Cells were stained with the indicated antibodies and DAPI.

Scale bar, 15 μm. C) Single step growth of rRV-NSP5/KO in MA-NSP5, MA104,

MA-ΔT and MA-Δ3 cells, as indicated. The experiment was terminated at 24 hpi.

**, $p < 0.01$; #, Viral titer < 300 FFU/ml.

Figure 11. Model of NSP5 hyperphosphorylation and assembly of

replication-competent viroplasms. (***a***) Interaction of non-phosphorylated NSP5

with either NSP2 or VP2 is required to (***b***) induce conformational changes that

make S67 available for CK1α phosphorylation (P). This initial step of the cascade

is then (***c***) sequentially completed by phosphorylation by CK2 of other residues

including serines in domain 4, to generate the NSP5 activator or, (***c'***) a step of

interaction with non-phosphorylated molecule precedes the involvement of CK2.

(***d***) NSP5 interactions to form dimers/multimers are mediated by the carboxy-

terminal tails (T). (***e***) The primed (S67-phosphorylated) activator molecules result

in substrate activation and fully hyper-phosphorylated NSP5. (***f***) Assembly of

viroplasms scaffolds containing NSP2 serve to (***g***) recruit the other viroplasmic

components, VP1, VP2, VP3 and VP6 to assemble replication-competent

viroplasms. P; phosphorylated amino acid. *This image was created using

BioRender.com.

Table 1. Schematic representation of NSP5 and NSP2 mutants or fusion proteins

used to generate rRV and/or stable MA104 transfectant cell lines, used in this
study.

<u>REFERENCES</u>

1.   Tate JE, Burton AH, Boschi-Pinto C, Parashar UD, World Health
Organization–Coordinated Global Rotavirus Surveillance Network. 2016.
Global, Regional, and National Estimates of Rotavirus Mortality in Children
<5 Years of Age, 2000–2013. Clin Infect Dis 62:S96–S105.

2.   Parashar UD, Hummelman EG, Bresee JS, Miller MA, Glass RI. 2003.
Global illness and deaths caused by rotavirus disease in children. Emerg
Infect Dis 9:565–572.

3.   Patton JT, Silvestri LS, Tortorici MA, Vasquez-Del Carpio R, Taraporewala
ZF. 2006. Rotavirus genome replication and morphogenesis: role of the
viroplasm. Curr Top Microbiol Immunol 309:169–87.

4.   Desselberger U. 2014. Rotaviruses. Virus Res 190:75–96.

5.   Estes MK, Greenberg HB. 2013. Rotaviruses, p. 1347–1401. *In* Health,
WK, Lippincott Williams Wilkins (eds.), Fields Virology, 6th ed. Philadelpia,
PA.

6.   Clapp LL, Patton JT. 1991. Rotavirus morphogenesis: domains in the
major inner capsid protein essential for binding to single-shelled particles
and for trimerization. Virology 180:697–708.

7.   Eichwald C, Rodriguez JF, Burrone OR. 2004. Characterization of rotavirus
NSP2/NSP5 interactions and the dynamics of viroplasm formation. J Gen
Virol 85:625–634.

8.   Fabbretti E, Afrikanova I, Vascotto F, Burrone OR. 1999. Two non-
structural rotavirus proteins, NSP2 and NSP5, form viroplasm-like
structures in vivo. J Gen Virol 80:333–339.

9.   Poncet D, Lindenbaum P, L'Haridon R, Cohen J. 1997. In vivo and in vitro
phosphorylation of rotavirus NSP5 correlates with its localization in
viroplasms. J Virol 71:34–41.

10.   Arnoldi F, Campagna M, Eichwald C, Desselberger U, Burrone OR. 2007.
Interaction of Rotavirus Polymerase VP1 with Nonstructural Protein NSP5
Is Stronger than That with NSP2. J Virol 81:2128–2137.

11.   Gallegos CO, Patton JT. 1989. Characterization of rotavirus replication
intermediates: a model for the assembly of single-shelled particles.
Virology 172:616–27.

12.   Campagna M, Eichwald C, Vascotto F, Burrone OR. 2005. RNA
interference of rotavirus segment 11 mRNA reveals the essential role of
NSP5 in the virus replicative cycle. J Gen Virol 86:1481–1487.

13.   Lopez T, Rojas M, Ayala-Breton C, Lopez S, Arias CF. 2005. Reduced
expression of the rotavirus NSP5 gene has a pleiotropic effect on virus
replication. J Gen Virol 6:1609–1617.

14.   Silvestri LS, Taraporewala ZF, Patton JT. 2004. Rotavirus replication: plus-
sense templates for double-stranded RNA synthesis are made in
viroplasms. J Virol 78:7763–74.

15.   Vascotto F, Visintin M, Cattaneo A, Burrone OR. 2005. Design and
selection of an intrabody library produced de-novo for the non-structural
protein NSP5 of rotavirus. J Immunol Methods 301:31–40.

16.   Gonzalez SA, Burrone OR. 1991. Rotavirus NS26 is modified by addition
of single O-linked residues of N-acetylglucosamine. Virology 182:8–16.

17.   Sotelo PH, Schumann M, Krause E, Chnaiderman J. 2010. Analysis of
rotavirus non-structural protein NSP5 by mass spectrometry reveals a
complex phosphorylation pattern. Virus Res 149:104–108.

18.   Campagna M, Marcos-Villar L, Arnoldi F, de la Cruz-Herrera CF, Gallego
P, Gonzalez-Santamaria J, Gonzalez D, Lopitz-Otsoa F, Rodriguez MS,
Burrone OR, Rivas C. 2013. Rotavirus Viroplasm Proteins Interact with the
Cellular SUMOylation System: Implications for Viroplasm-Like Structure
Formation. J Virol 87:807–817.

19.   Afrikanova I, Miozzo MC, Giambiagi S, Burrone O. 1996. Phosphorylation
generates different forms of rotavirus NSP5. J Gen Virol 9:2059–2065.

20.   Blackhall J, Fuentes A, Hansen K, Magnusson G. 1997. Serine protein
kinase activity associated with rotavirus phosphoprotein NSP5. J Virol
71:138–144.

21.   Eichwald C, De Lorenzo G, Schraner EM, Papa G, Bollati M, Swuec P, de
Rosa M, Milani M, Mastrangelo E, Ackermann M, Burrone OR, Arnoldi F.
2018. Identification of a Small Molecule That Compromises the Structural
Integrity of Viroplasms and Rotavirus Double-Layered Particles. J Virol 92.

22.   Contin R, Arnoldi F, Campagna M, Burrone OR. 2010. Rotavirus NSP5
orchestrates recruitment of viroplasmic proteins. J Gen Virol 7:1782–1793.

23.   Afrikanova I, Fabbretti E, Miozzo MC, Burrone OR. 1998. Rotavirus NSP5
phosphorylation is up-regulated by interaction with NSP2. J Gen Virol
79:2679–2686.

24.   Eichwald C, Jacob G, Muszynski B, Allende JE, Burrone OR. 2004.
Uncoupling substrate and activation functions of rotavirus NSP5:
phosphorylation of Ser-67 by casein kinase 1 is essential for
hyperphosphorylation. Proc Natl Acad Sci U S A 101:16304–16309.

25.   Campagna M, Budini M, Arnoldi F, Desselberger U, Allende JE, Burrone
OR. 2007. Impaired hyperphosphorylation of rotavirus NSP5 in cells
depleted of casein kinase 1alpha is associated with the formation of
viroplasms with altered morphology and a moderate decrease in virus

replication. J Gen Virol 10:2800–2810.

26.   Eichwald C, Vascotto F, Fabbretti E, Burrone OR. 2002. Rotavirus NSP5:
mapping phosphorylation sites and kinase activation and viroplasm
localization domains. J Virol 76:3461–3470.

27.   Martin D, Ouldali M, Menetrey J, Poncet D. 2011. Structural organisation of
the rotavirus nonstructural protein NSP5. J Mol Biol 413:209–221.

28.   Kanai Y, Komoto S, Kawagishi T, Nouda R, Nagasawa N, Onishi M,
Matsuura Y, Taniguchi K, Kobayashi T. 2017. Entirely plasmid-based
reverse genetics system for rotaviruses. Proc Natl Acad Sci U S A
114:2349–2354.

29.   Komoto S, Fukuda S, Ide T, Ito N, Sugiyama M, Yoshikawa T, Murata T,
Taniguchi K. 2018. Generation of Recombinant Rotaviruses Expressing
Fluorescent Proteins by Using an Optimized Reverse Genetics System. J
Virol 92:e00588-18.

30.   Pertel T, Hausmann S, Morger D, Züger S, Guerra J, Lascano J, Reinhard
C, Santoni FA, Uchil PD, Chatel L, Bisiaux A, Albert ML, Strambio-De-
Castillia C, Mothes W, Pizzato M, Grütter MG, Luban J. 2011. TRIM5 is an
innate immune sensor for the retrovirus capsid lattice. Nature 472:361–
365.

31.   Dull T, Zufferey R, Kelly M, Mandel RJ, Nguyen M, Trono D, Naldini L.
1998. A third-generation lentivirus vector with a conditional packaging
system. J Virol 72:8463–71.

32.   Yusa K, Zhou L, Li MA, Bradley A, Craig NL. 2011. A hyperactive piggyBac
transposase for mammalian applications. Proc Natl Acad Sci U S A
108:1531–6.

33.   Eichwald C, Arnoldi F, Laimbacher AS, Schraner EM, Fraefel C, Wild P,
Burrone OR, Ackermann M. 2012. Rotavirus viroplasm fusion and

perinuclear localization are dynamic processes requiring stabilized
microtubules. PLoS One 7:e47947.

34.   Vascotto F, Campagna M, Visintin M, Cattaneo A, Burrone OR. 2004.
Effects of intrabodies specific for rotavirus NSP5 during the virus replicative
cycle. J Gen Virol 85:3285–3290.

35.   Ito H, Minamoto N, Goto H, Rong LT, Sugiyama M, Kinjo T. 1996.
Expression of the major inner capsid protein, VP6, of avian rotavirus in
mammalian cells. Vet Microbiol 49:257–65.

36.   Tosser G, Labbé M, Brémont M, Cohen J. 1992. Expression of the major
capsid protein VP6 of group C rotavirus and synthesis of chimeric single-
shelled particles by using recombinant baculoviruses. J Virol 66:5825–31.

37.   McDonald SM, Patton JT. 2011. Assortment and packaging of the
segmented rotavirus genome. Trends Microbiol 19:136–144.

38.   Borodavka A, Desselberger U, Patton JT. 2018. Genome packaging in
multi-segmented dsRNA viruses: distinct mechanisms with similar
outcomes. Curr Opin Virol 33:106–112.

39.   Criglar JM, Anish R, Hu L, Crawford SE, Sankaran B, Prasad BVV, Estes
MK. 2018. Phosphorylation cascade regulates the formation and
maturation of rotaviral replication factories. Proc Natl Acad Sci
115:E12015–E12023.

40.   Liu C, Li Y, Semenov M, Han C, Baeg GH, Tan Y, Zhang Z, Lin X, He X.
2002. Control of beta-catenin phosphorylation/degradation by a dual-
kinase mechanism. Cell 108:837–47.

41.   Amit S, Hatzubai A, Birman Y, Andersen JS, Ben-Shushan E, Mann M,
Ben-Neriah Y, Alkalay I. 2002. Axin-mediated CKI phosphorylation of β-
catenin at Ser 45: A molecular switch for the Wnt pathway. Genes Dev.

42.   Verheyen EM, Gottardi CJ. 2010. Regulation of Wnt/β-catenin signaling by
995   protein kinases. Dev Dyn.

996   43.   Goonawardane N, Yin C, Harris M. 2018. A pivotal role of serine 225
997   phosphorylation in the function of hepatitis C virus NS5A revealed with the
998   application of a phosphopeptide antiserum and super-resolution
999   microscopy. bioRxiv.

1000   44.   Kim J, Lee D, Choe J. 1999. Hepatitis C virus NS5A protein is
1001   phosphorylated by casein kinase II. Biochem Biophys Res Commun
1002   257:777–781.

1003   45.   Ross-Thriepland D, Mankouri J, Harris M. 2015. Serine Phosphorylation of
1004   the Hepatitis C Virus NS5A Protein Controls the Establishment of
1005   Replication Complexes. J Virol 89:3123–3135.

1006   46.   Keating JA, Bhattacharya D, Lim P-Y, Falk S, Weisblum B, Bernard KA,
1007   Sharma M, Kuhn RJ, Striker R. 2013. West Nile virus methyltransferase
1008   domain interacts with protein kinase G. Virol J 10:242.

1009   47.   Spencer K-A, Dee M, Britton P, Hiscox JA. 2008. Role of phosphorylation
1010   clusters in the biology of the coronavirus infectious bronchitis virus
1011   nucleocapsid protein. Virology 370:373–381.

1012   48.   Bhattacharya D, Hoover S, Falk SP, Weisblum B, Vestling M, Striker R.
1013   2008. Phosphorylation of yellow fever virus NS5 alters methyltransferase
1014   activity. Virology 380:276–284.

1015

Figure 1

A

gs11-NSP5/S67A

| 5'UTR | NSP5 | 3'UTR |

S67A

NSP5-wt

P S D S A S N D P
CATCTGATTCTGCTTCAAACGATCCA
185   190   195   200   205   210

S67A

NSP5/S67A

CCATCTGATTCTGCTGCAAACGATCCA
P S D S A A N D P

B

C

Viral Titer (log$_{10}$FFU/ml) vs Time post infection (h)

— rRV-wt
— rRV-NSP5/S67A

***

D

rRV-wt   rRV-NSP5/S67A
U2OS CACO-2 MA104   U2OS CACO-2 MA104   kDa

NSP5
25
actin

E

rRV

wt   NSP5/S67A   wt   NSP5/S67A
kDa

VP2
100

NSP5
25

actin

5 hpi   10 hpi

F

untreated   Alkaline-Ppase   λ-Ppase
wt NSP5/S67A   wt NSP5/S67A   wt NSP5/S67A
kDa

NSP5

actin

Figure 5

B

C

A

B

C

Construct	MA104 stable cell line	rRV
NSP5 wt	MA-NSP5	rRV-wt
NSP5/S67A		rRV-NSP5/S67A
NSP5/Δ176-180		rRV-NSP5/Δ176-180
NSP5/Δ3	MA-Δ3	
NSP5/ΔT	MA-ΔT	rRV-NSP5/ΔT
NSP5/KO		rRV-NSP5/KO
NSP5-EGFP	MA-NSP5-EGFP	
NSP2-mCherry	MA-NSP2-mCherry	