Recombinant rotaviruses rescued by reverse genetics reveal the role of NSP5 hyperphosphorylation in the assembly of viral factories

Guido Papa[1,*], Luca Venditti[1], Francesca Arnoldi[1,2], Elisabeth M. Schraner[3], Christiaan Potgieter[4,5], Alexander Borodavka[6], Catherine Eichwald[3] and Oscar R. Burrone[1,*]

[1] International Centre for Genetic Engineering and Biotechnology, ICGEB, Padriciano 99, 34149 Trieste, Italy
[2] Department of Medicine, Surgery and Health Sciences, University of Trieste, Trieste, Italy
[3] Institute of Virology, University of Zürich, Zürich, Switzerland
[4] Department of Biochemistry, Centre for Human Metabolomics, North-West University, Potchefstroom, South Africa
[5] Deltamune (Pty) Ltd, 248 Jean Avenue, Lyttelton, Centurion, South Africa
[6] Astbury Centre for Structural Molecular Biology, School of Molecular and Cellular Biology, University of Leeds, Leeds, UK

* Corresponding authors, guido.papa@icgeb.org, burrone@icgeb.org

ABSTRACT

Rotavirus (RV) is a non-enveloped RNA virus that replicates in round-shaped cytoplasmic viral factories termed viroplasms. The formation of these viral factories requires the co-expression of the two non-structural viral proteins NSP5 and NSP2.

During RV infection, NSP5 undergoes hyperphosphorylation, which is primed by the phosphorylation of a single serine residue. Despite extensive phosphorylation, the role of this post-translational modification in the formation of viroplasms and its impact on the virus replication remains obscure. Here we shed light on the role of NSP5 during RV infection by taking advantage of a modified fully tractable reverse genetics system. An NSP5 trans-complementing cell line was used to generate and characterise several recombinant rotaviruses (rRVs) with mutations in NSP5. We demonstrate that a rRV lacking NSP5, was completely unable to assemble viroplasms and to replicate, confirming its pivotal role in rotavirus replication.

A number of mutants with impaired NSP5 phosphorylation were generated to further interrogate the function of this post-translational modification in the assembly of replication-competent viroplasms. We showed that the rRV mutant strains exhibit impaired viral replication and the ability to assemble round-shaped viroplasms in MA104 cells. Furthermore, we have investigated the mechanism of NSP5 hyper-phosphorylation during RV infection using NSP5 phosphorylation-negative rRV strains, as well as MA104-derived stable transfectant cell lines expressing either wt NSP5 or selected NSP5 deletion mutants. Our results indicate that NSP5 hyper-phosphorylation is a crucial step for the assembly of round-shaped viroplasms, highlighting the key role of the C-terminal tail of NSP5 in the formation of replication-competent viral factories. Such a complex NSP5 phosphorylation cascade may serve as a paradigm for the assembly of functional viral factories in other RNA viruses.

AUTHOR SUMMARY

Rotavirus (RV) is considered the most common cause of viral gastroenteritis in children worldwide. RV double-stranded RNA genome is replicated and packaged into virus progeny in cytoplasmic structures termed viroplasms. The non-structural protein NSP5, which undergoes a complex hyperphosphorylation process during RV infection, is required for the formation of these virus-induced organelles. However, its roles in viroplasm formation and RV replication have never been directly assessed due to the lack of a fully tractable reverse genetics (RG) system for rotaviruses.

Here we show a novel application of a recently developed RG system by establishing a stable trans-complementing NSP5-producing cell line required to rescue rotaviruses with mutations in NSP5. This approach allowed us to provide the first direct evidence of the pivotal role of this protein during RV replication. Furthermore, using recombinant RV mutants we shed light on the molecular mechanism of NSP5 hyperphosphorylation during infection and its involvement in the assembly and maturation of replication-competent viroplasms.

Considering that hyperphosphorylation of proteins appears to be a common mechanism regulating many cellular processes, our work suggests a key role of these multiple phosphorylation events in orchestrating assembly of structures required for replication of other virus pathogens.

INTRODUCTION

Rotavirus (RV) is the most common cause of viral gastroenteritis in young children and infants worldwide [1,2]. It is a non-enveloped RNA virus with a genome composed of 11 segments of double-stranded RNA (dsRNA), which replicates in cytoplasmic structures primarily composed of viral proteins [3–5]. During infection, the first steps of viral morphogenesis and genome replication occur within cytoplasmic viral replication factories known as viroplasms. [3,5–7]. The assembly of viroplasms requires co-expression of at least two non-structural proteins, NSP5 and NSP2 [8,9], however, how these virus-induced organelles are formed remains unknown.

Other viral proteins also found in viroplasms include RNA-dependent RNA polymerase (RdRp) VP1, the main inner-core protein VP2, guanyltransferase/ methylase VP3, and the middle layer (inner capsid) protein VP6 [10,11]. Biochemical evidence suggests that viroplasms are essential for RV replication since the virus production is highly impaired upon silencing of either NSP2 or NSP5 [12–15].

Rotavirus NSP5, encoded by genome segment 11, is a small serine (Ser)- and threonine (Thr)-rich non-structural protein that undergoes multiple post-translational modifications in virus-infected cells, including O-linked glycosylation [16], N-acetylation [17], SUMOylation [18] and crucially hyperphosphorylation that involves several distinct Ser residues [19,20]. The NSP5 hyperphosphorylation is a complex process, which gives rise to multiple phosphorylation states ranging from the most abundant 28 kDa phospho-isoform, up to the hyperphosphorylated 32-34 kDa states [19,20]. All these forms have been found to be more stable in viroplasms, while chemical disruption of viroplasms results in NSP5 de-phosphorylation [21]. The mechanism of NSP5 phosphorylation is not yet wholly understood, but it involves interactions with other viral proteins. When expressed alone in non-infected cells, NSP5 is not phosphorylated, while co-expression with NSP2 or VP2 results in NSP5 hyperphosphorylation and formation of viroplasm-like structures (VLS) [8,22,23]. NSP5 hyperphosphorylation involves the phosphorylation of Serine 67 (Ser67) by

Casein Kinase 1α (CK1α) to initiate the phosphorylation cascade [24,25] and it is
considered to be essential for the assembly of viroplasms [26].

Although the structure of NSP5 remains unknown, it readily forms higher
molecular weight oligomeric species in solution, potentially providing a larger
interface for interacting with multiple components of viroplasms [27].

In addition, the C-terminal region ('a tail' including amino acids 180-198) is
required for NSP5 decamerisation *in vitro* [27] and VLS formation *in vivo* [7].

However, due to the lack of a fully tractable reverse genetics (RG) system for
RVs until recently, previous studies on NSP5 have been carried out using the
mutants expressed in the absence of a complete set of viral proteins. Here, we
took advantage of the novel plasmid only-based, helper-virus free RG systems
for rotaviruses [28,29] to gain new insights into the mechanisms of NSP5
hyperphosphorylation and its role in viroplasm assembly and virus replication
during viral infection.

To achieve this, we generated and characterised several viable recombinant
rotaviruses (rRVs) with mutations in NSP5. Using these mutants, we show the
essential role of NSP5 hyperphosphorylation for viroplasm assembly and in
genome replication. These studies shed light on a complex hierarchical
mechanism of NSP5 hyperphosphorylation during rotaviral infection.

MATERIALS AND METHODS

Cells and viruses. MA104 (embryonic African green monkey kidney cells,
ATCC® CRL-2378), U2OS (Human bone osteosarcoma epithelial cells), Caco-2
(colorectal adenocarcinoma human intestinal epithelial cell line, ATCC®HTB-37)
and HEK293T (embryonic human kidney epithelial, ATCC®CRL-3216) cells were
cultured in Dulbecco's Modified Eagle's Medium (DMEM) (Life Technologies)
supplemented with 10% Fetal Bovine Serum (FBS) (Life Technologies) and 50
µg/ml gentamycin (Biochrom AG).

MA104-NSP5-EGFP cells (MA-NSP5-EGFP) [7] were cultured in DMEM
supplemented with 10% FBS (Life Technologies), 50 µg/ml gentamycin
(Biochrom AG) and 1 mg/ml geneticin (Gibco-BRL, Life Technologies).
MA104-NSP2-mCherry (MA-NSP2-mCherry), MA104-Δ3 (MA-Δ3), MA104-Δtail
(MA-ΔT) and MA104-NSP5wt (MA-NSP5) stable transfectant cell lines
(embryonic African green monkey kidney cells, ATCC® CRL-2378) were grown
in DMEM (Life Technologies) containing 10% FBS, 50 µg/ml gentamycin
(Biochrom AG) and 5 µg/ml puromycin (Sigma-Aldrich).
BHK-T7 cells (Baby hamster kidney stably expressing T7 RNA polymerase) were
cultured in Glasgow medium supplemented with 5% FBS, 10% Tryptose
Phosphate Broth (TPB) (Sigma-Aldrich), 50 µg/ml gentamycin (Biochrom AG),
2% Non-Essential Amino Acid (NEAA), 1% Glutamine.
Recombinant simian RV strain SA11 (rRV-wt), rescued using reverse genetics
system using cDNA clones encoding the wild-type SA11 (G3P[2]) virus [28], was
propagated in MA104 cells cultured in DMEM supplemented with 0.5 µg/ml
trypsin (Sigma Aldrich).
173
Recombinant RVs Titration
Recombinant NSP5 mutant RVs were grown in MA-NSP5 cells and the lysate
was 2-fold serially diluted and used to infect MA-NSP5 cells, seeded in 24-wells
plates with coverslips. After 1 hour of adsorption, virus was removed, and cells
were incubated at 37°C. At 5 hours post-infection (hpi), cells were fixed with 4%
paraformaldehyde (PFA) in phosphate buffer saline (PBS) [137 mM NaCl; 2.7
mM KCl; 8.1 mM Na_2HPO_4 and 1.74 mM KH_2PO_4 pH7.5] for 15 min at room
temperature and permeabilized for 5 min with PBS containing 0.01% Triton X-
100. Next, cells were incubated for 30 min with PBS supplemented with 1%
bovine serum albumin (PBS-BSA) at room temperature and then with anti-NSP5
(1:1000) or anti-VP2 (1:200) or anti-NSP2 (1:200) guinea pig serum diluted in
PBS-BSA. After washing three times with PBS, cells were incubated for 1 h at
room temperature with TRITC-conjugated Anti-guinea Pig IgG (Jackson
ImmunoResearch) (1:500) diluted in PBS-BSA.

Nuclei were stained with ProLong™ Diamond Antifade Mountant with DAPI
(Thermo Scientifc). Samples were imaged using a confocal setup (Zeiss Airyscan
equipped with a 63x, NA=1.3 objective). Each viroplasms-containing cell was
counted as one focus-forming unit (FFU). The average of cells with viroplasms of
six fields of view per each virus dilution was determined and the total number of
cells containing viroplasms in the whole preparation was estimated. The virus
titre was determined as:

195
$$Virus\ Titer\ \left(\frac{FFU}{ml}\right) = \frac{N * Dilution\ Factor}{V(ml)}$$

where N is a total number of cells containing 1 or more viroplasms, and V is the
volume of virus inoculum added.

Replication Kinetics of recombinant viruses
MA104 cell line (ATCC CRL-2378.1) or stably transfected MA104 cells (NSP5;
Δ3, ΔT) were seeded into 24-well plates and subsequently infected with
recombinant RVs at MOI (FFU/cell) of 0.5 for multi-step growth curve
experiments and MOI 5 for a single-step growth curve experiment. After
adsorption for 1 h at 37°C, cells were washed twice with PBS and the medium
replaced with DMEM without trypsin. After incubation at 37°C, cells were
harvested after 8, 16, 24, 36 hours post virus adsorption. The cell lysates were
freeze-thawed three times and activated with trypsin (1 μg/ml) for 30 min at 37°C.
The lysates were used to infect monolayers of MA-NSP5 cells seeded in μ-Slide
8 Well Chamber Slide-well (iBidi GmbH, Munich, Germany). The cells were then
fixed 5 hours post infection for 15 min with 4% paraformaldehyde and
permeabilized for 5 min with PBS containing 0.01% Triton X-100. Next, cells
were incubated for 30 min with PBS-BSA at room temperature and then with anti-
NSP5 serum (1:1000) diluted in PBS-BSA. After three washes with PBS, cells
were incubated for 1 h at room temperature with TRITC-conjugated Anti-guinea
Pig IgG (Jackson ImmunoResearch) (1:500) diluted in PBS containing 1% BSA
(PBS-BSA).

The number of infected cells was counted, and the virus titres were expressed in
Focus-Forming Units per mL (FFU/mL).
219

Plasmid construction.
RV plasmids pT7-VP1-SA11, pT7-VP2-SA11, pT7-VP3-SA11, pT7-VP4-SA11,
pT7-VP6-SA11, pT7-VP7-SA11, pT7-NSP1-SA11, pT7-NSP2-SA11, pT7-NSP3-
SA11, pT7-NSP4-SA11, and pT7-NSP5-SA11 [28] were used to rescue
recombinant RVs by reverse genetics. pT7-NSP5/S67A carrying a mutation in the
nucleotide T220G in the gs11 and pT7-NSP5/Tyr18Stop harbouring a nucleotide
substitution T75G were generated by QuikChange II Site-Directed Mutagenesis
(Agilent Technologies). pT7-NSP5/ΔT was generated from pT7-NSP5-SA11 by
deleting the last 18 C-terminal amino acids (FALRMRMKQVAMQLIEDL) using
substitution of the F181 encoding triplet with a stop codon. pT7-NSP5/Δ176-180
were obtained deleting the amino acids 176 to 180 (YKKKY). The described
deletions were performed using the QuikChange II Site-Directed Mutagenesis kit
(Agilent Technologies).
For the generation of lentiviral plasmids, NSP5 and NSP2-mCherry were
amplified by PCR and inserted into the plasmid pAIP (Addgene #74171; [30] at
the *NotI-EcoRI* restriction enzymes sites to yield pAIP-NSP5 and pAIP-NSP2-
mCherry, which were then used to generate lentiviruses for the MA104-stable
transfectant cell lines (MA-NSP5 and MA-NSP2-mCherry) [31]. NSP5/ΔT was
amplified from the pT7-NSP5/ΔT by PCR and inserted into the pPB-MCS (Vector
Builder) at restriction enzyme sites *NheI-BamHI* to generate pPB-NSP5/ΔT for
the production of the stable transfectant MA-ΔT.
pPB-NSP5/Δ3, for MA-Δ3 stable cell line establishment, was generated with a
GenParts™ DNA Fragment (GenScript) containing NSP5 ORF lacking the amino
acids 80-130 (VKTNADAGVSMDSSAQSRPSSNVGCDQVDFSLNKG
LKVKANLDSSISIST) and inserted into the *NheI-BamHI* restriction sites of pPB-
MCS vector.
246

Generation of stable cell lines

MA-NSP5-EGFP were generated as previously described [7].

MA-NSP2-mCherry and MA-NSP5 cell line were generated using lentiviral vector

system [31]. Briefly, HEK293T cells were maintained in DMEM (Life

Technologies) supplemented with 10% FBS (Life Technologies), and 50 µg/ml

gentamycin (Biochrom AG). Approximately 7×10^6 HEK293T cells were seeded

in 10 cm^2 tissue culture dishes 24 hours before transfection. For each well, 2.4

µg of pMD2-VSV-G, 4 µg of pMDLg pRRE, 1.8 µg of pRSV-Rev and 1.5 µg of

plasmid containing pAIP-NSP2-mCherry or pAIP-NSP5 and the human

immunodeficiency virus long terminal repeats were co-transfected with

Lipofectamine 3000 (Sigma-Aldrich) according to the manufacturer's instructions.

After 48 h, the virus was collected by filtration with a 0.45-µm polyvinylidene

fluoride filter and was immediately used or stored at −80 °C. For lentiviral

transduction, MA104 cells were transduced in six-well plates with 1 ml of lentiviral

supernatant for 2 days.

MA-Δ3 and MA-ΔT were generated using the PiggyBac Technology [32]. Briefly,

10^5 MA104 cells were transfected with the pCMV-HyPBase [32] and the

transposon plasmid pPB-NSP5/Δ3 and pPB-NSP5/ΔT using a ratio of 1:2.5 with

Lipofectamine 3000 (Sigma-Aldrich) according to the manufacturer's instructions.

The cells were maintained in DMEM supplemented with 10% FBS for 3 days and

then the cells were incubated with DMEM supplemented with 10% FBS and 5

µg/ml puromycin (Sigma-Aldrich) for 4 days to allow the selection of cells

expressing the gene of interest.

Rescue of recombinant RVs (rRVs) from cloned cDNAs.

To rescue recombinant RV strain SA11 (rRV-WT), monolayers of BHK-T7 cells

(4×10^5) cultured in 12-well plates were co-transfected using 2.5 µL of TransIT-

LT1 transfection reagent (Mirus) per microgram of DNA plasmid. Each mixture

comprised 0.8 µg of SA11 rescue plasmids: pT$_7$-VP1, pT$_7$-VP2, pT$_7$-VP3, pT$_7$-

VP4, pT$_7$-VP6, pT$_7$-VP7, pT$_7$-NSP1, pT$_7$-NSP3, pT$_7$-NSP4, and 2.4 µg of pT$_7$-

NSP2 and pT$_7$-NSP5 (Komoto et. al, 2018). Furthermore 0.8 µg of pcDNA3-

NSP2 and 0.8 µg of pcDNA3-NSP5, encoding NSP2 and NSP5 proteins, were

also co-transfected to increase rescue efficiency.

To rescue recombinant rRVs encoding NSP5 mutants, pT_7 plasmids encoding

NSP5/S67A, NSP5/Y18Stop, NSP5/Δ180-198 segments were used instead of

pT_7-NSP5. At 24 h post-transfection, MA-NSP5 cells (5×10^4 cells) were added

to transfected cells to provide a functional NSP5 for the virus rescue. The cells

were co-cultured for 3 days in FBS-free medium supplemented with trypsin (0.5

µg/mL) (Sigma Aldrich). After incubation, transfected cells were lysed by freeze-

thawing and 200 µl of the lysate was transferred to fresh MA-NSP5 cells. After

adsorption at 37°C for 1 hour, cells were washed three times with PBS and

further cultured at 37°C for 4 days in FBS-free DMEM supplemented with 0.5

µg/mL trypsin (Sigma Aldrich, 9002-07-7) until a clear cytopathic effect was

visible. The recombinant viruses were then checked by RT-PCR.

Immunofluorescence microscopy

Immunofluorescence experiments were performed using µ-Slide 8 Well Chamber

Slide-well (iBidi GmbH, Munich, Germany) and the following antibody dilutions:

anti-NSP5 guinea pig serum 1:1,000; anti-NSP2 guinea pig serum 1:200; anti-

VP2 guinea pig serum 1:500; anti-VP6 mouse monoclonal antibody 1:1,000;

Alexa Fluor 488-conjugated anti-mouse, 1:500 (Life Technologies), and TRITC-

conjugated anti-guinea pig, 1:500 (Life Technologies).

5-Ethynyl-Uridine (EU) labeling

Newly synthesized RNAs were labeled by including 2 mM 5-ethynyl uridine (EU)

into the cell culture medium, and modified incorporated nucleotides were reacted

with an azide-conjugated fluorophore Alexa-488 following the manufacturer's

protocol for Click-iT RNA Alexa Fluor 488 imaging kit (Thermo Fisher Scientific).

Cell nuclei were stained with ProLong™ Diamond Antifade Mountant with 4',6-

diamidino-2-phenylindole (DAPI, Thermo Scientific). Samples were imaged using

a confocal setup (Zeiss Airyscan equipped with a 63x, NA=1.3 objective), and the

images were processed using ZEN lite software.

RNA Fluorescence in situ Hybridization (FISH)

Rotavirus-infected MA104 cells were fixed with 4% (v/v) paraformaldehyde in nuclease-free Dulbecco's phosphate saline buffer (DPBS) for 10 min at room temperature. Samples were washed twice with DPBS, and then permeabilized with 70% (v/v) ethanol in RNAse-free water at +4°C for at least 1 hour prior to hybridization. Permeabilized samples were re-hydrated for 5 min in a pre-hybridization buffer (300 mM NaCl, 30 mM trisodium citrate, pH 7.0 in nuclease-free water, 10 % v/v formamide, supplemented with 2 mM vanadyl ribonucleoside complex). Re-hydrated samples were hybridized with 62.5 nM of an equimolar mixture of Cy3-labelled DNA probes designed to target the coding region of the gene segment 6 of simian rotavirus A/SA11 (Genbank Acc. AY187029.1) using Stellaris Probe Designer v2 software (LCG Biosearch Technologies), in a total volume of 200 μl of the hybridization buffer (Stellaris RNA FISH hybridization buffer, SMF-HB1-10, Biosearch Technologies, supplemented with 10% v/v of deionized formamide). After 4-8 hours of incubation at 37°C in a humidified chamber, samples were briefly rinsed with the wash buffer (300 mM NaCl, 30 mM trisodium citrate, pH 7.0, 10 % v/v formamide in nuclease-free water), after which a fresh aliquot of 300 μl of the wash buffer was applied to each sample and incubated twice at 37°C for 30 min. After 2 washes, nuclei were briefly stained with 300 nM DAPI solution in 300 mM NaCl, 30 mM trisodium citrate, pH 7.0, and the samples were finally rinsed with and stored in the same buffer without DAPI prior to imaging.

Transmission electron microscopy. MA104 cells were seeded at 1×10^5 cells in 2 cm^2 wells onto sapphire discs and infected at MOI of 75 FFU/cell. At 10 hpi, cells were fixed with 2.5% glutaraldehyde in 100 mM Na/K phosphate buffer, pH 7.4 for 1 h at 4°C and kept in that buffer overnight at 4°C. Afterward, samples were postfixed with 1% osmium tetroxide in 100 mM Na/K phosphate buffer for 1h at 4°C and dehydrated in a graded ethanol series starting at 70%, followed by two changes in acetone, and embedded in Epon. Ultrathin sections (60 to 80 nm)

340 were cut and stained with uranyl acetate and lead citrate [33]. Samples were
341 analysed in a transmission electron microscope (CM12; Philips, Eindhoven, The
342 Netherlands) equipped with a charge-coupled-device (CCD) camera (Ultrascan
343 1000; Gatan, Pleasanton, CA, USA) at an acceleration of 100 kV.
344

λ-Protein phosphatase and Calf Intestinal Alkaline Phosphatase Assays
Cellular extracts were incubated with 2,000 units of λ-protein phosphatase (λ-
Ppase) (New England Biolabs) in 50 mM Tris-HCl (pH 7.5), 0.1 mM EDTA, 5 mM
DTT, 0.01% Brij 35, and 2 mM $MnCl_2$. The mixture was incubated at 30°C for 2 h.
Samples were loaded in SDS-PAGE and analysed by Western blotting.
For the Calf Intestinal Alkaline Ppase (CIP) assay, cellular extracts were
incubated with 1,000 units of CIP (New England Biolabs) in CutSmart™ reaction
buffer (New England Biolabs) and incubated at 30°C for 2h. Samples were
subjected to SDS-PAGE and analysed by Western blotting.
354

Electrophoresis of Viral dsRNA Genomes.
Viral dsRNA was extracted with RnaZol® (Sigma-Aldrich) according the
manufacturer's protocol. Viral dsRNA genome segments from rRVs-infected cells
was mixed with Gel Loading Dye, Purple (6X) (New England Biolabs). The
dsRNAs segments were resolved on 10% (wt/vol) poly-acrylamide gels (PAGE)
for 2 hours at 180 Volts and visualized by ethidium bromide staining (1μg/ml).
361

Protein Analysis
Proteins derived from rRVs-infected cellular extract were separated on an SDS-
PAGE for 2 hours at 35 mA and transferred to polyvinylidene difluoride
membranes (Millipore; IPVH00010) for 1 hour and 30 minutes at 300 mA [24].
For protein analysis membranes were incubated with the following primary
antibodies: anti-NSP5 (1:5,000) [16], anti-VP2 (1:5,000) [10], anti-NSP2 guinea
pig sera (1:2,000). The membranes were then incubated with the corresponding
horseradish peroxidase (HRP)-conjugated goat anti-guinea pig (1:10,000;

Jackson ImmunoResearch). Mouse HRP-conjugated anti-actin mAb (1:35,000)

(clone AC-15, Sigma-Aldrich) was used as loading control.

Signals were detected using the enhanced chemiluminescence system (Pierce

ECL Western blotting substrate; Thermo Scientific).

Statistics used

Statistical analysis and plotting were performed using GraphPad Prism 6

software (GraphPad Prism 6.0, GraphPad Software Inc., La Jolla, CA, USA).

Error bars represent standard deviation. Data were considered to be statistically

significant when $p < 0.05$ by Student's t test.

RESULTS

Generation of gs11 mutated recombinant RVs

Recombinant rotaviruses (simian rotavirus A strain SA11) carrying different

mutations in the NSP5 coding region of gs11 were obtained using a recently

developed reverse genetics protocol [29]. Two essential modifications were

introduced to *trans*-complement the potential loss of NSP5 function in the rRV

mutants: i) an additional T7-driven plasmid encoding the ORF of wt NSP5 (i.e.,

without gs11 5' and 3' UTRs) was included in the transfection step of BHK-T7

cells; and ii) each rescued rRV was amplified in a stable transfectant cell line MA-

NSP5, supplying the wt NSP5 in trans (Fig. S1). Crucially, we established MA-

NSP5 cell line to support the replication of NSP5-deficient recombinant viruses to

enable their further in-depth characterisation. These mutants, as well as

additional stable cell lines generated for this study as described in Materials and

Methods, are summarised in Fig. S2 and S3A-B.

rRV-NSP5/KO

NSP5 expression and localisation to viroplasms in virus-infected cells have been

considered essential for virus replication [12,13,25,34]. Previous studies, using

siRNA targeting gs11 mRNA have shown strong impairment of RV replication

[12,13]. We, therefore, established MA104 cell line stably expressing NSP5 to

supply NSP5 in trans in order to investigate the effect of point mutations and
deletions within the NSP5 gene.   In addition, we also made two cell lines
expressing NSP2-mCherry and NSP5-EGFP fusions, which are rapidly and
efficiently recruited into viroplasms upon virus infection (Fig. S2) [7,22].

Here we provide direct demonstration of the essential role of NSP5 in RV
replication, using an NSP5 knock out rRV (termed rRV-NSP5/KO) generated by
reverse genetics. To rescue the NSP5/KO strain a premature stop codon at the
position 18 (Tyr18Stop) was introduced within the NSP5-coding mRNA (Fig.
S3A-B). Analysis of MA104-virus-infected cell extracts confirmed the presence of
NSP2 and VP2, but not of NSP5 (Fig. 1A). Moreover, we did not detect
viroplasms containing NSP2, VP2 or VP6 in both MA104-infected cells and in
stable transfectant cell lines expressing the fluorescent fusion proteins NSP2-
mCherry or NSP5-EGFP (Fig. S2 and Fig. 1B). Furthermore, both genomic
dsRNA synthesis and infectious progeny virus production were completely
abrogated in MA104 cells, but not in the trans-complementing MA-NSP5 cell line
(Fig. 1C). Together, these data confirm that NSP5 is essential for RV replication.

rRVs with impaired NSP5 phosphorylation

To address the role of NSP5 hyperphosphorylation, we then generated a number
of rRVs harbouring NSP5 mutations previously known to impact NSP5
phosphorylation. We first generated an rRV carrying an S67A mutation (rRV-
NSP5/S67A) (Fig. S2 and Fig. 2A-B). This rRV strain appeared to be genetically
stable after a number of passages in the MA-NSP5 cell line (Fig. 2A-B). Its
replication in wild type MA104 cells was strongly impaired, resulting in
approximately a 100-fold reduction of the infectious progeny virus titre at different
time points post-infection (Fig. 2C). Despite the overall reduction of replication
fitness, the rRV-NSP5/S67A mutant virus was stable after at least 10 passages
in the wild-type MA104 cells. Consistent with our previous results, the NSP5-
S67A mutant was not hyper-phosphorylated in all cell lines tested, including
MA104, U2OS and Caco-2 cells (Fig. 2D), further confirming the role of Ser67 in
the initiation of NSP5 phosphorylation cascade [24]. While the wt rRV yielded

multiple hyper-phosphorylated NSP5 isoforms, the NSP5/S67A mutant mostly
produced a single non-phosphorylated NSP5 form with an apparent mass of 26
kDa that could be detected in the virus-infected cell extracts at 5 or 10 hpi (Fig.
2E). Enzymatic de-phosphorylation with λ-Ppase and alkaline Calf Intestinal
Ppase (CIP) further confirmed the observed lack of NSP5 phosphorylation (Fig.
2F). Because of the differences in the molecular weight markers used, the NSP5
band with the fastest PAGE mobility has been traditionally described as 26 kDa,
and the most abundant one as 28 kDa. Since this nomenclature has been used
in many publications, we have preferred to maintain it, despite current PAGE
migrations do not correspond to the markers presently used.

We then generated two additional rRVs harbouring truncated versions of NSP5,
one lacking the 18 AA long C-terminal tail (rRV-NSP5/ΔT), and the second one
with a 5 AA deletion (176-YKKKY-180) just upstream of the tail region (rRV-
NSP5/Δ176-180) (Fig. S2 and S3A-B). Despite the presence of Ser67, and the
lack of Thr and Ser residues within the deletions, both mutants did not show a
classical hyperphosphorylation pattern (Fig. 3A) and failed to replicate, confirmed
by the absence of *de novo* synthesis of genomic dsRNA (Fig. 3B).

We also investigated viroplasm formation in cells infected with these rRV
mutants. At early infection (5 hpi), the rRV-NSP5/S67A mutant produced
structures resembling viroplasms that appeared smaller and more
heterogeneous in shape compared to the regular spherical shaped ones
produced during the wt rRV infection (Fig 5A upper panel). Remarkably, during
late infection (10-12 hpi), the rRV-NSP5/S67A mutant produced multiple NSP5-
containing aberrant structures, as well as fibre-like structures that became more
apparent during late infection (12 hpi) (Fig. 4A lower panel).

Interestingly, the aberrant and fibre-like structures produced by rRV-NSP5/S67A
were morphologically similar to those observed during wt RV infection of MA104
cells silenced for cellular kinase CK1α, previously shown to be required for NSP5
phosphorylation [25]. NSP2-mCherry and NSP5-EGFP fusion proteins were also
recruited to both types of these structures (Fig. 4C-D), suggesting that the
observed lack of phosphorylation does affect NSP2/NSP5 interactions.

Furthermore, the non-phosphorylated NSP5 form co-localised with viroplasmic

proteins NSP2, VP6 and VP2 (Fig. 4B) that also localised to aberrant and fibre-

like structures, suggesting that they could also represent sites of virus replication.

In contrast, no viroplasms containing NSP5, NSP2 or VP2 were observed when

MA104 cells were infected with the rRV-NSP5/ΔT and rRV-NSP5/Δ176-180

deletion mutants. Interestingly, these mutants yielded only few fibre-like

structures containing VP6 protein (Fig. 4D-E). Similar VP6-fibres are normally

formed when VP6 is over-expressed in cells in the absence of other viral proteins

[35,36].

Taken together, these results confirm the essential role of NSP5 hyper-

phosphorylation for controlling the assembly of regular-shaped viroplasms,

highlighting the key role of the C-terminal tail in the formation of RV viral

factories.

RNA accumulation in aberrant viroplasms

Having examined the viral protein composition of the aberrant viroplasms formed

during infection with rRVs exhibiting impaired NSP5 phosphorylation, we then

assessed their RNA content. Viral RNA transcripts were labelled by incorporation

of 5-ethynyl uridine (5-EU) in actinomycin D-treated RV-infected cells, and total

viral ssRNA was visualized by reacting with Alexa-488-azide, as described in

Materials and Methods. As expected, most viral transcripts localised in

viroplasms of wt rRV-infected cells (Fig. 5A), consistent with the roles of

viroplasms in supporting the viral replication and assembly. In contrast, no viral

RNA transcripts could be detected in both wt MA104 or MA-NSP2-mCherry cells

infected with the rRV-NSP5/S67A mutant (Fig. 5A). RNA accumulation in

viroplasms was restored to almost identical levels as the wt rRVs when the rRV-

NSP5/S67A mutant was used to infect the MA-NSP5 cell line that supplies NSP5

in *trans* (Fig. 5A).

Although we did not detect RNA in rRV NSP5/S67A-infected MA104 cells using

5-EU staining, this mutant could still replicate (Fig. 2C). We then explored

whether rRV-NSP5/S67A transcripts could accumulate in viroplasms, albeit with

much lower efficiency, i.e., beyond the sensitivity limit of 5-EU staining. We used

single molecule RNA fluorescence in situ hybridisation (smFISH) to identify the

sites of RV transcription. At 10 hpi, abundant gs6 transcripts could be detected in

MA-NSP5-EGFP cells infected with the wild-type virus (Fig. 5B). Conversely, the

rRV-NSP5/S67A-infected cells had sparse EGFP-tagged viroplasms, also

containing gs6 RNA (Fig. 5B). Interestingly, the less frequently occurring rod-like

aberrant viroplasms also sustained gs6 RNA accumulation, further suggesting

that these structures could represent replication-functional organelles.

In contrast, smFISH performed on cells infected with the rRV-NSP5/ΔT showed

diffuse distribution of gs6 RNA that did not localise to any structures resembling

viroplasms (Fig. 5B), also failing to support the virus genome replication (Fig.

3B).

We then examined the ultrastructures of viroplasms with altered morphologies in

the RV mutant rRV-NSP5/S67A using electron microscopy. Upon infection with

rRV-wt (Fig. 6A, left panel), multiple membrane-less electron dense inclusions

encircled by the well-defined endoplasmatic reticulum (ER) filled with triple-

layered particles were present in cells. At late infection points (10 hpi), filled with

triple-layered particles (TLPs) ER appeared to adopt a more tubular morphology,

suggesting a successive step in the virus egress. In contrast, the rRV-

NSP5/S67A-infected cells contained only few immature viroplasms that lacked

the ER network filled with TLPs (Fig. 6A, right panel). Only few immature

particles containing transient lipid membranes could be identified in cells infected

with rRV-NSP5/S67A mutant (Fig. 6A, left panel). Furthermore, the observed

immature viroplasms also appeared to be less-electron dense, consistent with

their decreased capacity to accumulate viral RNAs.

Together, these data strongly support the role of NSP5 phosphorylation in

maintaining the viral RNA production and genome replication in viroplasms.

The mechanism of NSP5 hyper-phosphorylation

We have previously proposed a model of the hierarchical NSP5

hyperphosphorylation associated with the assembly of viroplasms that involves a

three-step mechanism: i) initial interaction of non-phosphorylated NSP5 with
NSP2 (or VP2); ii) phosphorylation of Ser67 by CK1α. This step does not take
place when NSP5 is expressed alone; iii) hyper-phosphorylation of NSP5
triggered by Ser67 phosphorylation that requires the 18 AA long C-terminal tail
[24].

Here, we have investigated the phosphorylation mechanism of NSP5 during RV
infection using a number of NSP5 phosphorylation-negative rRV strains and
MA104-derived stable transfectant cell lines (Fig. S2). We demonstrated that
despite the presence of Ser67, deletion mutant NSP5/ΔT was not phosphorylated
and failed to form viroplasms. We have previously shown that co-expression of
NSP5/ΔT was also unable to trigger the phosphorylation cascade of NSP5/S67A,
while other NSP5 mutants referred hereafter as activators of phosphorylation,
e.g., NSP5/Δ3 did [24]. Interestingly, upon co-infection with two rRVs NSP5/S67A
and NSP5/ΔT, both mutated NSP5 variants were not phosphorylated (Fig. 7A)
[8,19,24,26]. This result was supported further by infecting the MA104
transfectant cell line stably expressing NSP5-ΔT (MA-ΔT) with the rRV-
NSP5/S67A strain (Fig. S2 and Fig. 7B, lanes 3-4). The NSP5-Δ176-180 mutant
was also unable to induce phosphorylation of NSP5/S67A following the co-
infection with the two rRVs, despite both containing Ser67 and the C-terminal tail.
This result suggests that the 'activator' NSP5 needs to be also hyper-
phosphorylated (Fig. 7C).

In MA-ΔT cells infected with rRV-NSP5/S67A mutant strain aberrant viroplasms
were produced (Fig. 7E), similar to those seen in wt MA104 cells, which did not
support virus replication compared to the wt virus (Fig. 7F).

In contrast to NSP5/ΔT, the NSP5 deletion mutant lacking amino acids 80-130,
(NSP5/Δ3) becomes hyper-phosphorylated when expressed alone and can
function as an activator of the NSP5/S67A phosphorylation [19,24]. We therefore
asked whether an MA104 stable cell line expressing the deletion mutant
NSP5/Δ3 (MA-Δ3) and infected with the rRV-NSP5/S67A strain was able to
trigger hyperphosphorylation of NSP5/S67A and as a consequence, sustain

replication of the mutant rRV strain. As shown in Fig. 7B (lanes 5-6) NSP5/S67A
was hyper-phosphorylated in presence of NSP5/Δ3 mutant and confirmed by the
λPpase treatment (Fig. 7D), although it did not completely rescue the
phosphorylation pattern of NSP5 normally observed in rRV-wt infection.
Interestingly, regular round-shaped viroplasms, containing NSP5, NSP2 and VP2
with peripheral localisation of VP6, were recovered in these cells infected with
rRV-NSP5/S67A, and yet viral replication was nevertheless impaired (Fig. 7E-F).
Consistently, the electron microscopy images showed structures of aberrant
viroplasms similar to those obtained in MA104 wt cells (Fig. 6B, right panel).
The NSP5 phosphorylation negative mutants of the two other rRVs (NSP5/ΔT
and NSP5/Δ176-180), did not undergo hyper-phosphorylation in MA-ΔT or MA-Δ3
cell lines (Figs. 8A-B). In both cases, viroplasms formation and virus replication
were not rescued (Figs. 8C-D and 8E-F). Interestingly, apparently normal round-
shaped structures were observed in MA-Δ3 cells, which, however, recruited
significantly less VP6 (Fig. 8C-D). In addition, the previously observed VP6 spiky
structures were not detected in these cells (Figs. 8C-D). Similar results were
obtained with the rRV-NSP5/KO virus strain, which in MA-Δ3 cells did not
replicate and also showed spherical structures that contained NSP5 and NSP2,
but not VP6 (Fig. 9A-C).
Taken together, our data support a model of NSP5 hyper-phosphorylation, which
absolutely requires the presence of the C-terminal tail, and the AA residues 176-
180. Furthermore, NSP5 hyperphosphorylation requires Ser67 phosphorylation
to initiate the phosphorylation cascade, thus playing a key role in the assembly of
replication-competent viral factories.
579
DISCUSSION
Rotaviruses replicate within cytoplasmic viral factories, or viroplasms. Most RV
assembly intermediates, i.e., single layered particles (cores) and double-layered
particles (DLPs) are primarily concentrated in viroplasms. Following budding of
DLPs into the lumen of the endoplasmic reticulum, the immature particles acquire
a transient envelope, as well as the outer capsid proteins VP4 and VP7, resulting

in a mature triple-layered virion. Moreover, downregulation of expression of the
most abundant viroplasm-forming proteins NSP5 and NSP2 severely impacts on
the formation of viroplasms and production of virus progeny [13,15,25,34]. In light
of these observations, viroplasms have long been recognised as essential
compartments supporting the RV replication.

Of the two non-structural proteins involved in viroplasm assembly, NSP5 appears
to play a crucial role by potentially providing a scaffold that allows for recruitment
of additional viral proteins. Only when NSP5 is co-expressed with NSP2 and/or
VP2, these proteins assemble into the viroplasm-like structures (VLS), which are
also capable to recruit additional structural proteins including VP1, VP2 and VP6
[8,10]. Given these observations, we hypothesised that complete removal of
NSP5 would be lethal for RV replication. Using a modified reverse genetics
system for rotaviruses [29], here we provide the first direct evidence of the
essential role of NSP5 in viroplasm formation and viral replication. In order to
characterise replication-deficient NSP5-negative mutants, we have established a
trans-complementing system that provides NSP5 to the virus both transiently in
BHK-T7 cells, and stably in NSP5-producing MA104 cell line (MA-NSP5), thus
enabling facile isolation of rRVs lacking functional NSP5. Using this approach,
we have demonstrated that NSP5-deficient rRV was unable to form viroplasms
and replicate in the wt MA104 cells, while the viroplasm formation and viral
replication were efficiently rescued in the trans-complementing MA-NSP5 cell
line. Interestingly, the rRVs generated using this method also failed to
incorporate dsRNA originating from the NSP5-encoding mRNA lacking the 5' and
3' untranslated regions (UTRs), further suggesting the essential roles of UTRs for
genome packaging in RVs [37,38].

NSP5 hyper-phosphorylation has been previously implicated in the regulation of
NSP5 assembly into viroplasms. This phosphorylation, however, requires the
interaction of NSP5 with, either NSP2 or VP2, as NSP5 is not phosphorylated
when expressed alone [8,22,23]. Previous studies suggest that activation of
NSP5 hyper-phosphorylation may require a conformational change that leads to
its efficient hyper-phosphorylation via a positive feedback loop mechanism

[8,23,24]. Two regions comprising the N-terminal amino acids 1-33 (region 1) and

the central region amino acids 81-130 (region 3) have been reported to prevent

NSP5 phosphorylation in the absence of other viral proteins, while the 18 amino

acids long C-terminal tail was found to be essential for its phosphorylation

[8,23,24].

Here, we have shown that all three rRV mutants S67A, ΔT and Δ176-180

expressing the phosphorylation-negative NSP5 variants were unable to form

round-shaped viroplasms upon infection of MA104 cells. Interestingly, further

analysis of these mutants reveals some key differences between each NSP5

variant. While rRV-NSP5/S67A strain formed aberrant structures resembling

viroplasms that poorly support RV replication, this variant was still capable of

producing the infectious progeny, in contrast to the two other rRV mutant strains.

Interestingly, the phenotype observed with the NSP5 mutant S67A was

essentially the same as the one previously reported with the wt virus infecting

MA104 cells silenced for expression of CK1α, which is involved in

phosphorylating Ser67 and initiating the hyper-phosphorylation cascade [24,25].

It has recently been shown that CK1α is also involved in phosphorylating NSP2,

controlling the formation of rotavirus viral factories [39]. Our data obtained with

the rRV-NSP5/S67A mutant strongly suggest that the lack of NSP5 hyper-

phosphorylation determines both the morphogenesis of viroplasms and their

capacity to support RV genome replication. We cannot rule out the role of NSP2

phosphorylation in assembly of viroplasms since NSP2 is also likely to be

phosphorylated by CK1α upon infection of the rRV-NSP5/S67A strain. Despite

the formation of abnormal viroplasms and fibre-like structures, the amount of

RNA produced within those structures in rRV-NSP5/S67A-infected MA104 cells

was practically below the detection limit of the 5-EU labelling, while the RNA

replication was fully rescued in the trans-complementing MA-NSP5 cell line (Fig.

5). Interestingly, we were able to detect viral RNA in some of these aberrant

structures using RNA FISH with single molecule sensitivity. This result is

consistent with the finding that the rRV-NSP5/S67A strain did replicate, albeit at

much lower levels than the rRV-wt. Thus, these structures are likely to sustain

virus replication with decreased efficiency, which was further confirmed by the

electron microscopy analysis of MA104 cells infected with the rRV-NSP5/S67A

strain.

The essential role of NSP5 hyper-phosphorylation was further supported by the

results obtained with the two phosphorylation-negative mutant strains NSP5/ΔT

and NSP5/Δ176-180 that possess Ser67 and yet failed to form viroplasms.

Surprisingly, the Δ176-180 mutant, containing the C-terminal tail has also failed

to form viroplasms, despite the absence of Ser or Thr within the chosen 176-180

region. Both phosphorylation-negative mutants tested did not replicate in MA104

cells and we could not detect any structures containing viral RNA.

Using the NSP5 mutant strains described above and the established stable

transfectant MA104 cells we were able to investigate the molecular mechanism

that leads to NSP5 hyperphosphorylation. We showed that the NSP5/S67A

mutant from the rRV was indeed hyperphosphorylated, albeit not completely,

when infecting MA-Δ3 cells, restoring round-shaped viroplasms with a complete

absence of the fibre-like structures observed in MA104 cells. This finding strongly

suggests that impairment of NSP5 phosphorylation is the direct cause of the

formation of both aberrant and fibre-like structures. Despite the fact that

viroplasms appeared morphologically normal in MA-Δ3 cells, recruitment of VP6

at the periphery of viroplasms was only observed with the hyperphosphorylated

NSP5/S67A mutant, in contrast to the other NSP5 phosphorylation negative

mutants. One possibility is that recruitment of VP6 to round-shaped viroplasms

requires a full-length hyperphosphorylated NSP5, as well as phosphorylation of

multiple serine residues likely by CK2 [17,26]. Moreover, the round-shaped

structures found in MA-Δ3 cells infected with rRV-NSP5/KO failed to recruit VP6.

The observed failure to rescue the replication of the rRV-NSP5/S67A strain to the

wt levels in MA-Δ3 cells could be the consequence of the incomplete recovery of

the complex pattern of phosphorylated isoforms of wt NSP5. This suggests that

some intermediate isoforms might be important for the formation of fully

functional replication-competent viroplasms.

We propose a model of the complex hierarchical mechanism of NSP5 hyper-
phosphorylation during RV infection (Fig. 10). It involves (a) interaction of NSP5
with either NSP2 or VP2, required to (b) make Ser67 available for CK1α
phosphorylation. This initial step is then sequentially completed (c) by CK2-
mediated phosphorylation of other serines to generate the NSP5 'activator',
during a step dependent (d) on the interaction of the Ser67-phosphorylated
molecules with the non-phosphorylated partners in the NSP5 oligomeric
complexes. Alternatively (c'), oligomers could be formed before the activation
step. NSP5 interactions mediated by the carboxy-terminal tails T result in (e)
substrate activation and a fully hyperphosphorylated NSP5. This process leads to
(f) the assembly of viroplasms scaffolds containing NSP2, and (g) recruitment of
VP6, as well as the other viroplasmic proteins, VP1, VP2, VP3 to assemble
replication-competent viroplasms.

The hierarchical phosphorylation of proteins appears to be a common
mechanism regulating many cellular processes. In mammalian cells, hierarchical
phosphorylation has been described for β-catenin, in which Ser 45
phosphorylated by CK1α primes it for hyperphosphorylation by glycogen
synthase kinase-3 (GSK-3) [40,41], which triggers its ubiquitination and
proteasomal degradation [42]. A similar phosphorylation mechanism has recently
been described for a non-structural protein NS5A of the Hepatitis C Virus (HCV),
with the hyper-phosphorylation cascade primed by the initial phosphorylation of
Serine 225 by CK1α, and the subsequent phosphorylation of neighbouring
residues involving other kinases [43,44]. Moreover, the NS5A phosphorylation
was shown to play a key role in controlling the establishment of replication
complexes during HCV infection [45]. Similarly, a number of other non-structural
viral proteins have been shown to undergo multiple phosphorylation events
during virus infection, suggesting that this complex post-translational
modifications play a pivotal role in orchestrating the assembly of replication-
competent viral factories [46–48].
707
708

ACKNOWLEDGEMENTS

We thank John T. Patton, Asha Ann Philipp and Jin Dai (Indiana University,

Bloomington, USA), Yuta Kanai and Takeshi Kobayashi (Fujita Health University,

Aichi) and Albie Van Dijik (North-West University, South Africa) for helpful

discussions on the establishment of the reverse genetics system. We are grateful

to Naoto Ito for providing BHK-T7 cells.

  FIGURE LEGENDS

  Fig. 1. Characterisation of rRV-NSP5/KO. A) Western blot of MA104 cells

  extracts infected with rRV-wt and rRV-NSP5/KO strains (MOI 1 FFU/cell). B)

  Confocal immunofluorescence microscopy of MA104, MA-NSP2-mCherry and

  MA-NSP5-EGFP cells infected with rRV-wt or rRV-NSP5/KO (MOI 1 FFU/cell),

  using antibodies for NSP5, NSP2, VP2 and VP6, as indicated. Scale bar, 15 μm.

  C) Electrophoretic migration pattern of dsRNAs extracted from rRV-NSP5/KO

  strain grown in MA-NSP5 or MA104 cells. Genome segments 1-11 are indicated

  to the left.

  Fig 2. Characterisation of rRV-NSP5/S67A. A) Schematic representation of

  rRV-NSP5/S67A gs11 and sequence of NSP5 wt and S67A mutant (highlighted).

  B) Electrophoretic pattern of dsRNA genome segments of rRV-wt and rRV-

  NSP5/S67A strains grown in MA-NSP5 cells. C) Replication kinetics of rRV-wt

  and rRV-NSP5/S67A in MA104 cells. Data are expressed as the means +/-

  standard deviations (n=3); ***, $p < 0.001$ (Student's t test). D) Western blot of

  extracts of U2OS, Caco-2 and MA104 cells infected with rRV-wt or rRV-

  NSP5/S67A strains. E) Western blot of NSP5 phosphorylation pattern in MA104

  cells infected with rRV-wt or rRV-NSP5/S67A (MOI 1 FFU/cell) at 5 and 10 hpi. F)

  Western blot of λ-Ppase and alkaline Ppase treatment of lysates of MA104 cells

  infected with rRV-wt or rRV-NSP5/S67A. The NSP5 proteins are shown.

  Fig 3. Characterisation of rRV-NSP5/ΔT and rRV-NSP5/Δ176-180. A) Western

  blot of NSP5 and NSP5 mutants from MA104 cells infected with rRV-wt, rRV-

  NSP5/ΔT or rRV- NSP5/Δ176-180 at 5 hpi. B) Electrophoretic dsRNA migration

  pattern of rRV-NSP5/ΔT and rRV-NSP5/Δ176-180 grown in MA-NSP5 and

  MA104 cells.

  Fig 4. Viroplasms morphology in MA104 cells infected with rRV-NSP5 wt

  and mutants. Representative confocal immunofluorescence micrographs of

MA104 cells infected with rRV-wt and rRV-NSP5/S67A (MOI 15 FFU/cell) at 5 hpi

(upper panels) or 10 hpi (lower panels). Cells were stained with anti-NSP5 and

DAPI. B) Confocal immunofluorescence micrographs of rRV-wt and rRV-

NSP5/S67A infected MA104 cells (MOI 15 FFU/cell) and stained at 10 hpi with

the indicated antibodies. C) Confocal immunofluorescence micrographs of MA-

NSP5-EGFP and MA-NSP2-mCherry cells infected with rRV-NSP5/S67A (MOI

15 FFU/cell). D), E), Confocal immunofluorescence of MA104 cells infected with

rRV-NSP5/Δ176-180 (D) or rRV-NSP5/ΔT (E) and stained with the indicated

antibodies and DAPI. Scale bar, 15 μm.

Fig 5. Viral RNA detection in rRV NSP5/S67A infected cells. A)

Representative confocal immunofluorescence micrographs of MA104 (upper

panel), MA-NSP5 (middle panel) and MA-NSP2-mCherry cells (lower panel)

infected with rRV-wt or rRV-NSP5/S67A strains and stained with anti-NSP5 (red,

in MA104 and MA-NSP5 cells) and EU (green). B) Single molecule RNA

Fluorescence In Situ Hybridisation (smFISH) on MA-NSP5-EGFP cells infected

with rRV-wt, rRV-NSP5/S67A or rRV-NSP5/ΔT strains. Viroplasms detected with

NSP5-EGFP (green) and viral RNA (probe specific for gs6 was Cy3-conjugated,

red). Co-localization of viroplasms and RNA is indicated by white arrows. Scale

bar, 15 μm.

Fig 6. Electron microscopy of cells infected with rRV-NSP5/S67A. High–

definition electron micrographs of MA104 (A) and MA-Δ3 (B) cells infected with

rRV-wt (left panel) and rRV-NSP5/S67A (right panel) (MOI, 75 FFU/cell). At 10

hpi, cells were fixed with glutaraldehyde and processed for transmission electron

microscopy. V, viroplasm; m, mitochondria; mt, microtubule-bundles; Nu, nucleus.

The white open box indicates the immediate right magnified image. White arrows

indicate the endoplasmic reticulum surrounding viroplasms; black arrowheads

indicate viral particles with an envelope. Scale bar, 500 nm.

Fig 7. Phosphorylation of NSP5/S67A. A) Western blot of extracts of MA104
cells infected with rRV-NSP5/S67A or co-infected with rRV-NSP5/S67A and rRV-
NSP5/ΔT. Blue arrowhead indicates NSP5/S67A. The faster migrating band
corresponds to NSP5-ΔT. B) Western blot of MA104, MA-ΔT and MA-Δ3 cells
infected with rRV-NSP5/S67A. Blue arrowhead indicates NSP5/S67A. C)
Western blot of extracts of MA104 cells co-infected with rRV-NSP5/S67A and
rRV-NSP5/Δ176-180. Blue arrowhead indicates NSP5/S67A. D) λ-Ppase
treatment of extracts of MA-Δ3 cells infected with the rRV-NSP5/S67A strain.
Filled and open arrowheads indicate de-phosphorylated NSP5-S67A and NSP5-
Δ3, respectively. E) Representative confocal immunofluorescence micrographs of
MA104, MA-ΔT and MA-Δ3 cells infected with rRV-NSP5/S67A. Cells were
stained with the indicated antibodies. Scale bar, 15 μm. F) Yield of infectious
virus of rRV-NSP5/S67A grown in MA-NSP5, MA104, MA-ΔT and MA-Δ3 cells at
24 hpi.*, $p < 0.05$; **, $p < 0.01$.

Fig 8. Phosphorylation of NSP5/-ΔT and NSP5/Δ176-180 and viroplasm
formation. Western blots (A, B) and confocal immunofluorescence micrographs
(C, D) of MA104, MA-ΔT and MA-Δ3 cells infected with rRV-NSP5/ΔT (A, C) or
with rRV-NSP5/Δ176-180 (B, D). Blue arrowhead indicates NSP5/ΔT and
NSP5/Δ176-180, respectively. Open arrowheads indicate NSP5-Δ3 and NSP5-
ΔT. Cells were stained with the indicated antibodies and DAPI. Scale bar, 15 μm.
E) Single step growth of rRV-NSP5/ΔT (left) and rRV-NSP5/Δ176-180 (right) in
MA-NSP5, MA104, MA-ΔT and MA-Δ3 cells, as indicated. The experiment was
terminated at 24 hpi. **, $p < 0.01$; #, Viral titer < 300 FFU/ml.

Fig 9. rRV NSP5/KO in infected cells. Western blot (A) and confocal
immunofluorescence (B) of MA104, MA-ΔT and MA-Δ3 cells infected with the
rRV-NSP5/KO strain. Cells were stained with the indicated antibodies and DAPI.
Scale bar, 15 μm. C) Single step growth of rRV-NSP5/KO in MA-NSP5, MA104,
MA-ΔT and MA-Δ3 cells, as indicated. The experiment was terminated at 24 hpi.
**, $p < 0.01$; #, Viral titer < 300 FFU/ml.

Fig 10. Model of NSP5 hyperphosphorylation and assembly of replication-competent viroplasms. (*a*) Interaction of non-phosphorylated NSP5 with either NSP2 or VP2 is required to (*b*) induce conformational changes that make S67 available for CK1α phosphorylation (P). This initial step of the cascade is then (*c*) sequentially completed by phosphorylation by CK2 of other residues including serines in domain 4, to generate the NSP5 activator or, (*c'*) a step of interaction with non-phosphorylated molecule precedes the involvement of CK2. (*d*) NSP5 interactions to form dimers/multimers are mediated by the carboxy-terminal tails (T). (*e*) The primed (S67-phosphorylated) activator molecules result in substrate activation and fully hyper-phosphorylated NSP5. (*f*) Assembly of viroplasms scaffolds containing NSP2 serve to (*g*) recruit the other viroplasmic components, VP1, VP2, VP3 and VP6 to assemble replication-competent viroplasms. P; phosphorylated amino acid. *This image was created using BioRender.com.

Fig S1. Schematic representation of the modified reverse genetics protocol used to rescue rRV with mutations in genome segment 11. gs; genome segment *Created using BioRender.com.

Fig S2. Schematic representation of the different NSP5 or NSP2 mutants or fusion proteins used to generate rRV or stable MA104 transfectant cell lines (or both) and used in this study. *Created using BioRender.com

Fig S3. A) Schematic representation of mutations in gs11 of the corresponding rRV strains. Sequences mutated or deleted in NSP5 are indicated. B), Profiles of viral dsRNAs of the different rRV strains generated. gs11, indicated in red with *.

REFERENCES

1.   Tate JE, Burton AH, Boschi-Pinto C, Parashar UD, World Health
Organization–Coordinated Global Rotavirus Surveillance Network. Global,
Regional, and National Estimates of Rotavirus Mortality in Children <5
Years of Age, 2000–2013. Clin Infect Dis. 2016;62: S96–S105.
doi:10.1093/cid/civ1013

2.   Parashar UD, Hummelman EG, Bresee JS, Miller MA, Glass RI. Global
illness and deaths caused by rotavirus disease in children. Emerg Infect
Dis.; 2003;9: 565–572.

3.   Patton JT, Silvestri LS, Tortorici MA, Vasquez-Del Carpio R, Taraporewala
ZF. Rotavirus genome replication and morphogenesis: role of the
viroplasm. Curr Top Microbiol Immunol. 2006;309: 169–87.

4.   Desselberger U. Rotaviruses. Virus Res; 2014;190: 75–96.
doi:10.1016/j.virusres.2014.06.016

5.   Estes MK, Greenberg HB. Rotaviruses. In: Health WK, Lippincott Williams
Wilkins, editors. Fields Virology. 6th ed. Philadelpia, PA; 2013. pp. 1347–
1401.

6.   Clapp LL, Patton JT. Rotavirus morphogenesis: domains in the major inner
capsid protein essential for binding to single-shelled particles and for
trimerization. Virology; 1991;180: 697–708.

7.   Eichwald C, Rodriguez JF, Burrone OR. Characterization of rotavirus
NSP2/NSP5 interactions and the dynamics of viroplasm formation. J Gen
Virol. 2004;85: 625–634. doi:10.1099/vir.0.19611-0

8.   Fabbretti E, Afrikanova I, Vascotto F, Burrone OR. Two non-structural
rotavirus proteins, NSP2 and NSP5, form viroplasm-like structures in vivo.
J Gen Virol. 1999;80: 333–339. doi:10.1099/0022-1317-80-2-333

9.   Poncet D, Lindenbaum P, L'Haridon R, Cohen J. In vivo and in vitro
phosphorylation of rotavirus NSP5 correlates with its localization in
viroplasms. J Virol. 1997;71: 34–41.

10.   Arnoldi F, Campagna M, Eichwald C, Desselberger U, Burrone OR.
Interaction of Rotavirus Polymerase VP1 with Nonstructural Protein NSP5
Is Stronger than That with NSP2. J Virol. 2007;81: 2128–2137.
doi:10.1128/JVI.01494-06

11.   Gallegos CO, Patton JT. Characterization of rotavirus replication
intermediates: a model for the assembly of single-shelled particles.
Virology. 1989;172: 616–27.

12.   Campagna M, Eichwald C, Vascotto F, Burrone OR. RNA interference of
rotavirus segment 11 mRNA reveals the essential role of NSP5 in the virus
replicative cycle. J Gen Virol; 2005;86: 1481–1487.
doi:10.1099/vir.0.80598-0

13.   Lopez T, Rojas M, Ayala-Breton C, Lopez S, Arias CF. Reduced
expression of the rotavirus NSP5 gene has a pleiotropic effect on virus
replication. J Gen Virol. 2005;6: 1609–1617.

14.   Silvestri LS, Taraporewala ZF, Patton JT. Rotavirus replication: plus-sense
templates for double-stranded RNA synthesis are made in viroplasms. J
Virol. American Society for Microbiology; 2004;78: 7763–74.
doi:10.1128/JVI.78.14.7763-7774.2004

15.   Vascotto F, Visintin M, Cattaneo A, Burrone OR. Design and selection of
an intrabody library produced de-novo for the non-structural protein NSP5
of rotavirus. J Immunol Methods; 2005;301: 31–40.

16.   Gonzalez SA, Burrone OR. Rotavirus NS26 is modified by addition of
single O-linked residues of N-acetylglucosamine. Virology. 1991;182: 8–16.

17.   Sotelo PH, Schumann M, Krause E, Chnaiderman J. Analysis of rotavirus

non-structural protein NSP5 by mass spectrometry reveals a complex

phosphorylation pattern. Virus Res. 2010;149: 104–108.

18.   Campagna M, Marcos-Villar L, Arnoldi F, de la Cruz-Herrera CF, Gallego

P, Gonzalez-Santamaria J, et al. Rotavirus Viroplasm Proteins Interact with

the Cellular SUMOylation System: Implications for Viroplasm-Like

Structure Formation. J Virol. 2013;87: 807–817. doi:10.1128/JVI.01578-12

19.   Afrikanova I, Miozzo MC, Giambiagi S, Burrone O. Phosphorylation

generates different forms of rotavirus NSP5. J Gen Virol. 1996;9: 2059–

2065.

20.   Blackhall J, Fuentes A, Hansen K, Magnusson G. Serine protein kinase

activity associated with rotavirus phosphoprotein NSP5. J Virol. 1997;71:

138–144.

21.   Eichwald C, De Lorenzo G, Schraner EM, Papa G, Bollati M, Swuec P, et

al. Identification of a Small Molecule That Compromises the Structural

Integrity of Viroplasms and Rotavirus Double-Layered Particles. J Virol.

American Society for Microbiology (ASM); 2018;92. doi:10.1128/JVI.01943-

17

22.   Contin R, Arnoldi F, Campagna M, Burrone OR. Rotavirus NSP5

orchestrates recruitment of viroplasmic proteins. J Gen Virol. 2010;7:

1782–1793.

23.   Afrikanova I, Fabbretti E, Miozzo MC, Burrone OR. Rotavirus NSP5

phosphorylation is up-regulated by interaction with NSP2. J Gen Virol.

1998;79: 2679–2686. doi:10.1099/0022-1317-79-11-2679

24.   Eichwald C, Jacob G, Muszynski B, Allende JE, Burrone OR. Uncoupling

substrate and activation functions of rotavirus NSP5: phosphorylation of

Ser-67 by casein kinase 1 is essential for hyperphosphorylation. Proc Natl

Acad Sci U S A. 2004;101: 16304–16309.

25.   Campagna M, Budini M, Arnoldi F, Desselberger U, Allende JE, Burrone
OR. Impaired hyperphosphorylation of rotavirus NSP5 in cells depleted of
casein kinase 1alpha is associated with the formation of viroplasms with
altered morphology and a moderate decrease in virus replication. J Gen
Virol. 2007;10: 2800–2810.

26.   Eichwald C, Vascotto F, Fabbretti E, Burrone OR. Rotavirus NSP5:
mapping phosphorylation sites and kinase activation and viroplasm
localization domains. J Virol. 2002;76: 3461–3470.

27.   Martin D, Ouldali M, Menetrey J, Poncet D. Structural organisation of the
rotavirus nonstructural protein NSP5. J Mol Biol. 2011;413: 209–221.

28.   Kanai Y, Komoto S, Kawagishi T, Nouda R, Nagasawa N, Onishi M, et al.
Entirely plasmid-based reverse genetics system for rotaviruses. Proc Natl
Acad Sci U S A. National Academy of Sciences; 2017;114: 2349–2354.
doi:10.1073/pnas.1618424114

29.   Komoto S, Fukuda S, Ide T, Ito N, Sugiyama M, Yoshikawa T, et al.
Generation of Recombinant Rotaviruses Expressing Fluorescent Proteins
by Using an Optimized Reverse Genetics System. J Virol. 2018;92:
e00588-18. doi:10.1128/JVI.00588-18

30.   Pertel T, Hausmann S, Morger D, Züger S, Guerra J, Lascano J, et al.
TRIM5 is an innate immune sensor for the retrovirus capsid lattice. Nature.
2011;472: 361–365. doi:10.1038/nature09976

31.   Dull T, Zufferey R, Kelly M, Mandel RJ, Nguyen M, Trono D, et al. A third-
generation lentivirus vector with a conditional packaging system. J Virol.
1998;72: 8463–71.

32.   Yusa K, Zhou L, Li MA, Bradley A, Craig NL. A hyperactive piggyBac
transposase for mammalian applications. Proc Natl Acad Sci U S A.
National Academy of Sciences; 2011;108: 1531–6.

doi:10.1073/pnas.1008322108

33.   Eichwald C, Arnoldi F, Laimbacher AS, Schraner EM, Fraefel C, Wild P, et

al. Rotavirus viroplasm fusion and perinuclear localization are dynamic

processes requiring stabilized microtubules. PLoS One. Public Library of

Science; 2012;7: e47947. doi:10.1371/journal.pone.0047947

34.   Vascotto F, Campagna M, Visintin M, Cattaneo A, Burrone OR. Effects of

intrabodies specific for rotavirus NSP5 during the virus replicative cycle. J

Gen Virol.; 2004;85: 3285–3290.

35.   Ito H, Minamoto N, Goto H, Rong LT, Sugiyama M, Kinjo T. Expression of

the major inner capsid protein, VP6, of avian rotavirus in mammalian cells.

Vet Microbiol. 1996;49: 257–65.

36.   Tosser G, Labbé M, Brémont M, Cohen J. Expression of the major capsid

protein VP6 of group C rotavirus and synthesis of chimeric single-shelled

particles by using recombinant baculoviruses. J Virol. American Society for

Microbiology (ASM); 1992;66: 5825–31.

37.   McDonald SM, Patton JT. Assortment and packaging of the segmented

rotavirus genome. Trends Microbiol. Elsevier Ltd; 2011;19: 136–144.

doi:10.1016/j.tim.2010.12.002

38.   Borodavka A, Desselberger U, Patton JT. Genome packaging in multi-

segmented dsRNA viruses: distinct mechanisms with similar outcomes.

Curr Opin Virol. Elsevier; 2018;33: 106–112.

doi:10.1016/J.COVIRO.2018.08.001

39.   Criglar JM, Anish R, Hu L, Crawford SE, Sankaran B, Prasad BVV, et al.

Phosphorylation cascade regulates the formation and maturation of

rotaviral replication factories. Proc Natl Acad Sci. 2018;115: E12015–

E12023. doi:10.1073/pnas.1717944115

40.   Liu C, Li Y, Semenov M, Han C, Baeg GH, Tan Y, et al. Control of beta-

catenin phosphorylation/degradation by a dual-kinase mechanism. Cell.
2002;108: 837–47.

41.   Amit S, Hatzubai A, Birman Y, Andersen JS, Ben-Shushan E, Mann M, et
al. Axin-mediated CKI phosphorylation of β-catenin at Ser 45: A molecular
switch for the Wnt pathway. Genes Dev. 2002; doi:10.1101/gad.230302

42.   Verheyen EM, Gottardi CJ. Regulation of Wnt/β-catenin signaling by
protein kinases. Developmental Dynamics. 2010. doi:10.1002/dvdy.22019

43.   Goonawardane N, Yin C, Harris M. A pivotal role of serine 225
phosphorylation in the function of hepatitis C virus NS5A revealed with the
application of a phosphopeptide antiserum and super-resolution
microscopy. bioRxiv. 2018; doi:10.1101/387407

44.   Kim J, Lee D, Choe J. Hepatitis C virus NS5A protein is phosphorylated by
casein kinase II. Biochem Biophys Res Commun; 1999;257: 777–781.

45.   Ross-Thriepland D, Mankouri J, Harris M. Serine Phosphorylation of the
Hepatitis C Virus NS5A Protein Controls the Establishment of Replication
Complexes. Diamond MS, editor. J Virol. 2015;89: 3123–3135.
doi:10.1128/JVI.02995-14

46.   Keating JA, Bhattacharya D, Lim P-Y, Falk S, Weisblum B, Bernard KA, et
al. West Nile virus methyltransferase domain interacts with protein kinase
G. Virol J. BioMed Central; 2013;10: 242. doi:10.1186/1743-422X-10-242

47.   Spencer K-A, Dee M, Britton P, Hiscox JA. Role of phosphorylation
clusters in the biology of the coronavirus infectious bronchitis virus
nucleocapsid protein. Virology. 2008;370: 373–381.
doi:10.1016/j.virol.2007.08.016

48.   Bhattacharya D, Hoover S, Falk SP, Weisblum B, Vestling M, Striker R.
Phosphorylation of yellow fever virus NS5 alters methyltransferase activity.
Virology. 2008;380: 276–284. doi:10.1016/j.virol.2008.07.013

Figure S3

Figure 1

Figure 2

A   rRV

37 —

25 —

NSP5

actin

B   rRV-NSP5/

ΔT

Δ176-180

Cells: MA-NSP5 MA104   MA-NSP5 MA104

1
2
3-4

7-8
9

A

rRV-wt   rRV-NSP5/S67A

5 hpi

10 hpi

15 µm

B

rRV-wt

NSP5   VP6   Merge

rRV-NSP5/S67A

NSP5   VP6   Merge

NSP2   VP6   Merge

VP2   VP6   Merge

C

rRV-NSP5/S67A

NSP5-EGFP

NSP2-mCherry

D

E

A

B

Figure 5

Figure 8

A

rRV
NSP5/KO:

MA104   MA-ΔT   MA-Δ3

−   +   −   +   −   +

kDa

100 —   VP2

37 —

25 —   NSP5

actin

B

rRV-NSP5/KO

MA104   NSP5 | VP6 | Merge

MA-ΔT   NSP5 | VP6 | Merge

MA-Δ3   NSP5 | VP6 | Merge

NSP2 | VP6 | Merge

C

rRV-NSP5/KO

Viral Titer (log₁₀FFU/ml)

**
**
**

#   #   #

MA-NSP5   MA104   MA-ΔT   MA-Δ3