Multiple PLCβ signaling pathways in taste receptor cells contribute to the detection of bitter, sweet and umami stimuli

Debarghya Dutta Banik[1], Eric D. Benfey[1], Laura E. Martin[2], Kristen E. Kay[2], Gregory C. Loney[2], Amy R. Nelson[1], Zachary C. Ahart[1], Barrett T. Kemp[1], Bailey R. Kemp[1], Aurelie Vandenbeuch[3], Ann-Marie Torregrossa[2,4] and Kathryn F. Medler*[1]

[1]Department of Biological Sciences, University at Buffalo, Buffalo, NY 14260, USA.
[2]Department of Psychology, University at Buffalo, Buffalo, NY 14260, USA.
[3]Department of Otolaryngology, University of Colorado, Aurora, CO, USA
[4]Center for Ingestive Behavior Research, University at Buffalo, Buffalo, NY 14260, USA

*Corresponding author: Email: kmedler@buffalo.edu

Summary: This study identifies a new role for PLCβ3 in the transduction of bitter, sweet and umami taste stimuli and demonstrates that multiple taste cell populations respond to these stimuli.

ABSTRACT

Taste receptor cells use multiple signaling pathways to detect chemicals. While salty and sour stimuli bind ion channels, bitter, sweet and umami stimuli activate a GPCR/PLCβ2/IP$_3$R3 signaling pathway. Current thinking is that all bitter, sweet and umami stimuli are detected by Type II cells via a PLCβ2/IP$_3$R3 pathway. Using live cell calcium (Ca^{2+}) imaging on isolated taste receptor cells from mice lacking IP$_3$R3 in Type II cells, we find many taste cells still robustly respond to bitter, sweet and umami stimuli, including a subset of Type III taste cells. Immunohistochemical analysis reveals that PLCβ3 is not expressed in Type II cells but is present in other taste cell populations, including a subset of Type III cells. Loss of either IP$_3$R3 or PLCβ3 in taste receptor cells cause comparable reductions in c-Fos labeling in the nucleus of the solitary tract following oral infusion of quinine. Short term behavioral assays also demonstrate that loss of either IP$_3$R3 from Type II cells or loss of PLCβ3 causes similar taste deficits, indicating that both molecules are required for normal taste transduction. Our data identify a critical, and previously unknown, role for PLCβ3 in taste transduction and demonstrate that input from multiple PLCβ pathways/taste receptor cell populations are required to transduce bitter, sweet and umami stimuli.

SIGNIFICANCE STATEMENT

It is currently thought that the transduction of bitter, sweet and umami stimuli in

taste cells depends solely on GPCR/PLCβ2/IP$_3$R3 signaling in Type II taste receptor

cells. However, in the absence of either PLCβ2 or IP$_3$R3, mice have reduced, but not

abolished, ability to detect these stimuli, suggesting that other taste cells also contribute

to taste transduction. When IP$_3$R3 is absent, Type II cells do not generate a Ca^{2+} signal

to these taste stimuli; however, other taste receptor cells respond to bitter, sweet and

umami stimuli, including a subset of Type III cells. We identified a PLCβ3 pathway that

is expressed in non-Type II taste cells, including a subset of Type III cells. Loss of

PLCβ3 significantly reduced taste cell responses, cFos-labeling in the nucleus of the

solitary tract, and caused significant behavioral deficits. We conclude that PLCβ3 has

an important role in taste transduction.

Chemicals in the oral cavity are detected by taste receptor cells which are

grouped together in taste buds found in epithelial specializations in the tongue called

papillae. There are three main papillae: Circumvallate (CV), foliate (Fol), and fungiform

(Fun) which differ in their localization on the tongue and innervation by the cranial

nerves. Despite these differences, taste papillae all house taste buds that are

comprised of distinct cell types that vary in their signaling functions within the bud (1).

The current view of taste transduction is that taste buds are comprised of Type I,

Type II and Type III taste receptor cells (2). Type I cells are thought to primarily function

as support cells and share some characteristics of glia cells (3), while Type II cells

detect bitter, sweet or umami stimuli through the activation of specific GPCRs for each

of these taste qualities. The subsequent transduction of these taste qualities within the

cells are due to a single signaling pathway that is comprised of phospholipase $C\beta2$

($PLC\beta2$) that activates the inositol 1,4,5-trisphosphate receptor type 3 (IP_3R3) on the

endoplasmic reticulum to cause Ca^{2+} release (4-6). This Ca^{2+} release activates the

transient receptor potential cation channel subfamily M members 4 and 5 (TRPM4 and

TRPM5) which depolarize the cell sufficiently to activate the release of ATP through the

calcium homeostasis modulator 1 (CALHM1) channel (5, 7-11). The expression of

$PLC\beta2$, IP_3R3 and TRPM5 is restricted to Type II taste cells within the bud and these

cells lack both voltage-gated calcium channels (VGCCs) and conventional chemical

synapses (model shown in Figure 1A)(12-16). Type III cells detect sour and salt stimuli

through ionotropic receptors that depolarize the cell to activate VGCCs and cause

vesicular neurotransmitter release (17-23) (see Figure 1A). It is currently thought that

Type III cells do not respond to bitter, sweet or umami stimuli.

While this is the widely accepted view of signal transduction in peripheral taste

cells, multiple genetic studies have reported that the deletion of individual signaling

proteins in Type II cells, including PLCβ2, IP$_3$R3 and TRPM5, does not abolish the

ability to detect all bitter sweet and umami stimuli (24-28). Similarly, the loss of Type II

cells leads to severely reduced, but not abolished, responses to bitter, sweet and

umami stimuli (29). These studies support the idea that additional signaling pathways

are present in taste cells, but to date, few studies have focused on identifying these

alternatives (30, 31). We previously reported that some taste cells generate Ca^{2+}

responses to a bitter stimulus and cell depolarization, suggesting that taste cells with

VGCCs, presumably Type III cells, respond to at least one bitter compound (32). A

separate report also found some taste cells with VGCCs can respond to bitter stimuli

(19). These studies led us to hypothesize that in addition to Type II cells, a separate

population of taste cells may contribute to the detection of bitter, sweet and umami

stimuli. To test this hypothesis, we used live cell imaging in mice that lack IP$_3$R3 and

found that many taste cells still responded to bitter, sweet and umami stimuli, including

a subset of Type III cells. These cells responded to taste stimuli through a distinct

PLCβ which we identified as PLCβ3. We conclude that multiple populations of taste

cells use distinct signaling pathways to respond to bitter, sweet and umami stimuli.

RESULTS

Bitter, sweet and umami taste evoked responses are still present in IP$_3$R3-KO

mice

IP$_3$R3 is part of the canonical signaling pathway in Type II cells that respond to

bitter, sweet and umami stimuli and its expression is localized in Type II cells (4, 12, 16,

33). One study using single cell PCR reported mRNA levels for IP$_3$R3 in both Type II

cells and some SNAP25+ cells (a Type III marker) but did not evaluate protein

expression for these compounds (34). In contrast, another study using an IP$_3$R3-GFP

mouse reported that GFP+ cells did not functionally express VGCCs, a physiological

marker for Type III cells (32). While these conflicting reports suggest that it is possible

that IP$_3$R3 expression is not restricted to Type II cells, for the purposes of our current

study, we are presuming that IP$_3$R3 is only expressed in Type II cells.

To evaluate the taste evoked signaling in taste receptor cells that lack the ability

to release Ca^{2+} via IP$_3$R3, we used a transgenic mouse in which GFP replaces the

coding region of IP$_3$R3. In these mice, GFP labels the cells that should express IP$_3$R3

but no longer do so. Initial immunohistochemical analyses found that taste cells from the

CV papillae of wild type mice were successfully labeled with anti-IP$_3$R3 antibody (n=3;

Figure S1A), while their KO littermates lacked IP$_3$R3 labeling but instead have GFP

expression (n=6; Figure S1B). Furthermore, anti-IP$_3$R3 labels the GFP expressing cells

in IP$_3$R3-heterzygous mice (n=4; Figure S1C); confirming that GFP expression identifies

the taste cells that normally express IP$_3$R3. To further characterize this mouse, we

evaluated the expression of other proteins that are part of this established signaling

pathway in Type II cells.   The GFP expression in the KO mice co-localized with both

PLCβ2 (n=4; Figure S1D) and gustducin (n=3; Figure S1E) labeling. As expected,

gustducin expression was restricted to the IP$_3$R3-GFP-KO cells but not all GFP

expressing cells had gustducin while there was complete overlap between PLCβ2

labeling and GFP. Co-localization analyses are reported in Figures S1F. Thus the

other components of the signaling pathway normally found in Type II cells are intact but

these cells no longer have the capacity to release Ca^{2+} in response to bitter, sweet or

umami stimuli.

We evaluated this IP_3R3-GFP-KO mouse using Fura2-AM to measure the taste-

evoked Ca^{2+} responses in taste receptor cells. All measurements were made in

isolated taste cells that were no longer associated with the taste bud to ensure that the

taste-evoked Ca^{2+} responses were due solely to the activity of the individual taste cells

and not cell-to-cell communication (example in Figure 1B, C). We applied multiple

bitter, sweet and umami stimuli as well as 50mM KCl to isolated cells from CV, Fol or

Fun papillae of the wild type and IP_3R3-KO mice. Since our goal was to identify all (or

most) responsive taste cells, we used the stimulus concentration that generated a

maximal Ca^{2+} signal. These concentrations were based on control experiments we

used to identify the lowest stimulus concentration that generated the maximal Ca^{2+}

signal (examples shown in Figure S2, A-C). This increased our likelihood of identifying

responsive taste cells but did not generate any non-specific responses (Figure S2, D,

E). As shown in Figure S2C, a GFP+ cell did not respond to any concentration of

stimulus applied. Thus, the taste cells respond to a wide range of stimulus

concentrations and the reported responses are not due to a concentration or osmotic

effect.

We reasoned that if the Type II cells were solely responsible for all bitter, sweet

and umami taste transduction, then the loss of IP_3R3 in Type II cells would abolish all

Ca^{2+} responses to bitter, sweet and umami stimuli. In the IP_3R3-KO mice, GFP-

expressing cells lack IP$_3$R3 and did not respond to any taste stimuli tested (n=80, Figure

2B, S2C); however, a large number of non-GFP expressing taste cells responded to the

stimuli. Initially, the response frequencies *(total number of responsive cells/total*

number of cells stimulated) were measured for all taste cells from the wild type (WT)

and IP$_3$R3-KO mice, including both GFP+ and GFP- cells (Figure 2A). The table in

Figure 2A reports the taste response frequencies in wild type (WT) and IP$_3$R3-KO mice

with the GFP+ cells included in the analysis. Not all cells were exposed to all stimuli

and some cells responded to multiple stimuli. These experimental conditions preclude

the data from being additive, so we compared the frequency of responses for a given

stimulus between the wild type and IP$_3$R3-KO mice. These data demonstrate that the

overall responsiveness to bitter, sweet, and/or umami stimuli was significantly reduced

(p<0.0001) in the KO mice compared to WT cells. The likelihood that taste cells from

CV would respond to taste stimuli was significantly reduced (p=0.0003). The

responsiveness of the taste cells from the Fol and Fun papillae was not significantly

affected by the loss of IP$_3$R3, even though the GFP+ cells were included in this

analysis.

To further characterize the response profiles of the non-Type II taste cells, we

analyzed the non-GFP expressing cells and plotted the response frequencies for each

papillae and stimulus (Figure 2B-E). The GFP+ cells were not included in these

response analyses and descriptions of the different taste cell populations are shown in

Supplementary Table 1. Figure 2B compares the overall percent responsiveness of WT

and KO mice for each papillae type (without the GFP+ cells) while the graphs in Figures

2C-E show the percentage of responsive cells in each papillae type for each specific

stimulus applied. The overall responsiveness of taste cells was only significantly

reduced in the CV papillae when IP_3R3 was absent (p<0.01). Analysis of the individual

stimuli for each papillae type (Figures 2C-E) identified significant reductions in the

responsiveness to the sweet stimuli tested in the KO mice. Individual cell numbers are

provided in Figure S3. Therefore, removing a key protein in the canonical

GPCR/PLCβ2/IP_3R3 signaling pathway in Type II cells does not eliminate the ability of

all taste cells to generate Ca^{2+} signals when stimulated by bitter, sweet or umami

stimuli. Based on these data, we hypothesize there are additional signaling pathways

that respond to these stimuli.

Bitter, sweet and umami stimuli are dependent on a PLCβ signaling pathway in

IP_3R3-KO mice

To identify the type of signaling pathway that generates the taste responses in

the non-GFP cells in the IP_3R3-KO mice, we analyzed bitter (5mM Den), sweet (20mM

sucrose) and umami (10mM MPG) stimuli and found they generated signals in Ca^{2+}-free

Tyrode's, indicating these signals depend on Ca^{2+} release from internal stores (Figure

3A,D,G). We confirmed these results by applying thapsigargin to disrupt the internal

Ca^{2+} stores which abolished the taste evoked Ca^{2+} signals (Figure 3B,E,H). These

responses were also eliminated when the general PLC blocker U73122 was applied

(Figure 3C,F,I), indicating that these taste evoked Ca^{2+} responses in the IP_3R3-KO mice

are due to the activation of another PLC signaling pathway. Individual data points for

each experiment are shown in Figure S4.

PLCβ3 expression in peripheral taste cells

To identify other PLCβ isoforms expressed in taste receptor cells, we analyzed

previously published RNAseq data (35). We found that PLCβ2 and PLCβ3 were

expressed at comparable levels and were the predominant PLCβ isoforms present in

taste cells. Therefore, we investigated the possibility that PLCβ3 is expressed in

peripheral taste cells. Immunohistochemical experiments in the IP$_3$R3-KO mouse

revealed strong PLCβ3 labeling that was separate from the Type II cells (identified with

GFP expression) in the CV, Fol and Fun papillae (Figure 4A-C). Colocalization analysis

is shown in Figure 4D. These data were confirmed using both the TRPM5-GFP mouse

as well as co-labeling with PLCβ2 in C57BL/6 mice (Type II cell markers, Figure S5A, B,

colocalization analysis shown in S5D). Co-labeling with the Type I taste marker

NTPDase2 found that PLCβ3 is also not expressed in Type I cells (Figure S5C). We

used qPCR to measure the relative amounts of mRNA for PLCβ3 and PLCβ2 in the

different papillae (Figure S5E,F) and found that both PLCβ2 and PLCβ3 are expressed

in CV, Fol and Fun papillae. We conclude that both PLCβ2 and PLCβ3 are highly

expressed in peripheral taste cells but that PLCβ3 is not expressed in either Type I or

Type II taste cells. Our next goal was to identify the cell population that expresses the

PLCβ3 signaling pathway and responds to bitter, sweet and umami stimuli.

Some Type III cells respond to bitter, sweet and umami stimuli

The only other identified taste receptor cell type is the Type III cell. Type III cells

are the only taste cell population that express VGCCs (36) and can be functionally

identified in live cell imaging by the ability to respond to cell depolarization with a Ca^{2+}

influx through the opening of VGCCs (19, 37). Earlier studies have reported that Type III

cells are sour sensitive and that a subset of the sour-sensitive Type III cells is also salt

sensitive (17, 19, 38). To functionally identify Type III cells, we applied 50mM KCl to

cause a cell depolarization which generated a Ca^{2+} signal if the cell expresses VGCCs.

Therefore, if a cell responded to 50mM KCl with a Ca^{2+} signal, we concluded that it

expresses VGCCs and identified it as a Type III cell. We found that some Type III taste

cells in the IP_3R3-KO mouse responded to bitter, sweet and umami stimuli (Figure 5A-

C). To confirm that the taste cells that responded to 50mM KCl and bitter, sweet or

umami stimuli were Type III cells, we tested a subset of cells with sour (citric acid) and

salt (NaCl) stimuli (Figure 5D-F). All of the taste cells that responded to 50mM KCl

responded to citric acid with a Ca^{2+} signal (CA) while some of these cells also

responded to bitter, sweet and/or umami stimuli (Figure 5D, F). Interestingly, a

separate subset of 50mM KCl/CA cells responded to NaCl with a Ca^{2+} signal but were

not sensitive to the bitter, sweet, umami stimuli (Figure 5E, F). This indicates that the

sour/salt sensitive Type III cells are separate from the sour sensitive Type III cells that

respond to bitter, sweet and/or umami stimuli with a Ca^{2+} signal. Thus a subset of Type

III cells are dual responsive (DR) because they express VGCCs and are sour sensitive

but also respond to bitter, sweet and/or umami stimuli.

To determine if the loss of IP_3R3 in Type II cells affected the activity of DR cells,

we compared the percentage of cells that responded to any stimulus (bitter, sweet,

umami or 50mM KCl) in wild type and IP_3R3-KO mice. These data include all taste cells

from all three papillae taken from Figure 2A. As shown before, the overall

responsiveness to taste stimuli in the IP_3R3-KO mice was significantly reduced

compared to wild type (Figure 5G); however, the percentage of DR cells in the IP_3R3-

KO was not different from wild type mice (Figure 5G). Therefore, the ability of the DR

cells to respond to taste stimuli does not depend on the presence of IP$_3$R3 in Type II

cells (Figure 5G, H).

We also performed control experiments using the GAD67-GFP mice as a cellular

marker to identify Type III taste cells (39) and found that both GAD67-GFP positive and

negative taste cells responded to taste stimuli + 50mM KCl with Ca^{2+} signals (Figure

S6A-B). DR cells were also identified in C57BL/6 mice (Figure S6C). A recent study

reported that some Type III cells from C57BL/6 mice responded to bitter stimuli (19) in

agreement with our findings. Thus, DR Type III taste cells are present in taste cells in

multiple mouse lines.

PLCβ3 contributes to taste evoked signaling in a subset of Type III cells

So far our data have identified that 1) there is a population of taste cells that do

not depend on IP$_3$R3 (in Type II cells) to respond to bitter, sweet and umami, 2) PLCβ3

is expressed in taste cells, and 3) some Type III cells are DR and can respond to bitter,

sweet and umami stimuli. We hypothesized that PLCβ3 is involved in the signaling of

the DR cells. We next performed live cell imaging on isolated cells from CV, Fol, and

Fun papillae of PLCβ3-KO and wild type mice to evaluate how the loss of PLCβ3 affects

taste cell responsiveness, in particular the responsiveness of the DR cells. Loss of

PLCβ3 resulted in significant reductions in the overall responsiveness of the PLCβ3-KO

Fol and Fun cells but not the CV cells (Figure 6A,B). Knocking out PLCβ3 had a two-

fold effect, it reduced the number of responses to bitter, sweet or umami stimuli and it

significantly reduced the number of DR taste cells in CV, Fol and Fun taste cells (Figure

6C-E). Control experiments found no change in the ability of Type III cells to respond to

sour or salt stimuli (Figure S7A-C). Type II cells were also still functional in PLCβ3-KO

mice (Figure S7D-F). Individual cell numbers are shown in Figure S8. These data

indicate that PLCβ3 is contributing to the activity of the DR cells.

We next determined if PLCβ3 is expressed in Type III cells using SNAP-25 which

is a synaptic protein that is a cellular marker for Type III taste cells (16, 40).

Immunohistochemical analysis found some co-expression of PLCβ3 and SNAP25 in

wild type mice (Figure S9A,B,E). There was no loss of SNAP-25 expression in the

PLCβ3-KO mouse compared to WT (Figure S9C,D). Thus, PLCβ3 is expressed in some

Type III cells and the loss of this protein did not affect the expression of SNAP25 in

Type III cells.

The significant reduction in the number of taste evoked Ca^{2+} responses in the DR

cells from PLCβ3-KO mice suggests that PLCβ3 is responsible for transducing most of

the taste-evoked responses in these cells. To determine if the loss of both IP_3R3 and

PLCβ3 would completely abolish the ability to respond to bitter, sweet and umami

stimuli, we attempted to generate a double knock out mouse by breeding the IP_3R3-KO

and PLCβ3-KO mice but the mice were not viable.

Loss of PLCβ3 causes deficits in the taste-evoked activity in the nucleus of the

solitary tract.

To determine if the loss of IP_3R3 or PLCβ3 in taste cells affected the taste signal

that is sent to the brain, we measured the neural activity in the nucleus of the solitary

tract (NTS) using c-Fos labeling as a marker for activated neurons. This approach is

commonly used to identify recently activated neurons, including neurons in the taste

pathway (41-45). The NTS receives input from the gustatory neurons and is the first

synaptic relay from the peripheral taste system. Mice (n=3 for each condition) were

orally infused with quinine (5mM) or water for 30 min. The level of c-Fos labeling for

both IP₃R3-KO and PLCβ3-KO mice were compared to WT and water controls. Loss of

either of these proteins resulted in a significant reduction in c-Fos activity to a level that

was comparable to water (Figure 7). Control experiments demonstrate that neither

IP₃R3 nor PLCβ3 are expressed in this region of the NTS (Figure S10).

Loss of PLCβ3 significantly affects taste-driven licking for bitter, sweet and

umami

We then performed brief-access licking experiments to determine if the loss of

taste-evoked signals in the PLCβ3-KO mice correlated with loss of taste sensitivity.

Measurements were also made with IP₃R3-KO mice for comparison. Since the PLCβ3-

KO mouse has a mixed genetic background while the IP₃R3-KO mice are in a C57BL/6

background, wild type littermates for both mouse lines were used as controls (Figure 8).

The wild type mice both demonstrated concentration dependent decreases in licking to

denatonium while PLCβ3-KO and IP₃R3-KO mice treated the solution similarly to water

except for the highest concentration (20mM Den, Figure 8A). Wild type mice treated

MSG (+ 10μM amiloride) solutions as more palatable than water up to 400mM and then

decreased licking at the highest concentrations (600 and 800mM, Figure 8B). Both of

the KO mice treated MSG like water. In the brief access test for the artificial sweetener,

Acesulfame K (AceK), wild type mice increased licking up to 2 mM and then decreased

licking at higher concentrations. Neither of the KO mice had a comparable concentration

dependent change in licking (Figure 8C). Because sucrose is hedonically positive under

wild type conditions, the lick score is calculated as licks to the stimulus minus licks to

water, therefore a score close to zero resembles water. Neither the PLCβ3-KO nor

IP$_3$R3-KO mice showed a concentration-dependent effect on licking to sucrose until

they were presented with the highest concentration.   In contrast, both sets of wild type

mice showed strong concentration dependent increases in licking to sucrose (Figure

8D). For the salt stimulus (NaCl), there were no significant differences in the licking

behavior for either of the KO mice compared to wild type mice, indicating that the ability

to respond to salt is not affected by loss of either PLCβ3 or IP$_3$R3 (Figure 8E). Thus

loss of either IP$_3$R3 or PLCβ3 caused comparable, and almost complete, loss of

preference for the bitter, sweet and umami stimuli tested but not salt taste.

Response profiles for the dual-responsive taste cells

Finally, we analyzed *how* DR cells respond to taste stimuli. We evaluated a

subset of DR cells that were stimulated with at least one bitter, one sweet and one

umami stimulus *and responded* to at least one of these stimuli. If the DR cells were not

stimulated with all three types of taste qualities, they were excluded from this analysis.

We created response profiles for bitter, sweet and umami in wild type and IP$_3$R3-KO

mice (Figure S11A-C) and found that the DR taste cells were most likely to be broadly

tuned (respond to all three types of stimuli). Few, if any, of the DR cells responded to a

single taste quality (in addition to 50mM KCl). We found that the loss of IP$_3$R3 did not

influence these response profiles.

Three populations of taste cells respond to bitter, sweet and umami stimuli

We repeated the same analysis on taste cells from wild type and IP$_3$R3-KO mice

that were stimulated with bitter, sweet, umami and 50mM KCl but *did not respond to*

50mM KCl (Figure S11D-F). Therefore, wild type taste cells analyzed included all the

Type II taste cells as well as any other taste cells that responded to bitter, sweet and/or

umami stimuli but lack VGCCs. We only analyzed taste cells that responded to at least

one of these stimuli and compiled a response profile for these taste cells from wild type

and IP$_3$R3-KO mice. These taste cells were much less likely to be broadly tuned in both

the WT and KO taste cell populations.

While the taste-only cells from wild type mice express IP$_3$R3 and therefore should

have Type II cells contributing to these response profiles, the analysis of the IP$_3$R3-KO

mice was focused on taste cells that were not functional Type II taste cells (due to the

absence of IP$_3$R3). These cells also did not respond to 50mM KCl with a Ca^{2+} signal

and are not Type III cells. Therefore, this taste cell population lacks functional Type II

cells and VGCCs but responds to bitter, sweet and/or umami stimuli. These functional

data suggest that there are at least three populations of taste cells that can respond to

bitter, sweet and umami stimuli: 1) Type II cells (identified as cells using IP$_3$R3), 2) a

subset of Type III cells (which respond to 50mM KCl with a Ca^{2+} signal) and 3) another

population of taste cells that are not Type II or Type III cells (neither reliant on IP$_3$R3 nor

responsive to 50mM KCl).   Future studies are needed to identify and characterize this

taste cell population.

DISCUSSION

Our data establish that PLCβ3 signaling in taste receptor cells is important in the

detection of bitter, sweet and umami taste stimuli. The presence of a second signaling

pathway that detects these stimuli resolves a long standing inconsistency in the taste

field. One study concluded that the PLCβ2 signaling pathway in Type II cells is entirely

responsible for the detection of bitter, sweet and umami stimuli (5). A second study

focused on understanding the role of Skn-1a in taste cell formation reported that the

loss of Skn-1a caused a significant remodeling of the taste bud. Using *in situ*

hybridization, the authors did not detect the canonical signaling molecules normally

found in Type II cells but did identify an increase in the Type III cell marker expression.

They also reported significant behavioral and gustatory nerve defects for bitter, sweet

and umami stimuli in the absence of Skn-1a, indicating that Type II cells are required to

detect these stimuli (29). However, other studies reported that knocking out any

component of the GPCR/PLCβ2/IP$_3$R3 pathway resulted in reduced, but not abolished,

signaling for these stimuli (24-28, 46-49). There are deficits in taste-evoked signaling

when any part of the GPCR/PLCβ2/IP$_3$R3 pathway is nonfunctional, but they are

variable and there was never complete abolishment of the responses. Nevertheless,

the current dogma within the taste field is that only this signaling pathway is responsible

for all bitter, sweet and umami transduction (5). While this is puzzling, it is likely due to

the lack of another identified signaling pathway that transmits taste signals. We have

now identified that a PLCβ3 signaling pathway also detects bitter, sweet and umami

stimuli.

Indeed, a closer inspection of the reported findings from these earlier studies

reveals differential deficits in taste responses when the PLCβ2/IP$_3$R3 signaling pathway

is nonfunctional. Evaluations of IP$_3$R3 loss in mice found impaired behavior and nerve

responses to different taste stimuli, but not abolishment of these signals (25). They

reported reduced behavioral sensitivity for bitter and sweet stimuli at some

concentrations but no differences from controls at other concentrations. The cranial

nerve recordings revealed deficits in signals that varied with the cranial nerve and taste

stimulus being tested (25). These findings confirm that IP$_3$R3 is required for normal

taste signaling but also suggest that it is not the only pathway involved. Similarly, work

focused on the loss of PLCβ2 reported reduced, but not abolished, avoidance of bitter

compounds by PLCβ2-KO mice (28). In this study, the PLCβ2-KO mice did not avoid

denatonium as well as wild type mice but still reduced licking as denatonium

concentration increased. The PLCβ2-KO mice did not lick normally for sucrose but did

lick normally for salt and sour stimuli. The authors concluded that a PLCβ2-

independent pathway also contributes to taste transduction (28) which agrees with the

IP_3R3-KO study (25). Other studies found that PLCβ2 and IP_3R3 co-localize (4, 12, 33)

in Type II taste cells, so it is not surprising that loss of either protein causes similar

deficits in taste-driven behaviors.

We measured the neural activity in the NTS using c-Fos labeling as a marker for

activated neurons to determine how the loss of PLCβ3 affects central processing of

taste information from the periphery (Figure 7). We used oral infusions of taste stimuli in

awake and behaving animals because it removes any potential confounds due to

anesthetization that is used in other approaches. Since the NTS receives input from the

gustatory neurons and is the first synaptic relay from the peripheral taste system, these

data confirm that loss of either IP_3R3 or PLCβ3 in the taste cells abolishes the taste

signal that is sent to the brain. These data from the NTS agrees entirely with our

behavioral data (Figure 8) which together support the idea that both IP_3R3 and PLCβ3

have important roles in taste transduction.

Loss of PLCβ3 caused significant behavioral impairments for the bitter, sweet

and umami stimuli that were comparable to the behavioral deficits when IP_3R3 was not

present in Type II cells. The loss of either protein was specific to these taste stimuli and

did not affect salt taste (Figure 8E). These data confirm that the lack of either IP$_3$R3

signaling in Type II or PLCβ3 signaling in taste cells had a comparable impact on taste

behaviors (Figure 8). Since PLCβ3 and IP$_3$R3 are not expressed in the NTS (Figure

S10) or the geniculate ganglia (50), these data strongly support the idea that input from

both of these signaling molecules in taste cells is required for normal taste behavior.

Therefore, if signaling from either population of taste cells (those using IP$_3$R3 or those

using PLCβ3) is impaired, taste information is not properly conveyed to the brain. This

conclusion agrees with other studies which suggest that Type II cells are not the only

cells that transduce bitter, sweet and umami stimuli (24, 25, 27, 28).

Our data support the idea that multiple taste receptor mechanisms, outside of

those canonically identified, are involved in the detection of different taste stimuli.

Multiple laboratories have independently concluded that the identified taste receptors

are not solely responsible for transducing all bitter, sweet and umami stimuli. For

instance, the detection of some carbohydrates is not impaired by the loss of T1R2 or

T1R3, the identified sweet receptors (47-49, 51), and umami stimuli appear to be

transduced by receptors in addition to the T1R1+T1R3 heterodimer (31, 52-58).

Moreover, because the identified T1Rs and T2Rs appear to have restricted expression

in Type II taste cells (5, 59), the presence of the PLCβ3 signaling pathway in non-Type

II cells bolsters the hypothesis that additional taste receptor mechanisms are likely

involved in transduction.

Our data clearly demonstrate that a subset of Type III cells (DR) are sensitive to

bitter, sweet and/or umami stimuli in contrast to the idea that Type III cells only detect

sour and salty stimuli (19, 38). Furthermore, it appears likely that PLCβ3 is expressed

in Type III cells and is responsible for the bitter, sweet and umami responses recorded

in a subset of Type III cells. We previously reported that a PLCβ3/IP$_3$R1 signaling

pathway is expressed in a subset of taste cells that are not Type II cells (32) which our

current data suggest is a subset of Type III cells. These physiological findings are

supported by recent RNA-sequencing analysis of individual Type II and Type III cells

(37) which revealed PLCβ3 expression in a subset of Type III cells.

We found that many of the DR Type III cells were broadly tuned and responded

to two or more taste qualities, including taste stimuli of different modalities (Figure S6.

S11). Importantly, our studies used isolated taste cells that were not in contact with

neighboring cells. Earlier studies using intact buds or taste epithelium reported that a

subset of taste cells responded to multiple stimuli with some of these studies concluding

these cells likely received input from neighboring taste cells (60-62). While we do not

rule out the possibility that taste cells receive input from other taste cells, our data show

that DR Type III cells do not require neighboring input to respond to bitter, sweet or

umami stimuli.

Our study has demonstrated that 1) Type II cells as defined by the expression of

the IP$_3$R3 are not the *only* taste cells in the bud that detect bitter, sweet and umami

stimuli; 2) taste cells with a PLCβ3 pathway also detect bitter, sweet and umami stimuli;

3) PLCβ3 is not expressed in Type II cells but is present in a subset of Type III cells and

another population of taste cells that lack VGCCs; and 4) loss of either IP$_3$R3 or PLCβ3

in taste cells causes similar deficits in taste-dependent c-Fos activity in the NTS as well

as deficits in taste driven behaviors. We conclude that peripheral taste transduction is

more complex than is currently appreciated.

EXPERIMENTAL PROCEDURES

Mice

Animals were cared for in compliance with the University at Buffalo Institutional Animal Care and Use Committee. Multiple mouse lines were used in this study. Both sexes were used and mice ranged in age from 1 to 6 months. Taste cells for qPCR analyses were collected from C57BL/6 mice. The IP$_3$R3-KO mouse was generated in a C57BL/6 background and was obtained from the Mutant Mouse Resources and Research Center (MMRRC:032884-JAX). This mouse has a targeted mutation in which exon 1 is replaced with a MAPT/GFP fusion (63) which results in the expression of GFP in place of a functional IP$_3$R3 receptor. Thus, these mice lack a functional IP$_3$R3 receptor and express GFP in cells that would otherwise have expressed IP$_3$R3. Heterozygous mice express both GFP and the IP$_3$R3 receptor. The PLCβ3-KO mouse was generously provided by Dr. Sang-Kyou Han (64). The mutation in these mice disrupts the catalytic domain of phospholipase C and was generated in a 129SV agouti mouse strain that was crossed with CD1 mice (65). These mice were maintained in this mixed background. Immunohistochemical analyses were also performed in the TRPM5-GFP mouse strain which expresses GFP in all taste cells that express TRPM5. These mice have been backcrossed into the C57BL/6 background and were used as another marker of Type II taste cells (16) to evaluate the expression patterns of PLCβ3. These mice were generously provided by Dr. Robert Margolskee.

Taste Receptor Cell Isolation

Taste receptor cells were harvested from CV, Fol and Fun papillae of adult mice as previously described (32, 66-71). Briefly, mice were sacrificed using carbon dioxide

and cervical dislocation. Tongues were removed and an enzyme solution containing

0.7mg/mL Collagenase B (Roche, Basel, Switzerland), 3mg/mL Dispase II (Roche), and

1mg/mL Trypsin Inhibitor (Sigma-Aldrich, St. Louis, MO) was injected beneath the

lingual epithelium. After the tongues were incubated in oxygenated Tyrode's solution

for approximately 17 min, the epithelium was peeled from the underlying muscle and

pinned serosal side up before it was incubated in Ca^{2+}-free Tyrode's for approximately

25 min. Cells were removed from taste papillae using capillary pipettes with gentle

suction and placed onto coverslips coated with Cell-Tak (Corning, Corning, NY).

Ca^{2+} Imaging

All measurements of intracellular calcium (Ca^{2+}) were performed in isolated taste

receptor cells that were no longer in contact with other taste cells. Cells were loaded for

20 minutes at room temperature (RT) with 2 µM Fura2-AM (Molecular Probes,

Invitrogen, Carlsbad, CA) containing Pluronic F-127 (Molecular Probes). Loaded cells

were then washed in Tyrode's solution under constant perfusion for 20min. Multiple

taste stimuli and high potassium (50mM KCl) solutions were individually applied and

Ca^{2+} responses were recorded. Cells were visualized using an Olympus IX73

microscope with a 40x oil immersion lens and images were captured with a Hamamatsu

ORCA-03G camera (Hamamatsu Photonics K.K., SZK Japan). Excitation wavelengths

of 340nm and 380nm were used with an emission wavelength of 510nm. Cells were

kept under constant perfusion using a gravity flow perfusion system (Automate

Scientific, San Francisco, CA). Images were collected every 4s using Imaging

Workbench 6.0 (Indec Biosystems, Santa Clara, CA). Experiments were graphed and

analyzed using Origin 9.2 software (OriginLab, Northhampton, MA).

Intracellular Ca^{2+} levels were measured as a ratio of fluorescence intensities.

Fluorescence values were calibrated using the Fura-2 Ca^{2+} Imaging Calibration kit

(Invitrogen). The effective dissociation constant K_d was calculated to be 180nM, which

was used in the following equation to calculate Ca^{2+} concentration:

$[Ca^{2+}] = K_d [(R-R_{min})/(R_{max}-R)](S_{f2}/S_{b2})$

$[Ca^{2+}] = 0.180 [(R-0.338)/(26.81-R)](22.19)$

R is the ratio value of fluorescence obtained after exciting cells at 340 and 380nm. Data

from cells were analyzed if the cell had a stable Ca^{2+} baseline within the range of 65nM

and 200nM. An evoked response was defined as measurable if the increase in

fluorescence was at least two standard deviations above baseline.

Immunohistochemistry

Mice were deeply anesthetized by intraperitoneal injections of sodium

pentobarbitol, 40 mg/kg (Patterson Veterinary, Mason, MI). Mice were then injected

intracardially with heparin (Sigma) and 1% sodium nitrate (Sigma) followed by perfusion

with approximately 30mL of 4% paraformaldehyde (Electron Microscopy Sciences, Ft.

Washington, PA) in 0.1M phosphate buffer (PB), pH 7.2. After perfusion, the tongues

were removed and placed in 4% paraformaldehyde/0.1M PB for 1-2h followed by a 4°C

overnight incubation in 20% sucrose/0.1M PB, pH 7.2. For some experiments, tongues

were immersion fixed overnight in 4% paraformaldehyde/0.1M PB, pH 7.2 at 4°C with

20% sucrose. Regardless of the fixation method, the next day, 40μm sections were cut

and washed in PBS 3X10 min at RT. For some experiments, antigen retrieval was

performed by placing sections in 10 mM sodium citrate, pH 8.5 at 80°C for 5 min. This

was done to disrupt the cross-bridges formed by fixation and expose antigen binding

sites.

Sections were incubated in blocking solution (0.3% Triton X-100, 1% normal goat

serum and 1% bovine serum albumin in 0.1M PB) for 1-2h at RT. Primary antibody was

added to the sections in blocking solution and incubated for 2 hours at RT followed by

overnight exposure to primary antibody at 4°C. Controls with no primary antibody were

included in each experiment. All primary antibodies were diluted in blocking solution.

Mouse anti-IP$_3$R3 (Transduction Labs, Lexington, KY) was used at 1:50 following

antigen retrieval. Rabbit anti-PLCβ3 (Abcam) was used at 1:200, rabbit anti-PLCβ2

(Santa Cruz Laboratories, Santa Cruz, CA) was used at 1:1000, rabbit anti-gustducin

(Santa Cruz) was used at 1:200 and anti-NTPDase2 (1:100(72). Mouse anti-SNAP-25

(Genway Biotech, San Diego, CA) was used at 1:200 following antigen retrieval.

Following overnight incubation in primary antibody, sections were washed in PBS 3X10

min at RT and then incubated with the appropriate secondary antibody (Cy5, 1:500;

Rhod, 1:250; Jackson ImmunoResearch Laboratories, West Grove, PA) at RT for 2h in

the dark. Controls were performed for double labeling experiments to ensure secondary

antibodies were not binding to primary antibodies raised in different organisms. After

secondary antibody incubation, sections were washed in PBS (3x10 min) and mounted

on Superfrost Plus slides (VWR, Radnor, PA) using Fluoromount G (Southern

Biotechnology Associates, Birmingham, AL) and coverslipped. All images were

obtained using a Zeiss LSM 710 Confocal Microscope (Zeiss, Oberkochen, Germany).

Stacks were collected using Zen software (Zeiss) and images were processed using

Adobe Photoshop CS5 software adjusting only brightness and contrast.

Real-Time PCR of Isolated Taste Cells

Taste receptor cells from CV, Fol or Fun papillae were isolated from the papillae as described above and then centrifuged for 20min at 13,000 RPM. RNA was purified using the NucleoSpin RNA XS Kit (Macherey-Nagel, Düren, Germany) according to kit instructions.   PCR analysis was performed for GAPDH to ensure sample quality and check for genomic contamination.   Contaminated samples were discarded and new samples were collected.   Real-Time PCR was performed using a BioRad MiniOpticon system (Bio-Rad Laboratories, Hercules, CA), with BioRad SYBR Green reagents (Bio-Rad Laboratories). Primers used for these experiments were: **PLCβ2**, *Fwd*: CAATTGAGGGGCAGCTGAGA *Rev*: TTCTAGGCTGCATCTGGGC; **PLCβ3**, *Fwd*: TCCTGGTGGTCAGGGAT *Rev*: CTGCCTGTCTCTGCTATCCG; **GAPDH** *Fwd*: ACAGTCAGCCGCATCTTCTT, *Rev*: ACGACCAAATCCGTTGACTC.

For real-time PCR analyses, each sample was run in triplicate. If there was more than 5% difference between the replicates, the data were discarded. Multiple biological repeats were used for each papillae type (CV, n=6; Fol, n=6; Fun, n=5). Data was normalized to GAPDH expression for each sample to correct for any loading differences and reported as fold differences.

Oral Infusions

Wild type, IP$_3$R3-KO and PLCβ3-KO mice (n=3 for each) were tested to measure the effects of quinine stimulation on the c-Fos immunoreactivity in the nucleus of the solitary tract (NTS). Water infusions (n=3) were performed to measure background responses. Surgical procedures were conducted using sterile technique while mice were anesthetized via 2% - 4% isoflurane anesthesia. Guided by a 21G needle, a small

length of PE-90 tubing was passed through the animal's cheek, emerging into the oral

cavity by the rear molars. The tubing was threaded through Teflon washers and heat

flared to secure on both the external and oral sides. Mice were given an analgesic

(0.5mg/ml of carprofen) and allowed to recover. After recovery, they were infused with

either quinine (5mM) or water into the oral catheter for 30 minutes (0.2ml/min, 30 min

infusion). Following the infusion, animals were returned to their cages and left

undisturbed for 45 minutes.

Brain Histology and Analysis

After the 45 minute post-infusion period, mice were perfused as described above

and the hindbrains were removed. The following day, hindbrains were sectioned into

40µm coronal sections which were washed in TBS (pH 7.5), 3x10 min each. Sections

were incubated in 1:100 H_2O_2 in TBS for 15 min, followed by 3X10 min TBS washes.

Sections were then incubated in anti-c-Fos (ab190289; Abcam, 1:1000) diluted in

blocking buffer (3% Normal Donkey Serum in TBS-TritonX) for 1 hr at RT. This was

followed by an overnight incubation at 4°C. The next day, sections were washed 3x10

min in TBS and then incubated for 2 hrs in anti-rabbit secondary antibody (711-065-152,

Jackson Immunoresearch, 1:1000) in blocking buffer. After incubation, sections were

kept in an avidin-biotin mixture (Elite kit; Vector Laboratories) in TBS-TritonX for 1 hr.

Tissue sections were washed (3x10 min in TBS) and then stained with DAB (3,3'-

diaminobenzidine-HCL; Vector Laboratories) for 5 min. Stained tissue sections were

washed (3x5 min in TBS) and mounted onto slides using Fluoromount-G (Southern

Biotechnology Associates). Sections were examined using a light microscope (10-100

X) equipped with a digital camera. Analyses of digital images were performed on the

intermediate rostral nucleus tractus solitarius (iRNTS; ~500 μm caudal to the dorsal

cochlear nucleus) which receives afferents from both the glossopharyngeal and chorda

tympani nerves and displays dense c-Fos-like immunoreactivity in response to intraoral

delivery of quinine (42, 73). A template tracing of the nucleus was made by an

experimenter blind to genotype and stimulus condition and this template was applied to

all sections. The number of FLI-positive neurons within the template was counted by

hand by an experimenter blind to genotype. Data were analyzed with a one way

ANOVA with follow up Student's t tests to identify individual differences. Significance

level was set at $p < 0.05$.

Analysis of Licking Behavior

Unconditioned licking responses to varying concentrations of taste stimuli were

recorded in a test chamber designed to measure brief-access licking (Davis MS80 Rig;

Dilog Instruments and Systems, Tallahassee, FL). This apparatus consisted of a

Plexiglas cage with a wire mesh floor. An opening at the front of the cage allowed

access to one of sixteen spill-proof glass drinking tubes that reside on a sliding platform.

A mechanical shutter opened and closed to allow the mouse access to one of the tubes

for a user-specified length of time. A computer controlled the movement of the platform,

order of tube presentation, opening and closing of the shutter, duration of tube access

and interval between tube presentations. Each individual lick was detected by a contact

lickometer and recorded on a computer via DavisPro collection software (Dilog

Instruments and Systems).

Mice were adapted to the test chamber and trained to drink from the sipper tubes

for 5 consecutive days as previously described (74, 75). During training, mice were 20-

h water deprived. On the first day of training, the mouse was presented with a single

stationary bottle of water for 30 min. On the second day, a tube containing water was

presented but this time the mouse was given 180s to initiate licking and once licking

was recorded the mouse was given 30s access to the tube. At the conclusion of either

the 30s access or the 180s limit, the shutter was closed again for 10s. Each of the 8

tubes, all containing water, was presented 3 times. During the remaining three days of

training, the mouse was given 30 min to initiate licking to one of eight tubes of water.

Once the mouse began licking, it was given 10s to lick before the shutter closed for 10s,

after which a new tube was presented.

During testing, animals were allowed to take as many trials as possible in 30 min.

Mice were tested on varying concentrations of sucrose (0,3,10,30,60,100,300,1000

mM), acesulfame K (0,0.5,2,6,8,16,20,32 mM), MSG with 10 μM amiloride

(0,25,50,75,100,400,600,800 mM), denatonium benzoate (0,0.1,0.3,1,3,5,10,20 mM),

and NaCl (0,3,10,30,60,100,300,1000 mM), in that order. Each stimulus was presented

in randomized blocks on Monday, Wednesday and Friday in a single week. Animals

were 22-h water deprived for all testing except sucrose, when animals were tested

water replete. Once the animal began licking the tube, they were allowed 10 seconds of

access before the shutter closed.

For stimuli tested in the water deprived condition (acesulfame K, MSG + amiloride,

denatonium benzoate, and NaCl), lick ratios were calculated by dividing the average

number of licks at each concentration by the average number of licks to water. For

stimuli tested while the animals were water replete (sucrose) licks scores were

calculated by subtracting the average number of licks at each concentration by the

average number of licks to water. These corrections are used to standardize for

individual differences in lick rate and are based on water need. Lick scores and licks

relative to water are compared by repeated measures ANOVA with genotype as

between factors variable and concentration as a repeated measures within factors

variable. Significant interaction terms were followed by Tukey's Honestly Significant

Difference tests. Statistical analyses were performed in Statistica.

Solutions

All chemicals were purchased from Sigma Chemical (Sigma-Aldrich, St. Louis,

MO) unless otherwise noted. Tyrode's solution contained 140mM NaCl, 5mM KCl, 3mM

$CaCl_2$, 1mM $MgCl_2$, 10mM HEPES, 10mM glucose, and 1mM pyruvate, pH 7.4. Ca^{2+}-

free Tyrode's contained 140mM NaCl, 5mM KCl, 2.7mM BAPTA, 2mM EGTA, 10mM

HEPES, 10mM glucose, 1mM pyruvate, pH 7.4. Nominal Ca^{2+}free Tyrode's contained

140mM NaCl, 5mM KCl, 10mM HEPES, 10mM glucose, 1mM pyruvate, pH 7.4. Hi KCl

solution contained 50mM KCl, 90mM NaCl, 3mM $CaCl_2$, 1mM $MgCl_2$, 10mM HEPES,

10mM glucose, 1mM pyruvate, pH 7.4. All taste-solutions, $2\mu M$ U73122 (Tocris, Bristol,

United Kingdom), and $2\mu M$ thapsigargin (Tocris) were prepared in normal Tyrode's.

Multiple taste stimuli were analyzed for bitter (5mM denatonium benzoate (Den), sweet

(20mM sucralose, 2mM saccharin, 50mM sucrose, 20mM Acesulfame K (Ace K)), and

umami (10mM monopotassium glutamate (MPG)). 50mM sucrose contained 90mM

NaCl instead of 140mM. Salt (250mM sodium chloride, NaCl) and sour (50mM citric

acid, CA, pH4) stimuli were tested using the same protocol as Lewandowski et al (19).

Statistics

Analysis of the taste responsiveness was performed using an interactive Chi-

square analysis with Yate's correction for continuity (76). Significant differences were

reported if $p<0.05$. For real-time PCR analyses, a one-way ANOVA with $p<0.05$ set as

the limit of significance was used to identify any significant differences between the

relative expression of PLCβ2 and PLCβ3 in the different papillae types. Comparisons

between two experimental conditions were made using a two-tailed Student's T test with

$p<0.05$ set as the limit of significance.

Colocalization analysis

Image stacks (n=5, 1 μm each) were acquired from labelled sections obtained

from three different animals and regions of interest (ROI) were drawn to include the

area inside clearly identifiable taste buds. Colocalization analysis was performed on the

ROIs using ImageJ Fiji software (NIH). Colocalization was determined based on

Pearson's coefficient using the Colocalization_Finder plugin. If the Pearson's coefficient

value was greater than 0.9, we considered it to be 100% colocalization due to the

variability in immunofluorescence intensity. If the Pearson's coefficient was less than

0.05, we considered this to be no colocalization. A Pearson's coefficient higher than

0.05 but lower than 0.9 was considered as partial colocalization.

ACKNOWLEDGMENTS
The authors thank Dr. Stefan Roberts at the University of Bristol for comments. This
work was supported by NSF1256950 to KFM. The authors thank Alan Siegel and the
UB North Campus Imaging facility funded by NSF-MRI DBI0923133 for the confocal
images. The authors thank Jhanna Flora for generating our model images.

AUTHOR CONTRIBUTIONS
DDB and EDB did live cell imaging, immunohistochemistry, data analysis and
contributed to writing the manuscript. DDB also did infusion experiments. LEM
performed brief-access lick behavior experiments, infusion experiments and data
analysis. KEK performed infusion experiments and data analysis. GCL conceived and

analyzed infusion experiments. ARN did immunohistochemistry, qPCR and data
analysis. ZCA and BRK did live cell imaging and data analysis. BTK did
immunohistochemistry. AV did experiments and analyzed data. AMT supervised brief-
access lick behavior, infusion experiments, analyzed and interpreted the data, and
edited the manuscript. KFM conceived and supervised the project, did live cell imaging,
data analysis and wrote the manuscript.
665
666   The authors declare no competing interests.
667
668

REFERENCES

1. Chaudhari N & Roper SD (2010) The cell biology of taste. *J Cell Biol* 190(3):285-296.
2. Finger TE (2005) Cell types and lineages in taste buds. *Chem Senses* 30 Suppl 1:i54-55.
3. Lawton DM, Furness DN, Lindemann B, & Hackney CM (2000) Localization of the glutamate-aspartate transporter, GLAST, in rat taste buds. *Eur J Neurosci* 12(9):3163-3171.
4. Miyoshi MA, Abe K, & Emori Y (2001) IP(3) receptor type 3 and PLCbeta2 are co-expressed with taste receptors T1R and T2R in rat taste bud cells. *Chem Senses* 26(3):259-265.
5. Zhang Y*, et al.* (2003) Coding of sweet, bitter, and umami tastes: different receptor cells sharing similar signaling pathways. *Cell* 112(3):293-301.
6. Yan W*, et al.* (2001) Bitter taste transduced by PLC-beta(2)-dependent rise in IP(3) and alpha-gustducin-dependent fall in cyclic nucleotides. *Am J Physiol Cell Physiol* 280(4):C742-751.
7. Dutta Banik D, Martin LE, Freichel M, Torregrossa AM, & Medler KF (2018) TRPM4 and TRPM5 are both required for normal signaling in taste receptor cells. *Proc Natl Acad Sci U S A* 115(4):E772-E781.
8. Taruno A*, et al.* (2013) CALHM1 ion channel mediates purinergic neurotransmission of sweet, bitter and umami tastes. *Nature* 495(7440):223-226.
9. Liu D & Liman ER (2003) Intracellular Ca2+ and the phospholipid PIP2 regulate the taste transduction ion channel TRPM5. *Proc Natl Acad Sci U S A* 100(25):15160-15165.
10. Zhang Z, Zhao Z, Margolskee R, & Liman E (2007) The transduction channel TRPM5 is gated by intracellular calcium in taste cells. *J Neurosci* 27(21):5777-5786.
11. Perez CA*, et al.* (2002) A transient receptor potential channel expressed in taste receptor cells. *Nat Neurosci* 5(11):1169-1176.
12. Clapp TR, Yang R, Stoick CL, Kinnamon SC, & Kinnamon JC (2004) Morphologic characterization of rat taste receptor cells that express components of the phospholipase C signaling pathway. *J Comp Neurol* 468(3):311-321.
13. Yee CL, Yang R, Bottger B, Finger TE, & Kinnamon JC (2001) "Type III" cells of rat taste buds: immunohistochemical and ultrastructural studies of neuron-specific enolase, protein gene product 9.5, and serotonin. *J Comp Neurol* 440(1):97-108.
14. Simon SA, de Araujo IE, Gutierrez R, & Nicolelis MA (2006) The neural mechanisms of gustation: a distributed processing code. *Nat Rev Neurosci* 7(11):890-901.
15. Roper SD (2007) Signal transduction and information processing in mammalian taste buds. *Pflugers Arch* 454(5):759-776.
16. Clapp TR, Medler KF, Damak S, Margolskee RF, & Kinnamon SC (2006) Mouse taste cells with G protein-coupled taste receptors lack voltage-gated calcium channels and SNAP-25. *BMC Biol* 4:7.

713   17.   Chang RB, Waters H, & Liman ER (2010) A proton current drives action
714   potentials in genetically identified sour taste cells. *Proc Natl Acad Sci U S A*
715   107(51):22320-22325.
716   18.   Ye W*, et al.* (2016) The K+ channel KIR2.1 functions in tandem with proton influx
717   to mediate sour taste transduction. *Proc Natl Acad Sci U S A* 113(2):E229-238.
718   19.   Lewandowski BC, Sukumaran SK, Margolskee RF, & Bachmanov AA (2016)
719   Amiloride-Insensitive Salt Taste Is Mediated by Two Populations of Type III Taste
720   Cells with Distinct Transduction Mechanisms. *J Neurosci* 36(6):1942-1953.
721   20.   Huang YA, Maruyama Y, Stimac R, & Roper SD (2008) Presynaptic (Type III)
722   cells in mouse taste buds sense sour (acid) taste. *J Physiol* 586(12):2903-2912.
723   21.   Huang AL*, et al.* (2006) The cells and logic for mammalian sour taste detection.
724   *Nature* 442(7105):934-938.
725   22.   Kataoka S*, et al.* (2008) The candidate sour taste receptor, PKD2L1, is
726   expressed by type III taste cells in the mouse. *Chem Senses* 33(3):243-254.
727   23.   Oka Y, Butnaru M, von Buchholtz L, Ryba NJ, & Zuker CS (2013) High salt
728   recruits aversive taste pathways. *Nature* 494(7438):472-475.
729   24.   Damak S*, et al.* (2006) Trpm5 null mice respond to bitter, sweet, and umami
730   compounds. *Chem Senses* 31(3):253-264.
731   25.   Hisatsune C*, et al.* (2007) Abnormal Taste Perception in Mice Lacking the Type 3
732   Inositol 1,4,5-Trisphosphate Receptor. *J Biol Chem* 282(51):37225-37231.
733   26.   Damak S*, et al.* (2003) Detection of sweet and umami taste in the absence of
734   taste receptor T1r3. *Science* 301(5634):850-853.
735   27.   Danilova V, Damak S, Margolskee RF, & Hellekant G (2006) Taste responses to
736   sweet stimuli in alpha-gustducin knockout and wild-type mice. *Chem Senses*
737   31(6):573-580.
738   28.   Dotson CD, Roper SD, & Spector AC (2005) PLCbeta2-independent behavioral
739   avoidance of prototypical bitter-tasting ligands. *Chem Senses* 30(7):593-600.
740   29.   Matsumoto I, Ohmoto M, Narukawa M, Yoshihara Y, & Abe K (2011) Skn-1a
741   (Pou2f3) specifies taste receptor cell lineage. *Nat Neurosci* 14(6):685-687.
742   30.   Yee KK, Sukumaran SK, Kotha R, Gilbertson TA, & Margolskee RF (2011)
743   Glucose transporters and ATP-gated K+ (KATP) metabolic sensors are present
744   in type 1 taste receptor 3 (T1r3)-expressing taste cells. *Proc Natl Acad Sci U S A*
745   108(13):5431-5436.
746   31.   Maruyama Y, Pereira E, Margolskee RF, Chaudhari N, & Roper SD (2006)
747   Umami responses in mouse taste cells indicate more than one receptor. *J*
748   *Neurosci* 26(8):2227-2234.
749   32.   Hacker K, Laskowski A, Feng L, Restrepo D, & Medler K (2008) Evidence for two
750   populations of bitter responsive taste cells in mice. *J Neurophysiol* 99(3):1503-
751   1514.
752   33.   Clapp TR, Stone LM, Margolskee RF, & Kinnamon SC (2001)
753   Immunocytochemical evidence for co-expression of Type III IP3 receptor with
754   signaling components of bitter taste transduction. *BMC Neurosci* 2:6.
755   34.   DeFazio RA*, et al.* (2006) Separate populations of receptor cells and presynaptic
756   cells in mouse taste buds. *J Neurosci* 26(15):3971-3980.

35. Shandilya J, Gao Y, Nayak TK, Roberts SG, & Medler KF (2016) AP1 transcription factors are required to maintain the peripheral taste system. *Cell Death Dis* 7(10):e2433.

36. Medler KF, Margolskee RF, & Kinnamon SC (2003) Electrophysiological characterization of voltage-gated currents in defined taste cell types of mice. *J Neurosci* 23(7):2608-2617.

37. Sukumaran SK*, et al.* (2017) Whole transcriptome profiling of taste bud cells. *Scientific reports* 7(1):7595.

38. Richter TA, Caicedo A, & Roper SD (2003) Sour taste stimuli evoke Ca2+ and pH responses in mouse taste cells. *J Physiol* 547(Pt 2):475-483.

39. Courtney E Wilson TF, Sue C Kinnamon (2017) Type III cells in anterior taste fields are more immunohistochemically diverse than those of posterior taste fields in mice. *Chemical Senses* bjx055.

40. Yang R, Crowley HH, Rock ME, & Kinnamon JC (2000) Taste cells with synapses in rat circumvallate papillae display SNAP-25-like immunoreactivity. *J Comp Neurol* 424(2):205-215.

41. King CT, Garcea M, & Spector AC (2000) Glossopharyngeal nerve regeneration is essential for the complete recovery of quinine-stimulated oromotor rejection behaviors and central patterns of neuronal activity in the nucleus of the solitary tract in the rat. *J Neurosci* 20(22):8426-8434.

42. King CT, Travers SP, Rowland NE, Garcea M, & Spector AC (1999) Glossopharyngeal nerve transection eliminates quinine-stimulated fos-like immunoreactivity in the nucleus of the solitary tract: implications for a functional topography of gustatory nerve input in rats. *J Neurosci* 19(8):3107-3121.

43. Stratford JM & Finger TE (2011) Central representation of postingestive chemosensory cues in mice that lack the ability to taste. *J Neurosci* 31(25):9101-9110.

44. Stratford JM & Thompson JA (2016) MSG-Evoked c-Fos Activity in the Nucleus of the Solitary Tract Is Dependent upon Fluid Delivery and Stimulation Parameters. *Chem Senses* 41(3):211-220.

45. Stratford JM, Thompson JA, & Finger TE (2017) Immunocytochemical organization and sour taste activation in the rostral nucleus of the solitary tract of mice. *J Comp Neurol* 525(2):271-290.

46. Ruiz-Avila L, Wong GT, Damak S, & Margolskee RF (2001) Dominant loss of responsiveness to sweet and bitter compounds caused by a single mutation in alpha -gustducin. *Proc Natl Acad Sci U S A* 98(15):8868-8873.

47. Zukerman S, Glendinning JI, Margolskee RF, & Sclafani A (2009) T1R3 taste receptor is critical for sucrose but not Polycose taste. *Am J Physiol Regul Integr Comp Physiol* 296(4):R866-876.

48. Treesukosol Y, Smith KR, & Spector AC (2011) Behavioral evidence for a glucose polymer taste receptor that is independent of the T1R2+3 heterodimer in a mouse model. *J Neurosci* 31(38):13527-13534.

49. Treesukosol Y & Spector AC (2012) Orosensory detection of sucrose, maltose, and glucose is severely impaired in mice lacking T1R2 or T1R3, but Polycose sensitivity remains relatively normal. *Am J Physiol Regul Integr Comp Physiol* 303(2):R218-235.

803   50.   Dvoryanchikov G*, et al.* (2017) Transcriptomes and neurotransmitter profiles of
804   classes of gustatory and somatosensory neurons in the geniculate ganglion. *Nat
805   Commun* 8(1):760.
806   51.   Treesukosol Y, Blonde GD, & Spector AC (2009) T1R2 and T1R3 subunits are
807   individually unnecessary for normal affective licking responses to Polycose:
808   implications for saccharide taste receptors in mice. *Am J Physiol Regul Integr
809   Comp Physiol* 296(4):R855-865.
810   52.   Chaudhari N, Landin AM, & Roper SD (2000) A metabotropic glutamate receptor
811   variant functions as a taste receptor. *Nat Neurosci* 3(2):113-119.
812   53.   Chaudhari N, Maruyama Y, Roper S, & Trubey K (2005) Multiple pathways for
813   signaling glutamate taste in rodents. *Chem Senses* 30 Suppl 1:i29-i30.
814   54.   Chaudhari N, Pereira E, & Roper SD (2009) Taste receptors for umami: the case
815   for multiple receptors. *Am J Clin Nutr* 90(3):738S-742S.
816   55.   Delay ER, Hernandez NP, Bromley K, & Margolskee RF (2006) Sucrose and
817   monosodium glutamate taste thresholds and discrimination ability of T1R3
818   knockout mice. *Chem Senses* 31(4):351-357.
819   56.   Eddy MC, Eschle BK, & Delay ER (2017) Comparison of the Tastes of L-Alanine
820   and Monosodium Glutamate in C57BL/6J Wild Type and T1r3 Knockout Mice.
821   *Chem Senses* 42(7):563-573.
822   57.   Pal Choudhuri S, Delay RJ, & Delay ER (2015) L-Amino Acids Elicit Diverse
823   Response Patterns in Taste Sensory Cells: A Role for Multiple Receptors. *PLoS
824   One* 10(6):e0130088.
825   58.   Pal Choudhuri S, Delay RJ, & Delay ER (2016) Metabotropic glutamate receptors
826   are involved in the detection of IMP and L-amino acids by mouse taste sensory
827   cells. *Neuroscience* 316:94-108.
828   59.   Stone LM, Barrows J, Finger TE, & Kinnamon SC (2007) Expression of T1Rs
829   and gustducin in palatal taste buds of mice. *Chem Senses* 32(3):255-262.
830   60.   Tomchik SM, Berg S, Kim JW, Chaudhari N, & Roper SD (2007) Breadth of
831   tuning and taste coding in mammalian taste buds. *J Neurosci* 27(40):10840-
832   10848.
833   61.   Gilbertson TA, Boughter JD, Jr., Zhang H, & Smith DV (2001) Distribution of
834   gustatory sensitivities in rat taste cells: whole-cell responses to apical chemical
835   stimulation. *J Neurosci* 21(13):4931-4941.
836   62.   Caicedo A, Kim KN, & Roper SD (2002) Individual mouse taste cells respond to
837   multiple chemical stimuli. *J Physiol* 544(Pt 2):501-509.
838   63.   Hegg CC, Jia C, Chick WS, Restrepo D, & Hansen A (2010) Microvillous cells
839   expressing IP3 receptor type 3 in the olfactory epithelium of mice. *Eur J Neurosci*
840   32(10):1632-1645.
841   64.   Han SK, Mancino V, & Simon MI (2006) Phospholipase Cbeta 3 mediates the
842   scratching response activated by the histamine H1 receptor on C-fiber
843   nociceptive neurons. *Neuron* 52(4):691-703.
844   65.   Xie W*, et al.* (1999) Genetic alteration of phospholipase C beta3 expression
845   modulates behavioral and cellular responses to mu opioids. *Proc Natl Acad Sci U
846   S A* 96(18):10385-10390.

66. Hacker K & Medler KF (2008) Mitochondrial calcium buffering contributes to the
maintenance of Basal calcium levels in mouse taste cells. *J Neurophysiol*
100(4):2177-2191.
67. Laskowski AI & Medler KF (2009) Sodium-calcium exchangers contribute to the
regulation of cytosolic calcium levels in mouse taste cells. *J Physiol* 587(Pt
16):4077-4089.
68. Maliphol AB, Garth DJ, & Medler KF (2013) Diet-induced obesity reduces the
responsiveness of the peripheral taste receptor cells. *PLoS One* 8(11):e79403.
69. Rebello MR, Maliphol AB, & Medler KF (2013) Ryanodine Receptors Selectively
Interact with L Type Calcium Channels in Mouse Taste Cells. *PLoS One*
8(6):e68174.
70. Rebello MR & Medler KF (2010) Ryanodine receptors selectively contribute to
the formation of taste-evoked calcium signals in mouse taste cells. *Eur J*
Neurosci 32(11):1825-1835.
71. Szebenyi SA, Laskowski AI, & Medler KF (2010) Sodium/calcium exchangers
selectively regulate calcium signaling in mouse taste receptor cells. *J*
Neurophysiol 104(1):529-538.
72. Bartel DL, Sullivan SL, Lavoie EG, Sevigny J, & Finger TE (2006) Nucleoside
triphosphate diphosphohydrolase-2 is the ecto-ATPase of type I cells in taste
buds. *J Comp Neurol* 497(1):1-12.
73. Harrer MI & Travers SP (1996) Topographic organization of Fos-like
immunoreactivity in the rostral nucleus of the solitary tract evoked by gustatory
stimulation with sucrose and quinine. *Brain Res* 711(1-2):125-137.
74. Martin LE*, et al.* (2017) Salivary proteins alter taste-guided behaviors and taste
nerve signaling in rat. *Physiol Behav* 184:150-161.
75. Torregrossa AM*, et al.* (2014) Induction of salivary proteins modifies measures of
both orosensory and postingestive feedback during exposure to a tannic acid
diet. *PLoS One* 9(8):e105232.
76. Preacher KJ (2001) Calculation for the chi-square test: An interactive calculation
tool for chi-square tests of goodness of fit and independence [Computer
software]. . p Available from http://quantpsy.org.
878

 FIGURE LEGENDS

Figure 1. Current taste cell models and experimental setup. A) Current models of

the signal transduction pathways in Type II cells (left) that respond to bitter, sweet, and

umami stimuli as well as Type III cells (right) that respond to sour and salty stimuli. B)

Example of an isolated taste cell that was stimulated with MPG (10mM) and KCl

(50mM). The DIC image is shown first, followed by image of Fura 2-AM loading. The

next three images show the ratio (340/380) at different time points as the cell is

stimulated. The corresponding calcium (Ca^{2+}) changes are shown in 1C.

Figure 2. Loss of IP$_3$R3 causes a reduction, but not abolishment, of taste cell

sensitivity to different taste stimuli. Chi square analysis with Yate's correction for

continuity was used to compare the frequency of evoked Ca^{2+} responses to different

taste stimuli between wild type (WT) and IP$_3$R3-KO (KO) mice. A) Table of the mouse

line, number of mice tested, number of cells tested and number of responses recorded.

The P value for each comparison between WT and IP$_3$R3-GFP-KO is shown as well as

the percent response value. These data include all cells from IP$_3$R3-GFP-KO mice,

both GFP+ and GFP-. B) Graph shows the percentage of overall responsiveness to

stimuli for taste cells from the CV, Fol and Fun papillae in WT and IP$_3$R3-KO. The

responsiveness to multiple stimuli was evaluated and is shown for CV (C), FOL (D) and

FUN (E), Specific numbers and P values for each are shown in Figure S3. (*, p<0.05; **,

p<0.01; ***, p<0.001).

Figure 3. Taste-evoked Ca^{2+} release in IP$_3$R3-KO mice is dependent upon PLC

activity and Ca^{2+} release from internal stores. Open columns represent the time that

the taste stimulus is presented. The application of Ca^{2+} free Tyrode's is indicated by the

dashed lines. The stimulus presented during this time is also in Ca^{2+} free Tyrode's.

The gray hatched columns represent the application of either thapsigargin or U73122,

both of which are irreversible inhibitors. A) Bitter-evoked taste responses (5mM Den)

persist in the absence of extracellular calcium (Ca^{2+}-free) and are abolished by the

SERCA pump inhibitor thapsigargin (B) as well as the PLC blocker U73122 (C). D)

Responses to sweet stimuli (20mM sucrose) persist in Ca^{2+}-free and are abolished by

thapsigargin (E) and U73122 (F). G) Umami stimuli (10mM MPG) persist in Ca^{2+}-free

and were abolished by thapsigargin (H) and U73122 (I). Individual data points are

shown in Figure S4.

Figure 4. Expression of PLCβ3 in taste cells. Laser scanning confocal micrographs

(LSCMs, stack of 5 slices, 1μm each) of PLCβ3 immunostaining in the IP$_3$R3-KO-GFP

mice reveals that PLCβ3 is expressed in a separate population from the GFP positive

taste cells in the CV (A, n=5), Fol (B, n=3) and Fun papillae (C, n=4). Scale bar=20μM.

D) Co-localization analysis is shown.

Figure 5. Some Type III cells respond to bitter, sweet and umami stimuli.

A-C) Representative traces of taste cells that responded to bitter, sweet and/or umami

stimuli and Hi KCl (50mM) depolarization with Ca^{2+} signals. Some cells responded to 1

stimulus while others responded to multiple stimuli. D) Representative trace of taste

cells from IP$_3$R3-KO mice that responded to 50mM citric acid (CA), 50mM KCl, bitter

(5mM denatonium), sweet (20mM sucralose) and umami (10mM monopotassium

glutamate, MPG) stimuli. E) Representative trace of a separate subset of cells that

responded to 250mM NaCl, 50mM citric acid (CA) and 50mM KCl but were not sensitive

to the bitter, sweet, umami stimuli tested. F) Summary of taste cells from IP$_3$R3-KO

mice that responded to 50mM KCl with a Ca^{2+} signal (n=20), 25% only responded to

CA, 35% responded to CA and NaCl, and 40% responded to CA and bitter, sweet

and/or umami stimuli (B,U,S) while no cells responded to all the taste stimuli. G) The

overall responsiveness of taste cells from the CV, FoI and Fun papillae of the IP$_3$R3-KO

mice and WT mice was measured. This is defined as the percentage of stimulated cells

that responded to at least one stimulus. The percentage of DR cells in both lines was

also measured using the criteria that the cell responded to at least one stimulus AND

50mM KCl with Ca^{2+} signals. The overall responsiveness to taste stimuli was

significantly reduced in the IP$_3$R3-KO mice compared to controls (**,p<0.01) while the

number of responsive DR cells did not change in the absence of IP$_3$R3. H) The

percentage of responsiveness in the DR taste cells to different taste stimuli in the WT

and IP$_3$R3-KO mice. Chi-square analysis found no significant differences.

Figure 6. PLCβ3 is required for the detection of bitter, sweet and umami stimuli.

Chi square analysis with Yate's correction for continuity was used to compare the

frequency of evoked Ca^{2+} responses to different taste stimuli between wild type (WT)

and PLCβ3-KO (KO) mice. A) Table of the mouse line, number of mice tested, number

of cells tested and number of responses recorded. The P value for each comparison

between WT and PLCβ3-KO is shown as well as the percent response value. B) The

overall responsiveness and the percentage of DR taste cells from the CV, Fol and Fun

papillae of the PLCβ3-KO mice and WT mice was measured. Both the overall

responsiveness to taste stimuli and the number of DR cells were significantly reduced in

the PLCβ3-KO mice compared to controls. The responsiveness to multiple stimuli was

evaluated and is shown for CV (C), FOL (D) and FUN (E). Specific numbers and P

values for each are shown in Figure S8. (*, $p<0.05$; **, $p<0.01$; ***, $p<0.001$).

Figure 7. Loss of PLCβ3 or IP$_3$R3 causes deficits in the taste-evoked neural

activity in the nucleus of solitary tract.

Oral infusion of bitter (quinine) and water elicited c-Fos immunoreactivity in the

intermediate rostral nucleus tractus solitarius (IRNTS). (A, B) Loss of either PLCβ3

(PLCβ3-KO) or IP$_3$R3 (IP$_3$R3-KO) caused a significant reduction in the c-Fos

immunoreactivity in the IRNTS compared to WT mice. The residual c-Fos labeling in

each of the KO mice was comparable to the c-Fos labelling generated by water infusion.

For all experiments, 3 mice of each genotype were used (***, $p<0.001$; **, $p<0.01$). The

red outlined area denotes the analyzed region for each. Data were analyzed with a one

way ANOVA with follow up Student's t tests to identify individual differences.

Significance level was set at $p<0.05$. C) A schematic diagram showing the analysis

region in the hindbrain that was analyzed.

Figure 8. Loss of PLCβ3 or IP$_3$R3 affects behavioral responses to taste stimuli.

Lick data (±standard deviation) from brief-access behavioral tests compare the

responses of IP$_3$R3-KO (red line) and PLCβ3-KO (pink line) to WT (IP$_3$R3-WT, black

line; PLCβ3-WT, blue line). A) Lick ratios (stimulus/water) of the WT mice for the bitter

stimulus denatonium (0, 0.1, 0.3, 1, 3, 5, 10, 20mM) were significantly different from the

IP$_3$R3-KO and PLCβ3-KO responses. B) Lick ratios of the WT mice for the umami

stimulus monosodium glutamate + 10μM amiloride (MSG, 0, 25, 50, 75, 100, 400, 600,

800mM) were significantly different from the IP$_3$R3-KO and PLCβ3-KO mice. C) Lick

ratios of the WT mice for the artificial sweetener acesulfame K (0, 0.5, 2, 6, 8, 16, 20,

32mM) were significantly different from the IP$_3$R3-KO and PLCβ3-KO mice. D) Lick

scores (stimulus-water) of the WT mice for sucrose (0, 3, 10, 30, 60,100, 300, 1000mM)

were significantly different from the IP$_3$R3-KO and PLCβ3-KO responses. E) No

significant differences were detected between the WT and KO mice for any

concentration of NaCl (0, 3, 10, 30, 60, 100, 300, 1000mM) tested. For all experiments,

5 mice of each genotype were used (***, $p < 0.001$; **, $p < 0.01$; *, $p < 0.05$). Data were

compared by repeated measures ANOVA. Significant interaction terms were followed

by Tukey's Honestly Significant Difference tests.

A

B

DIC   Fura 2   1   2   3

C

FIGURE 1

A

Papillae type	Mouse genotype	Cells tested	# of Resp.	P value	% value
All cells	WT (n=105 mice)	407	259	P<0.0001	63.6
	KO (n=115 mice)	690	213		30.9
CV	WT (n=63 mice)	250	159	P=0.0003	63.6
	KO (n=70 mice)	443	172		38.8
FOL	WT (n=28 mice)	53	29	P=0.44	54.7
	KO (n=37 mice)	123	52		42.3
FUN	WT (n=32 mice)	104	71	P=0.17	68.2
	KO (n=30 mice)	124	61		49.2

FIGURE 2

FIGURE 3

A

CV   IP₃R3-KO   PLCβ3

B

FOL   IP₃R3-KO   PLCβ3

C

FUN

D

Comparison	Average overlap, standard deviation
GFP vs anti-PLCβ3	0.03±0.02

FIGURE 4

FIGURE 5

A

Papillae type	Mouse genotype	Cells tested	# of Resp.	P value	% value
All cells	WT (n=53 mice)	577	344	P=0.002	59.6
	KO (n=68 mice)	375	155		41.3
CV	WT (n=40 mice)	221	137	P=0.246	61.9
	KO (n=55 mice)	138	68		49.2
FOL	WT (n=40 mice)	243	136	P=0.039	55.9
	KO (n=34 mice)	120	43		35.8
FUN	WT (n=35 mice)	113	71	P=0.035	62.8
	KO (n=48 mice)	117	44		37.6

FIGURE 6

FIGURE 7

FIGURE 8

Cell populations	Identifiers/Classification
IP$_3$R3-KO-GFP cells	Type II cells with GFP expression replacing IP$_3$R3 expression.
WT cells	All cells that were tested and/or responded to stimuli, including Type II cells, Type III cells, other cells.
IP$_3$R3-KO cells	Taste cells from mice that lack functional Type II cells and do not express GFP. These include Type III cells and other cells.
Dual-responsive cells	Taste cells that respond to 50mM KCl (indicating the presence of VGCC) and bitter, sweet and/or umami stimuli.
Taste-only cells	Taste cells from the IP$_3$R3-KO mouse that lack GFP expression (are not Type II cells) and do not respond to 50mM KCl (are not Type III cells).

Table S1. Definition of taste cell populations analyzed.

Figure S1. Characterization of the IP₃R3-KO mice. A) LSCMs (stack of 5 slices, 1µm each) from WT mice identified anti-IP₃R3 labeling in taste receptor cells from the CV (n=3). B) IP₃R3-KO mice express GFP in lieu of IP₃R3 and were not labeled by anti-IP₃R3 (n=6). C) LSCMs of the IP₃R3-het mouse identified strong co-localization between anti-IP₃R3 labeling and GFP expression. D) Anti-PLCβ2 labeling in IP₃R3-KO mice found that PLCβ2 co-localizes with GFP, indicating that IP₃R3-KO-GFP is specific to Type II cells (LSCMs: stack of 5 slices, 1µm each; n=4). E) α-gustducin is present in a subset of the IP₃R3-KO-GFP taste cells in the CV (LSCMs: stack of 10 slices, 1µm each; n=3). Asterisks identify some GFP expressing cells that do not express gustducin. Scale bars = 20 µm. F) Co-localization analysis identified the average (± standard deviation) overlapping expression for each target protein with the GFP expression, n=3 for each.

Figure S2. Taste-evoked Ca²⁺ signals are specific to the stimulus.

A concentration gradient analysis of (A) a bitter stimulus (denatonium), (B) an umami stimulus (MPG) and (C) a sweet stimulus (saccharin) demonstrates that DR cells respond to a wide range of stimulus concentrations. The GFP+ cell (lacks IP_3R3) in C did not respond to any concentration of saccharin applied. (D-E) Identical concentrations of the compound cellobiose did not generate a Ca²⁺ signal in DR cells, indicating that the measured responses are taste-activated and not due to non-specific osmotic effects.

IP$_3$R3

CV

CV Stimulus	MS	Cells tested	# of Resp.	P value
DEN	WT	238	102	0.06
	KO	341	107	
SUC	WT	72	16	0.0004
	KO	87	1	
SAC	WT	106	23	0.005
	KO	174	13	
SUCRAL	WT	121	47	p<0.001
	KO	137	10	
MPG	WT	81	13	0.82
	KO	301	42	
HI K	WT	245	75	0.68
	KO	384	108	

FOL

FOL Stimulus	MS	Cells tested	# of Resp.	P value
DEN	WT	46	18	0.93
	KO	100	38	
SAC	WT	15	1	0.69
	KO	50	4	
SUCRAL	WT	23	6	0.08
	KO	30	1	
MPG	WT	18	4	0.48
	KO	73	8	
HI K	WT	53	14	0.93
	KO	113	31	

FUN

FUN Stimulus	MS	Cells tested	# of Resp.	P value
DEN	WT	76	42	0.89
	KO	59	30	
SUC	WT	41	10	0.89
	KO	19	6	
SAC	WT	65	23	0.73
	KO	55	16	
SUCRAL	WT	85	25	0.04
	KO	99	13	
MPG	WT	75	17	0.4
	KO	81	12	
HI K	WT	104	33	0.7
	KO	103	28	

Figure S3. Tables of the mouse line, number of mice tested, number of cells tested and number of responses recorded from the CV, FOL, and FUN papillae of the WT and IP$_3$R3-KO mice data shown in Figure 2. Chi-square analysis with Yate's correction for continuity was used to identify significant differences in the response frequency between WT and KO mice for each stimulus in each papillae.

Figure S4. Taste-evoked Ca^{2+} release in IP$_3$R3-KO mice is dependent upon PLC activity and Ca^{2+} release from internal stores. Individual data points of the data presented in Figure 3. A) Bitter-evoked taste responses (5mM Den, n=6) persist in the absence of extracellular calcium (Ca^{2+}-free) and are abolished by the SERCA pump inhibitor thapsigargin (B, n=7) as well as the PLC blocker U73122 (C, n=7). D) Responses to sweet stimuli (50mM sucrose, n=6) persist in Ca^{2+}-free and are abolished by thapsigargin (E, n=4) and U73122 (F, n=6). G) Umami stimuli (10mM MPG) persist in Ca^{2+}-free (n=7) and were abolished by thapsigargin (H, n=7) and U73122 (I, n=4). A two-tailed Student's t test was used to identify significant differences for each experimental condition.

Figure S5. Expression of PLCβ3 in taste papillae. A) Anti-PLCβ3 labeling in the CV of TRPM5-GFP mice determined that PLCβ3 is expressed in taste cells lacking GFP expression (LSCMs: stack of 5 slices, 1μm each; n=4). B) Co-labeling with anti-PLCβ2 and anti-PLCβ3 in the CV of C57BL/6 mice revealed that these PLCβs are expressed in separate taste cell populations (LSCMs: stack of 5 slices, 1μm each; n=3). C) Co-labeling with anti-NTDPase2 and anti-PLCβ3 in the CV of C57BL/6 mice determined that these markers are expressed in separate taste cell populations (LSCMs: stack of 5 slices, 1μm each; n=3). Scale bar = 10μM. D) Co-localization analysis identified the average (± standard deviation) overlapping expression for PLCβ3 with TRPM5-GFP or anti-PLCβ2 expression, n=3 for each. mRNA was isolated from taste cells originating in the different papillae types from C57BL/6 mice. Taste cells were analyzed from at least five different mice for each. Values were normalized to GAPDH expression and are presented as a ratio to values from the CV papillae for (E) PLCβ2 and (F) PLCβ3. (***, p<0.001). A one-way ANOVA with p<0.05 set as the limit of significance was used to identify any differences in the relative mRNA levels between papillae types.

Figure S6. Dual responsive cells are present in multiple mouse lines. Control imaging experiments were performed using GAD67-GFP mice and C57BL/6 mice. GAD67-GFP is expressed in a large subset of Type III mouse taste cells (34). A-B) Representative traces of DR taste cells that responded to bitter, sweet and/or umami stimuli and Hi KCl (50mM) in GAD67-GFP mice. DR taste cells were present in both GAD67-GFP + (A) and -GFP- (B) taste cells. C) Experiments in C57BL/6 mice also identified the presence of DR taste cells.

Figure S7. Stimulus evoked Ca^{2+} responses in taste cells in PLCβ3-KO mice. A) Representative trace of taste cells from PLCβ3-KO mice that responded to 50mM citric acid (CA) and KCl (50mM) but did not respond to bitter (5mM denatonium), sweet (20mM sucralose) or umami (10mM monopotassium glutamate, MPG) stimuli. B) Representative trace of a separate subset of cells that responded to 250mM NaCl, CA and KCl but were not sensitive to the bitter, sweet, umami stimuli tested. C) Summary of responsive taste cells from PLCβ3-KO mice (n=38): CA and KCl (n=18); CA, NaCl and KCl (n=3); CA, KCl, and bitter, sweet and/or umami stimuli (B,U,S) (n=0); and B,U,S only (n=17). Representative taste-evoked responses for bitter (D), sweet (E) and umami (F) in the PLCβ3-KO mice. None of these cells responded to CA or KCl.

PLCβ3

CV

CV Stimulus	MS	Cells tested	# of Resp.	P value
DEN	WT	221	86	0.018
	KO	138	30	
SUC	WT	191	36	0.039
	KO	112	9	
SAC	WT	198	22	0.02
	KO	130	4	
ACEK	WT	181	27	0.04
	KO	111	6	
MPG	WT	182	32	0.031
	KO	116	8	
HI K	WT	221	82	0.06
	KO	138	32	
DR	WT	221	59	0.0008
	KO	138	12	

FOL

FOL Stimulus	MS	Cells tested	# of Resp.	P value
DEN	WT	233	85	.144
	KO	113	28	
SUC	WT	173	21	0.97
	KO	65	7	
SAC	WT	156	27	0.04
	KO	86	5	
ACEK	WT	163	28	0.04
	KO	76	4	
MPG	WT	179	34	0.21
	KO	67	3	
HI K	WT	243	68	0.08
	KO	120	20	
DR	WT	243	45	0.002
	KO	120	5	

FUN

FUN Stimulus	MS	Cells tested	# of Resp.	P value
DEN	WT	112	52	0.003
	KO	117	23	
SUC	WT	86	17	0.91
	KO	107	19	
SAC	WT	93	22	0.015
	KO	111	10	
ACEK	WT	84	12	0.015
	KO	108	3	
MPG	WT	81	10	0.016
	KO	107	2	
HI K	WT	113	15	0.053
	KO	116	5	
DR	WT	113	13	0.001
	KO	116	0	

Figure S8. Tables of the mouse line, number of mice tested, number of cells tested and number of responses recorded from the CV, FOL, and FUN papillae of the WT and PLCβ3-KO mice data shown in Figure 6. Chi-square analysis with Yate's correction for continuity was used to identify significant differences in the response frequency between WT and KO mice for each stimulus in each papillae.

E

Comparison	Average overlap, standard deviation
anti-PLCβ3 vs anti-SNAP25	0.62±0.03

Figure S9. PLCβ3 is expressed in some Type III taste cells. Immunohistochemical analyses (LSCMs: stack of 5 slices, 1μm each) using anti-PLCβ3 and anti-SNAP25 revealed some co-localization between PLCβ3 and SNAP25, confirming that PLCβ3 is expressed in some Type III cells but is not restricted in its expression to this taste cell type (A, CV papillae, n=3; B, Fun papillae, n=3) Scale bar=20μM. Labeling with anti-SNAP25 in the CV of PLCβ3-WT (**C**) and PLCβ3-KO (**D**) mice revealed no differences in the expression level of this synaptic marker (n=5 mice for each). Scale bars = 50 μm. E) Co-localization analysis identified the average (± standard deviation) overlapping expression for PLCβ3 with anti-SNAP25 expression, n=3.

	DIC	IP$_3$R3-GFP	Anti-PLCβ3

Figure S10. Neither IP$_3$R3 nor PLCβ3 is expressed in IRNTS.

CV papillae and IRNTS sections were evaluated in parallel. A) Analyses of the brain sections from the IP$_3$R3-KO mice revealed no GFP labelling in the IRNTS. Immunohistochemical analysis using anti-PLCβ3 also did not detect any PLCβ3 expression. Scale bar=50μm. B) LSCM analyses (n=5 sections, 1 μM each) of the tongues from the same mice identified the expression of IP$_3$R3-KO-GFP and PLCβ3 in the taste receptor cells from the CV papillae (n=3). Scale bar=10μm.

Figure S11. Analyses of the different taste cell populations that responded to taste stimuli. Graphs show the percent of the responses for each taste quality tested. A-C) Response profiles for bitter, sweet and umami stimuli from the DR taste cells in WT (n=29 mice, 62 cells) and IP$_3$R3-KO (n=18 mice, 31 cells) mice. Taste cells that were stimulated with bitter, sweet and umami stimuli as well as 50mM KCl and responded to at least one of the taste stimuli and KCl were analyzed. The relative percentage of DR taste cells that responded to 1, 2, or 3 taste qualities is shown for each taste quality analyzed. Chi-square analysis with Yate's correction for continuity found no significant differences between the WT and IP$_3$R3-KO mice for bitter (A), sweet (B), or umami (C) response profiles. The same analysis was applied to taste cells that responded to at least one bitter, sweet or umami stimulus but did not respond to 50mM KCl for WT (n=37 mice, 85 cells) and IP$_3$R3-KO (n=33 mice, 78 cells) mice. These data are shown for bitter (D), sweet (E), and umami (F). No significant differences were found between WT and IP$_3$R3-KO mice.