Morphology does not covary with predicted behavioural correlations of the

domestication syndrome in dogs

Christina Hansen Wheat[1*], Wouter van der Bijl[1,2] & Christopher West Wheat[1]

[1]*Department of Zoology, Stockholm University, SE-10961 Stockholm, Sweden*

[2]*Department of Zoology & Biodiversity Research Centre, University of British Columbia, Vancouver BC,*

Canada.

Corresponding author: christina.hansen@zoologi.su.se

Abstract

Domesticated animals display suites of altered morphological, behavioural and physiological

traits compared to their wild ancestors, a phenomenon known as the domestication syndrome

(DS). Because these alterations are observed to co-occur across a wide range of present day

domesticates, the traits within the DS are assumed to covary within species and a single

developmental mechanism has been hypothesized to cause the observed co-occurrence.

However, due to the lack of formal testing it is currently not well-resolved if the traits within

DS actually covary. Here we test the hypothesis that the presence of the classic morphological

domestication traits white pigmentation, floppy ears and curly tails predict the strength of

behavioural correlations in support of the DS in 78 dog breeds. Contrary to the expectations of

covariation among DS traits, we found that morphological traits did not covary among

themselves, nor did they predict the strength of behavioural correlations among dog breeds.

Further, the number of morphological traits in a breed did not predict the strength of

behavioural correlations. Our results thus contrast with the hypothesis that the DS arises due to

a shared underlying mechanism, but more importantly, questions if the morphological traits

embedded in the DS are actual domestication traits or post-domestication improvement traits.

For dogs, it seems highly likely that strong selection for breed specific morphological traits

only happened recently in relation to breed formation. Present day dogs therefore have limited

bearing of the initial selection pressures applied during domestication and we should re-

evaluate our expectations of the DS accordingly.

Key words: Domestication syndrome, dog domestication, correlated traits, morphology,

behaviour

Introduction

Domesticated animals display suites of altered morphological, behavioural and physiological

traits compared to their wild ancestors, a phenomenon known as the domestication syndrome

(DS). Key examples of components in the DS are increased tameness, reduced brain size,

white pigmentation and decreased hypothalamic-pituitary-adrenal (HPA) axis reactivity (

Kruska, 1996; Driscoll et al., 2009; Trut et al., 2009). Because these alterations are observed to

co-occur across a wide range of present day domesticates, such as dogs (*Canis familiaris)*, cats

(*Felis catus)*, rabbits (*Oryctolagus cuniculus*), horses (*Equus caballus)* and pigs (*Sus scrofa*)

(Sánchez-Villagra et al., 2016), the traits within the DS are assumed to covary within species (

Trut, 1998; Trut et al., 2009). Domestication experiments have demonstrated that selection for

tame behaviour alone can produce the myriad changes seen in the DS (Belyaev et al., 1985;

Trut et al., 2009). While the mechanistic origin of the DS in currently unresolved, these

findings have nurtured the hypothesis that the convergent patterns seen across domesticated

species arise via a singular developmental mechanism such as altered neuroendocrine control

of ontogenesis (Belyaev, 1979), or neural crest deficit during embryogenesis (Wilkins et al.,

2014). Both of these influential studies have led to the general assumption that morphological

changes, such as white pigmentation, floppy ears and curly tails, have arisen as by-products of

the physiological alterations caused by selection upon behaviour (Wilkins et al., 2014).

The hypotheses that the DS is founded in single developmental mechanism offer a coherent,

logical and satisfying explanation for the observed covariation among DS traits. However,

traits of the DS are not fully consistent with such hypotheses. First, DS traits are not evenly

distributed among domesticated animals (Sánchez-Villagra et al., 2016). Second, even though

rat (*Rattus norvegicus*) lines selected for tameness have an increased frequency of white spots

(Trut et al., 2000), there is no genomic association between white coat colouration and tame

behaviour (Albert et al., 2009). This is unexpected based on the hypothesis that white

pigmentation should arise as a by-product of selection on tameness and further, because

syndrome traits originating from a shared physiological origin should be difficult to decouple

(*sensu* Sih et al., 2004). Finally, while recent genomic studies in horses (Librado et al., 2017),

foxes (*Vulpes vulpes*, Wang et al., 2018), dogs (Pendleton et al., 2018) and cats (Montague et

al., 2014) find signatures of domestication selection pressures in genes associated with neural

crest development, and thus are argued to support the neural crest hypothesis, these genes are

only a subset of many showing selective signatures during domestication. Thus, while it is

generally assumed that DS traits covary, possibly due to a single developmental mechanism,

further quantitative testing of this hypothesis is warranted.

Only recently has a formal test of covariance of DS traits been conducted. In their study of the

behavioural component of the DS in more than 76,000 dogs, Hansen Wheat et al. (2019)

demonstrated that while correlations between fear, aggression, sociability and playfulness were

stronger in ancient breeds, these correlations were weaker or had been decoupled in modern

breeds. However, this study focused only upon behaviour, which was likely the focal trait in

dog domestication (*sensu* Belyaev et al., 1985; Trut et al., 2009). To date, no studies have

investigated the covariation of morphological traits, either among themselves, or with the

expected behavioural correlations of the DS. Such a formal investigation of the predicted

expectations of how behavioural and morphological components of domestication arise is

needed if we are to further our understanding of the DS.

Among domesticates, the dog has been argued to be the only species expressing the full DS

(Sánchez-Villagra et al., 2016). Dogs have been bred for highly breed-specific morphological

and behavioural traits (Svartberg, 2006; Mehrkam and Wynne, 2014), which are illustrated by

the extreme phenotypic variation expressed among the more than 400 present day dog breeds

(Lindblad-Toh et al., 2005; Parker et al., 2017). The result is dramatic phenotypic variation

expressed across breeds. However, while key DS traits of behaviour and morphology do not

qualitatively appear to occur simultaneously across breeds (Sánchez-Villagra et al., 2016), this

has never been tested quantitatively. Furthermore, though dogs express a range of traits not

present in wolves (Parker et al., 2009; Larson et al., 2014), it is currently not well resolved if

dog traits are original domestication traits, i.e. traits evolved under direct selection in the initial

stages of domestication, or so-called improvement traits that have been secondarily enhanced

post-domestication during breed formation (Larson and Fuller, 2014; *sensu* Olsen and Wendel,

2013).

With modern breeds created from intense breeding efforts only within the last 150-200 years

(Lindblad-Toh et al., 2005; vonholdt et al., 2010), it is possible that modern dogs provide a

suboptimal basis for the expectations embedded in the DS. Indeed, as noted earlier, modern

dogs lack the strong behavioural correlations expected of the DS (Hansen Wheat et al., 2019).

Nonetheless, because the foundation for the DS hypothesis is based on extant domesticates, it

remains unclear if we should expect the expression of the DS to vary across different stages of

domestication. Archaeological findings of early dogs provide limited information on

morphology (i.e. skeletal features), and none on behaviour, which impairs our ability to

compare trait expression in dogs at different stages of domestication. Pre-breed formation

domesticated dogs, i.e. village dogs, could be very informative, but unfortunately, the only

non-admixed village dog populations identified to date are found in Borneo (Shannon et al.,

2015) and have not been studied behaviourally. However, a small group of present day dogs

can be categorized as ancient breeds due to their a) detectable admixture with wolf, which is

not present in modern breeds, and b) an origin about 500 years ago (Lindblad-Toh et al., 2005;

vonholdt et al., 2010). Certainly, ancient breeds are expected to have improvement traits, but

importantly, these breeds have been shown to have stronger behavioural correlations expected

of the DS compared to modern breeds (Hansen Wheat et al., 2019). While acknowledging they

are an imperfect proxy, ancient breeds are arguably the only available representatives for

earlier stages of dog domestication, and thus the division of ancient and modern breeds

provides an opportunity for temporal comparisons among dogs on a domestication time scale.

Here we test the hypothesis that the presence of morphological traits predict the strength of

behavioural correlations in support of the DS in dogs. For the morphological component of our

study, we focused upon variation in the traits white pigmentation, floppy ears and curly tails,

which have been referred to as morphological markers of domestication (Trut et al., 2009). For

the behavioural component, we used estimates of effect sizes for behavioural correlations

associated with the DS, derived from data extracted from the Swedish Kennel Club's database

on 76,158 dogs completing a highly standardized behavioural test battery (Hansen Wheat et

al., 2019). We then matched these effect sizes of behavioural correlations with our estimates of

morphological traits from the 78 breeds. We further added a temporal component by assessing

7 ancient and 71 modern breeds separately. As predicted by the DS, we expected that the

presence of white pigmentation, floppy ears and curly tails would co-vary among breeds.

Additionally, we expected that the presence or absence of these morphological traits would

predict the strength of behavioural correlations of the DS. That is, we expected stronger

behavioural correlations of the DS when morphological traits of the DS are present, as well as

the converse, weaker behavioural correlations when morphological traits of the DS are absent.

We further predicted that behavioural correlations would be stronger with the number of

morphological traits present.

Methods

Morphological assessment

We based our study on the 78 dog breeds used in a recent study to test behavioural correlations

within the domestication syndrome (Hansen Wheat et al., 2019). Of the 78 breeds, seven were

ancient breeds and 71 modern breeds. This difference in sample sizes between breed groups

does not reflect a lack in sampling effort, but the natural limitation of only few breeds being

categorized as ancient. We carefully inspected the breed standards for those 78 breeds by

consulting the Fédèration Cynologique Internationale, the world's largest federation of kennel

clubs, to assess the presence or absence of our three chosen morphological traits; white

pigmentation, floppy ears and curly tails. We used both relaxed and conservative assessments

of the three morphological traits (Figure 1, Figure 2, Figure S1). We defined white

pigmentation as any form of white pigmentation in the breed, regardless of its placement or

shape. We also classified dogs with a white base colour, such as Dalmatians and Samoyeds, to

express white pigmentation. Breeds where "white" was not mentioned in the coat colour

description, such as Dobermann and Rottweiler, were assessed as not having white

pigmentation. For our conservative assessment of white pigmentation, only breeds specifically

described to have a white base colour or characteristic white coloration, or breeds where some

versions have white pigmentation (such as Schnauzers) were included. Breeds were a small

white spot or a few white hairs are "tolerated" or "undesirable" were not included as having

white pigmentation in our conservative assessment. For the relaxed assessment, we included

breeds were small white spots or a few white hairs, for instance on the chest, are "tolerated" or

"undesirable" (Figure 1A-D). Floppy ears were assessed based on whether a breed has ears

that are either erect or to some degree floppy (i.e. from just the tip to hanging straight down,

Figure 1E-H). Thereby the presence or absence of floppy ears was assessed as a completely

binary trait, and did not differ between the relaxed and conservative assessments. For our

conservative assessment of curly tails, only breeds described to specifically have their tail in a

permanent curl, i.e. with no option to let down the tail, as seen in Pugs, were included. For the

relaxed assessment breeds that are described to carry their tail in a "curl", "hook", "sabre",

"sickle" or "J", and even breeds carrying their tails in the slightest "curve", but can let their

tails straight down were assessed as having curly tails (Figure 1I-L). Breeds where the words

"curl", "hook", "sabre", "sickle", "J" or "curve" were not included in the description of the tail

were assessed as not having a curly tail in either assessment.

Behavioural assessment

For the behavioural component of our study, we used the dataset presented in Hansen Wheat et al. 2019, in which the strength and direction of behavioural correlations between aggression, fearfulness, sociability and playfulness across the 78 dog breeds were investigated. Behavioural data were provided by the Swedish Kennel Club for dog completing the Dog Mentality Assessment, a highly standardized behavioural test for dogs in Sweden. We refer to this paper for a full description of the methods used to estimate the effect sizes for these behavioural correlations.

Statistical analyses

To evaluate the relationship between breed morphology and agreement with the domestication syndrome hypothesis, we assessed the correlation between our morphology scores, treated as dichotomous variables. First, we estimated the phi coefficient (ϕ) for presence/absence of each of each trait in pairwise combinations with significance determined using Fisher's Exact Test, as implemented in the xtab_statistics function of the sjstats package v. 0.17.5 (Lüdeke, 2019). Second, we repeated this analysis using a Pearson's product-moment correlation with similar results. Third, we assess whether the presence/absence of traits were correlated while taking into account phylogenetic correction, using a pairwise bionomial phylogenetic glm.

To evaluate the relationship between breed morphology and agreement with the domestication syndrome hypothesis, as quantified by the strength and direction of behavioural correlations, we used a meta-analytic model. It is a multi-level model that uses the 1326 observed correlation coefficients (Hansen Wheat et al., 2019), and their associated uncertainty, as the

dependent variable. These correlations test multiple behavioural predictions by the DS, such as

a positive association between sociability and playfulness, or a negative association between

sociability and aggression (Trut et al., 2009; Himmler et al., 2013). The correlations test six

such DS predictions. For some predictions multiple correlations per breed were measured,

since the Dog Mental Assessment test provided multiple measurements for aggression and

fearfulness. 17 correlations were obtained per breed. Therefore, we treat the DS as a nested

compound hypothesis, with six predicted associations and 17 correlations. We aligned the sign

of the correlations with the predicted directions, i.e. we flipped the sign of correlations

expected to be negative, so that positive effect sizes represent support in favour of the DS.

To account for this nested structure, we included group level effects that allow the support for

the DS to vary between the different predicted associations and the measured correlations. We

additionally included group level effects of morphology for the associations and correlations,

so that the moderating effect of morphological traits could be stronger or weaker depending on

what behavioural correlations were measured. Since each breed was represented by multiple

correlations, we included a group level intercept for breed. And because breeds are non-

independent due to shared ancestry (Felsenstein, 1985), an additional group level effect was

added with the expected covariance matrix of the phylogeny. Morphology was modelled as

three additive binary effects, one each for the presence or absence of white pigmentation,

floppy ears and curly tails. We implemented the models in the probabilistic programming

language Stan (Carpenter et al., 2017), using the interfacing R (R Core Team, 2019) package

brms (Bürkner 2017, 2018). In brms syntax, the models were of the form: *Zr | se(vi, sigma =*

TRUE) ~ breed_category + pigmentation + ears + tails + (1 + breed_category +

pigmentation + ears + tails || prediction/correlation) + (1 | breed) + (1 | phylogeny), where Zr

is the z-transformed correlations coefficients, vi is the measurement error and sigma = TRUE

allows for the estimation of the residual standard deviation.

Inference about the effects of morphology was based on two approaches. We used the

posterior distributions for the parameters directly to evaluate the role of the three

morphological traits separately. Secondly, we assessed the role of the number of

morphological traits (regardless of which trait) by calculating the estimated mean response for

each trait combination, and then calculating the marginal mean for a breed having 0, 1, 2 or 3

traits present.

Posterior distributions for the parameters were obtained through MCMC sampling, using 16

chains of 2000 iterations each, of which 1000 were warmup. We adjusted the adapt delta to

0.995 and the maximum tree depth to 20 to eliminate any divergent transitions. For population

level effects, we used the default weak student-t prior with a mean of 0, scale parameter of 10

and 3 degrees of freedom. The same prior was used for standard deviations of group-level

effects and the residual standard deviation, but there it was restricted to be non-negative. Trace

plots indicated that the chains were well mixed, and we obtained an effective sample size of

more than 2500 for all parameters. The largest $\hat{R}$ was 1.01, indicating convergence.

All analyses were done for both relaxed and conservative assessments of morphological traits.

Results for the two different assessments were qualitatively similar, and below we present the

results for the relaxed assessment (see Supplemental Files for results for the conservative

assessment)

Results

We placed the morphological traits and average effect sizes for behavioural correlations onto

the latest dog phylogeny (Parker et al., 2017), revealing large variation among breeds in both

our morphological and behavioural traits (Figure 2, for conservative assessments see

Supplemental Files and Figure S1).

First, we used three different methods to test whether the presence of morphological DS traits

covary amongst themselves. Neither phi coefficients (φ), Pearson's product-moment

correlation (t) nor phylogenetically corrected correlations (z) for the three morphological

traits produced significant results: white pigmentation vs. floppy ears ($\varphi = 0.172$, $p_\varphi = 1$; $t = -$

0.115, $p_t = 0.909$; $z = -0.080$, $p_z = 0.937$), white pigmentation vs. curly tail ($\varphi = 0.013$, $p_\varphi =$

1; $t = -5.7071^{-20}$, $p_t = 1$; $z = 0.653$, $p_z = 0.514$), floppy ears vs. curly tail ($\varphi = 0$, $p_\varphi = 0.2063$;

$t = -1.5176$, $p_t = 0.1333$; $z = -0.49807$, $p_z = 0.618$).

Second, to test whether the presence of white pigmentation, floppy ears and curly tails predicts

the strength of any of the behavioural correlations, we evaluated these traits as binary

predictors of DS support. We found that there was no difference in the behavioural correlations

when any of the three morphological traits were present or absent (Table 1, Figure 3A and B,

Figure S2 and Supplemental Files). We emphasize that there is no support for even a very

small difference in effect size (most extreme effect within CI: 0.04, Table 1). We did not

confirm an effect of breed age, as the difference between ancient and modern breeds could not

be clearly distinguished from 0, although considerable uncertainty in this estimate remains and

most of the posterior favours stronger behavioural correlations in ancient breeds (Figure 3A

and B, Table 1, for non conservative measurements see Supplemental Files).

Lastly, we evaluated support for the DS based on the "morphology score" of each breed, which

ranged from 0 - 3 depending on how many, if any, of the three morphological traits is present

in a breed (Figure 3C, Supplemental Files). We found that the number of morphological traits

present in a breed did not predict the strength of behavioural correlations (0 traits: posterior

$mean_{slope}$ = 0.080, 95CI[-0.006 - 0.191]; 1 trait: posterior $mean_{slope}$ = 0.080, 95CI[-0.008 -

0.193]; 2 traits: posterior $mean_{slope}$ = 0.082, 95CI[-0.006 - 0.192]; 3 traits: posterior $mean_{slope}$ =

0.083, 95CI[-0.004 - 0.190]). Given the small number of ancient breeds, we were not able to

include breed age in this morphology score analysis.

Discussion

Here we tested the whether the presence of three traits referred to as the morphological

markers of domestication (white pigmentation, floppy ears and curly tails) predicted the

strength of behavioural correlations within the DS. Contrary to the expectations of covariation

among DS traits, we found that these morphological traits did not covary among themselves,

nor did they predict the strength of behavioural correlations among dog breeds. Further, the

number of morphological traits in a breed did not predict the strength of behavioural

correlations. Additionally, we found no effect of breed age, i.e. ancient and modern breeds, in

the predictive value of morphological traits on behavioural correlations. A high covariance

among DS traits suggests a strong, central role for their shared origin in a single developmental

source (e.g. white pigmentation arising as a by-product of increased tameness, Wilkins et al.,

2014), while a lack of covariance suggests a more complex genotype to phenotype

relationship. Thus, the lack of covariation among morphological and behavioural traits in our

study is not consistent with the hypothesis that trait alterations in the DS are founded in a

singular developmental source (Belyaev, 1979; Wilkins et al., 2014).

The DS in animals is primarily based on observations in present day domesticates. However,

the ability of phenotypes in extant domesticates to provide insights about selection during

initial domestication is complicated by post-domestication selection events, i.e. improvement

traits (Olsen and Wendel, 2013; Larson and Fuller, 2014). Initial domestication efforts likely

targeted existing variation at multiple loci across the genome (Larson and Fuller, 2014), but the

breed-specific morphology and behaviour expressed in present day dog breeds was likely

selected for post-domestication during breed formation. Many of the morphological traits seen

across modern dog breeds are therefore not likely to be by-products of initial selection for

domestication traits rather they are most likely improvement traits. Thus, while studies refer to

the phenotypes of modern dog breeds as evidence for the DS (Wilkins et al., 2014; Sánchez-

Villagra et al., 2016), whether these traits are relevant to domestication itself is questionable.

Thus, our findings of a lack of covariation among morphological and behavioural traits, rather

than providing insights into the DS, could be due to these traits being improvement traits, for

which no covariance is expected. Regardless, the phenotypes of modern dog breeds should be

interpreted with caution when trying to understand the domestication process.

One way to gain more insight into selection pressures during earlier stages of dog

domestication, rather than those of breed improvement, is to include a temporal comparison by

separating out ancient breeds and modern breeds. Here, we investigated whether the presence

of morphological traits predict the strength of behavioural correlations in each breed group, but

found no such effect. This finding contrasts with a recent study in which behavioural

correlations of the DS were demonstrated to be stronger in ancient breeds compared to modern

breeds (Hansen Wheat et al., 2019). Given that selection on tameness alone can generate the

DS in foxes (Trut et al., 2009), and that aggression shows selective signatures directly

associated with initial domestication efforts in these selection lines of foxes (Kukekova et al.,

2018), it is likely that initial selection during dog domestication acted upon behaviour, not

morphology (*sensu* Belyaev et al., 1985; Trut et al., 2009). Thus, with behaviours in the DS

likely representing domestication traits, behavioural domestication phenotypes might to a

larger extent be maintained in ancient compared to modern breeds. Morphology in dog breeds

on the other hand, is arguably linked to breed improvement (Larson and Fuller, 2014), as

reflected in the large variability in morphological trait combinations across dog breeds as

quantified here.

In sum, whether the lack of covariance between morphology and behaviour in dogs is due to

decoupling of independent domestication alleles (possibly caused by altered selection regimes

during breed formation), these traits never having covaried because of a singular

developmental mechanism or whether it is because we are applying a domestication hypothesis

on traits that are not actual domestication traits, but rather improvement traits, remains an open

question. If the latter is true, which seems likely for dogs, we must reevaluate our expectations

of the DS and thereby also our assessment of DS traits in present day domesticates, as they

have limited bearing of the initial selection pressures applied during domestication.

Author contributions

CHW and CWW conceived the study. CHW prepared the data and all authors discussed how

to analyse it. WvdB and CWW analysed the data. CHW prepared the manuscript draft and

WvdB and CWW provided comments to produce the final version.

Data availability statement

All data used for this study is available through Figure 2, the Supplemental Materials and

Hansen Wheat et al. 2019.

References

Albert, F. W., Carlborg, Ö., Plyusnina, I., Besnier, F., Hedwig, D., Lautenschlager, S., et al. (2009). Genetic
Architecture of Tameness in a Rat Model of Animal Domestication. *Genetics* 182:541–554.
doi:10.1534/genetics.109.102186.

Belyaev, D. K. (1979). Destabilizing selection as a factor in domestication. *Journal of Heredity* 70:301-308

Belyaev, D. K., Plyusnina, I. Z., and Trut, L. N. (1985). Domestication in the silver fox (Vulpes fulvus Desm):
Changes in physiological boundaries of the sensitive period of primary socialization. *Applied Animal*
Behaviour Science 13:359–370. doi:10.1016/0168-1591(85)90015-2.

Bürkner PC. (2017). brms: An R Package for Bayesian Multilevel Models Using. Stan. Journal of Statistical
Software. *80*:1-28.
355
356 Bürkner PC. (2018). Advanced Bayesian Multilevel Modeling with the R Package brms. The R Journal. *10*:395-
357 411.

358 Carpenter B, Gelman A, Hoffman MD, Lee D, Goodrich B, Betancourt M, Brubaker M, Guo J, Li P, Riddell A.
359 (2017). Stan: A probabilistic programming language. Journal of Statistical Software. *76*.
360
Driscoll, C. A., Macdonald, D. W., and O'Brien, S. J. (2009). From wild animals to domestic pets, an
evolutionary view of domestication. *PNAS* 106:9971–9978.

Fédération Cynologique Internationale, www.fci.be

Felsenstien J. (1985). Phylogenies and the Comparative Method. Am. Nat.*125:*1-15

Hansen Wheat, C., Fitzpatrick, J. L., Rogell, B., and Temrin, H. (2019). Behavioural correlations of the
domestication syndrome are decoupled in modern dog breeds. *Nature Communcations 10:*2422.
doi:10.1038/s41467-019-10426-3.

Himmler, B. T., Stryjek, R., Modlinska, K., Derksen, S. M., Pisula, W., and Pellis, S. M. (2013). How
domestication modulates play behavior: A comparative analysis between wild rats and a laboratory strain of
Rattus norvegicus. *Journal of Comparative Psychology* 127:453–464. doi:10.1037/a0032187.

Kruska, D. (1996). The effect of domestication on brain size and composition in the mink (*Mustela vison*). *J*
Zoology 239:645–661. doi:10.1111/j.1469-7998.1996.tb05468.x.

Kukekova, A. V., Johnson, J. L., Xiang, X., Feng, S., Liu, S., Rando, H. M., et al. (2018). Red fox genome
assembly identifies genomic regions associated with tame and aggressive behaviours. *Nat. ecol. evol.*, 1–16.
doi:10.1038/s41559-018-0611-6.

Larson, G., and Fuller, D. Q. (2014). The Evolution of Animal Domestication. *Annu. Rev. Ecol. Evol. Syst.*
45:115–136. doi:10.1146/annurev-ecolsys-110512-135813.

Larson, G., Piperno, D. R., Allaby, R. G., Purugganan, M. D., Andersson, L., Arroyo-Kalin, M., et al. (2014).
Current perspectives and the future of domestication studies. *Proceedings of the National Academy of*
Sciences 111:6139–6146. doi:10.1073/pnas.1323964111.

Librado L. E., et al. (2017). Ancient genomic changes associated with domestication of the horse. *Science*
356:442–445

Lindblad-Toh, K., Wade, C. M., Mikkelsen, T. S., Karlsson, E. K., Jaffe, D. B., Kamal, M., et al. (2005).
Genome sequence, comparative analysis and haplotype structure of the domestic dog. *Nature* 438:803–819.
doi:10.1038/nature04338.

Lüdecke D (2019. *sjstats: Statistical Functions for Regression Models (Version 0.17.5)*, https://CRAN.R-
project.org/package=sjstats

Mehrkam, L. R., and Wynne, C. D. L. (2014). Behavioral differences among breeds of domestic dogs (Canis
lupus familiaris): Current status of the science. *Applied Animal Behaviour Science* 155:12–27.
doi:10.1016/j.applanim.2014.03.005.

Montague, M. J., Li, G., Gandolfi, B., Khan, R., Aken, B. L., Searle, S. M. J., et al. (2014). Comparative analysis
of the domestic cat genome reveals genetic signatures underlying feline biology and domestication.
Proceedings of the National Academy of Sciences 111:17230–17235. doi:10.1093/nar/gks596.

Olsen, K. M., and Wendel, J. F. (2013). A Bountiful Harvest: Genomic Insights into Crop Domestication
Phenotypes. *Annu. Rev. Plant Biol.* 64:47–70. doi:10.1146/annurev-arplant-050312-120048.

Parker, H. G., Dreger, D. L., Rimbault, M., Davis, B. W., Mullen, A. B., Carpintero-Ramirez, G., et al. (2017).
Genomic Analyses Reveal the Influence of Geographic Origin, Migration, and Hybridization on Modern
Dog Breed Development. *CellReports* 19:697–708. doi:10.1016/j.celrep.2017.03.079.

Parker, H. G., vonholdt, B. M., Quignon, P., Margulies, E. H., Shao, S., Mosher, D. S., et al. (2009). An
expressed fgf4 retrogene is associated with breed-defining chondrodysplasia in domestic dogs. *Science*
325:995–998. doi:10.1126/science.1173275.

Pendleton, A. L., Shen, F., Taravella, A. M., Emery, S., Veeramah, K. R., Boyko, A. R., et al. (2018).
Comparison of village dog and wolf genomes highlights the role of the neural crest in dog domestication.
BMC Biology 16:64.

R Core Team (2019). R: A language and environment for statistical computing. R Foundation
for Statistical Computing, Vienna, Austria. www.R-project.org.

Sánchez-Villagra, M. R., Geiger, M., and Schneider, R. A. (2016). The taming of the neural crest: a
developmental perspective on the origins of morphological covariation in domesticated mammals. *R. Soc.*
open sci. 3:160107. doi:10.1093/icb/45.3.387.

Shannon, L. M., Boyko, R. H., Castelhano, M., Corey, E., Hayward, J. J., McLean, C., et al. (2015). Genetic
structure in village dogs reveals a Central Asian domestication origin. *Proceedings of the National*
Academy of Sciences 112:13639–13644. doi:10.1073/pnas.0909559107.

Sih, A., Bell, A., and Johnson, J. C. (2004). Behavioral syndromes: an ecological and evolutionary overview.
Trends in Ecology & Evolution 19:372–378. doi:10.1016/j.tree.2004.04.009.

Svartberg, K. (2006). Breed-typical behaviour in dogs—Historical remnants or recent constructs? *Applied*
Animal Behaviour Science 96:293–313. doi:10.1016/j.applanim.2005.06.014.

Trut, L. N. (1998). The evolutionary concept of destabilizing selection: status quo. *Journal of Animal Breeding*
and Genetics 115:415–431.

Trut, L. N., Pliusnina, I. Z., and Prasolova, L. A. (2000). Mutations hooded and nonagouti in the gray rat (Rattus
norvegicus): effect of selection on behavior and photoperiod. *Genetika* 36:813–822.

Trut, L., Oskina, I., and Kharlamova, A. (2009). Animal evolution during domestication: the domesticated fox as
a model. *BioEssays* 31:349–360. doi:10.1002/bies.200800070.

vonholdt, B. M., Pollinger, J. P., Lohmueller, K. E., Han, E., Parker, H. G., Quignon, P., et al. (2010). Genome-
wide SNP and haplotype analyses reveal a rich history underlying dog domestication. *Nature* 464:898–902.
doi:10.1038/nature08837.

Wang, X., Pipes, L., Trut, L. N., Herbeck, Y., Vladimirova, A. V., Gulevich, R. G., et al. (2018). Genomic
responses to selection for tame/aggressive behaviors in the silver fox (Vulpes vulpes). *Proceedings of the*
National Academy of Sciences 115:10398–10403. doi:10.1007/978-1-4939-0512-6_3.

Wilkins, A. S., Wrangham, R. W., and Fitch, W. T. (2014). The "Domestication Syndrome" in Mammals: A
Unified Explanation Based on Neural Crest Cell Behavior and Genetics. *Genetics* 197:795–808.
doi:10.1534/genetics.114.165423.

Photo references:

465   Rhodesian Ridgeback: By Bonnie van den Born, http://www.bonfoto.nl - Transferred from nl.wikipedia to
466   Commons., CC BY-SA 3.0, https://commons.wikimedia.org/w/index.php?curid=1747204
467

468   St. Bernard: By Cassie J - originally posted to Flickr as Harley-parents' dog, CC BY 2.0,
469   https://commons.wikimedia.org/w/index.php?curid=5120324
470

472   **Table 1. Predictive value of morphological traits.** Predictive value of the presence or absence of morphological

473   traits on the strength of behavioural correlations in the DS. Posterior mean, posterior standard deviation (sd) and

474   95% Credible Interval (CI) given for breed category (ancient and modern) and the three morphological traits

475   white pigmentation, floppy ears and curly tail.

Term	Posterior mean	Posterior sd	95CI $_{lower}$	95CI $_{upper}$
Intercept	0.113	0.053	0.02	0.22
Breed category	-0.06	0.034	-0.129	0.004
White pigmentation	0.001	0.014	-0.027	0.03
Floppy ears	0.004	0.019	-0.033	0.04
Curly tail	-0.001	0.014	-0.026	0.026

Figure 1. Morphological assessments. Examples of morphological variation across dog breeds and how this was taken into account when assessing the presence and absence of morphological traits in the DS. *White pigmentation (pigment, A-D):* Breeds where a small white spot or a few white hairs on the chest is tolerated or undesirable, here illustrated in a Rhodesian Ridgeback (A), were categorized as not having white pigmentation in the conservative assessment, but as having white pigmentation in the relaxed assessment. The presence of white pigmentation varies across breeds in size, shape and placement as illustrated in Bernese Mountain Dog (B), German Short-haired Pointer (C) and Dalmatian (D). *Floppy ears (ears, E-H)*: Floppiness of ears is binary and erect ears, as illustrated in the Shiba (E), can never be floppy. Other examples of breeds with erect ears are Siberian Husky (J) and Alaskan Malamute (K). The floppiness of ears can be graduated as illustrated by the Staffordshire Bull Terrier (F), Labrador Retriever (G) and English Springer Spaniel (H). Any degree of floppiness of the ears was assessed as presence of floppy ears. *Curly tail (tails, I-L):* Breeds, such as the St. Bernard (I), with tails hanging straight down and never carry their tail in a curl, curve, hook, sickle, sabre of J shaped express the absence of a curly tail (both assessments). Many breeds carry their tail in a curl, curve, hook, sickle, sabre of J shaped fashion, but can also let their tail straight down, here illustrated by Siberian Husky with a let down tail (J) and an Alaskan Malamute with a tail carried in a curl (K). For the conservative assessment such breeds were categorized as not having curly tails, while they were categorized as having curly tails in the relaxed assessment.

507   Other examples of breeds categorized like this are Rhodesian Ridgeback (A) and Dalmatian (D). A few breeds,

508   like Pugs (L), express the presence of a permanent curly tail (both assessments). All photos are from

509   wikicommons, please see references for specific credits.

Figure 2. Morphological scores placed onto the latest dog phylogeny. Morphological scores based on the presence or absence of curly tail, floppy ears and white pigmentation (relaxed assessment), and average effect sizes for behavioural correlations in ancient and modern dog breeds placed onto the latest dog phylogeny (Parker et al 2017). Average effect sizes were calculated by separate meta-analytic models per breed (not used for inference), and posterior means +- 95% credible intervals are depicted.

Figure 3. Morphological traits and the strength of behavioural correlations. A) Estimated support for the DS,

quantified as the strength of behavioural correlations (Z_r) depending on the presence or absence of morphological

traits (relaxed assessment) and trait category. B) Regression coefficients indicating the difference between binary

categories, as in A). C) The number a morphological traits present (relaxed assessment), i.e. morphological score,

related to the estimated strength of behavioural correlations within the DS. In all panels, density distributions

depict the full posterior distributions, with the thick lines covering the 66% credible interval, thin lines the 95%

credible interval and point estimate the posterior median. Scattered points in A) and C) are the estimated average

effect size per breed (as in Figure 2).