Mesenchymal stem cells suppress leukemia via macrophage-mediated functional restoration of bone marrow microenvironment

Chengxiang Xia[1, 3, 4,7†], Tongjie Wang[1†], Hui Cheng[2,5†], Yong Dong[1, 3], Qitong Weng[1, 3], Guohuan Sun[2,5], Peiqing Zhou[1, 3], Kaitao Wang[6], Xiaofei Liu[1], Yang Geng[1], Shihui Ma[2,5], Sha Hao[2,5], Ling Xu[8], Yuxian Guan[1], Juan Du[1], Xin Du[9,10], Yangqiu Li[8], Xiaofan Zhu[2,5], Yufang Shi[11], Sheng Xu[12], Demin Wang[13,14], Tao Cheng[2,5*], Jinyong Wang[1, 3, 4, 6, 7*]

[1]State Key Laboratory of Experimental Hematology, CAS Key Laboratory of Regenerative Biology, Guangzhou Institutes of Biomedicine and Health, Chinese Academy of Sciences, Guangzhou, China; [2]State Key Laboratory of Experimental Hematology & National Clinical Research Center for Blood Diseases, Institute of Hematology & Blood Diseases Hospital, Chinese Academy of Medical Sciences & Peking Union Medical College, Tianjin, China; [3]Guangzhou Regenerative Medicine and Health-Guangdong Laboratory (GRMH-GDL), Guangzhou, China; [4]Guangdong Provincial Key Laboratory of Stem cell and Regenerative Medicine, Guangzhou Institutes of Biomedicine and Health, Chinese Academy of Sciences, Guangzhou, China; [5]Center for Stem Cell Medicine & Department of Stem Cell and Regenerative Medicine, Chinese Academy of Medical Sciences & Peking Union Medical College, Tianjin, China; [6]Joint School of Life Sciences, Guangzhou Institutes of Biomedicine and Health, Guangzhou Medical University, Guangzhou, China; [7]University of Chinese Academy of Sciences, Beijing, China; [8]Department of Hematology, First Affiliated Hospital; Institute of Hematology, School of Medicine; Key Laboratory for Regenerative Medicine of Ministry of Education; Jinan University, Guangzhou, China; [9]Department of Hematology, Guangdong Provincial People's Hospital (Guangdong Academy of Medical Sciences), Guangzhou, Guangdong, China; [10]South China University of Technology, Guangzhou, China; [11]The First Affiliated Hospital of Soochow University, State Key Laboratory of Radiation Medicine and Protection, Institutes for Translational Medicine, Soochow University, Suzhou, China; [12]National Key Laboratory of Medical Immunology & Institute of Immunology, Second Military Medical University, Shanghai, China; [13]Blood Research Institute, Versiti, Milwaukee, WI, USA; [14]Biomedical Research Center of South China, College of Life Sciences, Fujian Normal University, Fuzhou, China.

[†]Equal contributors.

[*]Correspondences:

Jinyong Wang (J.W.), E-mail: wang_jinyong@gibh.ac.cn

Tao Cheng (T.C.), E-mail: chengtao@ihcams.ac.cn

33 **Key Points:**

34 *Key Point 1*: Intra-BM transfusion of MSCs restores the BM microenvironment, improves

35 thrombopoiesis, and suppresses MDS/MPN initiated by *Nras* mutation.

36 *Key Point 2*: Donor MSCs reprogram macrophages to restore the BM microenvironment, improve

37 thrombopoiesis, and suppress leukemia.

Abstract

Bone marrow (BM) mesenchymal stem cells (MSCs) are critical components of the BM microenvironment and play an essential role in supporting hematopoiesis. Dysfunction of MSCs is associated with the impaired BM microenvironment that promotes leukemia development. However, whether and how restoration of the impaired BM microenvironment can inhibit leukemia development remain unknown. Using an established leukemia model and the RNA-seq analysis, we discovered functional degeneration of MSCs during leukemia progression. Importantly, intra-BM instead of systemic transfusion of donor healthy MSCs restored the BM microenvironment, thus systemically altering cytokine expression patterns, improving normal hematopoiesis, reducing tumor burden, and ultimately prolonging survival of the leukemia-bearing mice. Donor MSC treatment restored the function of host MSCs and reprogrammed host macrophages to fulfill tissue-repair function. Transfusion of MSC-reprogrammed macrophages largely recapitulated the therapeutic effects of MSCs. Further, we found that donor MSCs reprogrammed macrophages to reduce leukemia burden through autocrine of IL-6. Taken together, our study reveals that donor MSCs reprogram host macrophages to restore the BM microenvironment and inhibit leukemia development, thus offering rationales for local MSC administration as a potentially effective therapy for leukemia.

Introduction

BM MSCs are key components of the BM stromal microenvironment, regulating homeostasis of the stromal niche and hematopoiesis[1]. Accumulating evidence uncovers that dysfunction of MSCs is associated with leukemia progression. Leukemic MSCs in a mouse T-ALL model suppressed normal hematopoiesis[2]. Deletion of *Dicer1* gene in MSCs leads to impaired osteogenic differentiation, thrombopenia, and secondary tumorigenesis[3]. In myeloproliferative neoplasms, the leukemic cells reprogram MSCs to support proliferation of leukemic stem cells[4]. Ablation of MSC-derived osteoblasts impairs the homeostasis of hematopoietic stem cells (HSC) and accelerates leukemogenesis in chronic myeloid leukemia[5]. In a myelodysplastic/myeloproliferative neoplasms (MDS/MPN) model, *Ptpn11* mutation in MSCs hyperactivates HSC by secreting CCL3, systemically resulting in myeloid-biased proliferation and promoting leukemia progression[6]. Chromosomal abnormalities in MSCs occur in over 16% MDS/acute myeloid leukemia (AML) patients, which leads to a shorter survival[7]. MSCs of every MDS patient show features of stemness loss, aging, and osteogenic differentiation defect, which results in insufficient hematopoiesis[8]. In a patient-derived xenograft (PDX) model, only co-transplantation of tumor cells and the same patient-derived MSCs successfully recapitulates the MDS phenotype, indicating that besides the loss of function of supporting hematopoiesis, MSCs in MDS patients may acquire a function of preferentially promoting tumorigenesis[9]. Despite these findings, the causal relationship between tumor cells and the BM microenvironment during leukemia development and progression remains elusive.

MSC therapy has been widely used in treating immune-related graft-vs-host disease (GVHD) and inflammation-related diseases[10,11]. Over the decades, a lot of evidence demonstrates that MSCs regulate innate and adaptive immune responses largely by secreting distinct sets of cytokines,

growth factors, and chemokines depending on different disease contexts[12-16]. Given the short

lifespan of donor MSCs after transfusion[17], the underlying molecular and cellular mechanisms by

which these cells produce therapeutic effects remain elusive. It is also completely unknown

whether donor MSCs can restore the impaired BM microenvironment and consequently suppress

disease progression in leukemia setting.

Macrophages are pivotal for maintenance of the tissue microenvironment, tissue repair, and even

the tumor microenvironment[18-22]. BM resident macrophages maintain the homeostasis of HSCs

and loss of these macrophages leads to mobilization of HSCs into peripheral blood (PB)[23]. The

functions of macrophages are plastic and can be reshaped by distinct sets of soluble factors. When

performing tissue repair, macrophages highly express arginase 1 ($Arg1$)[24], an enzyme that converts

L-arginine to urea and L-ornithine. After co-culture with MSCs, macrophages can be polarized

from pro-inflammation (M1) to anti-inflammation (M2) type, up-regulating IL-10 and CD206 and

down-regulating IL-6 and IL-1β[25]. Upon stimulated by LPS or TNF-α, MSCs can cross-talk with

lung macrophages and reprogram these macrophages to secrete IL-10 to alleviate sepsis[26]. Despite

these knowledge, whether healthy MSCs can reprogram macrophages from leukemia-bearing host

to repair the damaged BM microenvironment is not known.

Using the established mouse model mimicking chronic MPN/MDS diseases[27-29], we discovered

that the deteriorating BM microenvironment was associated with disease progression. Intra-BM

instead of systemic transfusion of healthy MSCs restored the local BM microenvironment,

improved thrombopoiesis, reduced tumor burden, and prolonged survival of leukemia-bearing

mice. Mechanistically, we found that MSCs suppress leukemia development through resident

macrophages and autocrine effect of IL-6. Our study demonstrates that intra-BM transfusion of

MSCs can restore the local BM niche to systemically prevent leukemia progression and can be a

novel therapy for leukemia.

Materials and Methods

Mice. All mouse strains were maintained on C57BL/6 genetic background. Mice expressing the

conditional oncogenic NrasG12D mutation (a gift from Dr. Jing Zhang lab at University of

Wisconsin-Madison, Wisconsin, USA) were crossed to Vav-Cre mice to generate *LSL Nras*/+;

Vav-Cre compound mice (NV mice). Genotyping of the adult mice was performed as described

previously[27]. Vav-Cre strain (CD45.2), wild-type CD45.2, CD45.1 strain (C57BL/6) and *Il6* knock

out strain (*Il6*$^{-/-}$, C57BL/6) were purchased from Jackson lab. GFP strain (CD45.2) was gifted by

Guangdong Laboratory Animals Monitoring Institute. *MLL-AF9* AML model mice were

maintained a specific pathogen-free animal facility at the State Key Laboratory of Experimental

Hematology. All mice were maintained within the SPF grade animal facility of Guangzhou

Institution of Biomedicine and Health, Chinese Academy of Science (GIBH, CAS, China). All

animal experiments were approved by the Institutional Animal Care and Use Committee of

Guangzhou Institutes of Biomedicine and Health (IACUC-GIBH).

NrasG12D leukemia model. White blood cells (CD45.2$^+$, 0.3 million) after depletion of stromal

cells from NrasG12D compound mice (*LSL Nras/+; Vav-Cre*) or control mice (CD45.2 strain)

were sorted and transplanted into sublethally (6.5 Gy, RS2000, Rad Source Inc) irradiated CD45.1

recipient by retro-orbital intravenous injection. Mice were fed with trimethoprim-

sulfamethoxazole-treated water for two weeks to prevent infection. Hematopoietic lineages in PB

were assessed monthly by flow cytometry. During the development of NrasG12D-induced

leukemia, the CD11b$^+$ percentage in PB indicated the tumor burden (CD11b$^+$%).

RNA-Seq and data analysis. For MSC library preparation, MSCs were sorted from wild type or

leukemia-bearing mice, and recovered MSCs were sorted from leukemia-bearing mice 8 weeks

post treatment with GFP[+] donor MSCs. MSCs were sorted from two mice of each group. 1000

target cells per sample were sorted into 500 µl DPBS-BSA buffer (0.5%BSA) using 1.5ml EP tube

and transferred into 250 µl tube to spin down with 500 g. The cDNA of sorted 1000-cell aliquots

were generated and amplified as described previously[30]. The qualities of the amplified cDNA were

examined by Q-PCR analysis of housekeeping genes (*B2m, Actb, Gapdh, Ecf1a1*). Samples passed

quality control were used for sequencing library preparation by illumina Nextera XT DNA Sample

Preparation Kit (FC-131-1096).

For macrophages (*in vivo*) library preparation, macrophages were sorted from BM of leukemia-

bearing mice before or after MSC treatment (12 hours post MSC treatment). Macrophages were

also sorted after 12 hours of co-culture with MSCs. 1×10^5 target cells per sample were sorted

and total RNA was extracted using the RNeasy micro kit with on-column DNase treatment

(Qiagen, 74004) according to manufacture's protocol. cDNA library was constructed using

VAHTSTM mRNA-seq V3 Library Prep Kit for Illumina (Vazyme, NR611) according to

manufacture's protocol. The qualities of the cDNA were examined by qPCR analysis of

housekeeping genes (*B2m, Actb, Gapdh, Ecf1a1*). Samples that passed quality control were used

for sequencing.

For data analysis, all libraries were sequenced by illumina sequencers NextSeq 500. The fastq files

of sequencing raw data samples were generated using illumina bcl2fastq software (version:

2.16.0.10) and were uploaded to Gene Expression Omnibus public database (GSE 125029). Raw

reads were aligned to mouse genome (mm10) by HISAT2[31] (version: 2.1.0) as reported. And raw

counts were calculated by featureCounts of subread[32] (version 1.6.0). Differential gene expression

analysis was performed by DESeq2[33] (R package version: 1.18.1). Unsupervised clustering

analysis was performed using facotextra (R package, version：1.0.5). Heatmaps were plotted

using gplots (R package, version 3.01). GSEA was performed as described[34], and gene-ontology

(GO)-enrichment analysis were performed by clusterProfiler[35] (R package, version: 3.6.0). MSC

stemness related genes and MSC osteogenesis related genes for heatmaps were from references as

follows: MSC stemness-related genes[36-38] and MSC osteogenesis-related genes[36,39]. The gene sets

for GSEA were from literatures as follows: angiogenesis-related genes in macrophages[40], cell

migration-related genes in macrophages (from MSigDB genesets), and secreted factors by

macrophages[41,42].

MSC treatment for leukemia-bearing mice. For MSC transfusion, multiple approaches

including retro-orbital, tail intravenous, and local intra-BM transfusion were applied

independently. For tail vein transfusion, each leukemia-bearing mouse was injected with $2.5 \times$

10^7 MSCs/kg (Passage 2) in 100 μl DPBS by tail vein transfusion. For retro-orbital transfusion,

each leukemia-bearing mouse was injected with 2.5×10^7 MSCs/kg (Passage 2) in 200 μl DPBS

by retro-orbital transfusion. For local intra-BM transfusion, tibia of each leukemia-bearing mouse

was injected with 2.5×10^7 MSCs/kg (Passage 2) in 20 μl DPBS by local intra-BM transfusion.

MSCs were injected once every two weeks and continued in a time window of 16 weeks. Every

tibia was treated once per month by switching the injection site every other dose. The control mice

were injected with DPBS following the same treatment procedure as MSCs. Analysis of platelets

and CD11b$^+$ cells in PB was performed monthly.

GFP-MSCs and BM macrophage co-culture assay. Short-term co-culture assay was performed,

with each well containing: 1×10^5 GFP-MSCs (passage 2; healthy MSCs were isolated from

GFP mice, *Il6*$^{-/-}$ MSCs were isolated from *Il6*$^{-/-}$ mice) and 2×10^6 CD11b$^+$ leukemic cells sorted

from leukemia-bearing mice in 2 mL culture medium of α-MEM, 10% FBS and 50 ng/ml SCF.

MSCs and CD11b$^+$ leukemic cells were incubated either by direct-contact culture or transwell

culture for 12 hours at 37^{o}C under 5% CO_2 in a humidified incubator. MSC-reprogrammed

macrophages from leukemia-bearing mice (CD11b$^+$F4/80$^+$) were sorted for detecting the gene

expression by Q-PCR.

Treatment for leukemia-bearing mice with MSC-reprogrammed macrophages. $1 \times 10^\wedge 5$

MSCs were seeded into each well of six-well plates. CD11b$^+$ leukemic cells were enriched from

BM of leukemia-bearing mice with severe tumor burden (CD11b$^+$% in PB > 60%). Then $2 \times 10^\wedge 6$

CD11b$^+$ leukemic cells were directly co-cultured with MSCs. After 12 hours, macrophages were

sorted for transfusion. Leukemia-bearing mice with severe tumor burden were treated by intra-BM

transfusion of PBS or MSC-reprogrammed macrophages from leukemia-bearing mice (E-Mac). A

dose of 1 million macrophages/mouse in 20 μl PBS were delivered into the tibia cavity using 29-

gauge needle. Every tibia was treated once per two weeks by switching the injection site every

other dose. Analysis of platelets and CD11b$^+$ cells in PB was performed monthly.

Statistical analysis. The data were represented as mean $\pm$ SD. Two-tailed independent Student's

t-tests were performed for comparison of two groups of data (SPSS v.23, IBM Corp., Armonk,

NY, USA). For the analysis of three groups or more, one-way ANOVA was used (SPSS v.23, IBM

Corp., Armonk, NY, USA), and further significance analysis among groups was analyzed by Post

Hoc Test (equal variances, Turkey-HSD; unequal variances, Games-Howell). Kaplan-Meier

method was used to calculate survival curves of leukemia, and Log-rank (Mantel-Cox) test was

performed to compare differential significance in survival rates. P values of less than 0.05 were

considered statistically significant (*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$).

Results

Deterioration of BM MSCs accompanies the development of *Nras*-mutant-induced leukemia

Mice carrying an endogenous mutant *Nras* allele develop myelodysplastic/myeloproliferative neoplasms (MDS/MPN)-like leukemia with a long latency[27-29,43]. Here we found the primary BM leukemic cells failed to accelerate the disease in the secondary recipient mice, implying a role of the BM microenvironment in disease etiology (Figure S1). We hypothesized that the BM microenvironment is impaired by *Nras*-mutant leukemic cells, which in return impedes normal hematopoiesis and accelerates leukemia progression. Indeed, we observed quantitative decreases and functional degeneration of MSCs (Ter119$^-$CD45$^-$CD31$^-$Sca1$^+$CD51$^+$CD146$^+$) during disease development and progression (Figure 1A-C, and Figure S2). To further characterize the residual MSCs in mice with leukemia, we performed RNA-Seq analysis of the residual MSCs from leukemia-bearing mice at an early disease phase (CD11b$^+$% in PB: 35%-45%). Consistent with the quantitative and functional reduction, the expression of the transcription factor *Gnl3*[36], an indicator of MSC self-renewal, was significantly down-regulated in MSCs from leukemia-bearing mice relative to wild-type mice (Figure 1D). The expression of *Nt5e* (CD73)*, Thy1* (CD90)*, Vcam1* (CD106)*, Cd81, Sdc4, Itgb1* and *Anpep*[36-38], encoding surface markers on three-lineage-potent MSCs but not on uni-lineage-primed MSCs, was markedly reduced in MSCs from leukemia-bearing mice (padj < 0.05, fold change > 1.6) (Figure 1D). Furthermore, the expression of *Bgn*, *Bmp4, Col1a1, Csf1, Dcn, Dkk2, Mmp13, Ogn, Wisp1*, and *Wisp2*[36,39], pivotal for osteogenic differentiation, was markedly suppressed in the residual MSCs (padj < 0.05, fold change > 2) (Figure 1E). *In vitro* differentiation assay confirmed that osteogenic and adipogenic differentiation potential in leukemic MSCs were impeded (Figure S3). MSCs fulfill their tissue-specific and condition-responsive regulatory functions through secreting distinct types of soluble factors[44]. Under leukemia condition, the residual MSCs indeed secreted much less soluble factors, including

Il6, Il11, Ccl2, Ccl7, Cxcl12, Cxcl13 and *Cxcl14* (padj < 0.05, fold change > 2), compared to MSCs

from normal wild-type mice (Figure 1F). These molecules are pivotal for tissue repairing[45-50].

Indirect immunofluorescence assay and intracellular flow cytometry staining confirmed that the

levels of IL-6 (Figure 1G-H), CCL2 (Figure 1I-J), and CXCL12 (Figure 1K-L) proteins in

leukemic MSCs were significantly lower than healthy MSCs (p < 0.001). In addition, *Ccl5,* a

chemokine involved in the pathogenesis of MPN[51] and the inhibition of thrombopoiesis[52], was

significantly up-regulated in the residual MSCs from leukemia-bearing mice (padj < 0.05, fold

change > 2). Gene set enrichment analysis (GSEA) further revealed features of inflammation in

the residual MSCs (Figure S4). Collectively, these results show that the MSCs dramatically

deteriorate during the disease development and progression of *Nras*-mutation-caused leukemia.

Intra-BM transfusion of healthy MSCs improves thrombopoiesis, reduces tumor burden and

improves survival of the leukemia-bearing mice

We hypothesized that restoration of the impaired BM microenvironment in leukemia-bearing mice

might suppress/delay the disease progression. To test this hypothesis, we attempted healthy MSC

treatment using GFP-tagged MSCs isolated from the tibias and femurs of healthy mice as

previously reported[53]. The isolated primary MSCs were expanded shortly *in vitro* to passage two

(P2) and cryopreserved. For MSC treatment, the cryopreserved P2 MSCs were recovered and

cultured   for   five   days,   phenotypically   identified   (CD45$^-$Ter119$^-$CD31$^-$

CD51$^+$CD105$^+$LepR$^+$PDGFRα$^+$PDGFRβ$^+$Sca1$^+$) (Figure S5), and suspended in DPBS (2.5 ×

10^7/ml) for transfusion. Initially, we adopted a direct delivery procedure by injecting donor

MSCs every two weeks either via tail vein (dose: 0.5 × 10^6/mouse) (Figure S6A) or retro-orbital

(dose: 0.5 × 10^6/mouse) transfusion (Figure S6B) into the leukemia-bearing mice at a late disease

phase (CD11b$^+$ cells > 60% in PB). However, these delivery approaches failed to produce

therapeutic effects. *In vitro* cultured MSCs lose their natural homing feature[54], the retro-orbital and

intravenous injection of cultured MSCs failed to home to bone marrow (Figure S6C-D). Thus, we

attempted intra-BM transfusion to overcome the homing defect caused by *in vitro* culture. A

sequential doses of MSCs (2.5×10^7 MSCs/kg per dose in 20 μL DPBS) were injected into the

tibia cavities of leukemia-bearing mice with two-week intervals for up to 16 weeks (Figure 2A).

Strikingly, the tumor burden continuously decreased during MSC treatment (Figure 2B and

Supplementary Figure S7). Consequently, the survival of treated leukemia-bearing mice was

significantly prolonged (Untreated: 51.5 days, MSC-treated: >115 days, $p < 0.001$) (Figure 2C).

Therefore, intra-BM transfusion of healthy donor MSCs improves the survival of leukemia-

bearing mice.

MSC-treatment systemically re-balances myelopoiesis and activates megakaryopoiesis

We next investigated the underlying mechanisms associated with the systemically decreased tumor

burden. We found that the hematopoiesis in the MSC-treated leukemia-bearing mice was re-

balanced, demonstrated by significant decreases of white blood cells (Untreated vs. MSC-treated:

23.04 vs 8.876, $p = 0.009$), and significant elevation of platelets (Untreated vs. MSC-treated: 2.64

vs. 6.01, $p = 0.004$) (Figure 2D) in PB. On the contrary, the PBS-treated leukemia-bearing mice

exhibited neither improved hematopoiesis (Figure S8) nor prolonged survival. High GM-CSF

levels in serum are associated with the tumor burdens of CMML in patients[55] and mouse models[27].

We analyzed the GM-CSF levels in serum of leukemia-bearing mice with or without MSC-

treatment. As expected, the GM-CSF levels in MSC-treated leukemia-bearing mice were

significantly decreased (> 7 folds) ($p < 0.001$) (Figure 2E), which was associated with the reduction

of GM-CSF secreting tumor cells (Figure S9). IL-6 promotes thrombopoiesis by increasing

systemic TPO levels[56]. Consistently, the MSC-treated leukemia-bearing mice exhibited elevated

IL-6 (> 25 folds) and TPO (> 2 folds) levels (Figure 2F) in PB serum. Collectively, these results

indicate that intra-BM transfusion of healthy donor MSCs systemically improves hematopoiesis

and prolongs the survival of leukemia-bearing mice.

To further investigate the systemic effects of the local MSC-treatment on hematopoiesis in

leukemia-bearing mice, we analyzed the ratios of myeloid progenitor subpopulations in MSC-

treated leukemia-bearing mice. Consistent with the elevated platelet levels and reduced myeloid

cells in PB, the MSC-treated leukemia-bearing mice showed increased proportions of

megakaryocyte-erythroid progenitors (MEP) (> 1.6 folds) (p < 0.001) and decreased ratios of

granulocyte-macrophage progenitors (GMP) (> 1.5 folds) (p < 0.001) in both injected and non-

injected sites than those sites in PBS-treated leukemia-bearing mice (Figure 3A-B, Figure S10A).

In addition, we observed increased (> 1.3 folds) ratios of mature megakaryocytes ($\geq$ 8N) in both

injected and non-injected sites in MSC-treated leukemia-bearing mice (Figure 3C-D, Figure S10B)

in comparison with PBS-treated leukemia-bearing control mice (p < 0.001). Thus, these data

demonstrate that MSC-treatment systemically re-balances myelopoiesis and activates

megakaryopoiesis in leukemia-bearing mice.

Recovered host MSCs are functional as healthy counterparts

Consistent with an early report[57], our data showed that MSCs indeed support normal

hematopoiesis. Interestingly, MSCs also promoted the growth of leukemic cells *in vitro*, which

suggests that donor MSCs indirectly inhibit the tumor development *in vivo*. To investigate whether

the improved hematopoiesis is associated with restoration of the BM microenvironment, we

analyzed the MSC-treated tibias eight weeks after MSC treatment. Interestingly, host MSCs (GFP

negative) were partially recovered (Figure 4A-B), but restricted to the locally treated tibias (Figure

S11). Functionally, the recovered host MSCs formed markedly more CFU-F colonies than the

residual MSCs from untreated leukemia-bearing mice (> 3.8 folds) (p < 0.001) (Figure 4C and

Figure S12). To characterize the recovered MSCs at the transcriptome level, we sorted the

recovered MSCs for RNA-Seq analysis. Unsupervised hierarchical clustering analysis showed that

the recovered MSCs clustered closer to healthy MSCs (Figure 4D). Further, the expression of

cytokines and chemokines, including *Il6, Ccl2, Ccl7, Ccl19, Cxcl12, Cxcl13,* and *Cxcl14*, was

restored in the recovered MSCs compared to that in MSCs from untreated leukemia-bearing

control mice (padj < 0.05, fold change > 2) (p < 0.05) (Figure 4E). Consistently, indirect

immunofluorescence assay and intracellular flow cytometry staining confirmed that the levels of

IL-6, CCL2, and CXCL12 proteins in recovered MSCs (from the injected sites) were significantly

improved to comparable levels as in healthy MSCs (p < 0.001) (Figure 4F-I and S13). Therefore,

donor MSC-treatment results in local functional restoration of host MSCs.

The donor MSCs reprogram macrophages to execute tissue-repair function

We further investigated the cellular mechanism underlying the restored BM microenvironment

mediated by donor MSCs under leukemia condition. BM macrophages play a pivotal role in

maintaining the BM niche[18]. To study whether donor MSCs reprogram BM macrophages, we

performed co-culture assay of healthy MSCs with BM macrophages (L-Mac) sorted from the

leukemia-bearing mice *in vitro* for twelve hours and re-sorted the macrophages (E-Mac) for RNA-

Seq analysis. GSEA illustrated that angiogenesis-related genes, including *Vegfa, Hif1a, Serpine1,*

Eng and Thbs1[40] (Figure S14A), were enriched among the differentially expressed genes in E-Mac

(Figure 5A). Genes associated with cell migration, including *Sirpa* and *Ccl5*[58,59], were also

enriched in E-Mac (Figure 5B and S14B). Further, gene-ontology analysis demonstrated features

of positive regulation of cell migration and angiogenesis in E-Mac (Figure 5C). Elevated

expression of genes encoding soluble factors, including *Tnf*, *Il1a*, *Ccr7*, *Ccl3*, and *Ccl5*, was also

observed in E-Mac (Figure 5D). Indirect immunofluorescence assay confirmed that E-Mac had

significantly higher levels of TNF-α, CCR7, and IL-1α proteins than L-Mac ($p < 0.001$) (Figure

5E and S15A). Furthermore, intracellular flow cytometry staining also showed that the levels of

TNF-α and CCR7 proteins in L-Mac were significantly lower than E-Mac ($p < 0.001$) (Figure

S15B-D). To characterize which subset of macrophages are involved in re-educating the bone

marrow microenvironment, we analyzed the classical M1 and M2 macrophage subtypes both *in*

vitro and *in vivo*. However, the results showed that MSC treatment caused no typical transition of

M1 and M2 (Figure S16), which indicates a functional alteration of leukemic macrophages

independent of M1 to M2 transition. RNA-Seq analysis showed that the expression of arginase 1

(*Arg1*), an indicator of tissue repair function[24], was dramatically up-regulated over thousand folds

in E-Mac (Figure 5F) after direct co-culture with MSCs *in vitro*. However, the *Arg1* expression in

macrophages after transwell co-culture was barely elevated (Figure 5F), indicating that direct cell-

cell interaction instead of MSC-secreted soluble factors is essential for the functional

reprogramming. Consistent with the observation *in vitro*, the expression of *Arg1* was also

significantly increased in BM macrophages directly isolated from MSC-treated leukemia-bearing

mice (Figure 5G). Furthermore, intracellular flow cytometry staining confirmed that the ratios of

Arg1^{high} macrophage subpopulation significantly increased in E-Mac at the MSC-treated sites (p

< 0.01) (Figure 5H-I). Collectively, these results indicate that the donor MSCs reprogram BM

macrophages from leukemia-bearing mice to execute tissue-repair function.

The E-Mac treatment largely recapitulates the therapeutic effects of MSC treatment

Given the short lifespan of donor MSCs *in vivo*[17], we speculated that MSCs mediate the restoration

of the BM microenvironment of leukemia-bearing mice by reprogramming macrophages. We

isolated macrophages from leukemia-bearing mice and co-cultured them with healthy MSCs for

12h, and then transplanted these E-Mac back into leukemia-bearing mice by intra-BM injection

(Figure 6A). We indeed found that the thrombopoiesis was significantly improved (> 6 folds) after

E-Mac treatment (p < 0.001) (Figure 6B-C). Host MSCs were also significantly increased (> 3

folds) in E-Mac-treated leukemia-bearing mice (p < 0.001) (Figure 6D-E). Consistent with the

MSC treatment, we also observed increased ratios of mature megakaryocytes (p < 0.001) (Figure

6F-G) and alleviated tumor burden (CD11b$^+$) (Figure S17) in E-Mac-treated leukemia-bearing

mice. Collectively, these results demonstrate that MSC-reprogrammed macrophages largely

recapitulate the therapeutic effects of MSCs.

MSCs reprogram macrophages and reduce leukemia burden through IL-6

Residual MSCs in leukemia-bearing mice lost the ability to secrete IL-6 (Figure 1G-H). IL-6 is

critical for maintaining the stemness and function of MSCs[60]. The recovered host MSCs after

donor MSC-treatment expressed comparable *Il6* mRNA as healthy MSCs (Figure 4E-G).

Consistently, BM plasma IL-6 levels in MSC-treated leukemia-bearing mice were significantly

elevated (> 3 folds) and were comparable to those in healthy mice (p < 0.001) (Figure 7A). To

investigate the biological consequence of reduced IL-6 on MSCs, we analyzed the MSCs in *Il6*$^{-/-}$

mice by *in vitro* proliferation assay. We observed that the *Il6*$^{-/-}$ MSCs proliferated at a significantly

slower speed than their WT counterparts *in vitro* (p < 0.001) (Figure. S18A). To investigate the

role of IL-6 in MSC-mediated therapeutic effects, we directly injected IL-6 proteins (40 μg/kg)

into leukemia-bearing mice. However, systemic IL-6 transfusion failed to suppress leukemia

(Figure S18B), which indicates that MSC-educated leukemic macrophages are distinct from the

conventional IL-6-dependent myeloid-derived immune-suppressor cells[61]. We speculated that IL-

6 might function as an autocrine factor for MSCs to reprogram macrophages in leukemia-bearing

mice. *In vitro* co-culture assay showed that *Il6*$^{-/-}$ MSCs compromised their ability to induce *Arg1*

expression in macrophages derived from leukemia-bearing mice (Figure 7B). In addition, intra-

BM injection of *Il6*[-/-] MSCs neither improved thrombopoiesis nor reduced tumor burden in

leukemia-bearing mice (Figure 7C-E). Furthermore, loss of *Il6* significantly compromised MSCs'

ability to reprogram leukemic macrophages, demonstrated by mild increase of the ratios of Arg1[high]

macrophage subpopulation in *Il6*[-/-] MSC-treated leukemic mice *in vivo* (Figure 7F-H). Taken

together, these results demonstrate that autocrine IL-6 is essential for MSC-mediated

reprogramming of macrophages and reduction of tumor burden in leukemia-bearing mice.

Discussion

Deteriorating BM microenvironment accompanies chronic leukemia progression. Here we unravel

a *de novo* approach of reverting the impaired BM microenvironment by intra-BM injection of

donor MSCs. Upon injection, the donor MSCs quickly reprogrammed local host BM macrophages

to repair the niche, thus improving normal hematopoiesis and suppressing leukemia development.

These effects of donor MSCs depend on the autocrine production of IL-6. Our studies reveal de

novo mechanisms underlying MSC-mediated local BM microenvironment restoration that

systemically suppress leukemia development.

Given the short-term lifespan of the exogenous MSC *in vivo*, it is surprising that local injection of

donor MSCs results in long-term improvement of thrombopoiesis and reduction of tumor burden.

Following injection, exogenous donor MSCs immediately reprogram host resident macrophages

that further organize the overhaul of local BM microenvironment, including restoring the functions

of host MSCs. There is a lot of evidence supporting the pivotal roles of macrophages in tissue

repair[62,63]. Donor MSCs can transiently release a key wave of tissue-repair factors, such as IL-6[45],

CCL7[50], and CXCL12[47], and reprogram host macrophages, subsequently resulting in the recovery

of host MSCs. Recovered host MSCs further secreted much higher level of CCL2, CCL7, and

CXCL12 that can further facilitate BM niche repair. Donor MSCs could also directly modulate the

other niche cells, in addition to macrophages, to participate in BM niche repair[12]. Consequently,

the restored local BM microenvironment outputs abundant hematopoiesis-improving cytokines,

including IL-6[56], and reduces tumor-growth-stimulating cytokines, such as GM-CSF[27,55]. Thus,

despite the short life-span, donor MSCs provide long-term thrombopoiesis improvement and

tumor burden reduction through the stepwise microenvironment restoration.

Of note, IL-6 deficiency in MSCs markedly compromised the abilities of these cells to reprogram

host resident macrophage. Transwell co-culture study has shown that direct interaction between

MSCs and macrophages is a necessity for reprogramming host macrophages. Both autocrine

production of IL-6 and reprogramming host macrophages by MSCs are required for MSC-

mediated microenvironment restoration and leukemia inhibition. However, it is not clear how

autocrine IL-6 controls the function of MSCs and how MSCs directly reprogram macrophages.

The underlying molecular mechanisms warrant further investigation. Besides IL-6, it deserves

additional investigations of whether other key molecules produced by MSCs are involved in the

MSC-mediated microenvironment restoration and leukemia inhibition, such as nitric oxide[64].

MSC treatment inhibits leukemia development in the *Nras* mutation-induced MPN/MDS-like

disease model. We also attempted to broaden MSC treatment for acute leukemia in the MLL-AF9-

initiated model (Figure S19A), in which impaired MSCs results in the reduction of osteogenesis

and CXCL12 production[65]. Despite a mild elevation of platelet level, the intra-BM transfusion of

donor MSCs failed to significantly improve normal hematopoiesis or suppress acute leukemia

development (Figure S19B-H). Therefore, the intra-BM MSC treatment might be beneficial for

MPN/MDS leukemias, such as JMML and CMML, but insufficient for suppressing acute

leukemia. Although MSC application could be an effective therapeutic regimen for patients with

MPN/MDS-subtype leukemia, combination therapy of conventional approaches with local MSC

transfusion might be required to achieve therapeutic outcomes for acute leukemia with impaired

BM microenvironment.

Acknowledgments

This work was supported by grants from the Strategic Priority Research Program of Chinese

Academy of Sciences (XDA16010601), the Chinese Ministry of Science and Technology

(2015CB964401,   2016YFA0100601,   2016YFA0100600,   2017YFA0103401,   and

2015CB964902), the Major Research and Development Project of Guangzhou Regenerative

Medicine and Health Guangdong Laboratory (2018GZR110104006, 2018GZR0201008), the CAS

Key Research Program of Frontier Sciences (QYZDB-SSW-SMC057), the Health and Medical

Care   Collaborative   Innovation   Program   of   Guangzhou   Scientific   and   Technology

(201803040017), CAMS Innovation Fund for Medical Sciences (2016-12M-1-002), the General

Program from Guangzhou Scientific and Technological Project (201707010157), the Science and

Technology Planning Project of Guangdong Province (2017B030314056, 2017B020230004), the

grants from the National Natural Science Foundation of China (Grant No 31471117, 31271457,

81470281, 81421002, 81730006, 31600948, and 81861148029), the CAMS Initiative for

Innovative Medicine (2016-I2M-1-017) and the grants from NIH, USA (AI079087, D.W. and

HL130724 D.W.)

Author Contributions

C.X.X. performed research, analyzed data and wrote the manuscript; D.Y. and Q.T.W. analyzed

RNA-Seq data; T.J.W., H.C., P.Q.Z., K.T.W., X.F.L., Y.G., S.H.M., L.X. and Y.X.G. performed

experiments; S.H., J.D., X.D., Y.Q.L., X.F.Z., Y.F.S. and S.X. discussed the manuscript; D.W.

discussed the project and wrote the manuscript. T.C. and J.Y.W. designed the research and wrote

the manuscript.

Conflict of Interest Disclosure

The authors declare no competing financial interests.

References

1.   Kfoury Y, Scadden DT. Mesenchymal cell contributions to the stem cell niche. *Cell Stem Cell*.
2015;16(3):239-253.

2.   Lim M, Pang Y, Ma S, et al. Altered mesenchymal niche cells impede generation of normal
hematopoietic progenitor cells in leukemic bone marrow. *Leukemia*. 2016;30(1):154-162.

3.   Raaijmakers MH, Mukherjee S, Guo S, et al. Bone progenitor dysfunction induces myelodysplasia
and secondary leukaemia. *Nature*. 2010;464(7290):852-857.

4.   Schepers K, Pietras EM, Reynaud D, et al. Myeloproliferative neoplasia remodels the endosteal
bone marrow niche into a self-reinforcing leukemic niche. *Cell Stem Cell*. 2013;13(3):285-299.

5.   Bowers M, Zhang B, Ho Y, Agarwal P, Chen CC, Bhatia R. Osteoblast ablation reduces normal
long-term hematopoietic stem cell self-renewal but accelerates leukemia development. *Blood*.
2015;125(17):2678-2688.

6.   Dong L, Yu WM, Zheng H, et al. Leukaemogenic effects of Ptpn11 activating mutations in the
stem cell microenvironment. *Nature*. 2016;539(7628):304-308.

7.   Blau O, Baldus CD, Hofmann WK, et al. Mesenchymal stromal cells of myelodysplastic syndrome
and acute myeloid leukemia patients have distinct genetic abnormalities compared with leukemic blasts.
Blood. 2011;118(20):5583-5592.

8.   Pandis N, Bardi G, Sfikas K, Panayotopoulos N, Tserkezoglou A, Fotiou S. Complex chromosome
rearrangements involving 12q14 in two uterine leiomyomas. *Cancer Genet Cytogenet*. 1990;49(1):51-56.

9.   Medyouf H, Mossner M, Jann JC, et al. Myelodysplastic cells in patients reprogram mesenchymal

stromal cells to establish a transplantable stem cell niche disease unit. *Cell Stem Cell*. 2014;14(6):824-837.

10.   Ren G, Zhang L, Zhao X, et al. Mesenchymal stem cell-mediated immunosuppression occurs via

concerted action of chemokines and nitric oxide. *Cell Stem Cell*. 2008;2(2):141-150.

11.   Prockop DJ. Inflammation, fibrosis, and modulation of the process by mesenchymal stem/stromal

cells. *Matrix Biol*. 2016;51:7-13.

12.   Shi Y, Wang Y, Li Q, et al. Immunoregulatory mechanisms of mesenchymal stem and stromal cells

in inflammatory diseases. *Nat Rev Nephrol*. 2018;14(8):493-507.

13.   Le Blanc K, Mougiakakos D. Multipotent mesenchymal stromal cells and the innate immune

system. *Nat Rev Immunol*. 2012;12(5):383-396.

14.   Mittal M, Tiruppathi C, Nepal S, et al. TNFalpha-stimulated gene-6 (TSG6) activates macrophage

phenotype transition to prevent inflammatory lung injury. *Proc Natl Acad Sci U S A*. 2016;113(50):E8151-

E8158.

15.   Wang G, Cao K, Liu K, et al. Kynurenic acid, an IDO metabolite, controls TSG-6-mediated

immunosuppression of human mesenchymal stem cells. *Cell Death Differ*. 2018;25(7):1209-1223.

16.   Du L, Lin L, Li Q, et al. IGF-2 Preprograms Maturing Macrophages to Acquire Oxidative

Phosphorylation-Dependent Anti-inflammatory Properties. *Cell Metab*. 2019.

17.   Eggenhofer E, Benseler V, Kroemer A, et al. Mesenchymal stem cells are short-lived and do not

migrate beyond the lungs after intravenous infusion. *Front Immunol*. 2012;3:297.

18.   Ehninger A, Trumpp A. The bone marrow stem cell niche grows up: mesenchymal stem cells and

macrophages move in. *J Exp Med*. 2011;208(3):421-428.

19.   Chen J, Yao Y, Gong C, et al. CCL18 from tumor-associated macrophages promotes breast cancer

metastasis via PITPNM3. *Cancer Cell*. 2011;19(4):541-555.

20.   Gubin MM, Esaulova E, Ward JP, et al. High-Dimensional Analysis Delineates Myeloid and

Lymphoid Compartment Remodeling during Successful Immune-Checkpoint Cancer Therapy. *Cell*.

2018;175(4):1014-1030 e1019.

21. Chen CC, Wang L, Plikus MV, et al. Organ-level quorum sensing directs regeneration in hair stem

cell populations. *Cell*. 2015;161(2):277-290.

22. Liu C, Wu C, Yang Q, et al. Macrophages Mediate the Repair of Brain Vascular Rupture through

Direct Physical Adhesion and Mechanical Traction. *Immunity*. 2016;44(5):1162-1176.

23. Winkler IG, Sims NA, Pettit AR, et al. Bone marrow macrophages maintain hematopoietic stem

cell (HSC) niches and their depletion mobilizes HSCs. *Blood*. 2010;116(23):4815-4828.

24. Bosurgi L, Cao YG, Cabeza-Cabrerizo M, et al. Macrophage function in tissue repair and

remodeling requires IL-4 or IL-13 with apoptotic cells. *Science*. 2017;356(6342):1072-1076.

25. Cho DI, Kim MR, Jeong HY, et al. Mesenchymal stem cells reciprocally regulate the M1/M2

balance in mouse bone marrow-derived macrophages. *Exp Mol Med*. 2014;46:e70.

26. Nemeth K, Leelahavanichkul A, Yuen PS, et al. Bone marrow stromal cells attenuate sepsis via

prostaglandin E(2)-dependent reprogramming of host macrophages to increase their interleukin-10

production. *Nat Med*. 2009;15(1):42-49.

27. Wang J, Liu Y, Li Z, et al. Endogenous oncogenic Nras mutation promotes aberrant GM-CSF

signaling in granulocytic/monocytic precursors in a murine model of chronic myelomonocytic leukemia.

Blood. 2010;116(26):5991-6002.

28. Wang J, Liu Y, Li Z, et al. Endogenous oncogenic Nras mutation initiates hematopoietic

malignancies in a dose- and cell type-dependent manner. *Blood*. 2011;118(2):368-379.

29. Wang JY, Kong GY, Liu YG, et al. Nras(G12D/+) promotes leukemogenesis by aberrantly

regulating hematopoietic stem cell functions. *Blood*. 2013;121(26):5203-5207.

30. Tang F, Barbacioru C, Nordman E, et al. RNA-Seq analysis to capture the transcriptome landscape

of a single cell. *Nat Protoc*. 2010;5(3):516-535.

31. Kim D, Langmead B, Salzberg SL. HISAT: a fast spliced aligner with low memory requirements.

Nat Methods. 2015;12(4):357-360.

32. Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for assigning

sequence reads to genomic features. *Bioinformatics*. 2014;30(7):923-930.

33.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq

data with DESeq2. *Genome Biol*. 2014;15(12):550.

34.   Subramanian A, Tamayo P, Mootha VK, et al. Gene set enrichment analysis: a knowledge-based

approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A*. 2005;102(43):15545-

15550.

35.   Yu G, Wang LG, Han Y, He QY. clusterProfiler: an R package for comparing biological themes

among gene clusters. *OMICS*. 2012;16(5):284-287.

36.   Freeman BT, Jung JP, Ogle BM. Single-Cell RNA-Seq of Bone Marrow-Derived Mesenchymal

Stem Cells Reveals Unique Profiles of Lineage Priming. *PLoS One*. 2015;10(9):e0136199.

37.   Song L, Webb NE, Song Y, Tuan RS. Identification and functional analysis of candidate genes

regulating mesenchymal stem cell self-renewal and multipotency. *Stem Cells*. 2006;24(7):1707-1718.

38.   Rostovskaya M, Anastassiadis K. Differential expression of surface markers in mouse bone marrow

mesenchymal   stromal   cell   subpopulations   with   distinct   lineage   commitment.   *PLoS   One*.

2012;7(12):e51221.

39.   Delorme B, Ringe J, Pontikoglou C, et al. Specific lineage-priming of bone marrow mesenchymal

stem cells provides the molecular framework for their plasticity. *Stem Cells*. 2009;27(5):1142-1151.

40.   Medina RJ, O'Neill CL, O'Doherty TM, et al. Myeloid angiogenic cells act as alternative M2

macrophages and modulate angiogenesis through interleukin-8. *Mol Med*. 2011;17(9-10):1045-1055.

41.   Arango Duque G, Descoteaux A. Macrophage cytokines: involvement in immunity and infectious

diseases. *Front Immunol*. 2014;5:491.

42.   Cavaillon JM. Cytokines and macrophages. *Biomed Pharmacother*. 1994;48(10):445-453.

43.   Li Q, Haigis KM, McDaniel A, et al. Hematopoiesis and leukemogenesis in mice expressing

oncogenic NrasG12D from the endogenous locus. *Blood*. 2011;117(6):2022-2032.

44.   Gnecchi M, Danieli P, Malpasso G, Ciuffreda MC. Paracrine Mechanisms of Mesenchymal Stem

Cells in Tissue Repair. *Methods Mol Biol*. 2016;1416:123-146.

45.   Gallucci RM, Simeonova PP, Matheson JM, et al. Impaired cutaneous wound healing in

interleukin-6-deficient and immunosuppressed mice. *FASEB J*. 2000;14(15):2525-2531.

46.   Castela M, Nassar D, Sbeih M, Jachiet M, Wang Z, Aractingi S. Ccl2/Ccr2 signalling recruits a

distinct fetal microchimeric population that rescues delayed maternal wound healing. *Nat Commun*.

2017;8:15463.

47.   Kato T, Khanh VC, Sato K, et al. SDF-1 improves wound healing ability of glucocorticoid-treated

adipose tissue-derived mesenchymal stem cells. *Biochem Biophys Res Commun*. 2017;493(2):1010-1017.

48.   Hayashi Y, Murakami M, Kawamura R, Ishizaka R, Fukuta O, Nakashima M. CXCL14 and MCP1

are potent trophic factors associated with cell migration and angiogenesis leading to higher regenerative

potential of dental pulp side population cells. *Stem Cell Res Ther*. 2015;6:111.

49.   Sims NA, Jenkins BJ, Nakamura A, et al. Interleukin-11 receptor signaling is required for normal

bone remodeling. *J Bone Miner Res*. 2005;20(7):1093-1102.

50.   Schenk S, Mal N, Finan A, et al. Monocyte chemotactic protein-3 is a myocardial mesenchymal

stem cell homing factor. *Stem Cells*. 2007;25(1):245-251.

51.   Kleppe M, Kwak M, Koppikar P, et al. JAK-STAT pathway activation in malignant and

nonmalignant cells contributes to MPN pathogenesis and therapeutic response. *Cancer Discov*.

2015;5(3):316-331.

52.   Pang L, Weiss MJ, Poncz M. Megakaryocyte biology and related disorders. *J Clin Invest*.

2005;115(12):3332-3338.

53.   Zhu H, Guo ZK, Jiang XX, et al. A protocol for isolation and culture of mesenchymal stem cells

from mouse compact bone. *Nat Protoc*. 2010;5(3):550-560.

54.   Rombouts WJ, Ploemacher RE. Primary murine MSC show highly efficient homing to the bone

marrow but lose homing ability following culture. *Leukemia*. 2003;17(1):160-170.

55.   Padron E, Painter JS, Kunigal S, et al. GM-CSF-dependent pSTAT5 sensitivity is a feature with

therapeutic potential in chronic myelomonocytic leukemia. *Blood*. 2013;121(25):5068-5077.

56. Kaser A, Brandacher G, Steurer W, et al. Interleukin-6 stimulates thrombopoiesis through

thrombopoietin: role in inflammatory thrombocytosis. *Blood*. 2001;98(9):2720-2725.

57. de Lima M, McNiece I, Robinson SN, et al. Cord-blood engraftment with ex vivo mesenchymal-

cell coculture. *N Engl J Med*. 2012;367(24):2305-2315.

58. Ohnishi H, Kobayashi H, Okazawa H, et al. Ectodomain shedding of SHPS-1 and its role in

regulation of cell migration. *J Biol Chem*. 2004;279(27):27878-27887.

59. Hocking AM. The Role of Chemokines in Mesenchymal Stem Cell Homing to Wounds. *Adv

Wound Care (New Rochelle)*. 2015;4(11):623-630.

60. Pricola KL, Kuhn NZ, Haleem-Smith H, Song Y, Tuan RS. Interleukin-6 maintains bone marrow-

derived mesenchymal stem cell stemness by an ERK1/2-dependent mechanism. *J Cell Biochem*.

2009;108(3):577-588.

61. Marigo I, Bosio E, Solito S, et al. Tumor-induced tolerance and immune suppression depend on

the C/EBPbeta transcription factor. *Immunity*. 2010;32(6):790-802.

62. Wynn TA, Vannella KM. Macrophages in Tissue Repair, Regeneration, and Fibrosis. *Immunity*.

2016;44(3):450-462.

63. Liu C, Wu CA, Yang QF, et al. Macrophages Mediate the Repair of Brain Vascular Rupture through

Direct Physical Adhesion and Mechanical Traction. *Immunity*. 2016;44(5):1162-1176.

64. Trento C, Marigo I, Pievani A, et al. Bone marrow mesenchymal stromal cells induce nitric oxide

synthase-dependent differentiation of CD11b(+) cells that expedite hematopoietic recovery.

Haematologica. 2017;102(5):818-825.

65. Hanoun M, Zhang D, Mizoguchi T, et al. Acute myelogenous leukemia-induced sympathetic

neuropathy promotes malignancy in an altered hematopoietic stem cell niche. *Cell Stem Cell*.

2014;15(3):365-375.

Figures and figure legends

Figure 1

Figure 1: Impaired BM MSCs in mice with NrasG12D mutation-induced leukemia

CD45.2$^+$ BMNC from LSL *Nras*/+; *Vav-Cre* mice (NV mice) were sorted and transplanted into

sublethally irradiated (6.5 Gy) individual recipients (CD45.1 strain) with a cell dose of 0.3 million

per recipient. For control recipient mice (CD45.1 strain), 0.3 million sorted CD45.2$^+$ BMNC from

WT mice were transplanted. **(A)** Gating strategies for BM MSCs. MSCs are defined as Ter119[-]

CD45[-]CD31[-]Sca1[+]CD51[+]CD146[+]. Plots from one representative control mouse (CD11b[+]% in PB

= 10%) and one leukemia-bearing mouse (CD11b[+]% in PB > 60%) of six mice of each group are

shown. The nucleated cell mixtures of BM and compact bones were prepared for flow cytometry

analysis of MSCs. **(B)** Statistical analysis of the absolute numbers of MSCs in tibias and femurs

from control and leukemia-bearing mice. Data are analyzed by unpaired student's t-test (two-

tailed). ***$p < 0.001$. Data are represented as mean ± SD (n = 6 mice for each group from multiple

independent experiments). **(C)** Kinetic analysis of functional MSCs in CFU-F assay. Total BMNC

equivalent to 100 MSCs calculated by the percentages of MSCs in individual BMNC samples by

flow cytometry analysis. Data are analyzed by unpaired student's t-test (two-tailed). *$p < 0.05$,

***$p < 0.001$. For each time point, three independent experiments were performed. Data are

represented as mean ± SD. **(D)** Heatmaps of MSC stemness-related genes differentially expressed

between healthy MSCs and MSCs from leukemia-bearing mice (padj < 0.05, fold change > 1.6).

One thousand sorted MSCs (Ter119[-]CD45[-]CD31[-]Sca1[+]CD51[+]) from the nucleated cell mixtures

of BM and compact bones of independent leukemia-bearing mice, control mice, and MSC-treated

leukemia-bearing mice were used as each cell sample input for RNA-Seq. Leukemia-bearing mice

(CD11b[+]% in PB = 35%-45%) and control mice (CD11b[+]% in PB = 10-15%) were accumulated

from multiple independent experiments. The expression value (DESeq2 normalized counts) of

each gene was converted to z-score values (red, high; blue, low), and the heatmaps were plotted

by gplots (heatmap.2). Columns represent the indicated cell subsets in nine MSC samples (Healthy

MSCs from control mice: n = 3, MSCs from leukemia-bearing mice: n = 6). **(E)** Heatmaps of

osteogenic differentiation-related genes differentially expressed between healthy MSCs and MSCs

from leukemia-bearing mice (padj < 0.05, fold change > 2). The expression value (DESeq2

normalized counts) of each gene was converted to z-score values (red, high; blue, low), and the

heatmaps were plotted by gplots (heatmap.2). Columns represent the indicated cell subsets in nine

MSC samples (Healthy MSCs: n = 3, MSCs from leukemia-bearing mice: n = 6). **(F)** Heatmaps of

MSC secreted factor-related genes differentially expressed between healthy MSCs and MSCs from

leukemia-bearing mice (padj < 0.05, fold change > 2). The expression value (DESeq2 normalized

counts) of each gene was converted to z-score values (red, high; blue, low), and the heatmaps were

plotted by gplots (heatmap.2). Columns represent the indicated cell subsets in nine MSC samples

(Healthy MSCs: n = 3, MSCs from leukemia-bearing mice: n = 6). **(G)** Single-cell imaging of

intracellular IL-6 proteins by indirect immunofluorescence assay (IFA) in primary MSCs sorted

from the control and leukemic mice. Images of three single representative cells of each group are

shown. **(H)** Statistical analysis of mean intensities of IL-6 fluorescence in control and leukemic

MSC samples. Each dot represents a single cell. Data are analyzed by unpaired student's t-test

(two-tailed). ***p < 0.001. Data are represented as mean ± SD. Control, n=30; Leukemia, n=30.

(I) Single-cell imaging of intracellular CCL2 proteins by IFA in primary MSCs sorted from the

control and leukemic mice. Images of three single representative cells of each group are shown.

(J) Statistical analysis of mean intensities of CCL2 fluorescence in control and leukemic MSC

samples. Each dot represents a single cell. Data are analyzed by unpaired student's t-test (two-

tailed). ***p < 0.001. Data are represented as mean ± SD. Control, n=30; Leukemia, n=30. **(K)**

Representative intracellular staining plots of CXCL12 proteins in control and leukemic MSCs

analyzed by flow cytometry. **(L)** Statistical analysis of mean fluorescence intensities (MFI) of

CXCL12 in control and leukemic MSCs. Data are analyzed by unpaired student's t-test (two-

tailed). ***p < 0.001. Plots from three representative control mice and leukemia-bearing mice are

shown. Data are represented as mean ± SD (n = 6 mice for each group).

Figure 2

Figure 2: Intra-BM transfusion of donor MSCs prolongs survival of leukemia-bearing mice

(A) Schematic diagram of MSC transfusion strategy. The donor MSCs prepared for transfusion were isolated from the compact bone and BMNC of three to four-week-old healthy GFP mice. The isolated MSCs were expanded *in vitro,* and the secondary passage products were used for transfusion. Leukemia-bearing mice with severe tumor burden (CD11b$^+$% in PB > 60%) were treated by intra-bone-marrow transfusion of donor MSCs. WT mice were treated by the same procedure as treatment control, and leukemia-bearing mice without MSC treatment were used as untreated controls. A sequential doses of MSCs ($2.5 \times 10^{\wedge}7$ MSCs/kg per dose in 20 μL DPBS) were delivered into the tibia cavity using 29-gauge needle. Every tibia was treated once per month by switching the injection site every other dose. **(B)** Kinetic analysis of tumor burden (CD11b$^+$) of

MSC-treated leukemia-bearing mice. The time window of MSC treatment is from 0 weeks (W0,

treatment starting time) to 16 weeks (W16). Flow cytometry analysis of tumor burden (CD11b$^+$)

in PB was performed monthly. These leukemia-bearing mice were accumulated from multiple

independent experiments. The individual leukemia-bearing mice were selected to perform

treatment once they reached the leukemic burden standard (CD11b$^+$% > 60% in peripheral blood),

but at different time from multiple experiments. Untreated leukemia-bearing mice were used as

disease control (red line), PBS-treated leukemia-bearing mice were used as injected control (green

line), and MSC-treated WT mice were used as treatment control (black line). Untreated leukemia-

bearing mice: n = 6; PBS-treated leukemia-bearing mice: n = 3; MSC-treated leukemia-bearing

mice (blue line): n = 6; MSC-treated WT mice: n = 4. **(C)** Kaplan-Meier survival of MSC-treated

leukemia-bearing mice. Kaplan-Meier survival curves of untreated (n = 6, Median survival = 51.5

days) and MSC-treated (n = 6, Median survival = 115 days) leukemia-bearing mice are shown.

MSC treatment was terminated after 16 weeks. The untreated leukemia-bearing mice from the

same batch were used as control (red line). Log-rank (Mantel-Cox) test: $p < 0.001$. **(D)** Statistical

analysis of white blood cells (WBC) and platelets (PLT) counts in PB of WT mice, untreated

leukemia-bearing mice, and MSC-treated leukemia-bearing mice at week nine since the MSC

treatment. Data are analyzed by one-way ANOVA test. **$p < 0.01$. Data are represented as mean

± SD (n =6-8 mice for each group). **(E)** ELISA of GM-CSF levels in PB serum. Serum prepared

from 200 μL PB of individual WT mice (n=8), untreated leukemia-bearing mice (Untreated, n=6)

and MSC-treated leukemia-bearing mice (MSC-treated, n=6) eight weeks post MSC treatment.

Data are analyzed by one-way ANOVA test. ***$p < 0.001$. Data are represented as mean ± SD.

(F) ELISA of IL-6 and TPO levels in PB serum. Serum prepared from 200 μL PB of individual

WT mice (n=6-8), untreated leukemia-bearing mice (Untreated, n=6) and MSC-treated leukemia-

bearing mice (MSC-treated, n=6) eight weeks post MSC treatment. Data are analyzed by one-way

ANOVA. **$p < 0.01$, ***$p < 0.001$. Data are represented as mean ± SD.

Figure 3

Figure 3: Systemically re-balanced myeloid lineage progenitor cells and systemically activated megakaryocytes in MSC-treated leukemia-bearing mice

(A) Ratios of myeloid progenitor subpopulations in MSC- and PBS-treated leukemia-bearing mice. Total bone marrow nucleated cells were from the MSC-injected site (tibia) and non-injected site (femur) of each leukemia-bearing mouse, or PBS-injected site (tibia) and non-injected site (femur) of each leukemia-bearing mouse four weeks post MSC/PBS treatment. GMP (granulocyte/macrophage progenitors): $Lin^-IL\text{-}7R^-Sca1^-c\text{-}Kit^+CD34^+CD16/32^{high}$; MEP (megakaryocyte/erythroid progenitors): $Lin^-IL\text{-}7R^-Sca1^-c\text{-}Kit^+CD34^-CD16/32^-$. **(B)** Statistical

667 analysis of myeloid progenitor components (GMP and MEP) in the injected site (tibia) and non-

668 injected site (femur) of each PBS-treated leukemia-bearing mouse and MSC-treated leukemia-

669 bearing mouse four weeks post-MSC treatment. Data are analyzed by one-way ANOVA test. ***p

670 < 0.001. Data are represented as mean ± SD (n = 5-6 mice for each group accumulated from

671 multiple independent experiments). **(C)** Activation analysis of megakaryocytes in MSC- and PBS-

672 treated leukemia-bearing mice. Plots from one representative PBS-treated leukemia-bearing

673 mouse and MSC-treated leukemia-bearing mouse four weeks post MSC treatment are shown.

674 Percentages of mature megakaryocytes with 8N and greater ploidy ($\geq 8N$) are shown. **(D)**

675 Statistical analysis of the percentages of mature megakaryocytes ($\geq 8N$). Data are analyzed by one-

676 way ANOVA test. ***$p < 0.001$, N.S indicates not-significant. Data are represented as mean ± SD

677 (n = 4 mice for each group accumulated from multiple independent experiments).

Figure 4

Figure 4: Characterization of recovered host MSCs from MSC-treated leukemia-bearing mice

(A) Flow cytometry analysis of recovered host MSCs in leukemia-bearing mice eight weeks post MSC treatment. Tibias of MSC-treated leukemia-bearing mice eight weeks after the first dose MSC treatment were analyzed. The leukemic mice are induced as described in Figure 1. The data of the MSC-treated tibias (injected sites) from MSC-treated leukemic mice and the control tibias from untreated leukemic mice are shown. Plots of one representative mouse from each group are shown. MSCs are defined as $Ter119^-CD45^-CD31^-Sca1^+CD51^+CD146^+$. The nucleated cell mixtures of BM and compact bones were prepared for flow cytometry analysis of MSCs. **(B)** Statistical analysis of the absolute numbers of host MSCs (GFP^-, host-derived MSCs) in tibias from untreated and MSC-treated leukemia-bearing mice. Data are analyzed by unpaired student's t-test (two-tailed). ***$p < 0.001$. Data are represented as mean ± SD (n = 5 mice for each group).

(C) Statistical analysis of CFU-F colonies. The numbers of colonies of each group were counted

after Giemsa staining. Data are analyzed by unpaired student's t-test (two-tailed). ***$p < 0.001$.

Data are represented as mean ± SD (n = 5 mice for each group). **(D)** Unsupervised hierarchical

clustering of RNA-Seq data of MSCs from leukemia-bearing mice, healthy MSCs, and recovered

MSCs (GFP⁻, recipient-derived). For each RNA-Seq sample, one thousand MSCs from leukemia-

bearing mice, healthy mice, and MSC-treated leukemia-bearing mice were sorted and analyzed (n

= 3-6). The raw reads (fastq files) from RNA-Seq were aligned to mouse genome by Tophat2

package, and further normalized by Cufflinks. Unsupervised hierarchical clustering was conducted

by factoextra R package. **(E)** Heatmaps of MSC secreted factor-related genes differentially

expressed between MSCs from leukemia-bearing mice and recovered MSCs (padj < 0.05, fold

change > 1.4). The expression value (DESeq2 normalized counts) of each gene was converted to

z-score values (red, high; blue, low), and the heatmaps were plotted by gplots (heatmap.2).

Columns represent the indicated cell subsets in ten MSC samples (Leukemic MSCs: n = 6,

Recovered MSCs: n = 4). **(F)** Single-cell imaging of intracellular IL-6 proteins by indirect

immunofluorescence assay (IFA) in primary MSCs sorted from non-injected sites (Leukemic

MSCs) and MSC-injected sites (Recovered MSCs) of MSC-treated leukemic mice. Images of three

single representative cells of each group are shown. **(G)** Statistical analysis of mean intensities of

IL-6 fluorescence in leukemic and recovered MSC samples. Each dot represents a single cell. Data

are analyzed by unpaired student's t-test (two-tailed). ***$p < 0.001$. Data are represented as mean

± SD. Control, n=30; Leukemia, n=30. **(H)** Single-cell imaging of intracellular CCL2 proteins by

IFA in primary MSCs sorted from non-injected sites (Leukemic MSCs) and MSC-injected sites

(Recovered MSCs) of MSC-treated leukemic mice. Images of three single representative cells of

each group are shown. **(I)** Statistical analysis of mean intensities of CCL2 fluorescence in leukemic

and recovered MSC samples. Each dot represents a single cell. Data are analyzed by unpaired

student's t-test (two-tailed). ***$p < 0.001$. Data are represented as mean ± SD. Control, n=30;

Leukemia, n=30.

Figure 5

Figure 5: Characterization of MSC-reprogrammed BM resident macrophages isolated from

leukemia-bearing mice

(A) Gene set enrichment analysis (GSEA) of the positive regulation of angiogenesis in L-Mac and

E-Mac. L-Mac indicates leukemic macrophages. E-Mac indicates MSC-reprogrammed leukemic

macrophages, which were co-cultured with MSCs *in vitro* for 12 h. DESeq2 normalized values of

the expression data were used for GSEA analysis. **(B)** Gene set enrichment analysis (GSEA) of

the cell migration-related genes in L-Mac and E-Mac. DESeq2 normalized values of the expression

data were used for GSEA analysis. **(C)** Gene ontology (GO)–enrichment analysis of the 3277

differentially expressed genes between L-Mac and E-Mac: each symbol represents a GO term

(noted in the plot); color indicates adjusted P value (padj (significance of the GO term)), and

symbol size is proportional to the number of genes. **(D)** Heatmaps of soluble factor-related genes

in MSC-reprogrammed leukemic macrophages. The expression value (DESeq2 normalized

counts) of each gene was converted to z-score value (red, high; blue, low). The heatmaps were

plotted by gplots (heatmap.2). Columns represent the indicated macrophage sample replicates (L-

Mac: n = 4 biological replicates; E-Mac: n = 4 biological replicates). **(E)** Imaging of intracellular

TNF-α, CCR7, and IL-1α proteins by indirect immunofluorescence assay (IFA) in sorted L-Mac

(Non-injected site) and E-Mac (MSC-injected site) from MSC-treated leukemic mice 12 h post-

treatment *in vivo*. The representative images of each group are shown. **(F)** RNA-Seq analysis of

Arg1 in leukemic macrophages co-cultured with MSCs *in vitro*. L-Mac indicates leukemic

macrophages. E-Mac (+MSCs) indicates MSC-reprogrammed leukemic macrophages, which were

co-cultured with MSCs *in vitro* for 12 h. L-Mac (+MSCs transwell) indicates leukemic

macrophages, which were co-cultured in transwell with MSCs *in vitro* for 12 h. Y-axis indicates

the expression value. The expression value (DESeq2 normalized counts) of each gene is illustrated

by graphpad. Each column represents a replicate. **(G)** RNA-Seq analysis of *Arg1* in leukemic

macrophages sorted from MSC-treated leukemic mice *in vivo*. Leukemic macrophages

(CD11b$^+$F4/80$^+$) were sorted from the injected site (MSC-treated tibia) and non-injected site

(untreated femur) of leukemic mice 12 h post-treatment. Y-axis indicates the expression value.

The expression value (DESeq2 normalized counts) of each gene is illustrated by graphpad. Each

column represents a replicate. **(H)** Representative intracellular staining plots of Arg1 proteins in

MSC-reprogrammed leukemic macrophages (E-Mac, MSC-injected site) and leukemic

macrophages (L-Mac, Non-injected site) from MSC-treated leukemic mice analyzed by flow

cytometry. The ratios of Arg1^{high} macrophage subpopulation are shown from L-Mac and E-Mac.

The bone marrow nucleated cells from leukemia-bearing mouse 12 h post-treatment were prepared

for intracellular flow cytometry staining of Arg1. Macrophages are identified as

CD45$^+$CD11b$^+$F4/80$^+$. **(I)** Statistical analysis of the ratios of Arg1^{high} macrophage subpopulation

in L-Mac and E-Mac. Data are analyzed by unpaired student's t-test (two-tailed). **p < 0.01. Data

are represented as mean ± SD (n = 4 mice for each group).

Figure 6

Figure 6: Intra-bone-marrow transfusion of MSC-reprogrammed macrophages largely rescues the therapeutic effects of MSC-treatment in leukemic mice

(A) Schematic diagram of MSC-reprogrammed macrophages transfusion strategy. 1×10^{5} GFP^{+} MSCs were seeded into each well of six-well plates. CD11b^{+} leukemic cells were enriched from bone marrow of leukemic mice with severe tumor burden (CD11b^{+}% in PB > 60%). Then 2×10^{6} CD11b^{+} leukemic cells were directly co-cultured with MSCs. After 12 hours, leukemic macrophages (CD11b^{+}F4/80^{+}) were sorted for transfusion. Leukemic mice with severe tumor

burden were treated by intra-bone-marrow transfusion of PBS or MSC-reprogrammed leukemic

macrophages (E-Mac). A sequential doses of E-Mac (3.3×10^7 E-Mac/kg per dose in 20 μl PBS)

were delivered into the tibia cavity using 29-gauge needle. Every tibia was treated once per two

weeks by switching the injection site every other dose. Analysis of platelets and CD11b$^+$ cells in

PB was performed monthly. **(B)** Representative dot plots of platelet populations and quantitative

gating, as identified by CD41 and CD61 staining in PB of WT mice and PBS/E-Mac treated

leukemic mice. After 4 weeks of PBS/E-Mac treatment, PB of leukemic mice with PBS/E-Mac

treatment was analyzed. **(C)** Kinetic analysis of platelets in PB of leukemia-bearing mice treated

with PBS or E-Mac (n = 6 mice for each group). **(D)** Flow cytometry analysis of MSCs in

leukemia-bearing mice post PBS/E-Mac treatment. Tibias of PBS/E-Mac treated leukemic mice at

week five since the first dose of PBS/E-Mac treatment were analyzed. MSCs were defined as

Ter119$^-$CD45$^-$CD31$^-$Sca1$^+$CD51$^+$CD146$^+$. The nucleated cell mixtures of BM and compact bones

were prepared for flow cytometry analysis of MSCs. **(E)** Statistical analysis of the absolute

numbers of host MSCs in tibias from PBS/E-Mac treated leukemic mice. Data are analyzed by

unpaired student's t-test (two-tailed). ***$p < 0.001$. Data are represented as mean ± SD (n = 6

mice for each group). **(F)** CD41$^+$ megakaryocytes from BM of PBS/E-Mac-injected site (tibia) and

non-injected site (femur) were analyzed for DNA content. Plots from representative PBS-treated

and E-Mac-treated leukemic mice four weeks post PBS/E-Mac treatment are shown. Percentages

of mature megakaryocytes with 8N and greater ploidy (≥8N) are shown. **(G)** Statistical analysis

of the percentages of mature megakaryocytes (≥8N) of CD41$^+$ BM megakaryocytes. Data are

analyzed by one-way ANOVA test. ***$p < 0.001$. Data are represented as mean ± SD (n = 4 mice

for each group).

Figure 7: *Il6^-/-* MSCs neither reprogram macrophages nor suppress leukemia

(A) ELISA of IL-6 levels in BM plasma. The BM plasma from WT mice, untreated leukemia-bearing mice, and MSC-injected sites (tibias) and non-injected sites (femurs) of MSC-treated leukemia-bearing mice eight weeks post MSC treatment were analyzed. The BMNC of each tibia or femur were flushed out using 1 ml PBS, then the supernatants of each sample were collected for ELISA. Data are analyzed by one-way ANOVA test. ***$p < 0.001$. Data are represented as mean ± SD (n = 6 mice). **(B)** Q-PCR of *Arg1* expression levels in macrophages from leukemia-bearing mice after co-culture with different sources of MSCs *in vitro*. Macrophages from leukemia-bearing mice were co-cultured with MSCs from leukemia-bearing mice, *Il6^-/-* MSCs or GFP-MSCs *in vitro* for 12 h, separately. Then, one hundred thousand E-Mac (CD11b⁺F4/80⁺) were sorted and were analyzed the expression of *Arg1*. Y-axis shows the relative expression of *Arg1* in

each group. Gapdh was used as a reference. Data are analyzed by one-way ANOVA test. **p <

0.01. Data are represented as mean ± SD (n = 3 repeats for each group). **(C)** Kinetic analysis of

tumor burden (CD11b$^+$) in PB of leukemia-bearing mice treated with GFP-MSCs or $Il6^{-/-}$ MSCs

(n = 5 mice for each group). **(D)** Flow cytometry analysis of platelets in $Il6^{-/-}$ MSC-treated

leukemia-bearing mice. Representative dot plots of platelet populations and quantitative gating, as

identified by CD41 and CD61 staining in PB of GFP-MSC- or $Il6^{-/-}$ MSC-treated leukemia-bearing

mice. PB of leukemia-bearing mice was analyzed at week-0 and week-4 post-treatment with GFP-

MSCs or $Il6^{-/-}$ MSCs. **(E)** Kinetic analysis of platelets (PLT) in PB of leukemia-bearing mice

treated with GFP-MSCs or $Il6^{-/-}$ MSCs (n = 5 mice for each group). **(F)** Q-PCR of *Arg1* expression

levels in macrophages isolated from $Il6^{-/-}$ MSC-treated leukemia-bearing mice. One hundred

thousand leukemic macrophages (CD11b$^+$F4/80$^+$) were sorted from the MSC-injected site (MSC-

treated tibia) and non-injected site (femur) of leukemia-bearing mice 12h post-treatment. Y-axis

shows the relative expression of *Arg1* in each group. Gapdh was used as reference. Data are

analyzed by one-way ANOVA test. ***p < 0.001. Data are represented as mean ± SD (n = 6 mice

for each group). **(G)** Representative intracellular staining plots of Arg1 proteins in MSC-

reprogrammed leukemic macrophages (E-Mac, MSC-injected sites) and leukemic macrophages

(L-Mac, Non-injected site) from GFP-MSC- or $Il6^{-/-}$ MSC-treated leukemic mouse analyzed by

flow cytometry. The ratios of Arg1^{high} macrophage subpopulation are shown from L-Mac and E-

Mac. The bone marrow nucleated cells from leukemia-bearing mouse 12 h post-treatment were

prepared for intracellular flow cytometry staining of Arg1. Macrophages are identified as

CD45$^+$CD11b$^+$F4/80$^+$. **(H)** Statistical analysis of the ratios of Arg1^{high} macrophage subpopulation.

Data are analyzed by one-way ANOVA test. *p < 0.05. Data are represented as mean ± SD (n = 4

mice for each group).