A Single Cell Transcriptomic Atlas Characterizes Aging Tissues in the Mouse

The Tabula Muris Consortium

Abstract

Aging is characterized by a progressive loss of physiological integrity, leading to impaired function and increased vulnerability to death[1]. Despite rapid advances over recent years, many of the molecular and cellular processes which underlie progressive loss of healthy physiology are poorly understood[2]. To gain a better insight into these processes we have created a single cell transcriptomic atlas across the life span of Mus musculus which includes data from 23 tissues and organs. We discovered cell-specific changes occurring across multiple cell types and organs, as well as age related changes in the cellular composition of different organs. Using single-cell transcriptomic data we were able to assess cell type specific manifestations of different hallmarks of aging, such as senescence[3], genomic instability[4] and changes in the organism's immune system[2]. This Tabula Muris Senis provides a wealth of new molecular information about how the most significant hallmarks of aging are reflected in a broad range of tissues and cell types.

We performed single cell RNA sequencing on 529,823 cells from male and female C57BL/6JN mice belonging to six age groups ranging from one month (human early childhood equivalent) to thirty months (human centenarian equivalent) (Figure 1a). We prepared single cell suspensions of the bladder, bone marrow, brain (cerebellum, cortex, hippocampus and striatum), fat (brown, gonadal, mesenteric and subcutaneous), heart and aorta, kidney, large intestine, limb muscle and diaphragm, liver, lung, mammary gland, pancreas, skin, spleen, thymus, tongue and trachea for all mice. Data were collected for all six age groups using microfluidic droplets (droplet), while the 3m, 18m and 24m time points were also analyzed using single cells sorted in microtiter well plates (FACS) (Extended Data Figure 1, Extended Data Figure 2a; Supplementary Tables 1&2; Supplementary Figures 1-3). The droplet data allow large numbers of cells to be analyzed using 3' end counting, while the FACS data allow for higher sensitivity measurements over smaller numbers of cells as well as sequence information across the entire transcript length. Analyzing multiple organs from the same animal enabled us to create data controlled for age, environment, and epigenetic effects.

The previously published 3m time point, referred to as the *Tabula Muris*[5], represents ~20% of the cells in the entire dataset and was used as a basis to perform semi-automated cell type annotation of the additional time points (Figure 1b, Extended Data Figure 2b). Using this approach, we were able to automatically annotate over 70% of the cells. All the automated cell annotations were reviewed and approved by human experts, and the remaining cells were annotated by hand, creating one of the largest manually curated single cell transcriptomic resources in existence. Many of these cell types have not previously been obtained in pure populations, and these data provide a wealth of new information on their characteristic gene-expression profiles. To demonstrate that the

47   annotations performed separately for each tissue were consistent across the entire
48   organism, we clustered all cells using an unbiased, graph-based clustering approach[6,7]
49   (Figure 1c,e) and showed that cell types such as B cells and endothelial cells which are
50   shared across different organs and tissues occupy the same clusters irrespectively of the
51   tissue of origin (Figure 1d,f).
52
Tabula Muris Senis provides a powerful resource with which to explore aging related
changes in specific cell types. The entire dataset can be explored interactively at tabula-
muris-senis.ds.czbiohub.org.   Gene counts and metadata are available from figshare
(https://figshare.com/projects/Tabula_Muris_Senis/64982) and GEO (GSE132042), the
code used for the analysis is available from GitHub (https://github.com/czbiohub/tabula-
muris-senis)   and   the   raw   data   are   available   from   AWS   Public   Datasets
(https://s3.console.aws.amazon.com/s3/buckets/czb-tabula-muris-senis/).   An   important
use of the single cell data is to resolve whether gene expression changes observed in bulk
experiments are due to changes in gene expression in each cell of the population, or
whether the gene expression in each cell stays constant but the number of cells of that
type changes, or both. In a global analysis of gene expression changes using the Tabula
Muris Senis and bulk RNAseq from tissues[8], we observed that in many cases changes in
gene expression are due to both changes in the numbers of cells in a population and to
changes in the gene expression levels in each cell (Extended Data Figure 3). As a
specific example of this approach, we investigated how the expression of Cdkn2a
changes with age. As Cdkn2a/p16 is one of the most commonly used markers of
senescence[9] and an important hallmark of aging[10], we computed the fraction of cells
expressing Cdkn2a at each age. The fraction of cells expressing the gene more than
doubled in older animals in both FACS (Figure 2a) and droplet (Figure 2b), accompanied
by a 2-fold increase in the actual expression level of p16 by those cells that did express it
(Figure 2c,d).
74
As another example of how the dataset can be used, we investigated how the cellular
composition of each tissue changes with age by evaluating how the relative cell type
proportions within a tissue change with age (Supplementary Table 3). The overall cell
composition for all tissues is in Extended Data Figure   4. When interpreting
compositional data, one must bear in mind that dissociation does not affect all of the cell
types in a tissue equally, so changes in the relative composition of a given cell type with
age are more meaningful than trying to compare proportions of different cell types at a
single age[11–13]. Nonetheless, the changes in relative proportion of cell types provide
important information on the effects of aging in a variety of tissues.
84
The bladder has pronounced changes in cell type composition with age (Figure 2e).
While the mesenchymal compartment of this tissue decreases by a factor of three over the
lifetime of the mouse (Figure 2e left), the urothelial compartment increases by a similar
amount (Figure 2e right). The observation that the bladder urothelial cells increase with
age is concordant with known age-related urothelial changes[14]. Differential gene
expression analysis of overall tissue changes with age revealed that stromal-associated
genes (Col1a1, Col1a2, Col3a1, Dcn) are downregulated while epithelial-associated
genes (Krt15, Krt18, Sfn) are upregulated, supporting the compositional observations

(Figure 2f,g; Supplementary Table 4). The decline of the endothelial population suggests
that bladder aging in mice may be associated with lower organ vascularization, consistent
with recent findings[15,16] and with the observed downregulation of vasculature associated
genes Htra1 and Fos (Figure 2f,g; Supplementary Table 4). The increase in the leukocyte
population could be indicative of an inflammatory tissue microenvironment, a common
hallmark of aging which is consistent with literature on overactive bladders[17] and
supported by a significant overexpression of Lgals3, Igfbp2 and Ly6d (Figure 2f,g;
Supplementary Table 4).
101
Age-dependent changes in the kidney include a decrease in the relative abundance of
mesangial cells, capillary endothelial cells, loop of Henle ascending limb epithelial cells
and loop of Henle thick ascending limb epithelial cells (Figure 2h). Both mesangial cells
and capillary endothelial cells are core glomerular cells and their relative abundances
reduction (Figure 2h top panels), together with downregulation of Egf and Atp1a1
(Figure 2i,j; Supplementary Table 4) suggest impaired glomerular filtration rate[18,19]. This
finding is reinforced by the differential gene expression results indicating that
uromodulin (Umod), the most abundant protein in urine[20], is downregulated. Umod is
produced by the epithelial cells that line the thick ascending limb, and therefore given the
relative decrease in the proportion of epithelial cells in the ascending and thick ascending
limb, our results suggest that normal kidney functions are impaired[21] (Figure 2h bottom
panels, Figure 2i,j; Supplementary Table 4).
114
The liver is yet another tissue for which we observed changed tissue compositions with
age, namely that the relative amount of hepatocytes decreases with age (Extended Data
Figure 5a), which is supported by the reduction in the expression of albumin (Alb;
Extended Data Figure 5b,c; Supplementary Table 4). Differential gene expression
showed an increased immune signature, as illustrated by overexpression of H2-Aa, H2-
Ab1, H2-D1, H2-Eb1, Cd74, Lyz2 and others. Previous findings suggested that pro-
inflammatory macrophages drive cellular senescence and identified Il1b as a gene whose
liver expression was remarkably different with age[22]. We stained liver Kupffer cells
(Extended Data Figure 5d) with Clec4f (canonical Kupffer cell marker) and found the
number of Clec4f+ cells do not change with age, consistent with the results of the tissue
composition analysis (Supplementary Table 5; Extended Data Figure 5e). However, when
co-staining with Il1b, we found an increase with age in the number of cells expressing
Clec4f and Il1b (Extended Data Figure 5f,g). Il1b has low expression in normal
physiological conditions[23]. Specific blocking of IL1-RI (Il1b receptor) in hepatocytes has
been shown to attenuate cell death upon injury, supporting the idea that increased
expression of Il1b in Kupffer cells is typically a poor prognostic[24]. Regarding immune
defense within the liver, sinusoidal endothelial cells (LSECs) play a unique role, being
the main carriers of the mannose receptor (Mrc1) in the liver. Mrc1 expression in LSECs
mediates endocytosis of pathogen and damage related molecules. Our findings identify
increased Mrc1 age-related expression. Inflammatory signals have been found to up
regulate Mrc1 expression and endocytosis [25]. Staining for Mrc1 alongside classical LSEC
marker Pecam1 (Supplementary Table 5; Extended Data Figure 5h,i) found the number
of Mrc1 expressing LSECs increase over age (Extended Data Figure 5j,k). LSECs have a
been found to have a reduced endocytic capacity in aged livers, while it has been

suggested that LSECs proliferate after injury or that bone-marrow derived LSECs
progenitors are recruited to the liver. This suggests that changes in LSEC gene signatures
with age are linked closely with their function in immune response.
142
In the case of spleen our results show that with age the proportion of T cells decreases
while the relative amount of plasma cells increases (Figure 2k). This is supported by
upregulation of B cell/plasma cell markers (Cd79a, Igj; Figure 2l,m; Supplementary
Table 4) and downregulation of Cd3d (Figure 2m; Supplementary Table 4). Similarly, in
mammary gland we also observed a significant decline of the T cell population (Extended
Data Figure 6a). Age-related decline of T cell populations has been associated with an
increased risk of infectious disease and cancer[26] and our results suggest that this may also
happen in the spleen and mammary gland. We found that members of the AP1
transcription factors[27] (Junb, Jund and Fos) were upregulated with age (Extended Data
Figure 6b,c; Supplementary Table 4); this result is consistent with the observation that
normal involution of the mammary gland is accompanied by significantly increased
expression of many of these AP1-related transcription factors[28].
155
Genomic instability is among the most widely studied aging hallmarks[1] and the full-
length transcript data from the FACS data allows the analysis of somatic mutation
accumulation with age. We used the Genome Analysis ToolKit (GATK)[29] to perform
SNP discovery across all FACS samples simultaneously (Supplementary Table 6), using
GATK Best Practices recommendations[30,31]. We focused on genes expressed in at least
75% of cells for each age group within a particular tissue. We observed an age-related
increase in the number of mutations across all of the organs we analyzed (Figure 3;
Extended Data Figure 7a,8a,9a), with tongue and bladder being the most affected. Our
analysis controls for sequencing coverage and gene expression levels (Figure 7b, 8b, 9b).
The number of mutations observed at each age are larger than technical errors due to
amplification and sequencing errors, which can be estimated using ERCC controls that
were spiked into each well of the microtiter plates[32] (Figure 3; Extended Data Figure 7c-
d, 8c-d, 9c-d). Despite the fact that it is difficult to infer genome-wide mutation rates
from the transcriptome, which is known to inflate apparent mutational rates for a variety
of reasons[32], the observed trend is a useful indirect estimate of mutational frequency and
genome stability.
172
A final hallmark of aging which we investigated was the effect of age-induced changes
on the immune system[2]. Analyzing a complete set of tissues from the same individual
animal using the full-length transcripts obtained in the FACS data enabled us to analyze
clonal relationships between B-cells and T-cells throughout the organism. We
computationally reconstructed the sequence of the B-cell receptor (BCR) and T-cell
receptor (TCR) for B cells and T cells present in the FACS data using singlecell-ige and
TraCeR, respectively[33,34]. BCRs were assembled for 6,050 cells (Figure 4a) and TCRs for
6,000 cells (Figure 4b). The number of cells with assembled BCRs was 1,818 for 3m,
1,356 for 18m and 2,876 for 24m old mice. We parsed the singlecell-ige[33] output to
define B-cell clonotypes based on the sequence of the assembled BCR (Supplementary
Table 7; see Extended Methods) and found that while most of the cells at 3m were not
part of a clone (9% were part of a clonal family), the number of B-cells belonging to a

clonotype doubled at 18m (20%) when compared to 3m and doubled again from 18m to
24m (~38%).
187
The number of cells with assembled TCRs were roughly equal between 3m, 18m and
24m (2,076, 2,056 and 1,868 cells, respectively). Clonotype assignment is part of the
output obtained by TraCeR[34] (Supplementary Table 7). Interestingly, only 55 out of
1,895 cells at 3m were part of a clone. For 18m, 479 out of 2,056 cells were part of a
clone and for 24m, 348 out of 1,780 cells were part of a clone, indicating again an
increase in clonality of the T-cell repertoire at later ages. These changes in clonality for
both B and T cell repertoires are noteworthy because they suggest that the immune
system of a 24m mouse will be less likely to respond to new pathogens, corroborating
literature suggesting that older individuals have higher vulnerability to new infections
and lower benefits from vacination[35,36].
198
As a final example of how the Tabula Muris Senis can be used to discover how cell types
change with age, we computed an overall diversity score to identify which cell types
were more susceptible to changes with age (Extended Data Figure 10). The diversity
score is computed as the Shannon entropy of the cluster assignment and then regressed
against age to provide a p-value (see Methods). We observed significant changes in
diversity affecting cells of the immune system originating from the brain and in the
kidney (Figure 4c, Extended Data Figure 11a,b). These results were not confounded by
the number of genes expressed per cell (Extended Data Figure 11c,d). We found that in
brain myeloid microglial cells, the majority of young (3m) microglia occupy clusters 1
and 6, while old (18m, 24m) microglia constitute the vast majority of cells in clusters 10,
12 and 14 (Figure 4d). Trajectory analysis suggests that young microglia go through an
intermediate state, represented by the clusters mostly occupied by 18m microglial cells
before acquiring the signature of old microglia (Extended Data Figure 11e). Clusters 10,
12 and 14 are mainly comprised of 18- and 24-month old microglia. These cells up-
regulate MHC class I genes (H2-D1, H2-K1, B2m), along with genes associated with
degenerative disease (e.g. Fth1)[37,38]. When contrasting with clusters 1 and 6, which
contain mostly 3m microglia, clusters 10, 12 and 14 gene expression is enriched with
interferon responsive or regulatory genes (e.g. Oasl2, Oas1a, Ifit3, Rtp4, Bst2, Stat1, Irf7,
Ifitm3, Usp18, Ifi204, Ifit2), suggesting an expansion of this small pro-inflammatory
subset of microglia in the aging brain[39]. Moreover, the list of differentially expressed
genes between "young" and "old" clusters resembled the Alzheimer's disease specific
microglial signature previously reported[37], with 55 out of the top 200 differential
expressed genes being shared between the two differential gene expression lists (Figure
4e; Supplementary Table 8). Regarding kidney macrophages, we found two clusters that
remarkably changed their composition with age. Cluster 10 is primarily composed of
cells of 1m- and 3-month old mice while cluster 13 is mostly composed of cells of 18-,
21-, 24- and 30-month old mice (Figure 4f). Differential gene expression revealed that
cluster 10 is enriched for an M2-macrophage gene signature (e.g. Il10, H2-Eb1, H2-Ab1,
H2-Aa, Cd74, C1qa, Cxcl16, Hexb, Cd81, C1qb, Cd72) while cluster 13 resembles a M1-
proinflammatory macrophage state[40] (e.g. Hp, Itgal, Spex1, Gngt2) (Extended Data
Figure 11f; Supplementary Table 8).
230

The Tabula Muris Senis is a comprehensive resource for the cell biology community
which offers a detailed molecular and cell-type specific portrait of aging. We view such
a cell atlas as an essential companion to the genome: the genome provides a blueprint for
the organism but does not explain how genes are used in a cell type specific manner or
how the usage of genes changes over the lifetime of the organism. The cell atlas provides
a deep characterization of phenotype and physiology which can serve as a reference for
understanding many aspects of the cell biological changes that mammals undergo during
their lifespan.
239

Figure Legends
Figure 1. Overview of Tabula Muris Senis.
a, 23 organs from 19 male and 11 female mice were analyzed at 6 different time points. The bar plot shows the number of sequenced cells per organ prepared by FACS (n=23 organs) and microfluidic droplets (n=16 organs). For the droplet dataset the Fat subtissues were processed together (Fat = BAT+GAT+MAT+SCAT). **b**, Annotation workflow. Data were clustered together across all time points. We used the Tabula Muris (3m time point) as a reference for the automated pipeline and the annotations were manually curated by tissue experts. **c**, UMAP plot of all cells collected by FACS, colored by organ (Extended Data Figure 2c), overlaid with the Louvain cluster numbers; n = 110,824 individual cells. **d**, B cells (top) and endothelial cells (bottom) independently annotated for each organ cluster together by unbiased whole-transcriptome Louvain clustering, irrespectively of the organ they originate from. **e**, UMAP plot of all cells collected by droplet, colored by organ (Extended Data Figure 2c), overlaid with the Louvain cluster numbers; n = 245,389 individual cells. **f**, B cells (and endothelial cells) independently annotated for each organ cluster together by unbiased whole-transcriptome Louvain clustering, irrespectively of the organ where they were found.

Figure 2. Cellular changes during aging.
a,b, Bar plot showing the fractions of cells expressing Cdkn2a at each age group for FACS (**a**) and droplet (**b**). **c,d**, Bar plot of the median expression of Cdkn2a for the cells that do express the gene at each age group for FACS (**c**) and droplet (**d**). The p-value was obtained using a Mann-Whitney-Wilcoxon rank-sum two-sided test. **e**, Bladder cell (**left**) and bladder urothelial cell (**right**) relative abundances change significantly with age (p-value<0.05 and r^2>0.7 for a hypothesis test whose null hypothesis is that the slope is zero, using two-sided Wald Test with t-distribution of the test statistic). **f,g**, Top 10 upregulated and downregulated genes in bladder FACS (**f**) and droplet (**g**) using age as a continuous covariate while controlling for sex. Genes were classified as significant under an FDR threshold of 0.01 and an age coefficient threshold of 0.005 (corresponding to ~10% fold change). **h**, Kidney capillary endothelial cell (**top-left**), mesangial cell (**top-right**), loop of Henle ascending limb epithelial cell (**bottom-left**) and loop of Henle thick ascending limb epithelial cell (**bottom-right**) relative abundances change significantly with age (p-value<0.05 and r^2>0.7 for a hypothesis test whose null hypothesis is that the slope is zero, using two-sided Wald Test with t-distribution of the test statistic). **i,j,** Top 10 upregulated and downregulated genes in kidney FACS (**i**) and droplet (**j**) using age as a continuous covariate while controlling for sex. Genes were classified as significant under an FDR threshold of 0.01 and an age coefficient threshold of 0.005 (corresponding to ~10% fold change). **k**, Spleen plasma cell (**left**) and T cell (**right**) relative abundances change significantly with age (p-value<0.05 and r^2>0.7 for a hypothesis test whose null hypothesis is that the slope is zero, using two-sided Wald Test with t-distribution of the test statistic). **l,m,** Top 10 upregulated and downregulated genes in spleen FACS (**l**) and droplet (**m**) using age as a continuous covariate while controlling for sex. Genes were classified as significant under an FDR threshold of 0.01 and an age coefficient threshold of 0.005 (corresponding to ~10% fold change).

Figure 3. Mutational burden across tissues in the aging mice.
Distribution of the difference of the mean mutation in the gene set (and ERCC spike-in controls) per cell between 24m and 3m and 18m and 3m for all tissues and cells (**a**) and with the cell types split in five functional groups, endothelial (**b**), immune (**c**), parenchymal (**d**), stem/progenitor cell (**e**) and stromal (**f**).

Figure 4. The aging immune system.
a, B-cell clonal families. The pie chart shows the proportion of singleton B cells and B cells that are part of clonal families at 3m, 18m and 24m. For each time point, the clonal families are represented in a tree structure for which the central node is age. Connected to the age node there is an additional node (dark gray) that represents each animal and the clonal families are depicted for each animal. For each clonal family, cells that are part of that family are colored by the organ of origin. **b**, T-cell clonal families. The pie chart shows the proportion of singleton T cells and T cells that are part of clonal families at 3m, 18m and 24m. For each time point, clonal families are represented in a tree structure for which the central node is age. Connected to the age node there is an additional node (dark gray) that represents each animal and the clonal families are depicted for each animal. For each clonal family, cells that are part of that family are colored by the organ of origin. **c**, Diversity score for the two cell types that significantly change with age. **d**, UMAP plot of the brain myeloid microglial cell Leiden clusters (numbers) colored by age. Faded clusters do not change their relative age cell composition; colored clusters change their relative cell composition. **e**, UMAP plot of the brain myeloid microglial cells when scored using the microglia Alzheimer's disease signature (Supplementary Table 8). **f**, UMAP plot of the kidney macrophage Leiden clusters (numbers) colored by age group.

Extended Data Figure Legends
Extended Data Figure 1. Overview of Tabula Muris Senis (cont.)
a,b, UMAP plot of all cells collected for FACS colored by tissue (**a**) or age (**b**). **c**, Pie chart with the summary statistics for FACS. **d,e**, UMAP plot of all cells collected for droplet colored by tissue (**d**) or age (**e**). **f**, Pie chart with the summary statistics for droplet.

Extended Data Figure 2. Overview of Tabula Muris Senis (cont.)
a, Balloon plot showing the number of sequenced cells per sequencing method per organ per sex per age. **b**, Schematic analysis workflow. **c,d,** Tabula Muris Senis color dictionary for organs and tissues (**c**) and ages (**d**).

Extended Data Figure 3. Comparison of bulk and single-cell datasets. Aging patterns
from bulk and single-cell data are consistent. Strong changes in bulk gene expression
with aging can be either explained by cell or read count-based changes in single-cell data
FACS (**a**) and droplet (**b**). Wilcoxon–Mann–Whitney indicates that single-cell data based
$\log_2$ fold-changes of cell or read counts distinguish between up and down regulated genes
in bulk data.

338
339
340   **Extended Data Figure 4. Tissue cell compositions. a-p**, Alphabetically sorted tissue bar
341   plot showing the relative abundances of cell types in each tissue across the entire age
342   range for the droplet dataset. The tissue cell composition is also available at our online
343   browser tabula-muris-senis.ds.czbiohub.org

344
345
346
Extended Data Figure 5. Cellular changes during aging in the liver.
a, Liver hepatocyte relative abundances change significantly with age (p-value<0.05 and
r^2>0.7 for a hypothesis test whose null hypothesis is that the slope is zero, using two-
sided Wald Test with t-distribution of the test statistic). **b,c**, Top 10 upregulated and
downregulated genes in liver FACS (**b**) and droplet (**c**) using age as a continuous
covariate while controlling for sex. Genes were classified as significant under an FDR
threshold of 0.01 and an age coefficient threshold of 0.005 (corresponding to ~10% fold
change). **d-g,** Staining of Kupffer cells across age (**d**) and respective quantification (**e-g**).
h-k, Staining of liver endothelial cells across ages (**h**) and respective quantification (**i-k**).
The white scale bar corresponds to 100μm.

357
358
Extended Data Figure 6. Cellular changes during aging (cont.)
a, Mammary gland T cell relative abundances change significantly with age (p-
value<0.05 and r^2>0.7 for a hypothesis test whose null hypothesis is that the slope is zero,
using two-sided Wald Test with t-distribution of the test statistic). **b,c**, Top 10
upregulated and downregulated genes in mammary gland FACS (**b**) and droplet (**c**) using
age as a continuous covariate while controlling for sex. Genes were classified as
significant under an FDR threshold of 0.01 and an age coefficient threshold of 0.005
(corresponding to ~10% fold change). **d**, Marrow precursor B cell relative abundances
change significantly with age (p-value<0.05 and r^2>0.7 for a hypothesis test whose null
hypothesis is that the slope is zero, using two-sided Wald Test with t-distribution of the
test statistic). **e,f**, Top 10 upregulated and downregulated genes in marrow FACS (**e**) and
droplet (**f**) using age as a continuous covariate while controlling for sex. Genes were
classified as significant under an FDR threshold of 0.01 and an age coefficient threshold
of 0.005 (corresponding to ~10% fold change). **g**, Skin keratinocyte stem cell relative
abundances change significantly with age (p-value<0.05 and r^2>0.7 for a hypothesis test
whose null hypothesis is that the slope is zero, using two-sided Wald Test with t-
distribution of the test statistic). **h**, Top 10 upregulated and downregulated genes in skin
FACS using age as a continuous covariate while controlling for sex. Genes were

classified as significant under an FDR threshold of 0.01 and an age coefficient threshold
of 0.005 (corresponding to ~10% fold change).
379
380

Extended Data Figure 7. Mutational burden across tissues in the aging mice (cont.
24m vs 3m).
a,b, Mean number of somatic mutations (**a**) and raw expression (**b**) across all tissues per
age group (3m and 24m). **c,d**, Mean number of mutations in ERCC spike-in (**c**) and
ERCC raw expression (**d**) across all tissues per age group (3m and 24m). Mutations are
presented as the mean number of mutations per gene per cell.
387
388

Extended Data Figure 8. Mutational burden across tissues in the aging mice (cont.
18m vs 3m).
a,b, Mean number of somatic mutations (**a**) and raw expression (**b**) across all tissues per
age group (3m and 18m). **c,d**, Mean number of mutations in ERCC spike-in (**c**) and
ERCC raw expression (**d**) across all tissues per age group (3m and 18m). Mutations are
presented as the mean number of mutations per gene per cell.
395
396

Extended Data Figure 9. Mutational burden across tissues in the aging mice (cont.
24m vs 18m).
a,b, Mean number of somatic mutations (**a**) and raw expression (**b**) across all tissues per
age group (18m and 24m). **c,d**, Mean number of mutations in ERCC spike-in (**c**) and
ERCC raw expression (**d**) across all tissues per age group (18m and 24m). Mutations are
presented as the mean number of mutations per gene per cell.
403
404

405 **Extended Data Figure 10. Diversity score summary.**
406 **a,b**, Heatmap summary of the overall tissue diversity score for FACS (**a**) and droplet (**b**).
407 **c,d**, Heatmap summary of the tissue cell-type diversity score for FACS (**c**) and droplet
408 (**d**).
409
410

Extended Data Figure 11. The aging immune system (cont.)
a,b, Diversity score at different cluster resolutions for FACS brain myeloid microglia cell
(**a**) and droplet kidney macrophage (**b**). **c,d**, Diversity score correlation with the number
of genes expressed per tissue (**c**) or tissue cell-type (**d**). **e**, PAGA[41] trajectory for brain
myeloid microglia cell. **f**, Differential gene expression analysis of cluster 10 (mostly
young macrophages) versus clusters 13 (mostly old macrophages). For the complete gene
list please refer to Supplementary Table 8.
418
419
420
421

Supplementary Figure Legends

Supplementary Figure 1. FACS sequencing statistics.

a, Box plot of the number of genes detected per cell for each organ and age. **b**, Box plot of the number of reads per cell (log-scale) for each organ and age.

Supplementary Figure 2. Droplet sequencing statistics.

Box plot of the number of genes detected per cell for each organ and age.

Supplementary Figure 3. Droplet sequencing statistics (cont.)

Box plot of the number of UMIs per cell (log-scale) for each organ and age.

Supplementary Tables

Supplementary Table 1. Summary of the FACS dataset.

a, Number of cells grouped by age, sex, mouse id and tissue. **b**, Number of cells grouped by tissue, cell ontology class and age. **c**, Number of cells grouped by Louvain cluster number, cell ontology class, tissue and age. **d**, Number of cells grouped by cell ontology class, Louvain cluster number, tissue and age. **e**, Fraction of cells in each Louvain cluster per cell ontology class and tissue. **f**, Fraction of cells in each Louvain cluster per tissue. **g**, Fraction of cells in each Louvain cluster per cell ontology class.

Supplementary Table 2. Summary of the droplet dataset.

a, Number of cells grouped by age, sex, mouse id and tissue. **b**, Number of cells grouped by tissue, cell ontology class and age. **c**, Number of cells grouped by Louvain cluster number, cell ontology class, tissue and age. **d**, Number of cells grouped by cell ontology class, Louvain cluster number, tissue and age. **e**, Fraction of cells in each Louvain cluster per cell ontology class and tissue. **f**, Fraction of cells in each Louvain cluster per tissue. **g**, Fraction of cells in each Louvain cluster per cell ontology class.

Supplementary Table 3. Cellular fraction changes. This supplementary table supports Figure 2e,h,k; Extended Data Figure 4; Extended Data Figure 5a and Extended Data Figure 6a,d,g.

Supplementary Table 4. Differential gene expression analysis. This supplementary table supports Figure 2f,g,i,j,l,m; Extended Data Figure 5b,c and Extended Data Figure 6b,c,e,f,h.

Supplementary Table 5. Quantification of Liver in-situ staining. This supplementary table supports Figure 5d-k. fov stands for field of view.

Supplementary Table 6. Summary statistics for the GATK analysis. Cell is the unique cell identifier; **ercc** is the average number of mutations per cell found in the ERCC spike-in, **adata** is the average number of mutations per cell in the gene set of the tissue; **ercc_raw_counts** are the average number of ERCC spike-in counts per cell and **ercc_counts** are the log(ercc_raw_counts+1); **adata_raw_counts** are the average number of gene counts per cell and **adata_counts** are the log(adata_raw_counts+1); **tissue**, **age** and **cell_ontology_class** are the metadata of the respective cell id and **agenum** is the age as a numerical variable; **functional_annotations** is a categorical variable binning each cell type as endothelial, immune, parenchymal, stem cell/progenitor or stromal.

Supplementary Table 7. B-cell and T-cell repertoire analysis raw data. This table supports Figure 4a,b.

Supplementary Table 8. Differential gene expression for the tissue cell type whose diversity significantly changes with age. a, FACS brain myeloid microglia differentially upregulated genes between clusters 10, 12 and 14 versus clusters 1 and 6. **b**, FACS brain myeloid microglia differentially upregulated genes between clusters 1 and 6 versus clusters 10, 12 and 14. **c**, Droplet kidney macrophage differentially upregulated genes between cluster 13 and cluster 10. **d**, Droplet kidney macrophage differentially upregulated genes between cluster 10 and cluster 13. **e,** Alzheimer's disease microglia signature from[37]. This table supports Figure 4d,e and Extended Data Figure 11f.

Methods

All data, protocols, analysis scripts and an interactive data browser are publicly available.

Experimental Procedures

Mice and organ collection

Male and virgin female C57BL/6JN mice were shipped from the National Institute on Aging colony at Charles River (housed at 67–73 °F) to the Veterinary Medical Unit (VMU; housed at 68–76 °F)) at the VA Palo Alto (VA). At both locations, mice were housed on a 12-h light/dark cycle and provided food and water *ad libitum*. The diet at Charles River was NIH-31, and Teklad 2918 at the VA VMU. Littermates were not recorded or tracked, and mice were housed at the VA VMU for no longer than 2 weeks before euthanasia, with the exception of mice older than 18 months, which were housed at the VA VMU beginning at 18 months of age. Before tissue collection, mice were placed in sterile collection chambers at 8 am for 15 min to collect fresh fecal pellets. After anaesthetization with 2.5% v/v Avertin, mice were weighed, shaved, and blood was drawn via cardiac puncture before transcardial perfusion with 20 ml PBS. Mesenteric adipose tissue was then immediately collected to avoid exposure to the liver and pancreas perfusate, which negatively affects cell sorting. Isolating viable single cells from both the pancreas and the liver of the same mouse was not possible; therefore, two males and two females were used for each. Whole organs were then dissected in the following order: large intestine, spleen, thymus, trachea, tongue, brain, heart, lung, kidney, gonadal adipose tissue, bladder, diaphragm, limb muscle (tibialis anterior), skin (dorsal), subcutaneous adipose tissue (inguinal pad), mammary glands (fat pads 2, 3 and 4), brown adipose tissue (interscapular pad), aorta and bone marrow (spine and limb bones). Organ collection concluded by 10 am. After single-cell dissociation as described below, cell suspensions were either used for FACS of individual cells into 384-well plates, or for preparation of the microfluidic droplet library. All animal care and procedures were carried out in accordance with institutional guidelines approved by the VA Palo Alto Committee on Animal Research.

Tissue dissociation and sample preparation

All tissues were processed as previously described[5].

Sample size, randomization and blinding

No sample size choice was performed before the study. Randomization and blinding were not performed: the authors were aware of all data and metadata-related variables during the entire course of the study.

Single-cell methods

All protocols used in this study are described in detail elsewhere[5]. Those include: i) preparation of lysis plates, ii) FACS sorting, iii) cDNA synthesis using the Smart-seq2 protocol[42,43], iv) library preparation using an in-house version of Tn5[44,45], v) library pooling and Quality control and vi) sequencing. For further details please refer to http://dx.doi.org/10.17504/protocols.io.2uwgexe

538
Microfluidic droplet single-cell analysis
Single cells were captured in droplet emulsions using the GemCode Single-Cell
Instrument (10x Genomics) and scRNA-seq libraries were constructed as per the 10x
Genomics protocol using GemCode Single-Cell 3′ Gel Bead and Library V2 Kit. In
brief, single cell suspensions were examined using an inverted microscope, and if
sample quality was deemed satisfactory, the sample was diluted in PBS with 2% FBS
to a concentration of 1000 cells per μl. If cell suspensions contained cell aggregates or
debris, two additional washes in PBS with 2% FBS at 300$g$for 5 min at 4 °C were
performed. Cell concentration was measured either with a Moxi GO II (Orflo
Technologies) or a haemocytometer. Cells were loaded in each channel with a target
output of 5,000 cells per sample. All reactions were performed in the Biorad C1000
Touch Thermal cycler with 96-Deep Well Reaction Module. 12 cycles were used for
cDNA amplification and sample index PCR. Amplified cDNA and final libraries were
evaluated on a Fragment Analyzer using a High Sensitivity NGS Analysis Kit
(Advanced Analytical). The average fragment length of 10x cDNA libraries was
quantitated on a Fragment Analyzer (AATI), and by qPCR with the Kapa Library
Quantification kit for Illumina. Each library was diluted to 2 nM, and equal volumes of
16 libraries were pooled for each NovaSeq sequencing run. Pools were sequenced with
100 cycle run kits with 26 bases for Read 1, 8 bases for Index 1, and 90 bases for Read
2 (Illumina 20012862). A PhiX control library was spiked in at 0.2 to 1%. Libraries
were sequenced on the NovaSeq 6000 Sequencing System (Illumina).
560
561
Computational methods
Data extraction
Sequences from the NovaSeq were de-multiplexed using bcl2fastq version 2.19.0.316.
Reads were aligned using to the mm10plus genome using STAR version 2.5.2b with
parameters TK. Gene counts were produced using HTSEQ version 0.6.1p1 with
default parameters, except 'stranded' was set to 'false', and 'mode' was set to
'intersection-nonempty'. Sequences from the microfluidic droplet platform were de-
multiplexed and aligned using CellRanger version 2.0.1, available from 10x Genomics
with default parameters.
571
Data pre-processing
Gene count tables were combined with the metadata variables using the Scanpy[46]
Python package version 1.4. We removed genes not expressed in at least 3 cells and
then cells that did not have at least 250 detected genes. For FACS we removed cells
with less than 5000 counts and for droplet cells with less than 2500 UMIs. The data
was then normalized using size factor normalization such that every cell has 10,000
counts and log transformed. We computed highly variable genes using default
parameters and then scaled the data to a maximum value of 10. After we computed
PCA, neighborhood graph and clustered the data using Louvain[6] and Leiden[7] methods.
The data was visualized using UMAP projection. Step-by-step instructions to
reproduce the pre-processing of the data are available from GitHub.
583

Cell type annotation
To define cell types we analyzed each organ independently but combining all ages. In a
nutshell, we performed principal component analysis on the most variable genes between
cells, followed by Louvain and Leiden graph-based clustering. Next we subset the data
for 3m (Tabula Muris[5]) and compute how many cell types map to each individual cluster.
For the clusters that we had a single 1:1 mapping (cluster:cell type) we propagate the
annotations for all ages; in case there is a 1:many mapping we flagged that cluster for
manual validation. Step-by-step instructions to reproduce this method are available from
GitHub. For each cluster, we provide annotations in the controlled vocabulary of the cell
ontology[47] to facilitate inter-experiment comparisons. Using this method, we were able to
annotate automatically (~1min per tissue) over 70% of the dataset. The automatic
annotations were then reviewed by each of the tissue experts leading to a fully curated
dataset for all the cell types in Tabula Muris Senis.
597
Tissue cell composition analysis
For each tissue and age, we computed the relative proportion of each cell type. Next we
used scipy.stats linregress to regress the relative tissue-cell type changes against age
and considered significant the changes with p-value<0.05 for a hypothesis test whose null
hypothesis is that the slope is zero, using two-sided Wald Test with t-distribution of the
test statistic and a r^2>0.5.
604
Differential gene expression
We performed differential gene expression analysis on each tissue with a well-powered
sample size (>100 cells in both young (1m and 3m) and old age group (18m, 21m, 24m
and 30m)). We use a linear model[48] treating age as a numerical variable while controlling
for sex. We apply a false-discovery rate (FDR) threshold of 0.01 and an age coefficient
threshold of 0.005 (corresponding to ~10% fold change).
611
In Situ RNA Hybridization and quantification.
In situ RNA hybridization was performed using the Advanced Cell Diagnostics
RNAscope® Multiplex Fluorescent Detection kit v2 (323110, Bio-techne) according to
the manufacturer's instructions. Staining of mouse liver specimens was performed using
5μm paraffin-embedded thick sessions. Mouse livers were fixed in 10% formalin buffer
saline (HT501128, Sigma Aldrich) for 24h at room temperature before paraffin
embedding. For multiplex staining the following probes were used; *Clec4f* (Mm-Clec4f
480421, *Il1b* (Mm-Il1b 316891-C2), *Pecam1* (Mm-Pecam-1 316721), *Mrc1* (Mm-Mrc1
437511-C3). Slides were counter stained with Prolong gold antifade reagent with DAPI
(P36931, Life technologies). Mounted slides were imaged on a Leica DM6 B fluorescent
microscope (Leica Biosystems). Image quantification was performed using the starfish
open source image-based transcriptomics pipeline (please refer to Starfish: Open Source
Image Based Transcriptomics and Proteomics Tools available from
http://github.com/spacetx/starfish and [49]).
626
Comparison between bulk and single-cell datasets
The differential gene analysis was defined on a per tissue basis. First, we investigated
genes based on the single-cell data. We only considered cells from male animals and

perform our analysis on the log (1 + CPM) transformed single-cell count matrices. Note
that normalization of the single-cell data was done on a per cell basis. We defined two
group of cells based on age: young cells with age <= 3 months (Y) and old cells with age
> 3 months (O). For each gene we compute the $\log_2$ fold-change of cell and read counts
between O and Y. We defined cell count as the fraction of cells that express the gene.
Similarly, we defined read count as the mean read count of the gene in the cells that
express it. The calculated $\log_2$ fold-changes of a gene reflect its expression changes with
aging within the single-cell data. Next we analyze each gene based on the bulk data. We
computed the Spearman (Sp) correlation of bulk DESeq2 normalized gene expression
with aging. We defined two groups of genes based on the bulk data, increasing with age
Sp > 0.7 (U) and decreasing with age Sp < -0.7 (D). Finally, we compared the single-cell
data based $\log_2$ fold-changes between the bulk data defined groups U and D. Specifically,
we run Wilcoxon–Mann–Whitney test in order to understand if $\log_2$ fold-changes of cell
or read counts could distinguish between the two groups. We used the U statistic for
effect size.
645
T-Cell processing
We used TraCeR[34] version 0.5 to identify T-Cell clonal populations. We ran tracer
assemble with --species Mmus set. We then ran tracer summarise with –species Mmus
to create the final results. We used the following versions for TraCeR dependencies:
igblast version 1.7.0, kallisto version v0.43.1, Salmon version 0.8.2, Trinity version
v2.4.0, GRCm38 reference genome. Step-by-step instructions to reproduce the
processing of the data are available from GitHub.
653
654   **B-Cell processing**
655   We used singlecell-ige[33] version eafb6d126cc2d6511faae3efbd442abd7c6dc8ef
656   (https://github.com/dcroote/singlecell-ige) to identify B-Cell clonal populations. We
657   used the default configuration settings except we set the species to mouse. Step-by-step
658   instructions to reproduce the processing of the data are available from GitHub.
659
Mutation analysis
We used samtools[50] version 1.9 and GATK[29] version v4.1.1.0 for mutation analysis.
We used samtools faidx to create our index file. Then we used GATK
CreateSequenceDictionary and GRCm38, as the reference, to create our sequence
dictionary. Next we used GATK AddOrReplaceReadGroups to create a single read
group using parameters -RGID 4 -RGLB lib1 -RGPL illumina -RGPU unit1 -RGSM
20. Finally we used GATK HaplotypeCaller to call the mutations. We disabled the
following read filters: MappingQualityReadFilter, GoodCigarReadFilter,
NotSecondaryAlignmentReadFilter, MappedReadFilter,
MappingQualityAvailableReadFilter, NonZeroReferenceLengthAlignmentReadFilter,
NotDuplicateReadFilter, PassesVendorQualityCheckReadFilter, and
WellformedReadFilter, but kept all other default settings. The results were
summarized per gene in the form of a mutation count per cell table. We started by
removing genes mutated in over 60% of cells, to eliminate the possible bias of
germline mutations. Then for each tissue we selected genes expressed in at least 75%
of the cells for all the time points to avoid confounding the mutation results with

differential gene expression associated with age. Next we computed the average
number of mutations in the gene set (or ERCC spike-in controls) per cell and also the
average number of raw counts (Supplementary Table 6) and plotted the different
distributions. Step-by-step instructions to reproduce the processing of the data are
available from GitHub.
681
Diversity score
The raw FACS or droplet dataset were used as the input. We filtered genes expressed
in fewer than 5 cells, filtered cells if expressing fewer than 500 genes and discarded
cells with total number of counts less than 5000. Next we performed size factor
normalization such that every cell had 1e4 counts and performed a log1p
transformation. This was followed by clustering, where we clustered every tissue and
every tissue-cell type for every mouse separately using 6 different configurations:
resolution parameters (0.3, 0.5, 0.7) * clustering method (Louvain, Leiden). This is to
provide a robust clustering result. For each combination (each tissue-mouse and each
tissue-cell_type-mouse), we computed the clustering diversity score as the Shannon
entropy of the cluster assignment. We then regressed the diversity score against age to
detect the systematic increase/decrease of clustering diversity with respect to age. FDR
was used to correct for multiple comparisons. A tissue or a tissue-cell type was
selected if the slope was consistent (having the same sign) in all 6 clustering
configurations and at least 2 out of 6 clustering configurations had FDR<0.3. For each
selected tissue or tissue-cell type, a separate UMAP was computed using cells from all
mice for visualization using Leiden clustering with resolution parameter 0.7.
699
700　**Code availability**
701　All code used for analysis is available on GitHub (https://github.com/czbiohub/tabula-
702　muris-senis)
703
704　**Interactive Data Browsers**
705　tabula-muris-senis.ds.czbiohub.org
706　https://tabula-maris-senis.cells.ucsc.edu
707
708
709

References

1. López-Otín, C., Blasco, M. A., Partridge, L., Serrano, M. & Kroemer, G. The Hallmarks of Aging. *Cell* **153,** 1194–1217 (2013).

2. Nikolich-Žugich, J. The twilight of immunity: emerging concepts in aging of the immune system. *Nat Immunol* **19,** 10–19 (2018).

3. Campisi, J. Aging, Cellular Senescence, and Cancer. *Annu Rev Physiol* **75,** 685 705 (2013).

4. Vijg, J. & Suh, Y. Genome Instability and Aging. *Physiology* **75,** 645–668 (2013).

5. The Tabula Muris Consortium. Single-cell transcriptomics of 20 mouse organs creates a Tabula Muris. *Nature* **562,** 367–372 (2018).

6. Blondel, V. D., Guillaume, J.-L., Lambiotte, R. & Lefebvre, E. Fast unfolding of communities in large networks. (2008). doi:10.1088/1742-5468/2008/10/p10008

7. Traag, V., Waltman, L. & van Eck, N. From Louvain to Leiden: guaranteeing well-connected communities. *Sci Rep-uk* **9,** 5233 (2019).

8. Schaum, N. *et al*. The murine transcriptome reveals global aging nodes with organ-specific phase and amplitude. *Biorxiv* 662254 (2019). doi:10.1101/662254

9. Rayess, H., Wang, M. B. & vatsan, E. S. Cellular senescence and tumor suppressor gene p16. *Int J Cancer* **130,** 1715–1725 (2012).

10. Hernandez-Segura, A., Nehme, J. & Demaria, M. Hallmarks of Cellular Senescence. *Trends Cell Biol* **28,** 436–453 (2018).

11. van den Brink, S. C. *et al*. Single-cell sequencing reveals dissociation-induced gene expression in tissue subpopulations. *Nat Methods* **14,** 935–936 (2017).

12. Tung, P.-Y. *et al*. Batch effects and the effective design of single-cell gene expression studies. *Sci Rep-uk* **7,** 39921 (2017).

13. Nguyen, Q. H., Pervolarakis, N., Nee, K. & Kessenbrock, K. Experimental Considerations for Single-Cell RNA Sequencing Approaches. *Frontiers Cell Dev Biology* **06,** 108 (2018).

14. Daly, D. M. *et al*. Age related changes in afferent pathways and urothelial function in the male mouse bladder. *J Physiology* **592,** 537–549 (2014).

15. Burmeister, D. M., AbouShwareb, T., Bergman, C. R., Andersson, K.-E. & Christ, G.

J. Age-Related Alterations in Regeneration of the Urinary Bladder after Subtotal
Cystectomy. *Am J Pathology* **183,** 1585–1595 (2013).

16. Andersson, K.-E., Boedtkjer, D. B. & Forman, A. The link between vascular
dysfunction, bladder ischemia, and aging bladder dysfunction. *Ther Adv Urology* **9,** 11–
27   (2017).

17. Suskind, A. M. The Aging Overactive Bladder: a Review of Aging-Related Changes
from the Brain to the Bladder. *Curr Bladder Dysfunct Reports* **12,** 42–47 (2017).

18. Zhang, D. *et al*. Downregulation of ATP1A1 promotes cancer development in renal
cell carcinoma. *Clin Proteom* **14,** 15 (2017).

19. Isaka, Y. Epidermal growth factor as a prognostic biomarker in chronic kidney
diseases. *Ann Transl Medicine* **4,** S62–S62 (2016).

20. Devuyst, O., Olinger, E. & Rampoldi, L. Uromodulin: from physiology to rare and
complex kidney disorders. *Nat Rev Nephrol* **13,** 525–544 (2017).

21. Tokonami, N. *et al*. Uromodulin is expressed in the distal convoluted tubule, where it
is critical for regulation of the sodium chloride cotransporter NCC. *Kidney Int* **94,** 701–
715   (2018).

22. Covarrubias, A. J. *et al*. Aging-related inflammation driven by cellular senescence
enhances NAD consumption via activation of CD38+ pro-inflammatory macrophages.
Biorxiv 609438 (2019). doi:10.1101/609438

23. Tan, Q. *et al*. The Role of IL-1 Family Members and Kupffer Cells in Liver
Regeneration. *Biomed Res Int* **2016,** 6495793 (2016).

24. Gehrke, N. *et al*. Hepatocyte-specific deletion of IL1-RI attenuatesliver injury by
blocking IL-1 driven autoinflammation. *J Hepatol* **68,** 986–995 (2018).

25. Liu, Y., Gardner, C. R., Laskin, J. D. & Laskin, D. L. Classical and alternative
activation of rat hepatic sinusoidal endothelial cells by inflammatory stimuli. *Exp Mol*
Pathol **94,** 160–167 (2013).

26. Palmer, S., Albergante, L., Blackburn, C. C. & Newman, T. Thymic involution and
rising disease incidence with age. *Proc National Acad Sci* **115,** 201714478 (2018).

27. Shen, Q. *et al*. The AP-1 transcription factor regulates postnatal mammary gland
development. *Dev Biol* **295,** 589–603 (2006).

28. Girnius, N., Edwards, Y. J. & Davis, R. J. The cJUN NH2-terminal kinase (JNK)
pathway contributes to mouse mammary gland remodeling during involution. *Cell Death*
Differ **25,** 1702–1715 (2018).

29. McKenna, A. *et al*. The Genome Analysis Toolkit: A MapReduce framework for
analyzing next-generation DNA sequencing data. *Genome Res* **20,** 1297–1303 (2010).
804

805 30. DePristo, M. A. *et al*. A framework for variation discovery and genotyping using
806 next-generation DNA sequencing data. *Nat Genet* **43,** 491 (2011).
807

808 31. Auwera, G. A. *et al*. From FastQ Data to High Confidence Variant Calls: The
809 Genome Analysis Toolkit Best Practices Pipeline. *Current Protocols in Bioinformatics*
810 11.10.1-11.10.33 (2013). doi:10.1002/0471250953.bi1110s43
811

812 32. Zook, J. M., Samarov, D., niel, J., Sen, S. K. & Salit, M. Synthetic Spike-in Standards
813 Improve Run-Specific Systematic Error Analysis for DNA and RNA Sequencing. *Plos*
814 *One* **7,** e41356 (2012).
815

816 33. Croote, D., Darmanis, S., Nadeau, K. C. & Quake, S. R. High-affinity allergen-
817 specific human antibodies cloned from single IgE B cell transcriptomes. *Science* **362,**
818 1306–1309   (2018).
819

820 34. Stubbington, M. J. *et al*. T cell fate and clonality inference from single-cell
821 transcriptomes. *Nat Methods* **13,** nmeth.3800 (2016).
822

823 35. Goronzy, J. J. & Weyand, C. M. Understanding immunosenescence to improve
824 responses to vaccines. *Nat Immunol* **14,** ni.2588 (2013).
825

826 36. Goronzy, J. J. & Weyand, C. M. Successful and Maladaptive T Cell Aging. *Immunity*
827 **46,** 364–378 (2017).
828

829 37. Keren-Shaul, H. *et al*. A Unique Microglia Type Associated with Restricting
830 Development of Alzheimer's Disease. *Cell* **169,** 1276-1290.e17 (2017).
831

832 38. Li, Q. *et al*. Developmental Heterogeneity of Microglia and Brain Myeloid Cells
833 Revealed by Deep Single-Cell RNA Sequencing. *Neuron* **101,** 207-223.e10 (2019).
834

835 39. Hammond, T. R. *et al*. Single-Cell RNA Sequencing of Microglia throughout the
836 Mouse Lifespan and in the Injured Brain Reveals Complex Cell-State Changes. *Immunity*
837 **50,** 253-271.e6 (2019).
838

839 40. Jablonski, K. A. *et al*. Novel Markers to Delineate Murine M1 and M2 Macrophages.
840 *Plos One* **10,** e0145342 (2015).
841

842 41. Wolf, A. F. *et al*. PAGA: graph abstraction reconciles clustering with trajectory
843 inference through a topology preserving map of single cells. *Genome Biol* **20,** 59 (2019).
844

42. Picelli, S. *et al*. Smart-seq2 for sensitive full-length transcriptome profiling in single
cells. *Nat Methods* **10,** nmeth.2639 (2013).

43. Darmanis, S. *et al*. A survey of human brain transcriptome diversity at the single cell
level. *Proc National Acad Sci* **112,** 7285–7290 (2015).

44. Picelli, S. *et al*. Tn5 transposase and tagmentation procedures for massively scaled
sequencing projects. *Genome Res* **24,** 2033–2040 (2014).

45. Hennig, B. P. *et al*. Large-Scale Low-Cost NGS Library Preparation Using a Robust
Tn5 Purification and Tagmentation Protocol. *G3 Genes Genomes Genetics* **8,** 79–89
(2018).

46. Wolf, A. F., Angerer, P. & Theis, F. J. SCANPY: large-scale single-cell gene
expression data analysis. *Genome Biol* **19,** 15 (2018).

47. Diehl, A. D. *et al*. The Cell Ontology 2016: enhanced content, modularization, and
ontology interoperability. *J Biomed Semant* **7,** 44 (2016).

48. Finak, G. *et al*. MAST: a flexible statistical framework for assessing transcriptional
changes and characterizing heterogeneity in single-cell RNA sequencing data. *Genome*
Biol **16,** 278 (2015).

49. McQuin, C. *et al*. CellProfiler 3.0: Next-generation image processing for biology.
Plos Biol **16,** e2005970 (2018).

50. Li, H. *et al*. The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25,**
2078–2079 (2009).

879 **Supplementary Information** is available in the online version of the paper.
880
881

Acknowledgements We thank Sony Biotechnology for making an SH800S instrument
available for this project. Some cell sorting/flow cytometry analysis for this project was
done on a Sony SH800S instrument in the Stanford Shared FACS Facility. Some
fluorescence activated cell sorting (FACS) was done with instruments in the VA Flow
Cytometry Core, which is supported by the US Department of Veterans Affairs (VA),
Palo Alto Veterans Institute for Research (PAVIR), and the National Institutes of Health
(NIH). We would like to thank Bruno Tojo for helping with the artwork.
889
890

891 **Author Contributions**
892 **Overall Coordination:** Angela Oliveira Pisco[1], Aaron McGeever[1], Nicholas Schaum[2],
893 Jim Karkanias[1], Norma F. Neff[1], Spyros Darmanis[1]*, Tony Wyss-Coray[3-5]*, and Stephen
894 R. Quake[1,6]*
895

896 * Correspondence to: steve@quake-lab.org, twc@stanford.edu,
897 spyros.darmanis@czbiohub.org
898
899

Organ collection and processing: Jane Antony[2], Ankit S. Baghel[2], Isaac Bakerman[2,7,8],
Ishita Bansal[3], Daniela Berdnik[5], Biter Bilen[3,4], Douglas Brownfield[9], Corey Cain[10],
Michelle B. Chen[4], Stephanie D. Conley[2], Spyros Darmanis[1], Aaron Demers[2], Kubilay
Demir[2,11], Antoine de Morree[3,4], Tessa Divita[1], Haley du Bois[5], Laughing Bear Torrez
Dulgeroff[2], Hamid Ebadi[1], F. Hernán Espinoza[9], Matt Fish[2,11,12], Qiang Gan[3,4], Benson M.
George[2], Astrid Gillich[9], Foad Green[1], Geraldine Genetiano[1], Xueying Gu[12], Gunsagar S.
Gulati[2], Yan Hang[12], Shayan Hosseinzadeh[1], Albin Huang[4], Tal Iram[4], Taichi Isobe[1],
Feather Ives[2], Robert Jones[3], Kevin S. Kao[2], Guruswamy Karnam[13], Aaron M. Kershner[2],
Nathalie Khoury[3], Bernhard M. Kiss[2,14], William Kong[2], Maya E. Kumar[15,16], Jonathan
Lam[12], Davis P. Lee[6], Song E. Lee[4], Olivia Leventhal[5], Guang Li[17], Qingyun Li[18], Ling
Liu[3,4], Annie Lo[1], Wan-Jin Lu[1,9], Maria F. Lugo-Fagundo[5], Anoop Manjunath[1], Andrew
P. May[1], Ashley Maynard[1], Marina McKay[1], M. Windy McNerney[32,33],Ross J.
Metzger[19,20], Marco Mignardi[1], Dullei Min[21], Ahmad N. Nabhan[9], Norma F. Neff[1],
Katharine M. Ng[3], Joseph Noh[2], Rasika Patkar[13], Weng Chuan Peng[12], Lolita Penland[1],
Robert Puccinelli[1], Eric J. Rulifson[12], Nicholas Schaum[2], Michael Seamus Haney[3],
Shaheen S. Sikandar[2], Rahul Sinha[2,22-24], Rene V. Sit[1], Daniel Staehli[3], Krzysztof
Szade[2,25], Weilun Tan[1], Cristina Tato[1], Krissie Tellez[12], Kyle J. Travaglini[9], Carolina
Tropini[26], Lucas Waldburger[1], Linda J. van Weele[2], Michael N. Wosczyna[3,4], Jinyi
Xiang[2], Soso Xue[3], Andrew C. Yang[3], Lakshmi P. Yerra[5], Justin Youngyunpipatkul[1],
Fabio Zanini[3], Macy E. Zardeneta[6], Fan Zhang[19], Hui Zhang[5], Lu Zhou[18]
920

921 **Library preparation and sequencing:** Spyros Darmanis[1], Shayan Hosseinzadeh[1],
922 Ashley Maynard[1], Norma F. Neff[1], Lolita Penland[1], Rene V. Sit[1], Michelle Tan[1], Weilun
923 Tan[1], Alexander Zee[1]
924

Computational Data Analysis: Oliver Hahn[3], Lincoln Harris[1], Andreas Keller[3,36], Benoit Lehallier[3], Aaron McGeever[1], Angela Oliveira Pisco[1], Róbert Pálovics[3], Weilun Tan[1], Martin Zhang[30,37]

Cell Type Annotation: Nicole Almanzar[21], Jane Antony[2], Biter Bilen[3,4], Weng Chuan Peng[12], Spyros Darmanis[1], Antoine de Morree[3,4], Oliver Hahn[3], Yan Hang[12], Mu He[31], Shayan Hosseinzadeh[1], Tal Iram[4], Taichi Isobe[1], Aaron M. Kershner[1], Jonathan Lam[12], Guang Li[17], Qingyun Li[18], Ling Liu[3,4], Wan-Jin Lu[1,9], Ashley Maynard[1], Dullei Min[21], Ahmad N. Nabhan[9], Patricia K. Nguyen[2,7,8,17], Angela Oliveira Pisco[1], Zhen Qi[2], Nicholas Schaum[2], Joe M. Segal[13], Shaheen S. Sikandar[2], Rahul Sinha[1,22-24], Rene Sit[1], Michelle Tan[1], Weilun Tan[1], Kyle J. Travaglini[9], Margaret Tsui[13], Bruce M. Wang[13], Linda J. van Weele[2], Michael N. Wosczyna[3,4], Jinyi Xiang[2], Alexander Zee[1], Lu Zhou[18]

Liver staining and data analysis: Rafael Gòmez-Sjöberg[1], Angela Oliveira Pisco[1], Joe M. Segal[13], Margaret Tsui[13], Kevin A Yamauchi[1]

Microbiome analysis: Bryan Merrill[26], Aaron McGeever[1], Katharine M. Ng[3], Angela Oliveira Pisco[1], Carolina Tropini[26], Brian Yu[1], Chunyu Zhao[1], Katherine Pollard[34], Justin Sonnenburg[26], Kerwyn Casey Huang[2,3]

Writing Group: Spyros Darmanis[1], Angela Oliveira Pisco[1], Stephen R. Quake[1,6], Tony Wyss-Coray[3-5]

Principal Investigators: Ben A. Barres[18], Philip A. Beachy[2,9,11,12], Charles K. F. Chan[28], Michael F. Clarke[2], Spyros Darmanis[1], Kerwyn Casey Huang[2,3,26], Jim Karkanias[1], Seung K. Kim[12,29], Mark A. Krasnow[9,11], Maya E. Kumar[15,16], Christin S. Kuo[9,11,21], Ross J. Metzger[19,20], Norma F. Neff[2], Roel Nusse[9,11,12], Patricia K. Nguyen[2,7,8,17], Thomas A. Rando[3-5], Justin Sonnenburg[26], Bruce M. Wang[13], Kenneth Weinberg[21], Irving L. Weissman[2,22-24], Sean M. Wu[2,7,17], James Zou[1,30,35], Stephen R. Quake[2,6], Tony Wyss-Coray[3-5]

[1] Chan Zuckerberg Biohub, San Francisco, California, USA
[2] Institute for Stem Cell Biology and Regenerative Medicine, Stanford University School of Medicine, Stanford, California, USA
[3] Department of Neurology and Neurological Sciences, Stanford University School of Medicine, Stanford, California, USA
[4] Paul F. Glenn Center for the Biology of Aging, Stanford University School of Medicine, Stanford, California, USA
[5] Center for Tissue Regeneration, Repair, and Restoration, V.A. Palo Alto Healthcare System, Palo Alto, California, USA
[6] Department of Bioengineering, Stanford University, Stanford, California, USA
[7] Stanford Cardiovascular Institute, Stanford University School of Medicine, Stanford, California, USA
[8] Department of Medicine, Division of Cardiology, Stanford University School of Medicine, Stanford, California, USA

[9] Department of Biochemistry, Stanford University School of Medicine, Stanford,
California, USA
[10] Flow Cytometry Core, V.A. Palo Alto Healthcare System, Palo Alto, California, USA
[11] Howard Hughes Medical Institute, USA
[12] Department of Developmental Biology, Stanford University School of Medicine,
Stanford, California, USA
[13] Department of Medicine and Liver Center, University of California San Francisco, San
Francisco, California, USA
[14] Department of Urology, Stanford University School of Medicine, Stanford, California,
USA
[15] Sean N. Parker Center for Asthma and Allergy Research, Stanford University School
of Medicine, Stanford, California, USA
[16] Department of Medicine, Division of Pulmonary and Critical Care, Stanford University
School of Medicine, Stanford, California, USA
[17] Department of Medicine, Division of Cardiovascular Medicine, Stanford University,
Stanford, California, USA
[18] Department of Neurobiology, Stanford University School of Medicine, Stanford, CA
USA
[19] Vera Moulton Wall Center for Pulmonary and Vascular Disease, Stanford University
School of Medicine, Stanford, California, USA
[20] Department of Pediatrics, Division of Cardiology, Stanford University School of
Medicine, Stanford, California, USA
[21] Department of Pediatrics, Stanford University school of Medicine, Stanford,
California, USA
[22] Department of Pathology, Stanford University School of Medicine, Stanford,
California, USA
[23] Ludwig Center for Cancer Stem Cell Research and Medicine, Stanford University
School of Medicine, Stanford, California, USA
[24] Stanford Cancer Institute, Stanford University School of Medicine, Stanford,
California, USA
[25] Department of Medical Biotechnology, Faculty of Biophysics, Biochemistry and
Biotechnology, Jagiellonian University, Poland
[26] Department of Microbiology & Immunology, Stanford University School of Medicine,
Stanford, California, USA
[27] Department of Biochemistry and Biophysics, University of California San Francisco,
San Francisco, California USA
[28] Department of Surgery, Division of Plastic and Reconstructive Surgery, Stanford
University, Stanford, California USA
[29] Department of Medicine and Stanford Diabetes Research Center, Stanford University,
Stanford, California USA
[30] Department of Electrical Engineering, Stanford University, Palo Alto, 94304 USA
[31] Department of Physiology, University of California, San Francisco, CA 94158
[32] Mental Illness Research Education and Clinical Center, V.A. Palo Alto Healthcare
System, Palo Alto, California, USA
[33] Department of Psychiatry, Stanford University School of Medicine, Stanford,
California, USA

[34] Department of Epidemiology and Biostatistics, University of California, San Francisco,
CA 94158
[35] Department of Biomedical Data Science, Stanford University, Palo Alto, 94304 USA
[36] Clinical Bioinformatics, Saarland University, Saarbrücken, Germany
[37] Department of Epidemiology, Harvard T.H. Chan School of Public Health,
Boston, Massachusetts, USA

Tabula Muris Senis Fig. 1

Tabula Muris Senis Fig.2

Tabula Muris Senis Fig.3

Pairwise difference of mean number of mutations in gene set/ERCC across all tissues

Tabula Muris Senis Fig.4

a

6050 cells with assembled BCR

- 3m (n=1818 cells)
  - 3m singleton B cell
  - 3m clonal families
- 18m (n=1356 cells)
  - 18m singleton B cell
  - 18m clonal families
- 24m (n=2876 cells)
  - 24m singleton B cell
  - 24m clonal families

91% · 9% · 20% · 62% · 38% · 80%

b

6000 cells with assembled TCR

- 3m (n=2076 cells)
  - 3m singleton B cell
  - 3m clonal families
- 18m (n=2056 cells)
  - 18m singleton B cell
  - 18m clonal families
- 24m (n=1868 cells)
  - 24m singleton B cell
  - 24m clonal families

97% · 3% · 23% · 77% · 81% · 19%

c

Diversity score change with age

Kidney Macrophage (droplet)
Brain Myeloid Microglial cell (FACS)

resolution 0.3 0.5 0.7
method Louvain Leiden Louvain Leiden Louvain Leiden

−0.025 / −0.015 / −0.005

d

FACS Brain Myeloid Microglial cell

Cluster 12 (98 cells)
3m:6%, 18m:48%, 24m:45%

Cluster 1 (2464 cells)
3m:78%, 18m:9%, 24m:11%

Cluster 10 (163 cells)
3m:8%, 18m:5%, 24m:85%

Cluster 6 (1004 cells)
3m:92%, 18m:5%, 24m:1%

Cluster 14 (88 cells)
3m:4%, 18m:0%, 24m:95%

Age
3m
18m
24m

UMAP2 / UMAP1

e

FACS Brain Myeloid Microglial cell

- Alzheimer's disease signature
- Healthy microglia signature

UMAP2 / UMAP1

f

Droplet Kidney Macrophage

Age
young (1m, 3m)
old (18m,21m,24m,30m)

Cluster 13 (513 cells)
Young:15%; Old:84%

Cluster 10 (753 cells)
Young:63%; Old:36%

UMAP2 / UMAP1

Tabula Muris Senis Extended Data Fig. 1

Tabula Muris Senis Extended Data Fig. 2

Tabula Muris Senis Extended Data Fig. 3

a

FACS

b

Droplet

Legend:
- bulk signal explained neither by single-cell read nor by cell count
- bulk signal explained by read count only
- bulk signal explained by cell count only
- bulk signal explained by both single-cell read and cell count

Tabula Muris Senis Extended Data Fig. 4

Tabula Muris Senis Extended Data Fig. 5

Tabula Muris Senis Extended Data Fig. 6

Tabula Muris Senis Extended Data Fig.7

Tabula Muris Senis Extended Data Fig.8

Tabula Muris Senis Extended Data Fig.9

Tabula Muris Senis Extended Data Figure 10

Tabula Muris Senis Extended Data Figure 11

Tabula Muris Senis Supplementary Fig. 1

Tabula Muris Senis Supplementary Fig. 2

Tabula Muris Senis Supplementary Fig. 3