Title:

Predicting transfer RNA gene activity from genome variation data

Authors and Affiliation:

Bryan Thornlow[1], Joel Armstrong[1,2], Andrew Holmes[1,2], Russell Corbett-Detig[1,2], Todd Lowe[1,2]

[1]Department of Biomolecular Engineering, University of California, Santa Cruz, CA 95064; [2]Genomics Institute, University of California, Santa Cruz, CA 95064

ABSTRACT:

Transfer RNA (tRNA) genes are among the most highly transcribed genes in the genome due to their central role in protein synthesis. However, there is evidence for a broad range of gene expression across tRNA loci. This complexity, combined with post-transcriptional modifications and high sequence identity across transcripts, has severely limited our collective understanding of tRNA gene expression regulation and evolution. We establish correlates to tRNA gene expression and develop a tRNA gene classification method that minimally requires a single reference genome per species of interest and achieves accuracy comparable to epigenomic and ChIP-Seq assays. We observe that GC content and CpG density surrounding tRNA loci is exceptionally well correlated with tRNA gene activity, supporting a prominent regulatory role of the local genomic context in combination with internal sequence features. We use our classifications in conjunction with a novel tRNA gene ortholog set spanning 29 placental mammals to infer that tRNA gene expression regulation evolves slowly. The success of our method has fundamental implications to tRNA annotation and variant prioritization. Its simplicity and robustness suggests facile application to other clades and timescales, as well as to explore functional diversification of members of large gene families.

INTRODUCTION:

Transfer RNAs (tRNAs) are essential for the translation of messenger RNA (mRNA) into protein across all domains of life. At the gene level, they are of particular interest for their high copy number, strong nucleotide sequence conservation, and variation in expression (Pan 2018; Kutter et al. 2011; Schmitt et al. 2014). tRNA molecules are required in large abundance to meet the dynamic metabolic needs of cells, and tRNA genes may therefore be among the most highly transcribed genes in the genome (Palazzo and Lee 2015; Boivin et al. 2018).

Despite high cellular demands, several individual tRNA genes have no evidence for expression. High duplication rates and consequent weakened purifying selection might lead to an abundance of pseudogenes. Additionally, many of these genes may be tRNA-derived short interspersed nuclear elements (SINEs), which often retain their promoter elements. However, even after removal of apparent pseudogenes and SINEs, more than 60 human tRNA genes and over 100 mouse tRNA genes have no epigenomic evidence for expression in any of more than 100 tissues and cell lines (Thornlow et al. 2018). Chromatin-immunoprecipitation sequencing (ChIP-Seq) data supports this conclusion, as a recent study detected expression of only 224 of 417 high-confidence tRNA genes in human liver, with other mammals showing similar patterns (Kutter et al. 2011).

The evolution of tRNA gene expression may underlie phenotypic differences between species. Data from prior studies suggests that the rate of evolution of protein coding gene expression levels differs by clade (Necsulea and Kaessmann 2014; Brawand et al. 2011; Li et al. 1996). The rate of evolution of gene expression also varies among non-coding RNA gene families (Meunier et al. 2013; Necsulea and Kaessmann 2014; Necsulea et al. 2014). Due to difficulties in quantifying tRNA gene expression, the conservation of tRNA gene expression across mammals is not well characterized. The expanding functional repertoire of tRNA transcripts

and fragments (Kirchner and Ignatova 2015; Goodarzi et al. 2015; Mleczko et al. 2014; Sun et al. 2018) indicates that changes in tRNA gene expression between species could have profound phenotypic effects.

Expression of tRNA genes has clear importance for organismal development and disease treatment, but our understanding of its regulation and evolution is severely lacking for several reasons. Measuring expression of unique mRNA transcripts is straightforward as these molecules can be directly sequenced. However, tRNA sequencing by traditional methods is impeded by post-transcriptional modifications. tRNAs are the most modified RNAs, and only very recently have specialized library preparation methods been developed to remove these modifications, enabling quantitative sequencing (Cozen et al. 2015; Zheng et al. 2015). Furthermore, because many tRNA transcripts are identical at the sequence level, perfect transcript abundance measurements may not be sufficient to annotate tRNA gene expression. In order to understand how individual tRNA genes are regulated, the study of the local genomic context is therefore essential.

Several alternative genome-wide methods examine tRNA loci in their genomic context to bypass the problem of identical mature tRNA transcripts. Such assays include chromatin immunoprecipitation (ChIP; Bogu et al. 2015; Roadmap Epigenomics Consortium et al. 2015; Thornlow et al. 2018), RNA Polymerase III (Pol3) ChIP-Seq (Kutter et al. 2011), and ATAC-Seq (Foissac et al. 2018), among others. These high-throughput assays are cost- and resource-intensive, so resulting data are often limited to few species and tissues. Nonetheless, these data indicate that identical tRNA genes vary in expression (Pan 2018; Kutter et al. 2011; Schmitt et al. 2014), suggesting that tRNA gene expression may be regulated by factors extrinsic to tRNA genes.

Here, we resolve many of these concerns by developing a model to predict which tRNA genes are actively transcribed and which are inactive. Our previous work has shown that relative tRNA expression may be inferred based on DNA variation driven by transcription-associated mutagenesis (Thornlow et al. 2018). We leverage this correlation and other genomic correlates to gene expression to infer expression of tRNA genes. Our method requires only one target genome in order to predict its constituent tRNA gene activity, so that it may be applied broadly. We demonstrate our method using 29 placental mammal genomes, many of which have no tRNA expression data. We also present a set of tRNA gene orthologs encompassing the tRNA genes of all 29 species. We use this in conjunction with our expression classifications to infer that tRNA gene function appears to evolve slowly and that changes between predicted activity states are rare across mammalian evolution.

RESULTS:

Our classifier uses only data obtainable from reference genomes and alignments. Our goal was to develop a model that could be applied to as many species as possible. We therefore used only features obtainable from a reference genome. We explored features correlated with gene activity, and then reduced this set to just those yielding the best predictions for our training data.

Towards this, we evaluated and incorporated three types of function-predictive statistics. First, mutation and selection should leave genomic signatures in genome variation data. Our previous study found that tRNA gene conservation is highest for actively transcribed tRNA genes presumably due to stronger purifying selection. Conversely, the putatively neutral flanking regions are more divergent in highly expressed tRNA genes due to transcription-associated mutagenesis (Thornlow et al. 2018). Thus, we designed our classifier to incorporate features of sequence variation in and around tRNA genes.

Second, we reasoned that highly expressed tRNA genes should encode strong internal promoter sequences and their transcripts should fold to yield a functional mature tRNA. Both primary sequence and secondary structure are included in tRNAscan-SE bit scores (Chan and Lowe 2019). We also assessed the correlation of the length of each pre-tRNA's 3' tail, measured by the distance from the end of the tRNA gene sequence to the beginning of its transcription termination sequence (Allison and Hall 1985; Koski et al. 1980), with tRNA gene activity. Multiple studies on tRNA transcription termination (Maraia et al. 1994; Hamada et al. 2000; Orioli et al. 2011; Arimbasseri et al. 2013) observed that transcription sites vary in sequence, length, and proximity to their associated gene, and that distantly located or weak termination sequences allow for

significant readthrough, which might lead to coupling of tRNA maturation and production of downstream-encoded RNA transcripts. We found that both tRNAscan-SE general bit scores (Chan and Lowe 2019) and distance to transcription termination sites are significantly correlated with tRNA gene activity based on epigenomic data (Spearman rank correlation, p < 1e-5 for all comparisons).

Third, because mRNA expression depends on local chromatin context, we explored features of the genomic environment. Protein coding genes in regions rich in CpG (or CG) dinucleotides, are more frequently expressed (Gardiner-Garden and Frommer 1987; Krinner et al. 2014). Gardiner-Garden and Frommer (1987) defined CpG Islands scores, which measure the observed frequency of CpG dinucleotides compared to their expected frequency given the GC content of the region. We found that these scores are significantly correlated with tRNA gene expression (Spearman rank correlation, p < 8e-30). We also found that tRNA genes are more likely to be expressed if adjacent to protein-coding genes (p < 4.7e-5) or other tRNA genes (p < 1.36e-6). We therefore expected that a combination of both intrinsic and extrinsic features could enable reliable inference of tRNA gene expression.

To choose a model, we tested several common frameworks, including random forest (RF), logistic regression, and support vector machines. The RF classifier was most effective, achieving the greatest area under the Receiver Operating Characteristic (ROC) curve (AUC, Figure 1) based on ten-fold cross-validation of the human tRNA gene data and application to mouse tRNA gene data (see Methods). We developed our initial model with 30 features (Supplementary Table 1) and gradually reduced it to 10 (Table 1), using the correlation between each feature and the activity labels, as well as intercorrelations between features (Hall 1998; Hall et al. 2009).

Figure 1: Random forest outperforms other standard binary classification methods. Receiver operating characteristic (ROC) curves for random forest (blue), logistic regression (red) and support vector machine (yellow) upon application to human training data with ten-fold cross-validation (A) and mouse test data (B).

Features derived from CpG islands are most informative. To better understand our classifier, we determined the relative importance of each feature, in terms of the normalized mean decrease in node impurity across all decision trees in the model (Table 1; Pedregosa et al. 2011). We also used the Spearman rank correlation between each feature and the activity measurements to assess the strength of each feature. While all features contribute to model accuracy and are significantly correlated with the activity labels (Spearman rank correlation, p < 1e-5 for all features), the most informative features are derived from CpG islands at each

tRNA locus, extolling the importance of chromosomal context and the ability to regulate expressed tRNA genes by altering methylation state (Gardiner-Garden and Frommer 1987).

While tRNAscan-SE bit scores reflect elements of primary sequence and secondary structure known to be associated with functional tRNA genes, other intrinsic features are comparably informative of tRNA activity (Table 1). However, tRNAscan-SE bit scores are used in the tRNAscan-SE high-confidence filter, which was applied to each tRNA gene set prior to classification (Chan and Lowe 2019). This may lead to underestimation of its relative importance to the classifier. Nonetheless, the importance of CpG density suggests that it is the chromatin environment more than the tRNA gene sequence that regulates tRNA gene expression, although tRNA sequence features likely also play important roles.

	Feature	Importance
Extrinsic	Average PhyloP Score in tRNA Gene Sequence	0.068
	Average PhyloP Score in 5' Flanking Region	0.087
	CpG Density Across tRNA Locus	0.322
	Exons Within 75 Kilobases	0.017
	Observed/Expected CpG Islands Score Upstream of tRNA Gene	0.232
	tRNA Genes Within 10 Kilobases	0.020
Intrinsic	Canonical tRNA Secondary Structure	0.069
	Distance To Nearest TTTT Transcription Termination Sequence	0.053
	tRNA Decoding Redundancy	0.077
	tRNAscan-SE General Bit Score	0.055

Table 1: Both intrinsic (tRNA-specific) and extrinsic (genome context) features are integral to the model. The relative importance of each feature as measured by decrease in node impurity by scikit-learn (Pedregosa et al. 2011) is shown (see Methods). Higher importance scores indicate greater contribution to discrimination between active and inactive tRNA genes by the model. tRNA decoding redundancy refers to the number of tRNA genes sharing the same anticodon, and canonical tRNA secondary structure is based on constrained minimum free energy (Lorenz et al. 2011).

Our classifier is at least 90% accurate in classifying mouse tRNA genes. We tested the accuracy of our classifier using epigenomic and tRNA expression data. Our activity classifications were based on chromatin-IP data for all human and mouse tRNAs (Roadmap Epigenomics Consortium et al. 2015; Bogu et al. 2015). For both species, we excluded tRNA genes for which epigenomic data was not available, and tRNA genes contained within large segmental duplications (see Methods). Our training set includes 371 human tRNA genes, 304 active and 67 inactive. The activity labels were determined using chromatin-IP (ChIP) data from Roadmap Epigenomics Consortium et al. (2015) for human tissues and analogous mouse data from Bogu et al. (2015). All tRNA genes located near active transcription start sites in at least one tissue were considered active. All others were considered inactive. Using ten-fold cross validation, we achieved 93.0% accuracy, correctly classifying 345 of 371 human tRNA genes. Upon applying this same model, without re-training, to the set of 387 mouse tRNA genes (262 active, 125 inactive), we achieved 92.0% accuracy, correctly classifying 356 tRNA genes (Figure 2A-C). We did not identify a pattern connecting the 26 misclassified human tRNA genes or the 31 misclassified mouse tRNA genes (Supplementary Table 2).

Classification without alignment or annotation is similarly accurate. We ensured that our approach is applicable to any species with a sequenced genome. Here, we used Cactus graphs to generate alignments, but we recognize that Cactus graphs are not yet ubiquitous among genomic studies. To accommodate species for which no alignments or annotations have been developed, we have included an option to remove alignment- and annotation-based features. Use of this model led to decreases in accuracy in

both human (AUC = 0.932, 92.5% accuracy compared to AUC = 0.942 and 93.0% accuracy in the full model) and mouse (AUC = .935, 90.1% accuracy compared to AUC = 0.964 and 92.0% accuracy in the full model), which may be exacerbated upon application to more phylogenetically distant species.

Figure 2: Classification of expression based on genomic data achieves similar results to ChIP-Seq expression analysis in four species. The number of mouse tRNA genes predicted active (A) and inactive (B) compared to the number of tissues in which they are actively transcribed according to Bogu et al. (2015). We consider any mouse tRNA gene active if it is actively transcribed in at least one tissue. Probability scores output by the classifier are shown on the x-axis where tRNA genes further left are predicted inactive with greater probability and tRNA genes further right are predicted active with greater probability. The y-axis shows Pol3 ChIP-Seq read counts from Kutter et al. (2011) for each tRNA gene in mouse (C), macaque (D), rat (E) and dog (F).

Previously collected ChIP-Seq and ATAC-Seq data validate our classifications. We used ChIP-Seq data from Kutter et al. (2011) to validate our model. Kutter et al conducted ChIP-Seq on the livers of six species using RNA Polymerase III (Pol3). We compared our tRNA gene classifications for four of these species (*Mus musculus*, *Macaca mulatta*, *Rattus norvegicus* and *Canis lupus familiaris*) to their Pol3 ChIP-Seq data. We found comparable agreement between our classifications, the activity states derived from epigenomic ChIP data, and the Pol3 ChIP-Seq read counts (Figure 2). We also found that the relationship between the probability scores in our classifications and the Pol3 ChIP-Seq read counts is consistent across all four species, suggesting similar accuracy in classification (Figure 2C-F). Our predictions are also consistent and similarly accurate when compared to mouse muscle and testes ChIP-Seq data (Supplementary Figure 1).

We predicted many tRNA genes as active despite a lack of Pol3-based expression in the liver. This may be because our model does not account for differential expression across tissues and is instead trained to

predict tRNA genes as active if epigenomic data indicates expression in at least one tissue. For example, in the mouse genome, 262 total tRNA genes are active in at least one tissue based on the epigenomic data, but 89 of these are not expected to be active in the liver based on the same data. Likewise, our model predicted that 122 macaque tRNA genes, 149 dog tRNA genes and 68 rat tRNA genes are active despite no recorded Pol3 activity in the liver (Kutter et al. 2011). Although chromatin-IP has not been performed on macaque, rat and dog tRNA loci, we find that virtually all tRNA genes with measured Pol3-based expression are predicted to be active by our classifier. Among tRNA genes with Pol3 ChIP-Seq read-counts greater than zero, we predicted that 95.3% are active in mouse, 97.7% in macaque, 98.9% in rat and 98.5% in dog. This consistency in tRNA distributions and classifier behavior across species suggests that the classifier is similarly accurate in mouse, macaque, rat and dog.

To further validate our predictions, we used ATAC-seq data (Foissac et al. 2018), captured in liver, CD4 and CD8 cells for the cow, pig and goat genomes. We measured the ATAC-Seq peak across these tissues for the regions up to 250 bases up- and down-stream of each tRNA gene (Supplementary Figure 2). This data spans only a small subset of tissues, and many tRNA loci that do not show activity in these tissues but were predicted active by our model may be active in other tissues. Among tRNA genes with ATAC-Seq read-counts greater than zero, we predicted 90.4% as active in cow, 95.1% in goat and 90.8% in pig. These results are comparable to measurements obtained from ChIP-Seq data in mouse, macaque, rat and dog. This slight decrease in accuracy may be due to differences in resolution between ATAC-Seq and ChIP-Seq, or the increased phylogenetic distance from human to the ungulate species, relative to rodent species.

tRNA gene classifications follow similar distributions across the eutherian phylogeny. Following validation upon comparison to previously collected data, we applied our model to 29 mammalian species (Figure 3). We also determined the distributions of active and inactive tRNA genes by anticodon across these species (Supplementary Table 3). Most species have approximately 250-350 predicted active genes, comprising roughly 75% of their tRNA gene sets. We observe similar distributions by clade, with few exceptions. Notably, *Bos taurus*, *Capra hircus* and *Orcinus orca* (cow, goat and orca, respectively) have more than 300 tRNA genes predicted inactive while no other species has more than 156. This is most likely the result of a decreased ability of tRNAscan-SE to discriminate tRNA genes from tRNA-derived SINEs (short interspersed nuclear elements) in these species (Chan and Lowe 2019). Consistent with Tang et al. (2009), we identified 406 active cow tRNA genes, and 437 when including those found in segmental duplications (Supplementary Table 4), the most of any species in our phylogeny. Of the 439 cow tRNA genes identified by (Tang et al. 2009), we find 396 genes with exactly matching sequences in our tRNA gene set, and classify 346 of these as active. tRNA gene sequences found by Tang et al. (2009) that were not found in our tRNA set are likely due to differences in genome assembly.

We also ensured that all species had at least one tRNA predicted active gene for each expected anticodon (Supplementary Table 3). This excludes anticodons with no known active tRNA genes in human or mouse. We observed three exceptions to this rule. *C. hircus* is the only species without a predicted active tRNA-Ser-TGA gene. *C. hircus* is also one of only three species that has any tRNA-Ser-GGA genes, and it has 27 such genes, one of which was predicted active. *C. hircus* may use one or several of these genes to translate TCA codons through wobble pairing. We also predict that the lone tRNA-Leu-TAA gene in *Erinaceus europaeus* (hedgehog) is inactive. The probability score for this prediction is 0.51, nearly the minimum possible score, indicating that it may be active. Additionally, the *Chinchilla lanigera* (chinchilla) genome does not contain any high-confidence selenocysteine (SeC) tRNA genes. This is likely due to an incomplete genome assembly, as selenocysteine tRNAs are required in mammals (Johansson et al. 2005).

		Final		
A		Missing	Inactive	Active
Initial	Missing	.871	.111	.017
	Inactive	.092	.859	.049
	Active	.012	.014	.974

B Species	Active	Inactive
Homo sapiens	314	57
Pan troglodytes	339	56
Gorilla gorilla gorilla	325	46
Pongo abelii	351	50
Macaca mulatta	337	54
Aotus nancymaae	285	43
Microcebus murinus	284	40
Octodon degus	240	58
Chinchilla lanigera	250	51
Cavia porcellus	266	54
Heterocephalus glaber	238	56
Mus musculus	275	112
Rattus norvegicus	245	102
Microtus ochrogaster	343	127
Jaculus jaculus	246	72
Marmota marmota	253	156
Oryctolagus cuniculus	377	70
Bos taurus	406	369
Capra hircus	216	317
Orcinus orca	328	333
Sus scrofa	294	104
Canis lupus familiaris	278	90
Mustela putorius furo	291	87
Equus caballus	351	65
Pteropus alecto	222	123
Eptesicus fuscus	199	15
Erinaceus europaeus	343	104
Dasypus novemcinctus	379	149
Chrysochloris asiatica	275	97

20 million years

Figure 3: Changes between activity states are rare among placental mammal tRNA genes. Transition probabilities between each predicted activity state over a branch length of 10 million years (A). The number of tRNAs in each class are shown alongside the phylogenetic tree of species involved in this study (B; Hedges et al. 2006).

Outlier tRNA genes may become foci of future experimentation. We identified several anomalous lineage-specific expansions. Notably, *E. europaeus* has 123 tRNA-Lys-CTT genes. All have high tRNAscan-SE general bit scores (Supplementary Figure 3) and are unlikely to be pseudogenes. We classified 98 as active. tRNA-derived SINEs can proliferate rapidly via retrotransposition, and a rapid expansion may have occurred in this species, targeting areas within or near regions of high transcription. While few of these tRNA genes have completely identical sequences, 43 have identical 3' flanking regions, extending 10 bases after the tRNA gene (CCATTTGTTG). An additional 24 genes share a different 3' flanking motif (CCAGATGTTG). The immediate flanking regions of actively transcribed tRNA genes are expected to be highly divergent due to transcription-associated mutagenesis (Thornlow et al. 2018). One possible explanation for this discrepancy is that these motifs are part of a fused SINE element responsible for the proliferation of the tRNA-derived SINE element. A similar expansion has occurred in *Microtus ochrogaster* (prairie vole), whose genome contains 131 tRNA-Lys-CTT genes, 66 of which we classified as active. 25 of these genes are exactly identical in tRNA gene sequence, suggesting a similarly recent expansion.

To our knowledge, no actively transcribed suppressor tRNA gene has been demonstrated in a primate genome. However, we have predicted a tRNA-Sup-CTA in *Pan troglodytes* (chimpanzee) as active, although experimentation is necessary to support that this tRNA gene is actively transcribed. While nonsense suppression via readthrough by near-cognate tRNAs has been demonstrated several times in metazoa (Jungreis et al. 2011; Yacoubi et al. 2012; Loughran et al. 2018; Roy et al. 2015) and suppressor tRNAs have been found in other species (Beier and Grimm 2001; Valle et al. 1987), evidence for an actively transcribed suppressor tRNA in a primate species is of great interest for future experimentation. This tRNA gene is a conserved ortholog of the human tRNA-Gln-CTG-6-1 gene, and exhibits several chimpanzee-specific nucleotide substitutions (Supplementary Figure 4) that are not expected to significantly impair the function of its transcript. The similarity of the sequence of this gene to other tRNA-Gln-TTG genes suggests an additional anticodon shift, prior to the divergence of placental mammals. Although our classifier was created in order to better understand tRNA gene expression, it may also be used to screen for outliers that may exhibit unique biology. This tRNA-Sup-CTA gene will be a focus of future experimentation and represents one of many examples of outliers brought to light by our method.

Figure 4: Cysteine is the only standard amino acid without a "core" tRNA. All human active and inactive high-confidence tRNA genes are shown in gray for each amino acid, and those found in the core set of 97 tRNA genes are shown in black.

tRNA genes predicted to be active are more deeply conserved. In addition to application of our classifier to placental mammal genome data, we have developed a complete set of placental mammal tRNA gene orthologs using a Cactus graph (Paten et al. 2011a, 2011b; Nguyen et al. 2015; see Methods). Of the 11,752 tRNA genes in our 29-species alignment, 3,582 genes in total, or about 122 per species, appear to be species-specific. The remaining 8,170 genes were condensed into 1,097 ortholog sets. On average, each of these tRNA genes is present in 7.4 species, indicating that tRNA genes are generally either fairly deeply conserved or recently evolved (Supplementary Figure 5). This is consistent with prior studies in *Drosophila* showing that tRNA genes can be "core" or "peripheral" (Rogers et al. 2010).

To investigate the relationship between activity and conservation, we identified a "core" set of tRNA genes found in all primates (Supplementary Figure 6). We found 97 tRNA genes for which all seven primate species (human, chimpanzee, gorilla, orangutan, macaque, *Microcebus murinus* and *Aotus nancymaae*) have a syntenic ortholog. This most likely represents the set of tRNA genes present in the primate common ancestor that have not been lost in any lineage leading to the sampled genomes. These genes encode 19 amino acids (Figure 4). The only standard amino acid not represented is cysteine. tRNA-Cys genes are often present in

high numbers, and every species in the phylogeny has at least 17 such genes. However, these genes are prone to accumulating substitutions in their sequences. Therefore, the lack of a "core" eutherian tRNA-Cys gene may be due to relatively rapid evolution of this gene family, or difficulty in alignment due to their high variation in sequence.

In 15 of these 97 "core" ortholog sets, we predicted at least one gene as active and at least one gene as inactive. In all cases, we predict more genes as active than inactive, further suggesting that tRNA genes predicted to be active are more deeply conserved. Out of all 1,097 ortholog sets, 750 contain only tRNA genes predicted to be active, approximately mirroring the distribution of active to inactive tRNA genes predicted at the species level and further indicating that active tRNA genes tend to be more strongly conserved.

Substitutions in tRNA anticodons accompany changes in predicted activity state. Among our 1,097 ortholog sets, we identified 111 sets containing genes corresponding to multiple different anticodons. 52 of these 111 anticodon shifts are non-synonymous, resulting in a change in the corresponding amino acid. Ortholog sets containing anticodon shifts are significantly enriched for changes in predicted activity, as 36 of 111 such ortholog sets contain at least one active and at least one inactive tRNA gene (Hypergeometric test, p < 2.3e-4). Likewise, of the 52 ortholog sets containing non-synonymous anticodon substitutions, 18 contain at least one active and at least one inactive tRNA gene (Hypergeometric test, p < 2.8e-3).

Transitions between active and inactive are rare. We fit our ortholog sets and their predicted activity states to a Markov model of evolution of discrete characters using RevBayes (Höhna et al. 2016; Figure 3, see Methods). By fitting our data to the model, we estimated transition probabilities to and from three states - active, inactive, and missing (no detected ortholog). We held the phylogeny constant and solved only for the transition rate parameters. Consistent with the premise that local gene duplication is the dominant mode of tRNA gene evolution, our model suggests that active tRNA genes overwhelmingly tend to remain active, inactive tRNA genes tend to remain inactive (Figure 3).

Although transition events are rare, our classifier does well to detect them. There are 188 human/mouse ortholog pairs for which we have epigenomic data, and in 173 of them, human and mouse have the same activity state. However, we correctly classify 183 human and 181 mouse tRNA genes within this set, indicating that most activity state changes between these species are detectable. If inputs to the model are unobtainable and true activity states are available for only one species, assuming that the activity state of orthologous tRNA genes remains constant across closely related species would yield mostly accurate results. However, our classifier represents an improvement over this assumption.

Generally, tRNA genes in a given activity state are most likely to remain in that state, over branch lengths of 10 million years (Figure 3A). Overall, transitions of any kind are rare, and the active state appears to be most stable, but, surprisingly, inactive tRNA genes most often remain inactive as well. We observed some variation in the relative transition probabilities within clades. Primate tRNA genes are less likely to remain in their initial predicted activity state than rodent tRNA genes. This is consistent with prior studies on the rate of evolutionary change of protein coding gene expression between clades (Brawand et al. 2011; Necsulea and Kaessmann 2014) but may also reflect differences in sample size between clades. Based on our results, turnover in tRNA gene expression class generally appears to be slow, similar to protein coding genes (Brawand et al. 2011).

DISCUSSION:

Greater understanding of tRNA regulation is a difficult and unmet challenge. There are many obstacles preventing direct measurement of expression at the gene level. Nonetheless, we show that, in order to determine relative levels of expression of tRNA genes, direct measurement can be effectively replaced by inference through data derived from genome sequences and alignments. We show that binary expression levels can be reliably predicted through features both intrinsic to tRNA genes, which relate directly to tRNA function and processing, and extrinsic, which relate to the genomic environment.

Our classifier demonstrates remarkable accuracy in predicting tRNA gene activity. However, there are several caveats to our method. Importantly, the epigenomic ChIP data used to create the "active" and "inactive"

labels for both human and mouse data is imperfect. It is possible that many inactive tRNAs are in fact very carefully regulated, such that their activity is not captured by previously conducted assays. ChIP captures the chromatin environment across many tissues and is by far the most comprehensive data available for tRNA gene activity, yet is still an indirect approximation of gene transcription. However, that comprehensive tRNA gene expression data is unavailable motivates the creation of our classifier. Available data indicate that our estimates are comparable to experimental results, but with much greater ease of use and cost-effectiveness.

We have focused on tRNA genes specifically, but the basic principles underlying our model might be applied to any duplicate gene family. For example, ribosomal RNA (rRNA) genes are highly transcribed and high in copy number, and many rRNA genes have no detected expression in plants and yeast. rRNA gene copy number and relative transcription appear to have fundamental implications for organismal fitness (Ide et al. 2010). We have also previously shown that histone protein coding genes exhibit similar genetic variation to tRNA genes (Thornlow et al. 2018). If one was interested in inferring relative levels of expression of rRNA or histone protein coding genes, a model similar to ours could be developed. This core idea therefore has the potential to revolutionize functional and evolutionary genomic inference for large multi-copy gene families.

Our methods may also be applied to other species. We have previously demonstrated that transcription-associated mutagenesis creates signatures at plant and insect tRNA gene loci that are qualitatively similar to those in human and mouse (Thornlow et al. 2018). While some features may not extend quite so far phylogenetically, and others may require recalibration or processing, a similar model could be developed for other eukaryotic clades. For the present study, we chose to work with placental mammals, as the corresponding data is of relatively high quality.

Classification using our approach may expand in several directions as more data is collected. For example, variation within populations, and potentially within highly proliferative tissues, may be useful for predicting relative transcript expression among gene families. We previously determined that actively transcribed tRNA genes accumulate more rare single nucleotide polymorphisms (SNPs) in both their flanking regions and gene sequences (Thornlow et al. 2018). Therefore, we expect that when population variation data is available for more species, we might adapt our approach to infer expression differences at narrower timescales. Our model may also be expanded to accommodate non-binary classification of expression levels and capture the nuance of tRNA gene expression regulation.

Our classifier also handles strongly intercorrelated features well. When modeling a single biological process, all parts of the process are inevitably correlated. This is visible in the current model, as we use two features related to CpG dinucleotides and two features derived from PhyloP data (Table 1). We also include tRNAscan-SE general bit scores and canonical secondary structure, although the general bit scores account for both primary and secondary structure (see Methods). Using highly correlated features leads to underestimation of their importance to the model (Strobl et al. 2008; Tolosi and Lengauer 2011). Nevertheless, we have included intercorrelated features to achieve optimal model performance.

While tRNA genes predicted to be active are most well conserved, it is surprising that inactive tRNA genes are also strongly conserved. This is true not only in their sequence but in their expression, as tRNA genes predicted to be inactive are more likely to remain inactive than to become pseudogenes or active genes. Our epigenomic data captures over 100 tissues, but is temporally limited, and some of these genes may be transcribed only at specific stages in development not captured by these assays. Other tRNA genes may be biologically important for functions other than production of mature transcripts. A subset of apparently silent human tRNA genes are present in the introns of protein coding genes, and may regulate their transcription. Prior studies have also demonstrated that tRNA genes act as insulators bounding chromatin regulatory programs (Raab et al. 2012).

Here we demonstrate reliable classification of tRNA genes using open-source software that requires little input data and can easily be expanded in the future. Annotations created by our method will be useful in aiding design of tRNA characterization experiments, as well as for variant effect interpretation, as polymorphic sites within or near more highly active tRNA genes may have more significant fitness consequences. Our method demonstrates the importance of the local genomic context in gene-level expression, and its success

suggests that the same strategy may be used to reliably predict expression of other multi-paralog gene families. This work informs the broader question of tRNA gene function evolution, illustrating that tRNA gene expression regulation is likely dependent on many factors, including those intrinsic to the tRNA gene sequence and those derived from the varied genomic environment.

METHODS

Developing the classifier. For the training data, we used coordinates from the high-confidence tRNA set in the hg38 genome from GtRNAdb (Chan and Lowe 2016). For all test species, we extracted the genomes from our Cactus alignment, ran tRNAscan-SE 2.0 using the default parameters and applied the EukHighConfidenceFilter to exclude tRNA pseudogenes and tRNA-derived SINEs (Chan and Lowe 2019). We used custom Python scripts to find tRNA loci that were identical from 80 nucleotides upstream to 40 nucleotides downstream of the gene start and end. We considered these segmental duplications and excluded them from classification. If any of these non-unique loci did not align to any tRNA loci in any other species, they were also removed from our ortholog calls, as these loci are most likely assembly errors. For genomes in which at least 85% of nucleotides were found on chromosomes, we also excluded all tRNA genes not found on chromosomes. For the human tRNA gene set, because our epigenomic data is derived from hg19 annotations, we also removed any tRNAs in segmental duplications in the hg19 genome. We used the PHAST (Hubisz et al. 2011) and HAL (Hickey et al. 2013) toolkits, and custom Python scripts to collect feature data and scikit-learn to test classification methods (Pedregosa et al. 2011).

Eliminating uninformative features. We used the Spearman rank correlation test to ensure that no features were perfectly correlated (Guyon and Elisseeff 2003). We used CfsSubsetEval (Hall et al. 2009) to remove uninformative features and scikit-learn to determine feature importance (Pedregosa et al. 2011).

When PhyloP data were unobtainable due to lack of alignment, we replaced feature values for each tRNA gene with the mean value for that feature across all tRNA genes in that species. The number of unalignable tRNA genes is correlated directly with the phylogenetic distance to the species' nearest neighbor (p < 8.8e-6, Spearman rank, Supplementary Figure 8), and indirectly with contig N50 (p < 3.6e-5).

We used output from RNAfold from ViennaRNA (Lorenz et al. 2011) to estimate constrained minimum free energy (Table 1). RNAfold output is significantly correlated with tRNAscan-SE secondary structure scores (Spearman rank correlation, p < 2e-4), indicating interchangeability. To determine which to include in the model, we created a separate model that included both scores, and found that RNAfold output had greater feature importance. We also tested a model that includes tRNAscan-SE secondary structure, HMM and isotype-specific bit scores. Upon their inclusion, the importance of the tRNAscan-SE general bit score decreases slightly, but it remains more important to the model than any of the other tRNAscan-SE scores, supporting its usage in the model.

Our final model uses 250 estimators, a minimum of 2 samples required to split an internal node, a maximum depth of 4 nodes, and is trained through 10-fold cross-validation. For all species, we used the pipeline on GitHub (github.com/bpt26/tRNA_classifier/; Supplementary Figure 9). For human and mouse, we used epigenomic data from Roadmap Epigenomics Consortium et al. (2015) and Bogu et al. (2015). Our alignment uses the hg38 genome, and we excluded annotated human tRNA genes found in hg38 that did not liftover (Casper et al. 2018) to hg19.

Creating an ortholog set. We used hal2maf to generate alignments (Hickey et al. 2013). For each tRNA locus, we considered the best aligning tRNA locus from all other species as orthologous, allowing only one ortholog per species per locus. We augmented our ortholog sets with syntenic human/mouse, human/dog and human/macaque tRNA gene ortholog pairs from Holmes et al (in prep). For all instances in which each tRNA gene in a Holmes et al ortholog pair aligned to mutually exclusive sets of species in the Cactus graph, we combined them into one ortholog set. We found that 29 Holmes et al human-mouse ortholog pairs align to each other in the Cactus graph, 152 align to mutually exclusive sets of species, and 17 align to overlapping sets of species. Therefore, we combined the 152 human-mouse tRNA gene pairs into larger ortholog sets.

Fitting a Markov model. We used a phylogeny from TimeTree (Kumar et al. 2017) and fit our data to a Markov model using RevBayes (Höhna et al. 2016). We held the phylogeny constant and allowed RevBayes to optimize only the Q matrix using our tRNA data. We then determined transition probabilities over 10 million years using the RevBayes function getTransitionProbabilities() across all species and by clade (Supplementary Figure 7).

DATA ACCESS

Our pipeline is available at https://github.com/bpt26/tRNA_classifier/. Classification distribution data are available in Supplemental Materials. Alignments generated by the Cactus graph are available upon request. The guide tree for the Cactus graph was:

(((((((((Homo_sapiens:0.00655,Pan_troglodytes:0.00684)Anc32:0.00422,Gorilla_gorilla_gorilla:0.008964)Anc29:0.009693,Pongo_abelii:0.01894)Anc25:0.015511,Macaca_mulatta:0.043601)Anc20:0.08444,Aotus_nancymaae:0.08)Anc15:0.08,Microcebus_murinus:0.10612)Anc11:0.083494,((((Jaculus_jaculus:0.1,(Microtus_ochrogaster:0.14,(Mus_musculus:0.084509,Rattus_norvegicus:0.091589)Anc30:0.047773)Anc26:0.06015)Anc21:0.122992,(Heterocephalus_glaber:0.1,(Cavia_porcellus:0.065629,(Chinchilla_lanigera:0.06,Octodon_degus:0.1)Anc31:0.06)Anc27:0.05)Anc22:0.06015)Anc16:0.05,Marmota_marmota:0.1)Anc12:0.05,Oryctolagus_cuniculus:0.21569)Anc08:0.04)Anc04:0.040593,(((Sus_scrofa:0.12,(Orcinus_orca:0.069688,(Bos_taurus:0.04,Capra_hircus:0.04)Anc23:0.09)Anc17:0.045488)Anc13:0.02,((Equus_caballus:0.109397,(Felis_catus:0.098612,(Canis_lupus_familiaris:0.052458,Mustela_putorius_furo:0.08)Anc28:0.02)Anc24:0.049845)Anc18:0.02,(Pteropus_alecto:0.1,Eptesicus_fuscus:0.08)Anc19:0.033706)Anc14:0.03)Anc09:0.025,Erinaceus_europaeus:0.278178)Anc05:0.021227)Anc02:0.023664,((Loxodonta_africana:0.098842,Chrysochloris_asiatica:0.04)Anc06:0.05,Dasypus_novemcinctus:0.169809)Anc03:0.02)backbone_root;

ACKNOWLEDGMENTS

We thank the R.B.C.-D. and T.M.L. laboratories for suggestions and feedback. This work was supported by National Human Genome Research Institute/NIH Grant 2R01HG006753-04A1 (to T.M.L.) and National Institute of General Medical Sciences/NIH Grant R35GM128932-01 (to R.B.C.-D.). B.P.T. was funded by NIH Grants T32HG008345 and 5F31HG010584-02.

Author contributions: Study design was by B.T., R.B.C.-D. and T.M.L.. J.A. provided the Cactus graph. A.H. provided mouse tRNA gene activity labels and ortholog data. B.T. wrote pipeline for extracting feature data and classifying genes. B.T., R.B.C.-D. and T.M.L. wrote the manuscript.

DISCLOSURE DECLARATION

We declare no competing interests.

REFERENCES

Allison DS, Hall BD. 1985. Effects of alterations in the 3' flanking sequence on in vivo and in vitro expression of the yeast SUP4-o tRNATyr gene. *EMBO J* **4**: 2657–2664.

Arimbasseri AG, Rijal K, Maraia RJ. 2013. Transcription termination by the eukaryotic RNA polymerase III. *Biochim Biophys Acta* **1829**: 318–330.

Beier H, Grimm M. 2001. Misreading of termination codons in eukaryotes by natural nonsense suppressor tRNAs. *Nucleic Acids Res* **29**: 4767–4782.

Bogu GK, Vizán P, Stanton LW, Beato M, Di Croce L, Marti-Renom MA. 2015. Chromatin and RNA Maps Reveal Regulatory Long Noncoding RNAs in Mouse. *Mol Cell Biol* **36**: 809–819.

Boivin V, Deschamps-Francoeur G, Couture S, Nottingham RM, Bouchard-Bourelle P, Lambowitz AM, Scott MS, Abou-Elela S. 2018. Simultaneous sequencing of coding and noncoding RNA reveals a human

transcriptome dominated by a small number of highly expressed noncoding genes. *RNA* **24**: 950–965.

Brawand D, Soumillon M, Necsulea A, Julien P, Csárdi G, Harrigan P, Weier M, Liechti A, Aximu-Petri A, Kircher M, et al. 2011. The evolution of gene expression levels in mammalian organs. *Nature* **478**: 343–348.

Casper J, Zweig AS, Villarreal C, Tyner C, Speir ML, Rosenbloom KR, Raney BJ, Lee CM, Lee BT, Karolchik D, et al. 2018. The UCSC Genome Browser database: 2018 update. *Nucleic Acids Res* **46**: D762–D769.

Chan PP, Lowe TM. 2016. GtRNAdb 2.0: an expanded database of transfer RNA genes identified in complete and draft genomes. *Nucleic Acids Res* **44**: D184–9.

Chan PP, Lowe TM. 2019. tRNAscan-SE: Searching for tRNA Genes in Genomic Sequences. *Methods Mol Biol* **1962**: 1–14.

Cozen AE, Quartley E, Holmes AD, Hrabeta-Robinson E, Phizicky EM, Lowe TM. 2015. ARM-seq: AlkB-facilitated RNA methylation sequencing reveals a complex landscape of modified tRNA fragments. *Nat Methods* **12**: 879–884.

Foissac S, Djebali S, Munyard K, Villa-Vialaneix N, Rau A, Muret K, Esquerre D, Zytnicki M, Derrien T, Bardou P, et al. 2018. Livestock genome annotation: transcriptome and chromatin structure profiling in cattle, goat, chicken and pig. *bioRxiv* 316091. https://www.biorxiv.org/content/early/2018/05/11/316091.abstract (Accessed November 20, 2018).

Gardiner-Garden M, Frommer M. 1987. CpG islands in vertebrate genomes. *J Mol Biol* **196**: 261–282.

Goodarzi H, Liu X, Nguyen HCB, Zhang S, Fish L, Tavazoie SF. 2015. Endogenous tRNA-Derived Fragments Suppress Breast Cancer Progression via YBX1 Displacement. *Cell* **161**: 790–802.

Guyon I, Elisseeff A. 2003. An Introduction to Variable and Feature Selection. *J Mach Learn Res* **3**: 1157–1182.

Hall MA. 1998. Correlation-based feature subset selection for machine learning. *Thesis submitted in partial fulfillment of the requirements of the degree of Doctor of Philosophy at the University of Waikato*. https://ci.nii.ac.jp/naid/10018668219/.

Hall M, Frank E, Holmes G, Pfahringer B, Reutemann P, Witten IH. 2009. The WEKA data mining software: an update. *ACM SIGKDD Explorations Newsletter* **11**: 10–18.

Hamada M, Sakulich AL, Koduru SB, Maraia RJ. 2000. Transcription termination by RNA polymerase III in fission yeast. A genetic and biochemically tractable model system. *J Biol Chem* **275**: 29076–29081.

Hedges SB, Dudley J, Kumar S. 2006. TimeTree: a public knowledge-base of divergence times among organisms. *Bioinformatics* **22**: 2971–2972.

Hickey G, Paten B, Earl D, Zerbino D, Haussler D. 2013. HAL: a hierarchical format for storing and analyzing multiple genome alignments. *Bioinformatics* **29**: 1341–1342.

Höhna S, Landis MJ, Heath TA, Boussau B, Lartillot N, Moore BR, Huelsenbeck JP, Ronquist F. 2016. RevBayes: Bayesian Phylogenetic Inference Using Graphical Models and an Interactive Model-Specification Language. *Syst Biol* **65**: 726–736.

Hubisz MJ, Pollard KS, Siepel A. 2011. PHAST and RPHAST: phylogenetic analysis with space/time models. *Brief Bioinform* **12**: 41–51.

Ide S, Miyazaki T, Maki H, Kobayashi T. 2010. Abundance of ribosomal RNA gene copies maintains genome integrity. *Science* **327**: 693–696.

Johansson L, Gafvelin G, Arnér ESJ. 2005. Selenocysteine in proteins—properties and biotechnological use.

Biochimica et Biophysica Acta (BBA) - General Subjects **1726**: 1–13.

Jungreis I, Lin MF, Spokony R, Chan CS, Negre N, Victorsen A, White KP, Kellis M. 2011. Evidence of abundant stop codon readthrough in Drosophila and other metazoa. *Genome Res* **21**: 2096–2113.

Kirchner S, Ignatova Z. 2015. Emerging roles of tRNA in adaptive translation, signalling dynamics and disease. *Nat Rev Genet* **16**: 98–112.

Koski RA, Clarkson SG, Kurjan J, Hall BD, Smith M. 1980. Mutations of the yeast SUP4 tRNATyr locus: transcription of the mutant genes in vitro. *Cell* **22**: 415–425.

Krinner S, Heitzer AP, Diermeier SD, Obermeier I, Längst G, Wagner R. 2014. CpG domains downstream of TSSs promote high levels of gene expression. *Nucleic Acids Res* **42**: 3551–3564.

Kumar S, Stecher G, Suleski M, Hedges SB. 2017. TimeTree: A Resource for Timelines, Timetrees, and Divergence Times. *Mol Biol Evol* **34**: 1812–1819.

Kutter C, Brown GD, Gonçalves A, Wilson MD, Watt S, Brazma A, White RJ, Odom DT. 2011. Pol III binding in six mammals shows conservation among amino acid isotypes despite divergence among tRNA genes. *Nat Genet* **43**: 948–955.

Li WH, Ellsworth DL, Krushkal J, Chang BH, Hewett-Emmett D. 1996. Rates of nucleotide substitution in primates and rodents and the generation-time effect hypothesis. *Mol Phylogenet Evol* **5**: 182–187.

Lorenz R, Bernhart SH, Höner Zu Siederdissen C, Tafer H, Flamm C, Stadler PF, Hofacker IL. 2011. ViennaRNA Package 2.0. *Algorithms Mol Biol* **6**: 26.

Loughran G, Jungreis I, Tzani I, Power M, Dmitriev RI, Ivanov IP, Kellis M, Atkins JF. 2018. Stop codon readthrough generates a C-terminally extended variant of the human vitamin D receptor with reduced calcitriol response. *J Biol Chem* **293**: 4434–4444.

Maraia RJ, Kenan DJ, Keene JD. 1994. Eukaryotic transcription termination factor La mediates transcript release and facilitates reinitiation by RNA polymerase III. *Mol Cell Biol* **14**: 2147–2158.

Meunier J, Lemoine F, Soumillon M, Liechti A, Weier M, Guschanski K, Hu H, Khaitovich P, Kaessmann H. 2013. Birth and expression evolution of mammalian microRNA genes. *Genome Res* **23**: 34–45.

Mleczko AM, Celichowski P, Bąkowska-Żywicka K. 2014. Ex-translational function of tRNAs and their fragments in cancer. *Acta Biochim Pol* **61**: 211–216.

Necsulea A, Kaessmann H. 2014. Evolutionary dynamics of coding and non-coding transcriptomes. *Nat Rev Genet* **15**: 734–748.

Necsulea A, Soumillon M, Warnefors M, Liechti A, Daish T, Zeller U, Baker JC, Grützner F, Kaessmann H. 2014. The evolution of lncRNA repertoires and expression patterns in tetrapods. *Nature* **505**: 635–640.

Nguyen N, Hickey G, Zerbino DR, Raney B, Earl D, Armstrong J, Kent WJ, Haussler D, Paten B. 2015. Building a pan-genome reference for a population. *J Comput Biol* **22**: 387–401.

Orioli A, Pascali C, Quartararo J, Diebel KW, Praz V, Romascano D, Percudani R, van Dyk LF, Hernandez N, Teichmann M, et al. 2011. Widespread occurrence of non-canonical transcription termination by human RNA polymerase III. *Nucleic Acids Res* **39**: 5499–5512.

Palazzo AF, Lee ES. 2015. Non-coding RNA: what is functional and what is junk? *Front Genet* **6**: 2.

Pan T. 2018. Modifications and functional genomics of human transfer RNA. *Cell Res* **28**: 395–404.

Paten B, Diekhans M, Earl D, John JS, Ma J, Suh B, Haussler D. 2011a. Cactus graphs for genome comparisons. *J Comput Biol* **18**: 469–481.

Paten B, Earl D, Nguyen N, Diekhans M, Zerbino D, Haussler D. 2011b. Cactus: Algorithms for genome multiple sequence alignment. *Genome Res* **21**: 1512–1528.

Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, Blondel M, Prettenhofer P, Weiss R, Dubourg V, et al. 2011. Scikit-learn: Machine Learning in Python. *J Mach Learn Res* **12**: 2825–2830.

Raab JR, Chiu J, Zhu J, Katzman S, Kurukuti S, Wade PA, Haussler D, Kamakaka RT. 2012. Human tRNA genes function as chromatin insulators. *EMBO J* **31**: 330–350.

Roadmap Epigenomics Consortium, Kundaje A, Meuleman W, Ernst J, Bilenky M, Yen A, Heravi-Moussavi A, Kheradpour P, Zhang Z, Wang J, et al. 2015. Integrative analysis of 111 reference human epigenomes. *Nature* **518**: 317–330.

Rogers HH, Bergman CM, Griffiths-Jones S. 2010. The evolution of tRNA genes in Drosophila. *Genome Biol Evol* **2**: 467–477.

Roy B, Leszyk JD, Mangus DA, Jacobson A. 2015. Nonsense suppression by near-cognate tRNAs employs alternative base pairing at codon positions 1 and 3. *Proc Natl Acad Sci U S A* **112**: 3038–3043.

Schmitt BM, Rudolph KLM, Karagianni P, Fonseca NA, White RJ, Talianidis I, Odom DT, Marioni JC, Kutter C. 2014. High-resolution mapping of transcriptional dynamics across tissue development reveals a stable mRNA–tRNA interface. *Genome Res* **24**: 1797–1807.

Strobl C, Boulesteix A-L, Kneib T, Augustin T, Zeileis A. 2008. Conditional variable importance for random forests. *BMC Bioinformatics* **9**: 307.

Sun C, Fu Z, Wang S, Li J, Li Y, Zhang Y, Yang F, Chu J, Wu H, Huang X, et al. 2018. Roles of tRNA-derived fragments in human cancers. *Cancer Lett* **414**: 16–25.

Tang DTP, Glazov EA, McWilliam SM, Barris WC, Dalrymple BP. 2009. Analysis of the complement and molecular evolution of tRNA genes in cow. *BMC Genomics* **10**: 188.

Thornlow B, Hough J, Roger J, Gong H, Lowe T, Corbett-Detig R. 2018. Transfer RNA genes experience exceptionally elevated mutation rates. *Proceedings of the National Academy of Sciences* **115**: 8996–9001.

Tolosi L, Lengauer T. 2011. Classification with correlated features: unreliability of feature ranking and solutions. *Bioinformatics* **27**: 1986–1994.

Valle RP, Morch MD, Haenni AL. 1987. Novel amber suppressor tRNAs of mammalian origin. *EMBO J* **6**: 3049–3055.

Yacoubi BE, El Yacoubi B, Bailly M, de Crécy-Lagard V. 2012. Biosynthesis and Function of Posttranscriptional Modifications of Transfer RNAs. *Annual Review of Genetics* **46**: 69–95. http://dx.doi.org/10.1146/annurev-genet-110711-155641.

Zheng G, Qin Y, Clark WC, Dai Q, Yi C, He C, Lambowitz AM, Pan T. 2015. Efficient and quantitative high-throughput tRNA sequencing. *Nat Methods* **12**: 835–837.

	Feature
Extrinsic	**Average PhyloP Score in tRNA Gene Sequence**
	Average PhyloP Score in 3' Flanking Region
	Average PhyloP Score in 5' Flanking Region
	CpG Density Across tRNA Locus
	Distance To Nearest Exon
	Exons Within 2 Megabases
	Exons Within 75 Kilobases
	Minimum PhyloP Score in tRNA Gene Sequence
	Minimum PhyloP Score in 5' Flanking Region
	Minimum PhyloP Score in 3' Flanking Region
	Observed/Expected CpG Islands Score Across tRNA Locus
	Observed/Expected CpG Islands Score Downstream of tRNA Gene
	Observed/Expected CpG Islands Score Upstream of tRNA Gene
	Percentage of CpG Dinucleotides Upstream of tRNA Gene
	Substitutions in tRNA Gene Sequence Relative to Outgroup
	Substitutions in 3' Flanking Sequence Relative to Outgroup
	Substitutions in 5' Flanking Sequence Relative to Outgroup
	tRNA Genes Within 10 Kilobases
	tRNA Genes Within 400 Kilobases
Intrinsic	**Canonical tRNA Secondary Structure**
	Distance To Nearest Transcription Termination Sequence of Any Kind
	Distance To Nearest TTTT Transcription Termination Sequence
	Distance To Nearest TTTTT Transcription Termination Sequence
	tRNA Genes Corresponding to the Same Amino Acid
	tRNA Genes Corresponding to the Same Codon
	tRNAscan-SE General Bit Score
	tRNAscan-SE HMM Bit Score
	tRNAscan-SE Isotype-Specific Bit Score
	tRNAscan-SE Secondary Structure Bit Score
	tRNA Sequence Unique In Genome

Supplementary Table 1: Our initial model used many features both intrinsic to tRNA structure and function as well as extrinsic and derived solely from the genomic context. All features considered for the model are shown, with those in the final model bolded. Features were removed either for lack of contribution to the model, overlapping contribution with other features, or difficulty in obtaining data due to lack of alignment. "Transcription Termination Sequence of Any Kind" refers to "TTTT", "TTTTT", as well as alternative sequences demonstrated in (Orioli et al. 2011).

Supplementary Figure 1: Comparison of tRNA gene classifications to ChIP-Seq data is consistent across tissues. Mouse ChIP-Seq data compared to tRNA gene classifications and ChIP data for both testes replicates and both muscle replicates. Probability scores output by the classifier are shown on the horizontal axis where tRNA genes furthest left are predicted inactive with greater probability and tRNA genes furthest right are predicted active with greater probability. The vertical axis indicates read counts in testes (A, B) and muscle (C, D) from Kutter et al. (2011). Purple indicates measured activity in at least one tissue from Bogu et al. (2015) while orange indicates no measured activity.

Supplementary Figure 2: Comparison of classifications to ATAC-Seq data suggests similar accuracy across clades. The mean ATAC-Seq peaks across liver, CD4 and CD8 cells in cow (A), goat (B) and pig (C) within 250 base pairs of each tRNA gene are shown against the classification probabilities for each gene as in Supplementary Figure 1.

Supplementary Figure 3: Most *Erinaceus europaeus* tRNA-Lys-CTT genes have high tRNAScan-SE general bit scores. Histogram showing the number of tRNA-Lys-CTT genes in the *E. europaeus* genome by tRNAScan-SE general bit score. For reference, the average active human tRNA gene has a bit score of 76.2 and the average inactive human tRNA gene has a bit score of 69.3.

Supplementary Figure 4: Chimpanzee ortholog of human tRNA-Gln-CTG-6-1 may be an actively transcribed suppressor tRNA gene. Gene browser screenshot showing the PhyloP and Multiz Alignment tracks of the human tRNA-Gln-CTG-6-1 gene, for which the chimpanzee genome has a tRNA-Sup-CTA gene. The sequence of the tRNA gene is highlighted in cyan. The chimpanzee sequence (second line from top) exhibits several unique substitutions. Most notably, the CTG anticodon (here from right to left in the center of the highlighted region) has changed to CTA in chimp, accompanied by other changes to the body of the tRNA. The PhyloP track also shows strong divergence upstream and downstream of the tRNA gene sequence, consistent with high levels of expression.

Supplementary Figure 5: tRNA genes are generally either fairly deeply conserved or recently evolved.
Size distribution of all ortholog sets, with genes present in at least 2 species. 3,582 of the 11,752 tRNA genes in our alignment are species-specific, excluded from this figure. The remaining 8,170 tRNA genes are condensed into 1,097 ortholog sets.

Supplementary Figure 6: Core tRNA genes are very deeply conserved. Histogram of sizes of the 97 tRNA gene ortholog sets present in at least all 7 primate species.

A

		Final		
		Missing	Inactive	Active
Initial	Missing	.871	.111	.017
	Inactive	.092	.859	.049
	Active	.012	.014	.974

B

		Final		
		Missing	Inactive	Active
Initial	Missing	.403	.341	.256
	Inactive	.116	.845	.040
	Active	.144	.419	.437

C

		Final		
		Missing	Inactive	Active
Initial	Missing	.723	.229	.049
	Inactive	.163	.656	.182
	Active	.106	.365	.528

D

		Final		
		Missing	Inactive	Active
Initial	Missing	.957	.022	.021
	Inactive	.165	.821	.015
	Active	.150	.188	.663

E

Species	Active	Inactive
Homo sapiens	314	57
Pan troglodytes	339	56
Gorilla gorilla gorilla	325	46
Pongo abelii	351	50
Macaca mulatta	337	54
Aotus nancymaae	285	43
Microcebus murinus	284	40
Octodon degus	240	58
Chinchilla lanigera	250	51
Cavia porcellus	266	54
Heterocephalus glaber	238	56
Mus musculus	275	112
Rattus norvegicus	245	102
Microtus ochrogaster	343	127
Jaculus jaculus	246	72
Marmota marmota	253	156
Oryctolagus cuniculus	377	70
Bos taurus	406	369
Capra hircus	216	317
Orcinus orca	328	333
Sus scrofa	294	104
Canis lupus familiaris	278	90
Mustela putorius furo	291	87
Equus caballus	351	65
Pteropus alecto	222	123
Eptesicus fuscus	199	15
Erinaceus europaeus	343	104
Dasypus novemcinctus	379	149
Chrysochloris asiatica	275	97

20 million years

Supplementary Figure 7: Clades vary in their activity state transition rates. Transition matrices shown as in Figure 3 for all descendants of labeled nodes (A-D). Phylogeny, species names and predicted active and inactive tRNA gene counts reproduced from Figure 3 (E).

Supplementary Figure 8: Unalignable tRNA genes are likely due to a combination of decreased genome quality and increased phylogenetic distance. The proportion of tRNA genes in a given genome is shown compared to the phylogenetic distance to the nearest neighbor in the phylogeny (A), as well as the log of the Contig N50 for that species (B). Branch length in the phylogeny is measured by expected number of substitutions between species in neutral regions.

Supplementary Figure 9: Classification pipeline uses tRNAscan-SE, HAL toolkit and PHAST. Workflow for classifying tRNAs with instructions for usage found at https://github.com/bpt26/tRNA_classifier. Files outlined in red are the inputs used by classifytRNAs.py, the program responsible for producing the final predictions.