Analysis of meiosis in *Pristionchus pacificus* reveals plasticity in homolog pairing and synapsis in the nematode lineage

Regina Rillo-Bohn[1,2*], Renzo S. Adilardi[1,2*], Therese Mitros[1], Bariş Avşaroğlu[1,2], Lewis Stevens[3], Simone Köhler[1,2 4], Joshua Bayes[1], Clara Wang[1,2], Sabrina Lin[1,2], K. Alienor Baskevitch[1,2], Daniel S. Rokhsar[1,5,6,7], and Abby F. Dernburg[1,2,8,9]

[1]Department of Molecular and Cell Biology, University of California, Berkeley, Berkeley, CA USA; [2]Howard Hughes Medical Institute, Chevy Chase, MD United States; [3]Darwin Tree of Life Project, Wellcome Sanger Institute, Cambridge, UK; [4]Present address: Cell Biology and Biophysics Unit, European Molecular Biology Laboratory, Heidelberg, Germany; [5]Department of Energy Joint Genome Institute, Berkeley, California 94720, USA. [6]Okinawa Institute of Science and Technology Graduate University, Onna, Okinawa 9040495; Japan, [7]Chan Zuckerberg Biohub, San Francisco, CA, USA [8]Biological Systems and Engineering Division, Lawrence Berkeley National Laboratory, Berkeley, CA USA, [9]California Institute for Quantitative Biosciences, Berkeley, CA USA

*Equal contribution

ABSTRACT

Meiosis is conserved across eukaryotes yet varies in the details of its execution. Here

we describe a new comparative model system for molecular analysis of meiosis, the

nematode *Pristionchus pacificus*, a distant relative of the widely studied model organism

Caenorhabditis elegans. *P. pacificus* shares many anatomical and other features that

facilitate analysis of meiosis in *C. elegans*. However, while *C. elegans* has lost the

meiosis-specific recombinase Dmc1 and evolved a recombination-independent

mechanism to synapse its chromosomes, *P. pacificus* expresses both DMC-1 and RAD-

51. We find that SPO-11 and DMC-1 are required for stable homolog pairing, synapsis,

and crossover formation, while RAD-51 is dispensable for these key meiotic processes.

RAD-51 and DMC-1 localize sequentially to chromosomes during meiotic prophase and

show nonoverlapping functions. We also present a new genetic map for *P. pacificus* that

reveals a crossover landscape very similar to that of *C. elegans*, despite marked

divergence in the regulation of synapsis and crossing-over between these lineages.

INTRODUCTION

All sexually reproducing organisms rely on the specialized cell division process of

meiosis to generate haploid gametes from diploid precursors. Upon fertilization,

gametes fuse and restore the diploid chromosome complement in the zygote. This

sexual cycle of meiosis and fertilization was likely present in the last eukaryotic common

ancestor (Goodenough and Heitman, 2014). Studies in plants, animals, fungi, and

protists have revealed intriguing diversity in the details of meiotic mechanisms.

A defining feature of meiosis is a "reductional" segregation in which homologous

chromosomes are separated, usually during the first of two nuclear divisions. A highly

choreographed series of chromosome transactions precedes this division and ensures

faithful homolog segregation: (1) pairing, in which chromosomes contact and recognize

their homologous partners; (2) synapsis, defined as the assembly of a protein ensemble

called the synaptonemal complex (SC) between homologs, which stabilizes pairing and

leads to their lengthwise alignment; and (3) crossover (CO) recombination, which

creates physical links between chromosomes that promote proper bi-orientation during

anaphase I. Failure to form at least one "obligate" CO between each pair of homologs

typically results in missegregation and aneuploid gametes (Zickler and Kleckner, 2015).

Although homolog pairing, synapsis and CO recombination during meiosis are

nearly ubiquitous among eukaryotes, key aspects of these chromosomal transactions

show remarkable diversity across different lineages. In most model fungi, plants, and

animals, synapsis is thought to be triggered by early steps in recombination. Meiotic

recombination is initiated by double-strand breaks (DSBs) catalyzed by the conserved

topoisomerase-like enzyme Spo11 (Bergerat et al., 1997; Keeney et al., 1997). DSBs

are resected to form 3' single-stranded DNA overhangs. Dmc1, a meiosis-specific

recombinase, forms filaments along the resulting ssDNA segments and promotes

interhomolog strand invasion to form joint molecules (JMs). Rad51 is often required as a

cofactor for Dmc1. A subset of these JMs is ultimately repaired through a meiosis-

specific pathway that generates COs, while the rest are mostly repaired through a more

"mitotic-like" homologous recombination pathway to yield non-CO products (Hunter,

2015; Kohl and Sekelsky, 2013).

Spo11-dependent induction of DSBs and Dmc1-dependent strand invasion are

crucial for synapsis, and thus for stable homolog pairing, in the budding yeast

Saccharomyces cerevisiae, the flowering plant *Arabidopsis thaliana,* and in mice

(Bishop et al., 1992; Couteau et al., 1999; Grelon, 2001; Pittman et al., 1998; Rockmill

et al., 1995; Yoshida et al., 1998). Thus, the formation of JMs is considered to be an

intrinsic part of the pairing process by which chromosomes recognize their partners. In

contrast, recombination-independent mechanisms of pairing and synapsis have been

characterized in other prominent model systems, including the dipteran insect

Drosophila melanogaster and the nematode *Caenorhabditis elegans* (Lake and Hawley,

2012; Rog and Dernburg, 2013)*.* While crossing-over is essential for successful

execution of meiosis in *C. elegans* and in female fruit flies, homolog pairing and

synapsis can be uncoupled from DSB formation and JM formation. In *Drosophila*

melanogaster females, pairing initiates in proliferating germline cells even before they

enter meiosis and is stabilized by SC formation during early prophase (Christophorou et

al., 2015, 2013). *D. melanogaster* males lack both recombination and SCs and have

apparently evolved a distinct mechanism to stabilize homolog pairing and enable

reductional segregation (McKee et al., 2012). In *C. elegans*, pairing and synapsis are

both mediated by Pairing Centers (PCs), specialized sites on each chromosome bound

by a family of zinc-finger proteins that interact with nuclear envelope proteins and

promote large-scale chromosome movement (MacQueen et al., 2005; Phillips et al.,

2005; Phillips and Dernburg, 2006; Sato et al., 2009). While interactions between

chromosomes and nuclear envelope proteins play important roles in meiotic pairing and

synapsis across eukaryotes, in *C. elegans* they have acquired a critical role in coupling

homolog pairing to synapsis initiation (Penkner et al., 2007, 2009; Sato et al., 2009).

To investigate how the meiotic program is modified during evolution, we have

established tools to investigate meiosis in the free-living nematode *Pristionchus*

pacificus, which has been established as a model for studies in development, evolution

and ecology (Sommer, 2015). Like its distant relative *C. elegans*, *P. pacificus* is an

androdioecious species, with populations consisting of self-fertilizing hermaphrodites

(XX) and a low frequency of spontaneous males (XO) (Sommer, 2015; Sommer et al.,

1996). *P. pacificus* also shares with *C. elegans* a short life cycle of 3.5 days at 20°C,

produces large broods of about 200 progeny by self-fertilization, and is easily cultured in

the lab (Hong and Sommer, 2006). Although *C. elegans* and *P. pacificus* diverged an

estimated 60-90 million years ago (Werner et al., 2018), they share the same number of

chromosomes (2n=12). Other than one major chromosomal translocation,

macrosynteny is largely maintained between the two species (Dieterich et al., 2008;

Rödelsperger et al., 2017) as well as other Rhabditids. Like other nematodes, *P.*

pacificus is thought to have holocentric chromosomes, which we confirm here through

cytological analysis. Recent improvements in the genome assembly (Rödelsperger et

al., 2017) and advances in genome editing (Lo et al., 2013; Namai and Sugimoto, 2018;

Witte et al., 2015) have facilitated investigation of developmental and cell biological

processes at a more mechanistic level.

We became interested in exploring meiosis in *P. pacificus* in part because of

differences in the inventory of meiotic genes predicted from its genome sequence

compared to *C. elegans.* In particular, genome sequencing and annotation have

revealed the presence of genes encoding an ortholog of Dmc1 as well as two Dmc1

cofactors, Mnd1 and Hop2, all of which are absent from the entire *Caenorhabditis* clade

(Figure 2 - Supplement 1 and (Dieterich et al., 2008)). Loss of Dmc1 correlates with the

adaptation of recombination-independent mechanisms for pairing and synapsis in

Drosophila and *Caenorhabditis* (Villeneuve and Hillers, 2001). Therefore, we were keen

to examine how homologous chromosomes pair and synapse in *P. pacificus.* In

addition, genetic linkage maps have suggested that multiple COs typically occur per

chromosome pair during meiosis in *P. pacificus* (Hong and Sommer, 2006; Srinivasan

et al., 2003, 2002), while only a single crossover per chromosome pair normally occurs

during meiosis in *C. elegans*, suggesting potential differences in crossover regulation.

Using genome editing, cytogenetic analysis, and recombination mapping, we

have characterized the early events of meiotic prophase in *P. pacificus*. We show that

homolog pairing, synapsis and CO recombination are dependent on *Ppa-spo-11* and

Ppa-dmc-1, while *Ppa-rad-51* is not essential for meiosis. We find that crossover sites

are designated very early during meiotic prophase, prior to completion of synapsis, and

are limited to one per chromosome pair. We also present a new genetic map, which

corroborates our cytological evidence that a single crossover normally occurs between

each pair of homologs per meiosis in in both spermatocytes and oocytes. The map also

provides evidence of crossover suppression and segregation distortion of specific

chromosomes in hybrids, indicative of reproductive barriers between strains. Our work

highlights both conserved and flexible features of the meiotic program within the

nematode lineage and establishes a platform for future investigation.

RESULTS

P. pacificus **as a comparative model system for meiosis**

The morphology and organization of the *P. pacificus* germline is very similar to

that of *C. elegans*. Hermaphrodites have two tubelike gonad arms in which sperm and

ova are produced sequentially, while males have a single arm (Rudel et al., 2005).

These are organized as a cylindrical monolayer of cells abutting a central core, or

rachis. The distal tip is populated by proliferating germline stem cells (Figure 1A, B, D),

as confirmed by the incorporation of microinjected fluorescently labeled nucleotides into

replicating DNA (Figure 1B). As in *C. elegans*, meiotic entry is marked by an obvious

change in nuclear organization in which fluorescently-stained chromosomes form a

conspicuous crescent-shaped mass (Figure 1A). Immediately proximal to this "transition

zone," DAPI staining reveals parallel tracks, indicative of paired and synapsed

homologous chromosomes at the pachytene stage. Gametogenesis switches from

spermatogenesis to oogenesis occurs during early adulthood. As oocytes approach

maturation, chromosomes undergo striking decondensation between diplotene and

diakinesis, a stage that has been referred to as the "growth zone" (Rudel et al., 2005).

Similar chromosome morphology is observed during a late "diffuse stage" of meiotic

prophase in many other eukaryotes (Zickler and Kleckner, 1999), but is rarely seen in *C.*

elegans. In the most proximal region of the hermaphrodite gonad, oocytes mature and

form a single row of large cells, and chromosomes condense dramatically as the nuclei

grow in size. Six bivalents can be detected as compact DAPI-staining bodies in oocytes

at diakinesis, the last stage of meiotic prophase preceding the first meiotic division

(Figure 1A, C, Rudel et al., 2005; Sommer et al., 1996).

It has been assumed that chromosomes in *P. pacificus* are holocentric, as in *C.*

elegans, but we are unaware of direct evidence to support this idea. We thus identified

a gene encoding CENP-C (HCP-4 in *C. elegans*), a conserved kinetochore protein, in

the *P. pacificus* genome and inserted a V5 epitope tag at its 3' end. The distribution of

CENP-C on mitotic chromosomes in embryos and mitotic germ cells confirmed their

holocentric organization (Figure 1D). Kinetochores appeared as linear structures along

the full length of chromosomes, rather than discrete foci. CENP-C also coated the

chromosomes during meiotic metaphase I (Figure 1C), as in *C. elegans* (Shakes et al.,

2009).

Stable homolog pairing requires early recombination factors

BLAST searches of the *P. pacificus* genome revealed an open reading frame

encoding an unambiguous ortholog of Dmc1, a meiosis-specific paralog of Rad51

(Table S1). Orthologs of the Dmc1 cofactors Mnd1 and Hop2 were also identified by

homology searches (Figure 2 – Supplement 1). By contrast, Dmc1/Mnd1/Hop2 are

absent from both *C. elegans* and *D. melanogaster*, two model organisms that have

evolved recombination-independent mechanisms of homolog pairing and synapsis

(Villeneuve and Hillers, 2001). We analyzed the genome sequences of other nematodes

to determine the evolutionary history of these genes within the nematode lineage. This

analysis revealed that Dmc1/Mnd1/Hop2 have been lost several times during the

evolution of nematodes, including the entire *Caenorhabditis* genus and all sequenced

members of Clade IV (Figure 2 –Supplement 1). As expected in light of its essential

function in DNA repair, the recombinase Rad51 was detected in all genomes examined

(data not shown).

In *C. elegans*, homolog pairing and synapsis require a family of zinc-finger

proteins (HIM-8, ZIM-1, ZIM-2, and ZIM-3) that bind to motifs enriched within Pairing

Center (PC) regions near one end of each chromosome (Phillips et al., 2009b). We

identified homologs of these proteins in many of the genome sequences from Clade V

nematodes, but not in *Pristionchus*. Because these homologs are present in species

that retain intact DMC-1 genes, it is unlikely that PC proteins arose as an alternative to

DMC-1 function in meiosis, although in *Caenorhabditis* they may have acquired a role

that eventually allowed degeneration and loss of DMC-1.

We tested whether homolog pairing in *P. pacificus* depends on double-strand

breaks or strand-exchange proteins. To generate *spo-11, dmc-1* and *rad-51* null

mutants, we first employed TALEN-mediated gene disruption, and later CRISPR/Cas9

genome editing techniques (this study; Lo et al., 2013; Witte et al., 2015). Genome

editing has thus far been less efficient in *P. pacificus* than in *C. elegans*, and techniques

such as co-CRISPR (in which another locus is edited to an allele with a visible

phenotype) have not been helpful to enrich for the desired edited progeny in our hands

or others' (Witte et al., 2015, and data not shown). Nevertheless, we were readily able

to isolate mutant alleles by screening a large number of F1 progeny from injected

hermaphrodites (see Materials and Methods). Independent alleles isolated from either

TALEN- or CRISPR-mediated genome editing resulted in identical mutant phenotypes.

All data presented here were based on alleles generated by CRISPR/Cas9. Because

balancer chromosomes are not currently available for *P. pacificus*, most mutations

described here were maintained in unbalanced heterozygotes, with frequent PCR-

based genotyping. Self-fertilization of heterozygotes results in broods with 25%

homozygous mutant animals, which were identified based on their phenotype and/or

immunostaining to determine the presence or absence of the protein of interest. We

also found that a freezing method developed for *C. elegans* that uses DMSO and

trehalose as cryoprotectants (Kevin F. O'Connell, Worm Breeders' Gazette, pers.

comm.) allowed robust recovery of frozen *P. pacificus*, facilitating strain maintenance

and archival storage.

As expected, disruption of either *spo-11* or *dmc-1* resulted in the detection of 12

DAPI-staining univalent chromosomes at diakinesis, indicative of a failure in CO

recombination (Figure 5). Surprisingly, *rad-51* mutants were fertile and displayed only

mild meiotic defects (see below). We thus validated the loss of RAD-51 function in

mutant animals by generating multiple alleles, which showed indistinguishable

phenotypes. We also confirmed the absence of RAD-51 protein by immunofluorescence

with a polyclonal antibody raised against recombinant Ppa-RAD-51 (Figure 4 –

Supplement 1B, see Materials and Methods). Mutations in *spo-11* or *dmc-1,* but not *rad-*

51, resulted in an obvious extension of the region of the germline displaying the

crescent-shaped nuclear morphology characteristic of early meiosis (Figure 2 –

Supplement 2). A similar "extended transition zone" phenotype is seen in *C. elegans*

mutants that fail to synapse their chromosomes during meiosis, suggesting that *spo-11*

and *dmc-1* might be required for synapsis in *P. pacificus.*

To visualize and quantify homolog pairing, we generated FISH probes against

two short tandem repeats found on Chromosomes X and IV (Figure 2A). We measured

the distance between pairs of homologous FISH signals in individual nuclei for each

genotype. To analyze pairing kinetics, we divided the distal gonads into five zones of

equal length. In zone 1, which contains mostly proliferating germ cells, pairs of FISH

signals remained far apart, with an average distance of 2.4 ± 1.0 µm (SD) and 2.5 ± 0.8

µm for Chromosomes X and IV, respectively (Figure 2B, D). In zone 2, which spans the

transition zone, the average distances decreased significantly in wild-type animals (1.2

± 1.1 µm and 1.1 ± 1.1 µm for probes on Chromosomes X and IV, respectively), and

homologous signals remained closely associated in the subsequent meiotic stages.

Notably, homologous FISH probes localized closer together, on average, in the two

zones immediately following meiotic entry, and showed some separation in zones 4 and

5 (Figure 2B, D). By contrast, in wild-type *C. elegans*, homologous loci remain closely

apposed in most throughout an extended pachytene stage spanning most of the distal

region (before the "loop") of the gonad (MacQueen, 2002). Together with analysis of

synapsis (below), this indicated that desynapsis initiates soon after completion of

synapsis in *P. pacificus*, resulting in partial separation of homologs.

We noted that the average distances between pairs of homologous FISH signals

in *spo-11* and *dmc-1* mutants also decreased markedly upon meiotic entry, although

clearly less so than in wild type (Figure 2D). In contrast, *rad-51* mutants showed

distributions of probe distances more similar to wild-type animals (Figure 2C, D). We

considered the possibility that the proximity between FISH signals might reflect the

clustering of all chromosomes during leptotene/zygotene, rather than specific

homologous interactions. If so, the extended transition zone morphology in *spo-11* and

dmc-1 might obscure a pairing defect that would be more apparent in the absence of

clustering (Figure 2 – Supplement 2). To address this, we measured the distances

between pairs of heterologous FISH signals in the premeiotic region (dispersed) versus

the transition zone (clustered). We observed that FISH signals on different

chromosomes were also significantly closer to each other in the transition zone

compared to premeiotic nuclei in both wild-type and mutant animals (Figure 2E). The

distances between heterologous versus homologous pairs of FISH loci were not

significantly different in *spo-11* and *dmc-1* mutants (p=0.1777 and p=0.6774,

respectively, by Student's *t*-test), but homologous signals were clearly closer than

heterologous signals in wild-type meiocytes (p<0.0001 by Student's *t*-test) (Figure 2E).

These data support the idea that clustering, rather than specific pairing, promote

proximity between both homologous and heterologous loci during leptotene/zygotene in

spo-11 and *dmc-1* mutants. Although we cannot conclude that transient homologous

pairing is absent in these mutants, it is evident that these early recombination factors

are required for stable pairing and extended association of homologous loci throughout

prophase. In contrast, *rad-51* is dispensable for homolog pairing, as in *C. elegans*.

SPO-11 and DMC-1 are required for homologous synapsis

To further investigate meiotic progression in *P. pacificus* and to probe the role of

early recombination factors in synapsis, we developed cytological markers for the

chromosome axis, which normally assembles upon meiotic entry, and the synaptonemal

complex (SC), which assembles between paired axes during early prophase.

Identification of a candidate axial element component was straightforward, due to the

presence of the easily recognized HORMA (Hop1, Rev7, Mad2) domain among

members of this family of proteins (Aravind and Koonin, 1998; Vader and Musacchio,

2014). We identified a gene encoding a HORMA domain protein that is most closely

related to *C. elegans* HIM-3 by reciprocal BLAST analysis (Table S1)*.* We refer to this

protein as Ppa-HOP-1, after the founding member of the meiotic HORMA proteins, *S.*

cerevisiae Hop1 (Hollingsworth and Byers, 1989, p. 1). Using genetic immunization, we

raised a polyclonal antibody against a 100-amino acid segment of the predicted protein

including part of the HORMA domain and found that it indeed recognized chromosome

axes from meiotic entry through late prophase (Figure 3A, B).

To enable cytological detection of synaptonemal complex (SC) assembly, we

searched for homologs of SC proteins. This was not straightforward due to rapid

divergence of these proteins in nematodes as well as their extensive regions of coiled-

coil potential, which is associated with a strongly skewed amino acid composition. *C.*

elegans expresses six known SC proteins, known as SYP-1–6. SYP-4 has a distinctive

C-terminal domain containing several unusual motifs enriched in glycine and

phenylalanine residues. We generated and sequenced an RNA library from isolated

gonads, which facilitated identification of full-length *Ppa-syp-4.* We inserted an HA

epitope tag at the C-terminus of the endogenous coding sequence, and found that

immunofluorescence with an epitope-specific antibody localized specifically between

paired meiotic chromosomes, confirming SYP-4::HA as a marker for the SC (Figure 3A,

B). The tagged protein supported normal meiosis, as indicated by the low percentage of

inviable embryos and males in the strain population (Figure 3 – Supplement 1).

HOP-1 was detected in the nucleoplasm in the premeiotic region of the germline

and formed linear structures along chromosomes upon meiotic entry. SYP-4 was

detected along chromosome segments shortly thereafter, and fully colocalized with

HOP-1 tracks during pachytene. Notably, the region of the germline containing nuclei

with fully aligned stretches of SYP-4 and HOP-1 was very short compared to *C.*

elegans, in which SC disassembly occurs close to the bend of the gonad arm, shortly

prior to diakinesis. In contrast, SC disassembly initiated much earlier in *P. pacificus*; the

major fraction of prophase nuclei are thus arrested at diplotene, since chromosomes are

only partially synapsed. Six short stretches of SYP-4 were apparent in these nuclei,

which persisted over an extended region (Figure 3A, B). This asymmetrical pattern of

SC disassembly is highly reminiscent of a more transient diplotene stage in *C. elegans*,

in which the SC remains associated with one "arm" of each homolog pair and

subsequently contributes to the step-wise loss of cohesion during the meiotic divisions

(Cloud et al., 2012; Da Ines et al., 2013).

HOP-1 localized normally to chromosome axes in *spo-11* and *dmc-1* mutants,

but extensive SC assembly failed. Instead, small, dispersed puncta of SYP-4 were

observed along chromosome axes, with occasional longer tracks (Figure 3C). In

contrast to *spo-11* and *dmc-1* mutants, *rad-51* mutants displayed robust synapsis with a

distribution of stages similar to that seen in wild-type hermaphrodites (Figure 3C).

These observations indicate that homologous synapsis depends on SPO-11 and DMC-

1 in *P. pacificus*, in contrast to the recombination-independent synapsis seen in *C.*

elegans.

DMC-1 and RAD-51 localize sequentially during distinct stages of meiotic

prophase

To investigate the functions of and interplay between DMC-1 and RAD-51 in *P.*

pacificus, we inserted a V5 epitope tag at the C-terminus of DMC-1 using CRISPR/Cas9

and raised a polyclonal antibody that recognizes RAD-51 (see Materials and Methods).

DMC-1::V5 supported normal meiosis, as evidenced by a normal brood size, high

embryonic viability and low percentage of males (Figure 3 – Supplement 1).

Surprisingly, the two proteins showed distinct and nonoverlapping patterns of

localization. DMC-1 localized very broadly along chromosomes, in transition zone nuclei

and disappeared immediately upon completion of synapsis. RAD-51 displayed a much

more restricted, punctate distribution along chromosomes, which was only detected in

nuclei in which DMC-1 no longer coated the chromosomes (Figure 4A, B). Occasional

nuclei at the border between the transition zone and pachytene region exhibited both

DMC-1 and RAD-51, although DMC-1 was very faint in these nuclei and did not overlap

with RAD-51 (Figure 4C). Additionally, DMC-1 remained strongly associated with

chromosomes in some late nuclei that retained clustered DAPI morphology, presumably

either "straggler" nuclei with delays in synapsis or CO designation, or apoptotic cells,

both of which are typically observed in the germlines of wild-type *C. elegans* (Figure

4D). Differences between the localization of DMC-1 and RAD-51 were further validated

by inserting a V5 epitope tag at the C-terminus of RAD-51. Staining of this strain with

anti-V5 recapitulated the sparse, punctate localization seen with anti-RAD-51

antibodies, demonstrating that the broader distribution of DMC-1 is not an artifact of the

V5 epitope tag or antibody (Figure 4 – Supplement 1A). RAD-51::V5 also supported

normal meiosis, with a normal brood size, high embryonic viability and low frequency of

male self-progeny (Figure 3 – Supplement 1).

We also tested the interdependence of DMC-1 and RAD-51 recombinases for

their localization. In *S. cerevisiae* and *A. thaliana*, Dmc1 functions as an essential

catalyst for interhomolog joint molecule (JM) formation during meiotic DSB repair, while

Rad51 acts as an accessory protein for Dmc1 nucleofilament formation (Cloud et al.,

2012; Da Ines et al., 2013). We did not detect RAD-51 in transition zone nuclei, where

DMC-1 was abundant on chromatin, and we found that DMC-1::V5 localization was

normal in *rad-51* mutants, indicating that RAD-51 does not play an essential role in the

recruitment of DMC-1 (Figure 4 – Supplement 1B). Conversely, in *dmc-1* mutants we

detected RAD-51 foci only in late prophase nuclei, proximal to the very extended

transition zone (Figure 4 – Supplement 1C). RAD-51 foci were more abundant and

larger in *dmc-1* mutants than in wild-type pachytene nuclei, perhaps due to persistent

unrepaired DSBs that accumulate in the absence of DMC-1 protein. Alternatively, the

bright foci of RAD-51 observed in late prophase nuclei could reflect an apoptotic

response to unrepaired breaks and/or extensive asynapsis.

The number of DMC-1 foci could not be quantified meaningfully due to their

density and wide variation in intensity, but the broad distribution suggests that DMC-1

associates with intact dsDNA in addition to sites undergoing recombination.

Nevertheless, in *spo-11* mutants DMC-1 was restricted to one or a few nuclear

aggregates rather than broadly along chromosomes (Figure 4 – Supplement 1D). It was

unclear whether these were associated with chromatin. This mislocalization may reflect

either an absence of potential binding sites due to an absence of DSBs, or *spo-11-*

dependent regulation of DMC-1 binding to chromatin, perhaps through activation of a

DNA damage signaling pathway.

In contrast, RAD-51 foci are much sparser along chromosomes, suggesting that

the protein localizes specifically to recombination intermediates. Thus, it was

unsurprising to see that RAD-51 foci were absent from meiotic nuclei in *spo-11* mutants

(Figure 4 – Supplement 1F). Some RAD-51 foci were observed in the mitotically

proliferating region the germline in *spo-11* mutants, as in wild-type animals, providing a

positive control for immunofluorescence (Figure 4 – Supplement 1E).

Together these observations indicate that DMC-1 and RAD-51 bind to chromatin

at distinct stages of meiotic prophase and are not interdependent, although both require

DSBs for their localization to chromosomes.

RAD-51 is not required for CO formation or completion of meiosis

To assess the roles of DMC-1 and RAD-51 in CO formation, we quantified the

number of DAPI-staining bodies at diakinesis in *dmc-1* and *rad-51* mutants. Wild-type

oocytes at this stage usually have six bivalents that can be resolved as discrete DAPI-

staining bodies (average = 5.6), while in *spo-11* mutants, ~12 DAPI-staining bodies

were detected (average= 11.5), consistent with an absence of COs (Figure 5A, B).

Interestingly, we frequently failed to detect oocytes at diakinesis in *dmc-1* mutant

germlines, indicative of a defect in meiotic progression and the likely activation of a

checkpoint in response to unrepaired DSBs. In cases when we did see nuclei at

diakinesis, we observed an average of 11.6 DAPI-staining bodies, reflecting an absence

of COs, as in *spo-11* mutants (Figure 5A-C).

Unexpectedly, disruption of *rad-51* resulted in homozygous mutant

hermaphrodites that were viable and fertile, although animals produced smaller broods

and their embryos showed greatly reduced viability (Figure 5D). This was surprising

because RAD-51 is essential for completion of meiosis in most organisms where it has

been examined. Homozygous *rad-51* mutant gonads also displayed diakinesis nuclei

more frequently than *dmc-1* mutants, although they were absent in 2 out of 20 gonads

scored, indicating that loss of DMC-1 function impairs meiotic progression more

severely than loss of RAD-51 (Figure 5C). Consistent with this observation, while self-

fertilizing *rad-51* mutants had a lower average brood size than wild-type

hermaphrodites, *dmc-1* mutants had even smaller broods, ranging from zero to 35

embryos laid per mutant homozygote (Figure 5D). In striking contrast to *C. elegans rad-*

51 mutants, which display chromatin aggregates and fragments at diakinesis (Martin et

al., 2005; Rinaldo et al., 2002), *Ppa-rad-51* mutants displayed an average of 6 DAPI-

staining bodies, similar to wild-type (Figure 5B). Together with the relatively high viability

of progeny of *rad-51* homozygous mutants, this indicates that RAD-51 does not play an

essential role in CO formation in *P. pacificus.*

COSA-1 marks designated CO sites during throughout early prophase

To further analyze CO formation in *P. pacificus*, we identified the gene encoding

the metazoan meiotic cyclin-related protein COSA-1 (Crossover Site Associated) /Cntd1

(Cyclin N-terminal Domain Homology) and inserted a 3xFLAG epitope tag at the C-

terminus of the coding sequence. The strain expressing COSA-1::3xFLAG yielded

progeny with high embryonic viability and few males, indicating that the tagged protein

supports normal meiosis (Figure 3 – Supplement 1). Immunostaining with anti-FLAG

antibodies revealed discrete foci along the SC, beginning as early as zygotene, which

decreased in number and became brighter within the short pachytene region (Figure

6A, B). Most pachytene nuclei displayed 6 COSA-1 foci, each of which was associated

with an individual SC, indicating the presence of a single designated crossover site

between each pair of homologs (Figure 6C and Supplemental Video 1). We stained J4

larval-stage hermaphrodites whose germlines had not yet undergone the switch from

spermatogenesis to oogenesis and found that pachytene spermatocytes also displayed

~6 COSA-1 foci (Figure 6- Supplemental Figure 1A). This suggests that *P. pacificus*,

like *C. elegans*, has robust chromosome-wide CO interference in both spermatogenesis

and oogenesis.

Intriguingly, SC disassembly appeared to be regulated by the position of the

designated CO site, as in *C. elegans.* By mid-prophase, 6 short stretches of SYP-4::HA

were observed, each associated with a single COSA-1::3xFLAG focus near one end

(Figure 6B). As meiosis progressed further, COSA-1::3xFLAG foci became

undetectable, although short stretches of SYP-4::HA could still be observed. We also

observed splaying of chromosome axes along the "long arms" on one side of the

COSA-1 focus upon disappearance of the SC from those regions (Figure 6D). HOP-1

was retained on both arms following SC disassembly, although the signal appeared

fainter along the long arms, perhaps due to separation of the two axes. At this stage,

short stretches of SYP-4::HA colocalize with corresponding bright stretches of HOP-1

(Figure 3B and Figure 6D). Bivalents at diakinesis and meiotic metaphase I also

displayed a cruciform structure similar to that seen in *C. elegans*, consistent with a

single chiasma per chromosome pair (Figure 1C).

In *C. elegans* the orientation of chromosomes during meiotic segregation is

stochastically determined by the position of the single crossover between each pair: the

"long arm," where SC first disassembles, is designated to retain cohesion and to lead

towards the poles while the "short arm" releases cohesion during MI, but then becomes

the leading end for segregation of chromatids during meiosis II (Albertson and

Thomson, 1993). Using FISH to mark specific chromosome regions together with

immunostaining of SYP-4::HA, we found that the same was true for *P. pacificus*: an

asymmetrically localized probe localized to either the long (desynapsed) arm or the

short arm in different nuclei (Figure 6- Supplemental Figure 2).

We examined the localization of COSA-1::3xFLAG in various mutant

backgrounds. As expected, *dmc-1* mutants showed a complete absence of COSA-1 foci

throughout prophase, while 6 foci were observed in pachytene nuclei in *rad-51* mutants

(Figure 7A), consistent with the number of DAPI-staining bodies observed at diakinesis

in these mutants (Figure 5B). A few bright COSA-1::3xFLAG foci were present

throughout prophase in *spo-11* mutants (Figure 7A). However, since ~12 DAPI-staining

bodies were observed during diakinesis, we conclude that these COSA-1 foci do not

mark designated COs. A similar phenomenon has been reported in C. elegans *spo-11*

mutants (Nadarajan et al., 2017; Pattabiraman et al., 2017), suggesting that COSA-1

and other crossover factors can coalesce at sites lacking bona fide recombination

intermediates when such structures are absent.

A genetic map for *P. pacificus* reveals conservation of the CO landscape

Prior work has led to divergent estimates of the meiotic recombination frequency

in *P. pacificus*. By summing over measured genetic intervals on the same linkage

group, the Sommer lab initially concluded that the map length of each chromosome

exceeded 100 cM, with a maximum length of 215 cM for Chromosome I, which is also

physically the longest (Hong and Sommer, 2006; Srinivasan et al., 2003, 2002).

However, a map derived by the same group based on RNA sequencing of ~F10

Recombinant Inbred Lines (RILs) yielded markedly shorter genetic lengths, below 100

cM per chromosome (Rödelsperger et al., 2017).

Since an understanding of the crossover landscape is an important reference for

analysis of meiosis, we addressed this ambiguity by constructing a new genetic map for

P. pacificus. We used three divergent parental strains from different geographic regions:

PS312 from California (CA), PS1843 from Washington State (WA), and RSB001 from

La Réunion island (LR) and mated them using a double-cross hybrid strategy. A total of

93 progeny from a cross between CA/WA hybrid males and CA/LR hybrid

hermaphrodites were sequenced (Figure 8A). This strategy allowed us to

simultaneously map crossovers that occur during hermaphrodite oogenesis and male

spermatogenesis from the same progeny (Figure 8B).

We draw several conclusions from the resulting crossover maps (Figure 8C).

First, the map lengths for each chromosome are close to 50 cM for both oocyte and

spermatocyte meiosis, very similar to genetic map lengths in *C. elegans*. This indicates

that a single crossover usually occurs between each pair of chromosomes per meiosis,

consistent with our cytological observations. Close to half of all chromatids inherited

from either parent are non-recombinant, as expected if a single crossover occurs

between two of the four possible chromatids. We did observe a few examples of double

crossovers in our data, but such events are rare (6/1012 chromatids). Second, the distal

regions ("arms") show higher recombination rates than the central regions of each

chromosome, as in *Caenorhabditis* (Barnes et al., 1995, Rockman and Kruglyak,

2009). This bias was also evident in a map constructed using RNA sequencing data

(Rödelsperger et al., 2017) and seems to be a widely conserved feature among

Rhabditids and perhaps other nematode clades (Doyle et al., 2018, Gonzalez de la

Rosa et al., 2021).

Third, the male spermatocyte map showed a pronounced asymmetry for

Chromosomes III and IV, with COs occurring predominantly on one arm, whereas the

corresponding oocyte maps show COs on both arms. Conversely, the oocyte-specific

map for the X chromosome shows an absence of crossovers on the right arm (the

single X chromosome does not undergo COs during male spermatogenesis) (Figure

8C). These findings suggest that crossovers were suppressed on one side of these

chromosomes in the hybrid parents. This suppression may reflect structural

rearrangements between the parental strains. Specifically, there are likely structural

differences between the Washington and California versions of Chromosomes III and

IV, and between the La Réunion and California versions of the X chromosome. A lack of

crossovers on the right arm of the X chromosome is also evident in data from RILs from

a cross between the same strains from La Réunion and California (Rödelsperger et al.,

2017). Crossover suppression on Chromosomes III and IV likely reflects intra- rather

than inter-chromosomal rearrangements, since we do not see evidence for

pseudolinkage, as would be expected if a III; IV translocation were present. Notably, the

asymmetry in crossover suppression also suggests that one arm of each chromosome

may contain a region that plays a dominant role in homolog pairing during meiosis,

perhaps related to the function of PCs in *C. elegans* (see discussion for more details).

This was unexpected given the absence of obvious orthologs of the zinc finger proteins

required for PC function in *C. elegans*, and the more "conventional" dependence of

pairing and synapsis on recombination that we have documented for *P. pacificus*.

Finally, we detected pronounced segregation distortion for chromosomes I and II

in the oocyte map, with inheritance of the La Réunion haplotype strongly favored

(Figure 8 – Supplement 1B). In each case non-recombinant chromosomes of the La

Reunion haplotype were about 8-fold more frequently inherited than the non-

recombinant California chromatids. These observations are consistent with either a bias

in meiosis segregation -- i.e., meiotic drive -- or the presence of loci on these

chromosomes that cause genetic incompatibilities (i.e., inviability of some F2 progeny).

Analysis of recombinant chromatids suggests that the latter scenario is more likely,

since a gradient in the severity of the distortion can be detected along the physical

length of the chromosomes, consistent with the idea that specific loci on the affected

autosomes may be toxic to hybrid progeny.

DISCUSSION

Distinct roles for DMC-1 and RAD-51

Comparison of the activities of Rad51 and Dmc1 *in vitro* has revealed similar

profiles: both RecA homologs bind preferentailly to single-stranded DNA and can

mediate strand-exchange interactions. However, Dmc1 is uniquely required during

meiosis. The distinct requirements for Rad51 and Dmc1 are thought to be due in part to

the activity of Dmc1 specific cofactors Mnd1 and Hop2, which confer different activities,

and/or a higher tolerance for mismatches by Dmc1, which may enable it to promote

recombination between nonidentical homologs (Steinfeld et al., 2019).

Our analysis of RAD-51 and DMC-1 in *P. pacificus* reveals their distinct

contributions during meiosis. The binding of these proteins to chromatin appears very

different; the large number of DMC-1 foci suggest that it binds to sites other than repair

intermediates. In contrast, RAD-51 displays a more restricted punctate localization, and

only after DMC-1 is largely removed from chromosomes upon completion of synapsis.

Intriguingly, both DMC-1 and RAD-51 depend on SPO-11 for their association with

meiotic chromosomes. We interpret these observations to indicate that DMC-1 is

positively regulated by a mechanism that responds to SPO-11 activity, likely through

activation of a DNA damage signaling pathway. In contrast, the association of RAD-51

with recombination intermediates may normally be inhibited until after the establishment

of CO intermediates between each pair of chromosomes.

The sequential localization of DMC-1 and RAD-51 first suggested that they

function independently, and this is further supported by our analysis of loss-of-function

mutations. In contrast to budding yeast and *A. thaliana*, we have found that RAD-51 in

P. pacificus is dispensable for the activity of DMC-1 in pairing, synapsis, and CO

formation. Instead, RAD-51 appears to play a supporting role in DSB repair during

pachytene, promoting repair of DSBs that remain after CO designation has occurred. In

C. elegans, which expresses only RAD-51, a similar switch between two modes of DSB

repair is nevertheless observed during meiotic prophase: association of RAD-51 with

repair intermediates is differentially regulated from the onset of meiosis until a mid-

pachytene transition that coincides with CO designation; at this time, cells' competence

to convert DSBs to interhomolog COs is also lost (Hayashi et al., 2007). An analogous

switch from a "meiotic" repair to a "somatic"-like repair pathway in mid-pachytene is also

observed in mouse spermatocytes (Enguita-Marruedo et al., 2019).

Taken together, it appears that DMC-1 and RAD-51 have specialized functions in

P. pacificus: formation of interhomolog CO intermediates by DMC-1 prior to completion

of synapsis, followed by a more generic mode of double-strand break repair by RAD-51.

Our observations that nuclei in *rad-51* mutant*s* display cruciform bivalents and lack

fragmented chromatin at diakinesis suggests that excess DSBs can be repaired through

an alternate pathway that does not depend on RAD-51 activity, such as non-

homologous end joining, or that DMC-1 can compensate for loss of RAD-51 but not vice

versa. Future studies may reveal how the activities of RAD-51 and DMC-1 are regulated

to accomplish an orderly hand-off during meiotic prophase.

Comparative analysis of meiosis reveals variations and similarities within the

nematode lineage

In addition to establishing key aspects of meiosis in *P. pacificus*, this work also

illuminates the evolutionary history of meiosis in *C. elegans*. A body of prior work has

revealed that recombination-independent homologous synapsis in *C. elegans* relies on

Pairing Centers (PCs), specialized chromosome regions that interact with nuclear

envelope and drive chromosome movement during early prophase. Similar roles in

chromosome movement and pairing during meiosis are typically mediated by telomeres,

but have shifted to a unique region on each chromosome in *C. elegans*. PCs in *C.*

elegans mediate synapsis-independent pairing and also act as the sites of synapsis

initiation (MacQueen 2005, Rog and Dernburg, 2013). In most other organisms,

telomere-led chromosome movement is thought to promote homologous interactions,

but stabilization of pairing and initiation of synapsis require and often occur at early

recombination intermediates, which depend on Spo11 and Dmc1 activity.

PC activity in *C. elegans* depends on and is largely defined by the recruitment of

a family of zinc finger proteins that bind to DNA sequence motifs in these regions

(Phillips et al., 2009b). These proteins, known as ZIM-1, ZIM-2, and ZIM-3, and HIM-8

in *C. elegans*, act as scaffolds to recruit a cascade of kinase activities required for

pairing and synapsis (Harper et al., 2011; Kim et al., 2015; Labella et al., 2011). Pairing

Centers have been regarded as an evolutionary alternative to "canonical" Dmc1-

mediated pairing mechanisms, so we found it intriguing that many of the sequenced

genomes of nematodes in Clade V include homologs of the HIM-8/ZIM family as well as

Dmc1, Hop2, and Mnd1 (Figure 2 – Supplement 1). The *Pristionchus* genus is unusual

within this Clade in that it lacks apparent homologs of the PC proteins, while

Caenorhabditis is among the few genera that lack Dmc1/Mnd1/Hop2, although these

genes were independently lost along the branch leading to *Oscheius tipulae*. This

suggests that the ability of PCs to trigger homologous synapsis in the absence of DSBs

and strand invasion may be recently derived, and perhaps restricted to *Caenorhabditis*.

The existence of PCs in *C. elegans* was initially inferred because chromosome

rearrangements often result in asymmetric crossover suppression: regions distal to

translocation breakpoints are inhibited from crossing over, while the chromosome region

containing the PC undergoes elevated recombination due to a feedback mechanism

that helps to ensure an "obligate crossover" (Rose and McKim 1992; Yu et al., 2016).

Based on the crossover suppression we observed in constructing a genetic map, it now

seems likely that *P. pacificus* may have a PC-like activity localized asymmetrically on its

chromosomes. This was unexpected given the absence of obvious homologs of the zinc

finger proteins required for PC function in *C. elegans*, and the more "conventional"

dependence of pairing and synapsis on recombination that we have documented for *P.*

pacificus.

If PCs are indeed conserved in *P. pacificus*, they clearly do not mediate

recombination-independent synapsis, as in *C. elegans*. Based on our cytological

observations, it is more likely that synapsis initiates at recombination intermediates in *P.*

pacificus, although this has not yet been tested rigorously. We speculate that Pairing

Centers may act as a source of a protein factor required for CO designation, and that

this factor must move from the source to the site of a recombination intermediate along

the SC. According to this model, the SC must eventually span the distance between

these two sites for a recombination intermediate to be designated as an eventual CO.

For example, in mammals Cdk2 is initially found at telomeres and later at crossover

intermediates; perhaps it moves along the SC, thereby ensuring that only recombination

intermediates linked in *cis* to homologous telomeres can eventually become crossovers.

Such a mechanism may help to account for the preservation of genetic linkage

units known as "Nigon elements" over long evolutionary periods, which has been

documented through genome sequencing of diverse nematodes (Foster et al., 2020;

Gonzalez de la Rosa et al., 2021; Tandonnet et al., 2019). We have previously

speculated that unique PC regions may have arisen in holocentric nematodes as part of

a meiotic system that promotes karyotype stability (MacQueen et al., 2005; Rog and

Dernburg, 2013), an idea that seems more viable in light of our indirect evidence for

their conservation in *Pristionchus*. When inversions arise that suppress local

recombination, they can lead to intraspecies reproductive barriers such as

toxin/antitoxin systems, and eventually to full speciation.

P. pacificus shares many experimental advantages with *C. elegans* for molecular

studies of meiosis. Naturally, it offers a more limited suite of experimental tools, since it

was more recently developed and is less widely studied. Genome engineering using

CRISPR/Cas9 is less efficient than in *C. elegans*, and has so far been refractory to

large insertions, such as transgenes encoding fluorescent proteins. However, this

barrier will likely be overcome, just as technical tweaks have greatly improved editing

efficiency in *C. elegans* in less than a decade. Another obstacle is the absence of

balancer chromosomes, which are invaluable for maintaining mutations in essential

genes in *C. elegans*. Notably, our genetic map data indicating the likely presence of

crossover-suppressing rearrangements between divergent strains of *P. pacificus*

suggests that it may be possible to propagate balancer chromosomes in this species,

which would be valuable additions to the experimental toolbox.

Our work has now established that the recombination landscape is remarkably

well conserved between *P. pacificus* and *C. elegans*. This was initially surprising in light

of the major differences in their regulation of synapsis and recombination. Indeed,

comparison of *P. pacificus* and *C. elegans* reveals that enrichment of COs on the

chromosome arms and robust chromosome-wide CO interference can be implemented

through divergent meiotic programs, *e.g.*, with synapsis either dependent on or

independent of crossover precursors, and with CO sites designated either before or

after completion of synapsis. Interestingly, other features of chromosome organization,

including enrichment for novel and developmentally-regulated genes on the arms, with

more ancestral and broadly expressed genes concentrated in the centers, as well as

chromosome-wide chromatin modification profiles, are also conserved across the

Rhabditina (Werner et al., 2018). How the recombination landscape is shaped by and/or

contributes to these genomic features is not well understood, but the conservation of

these features over tens of millions of years of evolution suggests that they are

important for genome transmission and/or function.

Nematodes are an ancient and widespread phylum in which most and perhaps

all species have holocentric chromosomes. The presence of spindle-attachment sites

along the length of chromosomes poses unique challenges for both mitosis and meiosis

(Dernburg, 2001; Melters et al., 2012). In monocentric organisms, the centromeres

and/or pericentromeric regions typically recruit factors such as Shugoshin to maintain

cohesion between sister chromatids during the first, reductional meiotic division, while

release of cohesion along chromosome arms allows segregation of recombinant

homologs (Watanabe and Kitajima, 2005). Holocentric chromosomes must use

alternate mechanisms to implement the stepwise loss of cohesion needed for two

orderly meiotic divisions. The asymmetric SC disassembly we observe in *P. pacificus*,

together with our evidence that the orientation of chromosomes during meiotic divisions

is stochastically determined by crossover position (Figure 6- Supplemental Figure 2)

likely reflects conservation of a mechanism discovered in *C. elegans* in which crossover

designation leads to asymmetric SC disassembly, retention of cohesion on one side of

the CO site during the first division, and release on the other side (Ferrandiz et al.,

2018; Martinez-Perez et al., 2008; Severson and Meyer, 2014; Tzur et al., 2012). Thus,

this particular solution to the problem of segregating holocentric chromosomes in

meiosis appears to predate the Rhabditine radiation. This pathway relies on the

formation of two bivalent "arms" with reciprocal functions, which in turn depends on

strict crossover interference; multiple crossovers "confuse" the system and lead to

frequent missegregation (Hollis et al., 2020). We expect that these two features of

meiosis are widely shared among and perhaps beyond Clade V nematodes. However,

they may not be universal among nematodes; for example, studies in Ascarids have

shown that heterochromatic regions near one end of a chromosome act as meiotic

centromeres and are eliminated in somatic cells (Goday and Pimpinelli, 1989; Melters et

al., 2012). This mechanism may allow for more relaxed CO control and/or a different

pattern of COs between homologous chromosomes.

Inexpensive genome sequencing and the adaptability of CRISPR/Cas-based

genome editing has enabled rapid progress in developing new experimental models

such as *P. pacificus* for molecular studies. Future studies in *P. pacificus* and other

nematodes will likely expand our understanding of core mechanisms and plasticity of

sexual reproduction.

MATERIALS AND METHODS

P. pacificus *strains and maintenance*

Animals were cultured on NGM media with *E. coli* OP50 at 20°C under the

standard conditions developed for *C. elegans* (Brenner, 1974). The wild-type strain in

which most mutations were generated was a derivative of PS312 designated as "97."

The Sommer lab found that this isolate was more amenable to genome editing by

CRISPR-Cas9 than the parental strain. Mutant alleles were maintained in unbalanced

heterozygotes. Every few generations and immediately prior to any analysis, single

adult hermaphrodites were picked to new plates and allowed to lay embryos for two

days, after which the genotype of the parent was determined by PCR genotyping.

Progeny from heterozygous mothers, 25% of which are homozygous for the meiotic

mutation, were analyzed using the assays described here; heterozygous and wild-type

siblings were frequently used as controls, in addition to unedited wild-type animals.

CRISPR/Cas9-mediated genome editing

To modify the *P. pacificus* genome, we adapted our preferred CRISPR/Cas9

protocol from *C. elegans* to *P. pacificus*. Equimolar quantities of Alt-R[®]CRISPR-Cas9

crRNA and tracrRNA molecules (Integrated DNA Technologies, Coralville, IA) were

hybridized using a thermocycler (95°C for 5 minutes, then 25°C for 5 minutes). 4µl of

100µM hybridized tracrRNA/crRNA was combined with 4µl of 40µM *S. pyogenes* Cas9-

NLS purified protein (QB3 MacroLab, UC Berkeley, Berkeley, CA) and incubated at

room temperature for 5 minutes. 2µl of 100µM stock of an Ultramer® DNA oligo (IDT)

repair template containing 50-60 bp homology arms and the desired epitope or mutation

sequence was added to the mixture, for a total volume of 10μl, and injected into the

gonads of young adult hermaphrodites aged 24 hours from the final larval (J4) stage.

Following a 2-4 hour recovery period, injected animals were allowed to lay embryos at

20°C for 16-20hr. Four days later, a fraction of the F1 population (typically 150-200

progeny from 6-8 injected P_0s) was screened for the presence of the desired mutation

or epitope tag sequence by PCR, and candidate alleles were verified by Sanger

sequencing. A complete list of crRNA, repair template, and genotyping primer

sequences used to generate alleles in this study is provided as Table S1.

TALEN constructs used in our initial analysis of *dmc-1* and *spo-11* were

generated using a protocol adapted from Zhang et al. (2011) and designed using the

TAL Effector Nucleotide Targeter 2.0 website (https://tale-nt.cac.cornell.edu/).

Brood analysis

To quantify embryonic viability, brood size, and male progeny of wild-type and

rad-51 mutants, J4 (virgin) hermaphrodites were picked onto individual plates and

transferred every 24 hours over 72 hours total. Embryos were counted each day, after

transferring the adult hermaphrodite to a new plate, and kept at 20°C. Three to four

days later, adults were counted on each plate. To analyze *spo-11* and *dmc-1* mutants,

24 individual J4 hermaphrodites were picked from progeny of a verified heterozygous

mutant hermaphrodite. Quantification was performed as in wild type, but after 72 hours,

the adult hermaphrodite was lysed and genotyped for the presence of the mutation.

Thus, although 24 animals total were quantified from a mixed population of *spo-11/+* or

dmc-1/+ animals, data from 5 homozygous *spo-11* and 7 homozygous *dmc-1* mutant

animals are reported in the data table, Figure 5D.

FISH probes

Probes targeting a central locus on Chromosome IV and the left end of

Chromosome X were designed based on two short tandem repeat motifs. Tandem

Repeat Finder v4.09 (Benson, 1999) was used to identify tandem repeats in *P. pacificus*

"El Paco" genome assembly (Rödelsperger et al., 2017) using default parameters and a

maximum periodicity of 200 bp. The output was then filtered to identify repeats that

spanned more than 8 kb. These were compared to the genome sequence using BLAST

to identify the subset of sequences restricted to a single major locus per genome. A

subset of these repeats was then tested for specific and robust hybridization with

oligonucleotide probes. The Chromosome IV probe

(TCATTGAAATGATCACAATCATTGA) targets a 30-base repeated motif

ATGATCATTGAAATGATCACAATCATTGAG, which spans 40.1 kb at a position 11.3

Mb from the left end of chromosome IV. The Chromosome X probe

(GGTGGTCGACGGCTGCGTCG) targets the 30-base repeat motif

GGTGGTCGACGGCTGCGTCGACTGAAGAGT, that spans two very close regions of

29.3 kb and 11.1 kb on the left end of the X chromosome. Single-stranded

oligonucleotides labeled at the 3' end with 6-FAM or Cy3 dyes were purchased from IDT

and used directly as FISH probes.

Antibodies

Antibodies against Ppa-RAD-51 were generated against a 6xHis-tagged N-

terminal fusion protein (aa 1-103) purified from *E. coli*. Four mice were immunized with

the antigen. Serum from one animal, designated S148, was used without purification at

1:300 dilution (Pocono Rabbit Farm and Laboratory, Canadensis, PA). Antibodies

against Ppa-HOP-1 were generated by genetic immunization against aa 177-276 (SDIX,

Newark, DE) and used in the following experiments at 1:300 dilution. Additional

antibodies were purchased from commercial sources and diluted as follows: mouse

anti-FLAG (1:500, Sigma #F1804), mouse anti-V5 (1:500, Thermo Fisher #R960-25),

rabbit anti-V5 (1:250, Millipore Sigma #V8137), goat anti-HA (1:500, Novus Biologicals

#NB600-362), and mouse anti α-tubulin, clone DM1A (1:400, Millipore #05-829).

Secondary antibodies raised in donkey and labeled with Alexa 488, Cy3, or Cy5 were

used at 1:400 dilution (Jackson ImmunoResearch Laboratories).

Cytological analysis

To stage animals for each experiment, 30-40 J4 larvae were picked from a PCR-

verified heterozygous mother onto a fresh plate and allowed to develop for an additional

24 or 48 hours at 20°C. Immunofluorescence and FISH methods for germline tissue

were adapted from *C. elegans* (Phillips et al., 2009a), with minor modifications. Briefly,

adult hermaphrodites were dissected on a clean coverslip in egg buffer containing

0.05% tetramisole and 0.1% Tween-20. Samples were fixed for 2 minutes in egg buffer

containing 1% formaldehyde and then transferred to a 1.5-ml tube containing PBST.

After 5 minutes, the buffer was replaced with ice-cold methanol and incubated at room

temperature for an additional 5 minutes. Worms were washed twice with PBST, blocked

with Roche blocking reagent, and stained with primary antibodies diluted in Roche

blocking solution for 1.5-2 hours at room temperature. Samples were washed with

PBST and incubated with secondary antibodies raised in donkeys and conjugated with

Alexa-488, Cy3 or Cy5 (Jackson ImmunoResearch Laboratories, West Grove, PA).

Worms were then incubated with 1µg/ml DAPI in PBST, washed with PBST, and

mounted in ProLong™ Gold (Invitrogen) before imaging.

For immunofluorescence in embryos, mixed-stage worms were washed off of 20

plates in distilled water, pelleted, and treated with household bleach diluted 8-fold with

distilled water for 5 minutes at room temperature. Embryos were collected by

centrifugation and washed twice with PBS. To dissolve the vitelline membrane, a

solution containing 2.6ml of n-heptane, 2ml of PBS, and 1% paraformaldehyde (final)

was added to the embryo pellet for 5 minutes at room temperature with shaking.

Treated embryos were collected by centrifugation, washed twice with 5ml MeOH, three

times with PBS, and incubated with Roche blocking reagent, primary, and secondary

antibodies as described above. Embryos were mounted on agarose pads for imaging.

To label replicating DNA in proliferating germline stem cells, gonads of age-

matched hermaphrodites were injected with 0.3nM Cy3-dUTP solution. After 30 minutes

of recovery on seeded plates, worms were dissected and fixed as for

immunofluorescence, stained with DAPI, and mounted for imaging.

For FISH experiments, age-matched animals were dissected and fixed as for

immunofluorescence experiments described above, except that the initial fixation was

for 4 minutes in 2% formaldehyde. After incubation in ice-cold methanol, worms were

washed with 2x SSCT twice and incubated in 50% 2x SSCT/50% formamide solution

overnight at 37°C. The next day, the worms were transferred to a small PCR tube,

excess solution was removed from the sample, and 40μl of hybridization mix containing

250ng of each probe in hybridization buffer (3.5x SSC, 58% formamide, 12.75% dextran

sulfate) was added. DNA was denatured by incubation in a thermocycler at 91°C for 2

minutes and incubated overnight at 37°C. Following overnight hybridization, samples

were transferred to a 1.5-ml tube and washed twice with 2x SSCT. After 5 minutes, the

solution was replaced with fresh 2x SSCT and mounted with ProLong™ Diamond

Antifade Mountant with DAPI (Invitrogen).

Images were acquired as stacks of optical sections at 0.2 μm z-spacing using a

DeltaVision Elite microscope (GE) with an Olympus 100X NA 1.4 objective. All data

were deconvolved using the constrained iterative algorithm included with the softWoRx

package (GE) using default cycle numbers and other settings, and maximum-intensity

projections were generated for most images presented here. Distances between FISH

probes were calculated using 3D peak finding and distance calculation functions within

softWoRx. Images were pseudocolored using softWoRx, Fiji (Schindelin et al., 2012)or

Adobe Photoshop.

Genetic map construction

We generated a genetic map of *P. pacificus* using a double-cross hybrid strategy

that allowed the simultaneous mapping of meiotic crossovers in male spermatocytes

and hermaphrodite oocytes (Figure 8A). We began with three parental lines (P0): "CA,"

the "wild type" reference strain PS312 from California; "WA," PS1843 from Washington

State; and "LR," RSB001 from La Réunion Island. These three strains have been

previously used to generate genetic linkage maps (Rödelsperger et al., 2017,

Srinivasan et al., 2003, 2002). We crossed these P0 lines to obtain F1 hybrid CA/WA

males and CA/LR hermaphrodites. These F1 hybrid types were then mated to produce

the F2 generation, so that each F2 inherits a recombinant genome from both of its F1

parents.

F2 progeny were genotyped using a custom sequencing approach: juvenile

(unmated) F2 hermaphrodites were placed on individual 60-mm plates and allowed to

self-fertilize for 2-3 generations until their bacterial food was depleted. Worms were

washed from each plates and genomic DNA was extracted using Qiagen DNeasy Blood

& Tissue Kit (Qiagen) and quantified by Qubit (Invitrogen). Because self- and cross-

progeny can be temporally intermingled among broods of *P. pacificus*, we genotyped

each sample using several SSLP markers (Dieterich et al., 2008) to confirm the

presence of the PS1843 (WA) haplotype (Table S3). A total of 93 validated F2 samples

were used to construct genomic libraries for whole-genome sequencing using the

plexWell LP384 Library Preparation Kit (seqWell), with ~5x shotgun sequence per F2

sample.

To obtain a set of reference variants for use in genotyping, we sequenced the

three parental lines: CA and LR at 86x depth and WA at 61x, and mapped to the CA "El

Paco" reference genome using bwa-mem2 (v2.0pre2-4-gf06fc0c) (Vasimuddin et al.,

2019). Using freebayes (v1.3.1-17-gaa2ace8) (Garrison and Marth, 2012), with the --

standard_filters option, we identified 1,395,065 polymorphic sites with that were (1)

homozygous for each P0 genotype and within three standard deviations of the mean

depth for all three (CA, 47-125, LR, 44-125, WA, 20-111) and (2) biallelic, with a

different nucleotide in one strain relative to the other two. There were 374,862 such

cross-informative sites where CA differed from both WA and LR; 490,484 where LR

differed from CA and WA; and 529,719 where WA differed from CA and LR. Importantly,

these sites were distributed across the genome.

We genotyped each F2 sample for these cross-informative variants in non-

overlapping 100 kb blocks spanning the genome. Our crossing scheme gives rise to

four possible genotypes at each position (CA/CA, CA/LR, CA/WA, and WA/LR) that can

be easily differentiated based on the relative total depth f_{CA}, f_{WA}, and f_{LR} at cross-

informative variant sites within a block, where $f_{CA} + f_{WA} + f_{LR} = 1$. Specifically, we called

CA/CA if $f_{CA} > 0.95$, CA/LR if $f_{WA} < 0.05$ and $|f_{CA} - f_{LR}| > 0.1$, etc. (Figure 8 – Supplemental

figure 1A). Genotypes not satisfying one of these inequalities were not called. Genomic

blocks that contained fewer than 100 diagnostic sites or for which more than 30% of F2

individuals could not be called were not genotyped. Of the 1,567 non-overlapping 100

kb blocks spanning the genome, 1,398 remained after filtering for too few sites, too few

individuals genotyped, and segregation distortion ($p \leq 0.001$).

By design, our cross generates sex-specific genetic maps, since WA haplotypes

can only be transmitted via the sperm of the CA/WA male parent and LR haplotypes

can only be transmitted to ova of the CA/LR hermaphrodite parent. Genetic linkage

maps were constructed with OneMap (Margarido et al., 2007) using the "F2 backcross"

population type and the "record" algorithm to order markers.

All sequencing data have been submitted to the NIH Sequence Read Archive

under accession number PRJNA734516.

Orthology analysis and phylogenetic inference

Accessions to all data used in the orthology analysis and phylogenetic inference

are available in Table S2. We downloaded the genomes and annotation files for 66

nematode species (selecting a maximum of two species per genus) from WormBase

ParaSite (Howe et al., 2017) and two tardigrade outgroup taxa (*Hysibius dujardini* and

Ramazzottius varieornatus) from tardigrades.org. We selected the longest isoform of

each protein-coding gene in each species using AGAT (version 0.5.1; (Dainat et al.,

2020). To infer the nematode phylogeny, we ran BUSCO (version 5.0.0; using the

'metazoa_odb10' dataset (Simão et al., 2015) on all filtered proteomes and used

busco2fasta.py (available at https://github.com/lstevens17/busc(Camacho et al.,

2009)o2fasta) to extract 534 single-copy orthologs that were present in at least 70% of

species. We aligned the orthologous sequences using MAFFT (version 7.475; (Katoh

and Standley, 2013)) and inferred gene trees using IQ-TREE (version 2.1.2; (Nguyen et

al., 2015)), allowing the substitution model to be automatically selected

(Kalyaanamoorthy et al., 2017). We provided the resulting gene trees to ASTRAL-III

(version 5.7.4; (Zhang et al., 2018) to infer a species tree. The resulting tree was

visualized using iTOL (Letunic and Bork, 2016).

To investigate the presence of meiotic pairing proteins across the nematode

phylogeny, we used OrthoFinder (version 2.5.2; (Emms and Kelly, 2015)) to cluster the

longest isoforms of each protein-coding gene into orthologous groups (OGs). We

identified the *P. pacificus* orthologs of DMC-1, MND-1, HOP-2, and RAD-51 by

searching the proteome using BLASTP (version 2.5.0+; with the orthologous protein

sequences from *Homo sapiens* as queries and identified the corresponding OGs. We

identified the OG containing the HIM-8/ZIM family using the *C. elegans* transcript IDs.

We aligned each OG using MAFFT and inferred a gene tree using IQ-TREE as

previously described. Each gene tree was visually inspected using iTOL and presence

or absence of a homolog was recorded for each species. For those species that lacked

an apparent MND-1 or HOP-2 ortholog, we searched the genome sequence using

TBLASTN with the *P. pacificus* orthologs as a query (recording evidence of a homolog

as one or more hits with an E-value of < 1e-5). For species which lacked a DMC-1

ortholog, we searched the genome sequence using both the *P. pacificus* DMC-1 and

RAD-51 protein sequences as query, and considered hits (E-value < 1e-5) for DMC-1

that were non-overlapping with those for RAD-51 as evidence of a DMC-1 ortholog. For

species which lacked a member of the HIM-8/ZIM family, we searched the genome

sequence using TBLASTN with the homologous sequence from its closest relative as a

query.

Acknowledgements

This work was supported by funding from the Howard Hughes Medical Institute

and the Miller Foundation to AFD, as well as a fellowship from the Helen Hay Whitney

Foundation to JB. This work used the Vincent J. Coates Genomics Sequencing

Laboratory at UC Berkeley, supported by NIH S10 OD018174 Instrumentation Grant.

We are grateful to Ralf J. Sommer, Christian Rödelsperger, Ray Hong, and other

members of the *P. pacificus* research community for providing strains and abundant

helpful advice. We thank members of the Dernburg Lab for helpful discussions about

this work and critical reading of the manuscript. We are grateful to the thoughtful

reviewers of the original version of our manuscript, particularly for spurring us to

produce a genetic map.

Figure Legends

Figure 1. Germline organization and meiotic nuclear morphology in *P. pacificus* are

superficially similar to *C. elegans.* A. Projection image of the distal arm of a *P. pacificus*

hermaphrodite gonad stained with DAPI. Scale bar, 30 µm. Insets show representative

nuclei from the premeiotic region (PM), transition zone (TZ), pachytene (Pach),

diplotene (Dip), diffuse stage (Diff) and diakinesis. Scale bar, 5 µm. B. Distal region of a

P. pacificus germline following injection of fluorescent nucleotides to label replicating

DNA. Scale bar, 30 µm. C. Metaphase I oocyte expressing CENP-C::V5, stained with

anti-V5, DAPI, and anti-tubulin. Scale bar, 10 µm. D. Mitotic chromosomes (DAPI) in the

premeiotic germline of adult hermaphrodites and a 2-4 cell stage embryo and

expressing CENP-C::V5 (magenta). Scale bar, 2 µm.

Figure 2. Stable homolog pairing requires DSBs and strand invasion. A. Diagram

showing the locations of tandem repeat sequences used to generate DNA FISH probes

for pairing analysis in *P. pacificus*. B. Representative images show the progression of

homolog pairing of Chromosome X (magenta) and Chromosome IV (yellow) during

meiotic prophase in wild-type hermaphrodites. Premeiotic region (PM), transition zone

(TZ), and pachytene (Pach). Scale bar, 5 µm. C. Representative images of FISH probe

signals in *spo-11*, *dmc-1*, and *rad-51* mutants during mid-prophase stage (roughly

equivalent to the pachytene stage in wild-type germlines). Scale bar, 5 µm. D. Temporal

progression of X and IV chromosome pairing in WT, *spo-11, dmc-1*, and *rad-51*

mutants. Graphs show the distribution of distances within each of 5 equally-sized zones

spanning meiotic prophase. E. Distance between pairs of heterologous FISH signals

were measured in premeiotic (PM) and transition zone (TZ) nuclei in WT, *spo-11* and

dmc-1 mutants (spanning zones 1 and 2 only). Distances between pairs of homologous

FISH signals (Chr. X and IV combined) in TZ nuclei are included for comparison.

***p<.0001, by Student's *t*-test.

Figure 2- figure supplement 1. Presence of meiotic pairing proteins across the

nematode phylogeny. D= Dmc1; M=Mnd1; H=Hop2; Z=HIM-8/ZIM-1,2,3. Circles

indicate the presence of a homolog; solid outlines indicate homologs identified via

orthology clustering; dashed outlines represent evidence of a homolog from TBLASTN

searches of the genome sequence. *C. elegans* and *P. pacificus* are highlighted in blue.

Major nematode clades are highlighted. Phylogeny rooted on two tardigrade species

(*Hysibius exemplaris* and *Ramazzottius varieornatus*); not shown. Branch lengths are

not shown.

Figure 2- figure supplement 2. A. Composite projection images of whole gonads

stained with DAPI from WT, *spo-11, dmc-1*, and *rad-51* mutants. The extent of the

transition zone of each gonad is indicated with dashed lines. Scale bar, 30 µm. B.

Quantification of transition zone length as a fraction of the distance from meiotic onset

to the end of pachytene in WT (n=7), *spo-11* (n=7), *dmc-1* (n=7), and *rad-51* (n=10).

Error bars indicate mean ± SD. *spo-11* and *dmc-1* mutants show significant differences

from WT (p<0.0001) but not *Ppa-rad-51* (p=0.8426) by ordinary one-way ANOVA.

Figure 3. SPO-11 and DMC-1 are required for homologous synapsis in *P. pacificus*,

while RAD-51 is dispensable. A. Composite projection image of a wild-type strain

expressing SYP-4::HA, stained with DAPI (gray), anti-HOP-1 (red), and anti-HA (green).

Meiosis progresses from left to right. Scale bar, 30 µm. B. Higher magnification images

of wild-type nuclei in the premeiotic region (PM), transition zone (TZ), pachytene (Pach),

and diplotene (D) stages. C. Localization of SYP-4::HA and HOP-1 in WT, *spo-11*, *dmc-*

1, and *rad-51* mutants during early and mid-prophase (roughly equivalent to the TZ and

pachytene regions in wild-type germlines, respectively). Synapsis fails in the absence of

spo-11 and *dmc-1* function but occurs normally in *rad-51* mutants. Scale bar, 5 µm.

Figure 3- figure supplement 1. All epitope-tagged proteins described in this study

support normal meiosis. Epitope-tagged alleles were generated by in-frame insertion

into the endogenous gene loci using CRISPR/Cas9 (see Materials and Methods), and

homozygosed by selfing of successfully edited progeny. Fidelity of meiotic segregation

in the resulting strains was analyzed by counting the frequency of viable embryos and

male progeny among whole broods from self-fertilizing hermaphrodites, as indicated.

Quantification of wild-type broods is also reported in Figure 5D.

Figure 4. DMC-1 and RAD-51 localize sequentially to meiotic chromosomes. A.

Composite projection image of a wild-type gonad expressing DMC-1::V5, stained with

DAPI (blue), anti-V5 (magenta), and anti-RAD-51 (yellow). Meiotic progression is from

left to right. Scale bar, 30 μm. Inset shows the distinct localization of DMC-1 (magenta)

and RAD-51 (yellow) in the transition zone and pachytene regions, respectively. Scale

bar, 5 μm. B. Higher magnification images of nuclei in the transition zone and pachytene

region. DMC-1 is present along chromatin in the transition zone and disappears at

pachytene. By contrast, RAD-51 localizes to discrete foci starting at pachytene. Scale

bar, 5 μm. C. Occasional nuclei at the transition from leptotene-zyogtene to pachytene

are positive for both DMC-1 and RAD-51. The signals do not completely overlap. Scale

bar, 2 μm. D. Example of a nucleus with polarized DAPI morphology and strong DMC-1

signal during later prophase. Such "straggler" cells may be delayed in completing

synapsis or undergoing apoptosis. Scale bar, 2 μm.

Figure 4- figure supplement 1. A. Localization of RAD-51::V5. Composite projection

image of a wild-type gonad expressing RAD-51::V5, stained with DAPI and anti-V5.

Meiotic progression is from left to right. Scale bar, 30 μm. DMC-1 and RAD-51 localize

to meiotic chromosomes independently of each other, but both require SPO-11. B.

DMC-1 (magenta) is abundant on chromosomes in transition zone nuclei of *rad-51*

mutants. Anti-RAD-51 immunofluorescence was used to identify homozygous mutants

among the progeny of heterozygotes. C. RAD-51 foci are observed in late pachytene

nuclei of *dmc-1* mutants, proximal to the extended transition zone. RAD-51 foci appear

larger and more numerous than in wild-type pachytene nuclei. D. DMC-1 forms nuclear

aggregates in *spo-11* mutants and does not localize along chromosomes. E and F.

RAD-51 foci are detected in premeiotic nuclei of *spo-11* mutants but are absent during

meiotic prophase. Scale bar, 5 μm.

Figure 5. CO formation requires SPO-11 and DMC-1, but not RAD-51. A.

Representative images of DAPI-staining bodies at diakinesis for each indicated

genotype. Scale bar, 5 μm. B. Quantification of DAPI-staining bodies in the "-1" oocyte

(immediately distal to the spermatheca) at diakinesis for each indicated genotype (*n*=

represents number of nuclei scored). C. Quantification of gonads which lacked nuclei

with DAPI-staining bodies at diakinesis stage. *n* is the number of germlines scored for

each genotype. D. Frequencies of viable embryos and male progeny of whole broods

from wild type, *spo-11*, *dmc-1* and *rad-51* mutant hermaphrodites.

Figure 6. COSA-1 accumulates at a single focus per chromosome pair. A. Composite

projection image of a wild-type strain expressing three epitope-tagged proteins (COSA-

1::3xFLAG, DMC-1::V5, and SYP-4::HA), stained with anti-FLAG, anti-V5 and anti-HA

antibodies. Scale bar, 30 μm. B. Higher magnification images of nuclei from the

transition zone (TZ), pachytene (pach), mid- and late diplotene (D). COSA-1::3xFLAG

(green) foci are detected in transition zone nuclei but do not colocalize with DMC-1::V5

(cyan). Foci peak in brightness in pachytene nuclei and gradually become dimmer until

they are no longer detected during late diplotene. In early to mid-diplotene nuclei, six

short stretches of SYP-4::HA (magenta) are observed per nucleus, each associated

with a single COSA-1::3xFLAG focus. Scale bar, 5 μm. C. Histogram showing the

number of COSA-1::3xFLAG foci observed per nucleus in the pachytene region.

Analysis was restricted to 15 nuclei per gonad immediately proximal to the transition

zone and lacking DMC-1::V5 signal. Five individual gonads were analyzed, for a total of

75 nuclei scored. D. Partial projection of a representative nucleus in mid to late

diplotene, stained with anti-HOP-1 (blue), anti-HA (marking the SC, magenta), and anti-

FLAG (marking COSA-1, green). A single COSA-1::3xFLAG focus is observed at a

junction (marked with a red arrowhead) between the "short arm," where SYP-4::HA is

retained, and splayed "long arms" lacking SC but positive for HOP-1. Scale bar, 2 µm.

1013

Figure 6 - figure supplement 1. COSA-1 accumulates at a single site per chromosome

pair during spermatogenesis. A. Composite projection image of a wild-type gonad from

a J4-stage hermaphrodite expressing COSA-1::3xFLAG (green) and SYP-4::HA

(magenta). Scale bar, 30 µm. At this stage the germline is still undergoing

spermatogenesis. Inset shows a higher magnification image of nuclei in the pachytene

region. As during oogenesis, ~6 COSA-1::3xFLAG foci are observed in pachytene

nuclei during spermatogenesis. Scale bar, 5 µm.

1021

1022 **Figure 6 - Supplemental Video 1.** COSA-1/Cntd1 accumulates at a single site per

1023 chromosome pair. 3D volume rendering of a single nucleus from the pachytene region

1024 of a wild-type worm expressing COSA-1::3xFLAG (green) and SYP-4::HA (magenta).

1025 Each frame is a maximum intensity projection from a region of a deconvolved 3D image.

1026 Each stretch of SYP-4::HA is associated with a COSA-1::3xFLAG focus.

1027

Figure 6 - figure supplement 2. Asymmetric disassembly of the synaptonemal

complex can occur along either arm of bivalent chromosomes. Projection images of

representative wild-type diplotene nuclei expressing SYP-4::HA, hybridized with a

fluorescent probe specific for the left end of chromosome X (magenta), and stained with

anti-HA (yellow) and DAPI (blue). A. Nuclei in which the FISH signal recognizes the

"long arm" lacking SYP-4::HA. B. Nuclei in which the probe recognizes the "short arm"

that retains SYP-4::HA. Scale bar, 2 μm.

1035

Figure 7. COSA-1::3xFLAG accumulates at sites of presumptive COs. A. Nuclei from

hermaphrodites of the indicated genotype displaying COSA-1::3xFLAG (green) in early

and mid-prophase (roughly equivalent to the transition zone and pachytene regions in

wild-type germlines, respectively). COSA-1 foci are absent in *dmc-1* mutants, but 6 foci

per nucleus are detected in wild type and *rad-51* mutants. Occasional foci are detected

in s*po-11* mutants. Scale bar, 5 μm.

1042

Figure 8. A. Crossing scheme to generate a recombination map using three parental

strains. California PS312 (CA), Washington PS1843 (WA), and La Réunion Island

RSB001 (LR) strains were crossed to obtain F1 hybrids, which were then crossed to

each other. Whole genome sequencing of progeny from crosses between hybrid F1s

enabled the analysis of meiotic recombination events in each F1 parent. B. Genotype

transitions along a chromosome in F2 correspond to recombination in the male or

hermaphrodite F1 parent. C. Marey plots show genetic map position in centimorgans vs.

the physical position in megabases for male (green) and hermaphrodite (magenta)

meiosis. Each bin was treated as a single locus and dots were plotted at the center of

each marker bin. Map positions were computed with OneMap as described. The

observed map length of ~50 cM indicates that chromosomes undergo an average of 1

CO per meiosis. The X chromosome lacks a homolog in males, so there is no male-

specific map for the X in our data.

1056

Figure 8 – figure supplement 1. A. Genotyping. For each 100 kb genomic block,

progeny were genotyped based on the number of CA-, WA-, and LR-specific variants

they contained, given the expectation of four possible genotypes CA/CA, CA/LR,

WA/CA, WA/LR. In scatterplot each point represents s a single bin in a specific

individual, with $X = f_{WA} + f_{LR} - 2f_{CA}$, and $Y = f_{LR} - f_{WA}$. Gray dots represent genotypes that

are conservatively left as "not called." B. Genotypes of the mapping panel. F2

genotypes inherited from male (left) and hermaphrodite (right) parents. Each thin row is

one progeny. White vertical stripes represent 100 kb windows that could not be reliably

genotyped due to low numbers of P0-specific variant sites and/or sequencing depth of

progeny.

1067

1068 **Table S1.** Genes, alleles, and genome editing reagents used in this manuscript.

1069 **Table S2.** Species names and sequence accession numbers used for the phylogenetic

1070 analysis summarized in Figure 2 – Figure Supplement 1.

1071

1072 **Table S3.** SSLP markers and genotyping primers used to confirm cross-progeny for

1073 genetic map construction.

1074

1075   **Table S4.** Software version numbers and parameters used in this work.

1076

1077   **Table S5.** Genotype calls for the 93 individual F2 progeny used for mapping crossovers

1078   in hermaphrodite oocytes. Each column represents one hybrid progeny in arbitrary

1079   order. "A" genotypes are inherited from the California PS312 (CA), F0 parent, "R"

1080   genotyes are inherited from the La Réunion Island RSB001 F0 parent

1081

1082   **Table S6**. Genotype calls for the 93 individual F2 progeny used for mapping crossovers

1083   in male spermatocytes. Each column represents one hybrid progeny in arbitrary order

1084   (same order as in Table S5). The "A" genotypes were inherited from the California

1085   PS312 (CA), F0 parent, "W" genotypes are inherited from the Washington PS1843 F0

1086   parent.

1087

1088   **Table S7.** Map distances in centimorgans (cM) generated with OneMap based on the

1089   data in Tables S5 and S6, using the 'record' algorithm.

1090

1091   **Table S8.** Summary of crossovers detected on each chromosome in oocytes (XX) and

1092   spermatocytes (XO), including the number of nonrecombinant chromatids and

1093   chromatids with 1 or 2 crossovers. No higher-order recombinant chromatids were

1094   observed.

1095

Literature and software cited

1096

Albertson DG, Thomson JN. 1993. Segregation of holocentric chromosomes at meiosis in the
nematode,Caenorhabditis elegans. *Chromosome Res* **1**:15–26.
doi:10.1007/BF00710603
Aravind L, Koonin EV. 1998. The HORMA domain: a common structural denominator in mitotic
checkpoints, chromosome synapsis and DNA repair. *Trends in Biochemical Sciences*
23:284–286. doi:10.1016/S0968-0004(98)01257-2
Barnes TM, Kohara Y, Coulson A, Hekimi S. 1995. Meiotic recombination, noncoding DNA and
genomic organization in Caenorhabditis elegans. *Genetics* **141**:159.
Bergerat A, de Massy B, Gadelle D, Varoutas P-C, Nicolas A, Forterre P. 1997. An atypical
topoisomerase II from archaea with implications for meiotic recombination. *Nature*
386:414–417. doi:10.1038/386414a0
Bishop DK, Park D, Xu L, Kleckner N. 1992. DMC1: A meiosis-specific yeast homolog of E. coli
recA required for recombination, synaptonemal complex formation, and cell cycle
progression. *Cell* **69**:439–456. doi:10.1016/0092-8674(92)90446-J
Brenner S. 1974. The genetics of Caenorhabditis elegans. *Genetics* **77**:71–94.
Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009.
BLAST+: architecture and applications. *BMC Bioinformatics* **10**:421. doi:10.1186/1471-
2105-10-421
Christophorou N, Rubin T, Bonnet I, Piolot T, Arnaud M, Huynh J-R. 2015. Microtubule-driven
nuclear rotations promote meiotic chromosome dynamics. *Nat Cell Biol* **17**:1388–1400.
doi:10.1038/ncb3249
Christophorou N, Rubin T, Huynh J-R. 2013. Synaptonemal Complex Components Promote
Centromere Pairing in Pre-meiotic Germ Cells. *PLoS Genet* **9**:e1004012.
doi:10.1371/journal.pgen.1004012
Cloud V, Chan Y-L, Grubb J, Budke B, Bishop DK. 2012. Rad51 Is an Accessory Factor for
Dmc1-Mediated Joint Molecule Formation During Meiosis. *Science* **337**:1222–1225.
doi:10.1126/science.1219379
Couteau F, Belzile F, Horlow C, Grandjean O, Vezon D, Doutriaux M-P. 1999. Random
Chromosome Segregation without Meiotic Arrest in Both Male and Female Meiocytes of
a dmc1 Mutant of Arabidopsis. *The Plant Cell* **11**:1623. doi:10.2307/3871042
Da Ines O, Degroote F, Goubely C, Amiard S, Gallego ME, White CI. 2013. Meiotic
Recombination in Arabidopsis Is Catalysed by DMC1, with RAD51 Playing a Supporting
Role. *PLoS Genet* **9**:e1003787. doi:10.1371/journal.pgen.1003787
Dainat J, Hereñú D, Pucholt P. 2020. NBISweden/AGAT: AGAT-v0.5.1. Zenodo.
doi:10.5281/ZENODO.4205393
Dernburg AF. 2001. Here, there, and everywhere: kinetochore function on holocentric
chromosomes. *J Cell Biol* **153**:F33-38. doi:10.1083/jcb.153.6.f33
Dieterich C, Clifton SW, Schuster LN, Chinwalla A, Delehaunty K, Dinkelacker I, Fulton L, Fulton
R, Godfrey J, Minx P, Mitreva M, Roeseler W, Tian H, Witte H, Yang S-P, Wilson RK,
Sommer RJ. 2008. The Pristionchus pacificus genome provides a unique perspective on
nematode lifestyle and parasitism. *Nat Genet* **40**:1193–1198. doi:10.1038/ng.227
Doyle SR, Laing R, Bartley DJ, Britton C, Chaudhry U, Gilleard JS, Holroyd N, Mable BK,
Maitland K, Morrison AA, Tait A, Tracey A, Berriman M, Devaney E, Cotton JA, Sargison
ND. 2018. A Genome Resequencing-Based Genetic Map Reveals the Recombination
Landscape of an Outbred Parasitic Nematode in the Presence of Polyploidy and
Polyandry. *Genome Biology and Evolution* **10**:396–409. doi:10.1093/gbe/evx269
Emms DM, Kelly S. 2015. OrthoFinder: solving fundamental biases in whole genome
comparisons dramatically improves orthogroup inference accuracy. *Genome Biol*
16:157. doi:10.1186/s13059-015-0721-2

Enguita-Marruedo A, Martín-Ruiz M, García E, Gil-Fernández A, Parra MT, Viera A, Rufas JS,
Page J. 2019. Transition from a meiotic to a somatic-like DNA damage response during
the pachytene stage in mouse meiosis. *PLoS Genet* **15**:e1007439.
doi:10.1371/journal.pgen.1007439
Ferrandiz N, Barroso C, Telecan O, Shao N, Kim H-M, Testori S, Faull P, Cutillas P, Snijders
AP, Colaiácovo MP, Martinez-Perez E. 2018. Spatiotemporal regulation of Aurora B
recruitment ensures release of cohesion during C. elegans oocyte meiosis. *Nat Commun*
9:834. doi:10.1038/s41467-018-03229-5
Foster JM, Grote A, Mattick J, Tracey A, Tsai Y-C, Chung M, Cotton JA, Clark TA, Geber A,
Holroyd N, Korlach J, Li Y, Libro S, Lustigman S, Michalski ML, Paulini M, Rogers MB,
Teigen L, Twaddle A, Welch L, Berriman M, Dunning Hotopp JC, Ghedin E. 2020. Sex
chromosome evolution in parasitic nematodes of humans. *Nat Commun* **11**:1964.
doi:10.1038/s41467-020-15654-6
Garrison E, Marth G. 2012. Haplotype-based variant detection from short-read sequencing.
arXiv:12073907 [q-bio].
Goday C, Pimpinelli S. 1989. Centromere organization in meiotic chromosomes of Parascaris
univalens. *Chromosoma* **98**:160–166. doi:10.1007/BF00329679
Gonzalez de la Rosa PM, Thomson M, Trivedi U, Tracey A, Tandonnet S, Blaxter M. 2021a. A
telomere-to-telomere assembly of *Oscheius tipulae* and the evolution of rhabditid
nematode chromosomes. *G3 Genes|Genomes|Genetics* **11**:1–17.
doi:10.1093/g3journal/jkaa020
Gonzalez de la Rosa PM, Thomson M, Trivedi U, Tracey A, Tandonnet S, Blaxter M. 2021b. A
telomere-to-telomere assembly of *Oscheius tipulae* and the evolution of rhabditid
nematode chromosomes. *G3 Genes|Genomes|Genetics* **11**:1–17.
doi:10.1093/g3journal/jkaa020
Goodenough U, Heitman J. 2014. Origins of Eukaryotic Sexual Reproduction. *Cold Spring
Harbor Perspectives in Biology* **6**:a016154–a016154. doi:10.1101/cshperspect.a016154
Grelon M. 2001. AtSPO11-1 is necessary for efficient meiotic recombination in plants. *The
EMBO Journal* **20**:589–600. doi:10.1093/emboj/20.3.589
Harper NC, Rillo R, Jover-Gil S, Assaf ZJ, Bhalla N, Dernburg AF. 2011. Pairing Centers Recruit
a Polo-like Kinase to Orchestrate Meiotic Chromosome Dynamics in C. elegans.
Developmental Cell **21**:934–947. doi:10.1016/j.devcel.2011.09.001
Hayashi M, Chin GM, Villeneuve AM. 2007. C. elegans Germ Cells Switch between Distinct
Modes of Double-Strand Break Repair During Meiotic Prophase Progression. *PLoS
Genet* **3**:e191. doi:10.1371/journal.pgen.0030191
Hollingsworth NM, Byers B. 1989. HOP1: a yeast meiotic pairing gene. *Genetics* **121**:445–462.
Hollis JA, Glover ML, Schlientz AJ, Cahoon CK, Bowerman B, Wignall SM, Libuda DE. 2020.
Excess crossovers impede faithful meiotic chromosome segregation in C. elegans. *PLoS
Genet* **16**:e1009001. doi:10.1371/journal.pgen.1009001
Hong RL, Sommer RJ. 2006. Pristionchus pacificus: a well-rounded nematode. *Bioessays*
28:651–659. doi:10.1002/bies.20404
Howe KL, Bolt BJ, Shafie M, Kersey P, Berriman M. 2017. WormBase ParaSite – a
comprehensive resource for helminth genomics. *Molecular and Biochemical
Parasitology* **215**:2–10. doi:10.1016/j.molbiopara.2016.11.005
Hunter N. 2015. Meiotic Recombination: The Essence of Heredity. *Cold Spring Harb Perspect
Biol* a016618. doi:10.1101/cshperspect.a016618
Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. 2017. ModelFinder: fast
model selection for accurate phylogenetic estimates. *Nat Methods* **14**:587–589.
doi:10.1038/nmeth.4285

Katoh K, Standley DM. 2013. MAFFT Multiple Sequence Alignment Software Version 7:
Improvements in Performance and Usability. *Molecular Biology and Evolution* **30**:772–
780. doi:10.1093/molbev/mst010

Keeney S, Giroux CN, Kleckner N. 1997. Meiosis-Specific DNA Double-Strand Breaks Are
Catalyzed by Spo11, a Member of a Widely Conserved Protein Family. *Cell* **88**:375–384.
doi:10.1016/S0092-8674(00)81876-0

Kim Y, Kostow N, Dernburg AF. 2015. The Chromosome Axis Mediates Feedback Control of
CHK-2 to Ensure Crossover Formation in C. elegans. *Developmental Cell* **35**:247–261.
doi:10.1016/j.devcel.2015.09.021

Kohl KP, Sekelsky J. 2013. Meiotic and Mitotic Recombination in Meiosis. *Genetics* **194**:327–
334. doi:10.1534/genetics.113.150581

Labella S, Woglar A, Jantsch V, Zetka M. 2011. Polo Kinases Establish Links between Meiotic
Chromosomes and Cytoskeletal Forces Essential for Homolog Pairing. *Developmental*
Cell **21**:948–958. doi:10.1016/j.devcel.2011.07.011

Lake CM, Hawley RS. 2012. The Molecular Control of Meiotic Chromosomal Behavior: Events
in Early Meiotic Prophase in *Drosophila* Oocytes. *Annu Rev Physiol* **74**:425–451.
doi:10.1146/annurev-physiol-020911-153342

Letunic I, Bork P. 2016. Interactive tree of life (iTOL) v3: an online tool for the display and
annotation of phylogenetic and other trees. *Nucleic Acids Res* **44**:W242–W245.
doi:10.1093/nar/gkw290

Lo T-W, Pickle CS, Lin S, Ralston EJ, Gurling M, Schartner CM, Bian Q, Doudna JA, Meyer BJ.
2013. Precise and Heritable Genome Editing in Evolutionarily Diverse Nematodes Using
TALENs and CRISPR/Cas9 to Engineer Insertions and Deletions. *Genetics* **195**:331–
348. doi:10.1534/genetics.113.155382

MacQueen AJ. 2002. Synapsis-dependent and -independent mechanisms stabilize homolog
pairing during meiotic prophase in C. elegans. *Genes & Development* **16**:2428–2442.
doi:10.1101/gad.1011602

MacQueen AJ, Phillips CM, Bhalla N, Weiser P, Villeneuve AM, Dernburg AF. 2005.
Chromosome Sites Play Dual Roles to Establish Homologous Synapsis during Meiosis
in C. elegans. *Cell* **123**:1037–1050. doi:10.1016/j.cell.2005.09.034

Margarido GRA, Souza AP, Garcia AAF. 2007. OneMap: software for genetic mapping in
outcrossing species: OneMap. *Hereditas* **144**:78–79. doi:10.1111/j.2007.0018-
0661.02000.x

Martinez-Perez E, Schvarzstein M, Barroso C, Lightfoot J, Dernburg AF, Villeneuve AM. 2008.
Crossovers trigger a remodeling of meiotic chromosome axis composition that is linked
to two-step loss of sister chromatid cohesion. *Genes & Development* **22**:2886–2901.
doi:10.1101/gad.1694108

Melters DP, Paliulis LV, Korf IF, Chan SWL. 2012. Holocentric chromosomes: convergent
evolution, meiotic adaptations, and genomic analysis. *Chromosome Res* **20**:579–593.
doi:10.1007/s10577-012-9292-1

Nadarajan S, Lambert TJ, Altendorfer E, Gao J, Blower MD, Waters JC, Colaiácovo MP. 2017.
Polo-like kinase-dependent phosphorylation of the synaptonemal complex protein SYP-4
regulates double-strand break formation through a negative feedback loop. *eLife*
6:e23437. doi:10.7554/eLife.23437

Namai S, Sugimoto A. 2018. Transgenesis by microparticle bombardment for live imaging of
fluorescent proteins in Pristionchus pacificus germline and early embryos. *Dev Genes*
Evol **228**:75–82. doi:10.1007/s00427-018-0605-z

Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. 2015. IQ-TREE: A Fast and Effective
Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Molecular Biology*
and Evolution **32**:268–274. doi:10.1093/molbev/msu300

Pattabiraman D, Roelens B, Woglar A, Villeneuve AM. 2017. Meiotic recombination modulates
the structure and dynamics of the synaptonemal complex during *C. elegans* meiosis
(preprint). Genetics. doi:10.1101/110064
Penkner A, Tang L, Novatchkova M, Ladurner M, Fridkin A, Gruenbaum Y, Schweizer D, Loidl
J, Jantsch V. 2007. The Nuclear Envelope Protein Matefin/SUN-1 Is Required for
Homologous Pairing in C. elegans Meiosis. *Developmental Cell* **12**:873–885.
doi:10.1016/j.devcel.2007.05.004
Penkner AM, Fridkin A, Gloggnitzer J, Baudrimont A, Machacek T, Woglar A, Csaszar E,
Pasierbek P, Ammerer G, Gruenbaum Y, Jantsch V. 2009. Meiotic Chromosome
Homology Search Involves Modifications of the Nuclear Envelope Protein Matefin/SUN-
1. *Cell* **139**:920–933. doi:10.1016/j.cell.2009.10.045
Phillips CM, Dernburg AF. 2006. A Family of Zinc-Finger Proteins Is Required for Chromosome-
Specific Pairing and Synapsis during Meiosis in C. elegans. *Developmental Cell* **11**:817–
829. doi:10.1016/j.devcel.2006.09.020
Phillips CM, McDonald KL, Dernburg AF. 2009a. Cytological Analysis of Meiosis in
Caenorhabditis elegans In: Keeney S, editor. Meiosis, Methods in Molecular Biology.
Totowa, NJ: Humana Press. pp. 171–195. doi:10.1007/978-1-60761-103-5_11
Phillips CM, Meng X, Zhang L, Chretien JH, Urnov FD, Dernburg AF. 2009b. Identification of
chromosome sequence motifs that mediate meiotic pairing and synapsis in C. elegans.
Nat Cell Biol **11**:934–942. doi:10.1038/ncb1904
Phillips CM, Wong C, Bhalla N, Carlton PM, Weiser P, Meneely PM, Dernburg AF. 2005. HIM-8
Binds to the X Chromosome Pairing Center and Mediates Chromosome-Specific Meiotic
Synapsis. *Cell* **123**:1051–1063. doi:10.1016/j.cell.2005.09.035
Pittman DL, Cobb J, Schimenti KJ, Wilson LA, Cooper DM, Brignull E, Handel MA, Schimenti
JC. 1998. Meiotic Prophase Arrest with Failure of Chromosome Synapsis in Mice
Deficient for Dmc1 , a Germline-Specific RecA Homolog. *Molecular Cell* **1**:697–705.
doi:10.1016/S1097-2765(00)80069-6
Rockman MV, Kruglyak L. 2009. Recombinational Landscape and Population Genomics of
Caenorhabditis elegans. *PLoS Genet* **5**:e1000419. doi:10.1371/journal.pgen.1000419
Rockmill B, Sym M, Scherthan H, Roeder GS. 1995. Roles for two RecA homologs in promoting
meiotic chromosome synapsis. *Genes & Development* **9**:2684–2695.
doi:10.1101/gad.9.21.2684
Rödelsperger C, Meyer JM, Prabh N, Lanz C, Bemm F, Sommer RJ. 2017. Single-Molecule
Sequencing Reveals the Chromosome-Scale Genomic Architecture of the Nematode
Model Organism Pristionchus pacificus. *Cell Reports* **21**:834–844.
doi:10.1016/j.celrep.2017.09.077
Rog O, Dernburg AF. 2013. Chromosome pairing and synapsis during Caenorhabditis elegans
meiosis. *Current Opinion in Cell Biology* **25**:349–356. doi:10.1016/j.ceb.2013.03.003
Rose AM, McKIM KS. 1992. Meiotic Recombination in Caenorhabditis elegansMechanisms of
Eukaryotic DNA Recombination. Elsevier. pp. 113–124. doi:10.1016/B978-0-12-293445-
2.50014-4
Rudel D, Riebesell M, Sommer RJ. 2005. Gonadogenesis in Pristionchus pacificus and organ
evolution: development, adult morphology and cell–cell interactions in the hermaphrodite
gonad. *Developmental Biology* **277**:200–221. doi:10.1016/j.ydbio.2004.09.021
Sato A, Isaac B, Phillips CM, Rillo R, Carlton PM, Wynne DJ, Kasad RA, Dernburg AF. 2009.
Cytoskeletal Forces Span the Nuclear Envelope to Coordinate Meiotic Chromosome
Pairing and Synapsis. *Cell* **139**:907–919. doi:10.1016/j.cell.2009.10.039
Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S,
Rueden C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K,
Tomancak P, Cardona A. 2012. Fiji: an open-source platform for biological-image
analysis. *Nat Methods* **9**:676–682. doi:10.1038/nmeth.2019

Severson AF, Meyer BJ. 2014. Divergent kleisin subunits of cohesin specify mechanisms to
tether and release meiotic chromosomes. *eLife* **3**:e03467. doi:10.7554/eLife.03467
Shakes DC, Wu J, Sadler PL, LaPrade K, Moore LL, Noritake A, Chu DS. 2009.
Spermatogenesis-Specific Features of the Meiotic Program in Caenorhabditis elegans.
PLoS Genet **5**:e1000611. doi:10.1371/journal.pgen.1000611
Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. 2015. BUSCO:
assessing genome assembly and annotation completeness with single-copy orthologs.
Bioinformatics **31**:3210–3212. doi:10.1093/bioinformatics/btv351
Sommer RJ, editor. 2015. Pristionchus pacificus: a nematode model for comparative and
evolutionary biology, Nematology monographs and perspectives. Leiden ; Boston: Brill.
Sommer RJ, Carta LK, Kim S-Y, Sternberg PW. 1996. Morphological, genetic and molecular
description of Pristionchus pacificus sp. N. (Nematoda: Neodiplogastridae). *Fundam*
appl Nematol **19**:511–521.
Srinivasan J, Sinz W, Jesse T, Wiggers-Perebolte L, Jansen K, Buntjer J, van der Meulen M,
Sommer RJ. 2003. An integrated physical and genetic map of the nematode
Pristionchus pacificus. *Mol Gen Genomics* **269**:715–722. doi:10.1007/s00438-003-0881-
8
Srinivasan J, Sinz W, Lanz C, Brand A, Nandakumar R, Raddatz G, Witte H, Keller H, Kipping I,
Pires-daSilva A, Jesse T, Millare J, de Both M, Schuster SC, Sommer RJ. 2002. A
bacterial artificial chromosome-based genetic linkage map of the nematode Pristionchus
pacificus. *Genetics* **162**:129–134.
Steinfeld JB, Beláň O, Kwon Y, Terakawa T, Al-Zain A, Smith MJ, Crickard JB, Qi Z, Zhao W,
Rothstein R, Symington LS, Sung P, Boulton SJ, Greene EC. 2019. Defining the
influence of Rad51 and Dmc1 lineage-specific amino acids on genetic recombination.
Genes Dev **33**:1191–1207. doi:10.1101/gad.328062.119
Tandonnet S, Koutsovoulos GD, Adams S, Cloarec D, Parihar M, Blaxter ML, Pires-daSilva A.
2019. Chromosome-Wide Evolution and Sex Determination in the Three-Sexed
Nematode <i/> *Auanema rhodensis*. *G3* g3.0011.2019. doi:10.1534/g3.119.0011
Tzur YB, Egydio de Carvalho C, Nadarajan S, Van Bostelen I, Gu Y, Chu DS, Cheeseman IM,
Colaiácovo MP. 2012. LAB-1 Targets PP1 and Restricts Aurora B Kinase upon Entrance
into Meiosis to Promote Sister Chromatid Cohesion. *PLoS Biol* **10**:e1001378.
doi:10.1371/journal.pbio.1001378
Vader G, Musacchio A. 2014. HORMA Domains at the Heart of Meiotic Chromosome
Dynamics. *Developmental Cell* **31**:389–391. doi:10.1016/j.devcel.2014.11.009
Vasimuddin Md, Misra S, Li H, Aluru S. 2019. Efficient Architecture-Aware Acceleration of BWA-
MEM for Multicore Systems2019 IEEE International Parallel and Distributed Processing
Symposium (IPDPS). Presented at the 2019 IEEE International Parallel and Distributed
Processing Symposium (IPDPS). Rio de Janeiro, Brazil: IEEE. pp. 314–324.
doi:10.1109/IPDPS.2019.00041
Villeneuve AM, Hillers KJ. 2001. Whence meiosis? *Cell* **106**:647–650. doi:10.1016/s0092-
8674(01)00500-1
Watanabe Y, Kitajima TS. 2005. Shugoshin protects cohesin complexes at centromeres. *Phil*
Trans R Soc B **360**:515–521. doi:10.1098/rstb.2004.1607
Werner MS, Sieriebriennikov B, Prabh N, Loschko T, Lanz C, Sommer RJ. 2018. Young genes
have distinct gene structure, epigenetic profiles, and transcriptional regulation. *Genome*
Res **28**:1675–1687. doi:10.1101/gr.234872.118
Witte H, Moreno E, Rödelsperger C, Kim J, Kim J-S, Streit A, Sommer RJ. 2015. Gene
inactivation using the CRISPR/Cas9 system in the nematode Pristionchus pacificus. *Dev*
Genes Evol **225**:55–62. doi:10.1007/s00427-014-0486-8

Yoshida K, Kondoh G, Matsuda Y, Habu T, Nishimune Y, Morita T. 1998. The Mouse RecA -like
Gene Dmc1 Is Required for Homologous Chromosome Synapsis during Meiosis.
Molecular Cell **1**:707–718. doi:10.1016/S1097-2765(00)80070-2
Yu Z, Kim Y, Dernburg AF. 2016. Meiotic recombination and the crossover assurance
checkpoint in Caenorhabditis elegans. *Seminars in Cell & Developmental Biology*
54:106–116. doi:10.1016/j.semcdb.2016.03.014
Zhang C, Rabiee M, Sayyari E, Mirarab S. 2018. ASTRAL-III: polynomial time species tree
reconstruction from partially resolved gene trees. *BMC Bioinformatics* **19**:153.
doi:10.1186/s12859-018-2129-y
Zickler D, Kleckner N. 2015. Recombination, Pairing, and Synapsis of Homologs during Meiosis.
Cold Spring Harb Perspect Biol **7**:a016626. doi:10.1101/cshperspect.a016626
Zickler D, Kleckner N. 1999. Meiotic Chromosomes: Integrating Structure and Function. *Annu*
Rev Genet **33**:603–754. doi:10.1146/annurev.genet.33.1.603
1358

Figure 1

A

DAPI

PM TZ Pach Dip Diff Diakinesis

B

DAPI

Cy3-dUTP

DAPI Cy3-dUTP

C CENPC::V5 DAPI TUBULIN merge

D CENPC::V5 DAPI merge

premeiotic germline

embryo

Figure 2

Figure 2 - figure supplement 1

Figure 2- figure supplement 2

A

B

Figure 3

Figure 3- figure supplement 1

Genotype	% Egg viability (±SD)	% Male progeny (±SD)	Eggs laid (± SD)
WT (n=30)	92.9 (±16.3)	0.9 (±1.0)	205 (±55)
syp-4::HA (n=7)	97.7 (±11.0)	0.5 (±0.6)	164 (±15)
dmc-1::V5 (n=8)	95.3 (±4.7)	1.1 (±1.4)	170 (±14)
rad-51::V5 (n=8)	107.1 (±18.9)	1.1 (±1.5)	157(±15)
syp-4::HA; dmc-1::V5; cosa-1::3xFLAG (n=8)	87.0 (±10.6)	0.4 (±0.6)	170 (±37)

Figure 4

A

DAPI   DMC-1::V5   RAD-51

TZ ⟶ Pachytene

B

DAPI | DMC-1::V5 | RAD-51 | merge

TZ

Pach

C

RAD-51
DMC-1::V5

D

DAPI
DMC-1::V5

Figure 4- figure supplement 1

Figure 5

A

B

Number of DAPI-staining bodies at diakinesis

Legend:
- WT (n= 39)
- spo-11 (n=15)
- dmc-1 (n=11)
- rad-51 (n=18)

C

% germlines without diakinesis nuclei

D

Genotype	% Egg viability (±SD)	% Male progeny (±SD)	Eggs laid (± SD)
WT (n=30)	92.9 (±16.3)	0.9 (±1.0)	205 (±55)
spo-11 (n=5)	0.3 (± .4)	20 (±44.7)	125 (±40)
dmc-1 (n=7)	0 (±0)	N/A	12 (±12)
rad-51 (n=14)	15.0 (±17.9)	4.5 (±5.1)	121 (±70)

Figure 6

Figure 6- figure supplement 1

A

Figure 6- figure supplement 2

Figure 7

A

Figure 8

A

P0: CA/CA ♂ × WA/WA ♀   CA/CA ♂ × LR/LR ♀
    (PS312)   (PS1843)   (PS312)   (RSB001)

F1: CA/WA ♂   ×   CA/LR ⚥

F2: ⚥

Isolate and self

Extract genomic DNA
and sequence

B

CA/LR

CA/CA   WA/LR

WA/CA

● ♂ Spermatocyte
● ⚥ Oocyte

Figure 8 - figure supplement 1

Table S1

Strain	El Paco genome reference, v1, 2017	Wormbase designation	Allele	Guide RNA target sequence (PAM sequence underlined)	ssDNA template	Genotyping primer sequences
cenpc::V5	*UMM-S71-6.7mRNA-1*	*PPA37734*	*ie1007*	ATGAAGAGATGGATTATAGT<u>A</u> <u>GG</u>	GGATATTAAAAAAAGGGGGTAAAATTGTACATCG CAAAAGGATGAGGAGTCATTTAATTCATCTTACTA TAATCCATCTCTTCAGGTGGAGTCGAGTCCAAGA AGTGGGTTTGGAATTGGCTTTCCAGATCCGGCT CCTTCATTCTCTTCATCGAACTCTTCCACTTCCTT CTCCTTCTGCTCTTTTCTCTTCTTGTT	(f)TTTCTCCAGGAGTGGTTATCG (r)ACATCGCAAAAGGATGAGGAG
cosa-1::3xflag	*UMM-S57-3.22mRNA-1*	*PPA23791*	*ie1003*	CTTTATTCTTCATTTTACAG<u>TG</u> <u>G</u>	GATCATCCCAGGGAGAGAACGACCTACCCAAAC AGAGATCATAGATCTAATTATCCACTGGGAGCCG GATCTGATTATAAAGACCATGATGGAGACTATAAG GATCACGATATTGATTACAAAGACGATGATGATAA ATAAAATGAAGAATAAAGAGTATTAAATTTATGTTT GTGTTCGTTTTTGTAATTACTGCTTTG	(f)ACGACCTACCCAAACAGAGA (r)CGGATGTGGAAAGACGTACC
dmc-1 mutant	*UMM-S442-1.74mRNA-1*	—	*ie1005*	TTCGATAAGCTGCTTGGAGG<u>T</u> <u>GG</u>	ATTTAATTGTAACATTCAGACATCCAGTTGATCAA CTAACCTCCGTTATTGCCTGACTTTCGATTCCAC CactagtgTCCAAGCAGCTTATCGAATTCAACACTT CCAGTGGAGATTTTAAAGACTTGCTTGCGTCGT GAACACACT	(f)GGACTCTCGGAGGCTAAAGT (r)ATTCTCGAGCATTGCTTCCT
dmc-1::V5	*UMM-S442-1.74mRNA-1*	—	*ie1001*	GCAACGTTTGCCATTGCAGC <u>AGG</u>	GATAAAACATGATCTTTTGTCCTTCATAATTGATTA GGTGGAGTCGAGTCCAAGAAGTGGGTTTGGAAT TGGCTTTCCATCCTTTGCATCGACAATTCCTCCT GC**G**GC**G**ATGGCAAACGTTGCTTCGTTCTCAGGC ATATC	(f)TGCCTGAGAACGAAGCAACG (r)ACATGAGATGGCACAAAGGAC
hop-1	*UMM-S341-6.31mRNA-1*	*PPA10281*				
rad-51 mutant	*UMM-S442-1.74mRNA-1*	*PPA42255*	*ie1006*	GTCGAGAACGAGGAGAATGC <u>CGG</u>	ATTTATTGGTTACCTCAAGGGACATGATGGACTG GCAGGCGAGTCCGGCATACTAGTTTCTCCTCGT TCTCGACATCAGCGTCGACGTGCGCCATTTGAG CGGACAT	(f)TTCTAGTGACGCGTGTTGTT (r)ACGAATCCTCGTTGCTGAAG

rad-51::V5	*UMM-S442-1.74mRNA-1*	*PPA42255*	*ie1008*	GACCTTCTCCATCACAACAA<u>CGG</u>	GATGGTCAGATAACAGCTATGGAGTGTGTAAATTGGAGAGAGTTAGACGACTAGGTGGAGTCGAGTCCAAGAAGTGGGTTTGGAATTGGCTTTCCATTATCCTTTCCATCTTCGATGCCCGTGGTGGTGATGGAGAAGGTCGCCTCGCCCTCGGCCAGGCATGGAGATTG	(f)GATTCATCTCTGTTTCACAGG (r)GATCTCGGTTTCTCTCTTGGAAA
spo-11 mutant	*UMM-S230-10.9mRNA-1*	*PPA33054*	*ie1004*	ATTCAGAACTTGGCAGAGAT<u>CGG</u>	ACAAACATCTTTTTGCACGACAGGATTCTCTCAATCGATCAGTTTCCGATAACTAGTACTCTGCCAAGTTCTGAATATGCAAAGATGTCAAGTCAATGTGGTAAGT	(f)GGAAATCCTTCGTTCTCACTATGG (r)GTCTCAATATCAGACAATTTCATTCCG
syp-4::HA	*UMM-S245-8.16mRNA-1*	—	*ie1002*	GGAGGAGAATTCAACTTCTT<u>CGG</u>	CAAATGGAGGTGGCGGCGGGGGAGGAGGAGAGTTTAATTTTTTTGGTTTTTACCCCTACGATGTCCCAGATTATGCTTAAACCAATTTTTTCGAGCTTGGTGTAATGTATCCA	(f)CCCGTTGATGATGCTACCAG (r)GATACATTACACCAAGCTCGAA

Table S2

Species	Accession	Version	Source
Acanthocheilonema viteae	PRJEB1697	WBPS15	WormBase ParaSite
Acrobeloides nanus	PRJEB26554	WBPS15	WormBase ParaSite
Ancylostoma ceylanicum	PRJNA231479	WBPS15	WormBase ParaSite
Ancylostoma duodenale	PRJNA72581	WBPS15	WormBase ParaSite
Angiostrongylus cantonensis	PRJNA350391	WBPS15	WormBase ParaSite
Angiostrongylus costaricensis	PRJEB494	WBPS15	WormBase ParaSite
Anisakis simplex	PRJEB496	WBPS15	WormBase ParaSite
Ascaris suum	PRJNA62057	WBPS15	WormBase ParaSite
Brugia malayi	PRJNA10729	WBPS15	WormBase ParaSite
Brugia pahangi	PRJEB497	WBPS15	WormBase ParaSite
Bursaphelenchus xylophilus	PRJEA64437	WBPS15	WormBase ParaSite
Caenorhabditis elegans	PRJNA13758	WBPS15	WormBase ParaSite
Caenorhabditis parvicauda	PRJEB12595	WBPS15	WormBase ParaSite
Cylicostephanus goldi	PRJEB498	WBPS15	WormBase ParaSite
Dictyocaulus viviparus	PRJNA72587	WBPS15	WormBase ParaSite
Diploscapter coronatus	PRJDB3143	WBPS15	WormBase ParaSite
Diploscapter pachys	PRJNA280107	WBPS15	WormBase ParaSite
Dirofilaria immitis	PRJEB1797	WBPS15	WormBase ParaSite
Ditylenchus destructor	PRJNA312427	WBPS15	WormBase ParaSite
Ditylenchus dipsaci	PRJNA498219	WBPS15	WormBase ParaSite
Dracunculus medinensis	PRJEB500	WBPS15	WormBase ParaSite
Enterobius vermicularis	PRJEB503	WBPS15	WormBase ParaSite
Globodera pallida	PRJEB123	WBPS15	WormBase ParaSite
Gongylonema pulchrum	PRJEB505	WBPS15	WormBase ParaSite
Haemonchus contortus	PRJEB506	WBPS15	WormBase ParaSite
Haemonchus placei	PRJEB509	WBPS15	WormBase ParaSite
Halicephalobus mephisto	PRJNA528747	WBPS15	WormBase ParaSite
Heligmosomoides polygyrus	PRJEB15396	WBPS15	WormBase ParaSite
Heterorhabditis bacteriophora	PRJNA13977	WBPS15	WormBase ParaSite
Hypsibius dujardini	-	3.1.5	tardigrades.org
Litomosoides sigmodontis	PRJEB3075	WBPS15	WormBase ParaSite
Loa loa	PRJNA246086	WBPS15	WormBase ParaSite
Meloidogyne incognita	PRJNA340324	WBPS15	WormBase ParaSite
Meloidogyne javanica	PRJNA340324	WBPS15	WormBase ParaSite
Mesorhabditis belari	PRJEB30104	WBPS15	WormBase ParaSite
Micoletzkya japonica	PRJEB27334	WBPS15	WormBase ParaSite
Necator americanus	PRJNA72135	WBPS15	WormBase ParaSite
Nippostrongylus brasiliensis	PRJEB511	WBPS15	WormBase ParaSite
Oesophagostomum dentatum	PRJNA72579	WBPS15	WormBase ParaSite
Onchocerca ochengi	PRJEB1204	WBPS15	WormBase ParaSite
Onchocerca volvulus	PRJEB513	WBPS15	WormBase ParaSite
Oscheius tipulae	PRJEB15512	WBPS15	WormBase ParaSite

Species	Accession	Version	Source
Panagrellus redivivus	PRJNA186477	WBPS15	WormBase ParaSite
Panagrolaimus davidi	PRJEB32708	WBPS15	WormBase ParaSite
Panagrolaimus superbus	PRJEB32708	WBPS15	WormBase ParaSite
Parapristionchus giblindavisi	PRJEB27334	WBPS15	WormBase ParaSite
Parascaris univalens	PRJNA386823	WBPS15	WormBase ParaSite
Parastrongyloides trichosuri	PRJEB515	WBPS15	WormBase ParaSite
Plectus sambesii	PRJNA390260	WBPS15	WormBase ParaSite
Pristionchus entomophagus	PRJEB27334	WBPS15	WormBase ParaSite
Pristionchus exspectatus	PRJEB24288	WBPS15	WormBase ParaSite
Ramazzottius varieornatus	-	Rv101	tardigrades.org
Rhabditophanes kr3021	PRJEB1297	WBPS15	WormBase ParaSite
Romanomermis culicivorax	PRJEB1358	WBPS15	WormBase ParaSite
Setaria digitata	PRJNA479729	WBPS15	WormBase ParaSite
Soboliphyme baturini	PRJEB516	WBPS15	WormBase ParaSite
Steinernema carpocapsae	PRJNA202318	WBPS15	WormBase ParaSite
Steinernema feltiae	PRJNA204661	WBPS15	WormBase ParaSite
Strongyloides ratti	PRJEB125	WBPS15	WormBase ParaSite
Strongyloides stercoralis	PRJEB528	WBPS15	WormBase ParaSite
Strongylus vulgaris	PRJEB531	WBPS15	WormBase ParaSite
Thelazia callipaeda	PRJEB1205	WBPS15	WormBase ParaSite
Toxocara canis	PRJNA248777	WBPS15	WormBase ParaSite
Trichinella pseudospiralis	PRJNA257433	WBPS15	WormBase ParaSite
Trichinella spiralis	PRJNA12603	WBPS15	WormBase ParaSite
Trichuris muris	PRJEB126	WBPS15	WormBase ParaSite
Trichuris suis	PRJNA179528	WBPS15	WormBase ParaSite
Wuchereria bancrofti	PRJNA275548	WBPS15	WormBase ParaSite

Table S3

SSLP marker	Chromosome	Primer sequences
L2	I	(f)GGAATGAGGGTCGTGAGGTA (r)GCGTTTTCAAAATGAAGCAA
L23	III	(f)AATTCGCAATGACCTATGAGA (r)ACAAATTCGTCACTCTGGCA
L109	III	(f)GTCCACCGTACCTCTCACCT (r)TCTTGATCCATCGGAATGCT
L100	IV	(f)CGATCGCCGTTTACGAGG (r)GTCCCGCATAGCAACCAATA
L106	IV	(f)GAAAGGGACTGCGGAACTC (r)GTACAGCGTACCGTCCGATT
L16	V	(f)TGAACACTTTCAATGCCTCTTT (r)GTCGATCTTGATCTTGTCCG

Table S4

Software	Version	Relevant parameters
catfasta2phyml	-	-c -f
edgeR	3.18.0	
HTSeq	0.6.1	
IQ-TREE	1.6.10	-bb 1000, -bb -m GTR20+G
MAFFT	7.407	--auto
NCBI-BLAST+	2.5.0+	blastp
OrthoFinder	2.2.7	-og
PhyloTreePruner	20150918	35 0.9 u
Priism	4.7.1	
R	3.5.1	
softWorx	7.0.0	
STAR	2.5.2	--outFilterType BySJout --outFilterMultimapNmax 20 --alignSJoverhangMin 8 -alignSJDBoverhangMin 8 --outFilterMismatchNmax 999 --outFilterMismatchNoverReadLmax 0.04 --alignIntronMin 20 --alignIntronMax 1000000 --alignMatesGapMax 1000000
StringTie	1.3.3	-p 10 -m 50 --rf -f 0.1 -a 10 -j 5 -c 2.5
Transdecorder	3.0.1	
trimalAI	v1.4.rev15	-gt 0.8 -st 0.001 -resoverlap 0.75 -seqoverlap 80