Analysis of meiosis in *Pristionchus pacificus* reveals plasticity in homolog pairing and synapsis in the nematode lineage

Regina Rillo-Bohn[1,2,3,4], Renzo Adilardi[1,2,3,4], Barış Avşaroğlu[1,2,3,4], Lewis Stevens[5], Simone Köhler[1,2,3,4,6], Joshua Bayes[1,2,3,4], Clara Wang[1,2,3,4], Sabrina Lin[1,2,3,4], Kayla Baskevitch[1,2,3,4], Abby F. Dernburg[1,2,3,4]

[1]Department of Molecular and Cell Biology, University of California, Berkeley, Berkeley, CA United States; [2]Howard Hughes Medical Institute, Chevy Chase, MD United States; [3]Biological Systems and Engineering Division, Lawrence Berkeley National Laboratory, Berkeley, CA United States; [4]California Institute for Quantitative Biosciences, Berkeley, CA United States; [5]Institute of Evolutionary Biology, University of Edinburgh, Edinburgh, UK; [6]Cell Biology and Biophysics Unit, European Molecular Biology Laboratory, Heidelberg, Germany

ABSTRACT

Meiosis gives rise to haploid gametes from diploid progenitor cells. Here we describe a new,

comparative model system for molecular analysis of meiosis, the nematode *Pristionchus*

pacificus, a distant relative of the widely studied model organism *Caenorhabditis elegans*.

Despite superficial similarities in germline organization and meiotic progression between *P.*

pacificus and *C. elegans*, we identify fundamental differences in the molecular mechanisms

underlying homolog pairing, synapsis, and crossover regulation. Whereas *C. elegans* has lost

the meiosis-specific recombinase Dmc1, *P. pacificus* expresses both DMC-1 and RAD-51, which

localize sequentially to meiotic chromosomes during prophase. *Ppa-spo-11* and *Ppa-dmc-1* are

required for stable homolog pairing, synapsis, and crossover formation, while Ppa-rad-51 is

dispensable for these key processes during early prophase. Additionally, we show that

elevated crossover recombination in P. pacificus likely arises through a Class II pathway

normally inactive in *C. elegans*, shedding light on crossover control and the evolution of

recombination rates.

INTRODUCTION

All sexually reproducing organisms rely on the specialized cell division process of

meiosis to generate haploid gametes from diploid precursors. Upon fertilization, haploid

gametes fuse and restore the diploid chromosome complement in the zygote. Thus, meiosis is

essential for the survival of sexually reproducing species. This process was likely present in the

last eukaryotic common ancestor (LECA). Studies in plants, animals, fungi, and protists have

revealed intriguing diversity in the details of meiotic mechanisms.

A defining feature of meiosis is a "reductional" segregation in which homologous

chromosomes are separated, usually during the first of two nuclear divisions. A highly

choreographed series of chromosome transactions precedes this division and ensures faithful

homolog segregation: (1) pairing, in which chromosomes contact and recognize their

homologous partners; (2) synapsis, defined as the assembly of a protein ensemble called the

synaptonemal complex (SC) between homologs, which leads to their lengthwise alignment;

and (3) crossover (CO) recombination, which creates physical linkages between chromosomes

that promote proper bi-orientation during anaphase I. Failure to form at least a single CO

between paired homologs results in nondisjunction and aneuploid gametes (Zickler and

Kleckner, 1999).

Although homolog pairing, synapsis and CO recombination during meiosis are nearly

ubiquitous among eukaryotes, key aspects of these chromosomal transactions show

remarkable diversity among different lineages. In most model fungi, plants, and animals,

stable homologous chromosome pairing and synapsis depend on early steps of the

recombination pathway. Meiotic recombination is initiated by a conserved topoisomerase-like

enzyme called Spo11, which catalyzes programmed DNA double-strand breaks (DSBs) across

the genome, a subset of which are ultimately processed into COs (Keeney, 2008; Keeney et al.,

1997). DSBs are resected to form 3' single-stranded DNA overhangs. Dmc1, a meiosis-specific

recombinase, forms filaments along the resulting ssDNA segments and promotes

interhomolog strand invasion. Rad51 is often required as a cofactor for these eary steps in

recombination. Spo11-dependent induction of DSBs and Dmc1-dependent strand invasion are

crucial for stable homolog pairing and synapsis in the budding yeast *Saccharomyces cerevisiae*,

the flowering plant *Arabidopsis thaliana*, and mammalian model *Mus musculus* (Bishop et al.,

1992; Couteau et al., 1999; Grelon, 2001; Pittman et al., 1998; Rockmill et al., 1995; Yoshida et

al., 1998).

In contrast, recombination-independent mechanisms of pairing and synapsis have been

characterized in other prominent model systems, including the dipteran *Drosophila*

melanogaster and the nematode *Caenorhabditis elegans*. While recombination is essential for

successful execution of meiosis in *C. elegans* and in female fruit flies, homolog pairing and

synapsis can be uncoupled from early recombination events. In the female germline of *D.*

melanogaster, homolog pairing initiates in proliferating germline stem cells even before they

enter meiosis (Christophorou et al., 2013; Joyce et al., 2013; Rubin et al., 2016) and is stabilized

by SC formation during early prophase. *D. melanogaster* males lack recombination and SCs,

and have apparently evolved a distinct mechanism to stabilize homolog pairing and enable

reductional segregation (McKee et al., 2012). In *C. elegans*, pairing and synapsis are driven by

Pairing Centers, specialized sites on each chromosome bound by a family of zinc-finger

proteins that mediate nuclear envelope attachment and chromosome dynamics (MacQueen et

al., 2005; Phillips et al., 2005; Phillips and Dernburg, 2006; Sato et al., 2009). While nuclear

envelope attachment and chromosome movement play important roles in meiotic pairing and

synapsis across eukaryotes, in *C. elegans* they have acquired a critical role in coupling homolog

pairing to synapsis initiation (Penkner et al., 2009; Sato et al., 2009).

To investigate how the meiotic program is modified during evolution, we have

established tools to investigate meiosis in the free-living nematode *Pristionchus pacificus.* Like

its distant relative *C. elegans*, *P. pacificus* is an androdioecious species, characterized by a

population of mostly self-fertilizing hermaphrodites (XX) and a low frequency of males (XO)

(Sommer et al., 1996). Like *C. elegans, P. pacificus* has a short life cycle of 3.5 days, produces

large broods of about 200 progeny by self-fertilization, and is easily cultured in the lab (Hong

and Sommer, 2006). Although *C. elegans* and *P. pacificus* diverged an estimated 200-300

million years ago (Pires-daSilva, 2004), they share the same number of chromosomes (2n=12)

and, with the exception of one major chromosomal translocation, macrosynteny is maintained

between the two species (Dieterich et al., 2008; Rödelsperger et al., 2017). *P. pacificus* has been

established as a model for comparative studies in development, evolution and ecology

(Sommer, 2015). Recent improvements in the genome assembly (Rödelsperger et al., 2017)

and advances in genome editing (Lo et al., 2013; Namai and Sugimoto, 2018; Witte et al., 2015)

have facilitated investigation of cell biological processes at a more mechanistic level.

In addition to these general features that make *P. pacificus* a tractable model system,

previous studies have suggested interesting differences from *C. elegans.* First, genome

sequencing revealed the presence of the Dmc1 gene, which is absent from the entire

Caenorhabditis clade (Figure 2 - Supplement 1 and Dieterich et al., 2008). Loss of Dmc1

correlates with the adaptation of recombination-independent mechanisms for pairing and

synapsis in *Drosophila* and *Caenorhabditis* (Villeneuve and Hillers, 2001). Therefore, it was of

great interest to us to examine how homologous chromosomes pair and synapse in the

presence of Dmc1. Second, genetic linkage maps have revealed that multiple COs typically

occur per chromosome pair during meiosis in *P. pacificus* (Srinivasan et al., 2003, 2002). This is

in striking contrast to *C. elegans,* which exhibits complete CO interference; only a single Class I

CO normally occurs per chromosome pair (Martinez-Perez and Colaiácovo, 2009). These

observations suggest a major difference in the mechanism of CO control between these two

species.

By employing CRISPR/Cas9-mediated genome editing techniques, genetics,

immunocytochemistry and microscopy, we describe here the early events of meiotic prophase

in *P. pacificus*. We show that homolog pairing, synapsis and CO recombination are dependent

on *Ppa-spo-11* and *Ppa-dmc-1*, while *Ppa-rad-51* is not essential for meiosis. We also provide

evidence that a single Class I CO occurs per homolog pair in *P. pacificus*, implying that the

higher recombination rate is due to Class II COs. Our work establishes tools for future

investigation and highlights the flexibility of the meiotic program within the nematode lineage.

RESULTS

P. pacificus **as a comparative model system for meiosis**

The morphology and organization of the *P. pacificus* germline is very similar to that of

C. elegans. Hermaphrodites have two gonad arms in which sperm and ova are produced

sequentially, while males have a single arm (Rudel et al., 2005). The switch from

spermatogenesis to oogenesis occurs during early adulthood. DAPI staining of adult

hermaphrodite gonads reveals a cylindrical monolayer of cells; the distal tip is populated by

proliferating germline stem cells (Figure 1A-C). As germ cells enter meiosis, their nuclear

morphology abruptly becomes asymmetrical, as the chromosome mass adopts a conspicuous,

crescent-shaped morphology (Figure 1A). Immediately proximal to this "transition zone," DAPI

staining reveals parallel tracks, indicative of paired and synapsed homologous chromosomes at

the pachytene stage. Oocyte chromosomes in *P. pacificus* undergo a dramatic decondensation

between diplotene and diakinesis, a stage that has been referred to as the "growth zone"

(Rudel et al., 2005) or "diffuse stage" (Zickler and Kleckner, 1999). Similar diffuse chromosome

morphology is observed during late prophase in other eukaryotes, but is rarely seen in *C.*

elegans. As oocytes mature in the proximal region of the hermaphrodite gonad, they form a

single row of large cells, and chromosomes condense dramatically as the nuclei grow in size.

Six bivalents can be detected as compact DAPI-staining bodies during diakinesis, the stage

preceding the first meiotic division, during oogenesis (Figure 1A, Sommer et al., 1996, Rudel et

al., 2005).

It has been assumed that chromosomes in *P. pacificus* are holocentric, as in *C. elegans*,

but we are unaware of direct evidence to support this idea. We thus identified a gene encoding

CENP-C (HCP-4 in *C. elegans*), a conserved kinetochore protein, in the *P. pacificus* genome and

inserted a V5 epitope tag at its 3' end using CRISPR/Cas9. The appearance of mitotic

chromosomes in embryos and in the premeiotic germline confirmed their holocentric

organization (Figure 1C). Ppa-CENP-C::V5 forms linear structures, a hallmark of holocentric

chromosomes, instead of discrete foci as observed for monocentric chromosomes.

Stable homolog pairing requires early recombination factors

BLAST searches of the *P. pacificus* genome revealed an open reading frame encoding

an unambiguous ortholog of Dmc1, a meiosis-specific paralog of Rad51 (Table S1). An ortholog

of the Dmc1 cofactor Mnd1 was also readily identified; although its partner Hop2 was not

apparent among the predicted proteins or published nucleotide sequence (Figure 2 –

Supplement 1). By contrast, Dmc1/Mnd1/Hop2 are absent from both *C. elegans* and *D.*

melanogaster, two model organisms that have evolved recombination-independent

mechanisms of homolog pairing and synapsis (Villeneuve and Hillers, 2001). We analyzed the

genome sequences of other nematodes to determine the evolutionary history of these genes

within the nematode lineage (Bolt et al., 2018; https://parasite.wormbase.org). This analysis

revealed that Dmc1/Mnd1/Hop2 have been lost several times during the evolution of

nematodes, including the entire *Caenorhabditis* genus and all sequenced members of Clade IV

(Figure 2 –Supplement 1). Not surprisingly in light of its essential function in DNA repair, the

recombinase Rad51 was detected in almost all genomes with the exception of *Haemonchus*

contortus, *Parascaris univalens* and *Parascaris equorum* (data not shown). It is likely that the

absence of Rad51 orthologs reflects incomplete genome assembly and/or annotation in these

species.

Our analysis also identified homologs of four related zinc finger proteins required for

pairing center activity in *C. elegans* (HIM-8, ZIM-1, ZIM-2, and ZIM-3) in most of the genome

sequences from Clade V. Thus, many species within this clade have genes encoding both

pairing center proteins and Dmc1/Mnd1/Hop2, but *Caenorhabditids* have lost the latter while

Pristionchus apparently diverged before the emergence of the former.

We tested whether homolog pairing in *P. pacificus* is dependent on these early

recombination factors. To generate *spo-11*, *dmc-1* and *rad-51* null mutants, we employed

TALEN-mediated gene disruption, and later CRISPR/Cas9 genome editing techniques (this

study; Lo et al., 2013; Witte et al., 2015). Methods such as co-CRISPR that facilitate detection

of genome editing events in *C. elegans* have not been helpful for *P. pacificus* in our hands or

others' (Witte et al., 2015, and data not shown). Nevertheless, we were able to isolate mutant

alleles by screening a large number of F1 progeny from injected hermaphrodites (see Materials

and Methods). Independent alleles isolated from either TALEN or CRISPR-mediated genome

editing resulted in identical mutant phenotypes. All data presented here were based on alleles

generated by CRISPR/Cas9. Because balancer chromosomes are not currently available for *P.*

pacificus, most mutations described here were maintained in unbalanced heterozygotes, which

were identified by PCR-based genotyping. Self-fertilization of heterozygotes results in broods

with 25% homozygous mutant animals.

As expected, disruption of either *spo-11* or *dmc-1* resulted in the detection of 12 DAPI-

staining univalent chromosomes at diakinesis, indicative of a failure in CO recombination

(Figure 5). Surprisingly, *rad-51* mutants showed only mild meiotic defects (see below), and we

thus validated the loss of RAD-51 function in mutant animals by generating multiple alleles

that showed indistinguishable phenotypes, and also confirmed the absence of RAD-51 protein

by immunofluorescence with a polyclonal antibody against Ppa-RAD-51 (Figure 4 –

Supplement 1A, see Materials and Methods). Mutations in *spo-11* or *dmc-1,* but not *rad-51,* also

resulted in an obvious extension of the region of the germline displaying the crescent-shaped

nuclear morphology characteristic of early meiosis (Figure 2 – Supplement 2). A similar

"extended transition zone" phenotype is seen in *C. elegans* mutants that fail to synapse their

chromosomes during meiosis, suggesting that *spo-11* and *dmc-1* might be required for synapsis

in *P. pacificus*.

To visualize and quantify homolog pairing, we generated FISH probes against two short

tandem repeats found on Chromosome X and IV (Figure 2A). We measured the distance

between pairs of homologous FISH signals in individual nuclei for each genotype. To analyze

pairing kinetics, we divided the distal gonads into five zones of equal length. In zone 1, which

mostly consists of proliferating germ cells, pairs of FISH signals remained far apart, with an

average distance of 2.4 ± 1.0 µm (SD) and 2.5 ± 0.8 µm for Chrom X and IV, respectively (Figure

2B, D). In zone 2, which spans the transition zone, the average distances decreased

significantly in wild-type animals (1.2 ± 1.1 µm and 1.1 ± 1.1 µm for Chrom X and IV,

respectively), and homologous signals remained closely associated in the subsequent meiotic

stages. Notably, homologous FISH probes localized closer together, on average, in the two

zones immediately following meiotic entry, and showed some separation in zones 4 and 5

(Figure 2B, D). By contrast, in wild-type *C. elegans*, homologous loci remain closely apposed in

most of the distal region of the gonad (MacQueen et al., 2002). Together with analysis of

synapsis (below), this indicated that desynapsis initiates shortly after completion of synapsis in

P. pacificus, resulting in partial separation of homologs.

We noted that the average distances between pairs of homologous FISH signals in *spo-*

11 and *dmc-1* mutants also decreased markedly upon meiotic entry, although clearly less so

than in wild type (Figure 2D). In contrast, *rad-51* mutants showed distributions of probe

distances similar to wild-type animals (Figure 2C, D). We considered the possibility that the

proximity between FISH signals might reflect the clustering of all chromosomes within a

subregion of the nucleus, which is apparent during the "transition zone" (leptotene/zygotene),

rather than specific homologous interactions. If so, the extended transition zone morphology

in *spo-11* and *dmc-1* might obscure a pairing defect that would be more apparent in the

absence of clustering (Figure 2 – Supplement 2). To address this, we measured the distances

between pairs of heterologous FISH signals in the premeiotic region (dispersed) versus the

transition zone (clustered). We observed that heterologous FISH signals were also significantly

closer to each other in the transition zone compared to premeiotic nuclei in both wild-type and

mutant animals (Figure 2E). Furthermore, the distances between heterologous versus

homologous pairs of FISH loci were not significantly different in *spo-11* and *dmc-1* mutants

(p=0.1777 and p=0.6774, respectively, by Student's *t*-test), but were clearly different in wild-

type meiocytes (p<0.0001 by Student's *t*-test) (Figure 2E). These data support the idea that

clustering, rather than specific pairing, promote proximity between both homologous and

heterologous loci during leptotene/zygotene in *spo-11* and *dmc-1* mutants. Although we

cannot conclude that transient homologous pairing is absent in these mutants, it is evident

that these early recombination factors are required for stable pairing and extended association

of homologous loci throughout prophase. In contrast, *rad-51* is dispensable for homolog

pairing, as in *C. elegans*.

SPO-11 and DMC-1 are required for homologous synapsis

To further investigate meiotic progression in *P. pacificus* and to probe the role of early

recombination factors in synapsis, we developed cytological markers for the chromosome axis,

which normally assembles upon meiotic entry, and the synaptonemal complex (SC), which

assembles between paired axes during early prophase. Identification of a candidate axial

element component was straightforward, due to the presence of the easily recognized

HORMA (Hop1, Rev7, Mad2) domain among members of this family of proteins (Aravind and

Koonin, 1998; Vader and Musacchio, 2014). We identified a gene encoding a HORMA domain

protein that is most closely related to *Cel-him-3* by reciprocal BLAST analysis (Table S1). We

refer to this protein as Ppa-HOP-1, after the founding member of the meiotic HORMA

proteins, *S. cerevisiae* Hop1. We raised a polyclonal antibody by genetic immunization using a

fragment that encodes a peptide of 100 amino acids in length, including part of the HORMA

domain, and found that this antibody indeed recognized chromosome axes from meiotic entry

through late prophase.

To enable cytological detection of synaptonemal complex (SC) assembly, we searched

for homologs of SC proteins. This is notoriously challenging due to rapid divergence of these

proteins, and their extensive regions of coiled-coil potential, which is associated with a strongly

skewed amino acid composition. *C. elegans* expresses four known SC proteins, known as SYP-

1, SYP-2, SYP-3, and SYP-4 (Colaiácovo et al., 2003; MacQueen et al., 2002; Smolikov et al.,

2009, 2007). SYP-4 contains a distinctive C-terminal domain with several unusual motifs

enriched in glycine and phenylalanine residues. We generated and sequenced an RNA library

from isolated gonads, which facilitated identification of full-length *Ppa-syp-4*. We inserted an

HA epitope tag at the C-terminus of the protein via template-directed repair of a CRISPR/Cas9-

induced break, and found that immunofluorescence with an epitope-specific antibody

localized specifically between paired meiotic chromosomes, confirming SYP-4::HA as a marker

for the SC (Figure 3A, B). The tagged protein supported normal meiosis, as indicated by the

low percentage of inviable embryos and males in the strain population (Figure 3 – Supplement

1).

HOP-1 was detected in the nucleoplasm in the premeiotic region of the germline and

formed linear structures along chromosomes upon meiotic entry. SYP-4 was detected along

chromosome segments shortly thereafter, and fully colocalized with HOP-1 tracks during

pachytene. Notably, the region of the germline containing nuclei with fully aligned stretches of

SYP-4 and HOP-1 was very short compared to *C. elegans,* in which SC disassembly occurs close

to the bend of the gonad arm, shortly prior to diakinesis. In contrast, SC disassembly initiated

much earlier in *P. pacificus*; the major fraction of prophase nuclei is best described as "pachy-

diplotene," since chromosomes remain only partially synapsed. Six short stretches of SYP-4

were apparent in these nuclei, which persisted over an extended region (Figure 3A, B). This

asymmetrical pattern of the SC is highly reminiscent of a more transient stage in *C. elegans*, in

which the SC remains associated with the "short arm" of each homolog pair and subsequently

contributes to the step-wise loss of cohesion during the first and second meiotic divisions (Lui

and Colaiácovo, 2012; Zhang et al. 2018).

HOP-1 localized normally to chromosome axes in *spo-11* and *dmc-1* mutants, but

extensive SC assembly failed. Instead, small, dispersed puncta of SYP-4 were observed along

chromosome axes, with occasional longer tracks (Figure 3C). In contrast to *spo-11* and *dmc-1*

mutants, *rad-51* mutants displayed robust synapsis with a distribution of stages similar to that

seen in wild-type hermaphrodites (Figure 3C). These observations indicate that *spo-11* and

dmc-1 play crucial roles during homologous synapsis in *P. pacificus*, while *rad-51* is dispensable

for this process. This further suggests that synapsis initiation may be tightly coupled to

homolog pairing and that SPO-11 and DMC-1, but not RAD-51, play central roles in this

process.

DMC-1 and RAD-51 localize sequentially during distinct stages of meiotic prophase

To investigate the functions of and interplay between DMC-1 and RAD-51 in *P.*

pacificus, we inserted a V5 epitope tag at the C-terminus of DMC-1 using CRISPR/Cas9, and

raised a polyclonal antibody that recognizes Ppa-RAD-51 (see Materials and Methods). DMC-

1::V5 supported normal meiosis, as evidenced by a normal brood size, high embryonic viability

and low percentage of males (Figure 3 – Supplement 1). Surprisingly, the two proteins showed

distinct and nonoverlapping patterns of localization. DMC-1 localized broadly, appearing to

coat the chromosomes, in transition zone nuclei and disappeared immediately upon

completion of synapsis. RAD-51 displayed a much more restricted, punctate distribution along

chromosomes, and was only detected in nuclei in which DMC-1 no longer coated the

chromosomes (Figure 4A, B). Occasional nuclei at the border between the transition zone and

pachytene region exhibited both DMC-1 and RAD-51, although DMC-1 was very faint in these

nuclei and did not overlap with RAD-51 (Figure 4C). Additionally, DMC-1 remained strongly

associated with chromosomes in some late nuclei that retained clustered DAPI morphology,

presumably either "straggler" nuclei with delays in synapsis or CO designation, or apoptotic

cells, both of which are typcially observed in the germlines of wild-type *C. elegans* (Figure 4D).

We also tested the interdependence of DMC-1 and RAD-51 recombinases for their

localization. In *S. cerevisiae* and *A. thaliana*, Dmc1 functions as an essential catalyst for

interhomolog joint molecule formation during meiotic DSB repair, while Rad51 acts as an

accessory protein for Dmc1 nucleofilament formation (Cloud et al., 2012; Da Ines et al., 2013).

We did not detect RAD-51 along chromosomes in the transition zone, where DMC-1 was

abundant on chromatin, and we found that DMC-1::V5 localization was normal in *rad-51*

mutants, indicating that RAD-51 does not play an essential role in the recruitment of DMC-1

(Figure 4 – Supplement 1A). Conversely, RAD-51 was also observed in some nuclei in *dmc-1*

mutants, specifically in late prophase nuclei proximal to the extended transition zone (Figure 4

– Supplement 1B). RAD-51 foci were both more abundant and larger in *dmc-1* mutants

compared to wild-type pachytene nuclei, perhaps due to persistent unrepaired DSBs that

accumulate in the absence of DMC-1 protein. Alternatively, the bright foci of RAD-51 observed

in late prophase nuclei could be indicative of an apoptotic response to unrepaired breaks

and/or extensive asynapsis.

In *spo-11* mutants, which are expected to lack any meiotic DSBs, DMC-1 failed to

localize to chromosomes, and instead formed nuclear aggregates (Figure 4 – Supplement 1C).

It was unclear whether these DMC-1 aggregates were associated with chromatin. Since DMC-1

localizes broadly to chromosomes during early prophase, rather than specifically at

recombination intermediates, this suggests that the mislocalization reflects *spo-11*–dependent

regulation of DMC-1, likely through activation of a DNA damage signaling pathway, rather

than an absence of potential binding sites.

In contrast to DMC-1, RAD-51 does appear to associate specifically with recombination

intermediates. Thus, it was unsurprising to see that RAD-51 foci were absent from meiotic

nuclei in *spo-11* mutants (Figure 4 – Supplement 1E). Some RAD-51 foci were oberved in the

premeiotic region of *spo-11* mutants, as in wild-type animals, providing a positive control for

immunofluorescence (Figure 4 – Supplement 1D).

Together these observations indicate that DMC-1 and RAD-51 bind to chromatin at

different stages of meiotic prophase and are not interdependent, although both require DSBs

for their chromosome localization.

RAD-51 is not required for CO formation or completion of meiosis

To assess the roles of DMC-1 and RAD-51 in CO formation, we quantified the number of

DAPI-staining bodies at diakinesis in *dmc-1* and *rad-51* mutants. Wild-type oocytes at this

stage contained ~6 DAPI-staining bodies (average= 5.6), as expected, while in *spo-11* mutants,

~12 DAPI-staining bodies were present (average= 11.5), consistent with a complete failure of

CO formation in the absence of DSBs (Figure 5A,B). Interestingly, in *dmc-1* mutant germlines

we frequently failed to detect oocytes at diakinesis, indicative of a defect in meiotic

progression and the likely activation of a checkpoint in response to unrepaired DSBs. In cases

when we did observe nuclei at diakinesis, they contained an average of 11.6 DAPI-staining

bodies, reflecting a complete absence of COs, as in *spo-11* mutants (Figure 5C, D).

Somewhat surprisingly, disruption of rad-51 resulted in homozygous mutant

hermaphrodites that were viable and fertile, although animals produced smaller broods and

their embryos showed greatly reduced viability (Figure 5E). Homozygous rad-51 mutant

gonads also displayed diakinesis nuclei more frequently than dmc-1 mutants, although they

were absent in 2 out of 20 gonads scored, indicating that loss of DMC-1 function impairs

meiotic progression more severely than loss of RAD-51 (Figure 5C, D). Consistent with this

observation, while rad-51 mutants had a lower average brood size compared to wild-type

animals, dmc-1 mutants had even smaller broods, ranging from zero to 35 embryos laid per

mutant homozygote (Figure 5E). In striking contrast to Cel-rad-51 mutants, which display

fragmented chromatin aggregates at diakinesis (Martin et al., 2005; Rinaldo et al., 2002), Ppa-

rad-51 mutants displayed an average of 6 DAPI-staining bodies, similar to wild-type (Figure

5B). Together with the relatively high viability of progeny of rad-51 homozygous mutants, this

indicates that RAD-51 does not play an essential role in CO formation in P. pacificus.

Conserved CO factor COSA-1 marks Class I COs in P. pacificus

To further analyze CO formation in P. pacificus, we identified the gene encoding the

metazoan meiotic cyclin-related protein COSA-1/Cntd1 and inserted a 3xFLAG epitope tag at

the C-terminus of the coding sequence via CRISPR/Cas9. The strain expressing COSA-

1::3xFLAG yielded progeny with high embryonic viability and low percentage of males,

demonstrating that the tagged protein functions sufficiently to support normal meiosis (Figure

3 – Supplement 1). Immunostaining with anti-FLAG antibodies revealed discrete foci along the

SC, beginning as early as zygotene, which decreased in number and became more intense

during the brief pachytene region (Figure 6A, B). Most pachytene nuclei displayed 6 COSA-1

foci, each of which was associated with an individual SC between each pair of homologous

chromosomes (Figure 6C and Supplemental Video 1). Previous genetic mapping experiments

revealed that homolog pairs often experience multiple COs in *P. pacificus* (Srinivasan et al.,

2002). To investigate whether these extra COs might arise specifically during

spermatogenesis, we analyzed younger J4 hermaphrodites, which had not yet undergone the

switch from spermatogenesis to oogenesis, and found that pachytene nuclei undergoing

spermatogenesis also displayed ~6 COSA-1 foci (Figure 6- figure supplement 1). This suggests

that *P. pacificus*, like *C. elegans*, exhibits complete chromosome-wide Class I CO interference

and, combined with previous genetic data, implies that additional COs likely arise through the

Class II CO pathway, which does not require COSA-1.

Intriguingly, SC disassembly appeared to be regulated by the position of the lone Class I

CO between each chromosome pair, as in *C. elegans.* By mid-pachy-diplotene, 6 short

stretches of SYP-4::HA were observed, each associated with a single COSA-1::3xFLAG focus

near one end (Figure 6B). During late prophase, COSA-1::3xFLAG foci were no longer visible,

although short stretches of SYP-4::HA could still be observed. We observed splaying of

chromosome axes along the long arms upon removal of the central region proteins (Figure

6D). HOP-1 was retained on both arms upon disassembly of the SC, although the signal

appeared fainter along the long arms, perhaps due to splitting of the signal. At this stage, short

stretches of SYP-4::HA colocalize with corresponding bright stretches of HOP-1 (Figure 3B and

Figure 6D). Bivalents at diakinesis also show a cruciform structure similar to that seen in *C.*

elegans, indicative that only the Class I COs give rise to chiasmata that persist until diakinesis.

We next examined the localization of COSA-1::3xFLAG in various mutant backgrounds.

As expected, *dmc-1* mutants showed a complete absence of COSA-1 foci throughout prophase,

while 6 foci were readily observed in pachytene nuclei in *rad-51* mutants (Figure 7A). These

observations were consistent with the number of DAPI-staining bodies we observed at

diakinesis (Figure 5B). Surprisingly, a few bright COSA-1::3xFLAG foci were present throughout

prophase in *spo-11* mutants (Figure 7A). However, since ~12 DAPI-staining bodies were

observed during diakinesis, we conclude that these COSA-1 foci do not mark designated COs.

A similar phenomenon has been reported in *C. elegans spo-11* mutants (Nadarajan et al., 2017;

Pattabiraman et al., 2017), suggesting that COSA-1 can coalesce at sites lacking bona fide

recombination intermediates.

DISCUSSION

Distinct roles of Dmc1 and Rad51

Rad51 and Dmc1 show very similar activities *in vitro*; both are capable of binding to

single-stranded and double-stranded DNA *in vitro*, although both show a strong preference for

ssDNA binding (Hong et al., 2001; Li et al., 1997; Masson et al., 1999). However, these proteins

clearly play non-redundant functions in meiosis. This is thought to be due in part to the activity

of Dmc-1 specific cofactors Mnd1 and Hop2, as well as differential regulation of the timing and

activities of the two paralogs. Our analysis of the two RecA homologs in *P. pacificus*, RAD-51

and DMC-1, defines their distinct contributions during meiosis.

DMC-1 appears to coat much of the chromatin in transition zone nuclei, whereas RAD-

51 shows a much more punctate localization, and only after DMC-1 is largely removed from

chromosomes. These observations suggest that DMC-1 interacts extensively with double-

stranded DNA, while RAD-51 localizes specifically to single-stranded DNA at sites of active

recombination. Intriguingly, even though DMC-1 apparently localizes much more broadly, both

DMC-1 and RAD-51 depend on SPO-11 for their association with meiotic chromosomes. We

interpret these observations to indicate that the binding of DMC-1 to chromatin is regulated by

a mechanism that respods to SPO-11 activity, likely through activation of a DNA damage

response pathway. In contrast, the association of RAD-51 with recombination intermediates is

likely inhibited by this pathway, and is only permitted after DMC-1 has ensured the formation

of CO intermediates between each pair of chromosomes.

The sequential localization of DMC-1 and RAD-51 first suggested that they function

independently, and this is supported by our analysis of loss-of-function mutations. By contrast

to budding yeast and *A. thaliana*, RAD-51 is dispensable for the activity of DMC-1 in pairing,

synapsis, and CO formation. Instead, RAD-51 appears to play a supporting role in DSB repair

during pachytene, processing excess DSBs that remain after CO designation has occured. In *C.*

elegans, which expresses only RAD-51, a similar switch between two modes of double-strand

break repair is nevertheless observed during meiotic prophase (Hayashi et al., 2007).

Association of Ce-RAD-51 with repair intermediates is differentially regulated from the onset

of meiosis until a mid-pachytene transition that coincides with CO designation; at this time,

competence to convert DSBs to interhomolog COs is also lost. Recent work revealed an

analogous switch from a "meiotic" repair to a "somatic"-like repair pathway during the

transition from mid- to late pachytene in mouse spermatocytes (Enguita-Marruedo et al.,

2019). In *P. pacificus*, it is thus likely that DMC-1 and RAD-51 have highly specialized functions:

formation of interhomolog COs by DMC-1 in the transition zone and a more general mode of

double-strand break repair by RAD-51 during pachytene. Furthermore, our observation that

nuclei in *rad-51* mutants display cruciform bivalents and lack fragmented chromatin at

diakinesis suggests that excess DSBs can be repaired through an alternate pathway which does

not depend on RAD-51 activity, such as non-homologous end joining, or that DMC-1 can

compensate for RAD-51 but not the other way around. The nature of RAD-51-dependent DSB

repair and how the activities of RAD-51 and DMC-1 are regulated in early meiotic prophase will

be a focus of future investigation in this species.

CO regulation in *P. pacificus*

Genetic mapping in *P. pacificus* revealed lengths of ~100-250 centimorgans for each

chromosome, corresponding to 2-5 interhomolog COs per meiosis (Srinivasan et al., 2003,

2002). By contrast, the genetic map of each chromosome in *C. elegans* is 50 cM, reflecting a

single CO per pair (Hillers et al., 2017). Surprisingly, our analysis of Ppa-COSA-1 localization

revealed only a single COSA-1 focus per chromosome by pachytene, very similar to what is

seen in *C. elegans* (Yokoo et al., 2012). This suggests that either multiple Class I COs occur but

only one retains COSA-1, or that all but a single CO forms by an alternate, COSA-1-

independent pathway. In many eukaryotic systems, two CO pathways co-exist (Gray and

Cohen, 2016). Class I COs show spatial interference and depend on factors including MSH-4/5,

ZHP-3/Rnf212, and COSA-1/Cntd1. On the other hand, Class II COs do not exhibit interference

and undergo an alternate resolution pathway that requires the structure-specific endonuclease

complex Mus81-Eme1/Mms4. In some species, notably in *A. thaliana* and *S. cerevisiae*, Class II

COs can occur in the absence of Class I COs, but this may not be the case in all organisms.

Future work will examine whether all COs in *P. pacificus* arise only after the single, obligate

Class I CO has been designated, and whether the designation of Class I COs is necessary for the

formation and resolution of Class II COs. We further note that there appears to be only one

chiasma formed between each homolog pair, which is likely created by the lone COSA-1-

associated Class I CO. Our work reveals the potential of *P. pacificus* to address some long-

standing questions about mechanisms and regulation of CO recombination that are not

accessible in the meiotic model *C. elegans* due to the absence of Class II COs during normal

meiosis.

Comparative analysis of meiosis reveals major variations within the nematode lineage

In addition to establishing key aspects of meiosis in *P. pacificus*, this work also

illuminates the evolutionary history of meiosis in *C. elegans*. A body of prior work has revealed

that recombination-independent homologous synapsis in *C. elegans* relies on pairing centers,

specialized chromosome regions that interact with nuclear envelope and drive chromosome

movement during early prophase. These meiosis-specific dynamics are typically mediated by

telomeres, but have shifted to a unique region on each chromosome in *C. elegans*. Pairing

centers also act as the sites of synapsis initiation (Rog and Dernburg, 2013). By contrast, in

other organisms, telomere-led chromosome movement is thought to promote homologous

interactions, but stabilization of pairing and initiation of synapsis occur at early recombination

intermediates, which depend on Spo11 and Dmc1 activity. Pairing center activity depends on

and is largely defined by the recruitment of a family of zinc finger proteins that bind to DNA

sequence motifs in these regions (Phillips et al., 2009). These proteins, known as ZIM-1, ZIM-2,

and ZIM-3, and HIM-8 in *C. elegans*, also act as scaffolds to recruit a cascade of kinase activities

required for pairing and synapsis (Harper et al., 2011; Kim et al., 2015; Labella et al., 2011).

Surprisingly, most of the sequenced genomes of nematodes in Clade V include both homologs

of the HIM-8/ZIM family and orthologs of Dmc1, Hop2, and Mnd1 (Figure 2 – Supplement 1).

The *Pristionchus* genus is unusual within this Clade in that it lacks apparent homologs of the

pairing center proteins, while *Caenorhabditids* are among the few genera that have lost

Dmc1/Mnd1/Hop2, which were independently lost along the branch leading to *Oscheius tiuplae*

and *Auanema rhodensis*. This suggests that the dominant role of pairing centers in homologous

synapsis is likely to be recently derived, perhaps restricted to *Caenorhabditids*. Future studies

of species in which Dmc1 and pairing center proteins co-exist may illuminate how synapsis

initiation activity relocalized from sites of Dmc1-mediated strand exchange to pairing centers.

In addition to clarifying how pairing centers acquired their central meiotic roles in *C. elegans*,

further analysis may help to address the longstanding question of how recombination

intermediates trigger initiation of synapsis.

467   *C. elegans* is a popular and powerful model system for molecular studies of meiosis. *P.*

pacificus shares many of the same experimental advantages, although naturally has fewer

experimental tools, since it has been developed far more recently. Genome engineering using

CRISPR/Cas9 is somewhat more challenging, and has so far been refractory to large insertions,

such as fluorescent proteins. However, this barrier will likely be overcome through advances in

editing efficiency. Perhaps a greater obstacle is the absence of balancer chromosomes, which

are invaluable for maintaining mutations that reduce viability or fertility in *C. elegans*. Because

balancers are unavailable for *P. pacificus*, most of the mutations described in this work have

been maintained in unbalanced heterozygotes through frequent, labor-intensive genotyping

assays. Moreover, the analysis presented here suggests that it may not be possible to construct

crossover-suppressing balancer chromosomes, given that recombination is essential for

homolog pairing and synapsis. By contrast, the derived, recombination-independent

mechanism of pairing and synapsis in *C. elegans* makes it possible to propagate large-scale

chromosome rearrangements that suppress meiotic recombination over large genomic regions

while maintaining regular segregation of chromosomes.

482   The advent of broadly applicable techniques for genome editing has enabled rapid

progress towards developing *P. pacificus* for molecular studies, along with many other

experimental models. Future exploration of meiosis in *P. pacificus*, and perhaps in other

nematode models, will further expand our understanding of core mechanisms and plasticity of

sexual reproduction.

MATERIALS AND METHODS

P. pacificus *strains and maintenance*

490   Animals were cultured on NGM media with *E. coli* OP50 at 20°C under the same

standard conditions as *C. elegans* (Brenner, 1974). The wild-type strain is a derivative of PS312

designated as "97," which was provided by Ralf Sommer. This isolate was found to be more

amenable to genome editing by CRISPR-Cas9 than the parental strain. Mutant alleles were

maintained in a heterozygous state. Every few generations and before each

immunofluorescence or FISH experiment, single adult hermaphrodites were picked to new

plates and allowed to lay embryos for two days, after which the genotype of the parent was

determined by PCR genotyping. Progeny from heterozygous mothers, one-fourth of which are

homozygous for the meiotic mutation, were analyzed using the assays described here;

heterozygous and wild-type siblings were frequently used as controls, in addition to analysis of

unedited wild-type animals.

CRISPR/Cas9 genome editing

To modify the *P. pacificus* genome, we adapted our preferred CRISPR/Cas9 protocol

from *C. elegans* to *P. pacificus*. Equimolar quantities of Alt-R®CRISPR-Cas9 crRNA and

tracrRNA molecules (Integrated DNA Technologies, Coralville, IA) were hybridized using a

thermocycler (95°C for 5 minutes, then 25°C for 5 minutes). 4μl of 100μM hybridized

tracrRNA/crRNA was combined with 4μl of 40μM *S. pyogenes* Cas9-NLS purified protein (QB3

Macrolab, UC Berkeley, Berkeley, CA) and incubated at room temperature for 5 minutes. 2μl of

100μM stock of an Ultramer® DNA oligo (IDT) repair template containing 50-60 bp homology

arms and the desired epitope or mutation sequence was added to the mixture, for a total

volume of 10μl, and injected into the gonads of 24 hour-post J4 adult hermaphrodites.

Following a 2-4 hour recovery period, injected animals were allowed to lay embryos at 20°C for

16-20hr. Four days later, a fraction of the F1 population (typically 150-200 progeny from 6-8

injected P_0s) was screened for the presence of the mutation or epitope tag sequence by PCR

genotyping, and candidate alleles were verified by Sanger sequencing. A complete list of

crRNA, repair template, and genotyping primer sequences used to generate alleles in this

study is provided as Table S1.

TALEN constructs were generated using a protocol adapted from Zhang et al. (2011)

and designed using the TAL Effector Nucleotide Targeter 2.0 website (https://tale-

[nt.cac.cornell.edu/).](https://tale-nt.cac.cornell.edu/)

Viability and fertility

To quantify embryonic viability, brood size, and male progeny of wild-type and *rad-51*

mutants, J4 hermaphrodites were picked onto individual plates and transferred every 24 hours

over 72 hours total. Embryos were counted each day, after transferring the adult

hermaphrodite to a new plate, and kept at 20°C. Three to four days later, adults were counted

on each plate. To analyze *spo-11* and *dmc-1* mutants, 24 individual J4 hermaphrodites were

picked from progeny of a verified heterozygous mutant hermaphrodite. Quantification was

performed as in wild type, but after 72 hours, the adult hermaphrodite was lysed and

genotyped for the presence of the mutation. Thus, although 24 animals total were quantified

from a mixed population of *spo-11/+* or *dmc-1/+* animals, data from 5 homozygous *spo-11* and 7

homozygous *dmc-1* mutant animals are reported in the data table, Figure 5E.

Immunofluorescence and FISH

To stage animals for each experiment, 30-40 J4s were picked from a PCR-verified

heterozygous mother onto a fresh plate and allowed to develop for an additional 24 or 48

hours at 20°C. Young adult hermaphrodites were dissected on a clean coverslip in egg buffer

containing 0.05% tetramisole and 0.1% Tween-20. Samples were fixed for 2 minutes in egg

buffer with 1% formaldehyde and transferred to a 1.5-ml tube containing PBST. After 5

minutes, the PBST was replaced with ice-cold methanol and incubated at room temperature

for an additional 5 minutes. Worms were washed twice with PBST, blocked with Roche

blocking reagent, and stained with primary antibodies diluted in Roche blocking solution for

1.5-2 hours at room temperature. Samples were washed with PBST and incubated with

secondary antibodies raised in donkey and conjugated with Alexa-488, Cy3 or Cy5 (Jackson

ImmunoResearch Laboratories, West Grove, PA). Worms were then incubated with 1μg/ml

DAPI in PBST, washed with PBST, and mounted in ProLong™ Gold antifade mounting solution

(Invitrogen) before imaging.

For embryo staining, 20 plates of mixed-stage worms were harvested with water and

treated with 1:8 solution of bleach:water for 5 minutes at room temperature. Embryos were

collected by centrifugation and washed twice with PBS. To dissolve the vitelline membrane, a

solution containing 2.6 ml of n-heptane, 2 ml of PBS, and 1% paraformaldehyde (final) was

added to the embryo pellet for 5 minutes at room temperature with shaking. Treated embryos

were collected by centrifugation, washed twice with 5 ml MeOH, three times with PBS, and

incubated with Roche blocking reagent, primary, and secondary antibodies as described

above. Embryos were mounted on agarose pads for imaging.

For FISH experiments, age-matched animals were dissected and fixed as for

immunofluorescence experiments described above, except that the initial fixation was for 4

minutes in 2% formaldehyde. After incubation in ice-cold methanol, worms were washed with

2x SSCT twice and incubated in 50% 2x SSCT/50% formamide solution overnight at 37°C. The

next day, the worms were transferred to a small PCR tube, excess solution was removed from

the sample, and a 40 µl hybridization mix containing 250ng of each probe in hybridization

buffer (3.5xSSC, 58% formamide, 12.75% dextran sulfate) was added. The sample was

immediately denatured in a thermocycler at 91°C for 2 minutes and incubated overnight at

37°C. On the last day, the worms were transferred to a 1.5-ml tube and washed with 2xSSCT.

After 5 minutes, the solution was replaced with fresh 2xSSCT and mounted with ProLong™

Diamond Antifade Mountant with DAPI (Invitrogen).

FISH probes

Probes for a central locus on chromosome IV and the left end of chromosome X were

designed based on two short tandem repeat motifs. Tandem Repeat Finder v4.09 (Benson,

1999) was used to identify tandem repeats in *P. pacificus* "El Paco" genome assembly

(Rödelsperger et al., 2017) using default parameters and a maximum periodicity of 200 bp. The

output was then filtered to identify repeats that spanned more than 8 kb. These were

compared to the genome sequence using BLAST to identify the subset of sequences restricted

to a single major locus per genome. A subset of these repeats was then tested for specific and

robust hybridization with oligonucleotide probes. The Chromosome IV probe targets the 30-

base repeated motif TCATTGAAATGATCACAATCATTGA, which spans 40.1 kb at a position

11.3 Mb from the left end of chromosome IV. The Chromosome X probe

(GGTGGTCGACGGCTGCGTCG) targets a 30-base repeat motif that spans two very close

regions of 29.3kb and 11.1kb on the left end of the X chromosome. Single-stranded

oligonucleotides labeled at the 3' end with 6-FAM or Cy3 dyes were purchased from IDT and

used directly as FISH probes.

Antibodies

Antibodies against Ppa-RAD-51 were generated against a 6x His-tagged N-terminal

fusion protein (aa 1-103) expressed and purified from bacteria. Four mice were immunized with

the antigen. Serum from one animal, designated S148, was used without purification at 1:300

dilution (Pocono Rabbit Farm and Laboratory, Canadensis, PA). Antibodies against Ppa-HOP-1

were generated by genetic immunization against aa 177-276 (SDIX, Newark, DE) and used in

the following experiments at 1:300 dilution. Additional antibodies were purchased from

commercial sources and diluted as follows: mouse anti-FLAG (1:500, Sigma #F1804), mouse

anti-V5 (1:500, Thermo Fisher #R960-25), rabbit anti-V5 (1:250, Millipore Sigma #V8137), and

goat anti-HA (1:500, Novus Biologicals #NB600-362). Secondary antibodies raised in donkey

and labeled with Alexa 488, Cy3, or Cy5 were used at 1:400 dilution (Jackson ImmunoResearch

Laboratories).

Orthology analysis and phylogenetic inference

Accessions to all data used in orthology analysis are available in Table S2. We

downloaded the predicted protein sequences of 65 nematode species and five outgroup taxa

and filtered for the longest isoform of each gene. OrthoFinder (Emms and Kelly, 2015) was

used to cluster all protein sequences into putative orthologous groups (OGs) using the default

inflation value of 1.5. OGs containing loci which were present in at least 75% of species and

which were, on average, single copy (mean < 1.3) were selected. We aligned each selected OG

using MAFFT (Katoh and Standley, 2013) and generated a maximum likelihood tree along with

1000 ultrafast bootstraps (Hoang et al., 2018) using IQ-TREE (Nguyen et al., 2015), allowing the

best-fitting substitution model to be selected automatically (Kalyaanamoorthy et al., 2017).

Each tree was screened by PhyloTreePruner (Kocot et al., 2013); collapsing nodes with

bootstrap support <90), and any OGs containing paralogues were discarded. If two

representative sequences were present for any species (*i.e.*, "in-paralogs") after this paralog

screening step, the longest of the two sequences was retained and the other discarded. We

then realigned the remaining OGs using MAFFT and trimmed spuriously aligned regions using

trimAl (Capella-Gutiérrez et al., 2009). The trimmed alignments were subsequently

concatenated using catfasta2phyml (available at https://github.com/nylander/catfasta2phyml)

to form a supermatrix. We inferred the nematode species tree using IQ-TREE with the general

time reversible model (GTR) with gamma-distributed rate variation among sites. The resulting

tree was visualized using iTOL (Letunic and Bork, 2016).

We identified the OGs which contain orthologs of DMC-1, MND-1, HOP-2 and RAD-51

using BLASTP to search using the orthologous protein sequences from *Homo sapiens* as

queries. The OG containing the *C. elegans* proteins HIM-8, ZIM-1, ZIM-2, ZIM-3 was identified

using the appropriate transcript IDs. Each OG was aligned using MAFFT and a gene tree was

inferred using IQ-TREE, allowing the best-fitting substitution model to be selected

automatically. Each gene tree was visually inspected using iTOL.

Total RNAseq

Total RNA was isolated from 20 whole worms or 30 dissected gonads from 48 h post J4

animals using TRIzol (Invitrogen). TruSeq Stranded Total RNA (Illumina) sequencing libraries

were constructed following the manufacturer's instructions. For both conditions, three

independent libraries were constructed. Libraries were pooled and sequenced on a HiSeq4000

(150bp, PE, QB3 Vincent J. Coates Genomics Sequencing Laboratory). Reads were mapped to

the "El Paco" genome assembly including annotated splice sites (Rödelsperger et al., 2017)

using STAR. To correct misannotated splice sites, a transcriptome was then reconstructed de

novo using StringTie. Transdecoder was used to generate potential open reading frames. To

identify potential meiotic genes, we identified genes enriched in dissected gonads over whole

worms using HTSeq and edgeR.

Acknowledgements

This work was supported by funding from the Howard Hughes Medical Institute to AFD

and a fellowship from the Helen Hay Whitney Foundation to JB. This work used the Vincent J.

Coates Genomics Sequencing Laboratory at UC Berkeley, supported by NIH S10 OD018174

Instrumentation Grant. We are grateful to Ralf J. Sommer, Ray Hong, and other members of

the *P. pacificus* research community for providing us with *P. pacificus* strains and abundant

helpful advice. We thank members of the Dernburg lab for helpful discussions about this work

and critical reading of the manuscript.

Figure Legends

Figure 1. Germline organization and meiotic progression in *P. pacificus* is superficially similar to

C. elegans. A. Projection image of a *P. pacificus* hermaphrodite gonad stained with DAPI. Scale

bar, 30 μm. Insets show representative nuclei from the premeiotic region (PM), transition zone

(TZ), pachytene (Pach), diplotene (Dip), diffuse stage (Diff) and diakinesis. Scale bar, 5 μm. B.

Distal region of a *P. pacificus* germline injected with 0.3 nM Cy3-dUTP solution, dissected and

stained with DAPI after 30 minutes of recovery. Scale bar, 30 μm. C. Mitotic chromosomes

(DAPI) in a 2-4 cell stage embryo and the premeiotic germline of adult hermaphrodites

expressing CENP-C::V5 (magenta). Scale bar, 2 μm.

Figure 2. Stable homolog pairing requires early recombination factors. A. Diagram showing

the locations of tandem repeat sequences used to generate DNA FISH probes for pairing

analysis in *P. pacificus*. B. Representative images show the progression of homolog pairing of

Chromosome X (magenta) and Chromosome IV (yellow) during meiotic prophase in wild-type

hermaphrodites. Premeiotic region (PM), transition zone (TZ), and pachytene (Pach). Scale

bar, 5 μm. C. Representative images of FISH probe signals in *spo-11*, *dmc-1*, and *rad-51* mutants

during mid-prophase stage (roughly equivalent to the pachytene stage in wild-type germlines).

Scale bar, 5 μm. D. Temporal progression of X and IV chromosome pairing in WT, *spo-11*, *dmc-*

1, and *rad-51* mutants. Distance between pairs of corresponding FISH signals were measured in

3D using Softworx or Prism for three gonads of each genotype. Each gonad was divided into

five zones of equal length, from the distal tip to the bend of the gonad arm, and the distances

between pairs of homologous FISH signals are presented in a scatter plot diagram. E. Distance

between pairs of heterologous FISH signals were measured in premeiotic (PM) and transition

zone (TZ) nuclei in WT, *spo-11* and *dmc-1* mutants (spanning zones 1 and 2 only). Distances

between pairs of homologous FISH signals (Chrom X and IV combined) in TZ nuclei are

included for comparison. ***p<.0001, by Student's *t*-test.

Figure 2- figure supplement 1. Partial representation of the nematode lineage, including 65

nematode species, and five outgroup taxa showing the presence of meiotic proteins. D= Dmc1;

M= Mnd1; H= Hop2; Z= HIM-8/ZIM-1,2,3. *C. elegans* and *P. pacificus* are highlighted in blue.

Figure 2- figure supplement 2. A. Composite projection images of whole gonads stained with

DAPI from WT, *spo-11*, *dmc-1*, and *rad-51* mutants. The extent of the transition zone of each

gonad is underlined with dashed lines. Scale bar, 30 μm. B. Quantification of transition zone

length relative to the length from meiotic onset to the end of pachytene in WT (n=7), *spo-11*

(n=7), *dmc-1* (n=7), and *rad-51* (n=10). Error bars indicate mean ± standard deviation. *spo-11* and

dmc-1 mutants show significant differences from WT (p<0.0001) but not *Ppa-rad-51* (p=0.8426)

by ordinary one-way ANOVA.

Figure 3. SPO-11 and DMC-1 are required, while RAD-51 is dispensable, for homologous

synapsis in *P. pacificus*. A. Composite projection image of a wild-type strain expressing Ppa-

SYP-4::HA, stained with DAPI (gray), anti-HOP-1 (red), and anti-HA (green). Meiotic

progression is from left to right. Scale bar, 30 μm. B. Higher magnification images of wild-type

nuclei in the premeiotic region (PM), transition zone (TZ), pachytene (Pach), and pachy-

diplotene (P-D) stages. C. Localization of Ppa-SYP-4::HA and Ppa-HOP-1 in WT, *spo-11*, *dmc-1*,

and *rad-51* mutants during early and mid-prophase (roughly equivalent to the TZ and Pach

regions in wild-type germlines, respectively). Synapsis fails in *spo-11* and *dmc-1* mutants but

forms normally in the *rad-51* background. Scale bar, 5 μm.

Figure 3- figure supplement 1. All epitope-tagged proteins used in this study support normal

meiosis. Epitope-tagged alleles were generated by in-frame insertion into the endogenous

gene loci using CRISPR/Cas9 (see Materials and Methods), and homozygosed by selfing of

successfully edited progeny. Fidelity of meiotic segregation in the resulting strains was

analyzed by counting the frequency of viable embryos and male progeny among whole broods

from self-fertilizing hermaphrodites, as indicated. Quantification of wild-type broods is also

reported in Figure 5E.

Figure 4. DMC-1 and RAD-51 localize sequentially to meiotic chromosomes. A. Composite

projection image of a wild-type gonad expressing DMC-1::V5, stained with DAPI (blue), anti-V5

(magenta), and anti-RAD-51 (yellow). Meiotic progression is from left to right. Scale bar, 30

μm. Inset shows the distinct localization of DMC-1 (magenta) and RAD-51 (yellow) in the

transition zone and pachytene regions, respectively. Scale bar, 5 μm. B. Higher magnification

images of nuclei in the transition zone and pachytene region. DMC-1 is present along

chromatin in the transition zone and disappears at pachytene. By contrast, RAD-51 localizes to

discrete foci starting at pachytene. Scale bar, 5 μm. C. Occasional nucleus on the border of the

transition zone and pachytene region display both DMC-1 and RAD-51. The signals do not

completely overlap. Scale bar, 2 μm. D. Nucleus with clustered DAPI morphology and strong

association of DMC-1 during later prophase. DMC-1 localizes to chromatin in "straggler" cells

that presumably have not completed synapsis or CO designation, or are undergoing apoptosis.

Scale bar, 2 μm.

Figure 4- figure supplement 1. DMC-1 and RAD-51 do not depend on each other for their

localization to chromosomes, but both require DSBs. A. DMC-1 (magenta) is abundant on

chromosomes in transition zone nuclei of *rad-51* mutants. Anti-RAD-51 immunofluorescence

was used to identify homozygous mutants among the progeny of heterozygotes. B. RAD-51

foci are observed in late pachytene nuclei, proximal to the extended transition zone, of *dmc-1*

mutants. RAD-51 foci appear larger and more numerous than in wild-type pachytene nuclei. C.

DMC-1 forms nuclear aggregates in *spo-11* mutants and does not localize along chromosomes.

D and E. RAD-51 foci are detected in premeiotic nuclei of *spo-11* mutants but are absent during

meiotic prophase.

Figure 5. CO formation requires SPO-11 and DMC-1, but not RAD-51. A. Representative

images of DAPI-staining bodies at diakinesis for each indicated genotype. Scale bar, 5 μm. B.

Quantification of DAPI-staining bodies in the -1 oocyte at diakinesis for each indicated

genotype (*n*= represents number of nuclei scored). C. Quantification of gonads which lacked

nuclei with DAPI-staining bodies at diakinesis stage. *n* is the number of germlines scored for

each genotype. D. Representative images of wild type, *dmc-1*, and *rad-51* mutant proximal

germlines. In wild-type germlines, diakinesis nuclei with fully condensed DAPI-staining bodies

are present distal to the spermatheca. However, nuclei with late prophase stage DAPI

morphology are frequently found adjacent to the spermatheca in *dmc-1* mutants and more

rarely in *rad-51* mutants. Meiotic progression is from left to right. Scale bar, 5 μm. E.

Frequencies of viable embryos and male progeny of whole broods from wild type, *spo-11*, *dmc-*

1 and *rad-51* mutant hermaphrodites.

Figure 6. COSA-1/Cntd1 accumulates at a single site per chromosome pair. A. Composite

projection image of a wild-type strain expressing three epitope-tagged proteins (COSA-

1::3xFLAG, DMC-1::V5, and SYP-4::HA), stained with anti-FLAG, anti-V5 and anti-HA

antibodies. Scale bar, 30 μm. B. Higher magnification images of nuclei from the transition zone

(TZ), pachytene (pach), mid- and late pachy-diplotene (P-D). COSA-1::3xFLAG (green) foci are

detected in transition zone nuclei but do not colocalize with DMC-1::V5 (cyan). Foci peak in

brightness in pachytene nuclei and gradually become dimmer until they are no longer detected

during late pachy-diplotene. In early to mid- pachy-diplotene nuclei, six short stretches of SYP-

4::HA (magenta) are observed per nucleus, each associated with a single COSA-1::3xFLAG

focus. Scale bar, 5 μm. C. Histogram showing the number of COSA-1::3xFLAG foci observed

per nucleus in the pachytene region. Analysis was restricted to 15 nuclei per gonad

immediately proximal to the transition zone and lacking DMC-1::V5 signal. Five individual

gonads were analyzed, for a total of 75 nuclei scored. D. Partial projection of a representative

nucleus in mid to late pachy-diplotene, stained with anti-HOP-1 (blue), anti-HA (marking the

SC, magenta), and anti-FLAG (marking COSA-1, green). A single COSA-1::3xFLAG focus is

observed at a junction (marked with a red arrowhead) between the "short arm," where SYP-

4::HA is retained, and splayed "long arms" lacking SC but positive for HOP-1. Scale bar, 2 μm.

Figure 6 - figure supplement 1. COSA-1/Cntd1 accumulates at a single site per chromosome

pair during spermatogenesis. Composite projection image of a wild-type gonad from a J4-

stage hermaphrodite expressing COSA-1::3xFLAG (green) and SYP-4::HA (magenta). Scale

bar, 30 μm. At this stage the germline is still undergoing spermatogenesis. Inset shows a

higher magnification image of nuclei in the pachytene region. As during oogenesis, ~6 COSA-

1::3xFLAG foci are observed in pachytene nuclei during spermatogenesis. Scale bar, 5 μm.

Figure 6 - Supplemental Video 1. COSA-1/Cntd1 accumulates at a single site per chromosome

pair. 3D volume rendering of a single nucleus from the pachytene region of a wild-type worm

expressing COSA-1::3xFLAG (green) and SYP-4::HA (magenta). Each frame is a maximum

intensity projection from a region of a deconvolved 3D image. Each stretch of SYP-4::HA is

associated with a COSA-1::3xFLAG focus.

Figure 7. COSA-1::3xFLAG accumulates at sites of presumptive Class I COs. A. Nuclei from

hermaphrodites of the indicated genotype displaying COSA-1::3xFLAG (green) in early and

mid-prophase (roughly equivalent to the transition zone and pachytene regions in wild-type

germlines, respectively). COSA-1 foci are absent in *dmc-1* mutants, but 6 foci per nucleus are

detected in wild type and *rad-51* mutants. A few small foci are detected in *spo-11* mutants.

Scale bar, 5 μm.

REFERENCES

Aravind L, Koonin EV. 1998. The HORMA domain: a common structural denominator in mitotic
checkpoints, chromosome synapsis and DNA repair. *Trends Biochem Sci* **23**:284–286.
Benson G. 1999. Tandem repeats finder: a program to analyze DNA sequences. *Nucleic Acids*
Research. doi:10.1093/nar/27.2.573
Bishop DK, Park D, Xu L, Kleckner N. 1992. DMC1: A meiosis-specific yeast homolog of E. coli
recA required for recombination, synaptonemal complex formation, and cell cycle
progression. *Cell* **69**:439–456. doi:10.1016/0092-8674(92)90446-j
Bolt BJ, Rodgers FH, Shafie M, Kersey PJ, Berriman M, Howe KL. 2018. Using WormBase
ParaSite: An Integrated Platform for Exploring Helminth Genomic Data. *Methods Mol Biol*
1757:471–491. doi:10.1007/978-1-4939-7737-6_15
Brenner S. 1974. The genetics of Caenorhabditis elegans. *Genetics* **77**:71–94.
Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. 2009. trimAl: a tool for automated
alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**:1972–1973.
doi:10.1093/bioinformatics/btp348
Christophorou N, Rubin T, Huynh J-R. 2013. Synaptonemal complex components promote
centromere pairing in pre-meiotic germ cells. *PLoS Genet* **9**:e1004012.
doi:10.1371/journal.pgen.1004012
Cloud V, Chan Y-L, Grubb J, Budke B, Bishop DK. 2012. Rad51 is an accessory factor for Dmc1-
mediated joint molecule formation during meiosis. *Science* **337**:1222–1225.
doi:10.1126/science.1219379
Colaiácovo MP, MacQueen AJ, Martinez-Perez E, McDonald K, Adamo A, La Volpe A,
Villeneuve AM. 2003. Synaptonemal Complex Assembly in C. elegans Is Dispensable for
Loading Strand-Exchange Proteins but Critical for Proper Completion of Recombination.
Developmental Cell. doi:10.1016/s1534-5807(03)00232-6
Couteau F, Belzile F, Horlow C, Grandjean O, Vezon D, Doutriaux M-P. 1999. Random
Chromosome Segregation without Meiotic Arrest in Both Male and Female Meiocytes of a
dmc1 Mutant of Arabidopsis. *Plant Cell* **11**:1623. doi:10.2307/3871042
Da Ines O, Degroote F, Goubely C, Amiard S, Gallego ME, White CI. 2013. Meiotic
Recombination in Arabidopsis Is Catalysed by DMC1, with RAD51 Playing a Supporting
Role. *PLoS Genet* **9**:e1003787. doi:10.1371/journal.pgen.1003787
Dieterich C, Clifton SW, Schuster LN, Chinwalla A, Delehaunty K, Dinkelacker I, Fulton L,
Fulton R, Godfrey J, Minx P, Mitreva M, Roeseler W, Tian H, Witte H, Yang S-P, Wilson RK,
Sommer RJ. 2008. The Pristionchus pacificus genome provides a unique perspective on
nematode lifestyle and parasitism. *Nat Genet* **40**:1193–1198. doi:10.1038/ng.227
Emms DM, Kelly S. 2015. OrthoFinder: solving fundamental biases in whole genome
comparisons dramatically improves orthogroup inference accuracy. *Genome Biol* **16**:157.
doi:10.1186/s13059-015-0721-2
Enguita-Marruedo A, Martín-Ruiz M, García E, Gil-Fernández A, Parra MT, Viera A, Rufas JS,
Page J. 2019. Transition from a meiotic to a somatic-like DNA damage response during
the pachytene stage in mouse meiosis. *PLoS Genet* **15**:e1007439.
doi:10.1371/journal.pgen.1007439

Gray S, Cohen PE. 2016. Control of Meiotic Crossovers: From Double-Strand Break Formation
to Designation. *Annu Rev Genet* **50**:175–210. doi:10.1146/annurev-genet-120215-035111
Grelon M. 2001. AtSPO11-1 is necessary for efficient meiotic recombination in plants. *EMBO J*
20:589–600. doi:10.1093/emboj/20.3.589
Harper NC, Rillo R, Jover-Gil S, Assaf ZJ, Bhalla N, Dernburg AF. 2011. Pairing centers recruit a
Polo-like kinase to orchestrate meiotic chromosome dynamics in C. elegans. *Dev Cell*
21:934–947. doi:10.1016/j.devcel.2011.09.001
Hayashi M, Chin GM, Villeneuve AM. 2007. C. elegans germ cells switch between distinct
modes of double-strand break repair during meiotic prophase progression. *PLoS Genet*
3:e191. doi:10.1371/journal.pgen.0030191
Hillers KJ, Jantsch V, Martinez-Perez E, Yanowitz JL. 2017. Meiosis. *WormBook* **2017**:1–43.
doi:10.1895/wormbook.1.178.1
Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS. 2018. UFBoot2: Improving the
Ultrafast Bootstrap Approximation. *Mol Biol Evol* **35**:518–522. doi:10.1093/molbev/msx281
Hong EL, Shinohara A, Bishop DK. 2001. Saccharomyces cerevisiae Dmc1 protein promotes
renaturation of single-strand DNA (ssDNA) and assimilation of ssDNA into homologous
super-coiled duplex DNA. *J Biol Chem* **276**:41906–41912. doi:10.1074/jbc.M105563200
Hong RL, Sommer RJ. 2006. Pristionchus pacificus: a well-rounded nematode. *Bioessays*
28:651–659. doi:10.1002/bies.20404
Joyce EF, Apostolopoulos N, Beliveau BJ, Wu C-T. 2013. Germline progenitors escape the
widespread phenomenon of homolog pairing during Drosophila development. *PLoS Genet*
9:e1004013. doi:10.1371/journal.pgen.1004013
Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. 2017. ModelFinder: fast
model selection for accurate phylogenetic estimates. *Nat Methods* **14**:587–589.
doi:10.1038/nmeth.4285
Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. *Mol Biol Evol* **30**:772–780.
doi:10.1093/molbev/mst010
Keeney S. 2008. Spo11 and the Formation of DNA Double-Strand Breaks in Meiosis. *Genome*
Dyn Stab **2**:81–123. doi:10.1007/7050_2007_026
Keeney S, Giroux CN, Kleckner N. 1997. Meiosis-specific DNA double-strand breaks are
catalyzed by Spo11, a member of a widely conserved protein family. *Cell* **88**:375–384.
Kim Y, Kostow N, Dernburg AF. 2015. The Chromosome Axis Mediates Feedback Control of
CHK-2 to Ensure Crossover Formation in C. elegans. *Dev Cell* **35**:247–261.
doi:10.1016/j.devcel.2015.09.021
Kocot KM, Citarella MR, Moroz LL, Halanych KM. 2013. PhyloTreePruner: A Phylogenetic Tree-
Based Approach for Selection of Orthologous Sequences for Phylogenomics. *Evol*
Bioinform Online **9**:429–435. doi:10.4137/EBO.S12813
Labella S, Woglar A, Jantsch V, Zetka M. 2011. Polo kinases establish links between meiotic
chromosomes and cytoskeletal forces essential for homolog pairing. *Dev Cell* **21**:948–958.
doi:10.1016/j.devcel.2011.07.011
Letunic I, Bork P. 2016. Interactive tree of life (iTOL) v3: an online tool for the display and
annotation of phylogenetic and other trees. *Nucleic Acids Res* **44**:W242–5.
doi:10.1093/nar/gkw290

Li Z, Golub EI, Gupta R, Radding CM. 1997. Recombination activities of HsDmc1 protein, the
meiotic human homolog of RecA protein. *Proc Natl Acad Sci U S A* **94**:11221–11226.
Lo T-W, Pickle CS, Lin S, Ralston EJ, Gurling M, Schartner CM, Bian Q, Doudna JA, Meyer BJ.
2013. Precise and heritable genome editing in evolutionarily diverse nematodes using
TALENs and CRISPR/Cas9 to engineer insertions and deletions. *Genetics* **195**:331–348.
doi:10.1534/genetics.113.155382
Lui DY, Colaiácovo MP. 2012. Meiotic Development in Caenorhabditis elegansAdvances in
Experimental Medicine and Biology. pp. 133–170. doi:10.1007/978-1-4614-4015-4_6
MacQueen AJ, Colaiácovo MP, McDonald K, Villeneuve AM. 2002. Synapsis-dependent and -
independent mechanisms stabilize homolog pairing during meiotic prophase in C. elegans.
Genes Dev **16**:2428–2442. doi:10.1101/gad.1011602
MacQueen AJ, Phillips CM, Bhalla N, Weiser P, Villeneuve AM, Dernburg AF. 2005.
Chromosome sites play dual roles to establish homologous synapsis during meiosis in C.
elegans. *Cell* **123**:1037–1050. doi:10.1016/j.cell.2005.09.034
Martinez-Perez E, Colaiácovo MP. 2009. Distribution of meiotic recombination events: talking
to your neighbors. *Curr Opin Genet Dev* **19**:105–112. doi:10.1016/j.gde.2009.02.005
Martin JS, Winkelmann N, Petalcorin MIR, McIlwraith MJ, Boulton SJ. 2005. RAD-51-
dependent and -independent roles of a Caenorhabditis elegans BRCA2-related protein
during DNA double-strand break repair. *Mol Cell Biol* **25**:3127–3139.
doi:10.1128/MCB.25.8.3127-3139.2005
Masson JY, Davies AA, Hajibagheri N, Van Dyck E, Benson FE, Stasiak AZ, Stasiak A, West SC.
1999. The meiosis-specific recombinase hDmc1 forms ring structures and interacts with
hRad51. *EMBO J* **18**:6552–6560. doi:10.1093/emboj/18.22.6552
Nadarajan S, Lambert TJ, Altendorfer E, Gao J, Blower MD, Waters JC, Colaiácovo MP. 2017.
Polo-like kinase-dependent phosphorylation of the synaptonemal complex protein SYP-4
regulates double-strand break formation through a negative feedback loop. *Elife* **6**.
doi:10.7554/eLife.23437
Namai S, Sugimoto A. 2018. Transgenesis by microparticle bombardment for live imaging of
fluorescent proteins in Pristionchus pacificus germline and early embryos. *Dev Genes Evol*
228:75–82. doi:10.1007/s00427-018-0605-z
Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. 2015. IQ-TREE: a fast and effective
stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol*
32:268–274. doi:10.1093/molbev/msu300
Pattabiraman D, Roelens B, Woglar A, Villeneuve AM. 2017. Meiotic recombination modulates
the structure and dynamics of the synaptonemal complex during C. elegans meiosis.
doi:10.1101/110064
Penkner AM, Fridkin A, Gloggnitzer J, Baudrimont A, Machacek T, Woglar A, Csaszar E,
Pasierbek P, Ammerer G, Gruenbaum Y, Jantsch V. 2009. Meiotic chromosome homology
search involves modifications of the nuclear envelope protein Matefin/SUN-1. *Cell*
139:920–933. doi:10.1016/j.cell.2009.10.045
Phillips CM, Dernburg AF. 2006. A family of zinc-finger proteins is required for chromosome-
specific pairing and synapsis during meiosis in C. elegans. *Dev Cell* **11**:817–829.
doi:10.1016/j.devcel.2006.09.020
Phillips CM, Wong C, Bhalla N, Carlton PM, Weiser P, Meneely PM, Dernburg AF. 2005. HIM-8

binds to the X chromosome pairing center and mediates chromosome-specific meiotic
synapsis. *Cell* **123**:1051–1063. doi:10.1016/j.cell.2005.09.035
Pires-daSilva A. 2004. Conservation of the global sex determination gene tra-1 in distantly
related nematodes. *Genes Dev* **18**:1198–1208. doi:10.1101/gad.293504
Pittman DL, Cobb J, Schimenti KJ, Wilson LA, Cooper DM, Brignull E, Handel MA, Schimenti
JC. 1998. Meiotic Prophase Arrest with Failure of Chromosome Synapsis in Mice Deficient
for Dmc1 , a Germline-Specific RecA Homolog. *Mol Cell* **1**:697–705. doi:10.1016/s1097-
2765(00)80069-6
Rinaldo C, Bazzicalupo P, Ederle S, Hilliard M, La Volpe A. 2002. Roles for Caenorhabditis
elegans rad-51 in meiosis and in resistance to ionizing radiation during development.
Genetics **160**:471–479.
Rockmill B, Sym M, Scherthan H, Roeder GS. 1995. Roles for two RecA homologs in promoting
meiotic chromosome synapsis. *Genes Dev* **9**:2684–2695. doi:10.1101/gad.9.21.2684
Rödelsperger C, Meyer JM, Prabh N, Lanz C, Bemm F, Sommer RJ. 2017. Single-Molecule
Sequencing Reveals the Chromosome-Scale Genomic Architecture of the Nematode
Model Organism Pristionchus pacificus. *Cell Rep* **21**:834–844.
doi:10.1016/j.celrep.2017.09.077
Rog O, Dernburg AF. 2013. Chromosome pairing and synapsis during Caenorhabditis elegans
meiosis. *Current Opinion in Cell Biology*. doi:10.1016/j.ceb.2013.03.003
Rubin T, Christophorou N, Huynh J-R. 2016. How to pre-pair chromosomes for meiosis. *Cell*
Cycle **15**:609–610. doi:10.1080/15384101.2015.1131524
Rudel D, Riebesell M, Sommer RJ. 2005. Gonadogenesis in Pristionchus pacificus and organ
evolution: development, adult morphology and cell–cell interactions in the hermaphrodite
gonad. *Developmental Biology*. doi:10.1016/j.ydbio.2004.09.021
Sato A, Isaac B, Phillips CM, Rillo R, Carlton PM, Wynne DJ, Kasad RA, Dernburg AF. 2009.
Cytoskeletal forces span the nuclear envelope to coordinate meiotic chromosome pairing
and synapsis. *Cell* **139**:907–919. doi:10.1016/j.cell.2009.10.039
Smolikov S, Eizinger A, Schild-Prufert K, Hurlburt A, McDonald K, Engebrecht J, Villeneuve
AM, Colaiácovo MP. 2007. SYP-3 restricts synaptonemal complex assembly to bridge
paired chromosome axes during meiosis in Caenorhabditis elegans. *Genetics* **176**:2015–
2025. doi:10.1534/genetics.107.072413
Smolikov S, Schild-Prüfert K, Colaiácovo MP. 2009. A yeast two-hybrid screen for SYP-3
interactors identifies SYP-4, a component required for synaptonemal complex assembly
and chiasma formation in Caenorhabditis elegans meiosis. *PLoS Genet* **5**:e1000669.
doi:10.1371/journal.pgen.1000669
Sommer RJ. 2015. Pristionchus pacificus: A Nematode Model for Comparative and
Evolutionary Biology. BRILL.
Sommer RJ, Carta LK, Kim S-Y, Sternburg PW. 1996. Morphological, genetic and molecular
description of Pristionchus pacificus sp. N. (Nematoda: Neodiplogastridae). *Fundam. Appl.*
Nematol. **19:** 511-521
Srinivasan J, Sinz W, Jesse T, Wiggers-Perebolte L, Jansen K, Buntjer J, van der Meulen M,
Sommer RJ. 2003. An integrated physical and genetic map of the nematode Pristionchus
pacificus. *Mol Genet Genomics* **269**:715–722. doi:10.1007/s00438-003-0881-8
Srinivasan J, Sinz W, Lanz C, Brand A, Nandakumar R, Raddatz G, Witte H, Keller H, Kipping I,

Pires-daSilva A, Jesse T, Millare J, de Both M, Schuster SC, Sommer RJ. 2002. A bacterial
artificial chromosome-based genetic linkage map of the nematode Pristionchus pacificus.
Genetics **162**:129–134.
Vader G, Musacchio A. 2014. HORMA domains at the heart of meiotic chromosome dynamics.
Dev Cell **31**:389–391. doi:10.1016/j.devcel.2014.11.009
Villeneuve AM, Hillers KJ. 2001. Whence meiosis? *Cell* **106**:647–650.
Witte H, Moreno E, Rödelsperger C, Kim J, Kim J-S, Streit A, Sommer RJ. 2015. Gene
inactivation using the CRISPR/Cas9 system in the nematode Pristionchus pacificus. *Dev*
Genes Evol **225**:55–62. doi:10.1007/s00427-014-0486-8
Yokoo R, Zawadzki KA, Nabeshima K, Drake M, Arur S, Villeneuve AM. 2012. COSA-1 reveals
robust homeostasis and separable licensing and reinforcement steps governing meiotic
crossovers. *Cell* **149**:75–87. doi:10.1016/j.cell.2012.01.052
Yoshida K, Kondoh G, Matsuda Y, Habu T, Nishimune Y, Morita T. 1998. The Mouse RecA -like
Gene Dmc1 Is Required for Homologous Chromosome Synapsis during Meiosis. *Mol Cell*
1:707–718. doi:10.1016/s1097-2765(00)80070-2
Zhang L, Köhler S, Rillo-Bohn R, Dernburg AF. 2018. A compartmentalized signaling network
mediates crossover control in meiosis. *Elife* **7**. doi:10.7554/eLife.30789
Zickler D, Kleckner N. 1999. Meiotic chromosomes: integrating structure and function. *Annu*
Rev Genet **33**:603–754. doi:10.1146/annurev.genet.33.1.603

Figure 1

A

DAPI

PM TZ Pach Dip Diff diakinesis

B

DAPI Cy3-dUTP

C CENPC::V5 DAPI merge

embryo

premeiotic germline

Figure 2

A

Chrom IV

Chrom X

B WT

PM | TZ | Pach

C meiotic mutants during mid-prophase

spo-11 | dmc-1 | rad-51

D

WT Chrom X

spo-11 Chrom X

dmc-1 Chrom X

rad-51 Chrom X

distance between FISH signals (µm)

zone 1   zone 2   zone 3   zone 4   zone 5

WT Chrom IV

spo-11 Chrom IV

dmc-1 Chrom IV

rad-51 Chrom IV

distance between FISH signals (µm)

zone 1   zone 2   zone 3   zone 4   zone 5

E

heterologous vs. homologous FISH signals

distance between FISH signals (µm)

*** | ns | ns

WT PM heterologous
WT TZ heterologous
WT TZ homologous
spo-11 PM heterologous
spo-11 TZ heterologous
spo-11 TZ homologous
dmc-1 PM heterologous
dmc-1 TZ heterologous
dmc-1 TZ homologous

Figure 2- figure supplement 1

Figure 2- figure supplement 2

A

WT

spo-11

dmc-1

rad-51

B

Figure 3

Figure 4

A

DAPI DMC-1::V5 RAD-51

TZ ⟶ Pachytene

B

DAPI   DMC-1::V5   RAD-51   merge

TZ

Pach

C

RAD-51
DMC-1::V5

D

DAPI
DMC-1::V5

Figure 4- figure supplement 1

Figure 5

A WT spo-11 dmc-1 rad-51

B Number of DAPI-staining bodies at diakinesis

WT (n= 39)
spo-11 (n=15)
dmc-1 (n=16)
rad-51 (n=18)

C % germlines without diakinesis nuclei

WT (n=39) dmc-1 (n=39) rad-51 (n=20)

D WT dmc-1 rad-51

E

Genotype	% Egg viability (±SD)	% Male progeny (±SD)	Eggs laid (± SD)
WT (n=30)	92.9 (±16.3)	0.9 (±1.0)	205 (±55)
spo-11 (n=5)	0.3 (± .4)	20 (±44.7)	125 (±40)
dmc-1 (n=7)	0 (±0)	N/A	12 (±12)
rad-51 (n=14)	15.0 (±17.9)	4.5 (±5.1)	121 (±70)

Figure 6

Figure 6- figure supplement 1

A

Figure 7

A

Table S1

Strain	Allele	guide RNA target sequence (PAM sequence underlined)	ssDNA template	genotyping primer sequences
cenpc::V5	*ie1007*	ATGAAGAGATGGATTATAGTAGG	GGATATTAAAAAAAGGGGGTAAAATTGTACATCGCAAAAGGATGAGGAGTCATTTAATTCATCTTACTATAATCCATCTCTTCAGGTGGAGTCGAGTCCAAGAAGTGGGTTTGGAATTGGCTTTCCAGATCCGGCTCCTTCATTCTCTTCATCGAACTCTTCCACTTCCTTCTCCTTCTGCTCTTTTCTCTTCTTGTT	(f)TTTCTCCAGGAGTGGTTATCG (r)ACATCGCAAAAGGATGAGGAG
cosa-1::3xflag	*ie1003*	CTTTATTCTTCATTTTACAGTGG	GATCATCCCAGGGAGAGAACGACCTACCCAAACAGAGATCATAGATCTAATTATCCACTGGGAGCCGGATCTGATTATAAAGACCATGATGGAGACTATAAGGATCACGATATTGATTACAAAGACGATGATGATAAATAAAATGAAGAATAAAGAGTATTAAATTTATGTTTGTGTTCGTTTTTGTAATTACTGCTTTG	(f)ACGACCTACCCAAACAGAGA (r)CGGATGTGGAAAGACGTACC
dmc-1 mutant	*ie1005*	TTCGATAAGCTGCTTGGAGGTGG	ATTTAATTGTAACATTCAGACATCCAGTTGATCAACTAACCTCCGTTATTGCCTGACTTTCGATTCCACCactagtgTCCAAGCAGCTTATCGAATTCAACACTTCCAGTGGAGATTTTAAAGACTTGCTTGCGTCGTGAACACACT	(f)GGACTCTCGGAGGCTAAAGT (r)ATTCTCGAGCATTGCTTCCT
dmc-1::V5	*ie1001*	GCAACGTTTGCCATTGCAGCAGG	GATAAAACATGATCTTTTGTCCTTCATAATTGATTAGGTGGAGTCGAGTCCAAGAAGTGGGTTTGGAATTGGCTTTCCATCCTTTGCATCGACAATTCCTCCTGC**G**GC**G**ATGGCAAACGTTGCTTCGTTCTCAGGCATATC	(f)TGCCTGAGAACGAAGCAACG (r)ACATGAGATGGCACAAAGGAC
rad-51 mutant	*ie1006*	GTCGAGAACGAGGAGAATGCCGG	ATTTATTGGTTACCTCAAGGGACATGATGGACTGGCAGGCGAGTCCGGCATACTAGTTTCTCCTCGTTCTCGACATCAGCGTCGACGTGCGCCATTTGAGCGGACAT	(f)TTCTAGTGACGCGTGTTGTT (r)ACGAATCCTCGTTGCTGAAG
spo-11 mutant	*ie1004*	ATTCAGAACTTGGCAGAGATCGG	ACAAACATCTTTTTGCACGACAGGATTCTCTCAATCGATCAGTTTCCGATaactagtaCTCTGCCAAGTTCTGAATATGCAAAGATGTCAAGTCAATGTGGTAAGT	(f)GGAAATCCTTCGTTCTCACTATGG (r)GTCTCAATATCAGACAATTTCATTCCG
syp-4::HA	*ie1002*	GGAGGAGAATTCAACTTCTTCGG	CAAATGGAGGTGGCGGCGGGGGAGGAGGAGAGTTTAATTTTTTTGGTTTTTACCCCTACGATGTCCCAGATTATGCTTAAACCAATTTTTTCGAGCTTGGTGTAATGTATCCA	(f)CCCGTTGATGATGCTACCAG (r)GATACATTACACCAAGCTCGAA

Table S2

Species	Prefix	Source	BioProject
Acanthocheilonema viteae	AVITAE	WormBase ParaSite	PRJEB4306
Acrobeloides nanus	ANANUS	WormBase ParaSite	PRJEB26554
Ancylostoma caninum	ACANIN	WormBase ParaSite	PRJNA72585
Ancylostoma ceylanicum	ACEYLA	WormBase ParaSite	PRJNA231479
Angiostrongylus cantonensis	ACANTO	WormBase ParaSite	PRJEB493
Angiostrongylus costaricensis	ACOSTA	WormBase ParaSite	PRJEB494
Anisakis simplex	ASIMPL	WormBase ParaSite	PRJEB496
Ascaris lumbricoides	ALUMBR	WormBase ParaSite	PRJEB4950
Ascaris suum	ASSUUM	WormBase ParaSite	PRJNA62057
Auanema rhodensis	ARHODE	-	PRJEB29492
Brugia malayi	BMALAY	WormBase ParaSite	PRJNA10729
Brugia pahangi	BPAHAN	WormBase ParaSite	PRJEB497
Bursaphelenchus xylophilus	BXYLOP	WormBase ParaSite	PRJEA64437
Caenorhabditis elegans	CELEGA	WormBase ParaSite	PRJNA13758
Caenorhabditis monodelphis	CMONOD	caenorhabditis.org	PRJEB7905
Cylicostephanus goldi	CGOLDI	WormBase ParaSite	PRJEB498
Dictyocaulus viviparus	DVIVIP	WormBase ParaSite	PRJNA72587
Diploscapter coronatus	DCORON	WormBase ParaSite	PRJDB3143
Diploscapter pachys	DPACHY	WormBase ParaSite	PRJNA280107
Dirofilaria immitis	DIMMIT	WormBase ParaSite	PRJEB1797
Ditylenchus destructor	DDESTR	WormBase ParaSite	PRJNA312427
Dracunculus medinensis	DMEDIN	WormBase ParaSite	PRJEB500
Drosophilae melanogaster	DMELAN	Ensembl	BDGP6
Elaeophora elaphi	EELAPH	WormBase ParaSite	PRJEB502
Enterobius vermicularis	EVERMI	WormBase ParaSite	PRJEB503
Globodera pallida	GPALLI	WormBase ParaSite	PRJEB123
Globodera rostochiensis	GROSTO	WormBase ParaSite	PRJEB13504
Gongylonema pulchrum	GPULCH	WormBase ParaSite	PRJEB505
Paragordius varius	GORDSP	-	-
Haemonchus contortus	HCONTO	WormBase ParaSite	PRJEB506
Haemonchus placei	HPLACE	WormBase ParaSite	PRJEB509
Heligmosomoides bakeri	HBAKEI	-	PRJEB15396
Heterorhabditis bacteriophora	HBACTE	caenorhabditis.org	PRJNA13977
Hypsibius exemplaris	HEXEMP	tardigrades.org	PRJNA360553
Litomosoides sigmodontis	LSIGMO	WormBase ParaSite	PRJEB3075
Loa loa	LOALOA	WormBase ParaSite	PRJNA246086
Meloidogyne hapla	MHAPLA	WormBase ParaSite	PRJNA29083
Meloidogyne incognita	MINCOG	WormBase ParaSite	PRJEA28837
Mesorhabditis belari	MBELAR	caenorhabditis.org	PRJEB30104
Necator americanus	NAMERI	WormBase ParaSite	PRJNA72135
Nippostrongylus brasiliensis	NBRASS	WormBase ParaSite	PRJEB511
Oesophagostomum dentatum	ODENTA	WormBase ParaSite	PRJNA72579
Onchocerca volvulus	OVOLVO	WormBase ParaSite	PRJEB513
Oschieus tipulae	OTIPUL	caenorhabditis.org	PRJEB15512
Panagrellus redivivus	PREDIV	WormBase ParaSite	PRJNA186477
Parascaris equorum	PEQUOR	WormBase ParaSite	PRJEB514
Parascaris univalens	PUNIVA	WormBase ParaSite	PRJNA386823
Parastrongyloides trichosuri	PTRICH	WormBase ParaSite	PRJEB515
Plectus murrayi	PMURRA	ngenomes.org	-
Poikilolaimus oxycercus	POXYCE	caenorhabditis.org	-
Pristionchus exspectatus	PEXPEC	WormBase ParaSite	PRJEB6009
Pristionchus pacificus	PPACIF	WormBase ParaSite	PRJNA12644
Ramazzottius varieornatus	RVARIE	tardigrades.org	PRJDB1451
Rhabditophanes sp. KR3021	KR3021	WormBase ParaSite	PRJEB1297
Soboliphyme baturini	SBATUR	WormBase ParaSite	PRJEB516
Steinernema carpocapsae	SCARPO	WormBase ParaSite	PRJNA202318
Steinernema scapterisci	SSCAPT	WormBase ParaSite	PRJNA204942
Strongyloides ratti	SRATTI	WormBase ParaSite	PRJEB125

Species	Prefix	Source	BioProject
Strongyloides venezuelensis	SVENEZ	WormBase ParaSite	PRJEB530
Strongylus vulgaris	SVULGA	WormBase ParaSite	PRJEB531
Syphacia muris	SMURIS	WormBase ParaSite	PRJEB524
Teladorsagia circumcincta	TCIRCU	WormBase ParaSite	PRJNA72569
Tetranychus urticae	TURTI	WormBase ParaSite	PRJEA71041
Thelazia callipaeda	TCALLI	WormBase ParaSite	PRJEB1205
Toxocara canis	TCANIS	WormBase ParaSite	PRJEB533
Trichinella nativa	TNATIV	WormBase ParaSite	PRJNA179527
Trichinella spiralis	TSPIRA	WormBase ParaSite	PRJNA12603
Trichuris muris	TRMURI	WormBase ParaSite	PRJEB126
Trichuris suis	TRSUIS	WormBase ParaSite	PRJNA179528
Wuchereria bancrofti	WBANCR	WormBase ParaSite	PRJEB536

Table S3

Software	Version	Relevant parameters
softWorx	7.0.0	
Priism	4.7.1	
OrthoFinder	2.2.7	-og
MAFFT	7.407	--auto
IQ-TREE	1.6.10	-bb 1000, -bb -m GTR20+G
PhyloTreePruner	20150918	35 0.9 u
NCBI-BLAST+	2.5.0+	blastp
trimalAl	v1.4.rev15	-gt 0.8 -st 0.001 -resoverlap 0.75 -seqoverlap 80
catfasta2phyml	-	-c -f