The conspicuousness of the toxic *Heliconius* butterflies across time and

habitat

Denise Dalbosco Dell'Aglio[1,2*]

Jolyon Troscianko[3]

Martin Stevens[3]

W Owen McMillan[2]

Chris D Jiggins[1]

Affiliations

1. Butterfly Genetics Group, Department of Zoology, University of Cambridge,

Cambridge, United Kingdom

2. Smithsonian Tropical Research Institute, Panama City, Panama

3. Centre for Ecology and Conservation, College of Life and Environmental Sciences,

University of Exeter, Penryn, United Kingdom

* Corresponding author

E-mail: denise.ddd3@gmail.com

Abstract

Forests are a mosaic of light spectra, and colour signal efficiency might change in

different light environments. Local adaptation in *Heliconius* butterflies is linked to

microhabitat use and the colourful wing colour patterns may be adapted for signalling in

different light environments. These toxic butterflies exhibit conspicuous colours as a

warning to predators that they should be avoided, but also find and choose potential mates

based on colour signals. The two selection pressures of predation and mate preference are

therefore acting together. Colour conspicuousness should show habitat-specific contrast

for the butterflies, which would facilitate detection and species identification. On the

other hand, selection for signal stability would be stronger in the avian visual system. In

this study we analysed the contrast of two *Heliconius* mimicry rings in their natural

habitats under varying degrees of forest fragmentation and light conditions. We used

digital image analyses and mapped the bird and butterfly vision colour space in order to

examine whether warning colours have greater contrast and if they transmit a consistent

signal across time of the day and habitat in a tropical forest. We tested conspicuousness

using opponent colour channels against a natural green background. For avian vision,

colours are generally very stable through time and habitat. For butterfly vision, there is

some evidence that species are more contrasting in their own habitats, where

conspicuousness is higher for red and yellow bands in the border and for white in the

forest. Light environment affects *Heliconius* butterflies' warning signal transmission to a

higher degree through their own vision, but to a lesser degree through avian predator

vision. This work provides insight into the use of colour signals in sexual and natural

selection in the light of ecological adaptation.

Introduction

The success of a signal is related to its effectiveness in a specific environment and how strongly it influences the behaviour of the receiver (Endler 1978). Forests are a mosaic of light colours, and the same colour pattern can have an altered appearance in different light environments (Endler 1993). If an individual shows high reflectance of a specific wavelength, but the environment lacks light in that part of the spectrum, the region of high reflection will be unimportant as a signal (Stevens et al. 2007). Ambient light spectra also vary from dawn to dusk, hence species that signal only at certain times and places are expected to evolve characteristics and predictable combinations of colours for particular environments (Endler 1993). Therefore, ambient light characteristics should be included together with the receiver visual system to understand the microhabitat choice and behaviour of animals.

In the case of colour signals, perception can be altered by the habitat they live in by filtering wavelengths and altering visual backgrounds (Endler 1993; Lovell et al. 2005). Sensory drive explains the process of adaptation of signalling and sensory systems to the local environment (Endler 1992; Endler and Basolo 1998). Environment tuned spectral sensitivity is better known in aquatic habitats, such as in guppies (Endler 1980) and cichlid fish (Seehausen et al. 2008), as compared to terrestrial light environments. On land, colour depends on the reflection of the surroundings and has greater variability over time (Boughman 2002). Habitat signal transmission can favour diversification of mating signals through local adaptation, leading to reproductive isolation. Distinct *Anolis* lizards male dewlaps are found in different microhabitats (Fleishman et al. 1997). Male dewlap colours are more conspicuous in their own habitat than in other habitats, mainly because of the contrast against the background in the ultraviolet (UV) range (Leal and Fleishman 2002). Perception of colours in different light conditions can also influence attacks by predators, for example among butterflies in an environment with high UV light, birds aimed at the butterfly wings, more specifically the marginal white eyespots that have UV reflectance, instead of the head (Olofsson et al. 2010).

Local adaptation in *Heliconius* butterflies commonly involves adaptation to specific microhabitat use (Estrada and Jiggins 2002; Elias et al. 2008; Jiggins 2008). Mimicry rings are groups of unpalatable species that share the same warning colour, and these tend to be found in specific microhabitats such as forest or open areas. The *Heliconius* habitats are associated with the use of larval host-plants, adult food plants, sexual behaviour and gregarious roosting (Mallet and Gilbert 1995). Species that lay eggs

on *Passiflora* species that occur in second growth tend to be seen in open areas, while
species that lay eggs on canopy *Passiflora* vines are seen flying high in the forest. The
choice of microhabitat also might be connected with light differences between those
environments, such as the choice of using shady areas in communal roosting (Mallet and
Gilbert 1995; Finkbeiner 2014).

Therefore, different light environments should create microhabitats where
butterfly signals would be more efficient. Although mimicry rings differ in their
microhabitat, the light environment has not been measured to verify whether colour
patterns could be specifically adapted to particular light environments. The colourful
wing colours of *Heliconius* butterflies may also be subject to evolution caused by sensory
drive due to their potentially conflicting roles in predation and mate preference. Many
species exhibit Müllerian mimicry (Müller 1879), in which two or more species share the
same conspicuous colour as a warning to predators that they are toxic and should be
avoided (Benson 1972). Also, these butterflies find and choose potential mates based on
colour signals, which can lead to reproductive isolation (Jiggins et al. 2001; Sweeney et
al. 2003; Kronforst et al. 2006). Furthermore, communication between conspecifics might
be based on UV signals, since *H. erato* females express the duplicate UV opsin gene,
which allows a greater degree of discrimination of the UV-yellow wing patches (Briscoe
et al. 2010; Bybee et al. 2012; McCulloch et al. 2016; Dell'Aglio et al. 2018).

This microhabitat structuring allows mimicry rings to remain distinct. This may
be because there are sets of predators in different habitats, each of which perceive a
different mimicry ring as the most abundant pattern (Joron and Mallet 1998). Although
little is known of the specific predators that attack *Heliconius*, it seems likely that their
aposematic signals are directed at several predators with different visual abilities and
spectral sensitivities (Dell'Aglio et al. 2018). Ambient light together with predator
sensitivity can interfere with the interpretation of the information perceived from colour
signals.

Warning coloration should, therefore, be easy to detect and memorize even in
heterogeneous environments and light conditions (Guilford and Dawkins 1991; Endler
1992). Warning signals are often dominated by red, yellow and orange, frequently
contrasting with black, which are the main colours in *Heliconius*. The reason why these
long-wavelength colours are widely represented in aposematic coloration is that they are
highly conspicuous against natural backgrounds, are more stable across light conditions,
allowing long distance discrimination and detectability, and influence memorability

(Guilford and Dawkins 1991; Stevens and Ruxton 2012; Arenas et al. 2014; Dell'Aglio
et al. 2016).

Perception of colour depends on several neurophysiological mechanisms, such as
the presence of opponent colour channels. This chromatic mechanism involves
comparisons of receptors outputs, in which opposite neural pathways are either activated
or inhibited depending on the stimuli reaching the eye (Kelber et al. 2003; Renoult et al.
2015). This mechanism is useful especially regarding colour stability against spatial and
temporal variation in illumination (Lovell et al. 2005; Renoult et al. 2015). For example
birds, the major predator of aposematic butterflies, have tetrachromatic vision and
seemingly have at least three opponent channels, as found in domestic chicks (Vorobyev
et al. 1998; Osorio et al. 1999). Opponent channels have also been described for insects
(Chittka et al. 1992; Chittka 1996) and butterflies (Kelber 1999), and have been
hypothesized for *Heliconius* butterflies although more behavioural analyses are needed
to confirm which opponent channels are actually used (Swihart 1971, 1972; Bybee et al.
2012). Butterflies in the genus *Papilio* have duplicate LW opsin genes to see in the red
and green range (Kelber 1999; Briscoe 2008), while *Heliconius* has differences in the red
and green photoreceptor sensitivity associated with the presence of red filtering pigments
in the ommatidia (Zaccardi et al. 2006; McCulloch et al. 2016). Thus, we expect avian
predators and butterflies to rely on these high-contrast systems to process information
under a changing light environment.

The aim of this study was to analyse *Heliconius* warning colouration under
different light conditions in their natural habitats. In particular, to test conspicuousness of
the wing colouration against a natural green background, encoded by opponent colour
channels. Using digital image analyses, butterfly wings were photographed and mapped
to UVS and VS avian predator vision and to *Heliconius erato* vision (Figure 1). Our
predictions are that (1) signal contrast and conspicuousness for avian predators should
have stability, that should be reliable throughout the day and in different light
environments (Stevens and Ruxton 2012; Arenas et al. 2014). Warning signals might be
honest indicators of prey unprofitability to predators, and if signals fluctuate through the
day and between light environments, we would predict that this could delay learning by
predators and be costly to the prey. Similarly for internal contrasts (i.e. contrast between
black and the coloured bands), therefore conspicuousness would not rely totally on
background contrast but also on internal patterns which account for close-distance
conspicuousness (Endler 1978; Aronsson and Gamberale-Stille 2009). (2) From a

Heliconius butterfly perspective; we predict that signal contrast and conspicuousness

should show habitat-specific maximum background contrast and higher colour

differences in their own habitats (Table 1), which would facilitate detection and species

identification. We therefore predict that selection for signal stability will be much stronger

in the avian visual system as compared to the butterfly visual system.

Figure 1. Co-mimics used in this study. (a) *Heliconius erato demophoon*. (b) *H. melpomene rosina*. (c) *H. cydno*. (d) *H. sapho.*

Table 1. Colour patches for each co-mimic pair studied and typical microhabitats and light conditions where these co-mimics occur. Microhabitats descriptions are based on Estrada and Jiggins (2002) and light conditions based on Endler (1993) and personal observations.

Co-mimics	Colour	Microhabitat	Light conditions
H. erato demophoon & *H. melpomene rosina*	red & yellow	forest border	partial shade
		open area	no shade
H. sapho & *H. cydno*	white	closed forest	full shade
		canopy	partial shade

Material and Methods

Study site and species

Fieldwork was performed during the dry season, along Pipeline Road, a tropical

lowland rainforest in the Panama Canal Zone (Parque Nacional Soberanía, 9º7'33"N,

79º42'90"W). Pipeline Road makes a transect through the forest, creating a

heterogeneous habitat with open sunny areas and close canopy exceeding 30 m in height.

All specimens were collected in the area. Two pairs of co-mimics that live in sympatry

were selected, *H. erato demophoon* (n = 8) and *H. melpomene rosina* (n = 8), *H. sapho*

(n = 5) and *H. cydno* (n = 5), each pair belonging to a different mimicry ring, red and

yellow, and white, respectively (Figure 1 and Table 1).

Digital photography

The general approach and methodology for this work was based on previous work with colour stability using opponent signals (Lovell et al. 2005; Arenas et al. 2014). The spectral reflectance of mimetic pairs was investigated using digital photography. This provides a way to control for natural variation in luminance intensity (shadowing) that is not captured by spectrometry, and also allows non-invasive colour measurements easily applied in the field (Stevens et al. 2007; Troscianko and Stevens 2015). Therefore, through this method we could obtain colour measures under the sensitivity of all receiver photoreceptors (300-700 nm) in the actual viewing conditions of conspecifics and avian predators.

Fresh wings of each specimen were photographed following the same methods of image collection in Dell'Aglio et al. (2018). The camera was fitted to a tripod and pointed towards the ground (90º) at a height of approximately 80 cm. Each photo setup included two individuals, one of each species of the co-mimic pair, a 40% grey standard (Spectralon® Labsphere) used for calibration and a leaf freshly collected to make background measures. The species used was *Guazuma ulmifolia* (Sterculiaceae), a small abundant shrub across all Pipeline Road, which facilitated the collection of fresh leaves.

Photos were taken under three different arboreal canopy conditions, forest border, closed forest, and open area, where those butterflies are usually seen (Table 1). All photos were taken under sunny to part-cloudy days, with three replicates in each habitat making sure that the amount of light was similar. In order to standardize the replicates, light measures were taken with a digital light meter (Digital Lux meter, Tondaj LX-1010B), which measures the total amount of LW (555 nm) per square meter (Lux) (Figure 2). Also photographs of the canopy were taken in order to measure vegetative cover, which was 81.5% (SE ± 0.2) for closed forest, 59.8 % (SE ± 3.4) for forest border, and 0% for open area. The aim was to analyze how colour signals are perceived throughout the morning when butterflies are most active. Therefore, photos were taken at dawn (7 am), morning (9 am) and noon (12 pm) during a short period of 15min as light conditions change rapidly. Open area photographs were taken only at 12 pm to represent a highly used environment by butterflies at this period of day.

Figure 2. Amount of light differs across light environment and time of day. Showing the average log Lux between the three replicates of each habitat (± SE). Lux represents the amount of long-wave (555 nm) per square meter.

Image analysis and visual modelling

All images were processed and analysed into the imaging software ImageJ

(Rasband, 1997-2012). RAW human-visible and UV images were linearized and aligned

following the methodology of Troscianko and Stevens (2015) and Arenas et al. (2014).

Normally, photos would be normalized to the grey standard, which removes effects of

light conditions (Arenas et al. 2014). Since our main interest was to measure how

coloration changes in different light environments, the images were not normalized.

Instead, an average grey standard value was obtained from all photographs. Photon catch

values were obtained for each colour using the entire patch from linearized photos, and

subsequently these values were multiplied by each photo exposure time and normalized

with the average grey standard. With this methodology we were able to calculate how

particular environment and time varies from the natural average light (Stevens et al. 2007;

Arenas et al. 2014). We used the average photon catch results from the three habitats

replicates. Predicted photon catch values were obtained using spectral sensitivity for each

cone type of the blue tit (*Cyanistes caeruleus*) for the UV-sensitive vision (UVS) (Hart et

al. 2000), peafowl (*Pavo cristatus*) for the violet-sensitive vision (VS) (Hart 2002) and

the female *Heliconius erato* (Briscoe et al. 2010; McCulloch et al. 2016).

Background of many terrestrial habitats is dominated by greenish vegetation;

therefore, a green leaf was chosen to make contrast calculations. Differences between

light environment and time of the day were calculated using the contrast of warning

colours against an average green leaf. Channel activation in avian vision was calculated

using the Red-Green (RG), Blue-Yellow (BY) and Blue-UV opponent channel (Osorio et

al. 1999; Lovell et al. 2005; Stevens et al. 2009). For the achromatic signal, we used avian

double cones (DBL). Using a ratio-based approach suggested by Lovell et al. (2005), we

calculated the opponent channel responses as follows:

$$RG = LW – MW / LW + MW$$

$$BY = SW – (LW + MW) / SW + (LW + MW)$$

$$Blue\text{-}UV = SW – UV / SW + UV$$

$$Achromatic = DBL$$

The *Heliconius* opponent channels were calculated based on what has been

proposed in earlier studies regarding *Heliconius* vision (Swihart 1971, 1972; Bybee et al.

2012). The *H. erato* compound eye has red filtering pigments that shift LW photoreceptor

sensitivity from green to red and as the physiological mechanisms underlying these two

LW photoreceptors are not known, sensitivities in Green (560 nm) and Red (600 nm)

were used (McCulloch et al. 2016, 2017). To investigate differences between co-mimic

species, we calculated opponent channel activation based on the prediction that

Heliconius mating system might use UV and Red/Green light for mate choice (Bybee et

al. 2012; McCulloch et al. 2016, 2017). Opponent channels in *Heliconius* vision was

calculated using as follows:

$$RG = Red – Green / Red + Green$$

$$BY = Blue – (Red + Green) / Blue + (Red + Green)$$

$$Blue\text{-}UV2 = Blue – UV2 / Blue + UV2$$

$$UV2\text{-}UV1 = UV2 – UV1 / UV2 + UV1$$

To examine whether warning colours have greater contrast against green

background we calculated the Weber Contrast (Whittle 1994), which takes into account

the image value of the objects of interest as a fraction of background appearance using

the formula:

$$C = (object - background) / background$$

Where *background* corresponds to the green leaf opponent channel values, and

object corresponds to warning colour opponent channel values. This measure is suited to

comparisons between small objects against larger backgrounds, such as butterflies against

the green forest. For internal contrast, achromatic values of the warning colours were used

against the black of each individual wing as background (Arenas et al. 2014). We plotted

the mean absolute contrast of each colour signal as a function of time and light

environment for the three vision models.

To test contrast stability through time and habitat, we used the coefficient of

variation (CV) of each colour in each opponent channel for each visual system. The CV

is an effective measurement to determine how relatively stable a measurement is around

a mean value. The CV can be interpreted as greater its value, more variance and instability

the signal would be, following methodology from Arenas et al. (2014).

Statistical analyses

All statistical calculations were processed in the software R 3.5.3 (R Core Team

2020). Our approach was to model colour contrasts over the course of a day and under

different habitats in term of both predator and butterfly vision. General linear mixed-

effects models were performed using random effects (package *lme4*) followed by Tukey's

post-hoc (package *multicomp*). Normality tests showed that contrast data were not

normally distributed, therefore *glmer* tests were performed using *Gama* distribution. Raw

data was plotted to illustrate the results (package *ggplot2*). The models were fitted

accordingly to the predictions outlined above. Analyses were carried out using contrast

values as the dependent variable, and fixed and random factors varied depending on the

question. Factors were individuals, colour (red, yellow, white), habitat (border, forest,

area), time (7am, 9am, 12pm), and bird vision (UVS, VS). For *Heliconius* vision, we also

added side of the wing (dorsal, ventral).

Results

Signal contrast and conspicuousness for avian predators

Red was generally the most contrasting colour against a green background in the

RG opponent channel as compared to yellow ($z = 14.05$, $P < 0.001$, Table S1) and white

($z = 17.68$, $P < 0.001$, Table S1). In contrast, white had higher contrasts against a green

background in the BY channel, as compared to red ($z = -15.3$, $P < 0.001$, Table S1) and

yellow (z = 22.04, $P < 0.001$, Table S1) (Figure 3). Colours in open areas showed a higher

contrast, such in the RG channel for red band (t = -6.16, $P < 0.001$, Table S2) with no

difference between border and forest (z = -0.31, $P = 0.94$, Table S2) (Figure 3). In the

Blue-UV opponent channel, UVS and VS birds could perceive red and yellow with less

stability (Table 2), as red and yellow showed higher contrast in the morning than at 12

pm (Table S3). On the other hand, white colour in the Blue-UV channel did not show any

changes through the day (Table S3).

Internal achromatic contrast was higher for yellow, compared to red (z = -36.7, P

< 0.001, Table S1) and white (z = -22.9, $P < 0.001$, Table S1). Moreover, yellow had

more contrast in the border (Figure 3), which is the preferred habitat of yellow band

butterflies, than in the forest (z = 3.5, $P = 0.001$, Table S4).

Signal contrast and conspicuousness for Heliconius conspecifics

In some cases, contrasts followed our prediction that species would be more

contrasting in their own habitats (Figure 4). The yellow colour was more contrasting in

the border in the UV2-UV1 channel, especially early hours such as 7 am (t = 6.9, $P <$

0.001, Table S5) and 9 am (t = 354.3, $P < 0.001$, Table S5), but not at 12 pm (t = 2.2, $P =$

0.218, Table S5). In the Blue-UV2 channel, white was more contrasting in the forest than

in the border at 7 am (t = -3.9, $P < 0.001$, Table S5). Although, while white colour contrast

decreased during the day in the forest, it increased at 12 pm in the border (t = 4.77, $P <$

0.001, Table S5) (Figure 4).

The red colour showed large differences in the RG channel between dorsal and

ventral side, with dorsal side with the higher contrast (t = 10.1, $P < 0.001$, Table S6).

Same results were found for red in the Blue-Yellow and Blue-UV2 opponent channel

(Table S6) (Figure 4).

The contrast of signals against the background revealed greater differences across

habitat and time when seen through the *Heliconius* vision model (Table 2). *Heliconius*

vision had higher values of coefficient of variation, which revealed greater fluctuations

across habitat and time in most of the opponent channels. The only exception was in the

Blue-UV opponent channel for red and yellow colour, which showed more instability for

avian vision than for *Heliconius* vision (Table 2).

Figure 3. Colour conspicuousness for avian predators. Mean absolute contrast of colour signals (±SE, standard error) in the bird vision systems analyzed (circles, UVS; triangles, VS) through habitats (red, border; green, forest; blue, open) and time (7am, 9am, 12pm). Vertical panels show the three colour signals (red, yellow and white), horizontal panels show opponent channels against green leaf (top, Red-Green; middle, Blue-Yellow and Blue-UV) and against the black of the wing (bottom, Achromatic). Note: Channels have different y-axis values. Error bars smaller than data points are not shown.

Figure 4. Colour conspicuousness for *Heliconius* conspecifics. Mean absolute contrast of colour signals against green leaf in *Heliconius* vision through habitats (red, border; green, forest; blue, open), time (7 am, 9 am, 12 pm) and side of the wing (circles, dorsal; triangles, ventral). Vertical panels show colour signals (red, yellow and white), horizontal panels show opponent channels against green leaf (top, Red-Green; middle, Blue-Yellow and Blue-UV2; bottom, UV2-UV1). Error bars: ± 1 standard error (SE), error bars smaller than data points are not shown.

Table 2. *Heliconius* vision has higher coefficient of variation in most of the colours compared with bird vision across time and habitats. Coefficient of variation for each visual system, colour and opponent channel, higher values are in bold.

Opponent channel	Colour	Visual system		
		Heliconius	UVS	VS
Red-Green				
	Red	**1.053**	0.704	0.695
	Yellow	**0.384**	0.211	0.178
	White	**0.722**	0.375	0.314
Blue-Yellow				
	Red	**0.699**	0.576	0.606
	Yellow	**0.735**	0.667	0.528
	White	**0.177**	0.134	0.169
Blue-UV				
	Red	0.783	0.897	**1.024**
	Yellow	0.265	**0.788**	0.637
	White	**1.035**	0.549	0.341
UV2-UV1				
	Red	0.525	-	-
	Yellow	1.047	-	-
	White	1.061	-	-
Achromatic				
	Red	-	0.367	0.436
	Yellow	-	0.717	0.699
	White	-	0.348	0.349

Discussion

The bright and contrasting *Heliconius* wing patterns appear well adapted for
signalling distastefulness to predators. However, their colour constancy and appearance
in different light environments remains poorly studied. Here we have shown that colours
are indeed very stable for avian predator vision, but somewhat less so for *Heliconius*
vision. This is consistent with the idea that wing patterns are primarily selected for their
role in signalling distastefulness to predators.

Signal stability and conspicuousness to avian predators

Generally, warning signals involve combinations of long-wavelength colours such
as red, orange, and yellow which are highly conspicuous against natural backgrounds and
stable under different natural conditions (Lovell et al. 2005; Stevens and Ruxton 2012).
In our results, *Heliconius* red colouration has higher detectability against average green
background in the RG output and these results are consistent regardless of habitat and
time of the day. Previous work has investigated colour stability through opponent colour

channels and also showed that red coloration is more contrasting and stable against green backgrounds over the course of a day and across light conditions to the bird visual system (Lovell et al. 2005; Arenas et al. 2014). Moreover, the *Heliconius* yellow colouration also is highly conspicuous against its internal black pattern in the achromatic output. Achromatic information is one of the main cues used for motion detection (Hämäläinen et al. 2015). Therefore, our results suggest that red and yellow signals work together and are likely effective in stimulating avian opponent channels in order to be conspicuous in all light environments.

There is evidence that red and yellow colouration serve as reliable warning signal to avian predators (Ham et al. 2006; Svádová et al. 2009; Arenas et al. 2015), but that this is less true of white colouration. One explanation is that white is more variable across time and habitat, so provides a less reliable signal under varied light conditions (Stevens and Ruxton 2012; Arenas et al. 2014). As an example, field and aviary experiments with polymorphic yellow and white wood tiger moths, *Parasemia plantaginis*, showed that yellow males are avoided more than white males by predators, but white males have higher mating success (Nokelainen et al. 2012). Our results showed that white contrasts against green background were lower and rather variable for avian vision. The co-mimics *H. sapho* and *H. cydno* also contain iridescence blue that was not measured with this methodology. However, the lack of high contrast in white colouration might be balanced with the fact that polarized light might act as a signal, especially in forest habitats (Sweeney et al. 2003; Douglas et al. 2007; Pegram et al. 2015).

Highly conspicuous warning signals are expected to evolve to be stable in their appearance throughout the day and between light environments, in order to remain honest indicators of prey unpalatability (Blount et al. 2009; Cortesi and Cheney 2010; Stevens and Ruxton 2012; Arenas et al. 2015). If warning signals fluctuate through time and space this could alter bird foraging experiences and reduce the effectiveness of the aposematic signal. The final decision on whether or not to attack a prey results from a combination of information reaching the predator brain, and for greater efficiency, aposematic coloration needs to be easy to remember (Endler 1988). Our results support this prediction, as colours were generally stable through time and light environments in all opponent systems with only a few exceptions. Notably these occurred where contrasts were higher in open areas and in the early morning. This might also be favourable as the prey would be more conspicuous when they are most vulnerable to predation, since birds are more active and forage early in the morning (Buskirk et al. 1972; Poulin et al. 2001;

376 Steiger et al. 2009). In agreement with this, *Heliconius* predation and roost disturbance
377 has been observed in the early morning (Mallet 1986; Finkbeiner 2014).
378
Habitat and time influence conspicuousness in Heliconius conspecifics
Butterflies belonging to the two mimetic rings studied here tend to be segregated
between habitats, corresponding to areas where the photographs were taken, although
there is considerable overlap (Estrada and Jiggins 2002). We showed that the colours
were more unstable when seen through *Heliconius* vision as compared to avian vision and
some colours tend to be more contrasting in their respective habitats.
Our results provide some evidence that co-mimic rings are more conspicuous in
their own habitat as seen through *Heliconius* vision, reinforcing the idea that ecological
adaptation leads to spatial segregation to where detection would be facilitated. Some
colours had higher contrast against green backgrounds in their respective habitat, such as
yellow in the border and white in the forest. Nonetheless, red showed the opposite trend
and was generally more contrasting in the forest. Differences across light environments
could affect mating preferences by altering search costs for a specific colour pattern, and
perhaps changing the fitness of different colour patterns. Adaptation in different
microhabitats within the forest might have an influence on how closely related species
commonly differ in pattern, while convergence in pattern occurs between more distantly
related species (Joron and Mallet 1998). Ecological adaptation is attributed to habitat
preference and leads to assortative mating (Jiggins 2008). The two sister species studied
here, *H. melpomene rosina* and *H. cydno*, are known to rarely hybridise in the wild, hence
microhabitat segregation reduces potential mating encounters between these two species
and reduces gene flow (Mallet et al. 1998; Merrill et al. 2013). Subtle environmental
conditions could affect recognition in mating behaviour as seen in the jumping spider,
Habronattus pyrrithrix, which red males were more successful in approaching females in
the sunlight (Taylor and McGraw 2013).
The activation of opponent channels was often higher in the early hours of the
morning, at the time when the butterflies are more active and leave their roost or perches
to forage (Mallet 1986; Finkbeiner et al. 2012). This was especially the case for
Heliconius white and yellow wing colours in the UV channel, which might act in
intraspecific communication (Briscoe et al. 2010; Bybee et al. 2012). There is evidence
that distinct UV colour signals are being transmitted between co-mimics, which may
reduce costs of mating confusion (Dell'Aglio et al. 2018). Similarly, in two species of

newt, belly colour is distinct in the UV range and females often made mistakes choosing

the wrong males in the absence of UV light (Secondi and Théry 2014).

In this context, the duplicate genes encoding two distinct visual pigments with

sensitivity peaks in the UV range in *H. erato* females offer the potential for enhanced

spectral discrimination in light environments and time of the day where UV is more

prominent. A UV2-UV1 opponent channel was proposed by Bybee et al. (2012), who

showed that this receptor combination would have lower error rates for discrimination

between *Heliconius* and *Dryas* yellows. There is no direct evidence for such a mechanism

yet, but the fact that males and females show differences in the expression of the two UV

proteins suggests that UV2-UV1 contrasts could be an important opponent channel for a

female specific behaviour, perhaps mate recognition or host plant finding (McCulloch et

al. 2016, 2017). On the other hand, there is some evidence for a Red-Green opponent

channel seen in behavioural trials, which showed that *Heliconius erato* use both green

and red photoreceptor to detect colours in the 590-640 nm range (Zaccardi et al. 2006)

and the presence of red filters in their eyes means that this is a possibility (McCulloch et

al. 2016). Differences between species in Red-Green channel activity for red colouration

might have a role in mate recognition since *Heliconius* tend to be attracted to red (Merrill

et al. 2011).

The sensory drive hypothesis describes evolutionary relationships among visual

systems, conditions of the light environment and mating preferences (Endler and Basolo

1998). *Heliconius* mating preference is highly linked to colour and in *H. melpomene*, the

gene responsible for red colour pattern is genetically linked to the preference for the same

pattern (Jiggins et al. 2001; Naisbit et al. 2001; Merrill et al. 2011). Visual sensitivity data

used here is only from *H. erato* females whereas it might differ for other species in their

visual systems and perhaps match with colour preference or habitat (Frentiu et al. 2007;

Briscoe et al. 2010; McCulloch et al. 2017). In addition, mating behaviour might benefit

from some habitats in maximizing conspicuousness, such as in tropical dwelling birds

and wire-tailed manakins which visual contrast is increased during display by habitat

choice (Endler and Théry 1996; Heindl and Winkler 2003). Nevertheless, our results

suggest that selection for conspicuousness in the preferred habitat could explain in part

the divergence in colour pattern in these species.

Conclusion

443   In conclusion, the transmission of *Heliconius* warning signals varies due to light
environment to a much greater degree through their own visual system, but to a smaller
degree through avian predator vision. Selection for signal detectability under different
habitat conditions is a mechanism that is proposed to lead to evolution of signal diversity,
as seen in species of *Anolis* lizards that occupy habitats that match their visual system and
signal design (Leal and Fleishman 2002), in species of warblers which different cone
opsin gene expression correlate with sexual selection and habitat use (Bloch 2015) and
also colour patterns of guppies are more conspicuous to guppies at the times and places
of courtship and relatively less conspicuous at times and places of predator risk (Endler
1991). *Heliconius* butterfly warning colours are highly contrasting against the forest
background and stable through time and habitat in terms of predator avoidance but also
conspicuous to attract the attention of conspecifics. However, more extensive studies
considering spectral sensitivities of different *Heliconius* species and their responses to
environmental changes in their signal visibility are needed to confirm the
conspicuousness to mates. Opponent channel colour contrasts can predict behaviour of
perceivers, however, additional behavioural experiments on how light environment
influences prey detectability, such with poison frogs (Rojas et al. 2014), are necessary to
verify our results.

Funding
This study was supported by Smithsonian Tropical Research Institute, Cambridge Trust,
and CAPES Brazil (9423/11-7) to D.D.D, by European Research Council (Speciation
Genetics 339873) to C.D.J., and by Biotechnology and Biological Sciences Research
Council David Phillips Research Fellowship (BB/G022887/1) to M.S.

Author contributions
D.D.D. collected all the data, analysed, and wrote the manuscript; J.T. developed digital
image methodology and data analysis; W.O.M, M.S, and C.D.J conceived the study,
edited, and wrote the manuscript. We have no conflict of interest to declare.

References

Arenas, L. M., J. Troscianko, and M. Stevens. 2014. Color contrast and stability as key
elements for effective warning signals. Front. Ecol. Evol. 2:1–12.

Arenas, L. M., D. Walter, and M. Stevens. 2015. Signal honesty and predation risk
among a closely related group of aposematic species. Sci. Rep. 5:11021.

Aronsson, M., and G. Gamberale-Stille. 2009. Importance of internal pattern contrast
and contrast against the background in aposematic signals. Behav. Ecol. arp141.

Benson, W. W. 1972. Natural Selection for Mullerian Mimicry in Heliconius erato in
Costa Rica. Science 176:936–939.

Bloch, N. I. 2015. Evolution of opsin expression in birds driven by sexual selection and
habitat. Proc. R. Soc. B 282:20142321.

Blount, J. D., M. P. Speed, G. D. Ruxton, and P. A. Stephens. 2009. Warning displays
may function as honest signals of toxicity. Proc. R. Soc. Lond. B 276:871–877.

Boughman, J. W. 2002. How sensory drive can promote speciation. Trends Ecol. Evol.
17:571–577.

Briscoe, A. D. 2008. Reconstructing the ancestral butterfly eye: focus on the opsins. J.
Exp. Biol. 211:1805–1813.

Briscoe, A. D., S. M. Bybee, G. D. Bernard, Y. Furong, M. P. Sison-Mangus, R. D.
Reed, A. D. Warren, J. Llorente-bousquets, and C. Chiao. 2010. Positive selection
of a duplicated UV-sensitive visual pigment coincides with wing pigment evolution
in Heliconius butterflies. PNAS 107:3628–3633.

Buskirk, W. H., G. V. N. Powell, J. F. Wittenberger, R. E. Buskirk, and T. U. Powell.
1972. Interspecific Bird Flocks in Tropical Highland Panama. Am. Ornithol. Union
89:612–624.

Bybee, S. M., F. Yuan, M. D. Ramstetter, J. Llorente-Bousquets, R. D. Reed, D. Osorio,
and A. D. Briscoe. 2012. UV Photoreceptors and UV-Yellow Wing Pigments in
Heliconius Butterflies Allow a Color Signal to Serve both Mimicry and
Intraspecific Communication. Am. Soc. Nat. 179:38–51.

Chittka, L. 1996. Optimal Sets of Color Receptors and Color Opponent Systems for
Coding of Natural Objects in Insect Vision. J. Theor. Biol. 181:179–196.

Chittka, L., W. Beier, H. Hertel, E. Steinmann, and R. Menzel. 1992. Opponent colour
coding is a universal strategy to evaluate the photoreceptor inputs in Hymenoptera.
J. Comp. Physiol. A 170:545–563.

Cortesi, F., and K. L. Cheney. 2010. Conspicuousness is correlated with toxicity in
marine opisthobranchs. J. Evol. Biol. 23:1509–1518.

Dell'Aglio, D. D., M. Stevens, and C. D. Jiggins. 2016. Avoidance of an aposematically
coloured butterfly by wild birds in a tropical forest. Ecol. Entomol. 41:627–632.

Dell'Aglio, D. D., J. Troscianko, W. O. McMillan, M. Stevens, and C. D. Jiggins. 2018.
The appearance of mimetic Heliconius butterflies to predators and conspecifics.
Evolution 72:2156–2166.

Douglas, J. M., T. W. Cronin, T. Chiou, and N. J. Dominy. 2007. Light habitats and the
role of polarized iridescence in the sensory ecology of neotropical nymphalid
butterflies (Lepidoptera: Nymphalidae). J. Exp. Biol. 210:788–799.

Elias, M., Z. Gompert, C. D. Jiggins, and K. Willmott. 2008. Mutualistic Interactions
Drive Ecological Niche Convergence in a Diverse Butterfly Community. PLoS
Biol. 6:e300.

Endler, J. A. 1978. A predator's view of animal color patterns. Pp. 319 – 364 *in*
Evolutionary Biology.

Endler, J. A. 1988. Frequency-dependent predation, crypsis and aposematic coloration.
Philos. Trans. R. Soc. B Biol. Sci. 319:505–523.

Endler, J. A. 1980. Natural Selection on Color Patterns in Poecilia reticulata. Evolution
34:76–91.

Endler, J. A. 1992. Signals, Signal Conditions, and the Direction of Evolution. Am. Nat.
139:S125–S153.

Endler, J. A. 1993. The color of light in forests and its implications. Ecol. Monogr.
63:1–27.

Endler, J. A. 1991. Variation in the appearance of guppy color patterns to guppies and
their predators under different visual conditions. Vision Res. 31:587–608.

Endler, J. A., and A. L. Basolo. 1998. Sensory ecology, receiver biases and sexual
selection. Trends Ecol. Evol. 13:415–420.

Endler, J. A., and M. Théry. 1996. Interacting Effects of Lek Placement, Display
Behavior, Ambient Light, and Color Patterns in Three Neotropical Forest-Dwelling
Birds. Am. Nat. 148:421–452.

Estrada, C., and C. D. Jiggins. 2002. Patterns of pollen feeding and habitat preference
among Heliconius species. Ecol. Entomol. 27:448–456.

Finkbeiner, S. D. 2014. Communal roosting in heliconius butterflies (nymphalidae):
roost recruitment, establishment, fidelity, and resource use trends based on age and
sex. J. Lepidopterists Soc. 68:10–16.

Finkbeiner, S. D., A. D. Briscoe, and R. D. Reed. 2012. The benefit of being a social
butterfly: communal roosting deters predation. Proc. R. Soc. B 279:2769–2776.

Fleishman, L. J., M. Bowman, D. Saunders, W. E. Miller, M. J. Rury, and E. R. Loew.
1997. The visual ecology of Puerto Rican anoline lizards: Habitat light and spectral
sensitivity. J. Comp. Physiol. A 181:446–460.

Frentiu, F. D., G. D. Bernard, M. P. Sison-Mangus, A. Van Zandt Brower, and A. D.
Briscoe. 2007. Gene duplication is an evolutionary mechanism for expanding
spectral diversity in the long-wavelength photopigments of butterflies. Mol. Biol.
Evol. 24:2016–2028.

Guilford, T., and M. S. Dawkins. 1991. Receiver psychology and the evolution of
animal signals. Anim. Behav. 42:1–14.

Ham, A. D., E. Ihalainen, L. Lindström, and J. Mappes. 2006. Does colour matter? The
importance of colour in avoidance learning, memorability and generalisation.
Behav. Ecol. Sociobiol. 60:482–491.

Hämäläinen, L., J. Valkonen, J. Mappes, and B. Rojas. 2015. Visual illusions in
predator–prey interactions: birds find moving patterned prey harder to catch. Anim.
Cogn. 18:1059–1068.

Hart, N. S. 2002. Vision in the peafowl (Aves: Pavo cristatus). J. Exp. Biol. 205:3925–
3935.

Hart, N. S., J. C. Partridge, I. C. Cuthill, and A. T. D. Bennett. 2000. Visual pigments,
oil droplets, ocular media and cone photoreceptor distribution in two species of
passerine bird: the blue tit (Parus caeruleus L.) and the blackbird (Turdus merula
L.). J. Comp. Physiol. A 186:375–387.

Heindl, M., and H. Winkler. 2003. Interacting effects of ambient light and plumage
color patterns in displaying wire-tailed manakins (Aves, Pipridae). Behav. Ecol.
Sociobiol. 53:153–162.

Jiggins, C. D. 2008. Ecological Speciation in Mimetic Butterflies. BioScience 58:541–
548.

Jiggins, C. D., R. E. Naisbit, R. L. Coe, and J. Mallet. 2001. Reproductive isolation
caused by colour pattern mimicry. Nature 411:302–305.

Joron, M., and J. Mallet. 1998. Diversity in mimicry: paradox or paradigm? Trends
Ecol. Evol. 13:461–466.
Kelber, A. 1999. Ovipositing butterflies use a red receptor to see green. J. Exp. Biol.
202:2619–2630.
Kelber, A., M. Vorobyev, and D. Osorio. 2003. Animal colour vision – behavioural
tests and physiological concepts. Biol. Rev. 78:81–118.
Kronforst, M. R., L. G. Young, D. D. Kapan, C. McNeely, R. J. O'Neill, and L. E.
Gilbert. 2006. Linkage of butterfly mate preference and wing color preference cue
at the genomic location of wingless. PNAS 103:6575–6580.
Leal, M., and L. J. Fleishman. 2002. Evidence for habitat partitioning based on
adaptation to environmental light in a pair of sympatric lizard species. Proc. R. Soc.
Lond. B 269:351–359.
Lovell, P. G., D. J. Tolhurst, C. A. Párraga, R. Baddeley, U. Leonards, J. Troscianko,
and T. S. Troscianko. 2005. Stability of the color-opponent signals under changes
of illuminant in natural scenes. J. Opt. Soc. Am. 22:2060–2071.
Mallet, J. 1986. Gregarious roosting and home range in Heliconius butterflies. Natl.
Geogr. Res. 2:198–215.
Mallet, J., and L. E. Gilbert. 1995. Why are there so many mimicry rings? Correlations
between habitat, behaviour and mimicry in Heliconius butterflies. Biol. J. Linn.
Soc. 55:159–180.
Mallet, J., W. O. Mcmillan, and C. D. Jiggins. 1998. Estimating the Mating Behavior of
a Pair of Hybridizing Heliconius Species in the Wild. Evolution 52:503–510.
McCulloch, K. J., D. Osorio, and A. D. Briscoe. 2016. Sexual dimorphism in the
compound eye of Heliconius erato: a nymphalid butterfly with at least five spectral
classes of photoreceptor. J. Exp. Biol. 219:2377–2387.
McCulloch, K. J., F. Yuan, Y. Zhen, M. L. Aardema, G. Smith, J. Llorente-Bousquets,
P. Andolfatto, and A. D. Briscoe. 2017. Sexual Dimorphism and Retinal Mosaic
Diversification following the Evolution of a Violet Receptor in Butterflies. Mol.
Biol. Evol. 34:2271–2284.
Merrill, R. M., R. E. Naisbit, J. Mallet, and C. D. Jiggins. 2013. Ecological and genetic
factors influencing the transition between host-use strategies in sympatric
Heliconius butterflies. J. Evol. Biol. 26:1959–1967.

Merrill, R. M., B. Van Schooten, J. A. Scott, and C. D. Jiggins. 2011. Pervasive genetic
associations between traits causing reproductive isolation in Heliconius butterflies.
Proc. R. Soc. B 278:511–518.

Müller, F. 1879. Ituna and Thyridia; a remarkable case of mimicry in butterflies. Trans.
Linn. Soc. Lond. 1879:xx–xxix.

Naisbit, R. E., C. D. Jiggins, and J. Mallet. 2001. Disruptive sexual selection against
hybrids contributes to speciation between Heliconius cydno and Heliconius
melpomene. Proc. R. Soc. B 268:1849–1854.

Nokelainen, O., R. H. Hegna, J. H. Reudler, C. Lindstedt, and J. Mappes. 2012. Trade-
off between warning signal efficacy and mating success in the wood tiger moth.
Proc. R. Soc. B 279:257–265.

Olofsson, M., A. Vallin, S. Jakobsson, and C. Wiklund. 2010. Marginal Eyespots on
Butterfly Wings Deflect Bird Attacks Under Low Light Intensities with UV
Wavelengths. PLoS ONE 5:e10798.

Osorio, D., M. Vorobyev, and C. D. Jones. 1999. Colour vision of domestic chicks. J.
Exp. Biol. 202:2951–2959.

Pegram, K. V., H. A. Han, and R. L. Rutowski. 2015. Warning Signal Efficacy:
Assessing the Effects of Color, Iridescence, and Time of Day in the Field. Ethology
121:1–13.

Poulin, B., G. Lefebvre, R. Ibáñez, C. Jaramillo, C. Hernández, and A. S. Rand. 2001.
Avian predation upon lizards and frogs in a neotropical forest understorey. J. Trop.
Ecol. 17:21–40.

R Core Team. 2020. R: A Language and Environment for Statistical Computing. R
Foundation for Statistical Computing, Vienna, Austria.

Renoult, J. P., A. Kelber, and H. M. Schaefer. 2015. Colour spaces in ecology and
evolutionary biology. Biol. Rev., doi: 10.1111/brv.12230.

Rojas, B., P. Rautiala, and J. Mappes. 2014. Differential detectability of polymorphic
warning signals under varying light environments. Behav. Processes 109:164–172.

Secondi, J., and M. Théry. 2014. An ultraviolet signal generates a conflict between
sexual selection and species recognition in a newt. Behav. Ecol. Sociobiol.
68:1049–1058.

Seehausen, O., Y. Terai, I. S. Magalhaes, K. L. Carleton, H. D. J. Mrosso, R. Miyagi, I.
van der Sluijs, M. V. Schneider, M. E. Maan, H. Tachida, H. Imai, and N. Okada.
2008. Speciation through sensory drive in cichlid fish. Nature 455:620–627.

Steiger, S. S., J. P. Kelley, W. W. Cochran, and M. Wikelski. 2009. Low Metabolism

and Inactive Lifestyle of a Tropical Rain Forest Bird Investigated via Heart-Rate

Telemetry. Physiol. Biochem. Zool. 82:580–589.

Stevens, M., C. A. Párraga, I. C. Cuthill, J. C. Partridge, and T. S. Troscianko. 2007.

Using digital photography to study animal coloration. Biol. J. Linn. Soc. 90:211–

237.

Stevens, M., and G. D. Ruxton. 2012. Linking the evolution and form of warning

coloration in nature. Proc. R. Soc. B 279:417–426.

Stevens, M., M. C. Stoddard, and J. P. Higham. 2009. Studying Primate Color: Towards

Visual System-dependent Methods. Int. J. Primatol. 30:893–917.

Svádová, K., A. Exnerová, P. Štys, E. Landová, J. Valenta, A. Fučíková, and R. Socha.

2009. Role of different colours of aposematic insects in learning, memory and

generalization of naïve bird predators. Anim. Behav. 77:327–336.

Sweeney, A., C. D. Jiggins, and S. Johnsen. 2003. Polarized light as a butterfly mating

signal. Nature 423:31–32.

Swihart, C. A. 1971. Colour discrimation by the butterfly, Heliconius charitonius Linn.

Anim. Behav. 19:156–164.

Swihart, S. L. 1972. The neural basis of colour vision in the butterfly, Heliconius erato.

J. Insect Physiol. 18:1015–1025.

Taylor, L. a., and K. J. McGraw. 2013. Male ornamental coloration improves courtship

success in a jumping spider, but only in the sun. Behav. Ecol. 24:955–967.

Troscianko, J., and M. Stevens. 2015. Image Calibration and Analysis Toolbox – a free

software suite for objectively measuring reflectance, colour and pattern. Methods

Ecol. Evol. 6:1320–1331.

Vorobyev, M., D. Osorio, A. T. D. Bennett, N. J. Marshall, and I. C. Cuthill. 1998.

Tetrachromacy, oil droplets and bird plumage colours. J. Comp. Physiol. A

183:621–633.

Whittle, P. 1994. The psychophysics of contrast brightness. Pp. 35–110 *in* A. L.

Gilchrist, ed. Lightness, Brightness, and Transparency. Lawrence Erlbaum

Associates, Hillsdale, NJ.

Zaccardi, G., A. Kelber, M. P. Sison-mangus, and A. D. Briscoe. 2006. Color

discrimination in the red range with only one long-wavelength sensitive opsin. J.

Exp. Biol. 209:1944–1955.

Supplementary Information

Table S1. Colour contrast differences for avian vision per opponent channel. Generalized linear mixed-
effects models with random effects and Tukey's post-hoc (y ~ Colour + (1 | Individuals) + (1 |
Vision:Habitat:Time)).

Opponent channel	Colour	N	t value	p-value	Tukey's post-hoc		
					Contrasts	z value	p-value
Red-Green							
	Red (intercept)	1176	14.17	< 2E-16	White - Red	17.68	< 2E-16
	White		17.68	< 2E-16	Yellow - Red	14.05	< 2E-16
	Yellow		14.06	< 2E-16	Yellow - White	-9.09	< 2E-16
Blue-Yellow							
	Red (intercept)	1176	23.66	< 2E-16	White - Red	-15.30	<2E-16
	White		-15.3	< 2E-16	Yellow - Red	10.55	<2E-16
	Yellow		10.55	< 2E-16	Yellow - White	22.04	<2E-16
Blue-UV							
	Red (intercept)	1176	8.93	< 2E-16	White - Red	8.63	< 1E-05
	White		8.63	< 2E-16	Yellow - Red	-4.78	< 1E-05
	Yellow		-4.78	1.72E-06	Yellow - White	-11.43	< 1E-05
Achromatic							
	Red (intercept)	1176	33.53	< 2E-16	White - Red	-17.41	< 2E-16
	White		-17.41	< 2E-16	Yellow - Red	-36.76	< 2E-16
	Yellow		-36.76	< 2E-16	Yellow - White	-22.97	< 2E-16

690 **Table S2.** Habitat contrast differences for avian vision per colour for the Red-Green opponent channel.
691 Generalized linear mixed-effects models with random effects and Tukey's post-hoc (y ~ Habitat + (1 |
692 Individuals) + (1 | Vision:Time)).

Colour	Habitat	N	Linear mixed model		Tukey's post-hoc		
			t value	p-value	Contrasts	z value	p-value
Red							
	Border (intercept)	448	16.59	<2E-16	Forest - Border	-0.31	0.947
	Forest		-0.31	0.753	Open - Border	-6.16	<1E-05
	Open		-6.16	7.28E-10	Open - Forest	-5.91	<1E-05
Yellow							
	Border (intercept)	448	23.56	<2E-16	Forest - Border	-8.46	<2E-16
	Forest		-8.46	<2E-16	Open - Border	-16.69	<2E-16
	Open		-16.69	<2E-16	Open - Forest	-10.65	<2E-16
White							
	Border (intercept)	280	9.07	<2E-16	Forest - Border	2.76	0.015
	Forest		2.76	0.006	Open - Border	-12.39	0.001
	Open		-12.39	<2E-16	Open - Forest	-14.39	0.001

693
694
695
696

697   **Table S3.** Time contrast differences for avian vision per colour for the Blue-Yellow opponent channel.

698   Generalized linear mixed-effects models with random effects and Tukey's post-hoc (y ~ Time + (1 |

699   Individuals) + (1 | Vision:Habitat)).

Colour	Time	N	Linear mixed model		Tukey's post-hoc		
			t value	p-value	Contrasts	z value	p-value
Red							
	12PM (intercept)	448	12.73	<2E-16	7AM - 12PM	-5.11	<1E-06
	7AM		-5.11	3.2E-07	9AM - 12PM	-5.39	<1E-06
	9AM		-5.39	6.7E-08	9AM - 7AM	-0.33	0.939
Yellow							
	12PM (intercept)	448	8.94	<2E-16	7AM - 12PM	-10.38	<1E-04
	7AM		-10.38	<2E-16	9AM - 12PM	-11.64	<1E-04
	9AM		-11.64	<2E-16	9AM - 7AM	-2.05	0.092
White							
	12PM (intercept)	280	10.6	<2E-16	7AM - 12PM	0.4	0.909
	7AM		0.41	0.677	9AM - 12PM	-0.23	0.970
	9AM		-0.23	0.813	9AM - 7AM	-0.64	0.797

703 **Table S4.** Habitat contrast differences for avian vision for the achromatic opponent channel per colour.
704 Generalized linear mixed-effects models with random effects and Tukey's post-hoc (y ~ Habitat + (1 |
705 Individuals) + (1 | Vision:Time)).

Colour	Habitat	N	Linear mixed model		Tukey's post-hoc		
			t value	p-value	Contrasts	z value	p-value
Red							
	Border (intercept)	448	8.19	2.49E-16	Forest - Border	2.94	0.008
	Forest		2.94	0.003	Open - Border	6.30	0.001
	Open		6.30	2.93E-10	Open - Forest	4.64	0.001
Yellow							
	Border (intercept)	448	8.85	<2E-16	Forest - Border	3.52	0.001
	Forest		3.52	4.3E-04	Open - Border	5.75	0.001
	Open		5.75	8.5E-09	Open - Forest	4.07	0.001
White							
	Border (intercept)	280	20.06	<2E-16	Forest - Border	2.12	0.083
	Forest		2.12	0.034	Open - Border	-0.33	0.940
	Open		-0.33	0.741	Open - Forest	-1.77	0.174

726 **Table S5.** Habitat contrast differences for *Heliconius* vision for yellow and white colours, per opponent
727 channel and time. Generalized linear mixed-effects models with random effects and Tukey's post-hoc (y ~
728 Habitat + (1 | Individuals) + (1 | Side)).

| Colour | Channel | Time | Habitat | N | Linear mixed model | | Tukey's post-hoc | | |
					t value	p-value	Contrasts	z value	p-value
Yellow	UV2-UV1	7AM							
			Border (intercept)	64	3.40	0.001			
			Forest		6.90	4.9E-12			
		9AM							
			Border (intercept)	64	117.9	<2e-16			
			Forest		354.3	<2e-16			
		12PM							
			Border (intercept)	96	5.95	2.7E-09	Forest - Border	-1.23	0.425
			Forest		2.20	0.218	Open - Border	-4.70	1E-04
			Open		7.49	2.6E-06	Open - Forest	-3.91	2.8E-04
White	Blue-UV2	7AM							
			Border (intercept)	40	5.37	7.7E-08			
			Forest		-3.99	6.5E-05			
		9AM							
			Border (intercept)	40	6.07	1.2E-09			
			Forest		-1.54	0.123			
		12AM							
			Border (intercept)	60	3.97	7.1E-05	Forest - Border	4.77	1E-04
			Forest		4.77	1.8E-06	Open - Border	4.29	1E-04
			Open		4.29	1.7E-05	Open - Forest	-0.75	0.724

744 **Table S6.** Side of the wing contrast differences for *Heliconius* vision for the red colour, per opponent
745 channel. Generalized linear mixed-effects models with random effects and Tukey's post-hoc (y ~ Side + (1
746 | Individuals) + (1 | Habitat:Time)).

Opponent Channel	Side of the wing	N	Linear mixed model	
			t value	p-value
Red-Green				
	Dorsal (intercept)	224	6.16	7.2E-10
	Ventral		10.10	< 2E-16
Blue-Yellow				
	Dorsal (intercept)	224	6.21	5.3E-10
	Ventral		12.08	< 2E-16
Blue-UV2				
	Dorsal (intercept)	224	15.91	< 2E-16
	Ventral		12.90	< 2E-16
UV2-UV1				
	Dorsal (intercept)	224	10.58	< 2E-16
	Ventral		-1.71	0.086

747