Title: Case-control design identifies ecological drivers of endemic coral diseases

Jamie M. Caldwell[1,2*], Greta Aeby[3], Scott F. Heron[2,4,5], and Megan J. Donahue[1]

1. Hawaii Institute of Marine Biology, University of Hawaii at Manoa, Hawaii,

USA

2. ARC Centre of Excellence for Coral Reef Studies, James Cook University,

Townsville, Australia

3. Qatar University, Department of Biological & Environmental Sciences, Doha,

Qatar

4. Marine Geophysical Laboratory, Physics, College of Science and

Engineering, James Cook University, Townsville, Australia

5. NOAA Coral Reef Watch, College Park, Maryland, USA

* contact: jamie.sziklay@gmail.com

Abstract

Endemic disease transmission is an important ecological process that is

challenging to study because of low occurrence rates. Here, we investigate the

ecological drivers of two coral diseases -- growth anomalies and tissue loss --

affecting five coral species. We first show that a statistical framework called the

case-control study design, commonly used in epidemiology but rarely applied to

ecology, provided high predictive accuracy (67-82%) and disease detection rates

(60-83%) compared with a traditional statistical approach that yielded high

accuracy (98-100%) but low disease detection rates (0-17%). Using this

framework, we found evidence that 1) larger corals have higher disease risk; 2)

shallow reefs with low herbivorous fish abundance, limited water motion, and

located adjacent to watersheds with high fertilizer and pesticide runoff promote

low levels of growth anomalies, a chronic coral disease; and 3) wave exposure,

stream exposure, depth, and low thermal stress are associated with tissue loss

disease risk during interepidemic periods. Variation in risk factors across host-

disease pairs suggests that either different pathogens cause the same gross

lesions in different species or that the same disease may arise in different

species under different ecological conditions.

Introduction

Disease is an ecologically and evolutionarily important process in shaping

populations, communities, and ecosystems [1–5], but identifying factors that

promote disease at low endemic levels or between epidemics is challenging

because disease occurrences are rare in space and time. Disease can affect

populations, communities, and ecosystems through multiple pathways, such as

changing the physiology, behavior, distribution, abundance, and fitness of

organisms (e.g., [3,6,7]). For example, when the trematode *Plagioporous sp.*

infects the coral *Porites compressa* it reduces coral growth and causes infected

polyps to appear as bright swollen nodules that cannot retract into their skeletal

cups, thus increasing fish predation and affecting coral physiology, behavior, and

fitness [8]. As another example, infection in three-spined sticklebacks by the fish

parasite *Gyrodactylus* spp. has been shown to alter the zooplankton community

structure and nutrient cycling in streams and lakes, which in turn, affect the

survival and fitness of the subsequent fish generation [5]. Despite the ecological

and evolutionary importance of disease, understanding the factors that increase

individual disease risk for endemic diseases with low prevalence or during

interepidemic periods for diseases with epidemic cycles can be challenging

because of the low probability of observing diseased individuals during non-

outbreak periods.

With limited occurrence data relative to nonoccurrence data, commonly used

biostatistics such as logistic regression can significantly underestimate the

probability of rare events such as disease occurrence [9]. Further, such models

are likely to have high predictive accuracy (proportion of correctly identified event

and nonevent observations), which can be misleading: for example, if 3 of 100

individuals are diseased, a model where all observations are predicted healthy

would return 97% predictive accuracy but provide very limited information about

disease occurrence or causation. Ecologists interested in rare events such as

insect outbreaks and species invasions could benefit from adopting statistical

approaches from fields of inquiry concerned with rare phenomena. One statistical

framework that works well with low event-to-nonevent ratio datasets is called the

case-control study design [10]. The case-control study design is commonly used

in epidemiology where the number of people affected by a condition is very small

relative to the potential control population (e.g., few people are HIV+ compared

with the general population). In the case-control design, for each subject with

some condition, a control subject is selected that does not have the condition but

is otherwise similar in terms of factors such as age, sex, and occupation. This

approach naturally lends itself to many ecological studies with low event-to-

nonevent ratios such as wildlife disease occurrence, with some modifications. In

particular, the case-control design is typically applied to cohorts of individuals

who are followed through time. For wildlife diseases, cohort data are rarely

available. Instead, we propose that wildlife studies can apply the case-control

design by retrospectively sub-sampling cross-sectional data that are widely

available.

Here, we compare logistic regression models with and without a case-control

design (based on retrospective sub-sampling) to answer three ecological

questions for marine wildlife diseases: 1) how accurately can we predict disease

occurrence; 2) what are the ecological factors associated with maintaining low

levels of disease occurrence; and 3) are risk factors more similar for closely

related species? We used coral diseases for this case study because (i) they can

be extremely rare (<0.1% of disease occurrence observations); (ii) lesions can be

visually identified; (iii) they can cause high mortality or strongly reduce fecundity

with likely long-term population effects; and (iv) large datasets are publicly

available on coral health observations and hypothesized risk factors.

In this study, we investigate two types of coral diseases – growth anomalies and

tissue loss diseases – affecting five species of coral from two widely distributed

genera in the Indo-Pacific Ocean. Growth anomalies are chronic, protuberant

masses of coral skeletons (i.e., tumors) that reduce growth, fecundity, and

survival [11–14]. Growth anomalies in some coral genera (e.g. *Porites* spp.) have

been associated with larger coral colonies [15] located in shallow water (<3 m)

[12] with elevated nutrients [16] and in regions with high human populations [17].

Tissue loss diseases, or white syndromes, usually have very low endemic

prevalence levels (<1% in the Indo-Pacific) [18] but can be associated with rapid

outbreak events resulting in localized mass mortality [19–23]. This suite of

diseases is characterized by progressive tissue loss across the coral colony with

lesions progressing slowly (chronic to subacute) or rapidly (acute). Bacterial

pathogens cause many tissue loss diseases [24–27] and can be associated with

high coral cover (i.e., density dependent transmission) and temperature stress

(e.g. anomalously warm or cold ocean temperatures) [28–32]. Based on these

previous findings, we hypothesize that long-term, chronic stressors are more

likely to drive growth anomaly persistence whereas locations with pulses of short-

term, acute stressors are more likely to be associated with interepidemic tissue

loss occurrence. Growth anomalies and tissue loss appear on numerous coral

species and likely constitute multiple diseases. Thus, we investigate each host-

disease pair separately.

Results

Comparison between statistical approaches

Logistic regression models with and without a case-control design had high

overall predictive accuracy (proportion of correctly predicted healthy and disease

observations) for all host-disease pairs, but only the models with the case-control

design accurately predicted disease occurrence (proportion of disease

observations the model correctly predicted as diseased; i.e., the true positive

rate). Using colony health state as a binary response variable (0 indicating

healthy, 1 indicating diseased; Table 1) and a suite of hypothesized risk factors

(Tables 2-3), we found that mean predictive accuracies ranged from 98-100% for

logistic regression without the case-control design and 67-82% for logistic

regression with the case-control design (Fig. 1). True positive rate, or the

proportion of disease observations the model correctly predicted as diseased, is

of greatest interest for rare event prediction and mean values ranged from 0-17%

for logistic regression without the case-control design and 60-83% for logistic

regression with the case-control design (Fig. 1). Given the moderate-to-high

mean predictive accuracies (67-82%) and mean true positive rates (60-83%) for

the case-control design, we report only results using the case-control design

below.

Table 1: Healthy observations far exceeded disease observations for all

host-disease pairs. Each row corresponds to the total number of disease and

healthy colony observations used in this analysis for a specific disease

(GA=growth anomalies, TL=tissue loss) and host (species or genus) pair. Hosts

at the genus level include observations from all hosts species within that genus

(e.g., *Montipora* spp. includes all observations from *Montipora capitata* and

Montipora patula). In the analysis, we used 80% of observations (training data) to

create the models and the remaining 20% of observations (test data) for model

validation. We calculated accuracy and true positive rates based on the test data.

There was an equal number of disease observations in the test data for logistic

regression with and without the case-control design. In contrast, the number of

healthy observations differed between the test data for logistic regression with

and without the case control design: in the case-control design, the number of

healthy observations was equal to the number of disease observations whereas

without the case-control design the number of healthy observations in the test

data was equal to ~20% of the total number of healthy observations.

Disease type	Host	Total disease	Total healthy
GA	*Montipora capitata*	49	35864
GA	*Montipora patula*	35	23118
GA	*Montipora* spp.	84	58982
GA	*Porites compressa*	160	17889
GA	*Porites evermanni*	42	1779
GA	*Porites lobata*	1396	64312
GA	*Porites* spp.	1598	83980
TL	*Montipora capitata*	39	28669
TL	*Montipora patula*	12	17480
TL	*Montipora* spp.	51	46149
TL	*Porites compressa*	89	16915
TL	*Porites evermanni*	17	1567
TL	*Porites lobata*	351	61489
TL	*Porites* spp.	457	79971

152 **Table 2. Risk factors (predictor variables) included in candidate models for each disease type with the direction**

153 **and mechanism(s) of hypothesized relationships.** Relationships and their proposed mechanisms described previously

154 in the literature are referenced whereas hypothesized relationships based on personal observations are not referenced.

155 References followed by an asterisk indicate relationships found for other type(s) of coral disease and/or related risk

156 factors.

Risk factor	Hypothesized relationship with disease risk	Hypothesized mechanism(s) relating risk factor to disease risk	Disease Type
Colony size	Positive	• Larger corals have more tissue exposed to pathogens • Larger, older corals have a longer duration of exposure to pathogens/stress • Larger, older corals have reduced immune capacity	Growth anomalies [15] Tissue loss [33]
Host density	Positive Negative	• Disease spread increases among many, closely spaced colonies • Disease spread is reduced in high density reefs because colonies are smaller	Growth anomalies [17] Tissue loss [33,56]
Depth	Negative	• Multiple environmental conditions become less stressful to coral with increasing depth: deeper water absorbs more light, has cooler temperatures, and fewer re-suspended sediments	Growth anomalies [12,29] Tissue loss [29,56]
Water temperature (Hot Snap)	Positive Negative	• High temperature reduces coral immune capacity • High temperature increases pathogen survival, growth rate, and/or virulence • High temperature reduces pathogen survival, growth rate, and/or virulence	Growth anomalies [29,57] Tissue loss [28–30]
Human population size	Positive	• People increase reef degradation and physical damage to coral	Growth anomalies [17,58] Tissue loss [59]
Embayment	Positive	• Areas inside bays have poor water quality (e.g., higher levels of nitrogen, sediments, and land-based pollutants) and are more likely to retain pathogens	Growth anomalies [15] Tissue loss [19]
Herbivore fish abundance	Positive Negative	• Fish alter algae composition to favor algal species that harbor pathogens • Fish spread pathogens via defecation on corals • Fish reduce macroalgae that harbor pathogens • Fewer fish indicate higher fishing intensity and associated impacts	Growth anomalies Tissue loss [18] [60–63]*
Agriculture and	Positive	• Pesticides and fertilizers increase coral stress and susceptibility to infection	Growth anomalies

golf course runoff			[41,64]*
Irradiance	Positive	• Light increases oxidative stress in coral	Growth anomalies [58]
Wave energy (chronic)	Negative	• Locations with high water flow reduce temperature and light stress to coral	Growth anomalies [57]
Wave energy (acute)	Positive Negative	• Pulses of water flow increase contact rate between corals and pathogens • Pulses of water flow limit duration of coral exposure to pathogens • Pulses of water flow increase coral abrasion via sediment or physical damage	Tissue loss
Stream exposure	Positive	• Stream runoff increases contact rate between corals and pathogens • Stream runoff reduces coral immune capacity by decreasing salinity and increasing sedimentation and exposure to land-based pollutants	Tissue loss [65]
Rainfall anomaly	Positive	• Rainfall reduces coral immune capacity by decreasing salinity and increasing land-based runoff of pathogens or pollutants • Rainfall increases contact rate between corals and pathogens	Tissue loss [66]*
Chlorophyll-*a* (proxy for nutrients and phytoplankton)	Positive	• High chlorophyll-*a* concentration increases coral stress and susceptibility to infection • High chlorophyll-*a* concentration leads to increased zooplankton, which can serve as disease vectors	Tissue loss [18]

 Table 3: Characterization of biological, environmental, physical, and anthropogenic risk factors (predictor

 variables) used in candidate models.

Risk factor	Ecological category	Characteristic	Description	Minimum value	Maximum value	Source
Colony size	Biological	Site	Maximum diameter or average size of size class (cm).	<1	300	Survey data[A]
Host density	Biological	Site	Number of colonies divided by survey area (m^{-2}).	0.02	9.49	Survey data[A]
Herbivore fish abundance	Biological	Site	Pooled estimate of herbivorous fish calculated between 2012 and 2013 (g m^{-2}).	0.06	0.42	[67]
Chlorophyll-a	Environmental	Chronic	Maximum anomaly calculated within a 30-meter boundary of the coast between 2003 and 2013 from NASA MODIS satellite imagery (mg m^{-3}).	0.0004	0.3566	[68][B]
Irradiance	Environmental	Chronic	Photosynthetically active radiation (PAR) average annual frequency of anomalies calculated within a 30-meter boundary of the coast between 2003 and 2013 from NASA MODIS satellite imagery (mol m^{-2} day^{-1}).	0.13	0.27	[68][B]
Hot Snap	Environmental	Acute	Magnitude and duration of heat stress above a summertime threshold at 5 km resolution (°C-week).	0	3.6	NOAA CRW[C]
Rainfall	Environmental	Acute	Monthly rainfall anomaly interpolated from rain gauge data, expert knowledge, radar observations, meteorological model simulations and vegetation data. Anomalies were calculated as the rainfall variation in any given month and year from the 1978-2007 mean.	0.04	4.35	[69][D]
Depth	Physical	Site	Distance below sea surface (m).	0.6	26.0	Survey data[A]
Embayment	Physical	Site	Binary variable indicating whether survey site is outside or inside of semi-enclosed habitat.	0	1	Calculated in this study
Stream exposure	Physical	Acute	Inverse planar distance between survey site and stream mouth in nearest watershed (m^{-1}).	2.72	3.33	Calculated in this study
Wave energy	Physical	Acute	Average annual maximum anomaly of wave energy flux calculated between 2000 and 2013 (kW m^{-1}).	0.08	285.74	[68][B]
Wave energy	Physical	Chronic	Average annual maximum monthly mean of wave energy flux calculated between 2000 and 2013 (kW m^{-1}).	0.04	103.48	[68][B]
Agricultural and golf	Anthropogenic	Chronic	Proxy for fertilizer and chemicals like pesticides and herbicides discharge from agricultural land and golf	0.00	0.53	[70]

course runoff			courses calculated as area of cultivated land and golf course per watershed (km^2) with a distance-decay function.			
Human population size	Anthropogenic	Chornic	Number of people living within 1 km of coastline nearest to survey site.	0	4425	NASA SEDAC GPW v4[E]

Public datasets:
A. Hawaiʻi Coral Disease database: https://www.sciencedirect.com/science/article/pii/S2352340916304607
B. Ocean Tipping Points: http://www.pacioos.hawaii.edu/projects/oceantippingpoints/#data
C. NOAA Coral Reef Watch CoralTemp: https://coralreefwatch.noaa.gov/satellite/coraltemp.php
D. Rainfall Atlas of Hawaiʻi: http://rainfall.geography.hawaii.edu/
E. NASA SEDAC Gridded Population of the world v4: http://sedac.ciesin.columbia.edu/data/collection/gpw-v4

Figure 1: Logistic regression with case-control design has higher disease

detection rates compared with logistic regression without case-control

design. Boxplots of accuracy (percent correctly predicted healthy and disease

observations) and true positive rates (percent correctly predicted disease

observations) across 500 randomly sampled test datasets (see Methods) for

each host-disease pair included in this study. The top panel shows results for

logistic regression models without the case-control design and the bottom panel

shows results for logistic regression models with the case-control design. The left

side of the panels show accuracy results and the right side of the panels show

true positive rates. Boxplots are grouped by disease type and colored by host;

models at the genus level include observations from all hosts species within that

genus (e.g., *Montipora* spp. includes all observations from *Montipora capitata*

and *Montipora patula*).

Ecological risk factors of different host-disease pairs

Of the 19 risk factors (predictor variables) we initially examined (see Methods),

14 were included in our candidate models (with the risk factors differing by

disease type; Table 2), but the most predictive risk factors varied by host-disease

pair. The 14 risk factors can be broadly classified into four categories: biological

(colony size, host density, and herbivorous fish abundance), environmental

(chlorophyll-*a*, irradiance, Hot Snap, and rainfall), physical (depth, embayment,

acute and chronic wave exposure, and stream exposure), and anthropogenic

(agricultural and golf course runoff and human population size).

All the risk factors that we hypothesized could be associated with growth

anomalies were identified as important for at least one host-disease pair, with

more similar risk factors associated with more closely related hosts. We found

one biological risk factor (colony size; Fig. 2) common to all hosts, with an

approximate 36-137% increase in disease risk of for every 1 cm increase in

colony size above the host specific average size, given all other risk factors are

held at their average values (Table 4). Herbivore fish abundance was the next

most common risk factor, with an approximate 5-15% decrease in disease risk

for every gram increase in herbivorous fish per meter squared (Table 4).

Montipora species were positively associated with size, and negatively

associated with herbivore fish abundance (*M.* capitata) or host density (*M.*

patula); the genus model (*Montipora* spp.) consisted of all three biological risk

factors (Table 4). In contrast, the *Porites* genus model (*Porites* spp.) only partially

overlapped with the risk factors identified in the species-specific models (Table

4). *P. compressa* was associated with all risk factors other than irradiance,

whereas *P. evermanni* was only associated with irradiance (in addition to colony

size) (Table 4). *P. lobata* was associated with biological (host density and

herbivore fish abundance), environmental (Hot Snap), physical (depth), and

anthropogenic (agriculture and golf course runoff) risk factors, and most closely

overlapped with the genus model (Table 4), likely because the majority of *Porites*

observations were observations of *P. lobata* (Table 1). Agriculture and golf

course runoff increased disease risk for *P. lobata* and *P. compressa* by

approximately 23 and 80%, respectively, for every km^2 increase in cultivated

land.

Table 4: All hypothesized risk factors were associated with at least one

host-disease pair for growth anomalies. Average coefficient values based on

model runs across 500 case-control datasets for all host-disease pairs.

Coefficient values indicate the magnitude and direction for the expected change

in disease risk given a one-unit change in the risk factor, holding all other risk

factors at their average values. Since the slope of the logistic curve is steepest at

its center, coefficient values can be divided by 4 ("divide by 4 rule") to reasonably

approximate the upper bound of the predicted difference in disease risk

corresponding to a one-unit difference in the risk factor [71].

Host	Intercept	Size	Density	Fish	Hot Snap	PAR	Depth	Bay	Wave exposure	Ag & Golf	Human pop.
M. capitata	1.92	5.46		-0.60							
M. patula	0.49	2.40	-0.59								
Montipora spp.	1.00	3.65	-0.50	-0.47							
P. compressa	-0.82	1.22	-1.64	-0.26	-5.54		0.65	3.07	-0.80	3.18	-0.42
P. evermanni	4.05	2.39				-1.93					
P. lobata	-0.58	1.34		-0.19	-6.99		-0.62			0.93	
Porites spp.	0.56	1.43	0.13	-0.24	-6.31		-0.50				

Figure 2: Larger colonies increase disease risk for most host-disease pairs.

Lines indicate the expected probability of disease occurrence given colony size

for a) growth anomalies and b) tissue loss. *M. capitata* is not shown in b because

colony size is not a risk factor in the *M. capitata* tissue loss model (Table 5).

Lines are colored by host species. Using the same color scheme, average colony

size is indicated along the horizontal axis (*Montipora capitata* = 16 cm; *Montipora*

patula = 19 cm; *Porites compressa* = 23 cm; *Porites evermanni* = 22 cm; *Porites*

lobata = 15 cm). Fig. S1 shows individual plots for each host-disease pair

(excluding genus models for visibility) with data points overlaid for one randomly

sampled case-control training dataset.

For tissue loss diseases, we identified eight hypothesized risk factors associated

with disease occurrence (five associated with *Montipora* species and seven

associated with *Porites* species) (Table 5). Like growth anomalies, we found a

positive association with colony size for most hosts with an approximate 33-

102% increase in disease risk for every 1 cm increase in colony size (Table 5,

Fig. 2). We found no overlap in risk factors for *Montipora capitata* and *Montipora*

patula: *M. capitata* was associated with one biological (herbivore fish abundance)

and three physical (depth, stream exposure, and wave exposure) risk factors

whereas *M. patula* was only associated with one biological (colony size) risk

factor (Table 5). Further, all risk factors for *M. capitata* had strong negative

effects (Table 5). We identified the exact same risk factors for the *Porites* genus

model and the *P. lobata* model (Table 5), which included biological (colony size,

host density), environmental (Hot Snap), physical (depth, wave exposure), and

anthropogenic (human population size) risk factors. However, Hot Snap had a

stronger effect in the genus model compared with the *P. lobata* model, potentially

indicating differences in thermal sensitivities across morphologies, as the

coefficients for Hot Snap in the genus model and the *P. compressa* model were

similar (Table 5). *P. evermanni* overlapped with four of those risk factors (colony

size, depth, wave exposure, and human population size). However, the effect of

wave exposure differed by host: higher wave exposure was associated with an

increased risk of tissue loss in *Porites* spp. and *P. lobata* but reduced risk of

tissue loss in *P. evermanni* (Table 5). The *P. compressa* model was the most

distinct of the *Porites* models, with associations with colony size, Hot Snap, and

stream exposure (Table 5).

Table 5: Tissue loss associated with different biological, environmental,

physical, and anthropogenic risk factors across host-disease pairs.

Average coefficient values based on model runs across 500 case-control

datasets for all host-disease pairs. Coefficient values indicate the magnitude and

direction for the expected change in disease risk given a one-unit change in the

risk factor, holding all other risk factors at their average values. Since the slope of

the logistic curve is steepest at its center, coefficient values can be divided by 4

("divide by 4 rule") to reasonably approximate the upper bound of the predicted

difference in disease risk corresponding to a unit difference in the risk factor [71].

Host	Intercept	Size	Density	Fish	Hot Snap	Depth	Stream exposure	Wave exposure	Human pop.
M. capitata	-0.18			-1.96		-1.48	-1.77	-1.40	
M. patula	0.69	2.53							
Montipora spp.	0.60	2.71		-1.28			-0.77	-0.76	
P. compressa	-2.37	1.45			-3.49		0.43		
P. evermanni	6.84	4.07				-1.52		-1.34	-2.01
P. lobata	-8.45	1.32	-0.39		-1.42	-0.23		0.22	-0.25
Porites spp.	-3.69	1.42	-0.29		-4.71	-0.23		0.16	-0.27

Discussion

Understanding the ecological conditions that promote endemic diseases with low

prevalence and for epidemic diseases during interepidemic periods is central for

understanding disease dynamics but can be difficult to investigate using common

biostatistical approaches. In this study, we adopted statistical methods better

suited towards highly skewed occurrence data to investigate ecological

correlates of several coral diseases. We show that the case-control study design

improved our ability to predict disease and healthy colony observations with

relatively high specificity given a suite of ecological conditions. In contrast, a

more traditional biostatistical approach yielded high overall predictive accuracy

(heavily biased by successful predictions of healthy state) but often failed to

accurately predict the events of interest: disease occurrences. This study

highlights the usefulness of the case-control design for ecological questions

using infrequent occurrence data. In this study, we identified biological,

environmental, physical, and anthropogenic drivers that promote low levels of

coral disease and explored how those factors differ across host-disease pairs.

These results provide insight into factors that should be explored further through

experimental and observational studies and that may be favorable for

intervention strategies.

Among the suite of ecological risk factors that we included in our study, we found

that disease risk was consistently higher for larger colonies across host-disease

pairs. A positive size-disease relationship has been found in other coral disease

studies [15,33,34] and could be associated with exposure area or duration of

exposure to potential disease causing agents (biotic or abiotic), or immune

function, which is hypothesized to decline with age [35]. Further, a positive

relationship between disease prevalence and coral cover has been well

established for infectious diseases like tissue loss [13,28,30,31]. However,

because the relationship between coral size-frequency distributions and coral

cover is not straightforward [36,37], it is unknown whether disease prevalence is

higher in locations with high coral cover because of many small colonies (many

potential susceptible hosts and/or reservoir species) or fewer large colonies

(higher susceptibility of certain individuals). Our results provide evidence that

highly susceptible corals (i.e., large colonies) drive the positive relationship

between disease prevalence and coral cover. Although we found a positive linear

relationship between disease risk and size within the range of 0 to 300 cm, it is

possible that disease risk saturates or is reduced above some species-specific

size threshold (Fig. 2). Regardless of what the functional form of this relationship

looks like across the entire range of colony sizes, the consistent importance of

size across different host-disease pairs (effects ranging from 32 to 137%

increase in disease risk) provides biological evidence for evaluating disease risk

at the colony scale (also supported by simulation models [38]) rather than

evaluating prevalence at the transect or site scale as is currently common

practice. Such probabilistic models of colony health state could easily be

integrated into a hierarchical framework to predict local or regional disease

prevalence.

In addition to colony size, our findings suggest that shallow reefs with low

herbivorous fish abundance, limited water motion, and located adjacent to

watersheds with high fertilizer and pesticide runoff promote low levels of chronic

disease year-round. We found that growth anomalies were more common in

reefs with lower herbivorous fish abundance, potentially indicating indirect

relationships with macroalgal cover that can harbor pathogens [39] or with more

degraded reefs with high fishing intensity and associated impacts. Further, we

found that growth anomalies were more likely to occur in shallow water, in reefs

with low host density (likely because as colony sizes increase, host density

decreases), and inside bays and other areas with low wave exposure (possibly

because of longer residence times and associated concentration of pollutants).

Growth anomalies in *P. compressa* and *P. lobata* were more common in

locations with elevated levels of pesticide and fertilizer runoff from agriculture and

golf courses (increasing disease risk by 23-80%, Table 4) and support previous

research associating disease risk to nutrient enrichment, human populations, and

poor water quality [17,40,41]. This research indicates that reducing coastal

development and high-intensity land use activities such as farming and golf

adjacent to semi-protected reefs would be an effective management approach to

break transmission cycles of growth anomalies.

For tissue loss, we found that wave exposure, stream exposure, depth, and

thermal stress were commonly associated with disease risk during interepidemic

periods, but their precise effects sometimes differed across hosts. Tissue loss

was consistently more common in shallow reefs with low thermal stress exposure

(Table 5). The strong negative association with anomalously warm thermal stress

in the summer for *P. compressa, P. lobata*, and *Porites* spp. models appears

contrary to several previous studies on tissue loss in other regions [28,30,31,42].

Although there are no documented outbreaks of *Porites* tissue loss in Hawaii, this

result may shed some light on why several tissue loss outbreaks in *Montipora* in

Hawaii occurred in the winter [19,33] compared with other regions where

outbreaks occurred in the summer [30,31,43]. This result may suggest low

temperature stress restricts the tissue loss pathogens' growth rate, enabling the

pathogen(s) to persist during interepidemic periods without causing an outbreak.

Alternatively, disease risk could be highest in winter months because of a time

lag between exposure to heat stress and increased host susceptibility to disease.

We found that wave exposure and stream exposure were important risk factors

for tissue loss as well, but the direction and magnitude of these relationships

varied by host. For example, wave exposure was negatively associated with

tissue loss in *M. capitata, Montipora* spp.*,* and *P. evermanni* but positively

associated with tissue loss in *P. lobata and Porites* spp. (Table 5). Similarly,

stream exposure was negatively associated with *M. capitata* and *Montipora* spp.

but positively associated with *P. compressa*. We did not find evidence that these

opposing patterns were driven by differences in habitat associations across hosts

(Figs. S2-3)*.* One hypothesis that would explain this result is that water

movement may be important for transmitting pathogens for some hosts (e.g., *P.*

lobata and *Porites* spp.) but more stagnant water where pathogens can

proliferate may be more likely to promote disease in other hosts (e.g., *M.*

capitata, Montipora spp.*,* and *P. evermanni*). Alternatively, the association with

water movement may be related to water diffusion processes as colonies

exposed to high water flow minimize the diffusion boundary layer [44] and could

therefore limit exposure to pathogens in areas with high water movement and

vice versa. Colony morphology can also affect water diffusion and pathogen

exposure as branching morphologies (common in *Montipora* spp.) are more likely

to retain water within their branches compared with massive colonies that tend to

have smooth surfaces (common in *Porites* spp.). Similarly, stream exposure and

potentially associated changes in salinity, sedimentation, and/or land-based

pollutant runoff may promote disease occurrence in some hosts (e.g., *P.*

compressa) and decrease disease risk in other hosts (e.g., *M. capitata* and

Montipora spp.).

The variation in risk factors across host-disease pairs suggests that there may be

different pathogens causing the same gross lesions in different hosts or that the

same disease may arise in different hosts under different ecological conditions.

Corals have a limited repertoire of responses to insult; therefore, disease lesions

that are grossly identical, such as tissue loss, could be caused by different

pathogens of the same general type (e.g., bacterial pathogens), which has been

suggested in other studies [45]. Recent research supports this hypothesis as

multiple bacterial pathogens have been identified for tissue loss in *Montipora*

capitata in Hawaii [46]. Alternatively, the same causative agent(s) could be

triggered to infect different species under different conditions. Our understanding

of coral disease ecology is still limited but our results suggest that there may be

specific ecological conditions that support or restrict the spread of specific

diseases in each host. Thus, researchers should strongly consider how

combining data for multiple species to assess disease risk may impact our

understanding of factors that promote transmission. For example, combining

species into a single model may be reasonable if the researchers acknowledge

that the models will perform best for reefs that reflect the composition of species

used in model development.

This study demonstrates the usefulness of the case-control design to study rare

events in space and time using existing datasets with low probability of

occurrence, but also provides a path forward for more efficiently collecting new

data for future investigations. For example, for studies specifically focused on

coral disease, we recommend complementing more traditional ecological

assessments that characterize overall reef health, such as belt transect surveys

and rugosity measurements, with survey efforts that maximize information on

disease observations. Using experimental designs to collect data geared towards

rare and/or clustered populations would allow for targetted investigations into the

underlying mechanisms that drive disease processes. Therefore, for coral

diseases, collecting more extensive information such as size, morphology,

geographic location, clustering with nearby diseased colonies, severity of

infection, area of affected tissue, etc. allows the use of modeling techniques to

better explore the ecology of specific diseases. Such survey designs and

associated analyses have already been developed and widely used in other

fields such as epidemiology, plant pathology, and conservation biology (e.g.,

case-control design [47], cluster sampling [48,49], line transect surveys [50]) and

could be adapted for coral disease and other ecological studies. Rare occurrence

or abundance data is very common in ecology and approaching ecological

questions with this statistical design in mind will allow ecologists to more

rigorously quantify ecological relationships for infrequent disturbances.

Methods

Coral health observations

We used coral health observations from the Hawaii Coral Disease Database

(HICORDIS) [51] and additional surveys collected by the authors following the

same methodologies used in HICORDIS (e.g., 10 x 1 m^2 belt transects surveys)

resulting in 362,366 coral health observations collected between 2004 and 2015.

We separated the data by host (species or genus of interest) and disease type

(growth anomalies or tissue loss) and used only observations with available data

for all hypothesized risk factors (predictor variables). The final dataset excluded

observations from the Northwestern Hawaiian Islands because several

hypothesized risk factors were only available for the Main Hawaiian Islands. The

number of disease and healthy colony observations we used for each host-

disease pair are shown in Table 1.

Ecological disease drivers

For each coral health observation, we collected biological, environmental,

physical, and anthropogenic risk factors corresponding to the time and location of

each observation (Tables 2-3). We selected risk factors based on relationships

described in previous studies and based on personal observations (Table 2).

After removing variables that were highly colinear (see *Testing for*

multicollinearity below), our hypothesized risk factors were: (i) biological variables

of colony size, host density, and herbivore fish abundance; (ii) environmental

variables of anomalous temperature (Hot Snap), rainfall, chlorophyll-*a* (proxy for

water quality/eutrophication), and irradiance; (iii) physical variables of depth,

outside/inside of an embayment, acute and chronic wave exposure, and stream

exposure (proxies for potential terrestrial runoff and retention); and (iv)

anthropogenic variables of human population size and agriculture and golf

course runoff. Variables reflect short-term (acute, pulse) or long-term (chronic,

press) interactions or site characteristics (Table 3).

For each risk factor, we used survey data where recorded, or compiled externally

sourced data from the vicinity (and within the watershed) of where the survey

was conducted. We used measurements of colony size and depth that were

recorded in the coral health surveys and calculated host density from the survey

data. We extracted all other risk factors by survey location and by time (where

applicable; i.e., acute interactions) using ArcGIS v 10.4. Measurements of rainfall

were land-based, using gridded values from the land location nearest to the

survey site that contained data within the appropriate watershed and month;

human population size was similarly determined. For chlorophyll-*a*, wave

exposure, irradiance, Hot Snaps, and herbivore fish abundance, we used data

from the pixel containing the survey location when possible or else the nearest

pixel with available data over the appropriate time period. However, we used

herbivore fish abundance data compiled from only one year (2012-2013). We

expect this data to adequately represent the spatial structure of herbivore fish

abundance across the archipelago. Further, we expect year to year variation

within the study period to be minimal, as the major drivers of herbivore fish

dynamics – major disturbance events [52,53] and protection status [54] – were

minimized during the study period: there were no changes to marine protection

status across the study region and only one major disturbance event (mass coral

bleaching in 2015) at the end of the study period. We calculated agriculture and

golf course runoff for each ocean pixel as the value from the nearest watershed

decayed with distance from shore; therefore, we used the same technique as

described above acquiring data from the pixel containing the survey location

when possible or else the nearest pixel with available data. We manually

determined whether a site was outside or inside of an embayment by visualizing

the survey site locations relative to coastal boundaries downloaded from the

State of Hawaii Office of Planning. We calculated stream exposure as the inverse

planar distance between the site location and the nearest stream mouth within a

given watershed. Higher stream exposure values indicate sites closer to stream

mouths. For sites located in watersheds without streams, we set the stream

exposure value to zero.

We used a different combination of risk factors for growth anomalies and tissue

loss models because of hypothesized differences in their modes of transmission

and etiologies. We included ten hypothesized risk factors in the growth

anomalies models and eleven hypothesized risk factors in the tissue loss models;

seven risk factors were common to both diseases (Table 2).

Testing for multicollinearity

To test for multicollinearity among risk factors, we calculated the variance

inflation factor. The variance inflation factor (VIF) is a ratio that quantifies the

amount of multicollinearity among covariates in a regression model by dividing

the variance of a model with multiple terms by the variance of a model with a

single term. A VIF equal to one indicates a variable is not correlated with other

variables, a value between one and five indicates moderate correlation, and a

value above five indicates high correlation. We initially hypothesized that, in

addition to the risk factors listed in Table 2, total coral density, wintertime thermal

stress (Cold Snaps, Winter Condition), and coastal development should be

included as risk factors for both growth anomalies and tissue loss models, and

that an additional metric of urban runoff should be included in the growth

anomalies model. However, these risk factors had VIF values greater than five

indicating strong multicollinearity with other covariates (Fig. S4), so we excluded

these risk factors from our candidate models.

Logistic regression without the case-control design

We investigated the ecological drivers of each host-disease pair using logistic

regression with all available coral health data (Tables 1-3) and assessed model

predictive accuracies and true positive rates based on withheld portions of data

(see below). The response variable in our logistic regression models described

the colony observation health state as 0 for healthy and 1 for diseased. We

included the hypothesized risk factors listed in Table 2 (combination dependent

on disease type) for the full models with island and month as random effects and

conducted backward model selection to find the best fit model based on the

lowest Akaike Information Criterion (AIC). We included island and month as

random effects to make the model results as comparable as possible with the

case-control model results, as the case-control datasets were sub-sampled

accounting for island and survey month. We conducted this analysis using the

lme4 package [55] in R statistical software v 3.5.1. We used 80% of the data for

model development and model selection, and the remaining 20% of data to

quantify model predictive accuracy (proportion of total colony observations

correctly identified as healthy and diseased) and true positive rate (proportion of

disease colony observations correctly identified as diseased). We split the data

using the sample function in base R, which selects data in proportion to the raw

data (e.g., if the full data has a zero-to-one ratio of 90%-to-10%, then the sample

will also contain 90% zeros and 10% ones).

To ensure that the randomly sampled training and testing data did not bias our

results, we repeated our model development and selection and accuracy

assessments 500 times. Therefore, for each host-disease pair, we split the full

data into 500 paired training and testing datasets, conducted model selection

within each training dataset, and then used the most frequently selected best fit

model (Table S1) to predict health state on the 500 withheld test datasets. We

compared predicted health state with observed health state to calculate accuracy

and true positive rates within each withheld test dataset and report mean values

across all 500 withheld test datasets.

Logistic regression with the case-control design

For the logistic regression analysis with the case-control design, we followed the

approach described above except instead of creating 500 randomly selected

training and testing datasets from all available coral health data for each host-

disease pair, we created 500 case-control datasets each with matching numbers

of disease and healthy observations and randomly split those case-control

datasets into paired training data and testing data (80% training/20% testing). To

create each case-control dataset, we selected all disease colony observations for

a given host-disease pair, and for each disease colony observation we randomly

selected a healthy colony observation of the same host type (by species or

genus) surveyed in the same month (but not necessarily year) and from the

same island as the disease colony observation. We repeated this process to

create 500 case-control datasets per host-disease pair that all had the same

disease observations but varied in the paired healthy colony observations. We

followed the same process of model development, selection, and testing as

described above. Results for the frequency of model selection for the best fit

model are available in Table S1.

Acknowledgements: We thank the Fore-C research team for valuable feedback

on analyses and the following funding sources for this research: NASA Roses

Ecological Forecasting grant NNX17AI21G (MJD, SFH, JMC), NASA Earth and

Space Science Fellowship (JMC, MJD), and Stanford University Woods Institute

for the Environment Environmental Ventures Program (JMC). The scientific

results and conclusions, as well as any views or opinions expressed herein, are

those of the author and do not necessarily reflect the views of NOAA or the

Department of Commerce.

Author contributions: JMC, GA, and MJD designed experiments. JMC and

SFH assembled data. JMC conducted the analyses and wrote the manuscript.

JMC, GA, SFH, and MJD interpreted results and edited the manuscript.

Competing interests: The authors declare no competing interests in relation to

the work described.

591
592

References

1.   Hudson, P. & Greenman, J. Competition mediated by parasites: biological

and theoretical progress. *Trends Ecol. Evol.* **13**, 387–390 (1998).

2.   Kohler, S. L. & Hoiland, W. K. Population regulation in an aquatic insect:

the role of disease. *Ecology* **82**, 2294–2305 (2001).

3.   Lafferty, K. D. & Morris, A. K. Altered behavior of parasitized killifish

increases susceptibility to predation by bird final hosts. *Ecology* **77**, 1390–

1397 (1996).

4.   Holdo, R. M. *et al.* A disease-mediated trophic cascade in the Serengeti

and its implications for ecosystem C. *PLoS Biol.* **7**, e1000210 (2009).

5.   Brunner, F. S., Anaya-Rojas, J. M., Matthews, B. & Eizaguirre, C.

Experimental evidence that parasites drive eco-evolutionary feedbacks.

Proc. Natl. Acad. Sci. U. S. A. **114**, 3678–3683 (2017).

6.   Behringer, D. C., Butler, M. J. & Shields, J. D. Avoidance of disease by

social lobsters. *Nature* **441**, 421–421 (2006).

7.   Woodworth, B. L. *et al.* Host population persistence in the face of

introduced vector-borne diseases: Hawaii amakihi and avian malaria. *Proc.*

Natl. Acad. Sci. U. S. A. **102**, 1531–6 (2005).

8.   Aeby, G. S. Behavioral and ecological relationships of a parasite and its

hosts within a coral reef system. *Pacific Sci.* **45**, 263–269 (1991).

9.   King, G. & Zeng, L. Logistic Regression in Rare Events Data. *Polit. Anal.* **9**,

137–163 (2001).

10. Lewallen, S. & Courtright, P. Epidemiology in practice: case-control

studies. *Community eye Heal.* **11**, 57–8 (1998).

11.   Sutherland, K. P., Porter, J. W. & Torres, C. Disease and immunity in

Caribbean and Indo-Pacific zooxanthellate corals. *Mar. Ecol. Prog. Ser.*

266, 273–302 (2004).

12.   Stimson, J. Ecological characterization of coral growth anomalies on

Porites compressa in Hawai'i. *Coral Reefs* **30**, 133–142 (2010).

13.   Ruiz-Moreno, D. *et al.* Global coral disease prevalence associated with sea

temperature anomalies and local factors. *Dis. Aquat. Organ.* **100**, 249–261

(2012).

14.   Palmer, C. V. & Baird, A. H. Coral tumor-like growth anomalies induce an

immune response and reduce fecundity. *Dis. Aquat. Organ.* **130**, 77–81

(2018).

15.   Couch, C. S. *et al.* Spatial and temporal patterns of coral health and

disease along leeward Hawai'i Island. *Coral Reefs* **33**, 693–704 (2014).

16.   Kaczmarsky, L. & Richardson, L. L. Do elevated nutrients and organic

carbon on Philippine reefs increase the prevalence of coral disease? *Coral*

Reefs **30**, 253–257 (2011).

17.   Aeby, G. S. *et al.* Growth anomalies on the coral genera Acropora and

Porites are strongly associated with host density and human population

size across the Indo-Pacific. *PLoS One* **6**, e16887 (2011).

18.   Williams, G. J., Aeby, G. S., Cowie, R. O. M. & Davy, S. K. Predictive

modeling of coral disease distribution within a reef system. *PLoS One* **5**,

e9264 (2010).

19.   Aeby, G. S. *et al.* Emerging coral diseases in Kaneohe Bay, Oahu, Hawaii

(USA): two major disease outbreaks of acute Montipora white syndrome.

Dis. Aquat. Organ. **119**, 189–198 (2016).

20.   Brodnicke, O. B. *et al.* Unravelling the links between heat stress, bleaching

and disease: fate of tabular corals following a combined disease and

bleaching event. *Coral Reefs* **38**, (2019).

21.   Hobbs, J.-P. A., Frisch, A. J., Newman, S. J. & Wakefield, C. B. Selective

Impact of Disease on Coral Communities: Outbreak of White Syndrome

Causes Significant Total Mortality of Acropora Plate Corals. *PLoS One* **10**,

e0132528 (2015).

22.   Precht, W. F., Gintert, B. E., Robbart, M. L., Fura, R. & van Woesik, R.

Unprecedented Disease-Related Coral Mortality in Southeastern Florida.

Sci. Rep. **6**, 31374 (2016).

23.   Walton, C. J., Hayes, N. K. & Gilliam, D. S. Impacts of a Regional, Multi-

Year, Multi-Species Coral Disease Outbreak in Southeast Florida. *Front.*

Mar. Sci. **5**, (2018).

24.   Sussman, M., Willis, B. L., Victor, S. & Bourne, D. G. Coral pathogens

identified for White Syndrome (WS) epizootics in the Indo-Pacific. *PLoS*

One **3**, e2393 (2008).

25.   Ben-Haim, Y. *et al.* Vibrio coralliilyticus sp. nov., a temperature-dependent

pathogen of the coral Pocillopora damicornis. *Int. J. Syst. Evol. Microbiol.*

53, 309–315 (2003).

26.   Ushijima, B. *et al.* Vibrio coralliilyticus Strain OCN008 is an etiological

agent of acute Montipora White Syndrome. *Appl. Environ. Microbiol.* **80**,

(2014).

27. Ushijima, B. *et al.* Mutation of the toxR or mshA genes from Vibrio

coralliilyticus strain OCN014 reduces infection of the coral Acropora

cytherea. *Environ. Microbiol.* **18**, 4055–4067 (2016).

28. Bruno, J. F. *et al.* Thermal stress and coral cover as drivers of coral

disease outbreaks. *PLoS Biol.* **5**, e124 (2007).

29. Caldwell, J. M., Heron, S. F., Eakin, C. M. & Donahue, M. J. Satellite SST-

Based Coral Disease Outbreak Predictions for the Hawaiian Archipelago.

Remote Sens. **8**, 93 (2016).

30. Heron, S. F. *et al.* Summer hot snaps and winter conditions: modelling

white syndrome outbreaks on Great Barrier Reef corals. *PLoS One* **5**,

e12210 (2010).

31. Maynard, J. A. *et al.* Predicting outbreaks of a climate-driven coral disease

in the Great Barrier Reef. *Coral Reefs* **30**, 485–495 (2011).

32. Randall, C. J., Jordan-Garza, A. G., Muller, E. M. & Van Woesik, R.

Relationships between the history of thermal stress and the relative risk of

diseases of Caribbean corals. *Ecology* **95**, 1981–1994 (2014).

33. Caldwell, J. M., Donahue, M. J. & Harvell, C. D. Host size and proximity to

diseased neighbours drive the spread of a coral disease outbreak in

Hawai'i. *Proc. R. Soc. B* **285**, (2018).

34. Jolles, A. E., Sullivan, P., Alker, A. P. & Harvell, C. D. Disease

Transmission of Aspergillosis in Sea Fans: Inferring Process from Spatial

Pattern. *Ecology* **83**, 2373–2378 (2002).

35. Bythell, J. C., Brown, B. E. & Kirkwood, T. B. L. Do reef corals age? *Biol.*

Rev. **93**, 1192–1202 (2018).

36. Bak, R. & Meesters, E. Coral population structure:the hidden information of

colony size-frequency distributions. *Mar. Ecol. Prog. Ser.* **162**, 301–306

(1998).

37. Meesters, E. H. *et al.* Colony size-frequency distributions of scleractinian

coral populations: spatial and interspecific variation. *Marine Ecology*

Progress Series **209**, 43–54 (2001).

38. Jordán-Dahlgren, E., Jordán-Garza, A. G. & Rodríguez-Martínez, R. E.

Coral disease prevalence estimation and sampling design. *PeerJ* **6**, e6006

(2018).

39. Nugues, M. M., Smith, G. W., Hooidonk, R. J., Seabra, M. I. & Bak, R. P.

M. Algal contact as a trigger for coral disease. *Ecol. Lett.* **7**, 919–923

(2004).

40. Kaczmarsky, L. & Richardson, L. L. Transmission of growth anomalies

between Indo-Pacific Porites corals. *J. Invertebr. Pathol.* **94**, 218–221

(2007).

41. Yoshioka, R. M., Kim, C. J. S. S., Tracy, A. M., Most, R. & Harvell, C. D.

Linking sewage pollution and water quality to spatial patterns of Porites

lobata growth anomalies in Puako, Hawaii. *Mar. Pollut. Bull.* **104**, 313–321

(2016).

42. Randall, C. J. & Van Woesik, R. Contemporary white-band disease in

Caribbean corals driven by climate change. *Nat. Clim. Chang.* **5**, 375–379

(2015).

43. Jones, R., Bowyer, J., Hoegh-Guldberg, O. & Blackall, L. L. Dynamics of a

temperature-related coral disease outbreak. *Mar. Ecol. Prog. Ser.* **281**, 63–

77 (2004).

44. Lesser, M. P., Weis, V. M., Patterson, M. R. & Jokiel, P. L. Effects of

morphology and water motion on carbon delivery and productivity in the

reef coral, Pocillopora damicornis (Linnaeus): Diffusion barriers, inorganic

carbon limitation, and biochemical plasticity. *J. Exp. Mar. Bio. Ecol.* **178**,

153–179 (1994).

45. Work, T. M., Russell, R. & Aeby, G. S. S. Tissue loss (white syndrome) in

the coral Montipora capitata is a dynamic disease with multiple host

responses and potential causes. *Proc. R. Soc. B Biol. Sci.* **279**, 4334–4341

(2012).

46. Beurmann, S. *et al.* Pseudoalteromonas piratica strain OCN003 is a coral

pathogen that causes a switch from chronic to acute Montipora white

syndrome in Montipora capitata. *PLoS One* **12**, e0188319 (2017).

47. Breslow, N. E. Statistics in Epidemiology: The Case-Control Study. *J. Am.*

Stat. Assoc. **91**, 14–28 (1996).

48. Venette, R. C., Moon, R. D. & Hutchison, W. D. Strategies and Statistics of

Sampling for Rare Individuals. *Annu. Rev. Entomol.* **47**, 143–174 (2002).

49. Balmaseda, A. *et al.* Index Cluster Study of Dengue Virus Infection in

Nicaragua. *Am. J. Trop. Med. Hyg.* **83**, 683–689 (2010).

50.   Burnham, K., Anderson, D., & Laake, J. Estimation of Density from Line

740   Transect Sampling of Biological Populations. *Wildl. Monogr.* **72**, 3–202

741   (1980).

742   51.   Caldwell, J. M. *et al.* Hawaiʻi Coral Disease database (HICORDIS):

743   species-specific coral health data from across the Hawaiian archipelago.

744   *Data in Brief* **8**, 1054–1058 (2016).

745   52.   Taylor, B. M. *et al.* Synchronous biological feedbacks in parrotfishes

746   associated with pantropical coral bleaching. *Glob. Chang. Biol.* gcb.14909

747   (2019). doi:10.1111/gcb.14909

748   53.   Graham, N. A. J., Jennings, S., MacNeil, M. A., Mouillot, D. & Wilson, S. K.

749   Predicting climate-driven regime shifts versus rebound potential in coral

750   reefs. *Nature* **518**, 94–97 (2015).

751   54.   Stockwell, B., Jadloc, C., Abesamis, R., Alcala, A. & Russ, G. Trophic and

752   benthic responses to no-take marine reserve protection in the Philippines.

753   *Mar. Ecol. Prog. Ser.* **389**, 1–15 (2009).

754   55.   Douglas, B., Mächler, M., Bolker, B. & Walker, S. Fitting Linear Mixed-

755   Effects Models Using lme4 | Bates | Journal of Statistical Software. *J. Stat.*

756   *Softw.* **67**, (2015).

757   56.   Hobbs, J. & Frisch, A. Coral disease in the Indian Ocean: taxonomic

758   susceptibility, spatial distribution and the role of host density on the

759   prevalence of white syndrome. *Dis. Aquat. Organ.* **89**, 1–8 (2010).

760   57.   McClanahan, T. R., Weil, E. & Maina, J. Strong relationship between coral

761   bleaching and growth anomalies in massive Porites. *Glob. Chang. Biol.* **15**,

1804–1816 (2009).

58.   Aeby, G. S. *et al.* Patterns of Coral Disease across the Hawaiian

Archipelago: Relating Disease to Environment. *PLoS One* **6**, e20370

(2011).

59.   Lamb, J. B. *et al.* Plastic waste associated with disease on coral reefs.

Science (80-.). **359**, 460–462 (2018).

60.   Nugues, M. M., Smith, G. W., Hooidonk, R. J., Seabra, M. I. & Bak, R. P.

M. Algal contact as a trigger for coral disease. *Ecol. Lett.* **7**, 919–923

(2004).

61.   Raymundo, L. J., Halford, A. R., Maypa, A. P., Kerr, A. M. & Karl, D. M.

Functionally diverse reef-fish communities ameliorate coral disease.

62.   Garren, M., Smriga, S. & Azam, F. Gradients of coastal fish farm effluents

and their effect on coral reef microbes. *Environ. Microbiol.* **10**, 2299–2312

(2008).

63.   Lamb, J. B., Williamson, D. H., Russ, G. R. & Willis, B. L. Protected areas

mitigate diseases of reef-building corals by reducing damage from fishing.

Ecology **96**, 150422141041004 (2015).

64.   Bruno, J. F. *et al.* Nutrient enrichment can increase the severity of coral

diseases. *Ecol. Lett.* **6**, 1056–1061 (2003).

65.   Sheridan, C., Jm, B., Kushmaro, A., Frejaville, Y. & Eeckhaut, I. Terrestrial

runoff influences white syndrome prevalence in SW Madagascar. *Mar.*

Environ. Res. **101**, 44–51 (2014).

66.   Haapkylä, J., Seymour, A. S., Trebilco, J. & Smith, D. Coral disease

prevalence and coral health in the Wakatobi Marine Park, south-east

Sulawesi, Indonesia. *J. Mar. Biol. Assoc. UK* **87**, 403 (2007).

67. Heenan, A. *et al. Ecological Monitoring 2012-2013 - reef fishes and benthic*

habitats of the main Hawaiian Islands, American Samoa, and Pacific

Remote Island Areas. (2014). doi:10.13140/RG.2.1.4351.9122

68. Wedding, L. M. *et al.* Advancing the integration of spatial data to map

human and natural drivers on coral reefs. *PLoS One* **13**, e0189792 (2018).

69. Frazier, A. G., Giambelluca, T. W., Diaz, H. F. & Needham, H. L.

Comparison of geostatistical approaches to spatially interpolate month-

year rainfall for the Hawaiian Islands. *Int. J. Climatol.* **36**, 1459–1470

(2016).

70. Lecky, J. H. Ecosystem Vulnerability and Mapping Cumulative Impacts on

Hawaiian Reefs. (University of Hawaii at Manoa, 2016).

doi:10.13140/RG.2.2.33194.52162

71. Gelman, A. & Hill, J. *Data analysis using regression and*

hierarchical/multilevel models. (Cambridge, 2007).

Case-control design identifies ecological drivers of endemic coral diseases - Supplemental Materials

Jamie M. Caldwell[1,2*], Greta Aeby[3], Scott F. Heron[2,4,5], and Megan J. Donahue[1]

6. Hawaii Institute of Marine Biology, University of Hawaii at Manoa, Hawaii, USA
7. ARC Centre of Excellence for Coral Reef Studies, James Cook University, Townsville, Australia
8. Qatar University, Department of Biological & Environmental Sciences, Doha, Qatar
9. Marine Geophysical Laboratory, Physics, College of Science and Engineering, James Cook University, Townsville, Australia
10. NOAA Coral Reef Watch, College Park, Maryland, USA
* contact: jamie.sziklay@gmail.com

Figure S1: Disease risk as a function of colony size across host-disease pairs. Points indicate disease presence (P=1) and disease absence (P=0) from one random sample of a case control training dataset.

Figure S2: Distribution of risk factors (predictor variables) in growth anomaly models. Distribution of risk factors included in candidate growth anomaly models after removing observations with missing data. For the risk factor "Bay", zero indicates outside a bay and one indicates inside a bay.

Figure S3: Distribution of risk factors (predictor variables) in tissue loss models. Distribution of risk factors included in candidate tissue loss models after removing observations with missing data. For the risk factor "Bay", zero indicates outside a bay and one indicates inside a bay.

Figure S4: Variance inflation factor plots. Variance inflation factors for each predictor variable across host-disease pairs. Values greater than five (dashed horizontal line) indicates strong multicollinearity. For consistency, we removed predictor variables from all analyses if values exceeded five any host-disease pair.

Table S1: Number of times the best fit model was selected as the best model within the 500 training datasets.
Disease refers to growth anomalies (GA) and tissue loss (TL).

Disease	Species	Frequency of best fit model without case-control deign	Frequency of best fit model with case-control deign
GA	*Montipora capitata*	497	76
GA	*Montipora patula*	125	73
GA	*Montipora* spp	176	69
GA	*Porites compressa*	212	113
GA	*Porites evermanni*	111	36
GA	*Porites lobata*	278	87
GA	*Porites* spp	354	83
TL	*Montipora capitata*	187	23
TL	*Montipora patula*	92	43
TL	*Montipora* spp	154	47
TL	*Porites compressa*	125	86
TL	*Porites evermanni*	216	30
TL	*Porites lobata*	474	90
TL	*Porites* spp	139	63