Acid Experimental Evolution of the Extremely Halophilic Archaeon *Halobacterium* sp.

NRC-1 Selects Mutations Affecting Arginine Transport and Catabolism

by

Karina S. Kunka,[a] Jessie M. Griffith,[a] Chase Holdener,[a] Katarina M. Bischof,[a] Haofan Li,[a] Priya

DasSarma,[b] Shiladitya DasSarma,[b] and Joan L. Slonczewski[a#]

[a]Department of Biology, Kenyon College, Gambier, Ohio, USA.

[b]Institute of Marine and Environmental Technology, Department of Microbiology and

Immunology, University of Maryland School of Medicine, Baltimore, Maryland, USA

[#]Corresponding Author:

Joan L. Slonczewski, slonczewski@kenyon.edu

Department of Biology, Kenyon College, Gambier, Ohio, USA

ABSTRACT

Background

Halobacterium sp. NRC-1 (NRC-1) is an extremely halophilic archaeon that is adapted to

multiple stressors such as UV, ionizing radiation and arsenic exposure. We conducted

experimental evolution of NRC-1 under acid stress. NRC-1 was serially cultured in CM+

medium modified by four conditions: optimal pH (pH 7.5), acid stress (pH 6.3), iron amendment

(600 μM ferrous sulfate, pH 7.5), and acid plus iron (pH 6.3, with 600 μM ferrous sulfate). For

each condition, four independent lineages of evolving populations were propagated. After 500

generations, 16 clones were isolated for phenotypic characterization and genomic sequencing.

Results

Genome sequences of all 16 clones revealed 378 mutations, of which 90% were haloarchaeal

insertion sequences (ISH) and ISH-mediated large deletions. This proportion of ISH events in

NRC-1 was five-fold greater than that reported for comparable evolution of *E. coli*. One acid-

evolved clone had increased fitness compared to the ancestral strain when cultured at low pH.

Seven of eight acid-evolved clones had a mutation within or upstream of *arcD*, which encodes an

arginine-ornithine antiporter; no non-acid adapted strains had *arcD* mutations. Mutations also

affected the *arcR* regulator of arginine catabolism, which protects bacteria from acid stress by

release of ammonia. Two acid-adapted strains shared a common mutation in *bop*, which encodes

the bacteriorhodopsin light-driven proton pump. Unrelated to pH, one NRC-1 minichromosome

(megaplasmid) pNRC100 had increased copy number, and we observed several mutations that

eliminate gas vesicles and arsenic resistance. Thus, in the haloarchaeon NRC-1, as in bacteria,

pH adaptation was associated with genes involved in arginine catabolism and proton transport.

Conclusions

Our study is among the first to report experimental evolution with multiple resequenced genomes

of an archaeon. Haloarchaea are polyextremophiles capable of growth under environmental

conditions such as concentrated NaCl and desiccation, but little is known about pH stress.

Halobacterium sp. NRC-1 (NRC-1) is considered a model organism for the feasibility of

microbial life in iron-rich brine on Mars. Interesting parallels appear between the molecular basis

of pH adaptation in NRC-1 and in bacteria, particularly the acid-responsive arginine-ornithine

system found in oral streptococci.

INTRODUCTION
49
Halobacterium sp. NRC-1 (NRC-1) is a polyextremophile that grows optimally at NaCl

concentrations in excess of 4 molar (1). A genetically tractable model microbe (2), it was the first

halophilic Archaeon with a fully sequenced genome (3). Besides high salt, NRC-1 is capable of

surviving: high doses of ionizing radiation and dessication (4), UV radiation (5), temperature

extremes (6), and toxic ions such as arsenite (7). These traits have made NRC-1 a model for

studying the possibility of life outside Earth under conditions such as the stratosphere (8,9) or on

Mars (10–12).

Water on Mars contains high concentrations of salt, as well as acid and iron (13). The

Mars Exploration Rover Opportunity discovered substantial deposits of an iron hydrous sulfate

mineral known as jarosite [$KFe^{3+}_3(OH)_6(SO_4)_2$] which forms in acidic and iron-rich aqueous

environments. On earth such conditions occur in acid mine drainage and near volcanic vents.

Opportunity's discovery of jarosite on Mars was evidence of acidic, liquid water and an

oxidizing atmosphere in the Martian past (13,14). Occuring together, acid and metals can

amplify the stress associated with each condition (15). Thus, it is of interest to investigate how a

neutralophilic halophile such as NRC-1 (16) might adapt to conditions of acid and high iron.

An informative approach to examine the genomic basis of stress response is experimental

laboratory evolution (17–23). Experimental evolution of bacteria reveals changes in phenotype

and genotype in response to specific stressors in a controlled environment, such as carbon source

limitation or extreme pH. In bacterial adaptation to various kinds of pH stress, we find a

recurring pattern that dominant responses to short-term stress actually decrease fitness over many

generations of long-term exposure. For example, amino-acid transport and catabolism play

important roles in extreme-acid survival of *Escherichia coli* (24,25). However, 2,000 generations

of *E. coli* evolution at pH 4.8 select for loss of three acid-inducible amino-acid decarboxylase

systems, including arginine decarboxylase (21). As a membrane-permeant acid, benzoic acid

induces glutamate decarboxylase and drug resistance regulons, yet these systems are lost or

downregulated during experimental evolution (26),(20). At high external pH, *E. coli* survival

requires the stress sigma factor RpoS; however, generations of growth at high pH select against

RpoS expression and activity (27). It is therefore of interest to investigate whether similar

patterns of reversal occur in archaea.

Relatively few experimental evolution studies have been reported in archaea. In NRC-1,

serial application of lethal doses of ionizing radiation selected more resistant mutants that had

increased expression of a single-strand DNA binding protein (28). In the thermoacidophile

Sulfolobus solfatericus, serial passage in extreme acid yielded strains that grow below pH 1 (29).

These strains showed mutations in amino acid transporters, as well as upregulation of membrane

biosynthesis and oxidative stress response. In *Metallosphaera sedula*, serial passage led to a pH

0.9-adapted strain with four mutations, one of which is an amino-acid/polyamine transporter

(30). These findings are intriguing, given the role of amino-acid transport and catabolism in

extreme-acid survival of bacteria (24,25). For example, arginine transport and catabolism, which

yields CO_2 plus two ammonium ions, is a prominent response to acid stress of oral streptococci

(31),(32).

Archaea employ various processes that involve proton transport via primary pumps and

antiporters (24,33,34). *Halobacterium* strains possess the light-driven proton pump

bacteriorhodopsin (*bop*) that generates proton motive force (PMF) (35,36) as well as several

sodium-proton antiporters, which export sodium in exchange for protons (6).

We conducted experimental evolution of NRC-1 under conditions of low pH (pH 6.5-6.3)

and at optimal pH for growth (pH 7.5), with high iron versus low iron concentration. The NRC-1

genome includes a main chromosome and two minichromosomes or megaplasmids (3,37) . It

accumulates frequent IS mutations (38,39) which may mediate rapid adaptations to

environmental stress. Our study of experimental evolution in a haloarchaeon assesses which

mutations contribute to archaeal evolution in acid stress. Here we describe analysis of phenotypic

changes across evolved clones from each population, and then use genomic analysis to identify

potential underlying mutational bases of these phenotypic responses to selection. Genome

analysis of 16 clones revealed a remarkable proportion of events mediated by insertion

sequences (ISH). In acid-adapted strains, we found a high frequency of mutations in the arginine-

ornithine antiporter *arcD* (40) and in the associated *arcR* arginine catabolism regulator (41).

RESULTS

Experimental evolution under conditions of acid and iron stress.

Serial culture of evolving populations was conducted as described under Methods (Additional

File 1, Fig. S1). Populations of NRC-1 were founded from a single clone and cultured in

modified CM^+ medium (2,3) with appropriate buffers to maintain pH. Each population was

diluted 500-fold every four days (approximately 9 generations). Four independent populations

were maintained for each condition: the optimal growth condition, pH 7.5 (designation M); acid

stress, initially pH 6.5, later pH 6.3 (designated J); iron amendment, pH 7.5 with 600 µM ferrous

sulfate (designated S); and acid with iron amendment (designated K) for a total of 16

experimental populations. Populations evolved under acid stress were cultured at an initial pH of

6.5, which was then lowered to 6.3 at generation 250, as the populations adapted.

After all populations reached 500 doublings, two clones were isolated from each

population by three rounds of streaking on CM+ agar for a total of 32 evolved clones. Genomic

DNA was extracted from 16 of these clones, and from the founder stock of NRC-1. DNA

samples were sequenced by Illumina MiSeq, and mutations were identified by comparison of the

"evolved strain" sequences to that of the NRC-1 ancestral stock, assembled on the reference

genome (3) using the *breseq* pipeline (42–44). The strains we characterized are listed in **Table 1**.

Mutations in the genomes from evolving populations.

The genomes of the 16 clones were compared to those of the NRC-1 ancestor which we

resequenced from our lab stock (**Tables S1, S2, S3**). The genome of our NRC-1 stock was also

compared to that of the NCBI reference sequence *Halobacterium* sp. NRC-1 (3) as shown in

Additional File 1, **Table S4**. A small number of positions differed from that of the reference.

Some of these differences are consistent with those of later sequence reports (45,46). The

sequences differences shown in **Table S4** were excluded in our analysis of the evolved clones.

The genomes of the evolved clones had a total of 378 mutations, of which 349 were

unique to one strain at the base-pair level. Representative mutations of interest are summarized

in **Table 2**. Mutation frequencies were compared for the main replicon and minichromosomes. In

total across all resequenced genomes there were 120 mutations in minichromosome pNRC100,

and 171 mutations on minichromosome pNRC200. pNRC100 is about 10% as long as the main

chromosome, and pNRC200 is about 20% as long; thus, the two minichromosomes had a

mutation frequency more than ten-fold greater than that of the main chromosome, a finding

consistent with previous reports of plasmid or minichromosome mutation (3).

In the 16 clones, overall, 87 different mutations were found on the main chromosome. Of

these, 90% consisted of new ISH positions, or of deletions mobilized by existing ISH elements

(**Table 3**). Mutation distributions of the minichromosome replicons showed no significant

difference in ISH proportion (94 mutations out of 111, on pNRC100; 141 out of 156 on

pNRC200). For comparison, we considered a recent *breseq* analysis of 16 *E. coli* genomes

following 500 generations evolution with an organic acid (26). Only 18% of the *E. coli*

mutations were mediated by insertion sequences. Thus, NRC-1 evolution showed five-fold

greater proportion of insertion sequence activity than *E. coli*. Our quantitative analysis of

experimentally evolved genomes is consistent with earlier evidence of high ISH activity in

halobacterial genomes (38),(47–50),(45).

Haloarchaea including *Halobacterium salinarum* species are known for polyploidy (15-

25 genome copies per cell) and for ploidy variation among replicons within a cell (51). Our

evolved clones showed evidence for variable ploidy between and within replicons. Mean read

coverage by replicon was modeled by *breseq* (**Table 4**).

Overall, within the ancestor and the evolved clones, the read coverage for the main

chromosome was consistent with that of the minichromosome pNRC200. However, the mean

coverage of the shorter minichromosome pNRC100 (191 kb) was more than twice that of the

main chromosome, for our ancestral NRC-1 and for 12 of the 16 evolved clones. Clones J1, M3-

1, K3, S2, and S3 had mean coverage of pNRC100 more than four-fold greater than that of the

main chromosome. These high coverage ratios could indicate that our original NRC-1 stock has

a double copy number of minichromosome pNRC100, relative to the main chromosome; and that

some descendant clones have increased relative copy number. However, the calculations are

complicated by wide variation in read coverage between different segments of the same replicon,

especially in pNRC100. This variation in read coverage may be caused by internal repeats within

the replicon (35). Interpretation of the data is complicated by the presence of massive deletions

(Additional File 1, **Table S2**) which comprise up to 50% of the ancestral sequence (for example

in clone K1) (50). Variation in read coverage could indicate the presence of plasmid copies with

different deletion levels within a given polyploid cell.

Acid-evolved clone J3-1 has a growth advantage over a range of pH values.

After 500 generations of serial culture under four conditions, clones were isolated from the

evolving populations. The clones were tested for genetic adaptation under various growth

conditions. Each evolved clone was cultured in parallel with the ancestral strain NRC-1. The loss

of gas vesicles (Vac$^-$ phenotype) alters their OD_{600} reading (38,47); for this reason, clones that

had lost gas vesicles were cultured in parallel with a Vac$^-$ isolate of NRC-1 ancestor.

The growth of acid-evolved J-population clones was compared to that of the NRC-1

ancestor (Vac$^+$) (**Figs. 1 and 2**). Clone J3-1 reached a significant two-fold higher culture density

than did the ancestor when cultured at pH 6.1 or at 6.3 (**Fig. 1B**). Growth advantage was seen for

all four replicate cultures of J3-1 at pH 6.1 and at pH 6.3, whereas the difference from NRC-1

cultures disappeared at pH 7.2 and at pH 7.5. Thus, strain J3-1 exhibits an acid-specific fitness

advantage. The other acid-evolved J-population strains, however, had no significant growth

advantage compared to NRC-1, under the conditions tested (**Fig. 2**).

Acid-adapted clones shared mutations in *arcD* and in *arcR*.

We inspected the genomes of acid-adapted populations J and K (acid with iron supplement) for

mutations in specific genes that were not found in the populations evolved at pH 7.5. Seven out

of eight of the J and K clones (but no M or S clones) had ISH mutations in or upstream of gene

VNG_6313G (**Table 2**). This gene was originally classified as encoding a sodium-proton

antiporter (*nhaC3*) but was shown instead by physiological experiments to encode an arginine-

ornithine antiporter ArcD (52). PCR amplification and Sanger sequencing of the mutant *arcD*

alleles confirmed the presence of insertion sequences ISH2 (strains K1 and K4) and ISH4

(strains J1, K2-1, K3) (**Table 5**; Additional File 1, **Fig. S2**). Additionally, in J4-2, a partial

sequence confirms the presence of 1.1 kb ISH11 insertion flanked by a 10 bp direct repeat, while

a large 3000+ bp insertion in K3 returned a partial sequence of ISH4. The partial sequence

suggests multiple copies of ISH4, or possibly a composite transposon.

Four acid-evolved genomes (J-3, K-1, K2-1, K4-1) and one non-acid-evolved clone (M3-

1) possess ISH insertions at different sites in *arcR* on pNRC100 (Additional File 1, **Table S3**).

ArcR mediates transcriptional regulation of the *arcABDCR* operon for arginine catabolism

(40,41). These components include the arginine deiminase (*arcA*), ornithine

carbamoyltransferase (*arcB*), carbamate kinase (*arcC*), and the arginine-ornithine antiporter

(*arcD*).

For comparison, a remarkably similar system of arginine catabolism reverses acidification

for the periodontal bacterium *Streptococcus gordonii* (31,32). Arginine catabolism releases CO_2

and two molecules of ammonia, which cause net alkalization. The system mediates tooth biofilm

formation by *S. gordonii*. For *E. coli*, the arginine decarboxylase Adi reverses acidification at

extreme low pH (52). The *adi* system of *E. coli* is induced by acid stress but largely lost by

insertion-sequence mutations after long-term evolution in acid (20,22). This suggests a model for

acid adaptation in haloarchaea that is remarkably similar to that observed in *E. coli*, in which

acid-stress adaptations are knocked down by long-term acid exposure (21).

Acid-adapted clones shared mutations in bacteriorhodopsin (*bop*).

In NRC-1, our acid-evolved clones J3-1 and K1 each contained an ISH element in the gene *bop*

that encodes the light-driven proton pump bacteriorhodopsin (35). The J3-1 allele was confirmed

by Sanger sequence as a 1.1 kb insertion of ISH1 with an eight bp target site duplication in *bop*

(**Table 5**; Additional File 1, **Fig. S2**). This exact mutation has been previously studied in

bacteriorhodopsin mutants, and was in fact the first transposable element identified in

haloarchaea (35). This particular target site duplication was shared with acid-evolved clone K1.

At a different position, a *bop* ISH insertion was found in one of the M population clones (M3-1)

which had not undergone acid selection, consistent with previous spontaneous insertions in this

gene.

The *bop* and *arcD* mutations were found together in J3-1, but also in acid-adapted K1,

which did not show a significant phenotype under our conditions tested. We inspected strain J3-1

221 for candidate mutations that might be responsible for this strain's unique degree of adaptation at

222 low pH. Overall, the J3-1 genome had 16 mutations compared to the NRC-1 ancestor (**Table 6**).

223 Of these, only one mutation affected a gene not affected in any other evolved clone. This is a

224 missense mutation in a ferredoxin gene (*VNG1561*) resulting in a conservative change from

225 lysine to arginine. Mutations were also found affecting several proteins involved in

226 transcriptional regulation, which in combination might contribute to the acid fitness phenotype.

227   **Clones evolved at pH 7.5 show no increase in relative fitness.** All evolved clones from

228 generation 500 with Vac$^-$ phenotypes were grown over 200 hours in unbuffered CM$^+$ medium

229 without acid or iron amendment and compared to the growth phenotype of the NRC-1 Vac$^-$

230 control strain (**Fig. S2A**). Similarly, the growth phenotypes in unstressed medium of Vac$^+$ clones

231 from the 500-generation populations that retained them were compared to that of the NRC-1

232 Vac$^+$ ancestor (**Fig. S2B**). None of the M populations show a significant growth advantage

233 compared to the ancestral strain (Additional File 1, **Fig. S3A and B**).

234   Growth curves were also conducted for clones from the S populations (evolved with 600

235 µM FeSO$_4$). Media contained CM+ pH 7.5 with 100 mM MOPS and 600 µM FeSO$_4$. All

236 evolved clones were persistent Vac$^-$ mutants at generation 500 and were therefore compared to

237 an NRC-1 Vac$^-$ control (Additional File 1, **Fig. S4**). No significant differences were observed.

238   **Multiple clones lost gas vesicles and arsenic resistance.** Under laboratory conditions,

239 gas vesicle-producing (Vac$^+$) NRC-1 clones have high rates of spontaneous mutation to a

240 vesicle-deficient (Vac$^-$) phenotype due to mutations in *gvp* on pNRC100 (38,47). Twelve out of

241 sixteen of our evolved clones, including representatives of each selection type, had lost genes

242 required for gas vesicle nanoparticle production (*gvp*) (53–55). Cultures were oxygenated

243 continually by rotating in a bath, effectively eliminating the competitive advantage of producing

gas vesicles in oxygen-limiting environments. Thus, as expected, many insertions and deletions

were found that had eliminated gas vesicles (45,47). We characterized gas vesicle phenotypes

every 100 generations for the stressed condition populations. These Vac phenotypes (loss of gas

vesicle nanoparticles) are presented by population and organized by respective evolution

condition in **Table 7.** All evolving populations showed loss of gas vesicle production in some

cells. By generation 500, the Vac⁻ phenotype was prevalent in all populations. There was no

significant correlation with pH or with iron amendment.

In addition, 13/16 evolved clones had lost the major arsenic resistance operon (*ars*)

encoded on pNRC100 (7). Other mutations affecting transcriptional regulators and initiation

factors occurred in parallel across multiple populations. These and other parallel mutations are

summarized in **Table 2**. Various hot spots for mutation appear, many of which are caused by

ISH insertions or ISH-mediated deletions.

DISCUSSION

Here we report one of the first evolution experiments to be conducted on a haloarchaeon. A

previous evolution experiment involves selection of mutants resistant to ionizing radiation (28).

We compared four environmental conditions: low pH versus optimal pH 7.5, with or

without iron supplementation. Overall, in the 500-generation evolved strains, we found a striking

pattern of large ISH-mediated deletions, particularly in the two minichromosomes (Additional

File 1, **Tables S1-S3**). For comparison, in *E. coli*, experimental evolution for 2,000 generations

at low pH yields only occasional large deletions (20,21). However, after just 500 generations of

evolution in the haloarchaeon NRC-1, every evolved clone contained several large-scale

deletions. ISH insertion mutations greatly outnumbered SNPs. These types of changes reflect

frequent DNA rearrangements and genetic variability observed previously in NRC-1 (35,39,49).

The acid-adapted NRC-1 populations showed a striking prevalence of mutations affecting

the *arcD* and *arcR* components of arginine transport and catabolism. Arginine catabolism with

ammonia release plays a major role in reversing acidity for Gram-positive and Gram-negative

bacteria. It is striking to see how the role of acid-dependent arginine catabolism may extend to

haloarchaea. The arginase/arginine deiminase family (COG0010) represents a set of orthologs

proposed to be among those transferred horizontally to archaea from an ancient bacterial

ancestor (56).

The ISH insertions seen in acid-adapted clones would be expected to knock out the

arginine system, as seen in *E. coli* experimental evolution with acid (20,41). The reason for the

evolutionary loss is proposed to be a readjustment to long-term acid exposure, for which the

sustained induction of arginine catabolism becomes counterproductive. It is interesting to find

evidence for a similar evolutionary mechanism in a haloarchaeon.

In addition, the acid-evolved strains J3-1 and K1 show an identical insertion mutation

affecting the bacteriorhodopsin *bop* gene. The loss of *bop* may be neutral or advantageous under

low external pH, where a high proton motive force already exists. The bacteriorhodopsin pump

could be a source of proton leakage at high PMF.

The acid-fitness advantage of clone J3-1 could arise from a single mutation unique to J3-

1, such as the missense mutation in a ferredoxin that is unique to J3-1. More likely, however,

acid fitness arises from a cumulative effect of loss of function mutations in a number of other

genes including *arcA*, *arcR*, and *bop*. It is possible that some unknown factor accounts for the

acid-fitness phenotype exhibited by J3-1 under the conditions tested. Nonetheless, it is

interesting that the three genes with mutations prevalent in acid-evolved strains all encode

products involved in proton consumption or export.

Our findings support previous reports of the importance of ISH elements in haloarchaeal

evolution (45), and the observations in *Sulfolobus* that large deletions and loss of function

mutations are fitness tradeoffs for surviving in stressful environments (57). Large deletions and

IS insertions are also common in experimental evolution of bacteria (20,21,27,58). We also find

evidence for accumulation of ploidy changes for the shorter minichromosome, pNRC100 (51).

We show that experimental evolution is an effective approach to identify candidate genes for

environmental stress response in a haloarchaeon.

MATERIALS AND METHODS

***Halobacterium* strains and media.** All evolved clones were derived from a stock of

Halobacterium sp. NRC-1 from the laboratory of Shiladitya DasSarma (3). Liquid cultures were

grown in Complex Medium Plus Trace Metals (CM^+) based on Ref (2), Protocol 25: 250 g/l

NaCl, 20 g/l $MgSO_4 \cdot 7H_2O$, 2 g/l KCl, 3 g/l $Na_3C_6H_5O_7 \cdot 2H_2O$, 10 g/l Oxoid Peptone, and 100

μl/l Trace Metals (3.5 g/l $FeSO_4 \square 7H_2O$, 0.88 g/l $ZnSO_4 \cdot 7H_2O$, 0.66 g/l $MnSO_4 \cdot H_2O$, and 0.2 g/l

$CuSO_4 \cdot 5H_2O$ dissolved 0.1M HCl) with supplements as needed for the conditions examined

(59). CM^+ solid medium included addition of 20 g/l granulated agar. All cultures were incubated

at 42˚C with rotation. Cultures on solid media were incubated at 42˚C for 7–10 days until

colonies reached approximately 1 mm in diameter. A Vac^- mutant of our NRC-1 stock culture

was obtained by picking a Vac^- colony followed by three restreaks on CM^+ agar.

Liquid CM^+ media for experimental evolution was made with either 100mM PIPES

(pKa=6.8) or 100mM MOPS (pKa=7.2) buffer with pH adjusted using 5 M NaOH or 5 M HCl as

needed, followed by filter sterilization. 100 mM $FeSO_4$ stock was prepared in deionized water

and filter-sterilized before every other dilution during serial batch culture evolution. Sterilized

$FeSO_4$ stock was added to buffered CM^+ after filter sterilization. For freezer stocks, live cultures

were mixed 1:1 with a 50% glycerol, 50% complex medium basal salts mixture as a

cryoprotectant. Complex medium basal salts were 250 g/l NaCl, 20 g/l $MgSO_4 \cdot 7H_2O$, 2 g/l KCl,

3 g/l $Na_3C_6H_5O_7 \cdot 2H_2O$. Acidic, control, iron-rich and acidic, and iron-rich media used in the

evolution consisted of: CM^+ pH 6.5 with 100 mM PIPES (populations J1-J4), CM^+ pH 7.5 with

100 mM MOPS (populations M1-M4), CM^+ pH 6.5 (or pH 6.3) with 100 mM PIPES 600 μM

$FeSO_4$ (populations K1-K4), and CM^+ pH 7.5 with 100 mM MOPS 600 μM $FeSO_4$ (populations

S1-S4).

323     **Experimental evolution.** A total of 16 populations (four per evolution condition) were

324   founded from a 5 ml CM^+ tube culture (7-10 days incubation) of *Halobacterium* sp. NRC-1 that

325   was diluted 500-fold and incubated 4 days in a 42^oC shaker bath at 200 rpm. At the end of the

326   fourth day, 10 µl of the previous culture was diluted into 5 ml of fresh CM^+ media amended as

327   necessary for the respective stress condition. The resulting 1:500 dilutions yield approximately

328   nine generations per dilution cycle. If cultures did not reach a healthy cell density as qualitatively

329   evaluated for each dilution, 1:100 or 1:250 dilutions were performed to prevent loss of evolving

330   populations. Alternative dilution concentrations were factored into total generation counts at the

331   end of experimental evolution. When evolution was interrupted, the populations were revived by

332   1:250 dilutions from freezer stocks of the previous dilution. Freezer stocks comprised 1 ml

333   liquid, mature haloarchaea culture for each evolving population and 0.5 ml glycerol/basal salts

334   mixture, stored in 2 ml Wheaton brand vials and frozen at -80^oC for each dilution, totaling 16

335   freezer stocks every four days. A summary of the evolution procedure is presented in **Figure S1**.

336     **Clone selection.** Clones were isolated by plating 10 µl of culture from generation 100,

337   200, 300, 400, and 500 from freezer stocks for all 16 evolving populations on CM^+ agar plates,

338   followed by incubation in a sealed container at 42^oC for 7–10 days. Isolated colonies were then

339   selected for diverse Vac phenotypes, streaked on fresh CM^+ agar plates, and incubated a second

340   time. The process was repeated a third time to ensure isolation of select genetically pure clones.

341   Colonies from the third streak were grown in unbuffered CM^+ pH 7.2, and stocks were frozen for

342   later phenotype and genotype characterization. One clone was isolated from each population

343   every 100 generations. For populations that presented mixed gas vesicle production phenotypes,

344   we isolated both a Vac^+ clone and a Vac^- clone. In total, 75 clones were isolated from generation

345   100, 200, 300, and 400 of the evolution. Clones were similarly isolated from generation 500;

however, the first streak was taken directly from evolving populations, rather than from frozen

stock in Wheaton vials. Two clones were isolated from each population at 500 generations, for a

total of 32 clones.

Gas vesicle formation phenotype analysis. Vesicle formation phenotype was assessed

qualitatively based on the relative translucence of plated colonies and denoted as Vac$^+$ or Vac$^-$ as

appropriate (2,47). If more than one Vac phenotype was observed in a streak during strain

isolation, the phenotypic variant colonies were re-streaked and treated as separate clonal isolates.

Vac phenotypes were evaluated for persistence with each streak based on whether or not Vac$^+$

colonies yielded >1% Vac$^-$ progeny or vice versa.

Growth assays. The generation 500 clones used in these assays are summarized in **Table**

1. Clones were cultured in unbuffered CM+ at pH 7.2, and incubated for four days in a 42°C

shaker bath with 200 rpm orbital aeration. Over-week starter cultures were diluted 1:1000 into

new test tubes with 5 ml of the appropriate test condition media. A media blank was included for

each media condition, and each clone was tested with four to eight biological replicates,

depending on the assay. Immediately after inoculation, OD$_{600}$ values were recorded by a

Spectramax 384+ spectrophotometer at 600 nm using Softmax Pro version 6.4.2. Daily readings

were taken for nine days. Media for these tests included CM+ pH 6.3 100 mM PIPES and CM+

pH 6.1 100 mM PIPES for J clones. M clones were tested in CM+ pH 7.5 100 mM MOPS. K

clones were tested in CM+ pH 6.3 100 mM PIPES 600 µM FeSO$_4$ and CM+ pH 6.1 100 mM

PIPES 600 µM FeSO$_4$. S clones were tested in CM+ pH 7.5 100 mM MOPS 600 µM FeSO$_4$.

To test for pH-dependent growth advantages, evolved clones that showed growth

advantages over ancestor in their respective evolution stress conditions under which they were

evolved were also tested for growth advantages in pH conditions other than those in which they

evolved. For these experiments, J3-1 was cultured in CM+ pH 7.5 100 mM MOPS and compared

using a Vac$^+$ NRC-1 control, M3-1 was cultured in CM+ pH 6.1 100 mM PIPES and compared

using a Vac$^+$ NRC-1 control, and K2-1 was cultured in CM+ pH 7.5 100 mM MOPS 600 μM

FeSO$_4$ and compared to both Vac$^+$ and Vac$^-$ NRC-1 controls due to gas vesicle phenotype

ambiguity. Analysis was carried out with comparisons to an ancestral control expressing the

same Vac phenotype as the evolved clone.

All growth assays were evaluated for statistical significance using ANOVA test with

Tukey post-hoc or paired T-test using base R and agricolae package. Comparisons between

clones were made using post log-phase endpoint "E" values for optical density at six days post

inoculation.

DNA extraction and genome sequencing. Genomic DNA was isolated from the 16

evolved clones and the ancestor NRC-1 using an Epicentre MasterPure Gram Positive DNA

Extraction Kit and a modified procedure. Lysozyme was omitted, and DNA purity and

concentration was determined using a Thermo Scientific NanoDrop 2000. Genomic DNA was

sequenced at the Michigan State University Research Technology Support Facility (RTSF)

Genomics Core. Libraries were prepared using the Illumina TruSeq Nano DNA library

preparation kit for Illumina MiSeq sequencing and loaded on a MiSeq flow cell after library

validation and quantitation. Sequencing was completed using a 2- by 250-bp paired-end format

using Illumina 500 cycle V2 reagent cartridge. Illumina Real Time Analysis (RTA) v1.18.54

performed base calling, and the output of the RTA was demultiplexed and converted to FastQ

format with Illumina Bcl2fastq v1.8.4.

Sequence assembly and analysis using the *breseq* computational pipeline. The

computational pipeline *breseq* version 0.27.1 was used to assemble and annotate the resulting

Illumina reads of the evolved clones (42–44). The current *breseq* version is optimized to detect

IS element insertions and IS-mediated deletions, as well as SNPs and other mutations in *E. coli*

(19). Illumina reads were mapped to the *Halobacterium* sp. NRC-1 reference genome (NCBI

GenBank assembly accession GCA_000006805.1). Mutations were predicted by *breseq* through

sequence comparisons between the evolved and ancestral clones.

The Integrative Genomics Viewer (IGV) from the Broad Institute at Massachusetts

Institute of Technology was used to visualize the assembly and mutations in the evolved clonal

sequences mapped to the reference NRC-1 genome (60). Each replicon was mapped separately

using the following RefSeq IDs: NC_002607.1 (main chromosome), NC_001869.1 (pNRC100),

and NC_002608.1 (pNRC200). Sequence mean coverage in each evolved clone was estimated

using the *breseq* fit dispersion function.

PCR confirmation of ISH insertions. PCR primers (Table 5) were designed to confirm

the presence of insertion sequences at hypothetical target site duplications. Primers adhered to

the following specifications using Sigma Aldrich Oligo Evaluator: 19-22 bp in length, GC

content between 40-60%, no single bp runs >3, weak to no secondary structure, and no primer

dimer. Oligos were checked for sequence identity of ≤13 bp to any part of the NRC-1 genome

other than the target site using NCBI BLAST. We ran 50-µl PCR using Applied Biosystems

Amplitaq Gold 360 Master Mix according to the package insert with 50 µl reaction containing

GC enhancer. To assess insert length, 10 µl of PCR product was electrophoresed in a 1% agarose

gel. PCR products were then purified either by Qiagen QIAquick PCR Purification Kit or

QIAquick Gel Extraction Kit.

Accession number for sequenced genomes. Sequenced genomes are deposited under

SRA accession number SRP195828.

ACKNOWLEDGEMENTS

This project was supported by the National Science Foundation award MCB-1613278 to

Joan Slonczewski; by the 2016 NASA Astrobiology Program Early Career Collaboration Award

to Karina Kunka and Jessie Griffith; and by NASA Exobiology grants NNX15AM07G and

NNH18ZDA001N to Shiladitya DasSarma. We thank Friedhelm Pfeiffer for pointing out the

arcD annotation. We thank Landon Porter for expert technical assistance.

TABLES AND FIGURES
424

425   **Table 1. Strains used in this study**

Strain Name	Description	Generation	Evolution Condition	Vac +/- *	Source
NRC-1	Ancestor strain	0	--	+	S. DasSarma
NRC-1	Ancestor strain	0	--	-	S. DasSarma
JLSHA075	Clone J1	500	pH 6.3 100 mM PIPES	-	This study
JLSHA078	Clone J2-2	500		-	This study
JLSHA079	Clone J3-1	500		+	This study
JLSHA082	Clone J4-2	500		-	This study
JLSHA083	Clone M1	500	pH 7.5 100 mM MOPS	-	This study
JLSHA086	Clone M2-2	500		-	This study
JLSHA087	Clone M3-1	500		+	This study
JLSHA089	Clone M4-1	500		+	This study
JLSHA091	Clone K1	500	pH 6.3 100 mM PIPES 600 µM Fe^{2+}	-	This study
JLSHA093	Clone K2-1	500		+	This study
JLSHA095	Clone K3	500		-	This study
JLSHA097	Clone K4	500		-	This study
JLSHA099	Clone S1	500	pH 7.5 100 mM MOPS 600 µM Fe^{2+}	-	This study
JLSHA101	Clone S2	500		-	This study
JLSHA103	Clone S3	500		-	This study
JLSHA105	Clone S4	500		-	This study

426   *"+" indicates gas vesicle-forming, "-" indicates non gas vesicle-forming

Table 2. Selected mutations found in evolved clones.*□

Replicon	Start bp	pH 6.3				pH 7.5				pH 6.3, Fe				pH 7.5, Fe				Annotation	Gene Description
		J1	J2-2	J3-1	J4-2	M1	M2-2	M3-1	M4-1	K1	K2-1	K3	K4	S1	S2	S3	S4		
Chromosome	~15,266									■						■		ISH: intergenic	(vng0018H→) / (vng0019H→)
Chromosome	~23,074				■			■										ISH: intergenic	(vng0027H←) / (→vng0028C)
Chromosome	~25,216										■		■					ISH: intergenic	(vng0029H←) / (→vng0030H)
Chromosome	~29,154									■	■							ISH: coding	[vng0032H→]
Chromosome	~48,587		■		■	■				■						■		ISH8-3 mediated	(vng0053H→) / [→vng0053H]
Chromosome	~52,664												■		■	■		ISH: intergenic	(vng0056H←) / (→vng0057 O_antigen polymerase)
Chromosome	~181,512										■	■						ISH: coding	[vng0215C→]
Chromosome	414,229						■	■								■		A675A (GCG→GCT)	[vng0053H7←] (TRAP transporter permease)
Chromosome	~749,543			■			■						■					ISH: coding	[vng0985H←]
Chromosome	~753,552							■									■	ISH: intergenic and (T)9→8	(vng0986H→) / (→vng0987H)
Chromosome	~754,476						■										■	ISH: coding	[vng0989C←] xcd (integrase)
Chromosome	~772,459												■		■			ISH: intergenic	(vng1007H←) / (→ vng1008G) flaA1a (flagellin A1 precursor)
Chromosome	~1,089,129							■		■								ISH: coding	[vng1467G→] bop (rhodopsin).
Chromosome	~1,229,749		■										■					ISH: coding/ ISH2 deletion	[vng1650H←]
PNRC100	0			■						■	■							ISH7-1 deleted	[vng7001←] - [←vng_RS13745]
PNRC100	~9,546				■			■										ISH3-1	[vng_RS13755→] / (→vng_RS12260)
PNRC100	~14,052						■						■					ISH3-1	(vng_RS12260←) / (→vng7012)
PNRC100	~15,600	■	■		■	■				■		■	■	■	■	■		ISH8-3, 8-1	(vng7012→) - (←vng_RS12360) Gas vesicle protein cluster (deleted)
PNRC100	~41,820									■						■		ISH5-1*, 8-5	(vng7039→) - (→→vng_RS13770) yobE, XRE regulator, MFS transporter, thioredoxin, deoxyribonuclease, cell division control protein (all deleted)
PNRC100	~71,208			■			■			■								ISH2	(vng_RS13790→) - [←vng7073] SOS response peptidase (deleted)
PNRC100	~75,169				■			■										ISH2	(vng7074←) - (←vng7078)
PNRC100	~81,100						■					■	■					ISH8-3, 3-1	[vng_RS12615←] - [→vng7079]
PNRC100	~83,375						■					■	■					ISH3-1, 7-2	(vng7080←) - [←vng7085]
PNRC100	~87,224						■					■						V→L(GTG→CTG)	[vng7085←]
PNRC100	~133,744	■	■	■						■	■	■	■	■	■	■		ISH8-2, 3-3	(vng_RS12830←) - (→vng7127) Arsenic resistance operon (deleted)
PNRC100	~143,907	■	■	■						■	■	■	■	■	■	■		ISH3-3, 2	(vng_RS12880→) - [→vng7136] ssDNA-binding protein A (deleted)
PNRC100	~150,769									■	■		■					ISH2, 3-2	(vng7136←) / (←vng_RS12925)
PNRC100	~152,257									■		■						ISH2, 3-2	[vng_RS13845←]
PNRC100	~153,526									■		■	■			■		ISH3-2, 2	[←vng_RS13850]
PNRC100	~164,889									■						■		ISH8-5, 5-1	(vng_RS13865←) - (←vng7170) Thioredoxin reductase, boa3 (bacteriorhdopsin activator), phoT1 phosphate transporter), cydBA cytochrome d oxidase (all deleted).

Replicon	Start bp	J1	J2-2	J3-1	J4-2	M1	M2-2	M3-1	M4-1	K1	K2-1	K3	K4	S1	S2	S3	S4	Annotation	Gene Description
PNRC200	0			green				blue	blue	magenta	magenta							ISH 7-1 deleted	(vng6001H←) - [←vng6011H]
PNRC200	~7,477						blue									purple		ISH: coding	[vng6011H←]
PNRC200	~9,569				green		blue											ISH 3-1	[vng6013G→] / (←vng_RS10565)
PNRC200	~14052						blue			magenta			magenta					ISH 3-1	(vng_RS10565←) / (→vng6016H)
PNRC200	~15,594	green	green	green	green	blue				magenta		magenta	magenta	purple	purple	purple	purple	ISH 8-3, 8-1	(vng6016H→) - (→vng_RS10665)
PNRC200	~41,819									magenta	magenta	magenta	magenta			purple		ISH 5-1*, 8-5	(vng6053G→) - (→vng6079H) **Thioredoxin reductase, boa3 (bacteriorhdopsin activator), phoT1 phosphate transporter), cydBA cytochrome d oxidase (all deleted).**
PNRC200	~71,208			green			blue	blue		magenta								ISH 2	(vng6094H→) - [←vng6097C] *SOS response peptidase (deleted)*
PNRC200	~75,168																	ISH 2, 8-4	(vng6099H←) - (←vng6105H)
PNRC200	~81,101			green			blue					magenta	magenta					ISH 8-4, 3-1	[vng_RS10920←] / [→vng_RS10925]
PNRC200	~83,374						blue					magenta						ISH 3-1, 7-2	(vng6112H←) - [←vng6119H]
PNRC200	~87,224											magenta						V→L(GTG→CTG)	[vng6119H←]
PNRC200	~138,366	green						blue	blue						purple		purple	ISH 8-4	[vng6162H←] - [→vng6181H] *orc2* (cell division control protein 6), *nbp2* (nucleic acid binding protein), *srl1* (ATPase), *trkA2* (TRK K+ uptake system protein), *kdpABC* (K+ ATPase), *cat3* (cationic amino acid transporter) (all deleted).
PNRC200	~140,521				green											purple		ISH: coding	[vng6164G←] *orc2* (orc/cell division control protein 6)
PNRC200	~244,149	green		green						magenta	magenta	magenta	magenta					ISH: coding	*[vng6313G←] arcD* **Arginine:ornithine antiporter (formerly annotated nhaC3)**
PNRC200	~248,490	green								magenta	magenta		magenta					ISH: coding	*[vng6318G←] arcR* **(Arginine catabolism transcriptional regulator)**
PNRC200	~262,599			green				blue		magenta					purple			ISH 3-2, 8-3 / ISH 8-3 deleted	(vng_RS11685←) / (→vng6331H)
PNRC200	~272,000					blue	blue	blue										ISH 6, 3-2	(vng6341H→) / (←vng_RS13610)
PNRC200	~274,345						blue											ISH 3-2, 8-4	(vng_RS11735←) / (←vng6345H)
PNRC200	~278,031					blue	blue	blue	blue									ISH 8-4, 2	[vng6346H←] - (←vng6393H) **DNA polymerase I**, *cdc6* **cell division control protein , HNH endonuclease, MarR family transcriptional regulator, AbrB family transcriptional regulator, L-lactate permease (all deleted)**
PNRC200	~278,118										magenta							purple	ISH: coding	[vng6346H←]
PNRC200	~279,927										magenta							purple	ISH: intergenic	(vng6348H←) / (←vng6349C)
PNRC200	~293,402											magenta						purple	ISH: coding	[vng6364H←]
PNRC200	~309,253	green		green		blue	blue	blue		magenta								ISH 2, 8-3	(vng6393H→) - (←vng6395H)
PNRC200	~311,206	green	green	green		blue	blue	blue		magenta	magenta	magenta		purple	purple	purple	purple	ISH 8-3, 11	(vng6396H→) - (←vng6420H) **arsR transcriptional regulator, phzF phenazine biosynthesis**
PNRC200	~324,384	green		green		blue	blue	blue		magenta				purple	purple	purple	purple	ISH 11	(vng_RS12000→) - (→vng6441H)

* "Annotation" column code: "ISH *** mediated" = flanking ISH elements, if relevant. Mutation codes: blue = missense, green = silent.

"Gene" column code: (mutation starts or ends before this gene name), → ← indicates gene directionality, [mutation starts, ends, or is entirely contained within this gene name], "-" indicates intervening omitted genes found in description, "/" indicates mutation is between two genes. Highlight indicates mutations associated with acid evolution.

☐This chart does not indicate shared lineage through identical mutations. For a complete list of mutations, see **Tables S1-S3.**

Table 3. Classes of mutations found in evolved clones.*

Chromosome																
	Low pH				Control				Low pH and iron-rich				Iron-rich				
Mutation Type	J1	J2-1	J3-1	J4-2	M1	M2-2	M3-1	M4-1	K1	K2-1	K3	K4	S1	S2	S3	S4	Mutation Sum
TSD	4	5	3	5	1	4	4	1	7	6	5	5	1	6	6	1	64
Deletion	1	1	0	1	1	1	2	2	1	1	0	1	0	0	1	0	13
SNP	0	0	1	0	1	2	1	1	1	1	0	1	0	0	0	1	10
Insertion	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Chromosome Total	5	6	4	6	3	7	7	4	9	8	5	7	1	6	7	2	87

PNRC100																
Mutation Type	J1	J2-1	J3-1	J4-2	M1	M2-2	M3-1	M4-1	K1	K2-1	K3	K4	S1	S2	S3	S4	Mutation Sum
TSD	0	1	0	0	0	7	0	0	3	0	0	0	2	2	0	0	15
Deletion	5	3	4	7	2	0	6	3	17	8	8	10	4	3	5	4	89
SNP	2	0	0	0	0	6	0	0	0	0	3	5	0	0	0	0	16
Insertion	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
PNRC100 Total	7	4	4	7	2	13	6	3	20	8	11	15	6	5	5	4	120

PNRC200																
Mutation Type	J1	J2-1	J3-1	J4-2	M1	M2-2	M3-1	M4-1	K1	K2-1	K3	K4	S1	S2	S3	S4	Mutation Sum
TSD	1	1	2	1	1	4	0	0	3	7	3	7	2	0	3	5	40
Deletion	6	4	6	7	8	14	15	7	15	5	4	5	3	5	6	5	115
SNP	1	0	0	0	0	5	0	0	1	0	3	5	0	0	0	0	15
Insertion	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1
PNRC200 Total	9	5	8	8	9	23	15	7	19	12	10	17	5	5	9	10	171

Complete genome																
Mutation Type	J1	J2-1	J3-1	J4-2	M1	M2-2	M3-1	M4-1	K1	K2-1	K3	K4	S1	S2	S3	S4	Mutation Sum
TSD	5	7	5	6	2	15	4	1	13	13	8	12	5	8	9	6	119
Deletion	12	8	10	15	11	15	23	12	33	14	12	16	7	8	12	9	217
SNP	3	0	1	0	1	13	1	1	2	1	6	11	0	0	0	1	41
Insertion	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1
Complete Total	21	15	16	21	14	43	28	14	48	28	26	39	12	16	21	16	378

*TSD = target site duplication indicating ISH; SNP = single nucleotide polymorphism.

Table 4. Coverage depth for NRC-1 and evolved clones.

Strain	Main chromosome		pNRC100		pNRC200	
	Read depth*	SD	Read depth*	SD	Read depth*	SD
NRC-1	50	10	128	15	64	10
J1	46	9	338	29	53	9
J2-2	41	9	156	14	48	8
J3-1	57	11	87	13	91	14
J4-2	64	11	137	15	85	13
M1	49	9	187	18	44	7
M2-2	67	11	165	17	62	10
M3-1	49	10	275	26	54	9
M4-1	65	11	162	13	75	11
K1	74	13	72	10	63	11
K2-1	72	13	NA	NA	76	11
K3	52	10	363	29	92	13
K4	59	11	96	11	71	15
S1	39	8	125	15	44	8
S2	42	9	169	18	34	7
S3	47	10	202	16	55	9
S4	46	9	185	14	37	7

*Mean copy number of sequence across the replicon, according to the *breseq* fitted dispersion model. SD = standard deviation predicted by the model.

Table 5. ISH insertions confirmed by PCR in acid-adapted strains.

Strain	Gene Mutation	ISH	Primer 1	Primer 2
J1	*arcD* insertion	ISH4	GATAACGATGGACATGTACT	GTCGGTATCGTTCTTTTGGG
J3-1	*arcR* insertion	ISH8-2	ACTGGTGTGGAGTTTCCGTG	ATCTCACGATCAAGGACGGTGTT
J3-1	*bop* insertion	ISH1	GAGTTACACACATATCCTCG	GCGTAGAATTTCTTTGCATC
J4-2	*arcD* insertion	ISH11	GATAACGATGGACATGTACT	GTCGGTATCGTTCTTTTGGG
K1	*arcD* insertion	ISH2	GATAACGATGGACATGTACT	GTCGGTATCGTTCTTTTGGG
K1	*arcR*	ISH8-2	ACTGGTGTGGAGTTTCCGTG	ATCTCACGATCAAGGACGGTGTT
K2-1	*arcD* insertion	ISH4	GATAACGATGGACATGTACT	GTCGGTATCGTTCTTTTGGG
K3	*arcD* insertion	ISH4	GATAACGATGGACATGTACT	GTCGGTATCGTTCTTTTGGG
K4	*arcD* insertion	ISH2	GATAACGATGGACATGTACT	GTCGGTATCGTTCTTTTGGG
K4	*arcR*	ISH3-2	AGAAGTCGTTCAGAAACAGG	GATACGCGATCAACGACGA

Table 6. Acid-evolved clone J3-1 mutations.* [ ]

Replicon	Start bp	End bp	Mutation	Annotation	Description
Chromosome	749,943	749,954	(11 bp) 1→2	ISH: coding (562/2007 nt)	[vng0985H→]
Chromosome	1,089,129	1,089,137	(8 bp) 1→2	ISH: coding (15/789 nt)	[vng1467G→] **bop** (rhodopsin).
Chromosome	1,163,363		A→G	K197R (AAA→AGA)	[vng1561→] Ferredoxin
Chromosome	1,229,749	1,229,760	(11 bp) 1→2	ISH: coding (582/849 nt)	In vng1650.
PNRC100	0	7,788	Δ7788 bp	ISH7-1 deleted	[vng7001←] - [←vng_RS13745]
PNRC100	71,210	74,656	Δ3447 bp	ISH2 mediated	(vng_RS13790→) - [←vng7073] **SOS response peptidase (deleted)**
PNRC100	133,743	142,521	Δ8779 bp	ISH8-2, 3-3 mediated	(vng_RS12830←) - (→vng7127) **Arsenic resistance operon (deleted)**
PNRC100	143,909	150,253	Δ6345 bp	ISH3-3, 2 mediated	(vng_RS12880→) - [→vng7136] **ssDNA-binding protein A (deleted)**
PNRC200	0	7,760	Δ7760 bp	ISH 7-1 deleted	(vng6001H←) - [←vng6011H]
PNRC200	71,219	74,595	Δ3377 bp	ISH 2 mediated	(vng6094H→) - [←vng6097C] *SOS response peptidase (deleted)*
PNRC200	244,422	244,430	9 bp (1→2)	ISH: intergenic	(vng6313G←) / (vng6315G←) **arcD Arginine:ornithine antiporter / ornithine carbamoyltransferase**
PNRC200	249,147	249,157	11 bp (1→2)	ISH: coding	[vng6318G←] **arcR Arginine catabolism transcriptional regulator**
PNRC200	262,603	265,437	Δ2835 bp	ISH 3-2, 8-3 mediated/ ISH 8-3 deleted	(vng_RS11685←) / (→vng6331H)
PNRC200	309,256	309,812	Δ557 bp	ISH 2, 8-3 mediated	(vng6393H←) - (←vng6395H)
PNRC200	311,213	323,320	Δ12108 bp	ISH 8-3, 11 mediated	(vng6396H→) - (←vng6420H) **arsR transcriptional regulator,** *phzF* **phenazine biosynthesis**
PNRC200	324,386	332,792	Δ8407 bp	ISH 11 mediated	(vng_RS12000→) - (→vng6441H)

* "Annotation" column code: "ISH *** mediated" = flanking ISH elements, missense mutations in blue.

"Gene" column code: (mutation starts or ends before this gene name), → ← indicates gene directionality, [mutation starts, ends, or is entirely contained within this gene name], "-" indicates intervening omitted genes found in description, "/" indicates mutation is between two genes.

[ ]Highlight indicates mutations associated with acid evolution.

Table 7: Change in gas vesicle phenotype during evolution across populations*

Media condition	Strain	Generation				
		100	200	300	400	500
pH 6.3 100 mM PIPES	J1	+	Vac$^{+/-}$	-	-	-
	J2	+	Vac$^{+/-}$	-	-	Vac$^{+/-}$
	J3	>1% Vac$^-$	Vac$^{+/-}$	-	-	Vac$^{+/-}$
	J4	+	+	-	-	Vac$^{+/-}$
pH 7.5 100 mM MOPS	M1	+	+	+	Vac$^{+/-}$	Vac$^{+/-}$
	M2	+	+	+	Vac$^{+/-}$	Vac$^{+/-}$
	M3	+	+	+	Vac$^{+/-}$	Vac$^{+/-}$
	M4	+	>1% Vac$^-$	+	+	Vac$^{+/-}$
pH 6.3 100 mM PIPES 600 µM FeSO$_4$	K1	+	Vac$^{+/-}$	-	Vac$^{+/-}$	Vac$^{+/-}$
	K2	Vac$^{+/-}$	Vac$^{+/-}$	-	-	Vac$^{+/-}$
	K3	+	Vac$^{+/-}$	-	-	Vac$^{+/-}$
	K4	+	Vac$^{+/-}$	-	-	Vac$^{+/-}$
pH 7.5 100 mM MOPS 600 µM FeSO$_4$	S1	+	+	-	-	-
	S2	+	+	-	-	-
	S3	+	+	-	-	-
	S4	+	+	+	-	-

*   "+" indicates gas vesicle-forming, "–" indicates non gas vesicle-forming

Figure 1. Acid-evolved clone J3-1 shows a pH-dependent growth rate increase compared to NRC-1. Growth medium consisted of CM^+ buffered at (**A**) pH 6.1 with 100 mM PIPES; (**B**) pH 6.3 with 100 mM PIPES; (**C**) pH 7.2 with 100 mM MOPS; or (**D**) pH 7.5 with 100 mM MOPS. Representative curves of three replicates are shown. For J3-1 and NRC-1, the OD_{600} values at 144 h were compared by two-tailed t-test. At pH 6.1, P = 0.002; at pH 6.3, P = 0.01; at pH 7.2, P = 0.91; at pH 7.5, P = 0.45. "*" indicates significant endpoint growth increase from NRC-1 ancestor in at least 2 replicates.

Figure 2. Growth of acid-evolved clones J1, J2-2, J4-2. Growth medium consisted of CM^+ pH

6.3 with 100 mM PIPES, at (**A**) pH 6.1, (**B**) pH 6.3. Cultures were diluted from a 7-day culture

in CM^+ pH 7.2. Gas vesicle-deficient clones were compared to gas vesicle-deficient ancestral

mutant NRC-1 and cell density values post log-phase (OD_{600} at 6 days) were analyzed using

ANOVA with Tukey post-hoc. Representative curves of three replicates are shown.

REFERENCES

1.   Oren A. Life at high salt concentrations, intracellular KCl concentrations, and acidic proteomes. Front Microbiol. 2013;4(NOV):1–6.

2.   Reysenbach A, Pace NR. Archaea: A laboratory manual - Halophiles. DasSarma S, Fleischmann E, editors. Cold Spring Harbor: Cold Spring Harbor Laboratory Press; 1995.

3.   Ng W V., Kennedy SP, Mahairas GG, Berquist B, Pan M, Shukla HD, et al. Genome sequence of *Halobacterium* species NRC-1. Proc Natl Acad Sci [Internet]. 2000;97(22):12176–81. Available from: http://www.pnas.org/cgi/doi/10.1073/pnas.190337797

4.   Kish A, Kirkali G, Robinson C, Rosenblatt R, Jaruga P, Dizdaroglu M, et al. Salt shield: Intracellular salts provide cellular protection against ionizing radiation in the halophilic archaeon, *Halobacterium salinarum* NRC-1. Environ Microbiol. 2009;11(5):1066–78.

5.   Jones DL, Baxter BK. DNA repair and photoprotection: Mechanisms of overcoming environmental ultraviolet radiation exposure in halophilic archaea. Front Microbiol. 2017;8:1882.

6.   Coker JA, DasSarma P, Kumar J, Müller JA, DasSarma S. Transcriptional profiling of the model archaeon *Halobacterium* sp. NRC-I: Responses to changes in salinity and temperature. Saline Systems. 2007;3:doi:10.1186/1746-1448-3-6.

7.   Wang G, Kennedy SP, Fasiludeen S, Rensing C, DasSarma S. Arsenic resistance in *Halobacterium* sp. strain NRC-1 examined by using an improved gene knockout system. J Bacteriol [Internet]. 2004;186(10):3187–94. Available from: http://www.ncbi.nlm.nih.gov/pubmed/15126481%0Ahttp://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC400623

8.   DasSarma P, Laye VJ, Harvey J, Reid C, Shultz J, Yarborough A, et al. Survival of halophilic archaea in Earth's cold stratosphere. Int J Astrobiol. 2017;16(4):321–7.

9.   DasSarma P, DasSarma S. Survival of microbes in Earth's stratosphere. Curr Opin Microbiol. 2018;43:24–30.

10.   DasSarma P, Tuel K, Nierenberg SD, Phillips T, Pecher WT, Nrc- H, et al. Inquiry-driven teaching & learning using the archaeal microorganism Halobacterium. Am Biol Teach [Internet]. 2016;78(1):7–13. Available from: http://www.bioone.org/doi/pdf/10.1525/abt.2016.78.1.7

11.   DasSarma S. Extreme halophiles are models for astrobiology. Microbe [Internet]. 2006;1(3):120–6. Available from: https://www.asm.org/ccLibraryFiles/FILENAME/000000002127/znw00306000120.pdf

12.   Leuko S, Domingos C, Parpart A, Reitz G, Rettberg P. The survival and resistance of *Halobacterium salinarum* NRC-1, *Halococcus hamelinensis*, and *Halococcus morrhuae* to simulated outer space solar radiation. Astrobiology. 2015;15(11):987–97.

13.   Madden MEE, Bodnar RJ, Rimstidt JD. Jarosite as an indicator of water- limited chemical weathering on Mars. Nature [Internet]. 2004;431(August):821–3. Available from: https://www.nature.com/articles/nature02971.pdf

14.   Pritchett BN, Elwood Madden ME, Madden AS. Jarosite dissolution rates and maximum lifetimes in high salinity brines: Implications for Earth and Mars. Earth Planet Sci Lett. 2014;391:67–8.

15.   Dopson M, Ossandon FJ, Lövgren L, Holmes DS. Metal resistance or tolerance? Acidophiles confront high metal loads via both abiotic and biotic mechanisms. Front

Microbiol. 2014;5(157):doi: 10.3389/fmicb.2014.00157.

16. Moran-Reyna A, Coker JA. The effects of extremes of pH on the growth and transcriptomic profiles of three haloarchaea. F1000 Res. 2014;3(168):1–15.

17. Lenski RE, Travisano M. Dynamics of adaptation and diversification: A 10,000-generation experiment with bacterial populations. Proc Natl Acad Sci U S A [Internet]. 1994 Jul 19 [cited 2015 Sep 9];91(15):6808–14. Available from: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=44287&tool=pmcentrez&rendertype=abstract

18. Lenski RE, Rose MR, Simpson SC, Tadler SC. Long-term experimental evolution in *Escherichia coli*. I. Adaptation and divergence during 2,000 generations. Am Nat. 1991;138:1315–41.

19. Tenaillon O, Barrick JE, Ribeck N, Deatherage DE, Blanchard JL, Dasgupta A, et al. Tempo and mode of genome evolution in a 50,000 □ generation experiment. Nature [Internet]. 2016;536(7615):165–70. Available from: http://dx.doi.org/10.1038/nature18959

20. Creamer KE, Ditmars FS, Basting PJ, Kunka KS, Hamdallah IN, Bush SP, et al. Benzoate- and salicylate-tolerant strains of *Escherichia coli* K-12 lose antibiotic resistance during laboratory evolution. Appl Environ Microbiol. 2017;83(2):e02736.

21. He A, Penix SR, Basting PJ, Griffith JM, Creamer KE, Camperchioli D, et al. Acid evolution of *Escherichia coli* K-12 eliminates amino acid decarboxylases and reregulates catabolism. Appl Environ Microbiol. 2017;83:e00442-17.

22. Harden MM, He A, Creamer K, Clark MW, Hamdallah I, Martinez KA, et al. Acid-adapted strains of *Escherichia coli* K-12 obtained by experimental evolution. Schottel JL, editor. Appl Environ Microbiol [Internet]. 2015 Mar 15 [cited 2015 Sep 17];81(6):1932–41. Available from: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4345379&tool=pmcentrez&rendertype=abstract

23. Schou MF, Kristensen TN, Kellermann V, Schlötterer C, Loeschcke V. A *Drosophila* laboratory evolution experiment points to low evolutionary potential under increased temperatures likely to be experienced in the future. J Evol Biol. 2014;27(9):1859–68.

24. Slonczewski JL, Fujisawa M, Dopson M, Krulwich TA. Cytoplasmic pH measurement and homeostasis in bacteria and archaea. In: Advances in Microbial Physiology. 2009. p. 1–79, 317.

25. Kanjee U, Houry WA. Mechanisms of acid resistance in *Escherichia coli*. Annu Rev Microbiol. 2013;67:65–81.

26. Moore JP, Li H, Engmann ML, Bischof KM, Kunka KS, Harris ME, et al. Benzoate-tolerant *Escherichia coli* strains from experimental evolution lose the Gad regulon, multi-drug efflux pumps, and hydrogenases. Appl Environ Microbiol. 2019;

27. Hamdallah I, Torok N, Bischof KM, Majdalani N, Chadalavada S, Mdluli N, et al. Experimental evolution of *Escherichia coli* K-12 at high pH and RpoS induction. Appl Environ Microbiol. 2018;84(13):e00520.

28. DeVeaux LC, Müller JA, Smith J, Petrisko J, Wells DP, DasSarma S. Extremely radiation-resistant mutants of a halophilic archaeon with increased single-stranded DNA-binding protein (RPA) gene expression. Radiat Res. 2007;168(4):507–14.

29. McCarthy S, Johnson T, Pavlik BJ, Payne S, Schackwitz W, Martin J, et al. Expanding the limits of thermoacidophily in the archaeon *Sulfolobus solfataricus* by adaptive evolution.

Appl Environ Microbiol. 2016;82(3):857–67.

30. Ai C, McCarthy S, Eckrich V, Rudrappa D, Qiu G, Blum P. Increased acid resistance of the archaeon, *Metallosphaera sedula* by adaptive laboratory evolution. J Ind Microbiol Biotechnol. 2016;43(10):1455–65.

31. Dong Y, Chen YYM, Burne RA. Control of Expression of the Arginine Deiminase Operon of Streptococcus gordonii by CcpA and Flp. J Bacteriol. 2004;

32. Sakanaka A, Kuboniwa M, Takeuchi H, Hashino E, Amano A. Arginine-ornithine antiporter ArcD controls arginine metabolism and interspecies biofilm development of Streptococcus gordonii. J Biol Chem. 2015;

33. Krulwich TA, Sachs G, Padan E. Molecular aspects of bacterial pH sensing and homeostasis. Nat Rev Microbiol [Internet]. 2011;9:330–43. Available from: http://dx.doi.org/10.1038/nrmicro2549

34. Lund P, Tramonti A, De Biase D. Coping with low pH: Molecular strategies in neutralophilic bacteria. FEMS Microbiol Rev. 2014;38(6):1091–125.

35. Simsek M, DasSarma S, RajBhandary U, Khorana H. A transposable element from *Halobacterium halobium* which inactivates the bacteriorhodopsin gene. Proc Natl Acad Sci. 2006;79(23):7268–72.

36. Dummer AM, Bonsall JC, Cihla JB, Lawry SM, Johnson GC, Peck RF. Bacterioopsin-mediated regulation of bacterioruberin biosynthesis in *Halobacterium salinarum*. J Bacteriol. 2011;193(20):5658–67.

37. Ng W V., Ciufo SA, Smith TM, Bumgarner RE, Baskin D, Faust J, et al. Snapshot of a large dynamic replicon in a halophilic archaeon: Megaplasmid or minichromosome? Genome Res. 1998;8:1131–41.

38. DasSarma S. Mechanisms of genetic variability in *Halobacterium halobium*: The purple membrane and gas vesicle mutations. Can J Microbiol. 1989;35(1):65–72.

39. Sapienza C, Doolittle WF. Unusual physical organization of the *Halobacterium* genome. Nature. 1982;295(5848):384–9.

40. Wimmer F, Oberwinkler T, Bisle B, Tittor J, Oesterhelt D. Identification of the arginine/ornithine antiporter ArcD from *Halobacterium salinarum*. FEBS Lett. 2008;582:3771–5.

41. Ruepp A, Soppa J. Fermentative arginine degradation in *Halobacterium salinarium* (formerly *Halobacterium halobium*): Genes, gene products,and transcripts of the *arcRACB* gene cluster. J Bacteriol. 1996;178(16):4942–7.

42. Deatherage DE, Barrick JE. Identification of mutations in laboratory-evolved microbes from next-generation sequencing data using *breseq*. Sun L, Shou W, editors. Methods Mol Biol [Internet]. 2014;1151:165–88. Available from: http://link.springer.com/10.1007/978-1-4939-0554-6

43. Deatherage DE, Traverse CC, Wolf LN, Barrick JE. Detecting rare structural variation in evolving microbial populations from new sequence junctions using *breseq*. Front Genet. 2015;5(JAN):1–16.

44. Barrick JE, Colburn G, Deatherage DE, Traverse CC, Strand MD, Borges JJ, et al. Identifying structural variation in haploid microbial genomes from short-read resequencing data using *breseq*. BMC Genomics. 2014;15(1):1039.

45. Pfeiffer F, Schuster SC, Broicher A, Falb M, Palm P, Rodewald K, et al. Evolution in the laboratory: The genome of *Halobacterium salinarum* strain R1 compared to that of strain NRC-1. Genomics. 2008;91(4):335–46.

46.   Pfeiffer F, Marchfelder A, Habermann B, Dyall-Smith ML. The Genome Sequence of the Halobacterium salinarum Type Strain Is Closely Related to That of Laboratory Strains NRC-1 and R1 . Microbiol Resour Announc. 2019;

47.   DasSarma S, Halladay JT, Jones JG, Donovan JW, Giannasca PJ, de Marsac NT. High-frequency mutations in a plasmid-encoded gas vesicle gene in *Halobacterium halobium*. Proc Natl Acad Sci U S A [Internet]. 1988;85(18):6861–5. Available from: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=282078&tool=pmcentrez&rendertype=abstract

48.   Ng W, DasSarma S. Physical and genetic mapping of the unstable gas vesicle plasmid in Halobacterium halobium NRC-1. In: Rodriguez-Valera F, editor. General and Applied Aspects of Halophilic Microorganisms. Plenum Press; 1991. p. 305–11.

49.   Ng WL, Kothakota S, DasSarma S. Structure of the gas vesicle plasmid in Halobacterium halobium: Inversion isomers, inverted repeats, and insertion sequences. J Bacteriol. 1991;173:1958–64.

50.   Ng WL, Arora P, DasSarma S. Large deletions in class III gas vesicle-deficient mutants of *Halobacterium halobium*. Syst Appl Microbiol. 1993;16:560–8.

51.   Soppa J. Evolutionary advantages of polyploidy in halophilic archaea. Biochem Soc Trans. 2013;41(1):339–43.

52.   Gong S, Richard H, Foster JW. YjdE (AdiC) is the arginine:agmatine antiporter essential for arginine-dependent acid resistance in *Escherichia coli*. J Bacteriol. 2003;185(15):4402–9.

53.   DasSarma S, Arora P. Genetic analysis of the gas vesicle gene cluster in haloarchaea. FEMS Microbiol Lett. 1997;153(1):1–10.

54.   DasSarma P, Zamora RC, Müller JA, DasSarma S. Genome-wide responses of the model archaeon *Halobacterium* sp. strain NRC-1 to oxygen limitation. J Bacteriol. 2012;194(20):5530–7.

55.   DasSarma S, Arora P, Lin F, Molinari E, Yin LRS. Wild-type gas vesicle formation requires at least ten genes in the gvp gene cluster of *Halobacterium halobium* plasmid pNRC100. J Bacteriol. 1994;176(24):7646–52.

56.   Nelson-Sathi S, Dagan T, Landan G, Janssen A, Steel M, McInerney JO, et al. Acquisition of 1,000 eubacterial genes physiologically transformed a methanogen at the origin of Haloarchaea. Proc Natl Acad Sci. 2012;109(50):20537–42.

57.   Redder P, Garrett RA. Mutations and rearrangements in the genome of *Sulfolobus solfataricus* P2. J Bacteriol. 2006;188(12):4198–206.

58.   Griffith JM, Basting PJ, Bischof KM, Wrona EP, Kunka KS, Tancredi A, et al. Experimental evolution of *Escherichia coli* K-12 in the presence of proton motive force (PMF) uncoupler carbonyl cyanide m-chlorophenylhydrazone selects for mutations affecting PMF-driven drug efflux pumps. Appl Environ Microbiol. 2019;85(5):e02792-18.

59.   Berquist BR, Müller JA, DasSarma S. 27 genetic systems for halophilic archaea. Methods Microbiol. 2006;35:649–80.

60.   Thorvaldsdóttir H, Robinson JT, Mesirov JP, Thorvaldsdóttir H, Robinson JT, Mesirov JP. Integrative genomics viewer (IGV): High-performance genomics data visualization and exploration. Brief Bioinform. 2013;14(2):178–92.