The emergent Yo-yo movement of nuclei driven by collective cytoskeletal remodeling in pseudo-synchronous mitotic cycles

Zhiyi Lv[1, 7, 8, 9], Jan Rosenbaum[2], Stephan Mohr[3], Xiaozhu Zhang[4], Deqing Kong[1, 6], Helen Preiß[5], Sebastian Kruss[5], Karen Alim[3], Timo Aspelmeier[2], Jörg Großhans[1, 6]

1. Institut für Entwicklungsbiochemie, Universitätsmedizin, Georg-August-Universität Göttingen, Germany.
2. Institut für Mathematische Stochastik, Georg-August-Universität Göttingen, Germany.
3. Germany Max Planck Institute for Dynamics and Self-Organization, Göttingen, Germany,
4. Chair for Network Dynamics, Institute for Theoretical Physics and Center for Advancing Electronics Dresden (cfaed), Cluster of Excellence Physics of Life, TU Dresden, 01062 Dresden,
5. Institut für Physikalische Chemie, Georg-August-Universität Göttingen, Germany
6. Fachbereich Biologie, Philipps-Universität Marburg, Germany
7. Institute of Evolution and Marine Biodiversity, Ocean University of China, Qingdao, 266003, China.
8. Ministry of Education Key Laboratory of Marine Genetics and Breeding, College of Marine Life Sciences, Ocean University of China, Qingdao, 266003, China
9. Laboratory for Marine Biology and Biotechnology, Qingdao National Laboratory for Marine Science and Technology, Qingdao, 266237, China.

corresponding author:
Zhiyi Lv
lv.zhiyi@med.uni-goettingen.de
Jörg Großhans
jgrossh@gwdg.de

Abstract:

Many aspects in tissue morphogenesis are attributed to the collective behavior of the participating cells. Yet, the mechanism for emergence of dynamic tissue behavior is not understood completely. Here we report the "yo-yo"-like nuclear drift movement in *Drosophila* syncytial embryo displays typical emergent feature of collective behavior, which is associated with pseudo-synchronous nuclear division cycle. We uncover the direct correlation between the degree of asynchrony of mitosis and the nuclear collective movement. Based on experimental manipulations and numerical simulations, we find the ensemble of spindle elongation, rather than a nucleus' own spindle, is the main driving force for its drift movement. The cortical F-actin acts as viscoelastic medium to dampen the movements and plays a critical role in restoring the nuclear positions after a mitosis cycle. Our study provides insights into how the interactions between cytoskeleton as individual elements leads to collective movement of the nuclear array on a macroscopic scale.

Keywords: syncytium, nuclear array, cortex, microtubule, F-actin, formin, Dia, network, emergent behavior, ELMO, flow, biomechanics

Introduction

Collective behaviors emerge from many interacting individuals in the absence of central coordination and supervision. Birds flocks, fish schools, as well as bacterial colonies exhibit spontaneous synchronization and many other complex dynamic functional patterns originating from simple local interaction rules [1-3]. Within a single organism, self-organized collective behaviors are an inseparable part in maintaining its basic biological functions by efficiently driving numerous complex biological processes without external regulations such as neural activation in brain [4-6] and morphogenesis during embryonic development [7,8]. On subcellular scale, actin and microtubules form collective motion in the presence of motor proteins and ATP [9-11]. Yet, despite the ubiquitousness and the significance of emergent collective behavior in biology, our knowledge of the fundamental mechanism of functional collective behaviors are very limited. The major obstacle lies in the typically extremely complex interactions between the individuals in most biological collective behaviors. However, in syncytial insect embryo only direct cytoskeletal interactions exist, owing to the lack of membranes separating the nuclei, which enables dissecting and understanding how the interactions of individuals drive the formation of emergent features at tissue level.

To obtain insights into the molecular mechanism leading to the emergence of a collective flow movement, we study the dynamics of nuclear array in *Drosophila* syncytial blastoderm, where the direct interactions between individuals lead to at least three features of emergent collective behavior observed at tissue level. Firstly, the nuclei divide synchronously to their immediate neighbors but asynchronously to more distant nuclei. This leads to a wave front of mitosis sweeping over the embryo [12,13]. Secondly, the nuclei arrange in an ordered array in interphase following disturbance during nuclear division. Ordering involves interactions by the microtubule asters but also with F-actin at the cortex [14,15]. Thirdly, the nuclei and the cytoplasm undergo stereotypic flow movements following the mitotic wave, which is reminiscent of the toy "yo-yo". This arises the questions: how does the flow emerge and what are the underlying driving forces? Is there any physiological function of this flow?

The large-scale collective movements are driven by active elements of the cytoskeleton. For example, kinesin-1 and microtubules drive the cytoplasmic streaming during *Drosophila* oogenesis [16]. The cytoplasmic streaming in plant and algae is driven by myosins moving on F-actin [17,18]. Apical constriction mediated by actomyosin can generate the cytoplasmic flow, which

95 compels nuclear spreading in *Drosophila* pre-blastoderm [19] and cell
96 elongation in gastrulation [20].
97
In *Drosophila* syncytial blastoderm, the nuclei and their associated
centrosomes and microtubule asters form an extended two-dimensional array
[14]. The dynamics of this array is dominated by two different interactions [21].
Lateral interactions between the neighbor nuclei and centrosomes are
mediated by microtubules and the associated motor proteins. The interactions
between centrosomes and actin cortex constitute the cortical interactions.
How these interactions lead to the stereotypic nuclear movement is unknow.
105
106 Beside the force generating mechanism, the material properties of the
107 embryos [22-24] may influence the movement of the nuclei. The actin
108 cytoskeleton inhibits short time-scale movements [21] and promotes ordering
109 of the nuclear array [14]. The actin cortex may act as a viscoelastic medium,
110 to which the centrosomes and their associated nuclei are connected.
111
Here we found that the nuclear movement was isotropic for individual spindles
but anisotropic for the collective flow over several nuclear diameters away
from the mitotic wave front and back to the original position slightly later. We
comprehensively quantified nuclear trajectories in wild type and mutant
embryos and modelled the process by computational simulation. In this way,
we uncovered that the ensemble of spindle elongation, rather than a nucleus'
own spindle, is the main driving force for its drift movement. In addition, we
defined a function of cortical F-actin for the apparent viscoelastic material
properties to restore the nuclear position. Lastly, based on a simulation, we
proposed that such nuclear movement is prerequisite for nuclear division in
high density in Drosophila syncytial blastoderm.
123

Results

Collective flow and density changes follow the mitotic wave front of
pseudo-synchronous nuclear cycles
To elucidate the mechanisms that arise the emergent features of nuclear
array dynamics in *Drosophila* syncytial cleavage cycle, we first documented
nuclear division cycles using time-lapse microscopy. The nuclei divide slightly
asynchronous especially during the last syncytial division in NC13 with a time
lag of up to minutes, which is easily visible as a mitotic wave front sweeping
over the embryo (Fig. 1A, Supp. Data Fig. S1A, Movie 1). The mitotic wave is
driven by Cdk1 activity wave and the molecular processes of the how the
chemical wave propagates were dissected [12,25].

The mitotic pseudo-synchrony and its corresponding wave front are associated with a stereotypic nuclear movement, which can be readily observed in time lapse movies (Fig. 1B, C, Supp. Data Movie 2). To obtain a precise description of the nuclear movements, we determined the trajectories of all nuclei within the field of view from time lapse recordings of fluorescently labeled nuclei. From the trajectories, we extracted time courses for displacements, velocities, and nuclear density (Fig. 1D, E, Supp. Data Fig. S1B, C). Every nucleus is assigned an individual time axis with the splitting of daughter chromosomes (metaphase-anaphase transition) as reference time t=0.

Concerning displacement (Fig. 1D), the nuclei moved in average about 20 μm which corresponds to about 4–5 nuclear diameters away from the position of their mother nucleus at t=0. The maximal displacement was reached after about 2 min (Fig. 1D). Following maximal displacement, the nuclei then returned to almost their initial position. This movement resembles yo-yo ball, and hereafter we refer it as yo-yo movement. We calculated the speed of nuclear movement from the individual trajectories as the derivative of the trajectories. The averaged flow speed revealed three peaks (Fig. 1E). The first peak corresponds to the chromosome segregation in anaphase with about 0.4 μm/s. The second peak after about 1 to 2 min corresponds to the flow away from the mitotic wave front. The least pronounced, third peak corresponds to the return movement after about 4 min.

Asynchronicity of mitosis might be prerequisite for yo-yo flow. To test whether there is a correlation, we collected data from 50 embryos (Fig. S1A) for quantification of the speed of wave front and maximal displacement. Plotting corresponding parameter sets revealed a negative correlation (R^2=45%, Fig. 1F), which suggests a slow wave is associated with a large displacement.

As flow is linked to density changes, we next established spatial and temporal maps of nuclear density. Each nucleus was assigned an area and corresponding density according to Voronoi segmentation (Supp. Data Fig. S1B, Movie 3). In the case of synchronous divisions, the density would be expected to double at t=0 (metaphase anaphase transition) and remain constant throughout interphase. In contrast, but consistent with the observed nuclear flow, our measurements revealed a peculiar time of the density. Although initially doubling, the density dropped in telophase before finally reaching the doubled density again a few minutes later (Supp. Data Fig. S1B, C). Corresponding profiles for displacement and flows were detected in preceding nuclear cycles 11 and 12, although in a less pronounced manner (Supp. Data Fig. S2).

The forth and back movement of the nuclei is reminiscent of a spring (Supp.
Data Fig. S1D). To obtain a phenomenological description of this behavior, we
applied a simple mechanical model to the nuclear trajectories during the
period of maximal displacement. By fitting a square function to the
displacement curve, we obtained an apparent spring constant for each
nucleus (Supp. Data Fig. S1E). The actual value of the apparent spring
constant is not informative, since our model and assumptions are too simple.
Friction is not included, for example. Yet, the apparent constant helps to
compare experimental conditions and mutant phenotypes.

Isotropic individual behavior is associated with an anisotropic collective
flow
Our analysis revealed a collective directional flow of the nuclear array. Yet the
individual spindles are isotropically oriented. The axes between daughter
nuclei are uniformly distributed over the angles against the anterior-posterior
axis of the embryo (Fig. 2A–C). In contrast, the same nuclei almost
unidirectionally moved along the embryonic axis a minute later, indicating an
anisotropic behavior (Fig. 2A–C). The transition of isotropic individual to
anisotropic collective behavior is strikingly obvious in the extreme cases of
spindle orientation. In the case of a spindle oriented in parallel to the
embryonic axis, one set of chromosomes segregated towards, whereas the
daughter chromosomes moved away the wave front during anaphase. One
minute later both nuclei moved away from the wave front (Fig. 2D). Similarly,
in case of a perpendicular orientation of chromosome segregation, the
movement during anaphase was perpendicular to the embryonic axis but
along the axis a minute later during collective flow. (Fig. 2D).

The uncoupling of spindle behavior and nuclear behavior was seen not only in
the orientation of the movements, but also in the time course of the two
behaviors. The maximal displacement was reached only about 1.5 min after
the maximal distance between daughter nuclei (maximal spindle length) was
reached (Fig. 2E, F), suggesting the maximal displacement was achieved
when the nuclei are in telophase. These findings demonstrated that these two
processes were mechanistically not directly linked at the individual level, since
spindles were isotropically oriented and preceded the flow behavior (Fig. 2G).

During collective flow the nuclei may move as individuals characterized by
neighbor exchanges. Alternatively, nuclei may behave as an array, which
would be indicated by fixed neighbor relationships. To distinguish these
options, we labeled groups of cells before mitosis and followed them during
the course of chromosome segregation and collective flow. We found that the

nuclei moved as an array. The groups of nuclei did neither intersperse with
unlabeled nuclei nor nuclei of the other groups indicating that neighbor
relationships were maintained during mitosis and collective flow despite the
movement over several nuclear diameters (Supp. Data Fig. S3A). In addition,
we measured the displacement fields of nuclei using a particle image velocity
algorithm. We found that the nuclei motion is similar to a laminar flow (Supp.
Data Fig. S3B, Movie 4). In summary, our observations indicate that the
nuclear layer phenomenologically behaves like an elastic sheet with fixed
neighbor relationships.
229
Computational modeling of nuclear movement
To gain a better understanding for how the isotropic spindles gives rise to an
anisotropic collective flow, we conducted computational simulations. Starting
from a computer model for static nuclear interactions in interphase [15], we
added a time axis for the interactions. The model is based on active and
passive forces (Fig. 3A). Stochastic active forces repulse adjacent nuclei, thus
resembling the sliding activity of motor proteins, e. g. Kinesin-5, on antiparallel
aligned microtubules. In addition, a passive elastic force leads to repulsion
accounting for the embedding of the nuclei into the cytoplasm and
cytoskeleton. This may include the link of the nuclei to the cortex.
Chromosome segregation is triggered at t=0 by a separation force acting
between the daughter nuclei. The interaction forces are dynamic according to
the mitotic stage and interphase (Supp. Data Fig. S4). For example, the active
force is low in anaphase, since astral microtubules prominently appear only in
telo- and interphase. Similarly, the passive force increases in telo- and
interphase as cortical actin increases during these stages. The segregation
force decays in telo and interphase. Balancing the magnitude of passive and
active forces over time, model simulation reproduced the experimentally
observed stereotypic flow behavior (Fig. 3B, C, Supp. Data Movie 5).
249
Strikingly, the simulations reproduce other features of nuclear yo-yo
movement. Firstly, maximal spindle length precedes maximal displacement
with a time lag of 1.5 min (Fig. 3D, E vs Fig. 2E, F). Secondly, the speed of
the mitotic wave front is negatively correlated with the nuclear maximal
displacement (Fig. 3F vs Fig. 1F). In addition, the simulations predict no
collective movement of nuclei with asynchronous nuclear divisions in a "salt-
and-pepper" pattern (Fig. 3G). Importantly, the simulations predict that the
force for separation of the daughter nuclei positively correlates with maximal
distance between daughter nuclei and importantly with maximal displacement
(Fig. 3H). Thus, our simulations predict that spindle elongation is a major
driving force for nuclear movement. Mechanistically, this is not a simple

relationship because spindles are isotropically oriented, whereas the direction
of displacement is anisotropic.
263
Ensemble spindle elongation is the driving force for nuclear
displacement
To test the prediction that spindle elongation drives the yo-yo movement, we
preformed the laser cutting on single spindle. However, the spindle recovered
in second-scale and no effects on nuclear motion (Supp. Data Fig. S5). To
circumvent this problem, we developed a method to reduce the spindle length
globally, without hampering the nuclear separation. Spindle elongation in
anaphase B requires the four-headed microtubule motor Kinesin 5, which can
slide microtubules against each other [26]. We employed embryos, in which
endogenous Kinesin 5 was substituted by a version susceptible to TEV
protease [27]. We titrated the amount of TEV protease to achieve a partial
depletion which still allowed completion of mitosis. Spindles in these embryos
were short (Fig. 4A). The average maximal spindle length was 8 µm instead
of 10 µm in wild type embryos. Complementary, we employed embryos from
females homozygous for *Map60*, which also displayed short spindles [14] with
an average length of 9 µm (Fig. 4A). Quantification of nuclear movement
revealed a strongly reduced maximal displacement in both experimental
conditions and thus a positive correlation of spindle length and maximal
displacement (Fig. 4B, C, D). In summary, analysis of mutant embryos with
shorter division distance support an ensemble spindle elongation constitutes a
major driving force for the nuclear movement thereafter.
285
F-actin cortex is required for the return movement.
The simulation predicts a function of passive force in yo-yo movement.
Cortical F-actin is a promising candidate. It has been previously reported that
the nuclei are strongly attached to F-actin cortex [28]. The actin cortex
suppresses the fluctuation movements of centrosomes [21] and contributes to
an ordered nuclear array in interphase [14]. Cortical F-actin undergoes
stereotypic remodeling during the course of nuclear cycles [29] (Fig. 4E,
Supp. Data Fig. S6A). We quantified total F-actin with a Utrophin-GFP as a
marker [30]. We found that the signal dropped in mitosis, with lowest level
during anaphase and steadily increased afterwards (Supp. Data Fig. S6A, B).
Given the timing of this dynamics, it is conceivable that cortical actin plays an
important part in controlling nuclear movement.
298
299 To test this conceivable function of the cortical F-actin, we employed two
300 mutants to genetically interfere with the organization of the actin cortex (Fig4.
301 F, Supp. Data Fig. S6G). Firstly, we prevented the formation of actin caps with
302 the mutant *ELMO* [21,31]. ELMO forms part of an unconventional guanyl

nucleotide exchange factor, which activates Rac signaling in a complex with Sponge/DOCK [32]. *ELMO* mutants lack any actin caps and are characterized by a uniformly structured cortical F-actin [21] (Supp. Data Fig. S6G). Secondly, we employed *dia* mutants [33-35]. Dia is a founding member of the formin family, which nucleate and polymerize linear actin filaments. *dia* mutants lack metaphase furrows but contain actin caps [33] (Supp. Data Fig. S6G).

We applied our quantitative assay to *ELMO* embryos. A strongly increased nuclear mobility was obvious in time lapse movies (Fig. 4G). Quantification of nuclear trajectories revealed a maximal nuclear displacement of 60 µm as compared to 20 µm in wild type embryos (Fig. 4H, Supp. Data Fig. S6H). In addition to the threefold increased displacement, we observed as a second phenotype that the nuclei did not return to their initial position in *ELMO* mutants (Fig. 4G, H). The impaired back movement indicates a loss of the spring-like behavior. Consistently, we calculated an almost 10-fold reduced spring constant at the turning point of the nuclear trajectories (Supp. Data Fig. S6I). A similar behavior and profiles were detected in NC12 of ELMO embryos (Supp. Data Fig. S6J, K).

We also detected changes in nuclear movement in *dia* mutants. Similar to *ELMO* mutants, we observed a loss of the spring-like back movement. The nuclei did not return to their initial position and the spring constant was almost 10-fold reduced (Fig. 4I, J, Supp. Data Fig. S6H, I). In contrast to *ELMO*, the maximal displacement was similar to wild type indicating that the stabilizing/viscos function of the cortex does not depend on *dia*. The neighborhood relationships were largely maintained in *dia* and *ELMO* embryos (Supp. Data Fig. S7). In summary, by employing two mutants affecting F-actin organization, we identified distinct functions of the actin cortex. F-actin is required for the back movement as revealed by the reduced apparent spring constant and the permanent displacement. The *ELMO*-dependent organization into caps appears to be important for limiting nuclear movement.

Long mitotic spindles and distance between daughter nuclei require pseudo-synchronous nuclear cycles

To obtain the further insight into the mitotic wave and consequent yo-yo movement in syncytial embryo, we documented the nuclear separation profile. We found that in mitosis the daughter nuclei are separated by an overshooting spindle, which pushes apart the daughter nuclei more than the average inter-nuclear distance (Fig. 5A, B). With every division during the

syncytial blastoderm (nuclear cycles NC 10–13), the nuclear density at the
cortex doubles (Supp. Data Fig. S8). The nuclei divide in a wave manner
especially during the last syncytial division. The mitotic wave is driven by
Cdk1 activity wave, and the activity of DNA replication checkpoint is important
in the slowdown of the wave, which occurs in later cycles [12,25]. However,
the function of this pseudo-synchrony and the consequent yo-yo movement is
unknown. Given the increase of asynchrony with nuclear density, we
speculated that spindle overshooting with nuclear crowding may pose a
problem for synchronous divisions.
354
To test this hypothesis, we employed the *mei41 zld* double mutant embryos
[36], as well as the *grp nmk* double mutant embryos [12], to experimentally
reduce the nuclear division time in cycle 13. We planned to check how much
the spindles were able to elongate in a similar level of nuclear crowding as in
wild type embryo, but with a more synchronous division manner.
Unfortunately, the chromosomes were not separated at cycle 13 in these
mutant embryos (Supp. Data Fig. S9).   To circumvent this problem, we
simulated nuclear divisions within a limited area. We assigned each nucleus
and mitotic spindles a protected area representing the entity in real embryos
(Fig. 5C). After chromosome division, the daughter nuclei were pushed apart
until reaching the protected area of a neighboring spindles/pair of daughter
nuclei, thus assessing the maximal possible distance between daughter nuclei
(maximal spindle length). We assumed synchronous divisions and symmetric
spindles with isotropic orientations (Fig. 5D). Our simulations showed that the
maximal spindle length decreased with an increase in nuclear number and
thus marked the transition line between structurally allowed and forbidden
regime of combinations of nuclear density and spindle length (Fig. 5E). Next,
we measured the maximal distance between daughter nuclei and their
corresponding nuclear densities in wild type embryos. We also included data
from haploid embryos, which undergo an extra nuclear division. Only the
parameters of NC11 fell into the allowed area, whereas the parameters of
NC12, NC13, and NC14 in haploids fell into the forbidden regime. This
analysis indicated that a synchronous division with the observed spindle
lengths was possible only in the early cycles but impossible in later cycles.
Thus, pseudo-synchrony allows for the observed spindle length in NC12,
NC13 and NC14 in haploid embryos.
381
Discussion
The direct interactions between the nuclei and their associated cytoskeleton
are a special feature of syncytial embryos. Due to the lack of separating cell
membranes, microtubule asters originating from the centrosomes associated
with each nucleus form an extended network of hundreds to thousands of

elements. Emergent features arise in this network by summing up the
behavior of individual elements, such as fluctuations or duplication, and their
interactions, such as repulsion. The analysis of the mechanism underlying the
emergent features is essential for understanding how the individual cells
function collectively to form a tissue.
392
We identified an anisotropic flow of the nuclear array as an emergent feature.
Based on a morphodynamical analysis of the nuclear array in wild type and
mutant embryos together with computational simulations, we analyzed the
mechanism of the flow behavior. In this way we identified spindle elongation
drives nuclei moving away whereas cortical F-actin restores the nuclear
positions, which is necessary for the following development. The emergent
nature of the nuclear flow becomes obvious, since individual behavior is
strikingly different than the collective behavior of the nuclear array. Nuclei
divide with an isotropic orientation, whereas the flow direction is anisotropic.
Furthermore, the maximal division distance is about 10 μm, whereas the
maximal displacement is about 20 μm.
404
In addition to the driving force of the nuclear yo-yo movement, our "limited
area" simulation (Fig. 5C, D, E) demonstrates the necessity of such
movement. To complete the nuclear division in a limited space with high
nuclear density, two strategies could be utilized. The first strategy is to reduce
the spindle length, as what happens in *Map60* mutant and Kinesin-5 partially
depleted embryos. This might raise the risk that the genetic materials cannot
separate completely. The second strategy is that nuclei divide in an
asynchronous manner. Besides *Drosophila*, the nuclear division asynchrony
was observed in beetle *Tribolium castaneum* [37], implying this might be a
conserved mechanism in insect syncytial embryos. The nuclear directional
movement is the consequence of the asynchronous divisions. With strongly
synchronous divisions, the pushing forces of mitotic spindles would generate
a spatially isotropic force distribution. Consequently, the nuclei would not
move due to a balance in forces. However, in the case of pseudo-
synchronous divisions, the force balance is broken leading to an imbalance
and thus a flow away from the wave front. The repulsive force between
daughter nuclei increases in anaphase pushing the daughter nuclei apart,
followed by a drop in telophase due to spindle disassembly (Supp. Data Fig.
S11A). The summing up of all nuclei in an embryo at a given mitotic time
results in an asymmetric force field, which likely determines the directionality
of the nuclear flow in telophase (Supp. Data Fig. S11B).
426
Upon the nuclei reach the maximal displacements, they return to the starting
positions. Consistency of nuclear position among different cleavage cycles is

important for maintaining the positioning information provided by morphogens.
We identified a contribution of the actin cortex to the viscoelastic feature of the
nuclear movement, i.e. that nuclei return to their starting position. This is
consistent with previous findings that the elasticity of *Drosophila* embryonic
cortex in cellularization stage depends on the actin cytoskeleton [23,24]. In
addition, the structure of actin cortex is undergoing remodeling, which may
contribute to the returning movements by actively changing the cortical
material properties in time and space.

Collective behaviors driven by the integration of forces originated from
cytoskeletal networks are indispensable in biological systems. Some basic
ingredients of the system — the mechanical properties of the cytoskeleton
and the function of motor proteins have been well studied *in vitro*. The
morphology of early *Drosophila* embryos has also been extensively imaged.
However, the assembly of the puzzle to achieve a quantitative understanding
of the molecular mechanics behind the dynamical self-organization of the
rapidly developing embryo, has only begun to be explored. Our study of the
dynamical properties of the syncytial embryo is a first step towards our long-
term goal to understand how cells mechanically interact with each other and
collectively function as active matter forming a tissue.

Methods and Materials
***Drosophila* Genetics**
Fly stocks were obtained from the Bloomington Drosophila Stock Center
[38,39], unless otherwise noted. Fly strains used in this study are the
followings: w; Histone2Av-GFP. *w;* mCherry-Tubulin, Histone2Av-GFP. *w*;
sqh-Utr-GFP/CyO; ubi-His2Av-RFP[30]. *Map60^{KG00506}. w*; ubi-GFP-D-Cad *dia^5*
Frt^{2L} ubi-His2Av-RFP/CyO [33]. *w ; ELMO^{367}* Frt^{2L}/CyO [21]. *w Hira^{ssm}*/FM7c,
w^a B [40]. His2Av-RFP; Kinesin5-[TEV]-GFP [27]. *mei41 zld/*FM7 [36]. *grp*
nmk. Sqh-GFP, Histone2Av-GFP. Fly stocks were kept at 25°C on a standard
cornmeal food. Germline clones of *dia* and *ELMO* were induced by crossing
with corresponding Frt chromosomes and the following heat shock at 37˚C for
one hour on two consecutive days after hatching.
460
Phalloidin staining and imaging
Wild type embryos and embryos from *dia* and *ELMO* germline clones were
fixed with 8% formaldehyde according to standard procedures. The vitelline
membrane was manually removed. Fixed embryos were incubated with
phalloidin-Alexa 488 (1:500, Thermo Fisher) for 1.5 h. After rinsing three times
and washing three times for 15 min each with PBT (PBS(Phosphate-Buffered
Saline) with 0.1% Tween 20), embryos were stained with DAPI (4′,6-
Diamidine-2′-phenylindole dihydrochloride) (0.2 µg/ml) for 10 min, rinsed three
times in PBT, washed in PBT for 10 min and mounted in Aqua-Poly/Mount

(Polysciences). The images of fixed embryos were acquired using a Zeiss
LSM780 confocal microscope.
472

Microinjection
1–2 h old embryos were collected, dechorionated with 50% bleach solution for
90 s, rinsed thoroughly with deionized water. After aligning on a coverslip, the
embryos were desiccated for 10 min, and covered with halocarbon oil
(Voltalef 10S, Lehmann & Voss). TEV protease (a gift from Dirk Görlich) and
Histone1-Alexa-488 protein (2 mg/ml, Thermo Fisher) were injected to the
desired embryos using Microinjector FemtoJet® (Eppendorf) on an inverted
microscope. Short spindle was induced by TEV injection in to the embryos
expressing Histone2Av-RFP and Kinesin5-[TEV]-GFP. To get the right
concentration of TEV protease for injection, we injected TEV with a serial
dilution that covers a range of concentration from 10 µM to 0.1µM. We found
1µM was robust to achieve a partial depletion allowing mitosis but with shorter
spindles during the cleavage cycle.
486

Live imaging for nuclear dynamics
Nuclear dynamics was recorded by movies of embryos with the fluorescently
labeled nuclei, by expression of Histone2Av-GFP or injection of Histone1-
Alexa-488 protein. Embryos were attached on a coverslip coated with embryo
glue and covered with halocarbon oil. Time-lapse images were recorded on a
spinning disc microscope (Zeiss, 25x/NA0.7 multi immersion) with an emCCD
camera (Photometrics, Evolve 512). To ensure reliable tracking of the nuclei,
the frame rate was 0.5–0.2 Hz with 4 axial sections, covering 8 µm. Images
were merged maximal intensity projections (Fiji/ImageJ[41]).
496

Images process and quantification
498 Imaging segmentation and analysis were performed with custom-written
499 Python algorithms. The software code is available on request. Briefly, the
500 nuclear positions were detected as blob-like features of size σ_i at position
501 (x_i, y_i) by finding the maxima $(x_i, y_i \, \sigma_i)$ of a rescaled Laplacian of Gaussian
502 (LoG) function

$$L(x, y, \sigma) = \sigma^2 \big(\Delta(g_\sigma * f)\big)(x, y, \sigma),$$

504 where $f_t(x, y)$ is the nuclei gray-scale value at time t, $g_\sigma(x, y)$ is Gaussian
505 kernel of width σ, and "$g * f$" stands for the convolution of function g and f.
506 When multiple blobs were detected in a single nucleus, we deleted a
507 neighboring blob b_2 of b_1 with a heuristic test function T

$$T(b_1, b_2) = \frac{1}{2\big(f_t(b_1) + f_t(b_2)\big)} - \int_r f_t(x) e^{-\left(x - \frac{1}{2}\right)^2} dx,$$

where $\gamma : [0, 1] \to \mathbb{R}^2$ is the straight line from b_1 of b_2.

We tracked the nuclei across frames based on a proximity criterion. The

distance between nucleus k in frame i and l in frame $i+1$ was defined as

$$d_{k,l} = \parallel x_{k,i} - x_{l,i+1} \parallel_2.$$

We determined the interval of mitosis time using the k-means-clustering

algorithm on the observed nucleus positions at time t. If a new blob was

detected, we considered this nucleus and its nearest neighbor were daughter

nuclei from a recent mitosis, and set their internal nucleus clock to 0.

Calculations of nuclear displacement, speed, nuclear density, spindle length

and orientation were done for each nucleus in its own eigentime after mitosis.

Laser ablation

Stage embryos expressing Histone2Av-GFP and Cherry-Tubulin were used.

Cross-section images were recorded in the Cherry channel with a frame rate

of 1/s on a spinning disc microscope (100x/oil, NA1.4) with a CCD camera.

Spindle apparatus was ablated at spindle midzone by a line of 355 nm YAG

laser (DPSL-355/14, Rapp Opto Electonic) with the 15% of laser power, and

around 400 ms exposure time during the recording mode (100x oil, NA 1.4)

(Fig. S5).

Particle Imaging Velocimetry (PIV) analyses

Particle Imaging Velocimetry (PIV) of Histone2Av-GFP images (in Fig. S3, S7)

analysis was performed using square interrogation windows of side 16 pixels

with an overlap of 10 s "PIVlab" in MATLAB.

Quantification of F-actin over cell cycle

Embryos expressing Histone2Av-RFP; Utrophin-GFP were imaged with a

Zeiss LSM780 confocal microscope (25x/NA0.7 multi immersion). The frame

rate was 0.1 Hz, and 10 μm was covered in z direction. Utrophin-GFP stacks

were merged by average intensity projection (Fiji/ImageJ). F-actin was

quantified manually with Fiji/ImageJ.

"Limited area" simulation of synchronous mitosis

In the simulation, the nuclei are randomly placed in a 50um*50um square via

Poisson-disc sampling, which produces random tightly-packed locations with

pair-wise distances not smaller than a specified value $d_{disc} > 4um$. We assume

the nuclei divide simultaneously and form mitotic spindles with isotropic

orientations. As the spindles extend with a constant speed, we check at each

time step if any two of the spindles touch each other by scanning a restricted

area of 4um*4um in the vicinity of each spindle (see supplementary video). If

a spindle touches at least one other, we assume it stops extending and

reaches its maximal length due to limited space. When all spindles reach their

respective maximal lengths, the simulation is ended and we compute the
average maximal length l_{max} over all spindles in the simulation. For each fixed
d_{disc}, we run the simulation 50 times, producing 50 l_{max} values for various
nuclear density around $1/(4d_{disc}^2)$. The mean of the 50 l_{max} values and the
mean of the 50 nuclear densities provide the coordinates of one point in
Fig.5E. The x- and y- error bars indicate the respective standard deviations.
Varying the minimum distance d_{disc} between nuclei, we obtain the mean l_{max}
values for a range of nuclear densities. The data from simulation are fitted to a
power-law function (solid curve in Fig. 5E) with the method of least squares.
561
Computational modeling of the nuclear movement
We extend the model by Kaiser, et al [15], which has previously been
successful in modeling static nuclei ordering during interphase, to now
account for nuclei dynamics during mitosis also now incorporating the
spherical topology of the embryo: Nuclei, positioned at $\vec{r}_i$, move due to
active forces, $\vec{F}_{act,ij}$, exerted by motor-activated pushing apart of overlapping
microtubule asters, and due to passive repulsive, $\vec{F}_{pass,ij}$, arising from the
visco-elastic matrix embedding the nuclei, built mainly from cytoskeletal actin.
The overdamped equation of motion is given by $\dot{\vec{r}}_i = \frac{1}{\eta}\sum_{j\neq i} \left(\vec{F}_{act,ij} + \vec{F}_{pass,ij} \right)$,
where $\eta = 6\pi a\mu \approx 56 \times 10^{-6}$N m^{-1}s^{-1} denotes the effective drag coefficient
for the approximately circular nuclei [42], where $a \approx 3$ μm is the nuclear
radius and $\mu \approx 1$ Pa is the viscosity of the matrix [22,23]. Both forces decay in
space following $1 / r^4$. For the active force, this is justified because the
maximal force a single microtubule can exert scales like $1 / r^2$ and the
density of microtubuli decays with $1 / r^2$ in two dimensions. For a detailed
justification on the passive force, see   Kaiser, et al [15].
The time dependence of the forces results from time dependent individual force
amplitudes contributed by each nucleus. In detail, forces are given by $\vec{F}_{pass,ij} =$
$-A_{pass}(t_i)A_{pass}(t_j)\frac{\vec{e}_{r_{i,j}}}{r_{i,j}^4}$ and $\vec{F}_{act,ij} = -A_{act}(t_i)A_{act}(t_j)\frac{\vec{e}_{r_{i,j}}}{r_{i,j}^4}$, where $\vec{e}_{r_{i,j}}$ and
$r_{i,j}$ denote the unit vector and the distance between nuclei i,j, respectively.
Nuclei divide when their age reaches $t_i = t_{div}$, and ages are initialized as $t_i =$
$t_{div} - \frac{2}{\pi}\theta_i t_{wave} - t_{offset}$, where θ_i is the polar coordinate of nucleus i. This
initialization leads to the division wave. For numerical stability, upon division,
the two daughter nuclei are placed a short distance r_{div} apart with random
orientation, their center of mass coinciding with the position of their mother.
587

To capture the dynamics of cytoskeletal elements during mitosis we subdivided
the time course of events into the series: $t_{\text{spindle ass}} < t_{\text{spindle const}} < t_{\text{div}} <$
$t_{\text{sp diss}} < t_{\text{MT ass}} < t_{\text{actin ass}} < t_{\text{MT inter}} < t_{\text{actin inter}}$. Passive forces change only
between $t_{\text{spindle ass}} < t_{\text{spindle const}}$ when actin caps shrink while spindles
assemble generating balancing forces and during regrowth of actin caps
between $t_{\text{actin ass}} < t_{\text{actin inter}}$ before entering interphase. Active forces in
contrast are much more dynamic: Between $t_{\text{spindle ass}} < t_{\text{spindle const}}$ spindles
assemble and the active force grows to exert a maximal force of h_{spindle}. After
division $t_{\text{div}} < t_{\text{spindle diss}}$ spindles move daughter nuclei apart: The active force
between the division partners is now increased by an additional time-dependent

factor A_{div} that reaches values of up to 3, $\vec{F}_{act,i,i+1} =$

$-A_{\text{div}}(t)A_{act}(t)A_{act}(t)\dfrac{\vec{e}_{r_{i,i+1}}}{r^4_{i,i+1}}$, where t is the common time of both daughter

nuclei. After $t_{\text{spindle diss}}$, forces between paired daughter nuclei reduce linearly
to regular levels. Meanwhile, for all other nuclei, after division the active force
is halved since each nucleus is only associated with one centrosome, instead
of two as before. This is balanced by the effective increase of passive forces
as nuclei are closer packed due to division while the passive force amplitude
stays constant. Between $t_{\text{spindle diss}} < t_{\text{MT ass}}$ spindles disassemble before the
microtubule asters regrow during $t_{\text{MT ass}} < t_{\text{MT inter}}$. All dynamics are
interpolated linearly. In detail:

$A_{\text{pass}}(t) =$

609
$$
\begin{cases}
h_{\text{inter}} & t \leq t_{\text{spindle ass}} \\
h_{\text{inter}} + (h_{\text{mitosis}} - h_{\text{inter}})\frac{t - t_{\text{spindle ass}}}{t_{\text{spindle const}} - t_{\text{spindle ass}}} & t_{\text{spindle ass}} \leq t \leq t_{\text{spindle const}} \\
h_{\text{mitosis}} & t_{\text{spindle const}} \leq t \leq t_{\text{actin ass}} \\
h_{\text{mitosis}} + (h_{\text{inter}} - h_{\text{mitosis}})\frac{t - t_{\text{actin ass}}}{t_{\text{actin inter}} - t_{\text{actin ass}}} & t_{\text{actin ass}} \leq t \leq t_{\text{actin inter}} \\
h_{\text{inter}} & t_{\text{actin inter}} \leq t
\end{cases},
$$

610 $A_{\text{act}}(t) =$

611
$$
\begin{cases}
h_{\text{MT}} & t \leq t_{\text{spindle ass}} \\
h_{\text{MT}} + (h_{\text{sp}} - h_{\text{MT}})\frac{t - t_{\text{spindle ass}}}{t_{\text{spindle const}} - t_{\text{spindle ass}}} & t_{\text{spindle ass}} \leq t \leq t_{\text{spindle const}} \\
h_{\text{spindle}} & t_{\text{spindle const}} \leq t \leq t_{\text{div}} \\
\frac{1}{2}h_{\text{spindle}} & t_{\text{div}} \leq t \leq t_{\text{spindle diss}} \\
\frac{1}{2}h_{\text{spindle}} + \frac{1}{2}(h_{\text{spindle diss}} - h_{\text{spindle}})\frac{t - t_{\text{spindle diss}}}{t_{\text{MT ass}} - t_{\text{spindle diss}}} & t_{\text{spindle diss}} \leq t \leq t_{\text{MT ass}} \\
\frac{1}{2}h_{\text{spindle diss}} + \frac{1}{2}(h_{\text{MT}} - h_{\text{spindle diss}})\frac{t - t_{\text{MT ass}}}{t_{\text{MT inter}} - t_{\text{MT ass}}} & t_{\text{MT ass}} \leq t \leq t_{\text{MT inter}} \\
\frac{1}{2}h_{\text{MT}} & t_{\text{MT inter}} \leq t
\end{cases},
$$

$$612 \quad A_{\text{div}}(t) = \begin{cases} h_{\text{div}} \dfrac{t - t_{\text{div}}}{t_{\text{spindle diss}}/2 - t_{\text{div}}} & t_{\text{div}} \le t \le t_{\text{spindle diss}}/2 \\[2ex] h_{\text{div}} & t_{\text{spindle diss}}/2 \le t \le t_{\text{spindle diss}} \\[2ex] h_{\text{div}} - (h_{\text{div}} - 1) \dfrac{t - t_{\text{spindle diss}}}{t_{\text{MT ass}} - t_{\text{spindle diss}}} & t_{\text{spindle diss}} \le t \le t_{\text{MT ass}} \\[2ex] 1 & t_{\text{MT ass}} \le t \end{cases}$$

Note that changes in both passive and active forces only get out of balance - initiating nuclei motion - during spindle disassembly with significant time delay relative to the time point of nuclei division. At that point in time nuclei that have not divided yet exert a stronger repulsive force than the already divided nuclei, since their actin caps and microtubule asters are not fully reformed. Therefore, nuclei move toward the region of higher nuclei density, only returning back when actin caps and microtubule asters are forming again.

Choices for the model parameters are found in the table below. Their magnitudes are chosen to match the length of the cell cycle, 800s, and the maximal force exerted by a single microtubule, which is around 3pN according to [43]. In total, the forces on a single nucleus range between 10-100pN, the same order of magnitude as the force applied to a single magnetic microparticle by [24] to move it through the cellularizing tissue in an early drosophila embryo. Note that computing the force involves multiplying the two amplitudes and dividing by the distance to the power of four.

$t_{\text{spindle ass}}$	$t_{\text{spindle const}}$	t_{div}	$t_{\text{sp diss}}$	$t_{\text{MT ass}}$	$t_{\text{actin ass}}$	$t_{\text{MT inter}}$	$t_{\text{actin inter}}$
-200s	-80s	0s	70s	200s	240s	400s	600s

h_{inter}	h_{mitosis}	h_{MT}	h_{spindle}	$h_{\text{spindle diss}}$	h_{div}
4e-17 $\text{N}^{1/2}\,\text{m}^2$	1e-17 $\text{N}^{1/2}\,\text{m}^2$	4e-17 $\text{N}^{1/2}\,\text{m}^2$	8e-16 $\text{N}^{1/2}\,\text{m}^2$	0 $\text{N}^{1/2}\,\text{m}^2$	3

Spring constant fitting
The data sets consisted of 1–5 nuclear displacement curves for 2–4 embryos of each type (*dia*, *ELMO*, *Kinesin5* and *Map60* mutants, wild type). The nuclear displacement curves (in the first phase) are similar to the oscillation of a not-actively driven and non-damped harmonic oscillator. Therefore, the individual nuclear displacement curves were fitted to a sine curve of the form $y(t) = A \sin(\omega t + \varphi)$, where A is the amplitude, ω the angular frequency and φ the phase shift, using a self-written script in Python. The biological rational behind this approach is that the nuclei behave like they were linked to an elastic spring,

which could be e.g. linkages to the cytoskeleton. At t=0 the spring is stretched and the nuclei start to move until the spring is compressed and the nuclei move back.

The fit region was determined as follows. For all curves, the lower bound on the fitting range was set equal to the point in time where the nuclear displacement first exceeds 5 μm, as some curves show a small, reversible displacement in the beginning. The upper bound was chosen independently for each of the data sets since the elastic part of the curve depends on the stiffness and dampening of the spring and hence differs across data sets. For the *dia* and *Map60* mutants as well as the wild type, the upper bound was set equal to 100 s after the turning point, while it was set to 180 s after for the *ELMO* mutant and 240 s for the Kinesin5 mutants. The results of the fit parameter ω were averaged for each embryo to give a set of angular frequencies ω_i for each type, where i runs over the number of embryos. The spring constant was derived from the average ω_i *via* the relation $k = m\ (\omega_i)^2,$ in which m denotes the mass of the nucleus. It was assumed that the nuclei are spherical with a diameter of 4.9 μm and a density equal to that of water at room temperature. Error bars, which correspond to one standard deviation, were calculated in the frequency domain and then converted to the force domain by the analogue of the relation above.

Acknowledgements

We are grateful to D. Görlich, T. Lecuit, B. Loppin and the members of the Grosshans laboratory for stains, materials and discussions. We acknowledge service support from the Bloomington Drosophila Stock Center (supported by NIH P40OD018537). X.Z. was supported by Center for Advancing Electronics Dresden (cfaed) and by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy – EXC-2068 – 390729961– Cluster of Excellence Physics of Life of TU Dresden. This work was in part supported by the Göttingen Centre for Molecular Biology (funds for equipment repair) and the Deutsche Forschungsgemeinschaft (DFG, SFB-937/A10, SFB-937/A19 and equipment grant INST1525/16-1 FUGG).

Author contributions

ZL and DK conducted the experiments. ZL, JR, HP, SK analyzed the data. SM, XZ, KA conducted the simulations. JG and ZL conceived and JG, TA, SG, KA supervised the study. JG, ZL and XZ wrote the manuscript.

References:

[1]   Couzin ID. Synchronization: The Key to Effective Communication in

687    Animal Collectives. Trends in Cognitive Sciences 2018;22:844–6.
688    doi:10.1016/j.tics.2018.08.001.

689 [2] Partridge BL. Rigid definitions of schooling behaviour are inadequate.
690    Animal Behaviour 1982;30:298–9.

691 [3] Moussaid M, Garnier S, Theraulaz G, Helbing D. Collective
692    Information Processing and Pattern Formation in Swarms, Flocks, and
693    Crowds. Topics in Cognitive Science 2009;1:469–97.
694    doi:10.1111/j.1756-8765.2009.01028.x.

695 [4] Chialvo DR. Emergent complex neural dynamics. Nature Physics
696    2010;6:744–50. doi:10.1038/nphys1803.

697 [5] Levina A, Herrmann JM, Geisel T. Dynamical synapses causing self-
698    organized criticality in neural networks. Nature Physics 2007;3:857–
699    60. doi:10.1038/nphys758.

700 [6] Plenz D, Thiagarajan TC. The organizing principles of neuronal
701    avalanches: cell assemblies in the cortex? Trends in Neurosciences
702    2007;30:101–10. doi:10.1016/j.tins.2007.01.005.

703 [7] McMahon A, Supatto W, Fraser SE, Stathopoulos A. Dynamic
704    Analyses of Drosophila Gastrulation Provide Insights into Collective
705    Cell Migration. Science 2008;322:1546–50.
706    doi:10.1126/science.1167094.

707 [8] Farrell JA, Wang Y, Riesenfeld SJ, Shekhar K, Regev A, Schier AF.
708    Single-cell reconstruction of developmental trajectories during
709    zebrafish embryogenesis. Science 2018;360:eaar3131.
710    doi:10.1126/science.aar3131.

711 [9] Sumino Y, Nagai KH, Shitaka Y, Tanaka D, Yoshikawa K, Chaté H, et
712    al. Large-scale vortex lattice emerging from collectively moving
713    microtubules. Nature 2012;483:448–52. doi:10.1038/nature10874.

714 [10] Schaller V, Weber C, Semmrich C, Frey E, Bausch AR. Polar patterns
715    of driven filaments. Nature 2010;467:73–7. doi:10.1038/nature09312.

716 [11] Butt T, Mufti T, Humayun A, Rosenthal PB, Khan S, Khan S, et al.
717    Myosin motors drive long range alignment of actin filaments. The
718    Journal of Biological Chemistry 2010;285:4964–74.
719    doi:10.1074/jbc.M109.044792.

720 [12] Deneke VE, Melbinger A, Vergassola M, Di Talia S. Waves of Cdk1
721    Activity in S Phase Synchronize the Cell Cycle in Drosophila Embryos.
722    Developmental Cell 2016;38:399–412.
723    doi:10.1016/j.devcel.2016.07.023.

724 [13] Idema T, Dubuis JO, Kang L, Manning ML, Nelson PC, Lubensky TC,
725    et al. The Syncytial Drosophila Embryo as a Mechanically Excitable
726    Medium. PLoS ONE 2013;8:e77216.
727    doi:10.1371/journal.pone.0077216.

728 [14] Kanesaki T, Edwards CM, Schwarz US, Grosshans J. Dynamic

ordering of nuclei in syncytial embryos: a quantitative analysis of the
role of cytoskeletal networks. Integrative Biology (Camb)
2011;3:1112–9. doi:10.1039/c1ib00059d.
[15] Kaiser F, Lv Z, Marques Rodrigues D, Rosenbaum J, Aspelmeier T,
Grosshans J, et al. Mechanical Model of Nuclei Ordering in Drosophila
Embryos Reveals Dilution of Stochastic Forces. Biophysical Journal
2018;114:1730–40. doi:10.1016/j.bpj.2018.02.018.
[16] Quinlan ME. Cytoplasmic Streaming in the Drosophila Oocyte. Annual
Review of Cell and Developmental 2016;32:173–95.
doi:10.1146/annurev-cellbio-111315-125416.
[17] Woodhouse FG, Goldstein RE. Cytoplasmic streaming in plant cells
emerges naturally by microfilament self-organization. Proceedings of
the National Academy of Sciences of the United States of America
2013;110:14132–7. doi:10.1073/pnas.1302736110.
[18] Shimmen T, Yokota E. Cytoplasmic streaming in plants. Current
Opinion in Cell Biology 2004;16:68–72.
doi:10.1016/j.ceb.2003.11.009.
[19] Deneke VE, Puliafito A, Krueger D, Narla AV, De Simone A, Primo L,
et al. Self-Organized Nuclear Positioning Synchronizes the Cell Cycle
in Drosophila Embryos. Cell 2019. doi:10.1016/j.cell.2019.03.007.
[20] He B, Doubrovinski K, Polyakov O, Wieschaus E. Apical constriction
drives tissue-scale hydrodynamic flow to mediate cell elongation.
Nature 2014;508:392–6. doi:10.1038/nature13070.
[21] Winkler F, Gummalla M, Künneke L, Lv Z, Zippelius A, Aspelmeier T,
et al. Fluctuation Analysis of Centrosomes Reveals a Cortical Function
of Kinesin-1. Biophysical Journal 2015;109:856–68.
doi:10.1016/j.bpj.2015.07.044.
[22] Wessel AD, Gumalla M, Grosshans J, Schmidt CF. The mechanical
properties of early Drosophila embryos measured by high-speed video
microrheology. Biophysical Journal 2015;108:1899–907.
doi:10.1016/j.bpj.2015.02.032.
[23] Doubrovinski K, Swan M, Polyakov O, Wieschaus EF. Measurement
of cortical elasticity in Drosophila melanogaster embryos using
ferrofluids. Proceedings of the National Academy of Sciences of the
United States of America 2017;114:1051–6.
doi:10.1073/pnas.1616659114.
[24] D'Angelo A, Dierkes K, Carolis C, Salbreux G, Solon J. In Vivo Force
Application Reveals a Fast Tissue Softening and External Friction
Increase during Early Embryogenesis. Current Biology 2019;29:1564–
6. doi:10.1016/j.cub.2019.04.010.
[25] Vergassola M, Deneke VE, Di Talia S. Mitotic waves in the early
embryogenesis of Drosophila: Bistability traded for speed.

771    Proceedings of the National Academy of Sciences of the United States
772    of America 2018;115:E2165–74. doi:10.1073/pnas.1714873115.

773 [26] Scholey JM. Kinesin-5 in Drosophila embryo mitosis: Sliding filament

774    or spindle matrix mechanism? Cell Motility and the Cytoskeleton

776 [27] 2009;66:500–8. doi:10.1002/cm.20349.osshans J. A "molecular

777    guillotine" reveals an interphase function of Kinesin-5. Journal of Cell

778    Science 2018:jcs.210583. doi:10.1242/jcs.210583.

779 [28] Sullivan W, Theurkauf WE. The cytoskeleton and morphogenesis of

780    the early Drosophila embryo. Current Opinion in Cell Biology

781    1995;7:18–22. doi:10.1016/0955-0674(95)80040-9.

782 [29] Karr TL, Alberts BM. Organization of the cytoskeleton in early

783    Drosophila embryos. The Journal of Cell Biology 1986;102:1494–509.

784    doi:10.1083/jcb.102.4.1494.

785 [30] Rauzi M, Lenne P-F, Lecuit T. Planar polarized actomyosin contractile

786    flows control epithelial junction remodelling. Nature 2010;468:1110–4.

787    doi:10.1038/nature09566.

788 [31] Schmidt A, Lv Z, Grosshans J. ELMO and Sponge specify subapical

789    restriction of Canoe and formation of the subapical domain in

790    earlyDrosophilaembryos. Development 2018;145:dev157909.

791    doi:10.1242/dev.157909.

792 [32] Geisbrecht ER, Haralalka S, Swanson SK, Florens L, Washburn MP,

793    Abmayr SM. Drosophila ELMO/CED-12 interacts with Myoblast city to

794    direct myoblast fusion and ommatidial organization. Developmental

795    Biology 2008;314:137–49. doi:10.1016/j.ydbio.2007.11.022.

796 [33] Afshar K, Stuart B, Wasserman SA. Functional analysis of the

797    Drosophila diaphanous FH protein in early embryonic development.

798    Development 2000;127:1887–97. doi:10.1083/jcb.137.1.169.

799 [34] Grosshans J, Wenzl C, Herz H-M, Bartoszewski S, Schnorrer F, Vogt

800    N, et al. RhoGEF2 and the formin Dia control the formation of the

801    furrow canal by directed actin assembly during Drosophila

802    cellularisation. Development 2005;132:1009–20.

803    doi:10.1242/dev.01669.

804 [35] Wenzl C, Yan S, Laupsien P, Grosshans J. Localization of RhoGEF2

805    during Drosophila cellularization is developmentally controlled by

806    Slam. Mechanisms of Developmen 2010;127:371–84.

807    doi:10.1016/j.mod.2010.01.001.

808 [36] Blythe SA, Wieschaus EF. Zygotic genome activation triggers the DNA

809    replication checkpoint at the midblastula transition. Cell

810    2015;160:1169–81. doi:10.1016/j.cell.2015.01.050.

811 [37] Benton MA, Akam M, Pavlopoulos A. Cell and tissue dynamics during

812    Tribolium embryogenesis revealed by versatile fluorescence labeling

approaches. Development 2013;140:3210–20.
doi:10.1242/dev.096271.

[38] Gramates LS, Marygold SJ, Santos GD, Urbano J-M, Antonazzo G,
Matthews BB, et al. FlyBase at 25: looking to the future. Nucleic Acids
Research 2017;45:D663–71. doi:10.1093/nar/gkw1016.

[39] Cook KR, Parks AL, Jacobus LM, Kaufman TC, Matthews K. New
research resources at the Bloomington Drosophila Stock Center. Fly
2010;4:88–91. doi:10.4161/fly.4.1.11230.

[40] Loppin B, Docquier M, Bonneton F, Couble P. The Maternal Effect
Mutation sésame Affects the Formation of the Male Pronucleus in
Drosophila melanogaster. Developmental Biology 2000;222:392–404.
doi:10.1006/dbio.2000.9718.

[41] Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M,
Pietzsch T, et al. Fiji: an open-source platform for biological-image
analysis. Nature Methods 2012;9:676–82. doi:10.1038/nmeth.2019.

[42] Koke C, Kanesaki T, Grosshans J, Schwarz US, Dunlop CM. A
computational model of nuclear self-organisation in syncytial embryos.
Journal of Theoretical Biology 2014;359:92–100.
doi:10.1016/j.jtbi.2014.06.001.

[43] Vleugel M, Kok M, Dogterom M. Understanding force-generating
microtubule systems through in vitro reconstitution. Cell Adhesion and
Migration 2016;10:475–94. doi:10.1080/19336918.2016.1241923.

Figure 1. Quantitative assay for nuclear movement.

(A)**,** Live image of an embryo during mitosis. The left half is in metaphase, the right half in anaphase. The dotted line in red indicates metaphase-anaphase transition sweeping over the embryo from right to left as a wave front.

(B), Snapshots show the nuclear motion. The red line indicates the metaphase-anaphase transition. Blue line is the forward trajectories of nuclei. The right panel is the corresponding angle distribution of nuclear motion.

(C), The projection of the nuclear motion. Red arrow indicates the nuclear division.

(D, E), Time course of nuclear displacement with the position of the mother nucleus and nuclear speeds during metaphase-anaphase transition as a reference (t=0) (n=260 nuclei in one embryo, representative for all embryos which have been checked). The numbers indicate ①chromosome segregation, ②forth movement away and ③ back movement toward the mitotic wave front. The right panel is a snapshot of nuclear movement labelled with three numbers. The image is the end of the nuclear movement.

(F), The maximal displacement plotted against the corresponding speed of the mitotic wave (n=50 embryos).

Figure 2. Emergence of collective nuclear movement.

(A), Schematic drawing of an embryo with definition of angles.

(B), Image from live imaging after mitosis. Orientations of the corresponding spindle at t=0 s and the directions of nuclear movement at t = 60 s are indicated by a magenta bar and green arrow, respectively.

(C), Distribution of observed angles for spindle orientation at t=0 s and nuclear movement at t=60 s. (n=20 embryos including 6230 nuclei).

(D), Image series with previous trajectories showing cases of perpendicular and parallel mitosis.

(E), Time course of nuclear displacement (blue) and distance between corresponding daughter nuclei (orange). The time lag (τ) between maxima is indicated (n=260 nuclei in one embryo, representative for all embryos we have checked).

(F), Distribution of the time lag (τ) (n=21 embryos).

(G), The schematic drawing of the transition from individual movement to collective behavior. Data are mean±s.e.m. Scale bar: 10 μm

Figure 3. Numerical simulation of nuclear movement.

(A), Scheme of division, active and passive forces in syncytial embryo.

(B), Snapshots from the simulation. Nuclei were projections. Color code indicates speed of nuclear movement.

(C), Time course of nuclear displacement (n=300 nuclei).

(D), Time course of nuclear displacement (blue) and distance between
corresponding daughter nuclei (orange) (n=300 nuclei). The time lag (τ)
between maxima is indicated.
(E), Distribution of the time lag (τ) (n=21 embryos).
(F), The negative correlation between maximal displacement and the
corresponding speed of the mitotic wave.
(G), Time course of nuclear displacement in wave (wild type) embryo and
"salt-and-pepper pattern" embryo (n=300 nuclei in both cases, respectively).
(H), The positive correlation between maximal displacement and division
distance. Data are mean±s.e.m.
887

Figure 4. Forces mediating Yo-Yo movement of nuclei *in vivo*.
(A). Spindle length/distance between respective daughter nuclei in embryos
partially depleted of Kinesin-5 and *Map60* mutants (n=30 spindle in 3 embryos
for each genotype).
(B), Images from live image. The trajectories of two nuclei over 300 s were
plotted into the images. Images were the end of the movement.
(C), Time course of nuclear displacement.
(D), Maximal displacement was plotted against corresponding division
distance. (n=290 nuclei in wild type, 256 nuclei in *kin5*, 292 nuclei in *Map60* in
Fig. C and D).
(E), Scheme of cortical actin dynamics during nuclear division cycles.
(F), ELMO and Dia are involved in F-actin cortex formation.
(G), Images from movies of *dia* and *ELMO* mutants. Nuclear trajectories over
10 min are plotted into the images. Images were the end of the nuclear
movement.
(H, J), Time course of displacement in *ELMO* and *dia* mutants.
(J), The final distance between the current position and original position of the
nuclei after yo-yo movement, as indicated as "d" in H and I (n=260 nuclei in
wild type, 45 nuclei in *ELMO* mutant and 90 in *dia* mutant embryo in Fig. H, I,
J). The Data are mean±s.e.m. Scale bar: 10 μm.
908

Figure 5. Long mitotic spindles require pseudo-synchronous nuclear
cycles.
(A), Live image series of the process of nuclear dividing.
(B), Quantification of the distance between splitting nuclear pair in cleavage
cycles.
(C) The protected area between daughter nuclei containing the spindle is
indicated by an overlaid colored area.
(D), Illustration of the simulation. Green dots indicate nuclei. Blue arrows
indicate spindles. Red dots indicate the touching of the neighbor spindles,
leading to the spindle growth stops.

(E), Spindle lengths/distances between daughter nuclei are plotted against the
corresponding nuclear density. Data from simulation (blue) and
measurements in embryos in NC11, NC12, NC13 in diploids and NC14 in
haploids (orange) (n=15 spindles in each embryo and 3 embryos for each
type).

Figure S1. Characterization of nuclear dynamics in syncytial embryo.
(A), Embryo to embryo variation of the speed of the wave front (n=50
embryos).
(B), Image series with Voronoi maps with the mitotic wave front in images at
35 s and 55 s. Color code indicates nuclear density.
(C), Time course of nuclear density with metaphase-anaphase transition at
t=0. Numbers indicate the three stages of nuclear movement. (n=260 nuclei in
one embryo).
(D). Schematic drawing shows nuclei move like an elastic sheet.
(E), A simple square function (red) was fitted to the forth and back movement
around the maximal displacement. According to Hook's law an apparent
spring constant ($1.5\pm0.02\times10^{-18}$ Nm^{-1}) was calculated (n=15 nuclei from 3
embryos). Data are mean±s.e.m. Scale bar: 10 µm.

Figure S2. Nuclear displacement and flow speeds are less pronounced
in earlier cycles.
(A), The time course of nuclear displacement in NC11, 12 and 13 in one
embryo (n=58, 107 and 206 nuclei in NC11, 12 and 13, respectively).
(B), Maximal displacement distribution in NC11, 12 and 13 from (A).
(C), Time course of nuclear flow speed from the same embryo as shown in a,
with colour-coding for indicated nuclear cycles.
(D), The percentage of embryos showing the nuclear movements ①, ② and
③ in different cycles (n=6, 18, 26 embryos in NC11, 12 and 13, respectively).
Data are mean±s.e.m.

Figure S3. The nuclei move as a sheet.
(A), Groups of cells have been marked in color at metaphase. Their daughter
nuclei were labelled with the same color in the following images.
(B) Particle Imaging Velocimetry analysis show that the nuclei move
collectively like a sheet.

Figure S4. The parameters for Numerical simulation used in Fig. 3.
(A), Time course of the division, active force and passive force used in the
simulation.
(B), Increasing in spindle strength ($h_{spindle}$) leads to increasing of maximal
displacement and division distance. Data are mean±s.e.m.

Figure S5. Laser cutting on spindle in metaphase. The embryo expressing mCherry-Tubulin was used for monitor the spindle structure. The spindle recovers in second-scale after laser ablation.

Figure S6. Actin cortex represses nuclear movement.
(A), Live-images showing F-actin organization during nuclear cycle.
(B), Quantification of F-actin. (n=10 regions in 3 independent recordings).
(C, D), Injection of ROCK inhibitor Y-27632 leads to the reduction of nuclear motion as well as spindle elongation.
(E, F), Time course of single nuclear displacement and the local myosin/F-actin intensity.
(E), Fixed *dia* and *ELMO* mutants stained for F-actin (red) and DNA (blue).
(F), Maximal displacement and final distance of nuclear dynamics in *dia* and *ELMO* mutants.
(G), Apparent spring constant.
(H), Images from movies of wild type and *ELMO* mutants in NC12. Nuclear trajectories are plotted into the first images. (I), The time course of nuclear displacement in NC12. (n=130 nuclei in *ELMO* and 290 nuclei in wild type. Data are mean±s.e.m. Scale bar: 10 μm.

Figure S7. Nuclei move as a sheet in *dia* and *ELMO* mutant embryo.
(A), Snapshots from movies at indicated time. Groups of cells have been marked in color at metaphase. Their daughter nuclei were labelled with the same color in the following images.
(B). Particle Imaging Velocimetry analysis show that the nuclei move collectively.

Figure S8. Nuclear densities in different cycles.
(A), Images of Drosophila syncytial embryo expressing Histone2Av-GFP in different cycles.
(B), Quantification of nuclear density in the indicated cycles. n=10 embryos. Data are mean±s.e.m.

Figure S9. Daughter nuclei cannot separate in *nmk grp* and *mei41 zld* mutant embryos.

Figure S10. Mitotic wave speed in the indicated embryos.

Figure S11. The asymmetric force field leads to the nuclear directional movement.

(A), The repulsive force between daughter nuclei increases in anaphase
pushing the daughter nuclei apart, followed by a drop in telophase due to
spindle disassembly.
(B), The summing up of all nuclei in an embryo at a given mitotic time results
in an asymmetric force field, which likely determines the directionality of the
nuclear flow in telophase.
1008
1009   **Movie list:**
1010   Movie 1. Mitotic wave sweeps over the embryo.
1011
1012   Movie 2. Nuclei undergo stereotypical movement after metaphase-anaphase
1013   transition. Scale bar: 10 µm.
1014
1015   Movie 3. The time course of the nuclear Voronoi map over nuclear division.
1016   Scale bar: 10 µm.
1017
1018   Movie 4. Nuclei move as a laminar flow shown in PIV analysis. Scale bar: 10
1019   µm.
1020
1021   Movie 5. Computational simulation.
1022
1023   Movie 6. The nuclear displacement in wild type, dia and ELMO mutant
1024   embryos. Scale bar: 20 µm.
1025
1026

Figure 1

Figure 2

A mitotic wave 90° 0° 90°

B t=0 s t=60 s

C Probability density, Angle — spidle orientation, nuclear flow orientation

D perpendicular 0 s 6 s 50 s 168 s; parallel; 0.5 μm/s

E Experiment; Displacement [μm]; d (daughter nuclei) [μm]; Time [s]; t

F t [s]

G isotropic individual spindles → anisotropic collective movement

Figure 3

A

$\vec{F}_{div}$
mitosis

$\vec{F}_{act}$
$\vec{F}_{pass}$
interphase

B

C

Displacement [a.u.]
Time [x10²s]

D

Displacement [a.u.]
d (daughter nuclei) [a.u.]
Time [s]
t

E

t [s]

amplitude of active force

F

Max. displacement [<r_n>]
Mitotic wave speed [1/s]

G

Displacement [a.u.]
wave
salt-and-pepper
Time [x10²s]

H

Max. displacement [<r_n>]
Division distance [<r_n>]

Figure 4

A

p < 0.0001

p < 0.0001

Spindle length [µm]

WT   Kin5   Map 60

B 300 s

WT

Kin5

Map60

C

Nuclear displacement [µm]

wild type
Kin5
Map60

Time [s]

D

Max. displacement [µm]

wild type
Kin5
Map60

Division distance [µm]

E

inter-   pro-   meta-   ana-   telo-   inter-

F

ELMO/Sponge

Arp2/3   Dia

Cortical F-actin

G

wild type   *dia*

ELMO

H

Displacement [µm]

wild type
ELMO

d

Time [s]

I

Displacement [µm]

wild type
dia

d

Time [s]

J

Final distance [µm]

p < 0.0001

p < 0.0001

wild type   *dia*   *ELMO*

Figure 5

A His-GFP

Meta.   Ana.   Telo.   Inter.

B Distance between
[μm] splitting nuclear pair

— NC11
— NC12
— NC13

Time [s]

C

D

E

Maximal daughter distance [μm]

forbidden

allowed

— Simulation
— Experiment
— Fitting

Nuclear density x 10^{-3} [μm^{-2}]

Figure S1

A

Number of embryos

Wave speed [µm/s]

B

0 s   35 s   55 s   95 s   160 s   350 s

low   high   nuclear density

C

Nuclear density [um⁻²]

① ③ ②

Time [x10²s]

D

①Doubling   ②Stretching   ③Relaxing

E

Displacement [µm]

measurement
fitting

Time [x10²s]

Spring constant:
1.50±0.02x10⁻¹⁸Nm⁻¹

Figure S2

Figure S3

A

0 min   0.75 min   2 min   6 min

B

Anaphase   0 min

Interphase   4 min

Figure S4

Figure S5

mCherry-Tubulin

Figure S6

Figure S7

Figure S8

A

interphase 12

interphase 13

interphase 14

50 μm

B

Figure S9

Figure S10

Figure S11