Effectiveness of treatments for firework fears in dogs

Stefanie Riemer*

Companion Animal Behaviour Group, Division of Animal Welfare, Vetsuisse Faculty,

University of Bern, Länggassstrasse 120, 3012 Bern, Switzerland

* Corresponding author: riemer.stefanie@gmail.com

Highlights

- An online survey on treatment options for firework fears in dogs was performed
- Feeding or playing with dogs during fireworks was associated with fear improvement
- Success was highest for ad-hoc counterconditioning, relaxation training and medication
- Success was similar for pheromones, nutraceuticals and other alternative products
- Success rates for these alternative products are consistent with a placebo effect

Keywords

Noise fear, dog *Canis familiaris*, counterconditioning, anxiolytic medication, pheromones, placebo effect

Abstract

An online questionnaire survey investigated (1) management and (2) treatment methods for firework fears in dogs employed by dog owners and their (perceived) effectiveness. A PCA on data from 1225 respondents revealed four management strategies (i.e. interventions during firework exposure): the principal components *"Environmental modification"* (e.g. providing a hiding place, keeping windows and blinds closed, and playing music), *"Feed/Play"* (providing the dog with chews, play and food during fireworks in general, as well as contingent on loud bangs), *"Alternative"* (use of calming nutraceuticals, pheromones, herbal products, homeopathic products, Bach flowers, and essential oils) and *"Interaction"* (allowing body contact, petting and talking to the dog when loud bangs occurred). To explore possible effects of these management methods on fear development, the components were correlated with a score for fear progression. Of the four components, only *"Feed/Play"* was statistically associated with an improvement in fear responses to fireworks. To evaluate the effectiveness of various treatment strategies, owners were asked to select from a range of options which interventions they had used and whether they considered them as effective. With prescription medication (N=202), improvements were noted by 69% of owners, with high success rates reported for the most frequently prescribed drugs, alprazolam (91%) and Sileo® (74%). While individual products were not evaluated, the reported success rates for the categories "pheromones" (N=316), "herbal products" (N=282), "nutraceuticals" (N=211), "essential oils" (N=183), "homeopathic remedies" (N=250) and "Bach flowers" (N=281) were all in the range of 27-35%, which is not higher than would be expected based on a placebo effect. Pressure vests were deemed as effective by 44% of respondents (N=300). Counterconditioning (providing desirable stimuli after the occurrence of noises) was the most successful training technique according to the owners (N=694), with a reported effectiveness of over 70%. Relaxation training (N=433) was reported to be almost as successful at 69%, while noise CDs (N=377) were effective in 55% of cases. Thus, counterconditioning, relaxation training and anxiolytic medication appear to be the most effective strategies in the treatment of firework fears in dogs. On this basis it is recommended that ad-hoc counterconditioning and relaxation training should complement the standard behavioral technique of desensitization/ counterconditioning with noise recordings.

Introduction

Fear responses to fireworks and other loud noises (often referred to as 'noise phobia', 'noise sensitivity', 'noise reactivity', 'noise aversion' or 'noise stress') represent a significant welfare problem for pet dogs. According to surveys, up to half the pet dog population react fearfully to fireworks (Blackwell et al., 2013; Riemer, 2019; Storengen and Lingaas, 2015; Tiira et al., 2016), and one study indicated that over 15% of fearful dogs require several days or longer to recover behaviorally from a firework event, with over 3% exhibiting behavioral changes for weeks or months (Riemer, 2019). Thus, identifying effective interventions to prevent and treat firework fears in dogs is of wide concern. Owners' awareness of the issue seems to be increasing (c.f. Riemer, 2019), and a wide range of products are marketed as alleviating firework fears in dogs, ranging from pheromones to special garments and from homeopathy to prescription medication. However, as has been pointed out previously, many products lack scientific evidence for their effectiveness, and even most published studies in the field are based on low sample sizes and often lack placebo treatments (Sherman and Mills, 2008). Subsequently I give a brief overview over approaches to treating noise fears in dogs.

1) Behavioral techniques

Management

In order to minimize adverse experiences for the animal, management measures are important when exposure to fearful stimuli cannot be avoided (Sherman and Mills, 2008). Commonly given management advice for noise fears includes providing a safe place to retreat, ideally by associating it with positive experiences prior to any firework events, darkening the room putting music or white noise on, ignoring fearful behavior, distracting the dog with games, training or food, and refraining from any punishment (Mills, 2005; Pike et al., 2015; Sherman and Mills, 2008). However, management is likely not to be sufficient to generate long-lasting improvement, and behavior modification techniques, often with

adjunctive use of pheromones or medication, are usually recommended (Sherman and Mills,

2008). Given that fears and phobias can significantly compromise welfare and become

severe, wherever possible clinicians should aim to treat fears and attempt to resolve the

problem, rather than relying on management only (Horwitz and Mills, 2012).

Systematic desensitization and counterconditioning

'Desensitization" can be defined as "gradual and controlled exposure to the stimulus so as to

extinguish the manifestations of fearful behavior" (Horwitz and Mills, 2012, p. 177). It

involves repeated exposure to the feared stimulus, but at such a low intensity that it elicits no

distress response. For instance, a noise recording can be played at a very low level. While

the dog might initially alert to it, it should learn to ignore it with repeated exposure. Gradually,

over the course of multiple sessions, the volume can be increased, as long as the dog does

not show any distress upon exposure (McMillan, 2019). Generally, this approach is most

effective in alleviating fear responses when combined with counterconditioning (McMillan,

2019).

'Counterconditioning' usually refers to pairing desirable stimuli such as food or play with the

fear-eliciting stimulus ('respondent' or 'classical counterconditioning') in order to reduce the

animal's distress through replacing the association with a positive one (McMillan, 2019).

Delivery of the incentive is contiguous only with the feared stimulus, regardless of the

animal's behavior. A behavioral change should follow as a result of a change of the animal's

emotional or motivational state (McMillan, 2019).

Some authors also describe the process of 'operant' or 'instrumental counterconditioning",

defined as "reinforcing a substitute behavior that is incompatible with the unwanted behavior"

(Horwitz and Mills, 2012, p.217). In contrast to 'classical counterconditioning', upon exposure

to the aversive stimulus, the animal is required to perform an operant response in order to

receive an incentive (McMillan, 2019). Based on early research, classical

counterconditioning is considered as superior to operant counterconditioning when aiming to

change negative emotional states (McMillan, 2019). Hereafter, I use the classic definition of

the term 'counterconditioning' as meaning that the feared stimulus is paired with an

appetitive/positive outcome (e.g. food) with the goal of decreasing the fear response over

repeated pairings and replacing it with an appetitive response (c.f. Newall et al., 2017).

Counterconditioning is most effective when combined with desensitization, i.e. the feared

stimulus is initially presented at a very low level and followed by an incentive. Subsequently,

this process can be repeated, whereby stimulus intensity can be gradually increased.

Nonetheless, using a desensitization approach in combination with counterconditioning is

not always possible as not all fear-inducing stimuli can be under complete human control,

and some fear improvements can still be achieved using counterconditioning when stimuli

are presented at full intensity. Crucial to the success of counterconditioning is identifying a

highly valued incentive (McMillan, 2019).

The use of noise recordings for systematic desensitization and counterconditioning

(abbreviated as DSCC) is the most commonly advised behavioral technique in the treatment

of noise fears (e.g. (Horwitz and Mills, 2012; Levine et al., 2007; Levine and Mills, 2008; Mills

et al., 2003; Sherman and Mills, 2008). Studies indicate that most owners perceive an

improvement in their dogs' firework fears after following a DSCC program using noise CDs

and are very satisfied with the treatment (Levine et al., 2007; Levine and Mills, 2008;

Sheppard and Mills, 2003), even up to one year later (Levine and Mills, 2008). However, one

study reports that no effect of treatment could be discerned using objective behavioral

measures when dogs were exposed to a novel CD recording in the clinic setting (Levine et

al., 2007), and this finding was similar to that of a previous study on the use of noise CDs (in

combination with medication) on thunderstorm fear in dogs: Despite owner-reported

improvement, no change in behavioral signs was noted during post-treatment exposure to a

thunderstorm recording in the clinic (Crowell-Davis et al., 2003).

Relaxation training

Training animals to relax on cue is a less commonly recommended tool in behavior

modification (but see Mills, 2003; Overall, 2013; Talegón and Delgado, 2011). Relaxation

may be achieved by different approaches. Horwitz and Mills (2012) suggest to firstly, induce

relaxation by massage or long strokes. Secondly, this can be associated with a word in order

to classically condition a calm physiological state. After successful conditioning, this cue can

then be used to induce relaxation during stressful events even without the need to massage

continuously (Horwitz and Mills, 2012). While this method relies on classical conditioning, an

operant 'protocol for training relaxation' is given in Overall (2013). Here, the dog is

progressively rewarded for behaviors, facial or bodily expressions consistent with relaxation,

and for remaining still for increasing amounts of time and in the face of gradually increased

distractions (Overall, 2013b). To what extent such training is effective in reducing fear or

anxiety in dogs has not been investigated to date.

2) Pharmacological interventions

Prescription medication

While benzodiazepines (e.g. Alprazolam), MAOIs, SSRIs, trazodone and the α2-

adrenoceptor agonist clonidine can be indicated in the medical management of noise fears

in dogs (Horwitz and Mills, 2012; Overall, 2013), only a handful of studies have investigated

the effects of medications on noise fears in dogs, and surprisingly few were placebo-

controlled. Two recent high-quality randomized, double-blind, placebo-controlled, clinical

field studies demonstrated a high effectiveness of Sileo® Dexmedetomidine oromucosal gel

(Korpivaara et al., 2017) and imepitoin (Engel et al., 2019), respectively. For Sileo®, 72% of

owners reported a good excellent treatment effect compared to only 37% in the placebo

group (Odds ratio in favor of Sileo®: 3.4; 95% confidence interval 1.95 to 5.99; $p<0.0001$).

This medication is already licensed for noise fears in dogs in the European Union at the time

of publication (European Medicines Agency, 2015). For imepitoin, around two thirds of

owners considered the treatment effect good or excellent treatment compared to

approximately a quarter of owners from the placebo group (Odds ratio in favor of imepitoin

4.7; 95% confidence interval 2.79-7.89; p<0.0001), which was also corroborated by highly

significant differences between the two groups in an additional measure, (owner-reported)

anxiety scores (Engel et al., 2019).

Only a few small, non-placebo controlled studies exist for the effect of other types of

medication on the expression of noise fears in dogs. Two small open-label studies indicated

that trazodone and clonidine, respectively, were effective in alleviating storm or noise phobia

in dogs where other treatments (including other medications) had failed (Gruen and

Sherman, 2008; Ogata and Dodman, 2011). A small prospective open clinical trial

investigated the effect of a combination of daily clomipramine, added alprazolam before

storms, and desensitization/ counterconditioning at home with a thunderstorm audio

recording on the behavior of dogs affected by fear of thunderstorms. While almost all

caregivers indicated an improvement in their dogs, this could not be confirmed when

exposing dogs to the audio recording in the clinic, although it is not clear whether post-

treatment recordings were made under the influence of alprazolam (Crowell-Davis et al.,

2003). Another small retrospective study indicated that diazepam was considered very or

somewhat effective by 67% of owners, but many owners discontinued the treatment due to

the occurrence of side effects (Herron et al., 2008).

3) Alternative products

Pheromonatherapy

Pheromonatherapy utilizes analogues of animal pheromones, chemical signals normally

involved in intraspecific communication that are processed by the vomeronasal organ and

are assumed to have an intrinsic effect on the emotional processing of animals (Mills et al.,

2012). Dog appeasing pheromone (DAP, Adaptil®) mimics a pheromone produced by

lactating bitches after parturition that is believed to instill a sense of well-being in the

puppies. It is used in veterinary behavioral medicine as a calming agent also for adult dogs

(Mills et al., 2012).

An open-label follow-up study indicated that the use of Dog Appeasing Pheromone plug-in

diffusers resulted in a high owner satisfaction level and reported improvement in dogs'

clinical signs during firework exposure (Sheppard and Mills, 2003), although the inclusion of

behavioral advice in the treatment plan does not allow to distinguish effects of behavioral

modification and pheromones (c.f. Frank et al., 2010). DAP was also used as in the above-

cited studies in combination with noise CDs (Levine et al., 2007; Levine and Mills, 2008).

One study suggested that the use of DAP and noise CDs in combination had higher owner-

reported success rates than when either intervention was used alone (Mills et al., 2003).

While the above studies were unblinded and did not include a placebo group, a blinded

parallel-group placebo controlled study on the effects of DAP collars on the behavior of

laboratory beagles during playbacks of thunderstorm recordings concluded that DAP is of

potential benefit as an adjunct to a behavior management program (Landsberg et al.,

2015a). In this study, observations of active (such as startling, scanning, bolting, pacing,

running, circling, digging, climbing, jumping and barking) and global fear scores were

significantly lower in the pheromone-treated group compared to the placebo group. No

significant group differences emerged in passive scores (such decreased activity, freezing,

lower body postures, panting, trembling, salivating, lip licking etc.). Hiding in a hiding box

occurred significantly more in placebo-treated subjects (Landsberg et al., 2015a).

Nutraceuticals

Given that nutraceuticals do not underlie the same stringent approval processes as

medications and therefore enter the market much more easily, many non-prescription

nutraceuticals are available for treating fear and anxiety in dogs (Orlando, 2018). However,

little published research on their effects exists, especially in regard to noise fears. For

example, Zylkene (alpha-casozepine) may be recommended to aid with noise fears although

it has not specifically been tested for this indication (Horwitz and Mills, 2012). The effect of L-

theanine on fear of thunder in dogs was investigated in a small open-label trial. However, a

high drop-out rate (eight of originally 26 subjects could not be included in the analysis due to

early withdrawal or failure to complete the paperwork) hinders drawing valid conclusions. For

the remaining 18 dogs, a highly significant reduction in owner-reported global anxiety scores

and time to return to baseline following a storm was found, and treatment success (defined

as "an improvement in the behavior score after the 5th storm compared with baseline of at

least 1 in 50% of the behaviors identified, with no behavior getting worse" was achieved in

12 of the 18 dogs (Pike et al., 2015). A fish protein supplement appears to have beneficial

effects on cortisol reactivity in dogs exposed to a thunderstorm recording (Landsberg et al.,

2015b).

Many owners tend to prefer to try "natural treatments" – such as herbal remedies,

homeopathy or essential oils – in the management of behavioral problems first, with

medication being regarded as only a last resort (Notari and Gallicchio, 2008). However, for

most of these products, there is either no peer-reviewed research at all, results are

inconclusive or negative.

Herbal formulations

A placebo-controlled crossover study investigated the effect of Harmonease Chewable

Tablets, containing a blend of extracts of *Magnolia officinalis* and *Phellodendron amurense*,

on inactivity duration during thunder storm recordings in 20 beagle dogs (with higher

inactivity interpreted as reflecting greater fear intensity). While an analysis of variance

indicated no significant treatment or order effect, the number of dogs reducing levels of

inactivity during thunderstorm recordings was higher in the treatment group (60%) than in

the placebo group (25%) (DePorter et al., 2012). Note, however, that later studies found that

many individuals react with an increase, rather than a decrease, in activity to thunderstorm

recordings (Gruen et al., 2015; Landsberg et al., 2015a, 2015b).

Homeopathy

A double-blind placebo-controlled study on the use of a homeopathic remedy in the

treatment of firework fears in dogs showed high owner-reported improvement rates in both

the placebo group (65%) and the verum group (71%), with no significant differences in any

measures between the two groups. While the owners were also given simply management

advice on how to react during firework exposure, this is unlikely to explain the perceived high

rates of success, with substantial placebo effects being more likely (Cracknell and Mills,

2008). As such, open-label studies should be interpreted with caution.

Aromatherapy (essential oils)

Odors, in particular essential oils, may be used in a therapeutic context to alleviate stress in

animals (Mills et al., 2012). As pointed out in the BSAVA Manual of Canine and Feline

Behavioral Medicine, there are suggestions that lavender and chamomile may promote

calmness in shelter dogs, but no evidence is available in relation to noise fears (Horwitz and

Mills, 2012).

Bach flowers

Although there is no science-based evidence for their effectiveness, Bach flowers may be

advised by some animal behavior professionals based on owners' preferences (Notari and

Gallicchio, 2008). Moreover, like Dog Appeasing Pheromone, herbal formulations,

homeopathic products and essential oils, Bach flowers are freely available over the counter

and so do not require a consultation with a professional.

4) Pressure vests

Studies on humans and on cattle indicate that deep pressure and weighted vests, or

pressure in a squeeze chute, respectively, have calming effects. While the mechanism of

action is not clear, it is suggested that peripheral oxytocin in response to the skin contact

may contribute to relieving stress, reducing blood pressure and heart rate (reviewed in

Pekkin et al., 2016). So-called pressure vests, tight-fitting vests for dogs exerting deep

pressure when worn, utilize this principle and are postulated to exhibit calming effects on pet

dogs. While the available evidence is inconclusive, there are some indications of effects on

some behavioral parameters and heart rate during thunderstorm or firework recordings that

could be interpreted as reflecting lowered anxiety when wearing a vest (Fish et al., 2017;

Pekkin et al., 2016). Also owners considered the vest to be of benefit during real-life

thunderstorm events (Cottam et al., 2013). A review concluded that "pressure vests may

have small but beneficial effects on canine anxiety and that habituating the dog to the vest,

assessing for comfort and using repeatedly may improve the likelihood of any benefit"

(Buckley, 2018). Several vests are commercially available, such as the Thundershirt, Anxiety

Wrap and Lymed Animal[TM] Supporting Garments.

To conclude, there is a lack of high-quality evidence for the majority of commonly

recommended interventions in the treatment of noise fears in dogs. Since almost all existing

studies suffer from very low sample sizes, it was decided to take a cross-sectional approach

here using an online internet survey targeting a variety of different interventions. Like most

previous studies, the study is based on the owners' assessment of their dogs' wellbeing

during noise exposure and the effectiveness of behavioral techniques or products used.

Methods

An online questionnaire survey (in an English and a German version) was distributed to dog

owners via our research group's website and social media. The questions covered the

owners' consent for the use of their data, demographic data about the dogs (date of birth,

sex, neuter status, breed, country, source of dog, age at acquisition), dogs' health problems

and other potential behavioral problems (Riemer, 2019). Two scores were used to rate the

severity and progression of firework fears, respectively (see below). Furthermore, owners

were asked to indicate which management and treatment options they had used by selecting

a number of options and, for the latter, whether they considered these as effective.

Welfare impaired score

As a measure of the severity of firework fears, participants were asked "Please rate your

level of agreement with the following statement: The overall welfare of my dog is strongly

compromised by fireworks". This was rated on a 5-point Likert scale ranging from "disagree

strongly", "tend to disagree", "partly/partly", "tend to agree" to "agree strongly". The ensuing

"Welfare Impaired" score was converted into a numerical score from 1 ("disagree strongly")

to 5 ("agree strongly") for the analysis.

Fear progression score

The "Fear progression" score was based on the question "How has your dog's fear of

fireworks progressed in the last years?", with the following response options "My dog was

never afraid of fireworks", "The fear has improved greatly", "The fear tends to have

improved", "The fear has remained the same", "The fear tends to have become worse", "The

fear has become much worse" or "I don't know". For analysis, this was converted into a

score ranging from 1 ("The fear has become much worse") to 5 ("The fear has improved

greatly"), with "I don't know" treated as missing data.

Management strategies

Owners were asked to select which management approaches (i.e. strategies during firework

exposure, independent of any preparatory interventions) they had used to help their dogs

cope with fireworks from a list of options (Table 1). They were also asked to select how they

responded to their dogs when hearing loud bangs (Table 1).

Effectiveness of interventions

Owners were asked to select from a number of options which interventions to prevent or

treat their dog's fear of fireworks they had tried and whether they considered these as

effective (response options: effective - not effective - not used; Table 2).

Table 1. Questions related to owners' management of firework fears (i.e. strategies during

firework events).

Which measures do you take on New Year's eve/ on days with fireworks? (Please select all applicable answers).	
	Nothing
	I create a safe hiding place for my dog
	I close the curtains/ window shutters
	I play loud music to drown out the noise
	I put a thundershirt/ anxiety wrap on my dog
	I offer chews to my dog
	I play with my dog
	I give food to my dog
How do you react yourself (to your dog) when hearing loud bangs? (Please select all applicable answers).	
	I ignore him or her/ I don't pay attention to him or her
	I pet him or her
	I give him or her a treat

	I play with him or her
	I allow him or her to maintain body contact
	I speak to him or her in a calming way
Have you administered medication or other calming products to your dog during the last firework event?	Yes - No
If yes, which ones? (Please select all applicable products).	
	Nutraceuticals (e.g. Zylkene, tryptophan etc.)
	Pheromone products (Adaptil diffuser, spray or collar)
	Herbal products
	Homeopathic products
	Bach flowers
	Essential oils
	Drugs (prescription medication available only from a veterinarian)
	Other

316   Table 2. Question on the perceived effectiveness of interventions for management or

317   training.

Which measures to treat or prevent firework fears did you consider as effective?	Response options: effective - not effective - not used
	Noise CD
	Pheromone products (such as Adaptil diffuser, collar or spray)

	Herbal products
	Homeopathic products
	Nutraceuticals (e.g. Zylkene, tryptophan etc.)
	Bach flowers
	Essential oils
	Drugs (prescription medication available only from a veterinarian)
	Counterconditioning (for example, giving a treat or playing with the dog after a noise has occurred)
	Relaxation training
	Thundershirt/ anxiety wrap
	Other

Analysis

In order to identify patterns in management strategies, a non-linear principal components

analysis (Linting et al., 2007; Linting and van der Kooij, 2012) or Categorical Principal

Components Analysis (CATPCA) in IBM SPSS Statistics Version 23 (© IBM Corporation and

its licensors 1989, 2015) was performed over the questions on owners' management

strategies during fireworks, including measures taken during New Year's Eve/ on days with

fireworks (e.g. creating a hiding place, closing windows, playing music etc.), how owners

reacted to loud bangs themselves (e.g. ignoring the dog, petting the dog, giving the dog a

treat etc.), and which types of products they had administered (e.g. pheromone products,

prescription medication, pressure vests etc.). To assess relationships of management

factors and fear development, the ensuing components were correlated with the Fear

Progression Score using Spearman rank correlation tests, as requirements for parametric

analysis were not met, using Statistica 6.1 (Statsoft Inc. 1984–2004).

For the specific questions on different products and training techniques used, the percentage

of respondents considering a given intervention as effective was calculated from the total

number of respondents that indicated having used this intervention.

Results

A total of 1225 responses were analyzed, including 527 responses in English and 699 in

German. Dogs were of various breeds or crosses and included 588 females (of which 430

neutered, 157 intact and one of unknown neuter status) and 637 males (of which 424

neutered, 207 intact and 6 chemically castrated). Over half the dogs in the survey (N=639)

were given a rating of 3 or higher on the Welfare Impaired score and so are considered as

fearful of fireworks (see (Riemer, 2019) for more details).

Management strategies

Regarding owners' management during firework exposure, the CATPCA yielded four factors

(based on Eigenvalues >1 and Cronbach's Alpha >0.7) that had no cross-loadings and

accounted for 49% of the variance (Table 3).

Table 3. Results of a CATPCA on management strategies employed by owners during

fireworks, with Varimax rotation. Loadings >0.4 are bolded.

		Environmental Modification	Feed/ Play	Alternative	Interaction	Total
Measures during firework events	No measures	**-0.718**	-0.210	-0.087	-0.256	
	Hiding place	**0.654**	0.049	0.148	0.217	
	Windows closed	**0.803**	0.104	0.100	0.100	
	Music	**0.722**	0.131	0.164	0.030	
	Thundershirt	0.351	0.077	0.400	0.044	
	Chews	0.238	**0.684**	0.038	0.003	

	Play	0.130	**0.751**	-0.064	0.081
	Food	0.074	**0.749**	0.071	0.086
Reaction to bangs	Ignore	-0.057	-0.071	-0.021	**-0.629**
	Pet	0.040	0.208	0.104	**0.760**
	Feed	-0.001	**0.658**	0.035	0.323
	Play	-0.035	**0.718**	-0.005	0.193
	Body contact	0.313	0.115	0.092	**0.585**
	Talk	0.257	0.189	0.083	**0.701**
Products used	Nutraceuticals	0.244	0.024	**0.443**	0.030
	Pheromones	0.248	0.005	**0.586**	0.021
	Herbal products	0.123	0.059	**0.729**	0.000
	Homeopathy	0.078	-0.036	**0.748**	0.051
	Bach flowers	0.112	-0.057	**0.645**	0.094
	Essential oils	-0.044	0.053	**0.580**	0.086
	Prescription medication	0.394	-0.048	0.213	0.066
	Cronbach's alpha	**0.769**	**0.731**	**0.726**	**0.708**	**0.948**
	Eigenvalue	**2.795**	**2.735**	**2.693**	**2.112**	**10.335**
	% of variance	**13.309**	**13.021**	**12.825**	**10.058**	**49.214**

The first component, termed *"Environmental modification"* had high positive loadings for

providing a hiding place, keeping windows and blinds closed, and playing music and a high

negative loading for not taking any measures. The use of a thundershirt or anxiety wrap and

prescription medication also loaded > 0.35 on this component. The second component,

"Feed/Play", had high positive loadings for providing the dog with chews, play and food

during fireworks in general, as well as for feeding and playing with the dog in response to

loud bangs. The third component, labelled *"Alternative"* had high positive loadings for the

use of calming nutraceuticals, pheromones, herbal products, homeopathic products, Bach

flowers, and essential oils. The fourth component, *"Interaction"*, had high positive loadings

for allowing body contact, petting and talking to the dog when loud bangs occurred, and a

high negative loading for ignoring the dog.

Within those dogs with Impaired Welfare scores of 3 or higher, the principal component

"Environmental Modification" was significantly positively correlated with Fear Progression

(i.e. an increase in severity of firework fears, Spearman rank correlation test, Rho=0.181,

N=566, p=0.00001) whereas the component "Feed/ Play" was significantly negatively

correlated with Fear Progression (Spearman Rho=-0.16, N=566, p=0.00008). There was no

relationship between Fear Progression and the components "Alternative" (Spearman

Rho=0.026, N=566, p=0.527) and "Interaction" (Spearman Rho=-0.03, N=566, p=0.357).

Owner-reported effectiveness of training techniques and products used

Noise CDs were considered as effective in alleviating dogs' firework fears by 54.4% of

owners who had used them (N=377; Figure 1). The reported effectiveness was 70.8% for

counterconditioning (N=694) and 69.3% for relaxation training (N=433). A success rate of

44% was indicated for the use of a thundershirt/ anxiety wrap (N=300). For prescription

medication, the reported rate of effectiveness was 68.9% (N= 202; Figure 2). Of those

owners who had used pheromone products (Adaptil® diffuser, spray or collar, N=316),

28.8% considered these to be effective. The reported success rates were 27.0% for

nutraceuticals (N=211), 35.1% for herbal products (N=282), 31.2% for homeopathic products

(N=250), 33.5% for Bach flowers (N=33.5) and 31.1% for essential oils (N=183),

respectively.

Figure 1. Owner-reported success rates of different behavior modification approaches and

pressure vests.

Medication (N=202)
Pheromone products (N=316)
Nutraceuticals (N=211)
Herbal products (N=282)
Homeopathic products (N=250)
Bach flowers (N=281)
Essential oils (N=183)

0%　20%　40%　60%　80%　100%

■Effective　□Not effective

Figure 2. Owner-reported success rates of prescription medication, pheromones, calming

nutraceuticals, herbal remedies, homeopathic products, Bach flowers and essential oils.

Since the question on which specific products were used were optional, data on the type of

prescription medication used were available for 103 dogs. Of those, 86 received one type of

drug, 13 received a combination of two drugs and four received three drugs. Alprazolam

(N=32) and Dexmedetomidine (N=19) were the most frequently prescribed single drugs. The

394 most frequent combination was Trazodone & Alprazolam (N=7). Other combinations

395 included Dexmedetomidine combined with Diazepam or Alprazolam. Table 4 shows the

396 reported effectiveness of the drugs in the sample, bearing in mind that the sample size was

397 very small for most types of medication.

399 Table 4. Owner-reported effectiveness of different prescription medications

	N	Effective	Effective (%)
Alprazolam	32	29	90.63
Dexmedetomidine	19	14	73.68
Diazepam	8	4	50
Trazodone	8	6	75
Acepromazine	4	4	100
Benadryl	3	3	100
Fluoxetine	3	3	100
Clomipramine	2	2	100
Clonidine	2	0	0
Sertraline	1	1	100
Tramadol	1	1	100
>1 drug	17	13	76.47
Trazodone & Alprazolam	7	7	100

Discussion

402 An online survey was performed to explore how owners managed potential firework fears in

403 their dogs and which interventions they perceived as effective in improving dogs' firework

404 fears.

Management

In order to gauge whether management measures can affect the progression of firework

fears in dogs, the four management factors were correlated with the Fear progression score.

The results indicated that Environmental modification" (providing a hiding place, keeping

windows and blinds shut, and playing music) was associated with a significant deterioration

of firework fears, whereas provision of chews, play and food during fireworks in general, as

well as deliberately providing these rewards when loud bangs occurred (principal component

"Feed/Play") was associated with a significant improvement. No association was found

between fear progression and the "Alternative" component (use of nutraceuticals,

pheromones, herbal products, homeopathic products, Bach flowers, and essential oils) or

the "Interaction" component (allowing body contact, petting and talking to the dog when loud

bangs occurred).

The positive correlation between Fear progression and "Environmental Modification" is

unlikely to indicate that measures of environmental modification were *contributing* to

increased fear severity. Conversely it is probable that owners who perceived an increased

fear response in their dogs were more likely to take measures of environmental modification,

since these are commonly recommended measures for dogs fearful of fireworks (e.g. Mills et

al., 2003; Pike et al., 2015; Sherman and Mills, 2008). Of real interest, however, is the

significant negative association of Fear progression and the "Feed/Play" component. It is

reasonable to assume that this reflects a positive effect of the intervention on fear

progression rather than vice versa, since owners experiencing an improvement in their dogs'

fears should be less likely and not more likely to attempt a given intervention. It appears that

using appetitive stimuli during firework exposure not only helped to keep the fear at bay

during the event, but may even have contributed to a longer-lasting improvement in

fearfulness, possibly exerting some counterconditioning effects even without a systematic

desensitization/ counterconditioning approach. Alternatively, it is possible that dog owners

who used food and toys during fireworks also performed ad-hoc counterconditioning

(providing these incentives contingent on occurrence of noises) in everyday life.

A common recommendation is to ignore the dog when it shows fearful behavior (e.g. Mills et

al., 2003; Pike et al., 2015; Sherman and Mills, 2008). The current study indicated that

interacting with the dog is neither associated with improvement nor with deterioration of the

dogs' fear level over time. Similarly, Dreschel and Granger (2005) found no influence of the

owner's behavior on behavior and cortisol levels in dogs after a playback of a thunderstorm

recordings. Whether interacting with a fearful dog has beneficial or detrimental effects on

fear level in the firework situation itself requires further investigation. There is, however,

evidence from other stressful situations that being petted and talked to is associated with

lower physiological or behavioral stress indicators in dogs (e.g. Csoltova et al., 2017;

Hennessy et al., 1998; Lynch and McCarthy, 1967), and some physiological mechanisms for

this have been identified (e.g. Handlin et al., 2011; Kostarczyk and Fonberg, 1982; Odendaal

and Meintjes, 2003; Rehn et al., 2014). Thus, there might be potential beneficial effects of

attending to fearful dogs also in firework situations, even if it cannot be ruled out that this

reinforces some behavioral actions, such as attention-seeking, by instilling some fear relief

(negative reinforcement).

Behavior modification and training

The above-mentioned correlation between higher scores on the "Feed/Play" principal

component and a more favorable fear progression has indicated that using food and play

during fireworks is an effective way of alleviating noise fears in dogs. This is also confirmed

by the owners' own perceptions: Counterconditioning (defined in the questionnaire as "for

example, giving a treat or playing with the dog after a noise has occurred") was considered

to be effective by 70.8% of those who had tried this technique. It is commonly believed that

desensitization/ counterconditioning can only be successful in achieving a lasting emotional/

behavioral change if pets are kept under their threshold or tolerance level (Horwitz and Pike,

2014, but see a critical evaluation of this assumption in Klein, 1969; Wilson and Davison,

1971). While this would clearly be the ideal approach, the results of the current study

indicate that even ad-hoc, non-systematic, counterconditioning in everyday life (such as

giving a treat, playing with the dog or celebrating a little 'party') whenever a loud noise

occurs, as well as during firework events, can contribute to an improvement of firework fears

in dogs. While some dogs may be too stressed to eat during exposure to loud noises,

providing high-value food after any noise occurrence is a simple strategy that can be easily

implemented also by inexperienced owners and was actually used by over half the

respondents in this study (N=694). Using this technique as a preventative measure appears

to be extremely effective in preventing the development of firework fears in the first place

(Riemer, 2019). This method is thus highly recommendable in both the prevention and

treatment of noise fears in dogs.

Interestingly, with a success rate of a 69.3% according to the owners, relaxation training was

similarly effective as counterconditioning in alleviating firework fears in dogs. Although

relaxation was an inherent component of the original concept of "systematic desensitization"

in people sensu Wolpe, (1958, reviewed in Thomas et al., 2012; Wilson and Davison, 1971),

relaxation training is less prominently mentioned in the current literature on clinical animal

behavior. Nonetheless, a relatively high number of respondents in the sample (N=433) had

attempted this technique. The current questionnaire did not go into detail about the type of

training protocols that were used, i.e. whether a classically conditioned approach was mostly

used or whether respondents achieved a relaxed state through positive reinforcement of

relaxed behaviors. Further research is therefore warranted on the optimal relaxation

protocols. Similarly as with counterconditioning, it appears that respondents did not

necessarily use relaxation training in a systematic way with a gradual increase in stimulus

intensity; nonetheless, the technique seems to have been successful in a large proportion of

cases, at least according to the owners' perceptions.

The current gold standard and most studied behavioral approach in the therapy of noise

fears is to perform desensitization/ counterconditioning (DSCC) using noise recordings (e.g.

Horwitz and Mills, 2012; Levine et al., 2007; Levine and Mills, 2008; Mills et al., 2003;

Sherman and Mills, 2008). Nonetheless, at 54.4% reported effectiveness, the use of noise

recordings had a considerably lower success rate than ad-hoc counterconditioning and

relaxation training in the current study. The rate of success is also somewhat lower than

reported in previous studies on the use of noise CDs, possibly reflecting the more long-term

view. For example, in a follow-up study one year after the first implementation of DSCC

using noise CDs, 66% of owners indicated either "a moderate or great improvement" in their

dog's fear of fireworks, even though only few of the respondents had continued to use the

CD recording (Levine and Mills, 2008), and reported success rates were even higher in the

short-term (Levine et al., 2007; Sherman and Mills, 2008).

Perhaps owners who received the CD from professionals in veterinary behavior medicine

were more motivated to follow through the procedure, or they had received better

instructions on how to approach DSCC – although Mills et al. (2003) and Levine et al. (2007)

found that many owners did not necessarily adhere to the instructions very well, and that two

CDs with different types of instructions appeared to be similarly effective. The published

studies had relatively high dropout rates. While personal reasons were mostly given (Levine

et al., 2007), it cannot be ruled out that those owners who perceived no success of treatment

were more likely to discontinue and were thus not available for reporting on the perceived

success (or lack thereof) at the study end. Also with relatively small sample sizes in the cited

studies compared to the current one, more random variation can be expected in the former.

Additionally, treatment success might be over-reported when personal follow-up via phone

calls are made as in Levine et al. (2007), Levine and Mills (2008) and Mills et al. (2003),

since it is suggested that patients may report positive outcomes to the treating clinician out

of politeness, enhancing the placebo effect (Kienle and Kiene, 1997). It is furthermore

possible that the advice to introduce a "safe haven" ("a location in the home in which the dog

had only positive experiences – not the same location to which the dog normally hid when

fearful" (Levine et al., 2007) and the provision of Dog Appeasing Pheromone (Levine et al.,

2007) enhanced the treatment success compared to the use of CD recordings alone (c.f.

Mills et al., 2003).

Some difficulties with using noise recordings for DSCC have been recognized. They may not

be realistic enough, depending on the quality of the recording or the speakers, the setup of

the room influencing acoustics, and lacking associated stimuli such as flashes (Sheppard

and Mills, 2003; Shull-Selcer and Stagg, 1991). Even under optimal conditions, some dogs

will not respond to the simulations (Shull-Selcer and Stagg, 1991). Moreover, DSCC using

noise recording is a time-consuming process, and may need to be repeated periodically to

prevent a relapse, which some owners may be unwilling or unable to provide (Levine and

Mills, 2008; Sheppard and Mills, 2003).

Thus, while noise recordings clearly have a place in the treatment of noise fears, ad-hoc

counterconditioning to noises and relaxation training should complement this treatment

strategy. In particular, using treats or other incentives whenever a (loud) noise occurs may

be an easier strategy to implement in everyday life for many owners, as unlike both DSCC

with noise recordings and relaxation training, it requires no targeted training other than

having some food on hand – although clearly, the more structured training can be done, the

better the outcome is likely to be.

Products

While the reported success rates for counter-conditioning and relaxation training were

relatively high, behavioral methods on their own may often not be sufficient to provide

adequate fear relief. Accordingly, a wide range of products with putatively calming properties

are on the market; however, for most of these, there is a lack of good-quality evidence on

their effectiveness.

In the current study, the highest success rates of all product categories was achieved for

prescription medication, even though it can be presumed that dogs that were medicated

were affected by more severe noise fears than those receiving no medication. Since

medications undergo a rigorous testing process (albeit not necessarily for the indication

"noise fears") before they are licensed, and work at the level of the central nervous system,

this should be expected. While the overall sample size for users of medication was large in

the current study (N=202), only a smaller number of owners had indicated which

medications they had used. Of primary interest here are the two most commonly reported

medications, Alprazolam (N=32) and Dexmedetomidine (N=19). The success rate was very

high for Alprazolam at over 90%, while Dexmedetomidine was considered to be effective in

over 73% of cases. The latter figure is very similar to the published effectiveness of

Dexmedetomidine in a placebo-controlled study, in which a good or excellent treatment

effect was found in 72% in the dexmedetomidine group, compared to 37% in the placebo

group (Korpivaara et al., 2017). Too few respondents have named other types of

medications so that no definitive conclusions can be drawn regarding their effectiveness.

Overall, pharmacological treatment is no doubt recommendable for dogs with severe noise

fears, both as a management strategy to help them cope in the case of firework events and

possibly to complement training efforts and facilitate learning during behavior modification

treatment. Many owners, however, view medication only as a last resort and prefer to use

more "natural" treatments, due to concerns such as potential side effects (Notari and

Gallicchio, 2008; Sheppard and Mills, 2003). Unfortunately, very few of these products have

demonstrated any evidence of effectiveness (Landsberg et al., 2015b).

Remarkably, all types of 'alternative' products in the current survey – including herbal

products, nutraceuticals, homeopathic remedies, essential oils, but also pheromones – had

reported success rates in the range of 27 to 35 percent. This is approximately the rate of

success that would be expected based on a placebo effect. A placebo is "an inert medication

used for its psychological effect, or for purposes of comparison in an experiment", with a

placebo effect defined as "any improvement or change in subjective discomfort or illness

resulting from an intervention possessing no physical effect" (Tavel, 2014). When other

individuals (clinicians or caregivers) rate the effectiveness of a given intervention, we may

observe a caregiver placebo effect, which often has even larger effect sizes than those

reported by the patients themselves (Rief et al., 2009). A caregiver effect in companion

animals has also been defined as "improved ratings of outcomes in companion animals in

the absence of improvement in objective measures" (Gruen et al., 2017). Whether a

placebo-by-proxy effect exists in veterinary medicine – meaning that the caregiver's belief

alters their interaction with the pet which consequentially has a true beneficial effect on the

animal – is not known (Gruen et al., 2017). Instead it is concerning that many interventions

leading to better welfare according to the owners' subjective assessments may in fact have

no benefit on the animal at all.

In the medical literature, response rates to placebos are commonly in the range of around

35% (Muñana et al., 2010). In companion animals, caregiver placebo effects of up to 50-

70% have been reported (Gruen et al., 2017). In the study by (Korpivaara et al., 2017) on the

effect of dexmedetomidine vs placebo on firework fears in dogs, 37% of dogs in the placebo

groups had a good or excellent treatment effect as judged by their owners. In the study by

Cracknell and Mills (2008) evaluating the effect of a homeopathic remedy on firework fears

in dogs, the reported rates of improvement did not differ significantly between treatment

groups but were extremely high in both the placebo group (65%) and the treatment group

(71%), although the fact that simple behavioral advice had also been given might have

contributed to these high success rates in both groups (Cracknell and Mills, 2008). In a study

on fluoxetine in the treatment of canine separation anxiety, global separation anxiety scores

in the placebo-treated dogs improved in 44-51.3% of cases across the six weeks of

treatment (Landsberg et al., 2008).

Thus, it seems likely that a placebo effect accounts for the perceived effectiveness of those

products where the success rates were 35% or less in the current study, which was the case

for pheromone products, nutraceuticals, herbal remedies, essential oils, as well as

homeopathic remedies. In this context it is also notable that the many promising open-label

trials have not been followed up by placebo controlled ones. Possibly this reflects a

publication bias, with positive results from open-label trials potentially being more likely to be

published than negative results in controlled studies.

Of the above-mentioned products, only Dog Appeasing Pheromone (Adaptil®) has been

investigated in several studies on fear and anxiety in dogs. Some open-label trials reported

high treatment success rates, but these were often confounded by the parallel application of

the pheromones and behavioral management or CCSD training (Levine et al., 2007; Levine

and Mills, 2008; Sheppard and Mills, 2003). One study suggested that using a combination

of CDs and DAP resulted in better outcomes than when only one of these treatments was

used (Mills et al., 2003). However, the high success rates for placebo treatments observed in

placebo-controlled studies on treatments for firework fears in dogs (Cracknell and Mills,

2008; Korpivaara et al., 2017) call for a cautious interpretation of open-label studies. One

parallel-group placebo controlled study using objective behavioral measures did indicate

some beneficial effects of DAP collars on global and active fear scores during exposure to

thunderstorm recordings in laboratory beagles. On the other hand, DAP treated dogs also

spent more time hiding, so the results are not entirely conclusive, and the sample size was

small (12 beagles per treatment group, Landsberg et al., 2015a). A systematic review on

pheromone application in dogs and cats concluded that only a single study yielded sufficient

evidence for a reduction in fear and anxiety by DAP in dogs, while six studies yielded

insufficient evidence (Frank et al., 2010). Thus, also in view of the present study's results,

there is currently little evidence to support the effectiveness of dog appeasing pheromones

as a calming agent in dogs fearful of fireworks. Similarly, no recommendations can be made

for the use of nutraceuticals, herbal remedies, essential oils, and homeopathic products. Of

course, perhaps with the exception of pheromones where the same ingredient, Dog

Appeasing Pheromone, formed the basis of all products (diffusers, sprays or collars), the

study does not allow drawing conclusions for a specific product in this case, as all products

in a category were analyzed together.

Finally, the reported effectiveness for pressure vests (thundershirt, anxiety wrap) was 44%.

Although this could still be in a range where a placebo effect is possible, the proportion of

respondents considering pressure vests effective was considerably higher compared to all

the 'alternative' products, indicating that pressure vests may potentially have beneficial

effects in some dogs. Further research is needed, as previous studies indicated a possible

small beneficial effect but were not conclusive either (reviewed in Buckley, 2018).

Critical evaluation

Like most studies assessing the effectiveness of treatments for noise fears in dogs (e.g.

Cracknell and Mills, 2008; Crowell-Davis et al., 2003; Korpivaara et al., 2017; Levine et al.,

2007; Levine and Mills, 2008; Sheppard and Mills, 2003), the current study relied on owners'

assessments of their dogs' behavior. Therefore it cannot be avoided that the results are

affected by the owners' interpretation and may be subject to bias. However, studies

indicated that owners' ratings of their dogs' fear are reliable and demonstrate external

validity (e.g. Tiira and Lohi, 2014), although they are susceptible to caregiver placebo effects

(e.g. Cracknell and Mills, 2008). Based on previous placebo-controlled studies, we know

approximately in what range placebo effects can be expected, as discussed above. While

randomized double-blind placebo-controlled studies, and using video analysis to assess

dogs' behavior, are no doubt the gold standard for assessing the effect of a given

intervention, they are often limited by the costs and manpower required for both conducting

and administrating the study. Additionally, ethical concerns might be raised (c.f. Frank et al.,

2010). Using a large-scale questionnaire survey thus represents a useful approach to gather

data from a large sample of participants. Moreover, a broad survey like the current one may

be less likely than localized trials of specific products to attract volunteer participants with

high expectations that might lead to an overestimate of effectiveness (c.f. Muñana et al.,

2010), or to induce participants to over-report effectiveness of treatments out of politeness to

the treating doctor (Kienle and Kiene, 1997). It can thus be viewed as an advantage that the

prescribing clinicians and the researchers were not the same persons in the current study.

Owners were furthermore able to draw from their long-term experiences (i.e. not only for a

few weeks within a particular study). While recall biases cannot be ruled out, the possibility

to test a given product over a longer time period or during several firework events may lead

to a more accurate assessment than when owners are asked to make an assessment over

only a few weeks.

By asking owners to simply rate a given intervention as "effective" or "not effective", the

outcome is relatively crude. Nonetheless it is useful, since effectiveness of products is often

measured by the proportion in the population in which the treatment had an effect, and

together with the large sample size, it allows us to compare relative effectiveness of different

treatments – even if it does not allow conclusions in absolute numbers regarding the true

effectiveness of the interventions.

Summary

Owner-reported effectiveness of different treatment options for firework fears in dogs was

analyzed from a large-scale questionnaire survey, with over 180 respondents per treatment

category. Ad-hoc counter-conditioning (providing desirable stimuli after the occurrence of

noises) was associated with a significant improvement in the severity of firework fears and

was considered as effective by over 70% of owners. Relaxation training was reported to be

almost as successful, while noise CDs were considered effective by only 55%. Forty-four

percent of respondents deemed pressure vests to be effective. With prescription medication,

improvements were noted by 70% of owners. While individual products were not evaluated,

the reported success rates for the categories "pheromones", "herbal products",

"nutraceuticals", "essential oils", "homeopathic remedies" and "Bach flowers" (all in the range

of 27-35%) are not higher than would be expected based on a placebo effect. Thus, besides

use of anxiolytic medication, counterconditioning and relaxation training appear to be the

most effective strategies in the treatment of firework fears in dogs and should complement

the standard behavioral technique of desensitization/ counterconditioning with noise

recordings.

Acknowledgements

Many thanks to all the dog owners who took the time to fill in this questionnaire and to

everybody who helped to spread the survey. Further thanks go to Hanno Würbel and Alja

Mazzini for feedback on the manuscript. S.R. was supported by an SNF Ambizione Grant

Project PZ00P3_174221.

Conflict of interest statement

The author declares no competing interests.

Authorship statement

The idea for the paper was conceived by Stefanie Riemer. The questionnaire design,

distribution, analysis and writing was performed by Stefanie Riemer.

Data Availability

The raw data set can be accessed

under https://figshare.com/articles/Riemer_2019_dogs_fireworks_data_xls/9657395.

References

Blackwell, E.J., Bradshaw, J.W.S., Casey, R.A., 2013. Fear responses to noises in domestic

   dogs: Prevalence, risk factors and co-occurrence with other fear related behaviour.

   Appl. Anim. Behav. Sci. 145, 15–25.

Buckley, L.A., 2018. Are Pressure Vests Beneficial at Reducing Stress in Anxious and

   Fearful Dogs? Vet. Evid. 3.

Cottam, N., Dodman, N.H., Ha, J.C., 2013. The effectiveness of the Anxiety Wrap in the

   treatment of canine thunderstorm phobia: An open-label trial. J. Vet. Behav. Clin. Appl.

   Res. 8, 154–161.

Cracknell, N.R., Mills, D.S., 2008. A double-blind placebo-controlled study into the efficacy of

   a homeopathic remedy for fear of firework noises in the dog (*Canis familiaris*). Vet. J.

   177, 80–88.

Crowell-Davis, S.L., Seibert, L.M., Sung, W., Parthasarathy, V., Curtis, T.M., 2003. Use of

   clomipramine, alprazolam, and behavior modification for treatment of storm phobia in

   dogs. J. Am. Vet. Med. Assoc. 222, 744–748.

Csoltova, E., Martineau, M., Boissy, A., Gilbert, C., 2017. Behavioral and physiological

   reactions in dogs to a veterinary examination: Owner-dog interactions improve canine

   well-being. Physiol. Behav. 177, 270–281.

DePorter, T.L., Landsberg, G.M., Araujo, J.A., Ethier, J.L., Bledsoe, D.L., 2012. Harmonease

   Chewable Tablets reduces noise-induced fear and anxiety in a laboratory canine

   thunderstorm simulation: a blinded and placebo-controlled study. J. Vet. Behav. Clin.

   Appl. Res. 7, 225–232.

Dreschel, N.A., Granger, D.A., 2005. Physiological and behavioral reactivity to stress in

   thunderstorm-phobic dogs and their caregivers. Appl. Anim. Behav. Sci. 95, 153–168.

Engel, O., Müller, H.W., Klee, R., Francke, B., Mills, D.S., 2019. Effectiveness of imepitoin

for the control of anxiety and fear associated with noise phobia in dogs. J. Vet. Intern.

Med. 33, 2675–2684.

European_Medicines_Agency, 2015. Sileo Dexmedetomidine hydrochloride - EPAR

summary for the public.

Fish, R.E., Foster, M.L., Gruen, M.E., Sherman, B.L., Dorman, D.C., 2017. Effect of Wearing

a Telemetry Jacket on Behavioral and Physiologic Parameters of Dogs in the Open-

Field Test. J. Am. Assoc. Lab. Anim. Sci. 56, 382–389.

Frank, D., Beauchamp, G., Palestrini, C., 2010. Systematic review of the use of pheromones

for treatment of undesirable behavior in cats and dogs. J. Am. Vet. Med. Assoc. 236,

1308–1316.

Gruen, M.E., Case, B.C., Foster, M.L., Lazarowski, L., Fish, R.E., Landsberg, G., DePuy, V.,

Dorman, D.C., Sherman, B.L., 2015. The use of an open-field model to assess sound-

induced fear and anxiety-associated behaviors in Labrador retrievers. J. Vet. Behav.

10, 338–345.

Gruen, M.E., Dorman, D.C., Lascelles, B.D.X., 2017. Caregiver Placebo Effect in Analgesic

Clinical Trials for Painful Cats with Naturally-Occurring Degenerative Joint Disease.

Vet. Rec. 180, 473.

Gruen, M.E., Sherman, B.L., 2008. Use of trazodone as an adjunctive agent in the treatment

of canine anxiety disorders: 56 cases (1995--2007). J. Am. Vet. Med. Assoc. 233,

1902–1907.

Handlin, L., Hydbring-sandberg, E., Nilsson, A., Ejdebäck, M., Jansson, A., Uvnäs-moberg,

K., 2011. Short-Term Interaction between Dogs and Their Owners⊡: Effects on

Oxytocin, Cortisol, Insulin and Heart Rate — An Exploratory Study. Anthrozöos 24,

301–315.

Hennessy, M.B., T Williams, M., Miller, D.D., Douglas, C.W., Voith, V.L., 1998. Influence of

male and female petters on plasma cortisol and behaviour: can human interaction

reduce the stress of dogs in a public animal shelter? Appl. Anim. Behav. Sci. 61, 63–77.

Herron, M.E., Shofer, F.S., Reisner, I.R., 2008. Retrospective evaluation of the effects of

diazepam in dogs with anxiety-related behavior problems. J. Am. Vet. Med. Assoc. 233,

1420–1424.

Horwitz, D., Mills, D., 2012. BSAVA manual of canine and feline behavioural medicine,

second. ed. BSAVA.

Horwitz, D.F., Pike, A.L., 2014. Common sense behavior modification: a guide for

practitioners. Vet. Clin. Small Anim. Pract. 44, 401–426.

Kienle, G.S., Kiene, H., 1997. The powerful placebo effect: fact or fiction? J. Clin. Epidemiol.

50, 1311–1318.

Klein, B., 1969. Counterconditioning and fear reduction in the rat. Psychon. Sci. 17, 150–

151.

Korpivaara, M., Laapas, K., Huhtinen, M., Schöning, B., Overall, K., 2017. Dexmedetomidine

oromucosal gel for noise-associated acute anxiety and fear in dogs—a randomised,

double-blind, placebo-controlled clinical study. Vet. Rec. 180, 356.

Kostarczyk, E., Fonberg, E., 1982. Heart rate mechanisms in instrumental conditioning

reinforced by petting in dogs. Physiol. Behav. 28, 27–30.

Landsberg, G., Beck, A., Lopez, A., Deniaud, M., Araujo, J., Milgram, N., 2015a. Dog-

appeasing pheromone collars reduce sound-induced fear and anxiety in beagle dogs: a

placebo-controlled study. Vet. Rec. 177, 260.

Landsberg, G., Mougeot, I., Kelly, S., Milgram, N., 2015b. Assessment of noise-induced fear

and anxiety in dogs: modification by a novel fish hydrolysate supplemented diet. J. Vet.

Behav. 10, 391–398.

Landsberg, G.M., Melese, P., Sherman, B.L., Neilson, J.C., Zimmerman, A., Clarke, T.P.,

2008. Effectiveness of fluoxetine chewable tablets in the treatment of canine separation

anxiety. J. Vet. Behav. 3, 12–19.

Levine, E.D., Mills, D.S., 2008. Long-term follow-up of the efficacy of a behavioural treatment

programme for dogs with firework fears. Vet. Rec. 162, 657–9.

Levine, E.D., Ramos, D., Mills, D.S., 2007. A prospective study of two self-help CD based

desensitization and counter-conditioning programmes with the use of Dog Appeasing

Pheromone for the treatment of firework fears in dogs (*Canis familiaris*). Appl. Anim.

Behav. Sci. 105, 311–329.

Linting, M., Meulman, J.J., Groenen, P.J.F., van der Koojj, A.J., 2007. Nonlinear principal

components analysis: introduction and application. Psychol. Methods 12, 336.

Linting, M., van der Kooij, A., 2012. Nonlinear principal components analysis with CATPCA:

a tutorial. J. Pers. Assess. 94, 12–25.

Lynch, J.J., McCarthy, J.F., 1967. The effect of petting on a classically conditioned emotional

response. Behav. Res. Ther. 5, 55–62.

McMillan, F.D., 2019. Mental health and well-being in animals, 2nd ed. CABI.

Mills, D., 2005. Management of noise fears and phobias in pets. In Pract. 27, 248.

Mills, D., Dubé, M.B., Zulch, H., 2012. Principles of pheromonatherapy. Stress

Pheromonatherapy Small Anim. Clin. Behav. 127–145.

Mills, D.S., 2003. Medical paradigms for the study of problem behaviour: a critical review.

Appl. Anim. Behav. Sci. 81, 265–277.

Mills, D.S., Estelles, M.G., Coleshaw, P.H., Shorthouse, C., 2003. Retrospective analysis of

the treatment of firework fears in dogs. Vet. Rec. 153, 561–562.

Muñana, K.R., Zhang, D., Patterson, E.E., 2010. Placebo effect in canine epilepsy trials. J.

Vet. Intern. Med. 24, 166–170.

Newall, C., Watson, T., Grant, K.-A., Richardson, R., 2017. The relative effectiveness of

extinction and counter-conditioning in diminishing children's fear. Behav. Res. Ther. 95,

42–49.

Notari, L., Gallicchio, B., 2008. Owners' perceptions of behavior problems and behavior

therapists in Italy: A preliminary study. J. Vet. Behav. 3, 52–58.

Odendaal, J.S.J., Meintjes, R.A., 2003. Neurophysiological correlates of affiliative behaviour

between humans and dogs. Vet. J. 165, 296–301.

Ogata, N., Dodman, N.H., 2011. The use of clonidine in the treatment of fear-based behavior

problems in dogs: an open trial. J. Vet. Behav. Clin. Appl. Res. 6, 130–137.

Orlando, J.M., 2018. Behavioral nutraceuticals and diets. Vet. Clin. Small Anim. Pract. 48,

473–495.

Overall, K., 2013. Manual of clinical behavioral medicine for dogs and cats. Elsevier, St

Louis, MO.

Pekkin, A.-M., Hänninen, L., Tiira, K., Koskela, A., Pöytäkangas, M., Lohi, H., Valros, A.,

2016. The effect of a pressure vest on the behaviour, salivary cortisol and urine

oxytocin of noise phobic dogs in a controlled test. Appl. Anim. Behav. Sci. 185, 86–94.

Pike, A.L., Horwitz, D.F., Lobprise, H., 2015. An open-label prospective study of the use of l-

theanine (Anxitane) in storm-sensitive client-owned dogs. J. Vet. Behav. 10, 324–331.

Rehn, T., Handlin, L., Uvnäs-Moberg, K., Keeling, L.J., 2014. Dogs' endocrine and

behavioural responses at reunion are affected by how the human initiates contact.

Physiol. Behav. 124, 45–53.

Rief, W., Nestoriuc, Y., Weiss, S., Welzel, E., Barsky, A.J., Hofmann, S.G., 2009. Meta-

analysis of the placebo response in antidepressant trials. J. Affect. Disord. 118, 1–8.

Riemer, S., 2019. Not a one-way road – severity, progression and prevention of firework

fears in dogs. PLoS One 14, e0218150. https://doi.org/10.1371/journal.pone.0218150

Sheppard, G., Mills, D.S., 2003. Evaluation of dog-appeasing pheromone as a potential

treatment for dogs fearful of fireworks. Vet. Rec. J. Br. Vet. Assoc. 152, 432–436.

Sherman, B.L., Mills, D.S., 2008. Canine anxieties and phobias: an update on separation

anxiety and noise aversions. Vet. Clin. North Am. Small Anim. Pract. 38, 1081–1106.

Shull-Selcer, E.A., Stagg, W., 1991. Advances in the understanding and treatment of noise

phobias. Vet. Clin. North Am. Small Anim. Pract. 21, 353–367.

Storengen, L.M., Lingaas, F., 2015. Noise sensitivity in 17 dog breeds: Prevalence, breed

risk and correlation with fear in other situations. Appl. Anim. Behav. Sci. 171, 152–160.

Talegón, M.I., Delgado, B.A., 2011. Anxiety disorders in dogs, in: Anxiety Disorders. InTech,

pp. 261–280.

Tavel, M.E., 2014. The placebo effect: the good, the bad, and the ugly. Am. J. Med. 127,

484–488.

Thomas, B.L., Cutler, M., Novak, C., 2012. A modified counterconditioning procedure

prevents the renewal of conditioned fear in rats. Learn. Motiv. 43, 24–34.

Tiira, K., Lohi, H., 2014. Reliability and validity of a questionnaire survey in canine anxiety

research. Appl. Anim. Behav. Sci. 155, 82–92.

Tiira, K., Sulkama, S., Lohi, H., 2016. Prevalence, comorbidity, and behavioral variation in

canine anxiety. J. Vet. Behav. Clin. Appl. Res. 16, 36–44.

Wilson, G.T., Davison, G.C., 1971. Processes of fear reduction in systematic desensitization:

Animal studies. Psychol. Bull. 76, 1.

Wolpe, J., 1958. Psychotherapy by reciprocal inhibition. Stanford University Press., Palo

Alto.