High and asymmetric somato-dendritic coupling of V1 layer 5 neurons independent of behavioural state and visual stimulation

Valerio Francioni[1,2] and Nathalie L. Rochefort[1,2, †]

[1]Centre for Discovery Brain Sciences, School of Biomedical Sciences Edinburgh, Edinburgh, EH8 9XD, United Kingdom.

[2]Simons Initiative for the Developing Brain, University of Edinburgh, Edinburgh, EH8 9XD, United Kingdom.

[†] Corresponding Author:

Nathalie Rochefort

Centre for Discovery Brain Sciences, Biomedical Sciences Edinburgh, Edinburgh, EH8 9XD, United Kingdom.

n.rochefort@ed.ac.uk

Abstract

Active dendritic conductances impact the integration of synaptic inputs. However, the extent to which dendrites act as independent computational subunits *in vivo*, remains unclear. We imaged semi-simultaneously calcium signals in the soma, trunk and distal tuft dendrites of individual layer 5 pyramidal neurons in the awake primary visual cortex. We found highly correlated calcium transients throughout the whole apical tuft: branch-specific activity was rare and limited to the smallest calcium transients. While apical tuft calcium transients were highly correlated to trunk and somatic transients, the frequency of calcium transients was found to decrease in a distance- and amplitude-dependent manner from soma to apical tuft. Visual stimulation and locomotion increased global apical tuft activity but did not alter the coupling between soma and apical tuft. These results suggest that the apical tuft acts as a single computational unit and that its coupling to somatic signals is strong and asymmetric.

Introduction

Cortical layer 5 excitatory neurons are characterized by long apical dendrites receiving inputs from multiple long-range cortical and subcortical connections (Harris and Shepherd, 2015; Ramaswamy and Markram, 2015). In the primary visual cortex (V1), layer 5 pyramidal neurons display selective responses to physical features of visual stimuli, such as the orientation and direction of movement (Kim *et al.*, 2015). In addition, layer 5 neurons activity is modulated by locomotion both in darkness and during visual stimulation (Erisken *et al.*, 2014; Dadarlat and Stryker, 2017; Dipoppa *et al.*, 2018), suggesting that V1 layer 5 neurons integrate inputs from anatomically and functionally distinct pathways (visual and motor-related). However, the integration of visual and locomotion-related inputs and its relationship with somatic output remains unknown.

Dendritic properties were shown to support both distance-dependent filtering and local dendritic non linearities (Grienberger, Chen and Konnerth, 2015; Stuart and Spruston, 2015). Different types of active dendritic currents have been described to support local nonlinear events, primarily driven by NMDA-receptors calcium and sodium- conductances both *in vitro* and *in vivo* (Schiller *et al.*, 1997; Matthew E. Larkum, Zhu and Sakmann, 1999, for review, Grienberger, Chen and Konnerth, 2015; Stuart and Spruston, 2015). These dendritic events are generally associated with calcium influx both locally and globally in the apical tuft of layer 5 neurons (Xu *et al.*, 2012; Hill *et al.*, 2013; Palmer *et al.*, 2014; Cichon and Gan, 2015; Manita *et al.*, 2015; Takahashi *et al.*, 2016). Additionally, the coincident occurrence of back-propagating action potentials and tuft depolarization was shown to generate widespread calcium transients in the apical tuft dendrites (Jaffe *et al.*, 1992; Spruston *et al.*, 1995; Magee and Johnston, 1997; Svoboda *et al.*, 1997; Helmchen *et al.*, 1999; Larkum, Zhu and Sakmann, 1999; Waters *et al.*, 2003; Manita *et al.*, 2015) .

The frequency and dynamics of local dendritic nonlinearities *in vivo* have been a matter of debate. In the motor cortex, one study suggested that individual branches of densely labelled layer 5 neurons selectively respond to specific motor outputs and serve as independent computational subunits (Cichon and Gan, 2015). However, other studies suggest that the entire apical tuft acts as a single, independent computational subunit (Xu *et al.*, 2012; Hill *et al.*, 2013; Ujfalussy *et al.*, 2018). Events described as local dendritically-generated spikes were reported in layer II/III neurons of the visual and somatosensory cortex during sensory

stimulation (Smith *et al.*, 2013; Palmer *et al.*, 2014). In the visual cortex, these results suggested that dendritic spikes triggered by visual input enhance somatic tuning to the preferred orientation (Smith *et al.*, 2013). However, single dendrite recordings prevented conclusions about the spatial extent and the branch-specific nature of the detected events. Other studies reported dendritic calcium spikes in association with backpropagating action potentials during spontaneous activity in the motor cortex (Hill *et al.*, 2013) and sensory stimulation in somatosensory cortex (Svoboda *et al.*, 1997; Helmchen *et al.*, 1999; Kerlin *et al.*, 2018). So far however, the recruitment of local and global dendritic events and their coupling with somatic output, remains largely unexplored in awake behaving animals.

In this study, we used multiplane, two-photon calcium imaging to semi-simultaneously monitor calcium transients in different compartments of individual layer 5 pyramidal neurons in the primary visual cortex (V1) of head-fixed awake behaving mice. We imaged calcium signals in the soma, trunk and distal tuft dendrites, both in darkness and during the presentation of drifting gratings, while the animal was either running or stationary. In agreement with a recently published article (Beaulieu-Laroche *et al.*, 2019), we found that the apical trunk dendritic activity was highly correlated with somatic activity. We extend these results, by showing that the whole apical tuft was highly correlated to trunk activity: the vast majority of dendritic calcium transients in the apical tuft were coincident with global events, with rare cases of branch-specific activity, limited to the smallest amplitude events. In addition, we found that somato-dendritic coupling was asymmetric: while almost all events observed in the tuft were also visible in the soma, around 40% of somatic events attenuated in an amplitude and distance-dependent manner from the soma to the apical tuft. Finally, our results show that neither visual stimulation nor locomotion affected the coupling between somatic and apical tuft dendritic calcium signals.

Results

Highly correlated, widespread calcium signals in apical tuft dendrites of single layer 5 neurons

We imaged changes in fluorescence overtime in apical dendrites of individual layer 5 pyramidal neurons sparsely labelled with the calcium sensor GCaMP6f (Chen *et al.*, 2013) in the primary visual cortex of adult mice. Using single-plane two-photon imaging (at 120 Hz),

we monitored changes of GCaMP6f signals in individual apical tuft dendrites of head-fixed

awake behaving mice that were free to run on a cylindrical treadmill (Figure 1A). Our results

showed highly correlated calcium signals in all apical branches belonging to the same neuron

and imaged in a given field of view (Pearson's correlation coefficient ρ =0.92, SEM = 0.01, n =

25 fields of view recorded from 14 neurons, 6 animals) (Figure 1D). Because distal dendrites

are thinner, more excitable and more compartmentalised than proximal tuft dendrites, we

tested whether the correlation of calcium transients between apical tuft branches decreased

with distance from the apical trunk (nexus). We reconstructed each individual neuron for

which we imaged the apical tuft and quantified the correlation of fluorescence signals across

sibling branches. We found that calcium signals in sibling branches were highly correlated

(average Pearson's correlation coefficient ρ = 0.92) regardless of their branching order (One-

way ANOVA, p = 0.34) (Figure 1E,F). Using semi-simultaneous imaging of dendrites at different

focal planes within the apical tuft of individual layer 5 neurons, we also found highly

correlated calcium signals between proximal and distal tuft branches (average Pearson's

correlation coefficient ρ=0.88, Figure 2D(i)). In addition to the global, widespread calcium

transients, we also observed in individual spines, independent calcium signals that were not

correlated with dendritic signals (Figure 1- figure supplement 1), indicating that in our

experimental conditions, we could resolve independent local calcium signals.

We then assessed whether this widespread, highly correlated apical tuft activity was

modulated by visual stimulation and locomotion. We found that both visual stimulation and

locomotion increased calcium activity in the apical tuft of layer 5 neurons (average ΔF/F0;

repeated measure Two-way ANOVA, p = 0.005, 0.003 and 0.99 for visual stimulation,

locomotion and interaction between both conditions, respectively; n = 14 neurons) (Figure

1H). However, the high correlation of calcium signals between all imaged dendritic branches

was not significantly different between periods of darkness and visual stimulation (drifting

gratings) and between stationary and locomotion periods (Two-way ANOVA, p = 0.43, 0.62

and 0.97 for visual stimulation, locomotion and interaction between both conditions,

respectively). Finally, we found that the selectivity of the responses to the drifting gratings

orientation, quantified by an orientation selectivity index, also did not affect the high

correlation of calcium transients in apical tuft dendrites (Figure 1- figure supplement 3).

In line with these findings, we found that branch-specific activity in the apical tuft was

rare. We quantified the amount of branch-specific activity in all pairs of branches belonging

to the same neuron as the proportion of calcium transients present in one branch and absent

in the other pair's branch. Among the 70 imaged branches across 14 neurons, branch-

specific calcium transients represented less than 3% of the total number of transients. Among

all imaged calcium transients, these local signals were dominated by calcium transients of the

smallest amplitudes (Figure 1J). In addition, neither visual stimulation with drifting gratings

nor locomotion significantly affected this small proportion of branch-specific calcium signals

(Three-way ANOVA, $p < 10^{-15}$ for event amplitude and $p = 0.29$; 0.8 and 0.94, for visual

stimulation, locomotion and interaction between both conditions, respectively. No other

interaction effect was statistically significant) (Figure 1J). We tested the robustness of our

quantification by using different thresholds for the detection of individual calcium transients

(Figure 1- figure supplement 2). We found that varying the threshold did not change our

findings: the number of branch-specific events ranged between 0.8 and 3.8% and remained

limited to transients of the smallest amplitudes (Figure 1- figure supplement 2).

Altogether, our results show that apical dendritic calcium signals in V1 layer 5 neurons

are almost exclusively dominated by highly correlated calcium transients in the whole apical

tuft regardless of the dendritic branching order. Neither visual stimulation nor locomotion

modulated this high correlation of apical tuft dendritic calcium signals.

Calcium signals are highly correlated throughout all compartments of

individual layer 5 neurons

We then assessed whether the global, widespread apical tuft calcium signals were coupled

with somatic activity. For this, we semi-simultaneously imaged GCaMP6s calcium signals at

two focal planes, separated on average by 170 μm in depth, in 4 different pairs of neuronal

compartments from the soma to the apical tuft: Soma-Trunk, proximal Trunk-distal Trunk,

distal Trunk-apical Tuft, proximal Tuft-distal Tuft (Figure 2A,B). We used GCaMP6s that has

higher signal amplitude and signal to noise ratio than GCaMP6f in order to increase the

detection of somatic signals (Chen *et al.*, 2013). We found that calcium transients

simultaneously imaged in each pair of proximal and distal compartments were highly

correlated (Figure 2C,D). Since the rise and decay kinetics were slower in somatic signals than

in dendritic ones, we performed our quantification on isolated transients (see Methods). For

each pair of compartments, we found that the average Pearson's correlation value between

the amplitudes of individual transients was 0.88 between the proximal and distal parts of the

apical tuft, 0.92 between the trunk and the apical tuft and 0.85 between the proximal and

distal trunk. The lowest correlation value was found between the soma and the proximal

trunk (0.74, Figure 2D). For each pair of neuronal compartments, shuffling the calcium

transients in one compartment resulted in an average correlation value of zero (Figure 2D).

Altogether, these results show that calcium transients amplitudes were highly correlated

from the soma to the apical tuft of V1 individual layer 5 neurons.

Frequency of calcium transients decreases in a distance- and amplitude dependent manner from soma to apical tuft

We then tested whether the high correlation of calcium transients' amplitudes between soma

and apical tuft was associated with distance-dependent changes in the frequency of these

events.

We quantified the frequency of calcium events in each pair of proximal and distal

compartments imaged semi-simultaneously (Figure 3A). We found that the frequency of

calcium transients decreased from proximal to distal compartments (Paired t-test, p = $1.1e^{-6}$,

n = 31 pairs of compartments from 19 neurons) by an average of 14% from soma to proximal

trunk, 8% from proximal to distal trunk, 24% from distal trunk to apical tuft and 22% form

proximal tuft to distal tuft. These proportions indicate a decrease of about 40% of calcium

transients from soma to the distal part of the apical tuft (Figure 3B and Figure 3- figure

supplement 1). This result was confirmed by a second data set, in which somatic and apical

tuft calcium transients were imaged independently (at 120 Hz) in individual layer 5 neurons.

We found that the mean frequency of calcium transients in the apical tuft corresponded to

62% of the frequency of events in the corresponding soma (Figure 3- figure supplement 1).

We then checked whether this decrease in frequency depended on the amplitude of

the calcium transients. We calculated the percentage of compartment-specific calcium

transients as a function of their amplitude (Figure 3C). The vast majority of compartment-

specific events were dominated by calcium transients in proximal compartments that were

not detected in distal ones (red trace in figure 3C), while only few calcium transients were

found in distal compartments and not in the corresponding proximal one (blue trace in Figure

3C) (see also Supplementary video 1 and 2). In addition, our results show that smaller

amplitude calcium transients in the proximal compartment were more likely to attenuate

below detection level in the distal compartment compared to larger amplitudes events (red

trace in figure 3C). As a result, compartment-specific events were dominated by small

amplitude calcium transients in proximal compartments (Two-way ANOVA, $p < 10^{-15}$, $p < 10^{-15}$,

$p < 10^{-15}$, for compartment (proximal vs distal), amplitude, and interaction effect,

respectively. Figure 4A). Our results were robust to the threshold we used for detecting

calcium transients (Figure 1- figure supplement 2). When decreasing the threshold for events

detection to up to 30% in the distal compartment, we still observed an amplitude-dependent

decay of calcium events in the proximal compartment (Figure 1- figure supplement 2).

Similarly, when increasing the threshold by 30% for event detection in the proximal

compartment, almost all the events observed distally were also found proximally (Figure 1-

figure supplement 2). Due to the scattering nature of cortical tissue, signal to noise ratio

decreased with imaging depth. As a consequence, the distance-dependent decrease in

frequency may be underestimated.

These results indicate that almost all distal events were found in proximal

compartments. Additionally, these findings show that, at least 40% of somatic calcium

transients attenuate in a distance dependent manner along the apical trunk and distal

dendrites.

Visual stimulation and locomotion do not alter the coupling between soma and apical tuft calcium signals.

We then tested whether visual stimulation and locomotion altered the coupling between the

different compartments of individual layer 5 neurons. For each condition, we plotted the

percentage of compartment-specific calcium transients as a function of their amplitude and

found no significant difference between stationary and locomotion periods and between

darkness and visual stimulation (Two-way ANOVA, $p = 0.57$ and $p = 0.14$ for visual stimulation

and locomotion, respectively, Figure 4B).

Even though smaller events were attenuated with similar probabilities across the four

conditions, calcium transients that do propagate through different neuronal compartments,

may be amplified or attenuated in a condition-dependent manner (Figure 4C). To capture

these nonlinearities, we plotted the amplitude of each detected transient in each

compartment and calculated a residual value as the distance from the linear robust regression

fit (Figure 4C). We then plotted the cumulative distribution of the residual value for every

calcium transient in all 4 pairs of imaged compartments from soma to apical tuft. We found

that, on average, the non-linear changes in calcium transients' amplitude across neuronal

compartments were not significantly different between darkness and visual stimulation nor

between stationary and locomotion periods (Paired t-test p = 0.62 and 0.96 respectively, n =

31 pairs of compartments from 19 neurons, Figure 4D).

These results indicate that the coupling between somatic and apical tuft dendritic

calcium signals remains unaltered by visual stimulation and locomotion under our

experimental conditions.

Discussion

Our results show highly correlated calcium transients in all imaged dendrites of the apical tuft

of individual layer 5 neurons in the primary visual cortex of awake behaving mice: we found

that branch-specific calcium transients were rare and limited to small amplitude transients.

Calcium transients were found to be highly correlated throughout all compartments of

individual neurons (soma, trunk, tuft). However, their frequency was found to decrease in a

distance- and amplitude-dependent manner from soma to apical tuft. Whereas almost all

transients observed in the tuft were present in proximal compartments, the frequency of

calcium transients in apical tuft was about 60% of their frequency in the corresponding soma.

This indicates that a substantial fraction (40%) of somatic calcium signals attenuate toward

the apical tuft. Neither visual stimulation nor locomotion altered the coupling between soma

and apical tuft nor the proportion of branch-specific events.

Limitations of the use of calcium imaging to assess dendritic activity in awake behaving mice

244 Investigations of local dendritic activity in awake behaving mice have so far mainly relied on

245 the use of genetically-encoded calcium indicators (Xu *et al.*, 2012; Cichon and Gan, 2015;

246 Manita *et al.*, 2015; Takahashi *et al.*, 2016; Sheffield, Adoff and Dombeck, 2017; Kerlin *et al.*,

247 2018; Ranganathan *et al.*, 2018; Sheffield and Dombeck, 2019). The analysis of our data set

248 showed that several experimental constraints may mistakenly bias results towards signals

interpreted as local dendritic activity. Dense labelling was already demonstrated to strongly affect correlation values (Xu *et al.*, 2012); even in cases of relatively sparse labelling of individual neurons, contamination from axons or dendrites of other labelled neurons may lead to local signals detected in single dendrites. In our data set, we systematically reconstructed our imaged neurons and excluded regions of interest where overlapping dendrites from other neurons were observed. Movement artefacts may also lead to inaccurate detection of local dendrite signals; these artefacts are more likely to be found during specific behaviour, such as locomotion, and therefore can bias the interpretation of the results. In our data set we used the activity-independent marker tdTomato to correct for motion artefacts and excluded the recordings in which tdTomato marker was not detected. For most of our imaged neurons, baseline fluorescence of GCamP6 could also be used for the same purpose. Another parameter that may cause artefactual local dendritic signals, is the length of the imaged dendritic segments (and therefore the size of the regions of interest used to extract the changes of fluorescence over time). Small regions of interest could include spines not clearly distinguishable from the corresponding dendritic shaft, especially when the spine is located above the dendrite at the imaged focal plane. As a consequence, in small regions of interests, spine signals may be wrongly interpreted as local dendritic signals.

An additional source of bias is linked to the properties of calcium indicators. Differences in calcium buffering properties of individual compartments lead to different decay times of fluorescent calcium indicators between somatic and dendritic compartments (Kerlin *et al.*, 2018; Beaulieu-Laroche *et al.*, 2019). In order to account for this difference, we based our analysis on the detection of peak amplitudes of individual calcium transients. This detection depends on arbitrary thresholds: we thus showed the robustness of our results across different detection thresholds (Figure 1- figure supplement 2). Finally, our results may be biased by the affinity of GCaMP6s. Studies that combined *in vivo* dendritic electrophysiology with calcium imaging have shown that dendritic calcium spikes can reliably be detected with calcium indicators of lower sensitivity than GCaMP6f and GCaMP6s (Helmchen *et al.*, 1999; Xu *et al.*, 2012). However, while it was shown that GCaMP6 could detect individual action potentials in layer 2/3 neurons *in vivo* (Chen *et al.*, 2013), this is not the case for GCaMP6f in layer 5 neurons (Beaulieu-Laroche *et al.*, 2019). These results suggest that in our conditions, the detection of somatic action potentials in layer 5 neurons, may be

280 biased towards larger events, and as a consequence, the attenuation from soma to tuft may
281 be underestimated.

Widespread, correlated calcium transients in apical tuft dendritic branches of layer 5 neurons

Our results show that in the apical tuft, calcium transients were highly correlated regardless
of how distal the imaged branches were from the nexus. Branch-specific calcium
transients were rare and limited to transients of the smallest amplitudes. These findings
suggest that either branch-specific calcium activity is virtually absent and that the entire
apical tuft behaves as a single computational unit, or that branch-specific activity is
systematically coincident with global tuft calcium events and thus could not be detected with
the temporal resolution of GCaMP6f. In addition, these results indicate that active dendritic
integration in the apical tuft is widely recruited across different behavioural and sensory
conditions, not supporting the view of a selective and sparse activity of apical tuft dendrites.
Widespread calcium activity throughout the layer 5 apical tuft may strongly influence
plasticity and learning (Magee and Johnston, 1997; Golding, Staff and Sruston, 2002; Kampa,
Letzkus and Stuart, 2006; Sjöström and Häusser, 2006; Remy and Spruston, 2007; Bittner *et*
al., 2017; Bono and Clopath, 2017; Guerguiev, Lillicrap and Richards, 2017).

Distance dependent attenuation of calcium signals from soma to apical tuft

Our results show an asymmetry in the coupling between somatic and apical tuft calcium
signals. Previous studies have shown that dendritic signals attenuate in a distance-dependent
manner, suggesting that at least some of the calcium events generated in one compartment
would fully attenuate as they reach the next compartment (Stuart, Schiller and Sakmann,
1997; Svoboda *et al.*, 1997; Helmchen *et al.*, 1999; Vetter, Roth and Häusser, 2001; Larkum *et*
al., 2009). To estimate the lower bound proportion of calcium events generated in the tuft
and in the soma, we investigated the proportion of attenuated events in both centrifugal and
centripetal directions. Our results show that while around 40% of somatic calcium transients
decay by the time they reach the apical tuft, only 1.4% of apical tuft calcium events were not
associated with somatic events (Figure 4A). These results indicate that at least 40% of somatic
calcium transients were not triggered by apical tuft calcium events, and that these transients

attenuated from the soma to the apical tuft, suggesting that these calcium transients are associated with back-propagating somatic action potentials. A recent study reported a high somato-dendritic coupling in V1 layer 5 neurons concluding that GCaMP6f events occurred concurrently in soma and dendritic compartments without attenuation from soma to apical tuft (Beaulieu-Laroche *et al.*, 2019). This discrepancy may be explained by differences in imaging conditions including a sparse labelling, which reduces the amount of background signals, and the use of GCaMP6s which has a higher signal amplitude and signal to noise ratio than GCaMP6f indicator, leading to a higher probability of detecting small events in layer 5 somata.

The remaining 60% of somatic calcium signals were highly correlated with global apical tuft calcium signals. This result is consistent with two possible mechanisms: either global apical tuft activation systematically triggers somatic action potentials, or global tuft calcium transients are triggered by back-propagating action potentials alone or in conjunction with tuft synaptic inputs. The distance-dependent decrease in the frequency of calcium transients from soma to apical tuft, suggests that at least some of the transients detected in the tuft correspond to calcium signals associated with back-propagating action potentials. Local synaptic inputs in the tuft may then modulate calcium signals triggered by back-propagating action potentials and facilitate further somatic action potentials firing, consistent with Backpropagation-activated calcium spike (BAC) theory (Larkum, 2013; Manita *et al.*, 2015). Alternatively, apical tuft global calcium signals may be generated by tuft synaptic inputs (Xu *et al.*, 2012; Grienberger, Chen and Konnerth, 2015). Further experiments are needed to resolve these mechanisms, for example using voltage-sensitive dyes (Chavarha *et al.*, 2018; Roome and Kuhn, 2018; Adam *et al.*, 2019) or dendritic electrophysiological recordings (Moore *et al.*, 2017), that would provide the temporal resolution to resolve the different types of dendritic events.

Integration of visual and locomotion-related inputs in layer 5 neurons in awake behaving mice

The prevalence and the dynamics of synaptic inputs received by layer 5 pyramidal neurons strongly vary between *in vitro* and *in vivo* conditions, between anesthetized and awake animals and between passive sensory stimulation and active learning tasks (Xu *et al.*, 2012;

Hong *et al.*, 2018). It is known that *in vivo*, apical tuft dendrites of layer 5 neurons receive a barrage of thousands of synaptic inputs (Stuart and Spruston, 2015). In awake behaving mice, apical tuft dendrites of V1 layer 5 neurons receive synaptic inputs conveying information about external stimuli (visual stimuli) as well as information about internal variables (arousal/attention, motor activity, predictive signals) (Pakan, Francioni and Rochefort, 2018). Our results show that, in our passive viewing conditions, changes in visual inputs (darkness versus drifting gratings) and locomotion-related inputs do not affect the relationship between somatic and apical tuft calcium signals nor the prevalence of branch-specific dendritic events.

It is however possible that during the active learning of a behavioural task, the synaptic inputs associated with this learning process would lead to different mechanisms of dendritic integration than during passive viewing. For example, dendritic integration of multiplicative combinations of sensory and motor inputs in the tuft dendrites of layer 5 pyramidal neurons has been described during an active sensing task in mouse somatosensory cortex (Ranganathan *et al.*, 2018). In addition, it was proposed that changes in neuronal representations during learning would rely on active dendritic signals generating calcium plateau potentials (Bittner *et al.*, 2015; Grienberger *et al.*, 2017). In that case, somato-dendritic coupling may change during the course of learning (Sheffield, Adoff and Dombeck, 2017). This remains to be tested in the visual cortex. Similarly, somato-dendritic coupling may evolve during postnatal development when sensori-motor associations are formed. Recently, it was shown in the mouse primary visual cortex, that dendrites of layer 2/3 neurons increase their coupling during adulthood, as a consequence of decreased responsiveness of dendrite-targeting interneurons to locomotion-related inputs (Yaeger, Ringach and Trachtenberg, 2019). Further investigations are needed to reveal these mechanisms in layer 5 visual cortical neurons.

Materials and Method

Animals

All experiments and procedures involving animals were approved by the University of Edinburgh Animal Welfare and the ethical review board (AWERB) and performed under the appropriate PIL and PPL license from the UK Home Office in accordance with the Animal

(Scientific Procedures) act 1986 and the European Directive 86/609/EEC on the protection of

animals used for experimental purposes.

Adult males and females aged between 8 and 10 weeks obtained from Jackson Laboratory,

ME, USA (B6.Cg-Gt(ROSA)26Sor$^{tm14(CAG-tdTomato)Hze}$/J [RRID:IMSR_JAX:007914]) were used for

experiments. Animals were group caged in groups of 2-6 animals, with a running wheel on a

reverse 12:12 hour light: dark cycle. They were allowed ad libitum access to food and water.

Surgical procedures

Viral delivery of GCaMP6

To obtain sparse labelling of excitatory neurons we used a cre-dependent approach.

AAV1.CamKII.GCaMP6f.WPRE.hGH (Penn Vector core, catalogue No. V4673MI-R) was diluted

1:10000 – 1:20000 while either AAV1.Syn.Flex.GCaMP6f.WPRE.SV40 (Penn Vector Core,

catalogue No. CS0215) or AAV1.Syn.Flex.GCaMP6s.WPRE.SV40 (Penn Vector Core, catalogue

No. CS0408) were diluted 1:10 in the final solution. All dilutions were made in ACSF sterilized

using a 0.2 μm syringe filter. AAV1.Syn.Flex.GCaMP6f.WPRE.SV40 was used for experiments

in the apical tuft and for non-simultaneous recordings of somatic and dendritic activity, while

AAV1.Syn.Flex.GCaMP6s.WPRE.SV40 was used for semi-simultaneous two-planes imaging.

Animals were anaesthetized by inhalation of 4% Isoflurane and maintained anaesthetised

using 1-2% isoflurane. The animal's body temperature was maintained at physiological levels

using a closed-loop, adjustable heating pad monitoring the animal's temperature via a rectal

probe. Eye cream (Bepanthen, Bayer) was applied to protect the eyes from dryness and light

exposure.

After induction of anaesthesia, mice were shaved, mounted onto a stereotaxic frame

(David Kopf instruments, CA, USA) and head-fixed using ear bars. An analgesic was

administered subcutaneously (Vetergesic, buprenorphine, 0.1 mg/kg of body weight). Layer

5 was targeted stereotactically. Injections were performed on the left hemisphere of the brain

using a glass pipette (20 μm tip diameter. Nanoject, Drummond Scientific), backfilled with the

viral preparation, using a nanoject II (Drummond Scientific), at 2 different depths, 650 and

500 μm from brain surface. A total volume of 55.2 nl was injected across the two sites (6x 4.6

nl at each location). Single 4.6 nl injections were spaced 30s from one another to allow

sufficient time for diffusion. After each injection, pipettes were left in situ for an additional

5 min to prevent backflow. The scalp was then sutured (Ethicon, Ethilon polyamide size 6) and

the animal monitored during recovery in a heated cage before returning to its home cage for

2-3 weeks.

Headplate and imaging window

Under anaesthesia (isoflurane), mice were shaved, mounted onto a stereotaxic frame (David

Kopf instruments, CA, USA) and head-fixed using ear bars. Analgesic and anti-inflammatory

drugs were administered subcutaneously (Vetergesic, buprenorphine, 0.1 mg/kg of body

weight; Carpaphen, Carprieve, 5mg/kg of body weight; Dexamethasone, Rapidexon, 2mg/kg

of body weight). For a cranial window over V1, a section of scalp was removed, the underlying

bone was cleansed before a rectangular craniotomy of ~ 2x1.5 mm was made over the left

primary visual cortex. The dura was kept intact. The craniotomy was then sealed with a glass

cover slip and fixed with cyanoacrylate glue. To stabilize the brain during imaging, we used a

triple glass window by stacking three cover-slip glasses (Menzel-Glaser 24x32 mm # 0) on top

of each other. These were glued together using an optically clear UV-cured glue (Norland

Optical adhesive). Overall, the thickness of the glass window inserted through the cranial

window, was comparable to the thickness of the skull at the time of imaging (Smith *et al.*,

2017).

A custom-built metal headplate was fixed on top of the skull with glue and cemented with

dental acrylic (Paladur, Heraeus Kulzer). At the end of the procedure, a single dose of 25ml/kg

of Ringer's solution was injected subcutaneously to rehydrate the animal after the procedure.

The animal was then released from the head fixation and returned to a heated recovery cage,

until full motor capacity was recovered.

In vivo two-photon imaging data acquisition

425 Imaging was performed using a 25x Objective (Olympus). The excitation wavelength of the

426 laser was set to 920 nm. Layer 5 pyramidal neurons were recorded between 500 and 650 μm

427 below the brain surface and followed up to their distal tuft dendrites along the apical trunk.

428 Neurons which had their nucleus filled with GCaMP6 or had blobby-looking dendrites were

429 excluded for recording. At the end of each imaging session, a z-stack was acquired to allow a

more detailed, offline morphological reconstruction of the imaged neuron. Reconstructions were done using the ImageJ plugin Simple Neurite Tracer. Imaging was performed using a 570 nm short-pass dichroic mirror and two single-band pass filters, a 525/50 (Scientifica) for GCaMP6 light absorption, and a 620/60 (Scientifica) for TdTomato absorption.

Habituation and imaging started 2–3 weeks after AAV injection. Mice were habituated to head-fixation in the dark for 45 min and began to run freely on a polystyrene cylinder (20 cm diameter, on a ball-bearing mounted axis). The mice's running speed on the circular treadmill was continuously monitored using an optical encoder (E7P, 250cpr, Pewatron, Switzerland) connected to a data acquisition device (National Instrument, UK) with custom-written software in LabView (National Instrument, UK) and analysed in MATLAB (Mathworks, MA)

Single-plane two-photon imaging data were acquired using a custom-built galvo-resonant scanning system with a Ti:Sapphire pulsing laser (Chameleon Vision-S, Coherent, CA, USA; < 70 fs pulse width, 80 MHz repetition rate) at 120 Hz, with a custom-programmed LabVIEW-based software (version 8.2; National Instruments, UK). Multi-plane data were acquired using an ultra-fast, solid-state, single 100 fs pulse width laser (InSight DeepSee, SpectraPhysiscs, CA, USA) and a FemtoSmart Dual two-photon microscope (Femtonics, Budapest, Hungary). Two focal planes (512x165 pixels), with an average distance of 170 μm in Z, were imaged at a frequency of 96 frames/s (48 frames/s per plane) using a Piezo objective positioner kit (P725.4CA, Physic Instruments, Germany), switching between planes at 9.6 Hz (4.8 Hz per plane).

Visual stimulation

Visual stimuli were generated using the Psychophysics Toolbox package (Brainard, 1997) for MATLAB (Mathworks, MA) and displayed on a custom-modified backlit LED monitor (51 × 29 cm, Dell, UK) placed 20 cm from the right eye, covering 104° × 72° of the visual field. Visual stimulation trials consisted of drifting full-field square-wave gratings for 3 seconds (spatial frequency of 0.05 cycles per degree, 1.5 Hz, 8 equally spaced directions in randomized order, contrast 80%, mean luminance 37 cd/m^2). Drifting gratings were separated from one another by 4 seconds of isoluminant grey periods. Time stamps for the onset of every stimulus were recorded and aligned to imaging frames using custom-built Matlab scripts.

Data Analysis

<u>Image analysis and signal extraction</u>

To correct for brain motion after image acquisition, we used 2D plane translation-based image alignment (SIMA 1.2.0, sequential image analysis; Kaifosh et al., 2014). We 3D reconstructed each imaged neuron and defined regions of interest (ROIs) corresponding to neuronal cell body and dendritic segments manually. Due to sparse labelling, signal contamination was negligible in most cases. In cases ROIs were contaminated by labelled structures from other neurons, we excluded those ROIs from further analysis.

The pixel intensity within each ROI was averaged to create a raw fluorescence time series F(t). Baseline fluorescence F0 was computed for each neuron by taking the fifth percentile of the smoothed F(t) (1 Hz lowpass, zero-phase, 60th-order FIR filter) over each trial (F0(t)), averaged across all trials. As a consequence, the same baseline F0 was used for computing the changes in fluorescence in darkness and during visual stimulation. The change in fluorescence relative to baseline, ΔF/F0 was computed by taking the difference between F and F0(t) and dividing by F0.

Single plane data was acquired at 120 Hz, and subsequently downsampled to 5Hz for signal processing. Multi-plane data were analysed at 4.8 Hz.

<u>Calcium transients analysis</u>

We first estimated noise levels by filtering our ΔF/F0 signal using a 9^{th} order, zero-phase, high-pass filter at 0.6 Hz (Matlab function *filter*). We then estimated the standard deviation of the filtered signal and used a threshold of 2.8 of this standard deviation to detect individual calcium transients, using the built-in Matlab function *findpeaks.* We tested the robustness of our results to different thresholds +/- 30% of the selected threshold (Figure 1- figure supplement 2). For each peak found in any branch or compartment, we defined an event as coincident in another branch or compartment when it occurred in a time window of 3 seconds around the frame where the first peak was originally detected (2 seconds before, 1 after). 99% of coincident events were observed within +/- 0.73 ms. Each event could only be

considered coincident with one event. The amplitude of a calcium event was determined by

taking the difference between the ΔF/F0 amplitude at the frame in which a peak was detected

and a local minimum in a 2 seconds, backward sliding window. Correlation values were

calculated as the Pearson's correlation values between event amplitudes in pairs of branches

or compartments. Shuffling values were obtained by randomly shuffling the order of the

events in one branch or compartment of each pair analysed. Residuals were extracted by

calculating the robust linear regression line between normalised event amplitudes in two

compartments, and then extracting the distance, along the y-axis, of each individual point

from this robust line.

Locomotion analysis

Changes in the position of the cylindrical treadmill (sampled at 12,000 Hz) were interpolated

to match the rate of imaging. To define stationary and locomotion periods we used the

following criteria: Stationary corresponded to periods where the instantaneous speed (as

measured at the 40 Hz sampling rate) was less than 0.1 cm/s. Locomotion corresponded to

periods meeting three criteria: instantaneous speed ≥ 0.1 cm/s, 0.25 Hz lowpass filtered

speed ≥ 0.1 cm/s, and an average speed ≥ 0.1 cm/s over a 2 s window cantered at this point

in time. Any inter-locomotion interval shorter than 500 ms was also labelled as locomotion.

Stationary periods less than 3 s after or 0.2 s before a period of locomotion were removed

from the analysis.

Orientation Selectivity

511   The orientation selectivity index (OSI) was calculated as $(O_{pref}-O_{orth})/(O_{pref}-O_{orth})$ where O_{pref}

512   represents the mean ΔF/F0 value during the presentation of the preferred orientation across

513   trials and O_{orth} represents the mean ΔF/F0 value at the orthogonal orientation compared to

514   preferred. The preferred stimulus of each neuron was the orientation that elicited the

515   maximal response when averaged across all trials.

Statistics

517   All analysis was performed either in Matlab 2017a, or in GraphPad Prism 8. All error bars in

518   the figures represent standard error of the mean. The details of the statistical test we used,

as well the details about what we used as independent samples can be found in the relevant figure legends.

Acknowledgments

We thank the GENIE Program and the Janelia Research Campus, specifically V. Jayaraman, R. Kerr, D. Kim, L. Looger, and K. Svoboda, for making GCaMP6 available. This work was funded by the Wellcome Trust and the Royal Society (Sir Henry Dale fellowship to N.R.), the Marie Curie Actions of the European Union's FP7 program (MC-CIG 631770 to N.R.), the Shirley Foundation, the Patrick Wild Center and the RS MacDonald Charitable Trust Seedcorn Grant (to N.R.), the Simons Initiative for the Developing Brain (to N.R.) and the University of Edinburgh (PhD scholarship to V.F.).

Declaration of Interests

The authors declare no competing interests

References

Adam, Y. *et al.* (2019) 'Voltage imaging and optogenetics reveal behaviour- dependent changes in hippocampal dynamics', *Nature*. Springer US. doi: 10.1038/s41586-019-1166-7.

Beaulieu-Laroche, L. *et al.* (2019) 'Widespread and Highly Correlated Somato-dendritic Activity in Cortical Layer 5 Neurons', *Neuron*. Elsevier Inc., pp. 1–7. doi: 10.1016/j.neuron.2019.05.014.

Bittner, K. C. *et al.* (2015) 'Conjunctive input processing drives feature selectivity in hippocampal CA1 neurons', *Nature Neuroscience*, 18(8), pp. 1133–1142. doi: 10.1038/nn.4062.

Bittner, K. C. *et al.* (2017) 'Behavioral time scale synaptic plasticity underlies CA1 place fields', *Science*, 357(6355), pp. 1033–1036. doi: 10.1126/science.aan3846.

Bono, J. and Clopath, C. (2017) 'Modeling somatic and dendritic spike mediated plasticity at the single neuron and network level', *Nature Communications*. Springer US, 8(1), p. 706. doi: 10.1038/s41467-017-00740-z.

Chavarha, M. *et al.* (2018) 'Fast two-photon volumetric imaging of an improved voltage

indicator reveals electrical activity in deeply located neurons in the awake brain', *BioRXiv*.

Chen, T.-W. *et al.* (2013) 'Ultrasensitive fluorescent proteins for imaging neuronal activity.',
Nature, 499(7458), pp. 295–300. doi: 10.1038/nature12354.

Cichon, J. and Gan, W.-B. (2015) 'Branch-specific dendritic Ca2+ spikes cause persistent
synaptic plasticity', *Nature*, 520(7546), pp. 180–185. doi: 10.1038/nature14251.

Dadarlat, M. C. and Stryker, M. P. (2017) 'Locomotion Enhances Neural Encoding of Visual
Stimuli in Mouse V1', *The Journal of Neuroscience*, 37(14), pp. 3764–3775. doi:
10.1523/JNEUROSCI.2728-16.2017.

Dipoppa, M. *et al.* (2018) 'Vision and Locomotion Shape the Interactions between Neuron
Types in Mouse Visual Cortex.', *Neuron*. doi: 10.1016/j.neuron.2018.03.037.

Erisken, S. *et al.* (2014) 'Effects of Locomotion Extend throughout the Mouse Early Visual
System', *Current Biology*. Elsevier Ltd, 24(24), pp. 1–9. doi: 10.1016/j.cub.2014.10.045.

Golding, N. L., Staff, N. P. and Spruston, N. (2002) 'Dendritic spikes as a mechanism for
cooperative   long-term   potentiation',   *Nature*,   418(6895),   pp.   326–331.   doi:
10.1038/nature00854.

Grienberger, C. *et al.* (2017) 'Inhibitory suppression of heterogeneously tuned excitation
enhances spatial coding in CA1 place cells', *Nature Neuroscience*, 20(3), pp. 417–426. doi:
10.1038/nn.4486.

Grienberger, C., Chen, X. and Konnerth, A. (2015) 'Dendritic function in vivo', *Trends in
Neurosciences*. Elsevier Ltd, 38(1), pp. 45–54. doi: 10.1016/j.tins.2014.11.002.

Guerguiev, J., Lillicrap, T. P. and Richards, B. A. (2017) 'Towards deep learning with segregated
dendrites', *eLife*, 6, pp. 1–37. doi: 10.7554/eLife.22901.

Harris, K. D. and Shepherd, G. M. G. (2015) 'The neocortical circuit: themes and variations',
Nature Neuroscience, 18(2), pp. 170–181. doi: 10.1038/nn.3917.

Helmchen, F. *et al.* (1999) 'In vivo dendritic calcium dynamics in deep-layer cortical pyramidal
neurons', *Nature Neuroscience*. doi: 10.1038/14788.

Hill, D. N. *et al.* (2013) 'Multibranch activity in basal and tuft dendrites during firing of layer 5
cortical neurons in vivo', *Proceedings of the National Academy of Sciences*, 110(33), pp.
13618–13623. doi: 10.1073/pnas.1312599110.

Hong, Y. K. *et al.* (2018) 'Sensation, movement and learning in the absence of barrel cortex',
Nature. Springer US, pp. 542–546. doi: 10.1038/s41586-018-0527-y.

Jaffe, D. B. *et al.* (1992) 'The spread of Na+ spikes determines the pattern of dendritic Ca2+
entry into hippocampal neurons', *Nature*, 357(6375), pp. 244–246. doi: 10.1038/357244a0.

Kampa, B. M., Letzkus, J. J. and Stuart, G. J. (2006) 'Requirement of dendritic calcium spikes
for induction of spike-timing-dependent synaptic plasticity', *The Journal of Physiology*, 574(1),

pp. 283–290. doi: 10.1113/jphysiol.2006.111062.

Kerlin, A. *et al.* (2018) 'Functional clustering of dendritic activity during decision-making',
bioRxiv. doi: 10.1101/440396.

Kim, E. J. *et al.* (2015) 'Three Types of Cortical Layer 5 Neurons That Differ in Brain-wide
Connectivity and Function', *Neuron*. Elsevier Inc., 88(6), pp. 1253–1267. doi:
10.1016/j.neuron.2015.11.002.

Larkum, M. (2013) 'A cellular mechanism for cortical associations: An organizing principle for
the cerebral cortex', *Trends in Neurosciences*. Elsevier Ltd, 36(3), pp. 141–151. doi:
10.1016/j.tins.2012.11.006.

Larkum, M. E. *et al.* (2009) 'Synaptic integration in tuft dendrites of layer 5 pyramidal neurons:
A new unifying principle', *Science*, 325(5941), pp. 756–760. doi: 10.1126/science.1171958.

Larkum, M. E., Zhu, J. J. and Sakmann, B. (1999) 'A new cellular mechanism for coupling inputs
arriving at different cortical layers', *Nature*. doi: 10.1038/18686.

Magee, J. C. and Johnston, D. (1997) 'A synaptically controlled, associative signal for Hebbian
plasticity in hippocampal neurons', *Science*. doi: 10.1126/science.275.5297.209.

Manita, S. *et al.* (2015) 'A Top-Down Cortical Circuit for Accurate Sensory Perception', *Neuron*.
Elsevier Inc., 86(5), pp. 1304–1316. doi: 10.1016/j.neuron.2015.05.006.

Moore, J. J. *et al.* (2017) 'Dynamics of cortical dendritic membrane potential and spikes in
freely behaving rats', *Science*, 355(6331), p. eaaj1497. doi: 10.1126/science.aaj1497.

Pakan, J. M., Francioni, V. and Rochefort, N. L. (2018) 'Action and learning shape the activity
of neuronal circuits in the visual cortex', *Current Opinion in Neurobiology*, 52, pp. 88–97. doi:
10.1016/j.conb.2018.04.020.

Palmer, L. M. *et al.* (2014) 'NMDA spikes enhance action potential generation during sensory
input', *Nature Neuroscience*. Nature Publishing Group, 17(3), pp. 383–390. doi:
10.1038/nn.3646.

Ramaswamy, S. and Markram, H. (2015) 'Anatomy and physiology of the thick-tufted layer 5
pyramidal neuron', *Frontiers in Cellular Neuroscience*, 9(June), pp. 1–29. doi:
10.3389/fncel.2015.00233.

Ranganathan, G. N. *et al.* (2018) 'Active dendritic integration and mixed neocortical network
representations during an adaptive sensing behavior', *Nature Neuroscience*. Springer US. doi:
10.1038/s41593-018-0254-6.

Remy, S. and Spruston, N. (2007) 'Dendritic spikes induce single-burst long-term
potentiation', *Proceedings of the National Academy of Sciences*, 104(43), pp. 17192–17197.
doi: 10.1073/pnas.0707919104.

Roome, C. J. and Kuhn, B. (2018) 'Simultaneous dendritic voltage and calcium imaging and
somatic recording from Purkinje neurons in awake mice', *Nature Communications*. Springer

US, 9(1), p. 3388. doi: 10.1038/s41467-018-05900-3.

Schiller, J. *et al.* (1997) 'Calcium action potentials restricted to distal apical dendrites of rat
neocortical pyramidal neurons', *The Journal of Physiology*, 505(3), pp. 605–616. doi:
10.1111/j.1469-7793.1997.605ba.x.

Sheffield, M. E. and Dombeck, D. A. (2019) 'Dendritic mechanisms of hippocampal place field
formation', *Current Opinion in Neurobiology*. Elsevier Ltd, 54, pp. 1–11. doi:
10.1016/j.conb.2018.07.004.

Sheffield, M. E. J., Adoff, M. D. and Dombeck, D. A. (2017) 'Increased Prevalence of Calcium
Transients across the Dendritic Arbor during Place Field Formation', *Neuron*. Elsevier Inc.,
96(2), pp. 490-504.e5. doi: 10.1016/j.neuron.2017.09.029.

Sjöström, P. J. and Häusser, M. (2006) 'A Cooperative Switch Determines the Sign of Synaptic
Plasticity in Distal Dendrites of Neocortical Pyramidal Neurons', *Neuron*, 51(2), pp. 227–238.
doi: 10.1016/j.neuron.2006.06.017.

Smith, I. T. *et al.* (2017) 'Stream-dependent development of higher visual cortical areas',
Nature Neuroscience, 20(2), pp. 200–208. doi: 10.1038/nn.4469.

Smith, S. L. *et al.* (2013) 'Dendritic spikes enhance stimulus selectivity in cortical neurons in
vivo', *Nature*, 503(7474), pp. 115–120. doi: 10.1038/nature12600.

Spruston, N. *et al.* (1995) 'Activity-dependent action potential invasion and calcium influx into
hippocampal   CA1   dendrites',   *Science*,   268(5208),   pp.   297–300.   doi:
10.1126/science.7716524.

Stuart, G. J. and Spruston, N. (2015) 'Dendritic integration: 60 years of progress', *Nature
Neuroscience*, 18(12), pp. 1713–1721. doi: 10.1038/nn.4157.

Stuart, G., Schiller, J. and Sakmann, B. (1997) 'Action potential initiation and propagation in
rat neocortical pyramidal neurons', *The Journal of Physiology*, 505(3), pp. 617–632. doi:
10.1111/j.1469-7793.1997.617ba.x.

Svoboda, K. *et al.* (1997) 'In vivo dendritic calcium dynamics in neocortical pyramidal
neurons', *Nature*, 385(6612), pp. 161–165. doi: 10.1038/385161a0.

Takahashi, N. *et al.* (2016) 'Active cortical dendrites modulate perception', *Science*,
354(6319), pp. 1587–1590. doi: 10.1126/science.aah6066.

Ujfalussy, B. B. *et al.* (2018) 'Global and Multiplexed Dendritic Computations under In Vivo-
like Conditions', *Neuron*. doi: 10.1016/j.neuron.2018.08.032.

Vetter, P., Roth, A. and Häusser, M. (2001) 'Propagation of Action Potentials in Dendrites
Depends on Dendritic Morphology', *Journal of Neurophysiology*, 85(2), pp. 926–937. doi:
10.1152/jn.2001.85.2.926.

Waters, J. *et al.* (2003) 'Supralinear Ca 2+ Influx into Dendritic Tufts of Layer 2/3 Neocortical
Pyramidal Neurons In Vitro and In Vivo', *The Journal of Neuroscience*, 23(24), pp. 8558–8567.

doi: 10.1523/JNEUROSCI.23-24-08558.2003.

Xu, N. *et al.* (2012) 'Nonlinear dendritic integration of sensory and motor input during an
active sensing task.', *Nature*. Nature Publishing Group, 492(7428), pp. 247–51. doi:
10.1038/nature11601.

Yaeger, C. E., Ringach, D. L. and Trachtenberg, J. T. (2019) 'Neuromodulatory control of
localized dendritic spiking in critical period cortex', *Nature*. Springer US, pp. 100–104. doi:
10.1038/s41586-019-0963-3.

663
664

Figure 1. Highly correlated activity in the whole apical tuft of layer 5 pyramidal neurons.
A) Schematic of the methodological approach. Calcium transients were recorded either in multiple apical tuft dendrites in a single focal plane, or semi-simultaneously in different compartments of the neuron (Soma, Trunk and Tuft). At the end of each imaging session, a z-stack was recorded for post-hoc reconstruction of each imaged neuron. **B)** Single plane imaging of the apical tuft branches of an individual layer 5 neuron. Calcium transients of the apical tuft branches belonging to one neuron are shown in yellow and calcium transients of tuft branch belonging to a different neuron is shown in red. Scale bars, 0.3 ΔF/F0 (normalised to max), 20 s. **C)** Pearson's correlation matrix between the calcium activity of the branches shown in B. Branches that belong to the same neuron (neuron 1) have a mean Pearson's correlation of 0.96, while the branch that belongs to the different neuron (neuron 2) has a mean Pearson's

correlation value of 0.13 with neuron 1's branches. **D)** Mean Pearson's correlation value ρ for each imaged field of view and corresponding shuffled data (Paired t-test, p = $3.5e^{-15}$; n = 25 field of view; coming from 14 neurons; 6 animals; 70 branches). **E)** Schematic of our definition of branching orders. **F)** Two representative traces of sibling branches belonging to the 5th and 8th (left panel, labelled) branching order and a quantification of the Pearson's correlation ρ for sibling tuft branches of different branching orders (right panel). (For 1st, 2nd, 3rd, 4th and more than 4th branching order, One-way ANOVA, p = 0.34; n = 5; 6; 4; 4; 4 pairs of branches, respectively). Scale bars, 0.3 ΔF/F0 (normalised to max), 20 s. **G)** Representative traces of 3 tuft branches belonging to the same neuron while the animal was either stationary or locomoting (lower panel) during either darkness or the presentation of drifting grating (purple segment). Scale bars, 12 cm/s, 0.3ΔF/F0 (normalised to max), 20 s. **H)** Mean ΔF/F0 for apical tuft dendrites during darkness and visual stimulation while the animal was either stationary of locomoting. Bar graph is normalised to visual stimulation during stationary condition. Both visual stimulation and locomotion significantly increase the mean ΔF/F0 of tuft branches without any interaction effect (Repeated Measures Two-way ANOVA on log-transformed data, p = 0.0053, 0.0026 and 0.99 for the visual stimulation, locomotion and interaction effects respectively, n = 14 neurons). **I)** Pearson's correlation coefficients ρ for the apical tuft branches of individual neurons during the four conditions (stim and stationary, stim and loco, dark and stationary, dark and loco). (Two-way ANOVA, p = 0.43, 0.62 and 0.97 for visual stimulation, locomotion and interaction effects; n =16; 14; 5 and 7 field of views respectively). **J)** Weighted mean of the proportion of branch-specific events as a function of event's amplitude for the four different conditions (lower left panel). Neither visual stimulation nor locomotion nor an interaction effect, increased the number of branch specific events (Three-way ANOVA, p = 0.29; 0.8 and 0.94, respectively; p < 10^{-15} for event amplitude). *Upper left panel*, frequency distribution of the events' amplitudes detected in the apical tuft branches. *Right panel*, pie chart showing that on average 3% of the total events were detected in only one branch.

Figure 2. High correlation of calcium transients from soma to apical tuft dendrites of individual layer 5 neurons. A) Morphological reconstruction of a neuron imaged at two focal planes semi-simultaneously. *Left panel*, coronal view of the imaged neuron. Red and blue dotted lines indicate the planes of imaging. *Middle left panel*, horizontal view of the same neuron. Branches highlighted in red and blue, represent the same dendritic branches observed in the middle right panel. *Right panel*, each trace represents the calcium activity of an individual branch shown in the middle panels. **B**) Schemata of the compartments imaged simultaneously,

(i) tuft-tuft, (ii) trunk-tuft, (iii), trunk-trunk, (iv) soma-trunk. pTrunk, dTrunk and pTuft, dTuft

indicate proximal and distal portions of the trunk and the apical tuft, respectively. **C)** *Left panel*,

representative calcium events recorded in two neuronal compartments semi-simultaneously as

shown in B. *Right panel*, scatter plot showing the correlation of the amplitude of calcium

transients in each compartment imaged simultaneously, and shown in B. Each dot represents a

calcium transient. Filled dots are the same transients shown in the left panel and indicated by

numbers. Scale bars 0.3 ΔF/F0 (normalised to max), 10 s. **D)** Pearson's correlation values ρ for

each pair of compartments imaged semi-simultaneously and compared to its corresponding

shuffled data (Paired t-test, pTuft-dTuft(i), p = $1.6e^{-6}$, n = 6 pairs; Trunk-Tuft(ii), p = $8e^{-7}$, n =

9; pTrunk-dTrunk(iii), p = $4.4e^{-4}$, n = 5; Soma-Trunk(iv), p = $4.4e^{-6}$, n = 11).

Figure 3. Frequency of calcium transients decreases in a distance and amplitude-dependent dependent manner from the soma to the apical tuft. A) Schemata of the compartments imaged simultaneously. **B)** *Left panel*, representative ΔF/F0 traces of calcium activity recorded semi-simultaneously in two different compartments of individual neurons. Asterisks indicate calcium transients detected in the proximal compartment (red trace) and not detected in the distal one (blue trace). *Right panel*, frequency of detected calcium transients, normalised to the proximal compartment. Individual lines represent individual neurons. Scale bars 0.25 ΔF/F0 (normalised to max), 20 s. **C)** Proportion of compartment-specific events as a function of calcium transients' amplitude. In red, proportion of events detected in the proximal

compartment only. In blue, proportion of events detected in the distal compartment only. *Upper right panel*, in red and blue the frequency histogram of calcium transients amplitudes detected in the proximal and distal compartments, respectively. Thick line represents the weighted mean proportion. Shaded area represents the weighted SEM for each bin (0.05 size). For all compartments, event amplitude, compartment (proximal vs distal) and an interaction between these two factors significantly affected the proportion of compartment-specific events (Two-way ANOVA, pTuft-dTuft (i), $p < 10^{-15}$, $p = 7.8 \times 10^{-5}$, $p = 6.2 \times 10^{-8}$, for event amplitude, proximal versus distal compartment and interaction between amplitude and compartment, respectively, n = 6 pairs; Trunk-Tuft(ii), $p < 10^{-15}$, $p = 3.4 \times 10^{-9}$, $p < 10^{-15}$, n = 9; pTrunk-dTrunk(iii), $p = 1.3 \times 10^{-13}$, $p = 2.4 \times 10^{-4}$, $p = 7 \times 10^{-3}$, n = 5; Soma-Trunk(iv), $p < 10^{-15}$, $p < 10^{-15}$ and $p < 10^{-15}$, n = 11).

Figure 4. Locomotion and visual stimulation do not alter the relationship between somatic and dendritic activity in individual layer 5 pyramidal neurons. A) Mean proportion of compartment-specific events as a function of calcium transient amplitude. Data points from figure 3C(i-iv) have been averaged together. In red, events found in the proximal, but not in the distal compartment. In blue, events that are found in the distal, but not proximal compartment. Event amplitude, compartment (proximal vs distal) and an interaction between these two factors significantly affected the proportion of compartment-specific events (Two-way ANOVA, $p < 10^{-15}$, $p < 10^{-15}$, $p < 10^{-15}$, n = 31 pairs of compartments from 19 neurons). **B**) Neither visual stimulation nor locomotion alter the curve describing amplitude-dependent decay of events in the proximal-to-distal direction (Two-way ANOVA, p = 0.57 and p = 0.14

for visual stimulation and locomotion, respectively). **C**) *Upper panel*, representative traces of the proximal (red) and distal (blue) compartment imaged semi-simultaneously, during stationary and locomotion periods (lower trace shows speed). Scale bars represent 0.25 $\Delta F/F0$ (normalised to max), 12 cm/s, 20 s. *Lower panel*, scatter plot between the amplitudes of calcium transients for the $\Delta F/F0$ trace shown above. Individual dots represent individual calcium transients. Filled dots are the same as shown in the upper panel. The red line represents the robust linear regression fit. For each transient, a residual from the robust linear regression was calculated. **D**) Cumulative distribution function for the residuals calculated for each pairs of compartments and for each condition. Neither visual stimulation nor locomotion alter the relationship between somatic and dendritic activity in individual layer 5 neurons (Paired t-test for the effect of visual stimulation and locomotion on residuals: Soma-Trunk(iv), p = 0.33 and 0.30, respectively, n = 11 pairs; pTrunk-dTrunk(iii), p = 0.34 and 0.93, n = 5; Trunk-Tuft(ii), p = 0.5 and 0.95, n = 9; pTuft-dTuft(iv), p = 0.83 and 0.08, n = 6).

Figure 1- figure supplement 1. Independent single spine activity in the apical tuft dendrites of layer 5 neurons. A) Field of view showing the apical tuft branches of an individual GCamP6-labelled layer 5 neuron. **B**) ΔF/F0 traces for the individual spines shown in A during a single, 65 seconds-long trial. Error bars indicate 0.3 ΔF/F0 (normalised to max), 10 s.

Figure 1- figure supplement 2. Impact of threshold for calcium transients detection. A)
To assess the amount of branch / compartment specific events, for every transient detected in
one branch / compartment, we assessed whether there was another corresponding event in the
second branch / compartment (see Methods). *Upper panel*, ΔF/F0 traces of two sibling
branches in the apical tuft branches of a single layer 5 pyramidal neurons. Asterisks indicate
two events which were not detected using the selected threshold but were detected lowering
the threshold by 30%. *Lower panel*, zoomed image of the dotted area in the upper panel. To
assess the resilience of our observed effects to different thresholds, we assessed the amount of
branch / compartment specific events, keeping the selected threshold in one branch /
compartment (red-dotted line) and varying the threshold in the second branch / compartment
from +30% to -30% (light, medium and dark grey lines, respectively). Scale bars: upper panel,

0.3 ΔF/F0 (normalised to max), 5 s; lower panel: 0.03 ΔF/F0, 1s. **B**) Number of branch-specific

events detected as a function of the amplitude and threshold selected. **C**) Number of

compartment-specific events detected as a function of the amplitude and threshold selected.

Figure 1- figure supplement 3. High correlation of calcium transients in apical tuft

dendrites, independent of somatic orientation selectivity. A) Morphological reconstruction

of three apical tuft branches that were imaged during the presentation of eight drifting gratings.

B) ΔF/F0 traces of the branches shown in A, in response to drifting gratings of four different

orientations (3 s gratings interleaved by 4 s isoluminant grey screen). Scale bars 0.3 ΔF/F0

(normalised to max), 10 s. **C**) Polar plot quantifying the responses of these 3 branches to the

eight oriented gratings. **D**) Mean Pearson's correlation value ρ of all imaged branches per

neuron, as a function of the orientation selectivity of the corresponding soma (each dot

represents a neuron; n = 23 neurons). The straight black line represents the best fit.

Figure 3- figure supplement 1. Decreased frequency of calcium events from soma to apical tuft. A) Frequency distribution of the depths at which neurons have been imaged. The red dotted line indicates the median (528 μm from the dura). **B**) Calculation of the proportion of somatic events, decaying from the soma to the most distal part of the tuft. Pairs of compartments (soma-trunk, trunk-trunk, trunk-tuft, tuft-tuft) were recorded at a distance of 170 μm (see Methods). As a consequence, the average amplitude-dependent decay function (lower panel, same as figure 4A) represents the decay of calcium transients within 170 μm. To calculate how many events attenuate from the soma to the most distal part of the apical tuft (528 μm apart, on average), we projected the decay measured within 170 μm, to a distance of 528 μm. We multiplied the amplitude-dependent decay function by the scaling factor 3.1 (528/170) and we applied this scaled function to the event distribution (upper panel) at each bin (0.05 ΔF/F0). The event distribution was derived by averaging all the calcium events in all the proximal compartments (see distribution of event amplitudes for each proximal compartment in red in figure 3). Using this approach, we calculated that the proportion of events that decay from the soma to the tuft was 39%, for the data set in which pairs of compartments were imaged semi-simultaneously. **C**) Proportion of events in the tuft, relative to the corresponding soma, for neurons in which both compartments were imaged non-simultaneously.